



**CHALMERS**  
UNIVERSITY OF TECHNOLOGY

## Effects of nitrogen on growth and carbohydrate formation in *Porphyridium cruentum*

Downloaded from: <https://research.chalmers.se>, 2024-04-10 23:09 UTC

Citation for the original published paper (version of record):

Razaghi, A., Godhe, A., Albers, E. (2014). Effects of nitrogen on growth and carbohydrate formation in *Porphyridium cruentum*. *Central European Journal of Biology*, 9(2): 156-162.  
<http://dx.doi.org/10.2478/s11535-013-0248-z>

N.B. When citing this work, cite the original published paper.

# Effects of nitrogen on growth and carbohydrate formation in *Porphyridium cruentum*

## Research Article

Ali Razaghi<sup>1</sup>, Anna Godhe<sup>1</sup>, Eva Albers<sup>2\*</sup>

<sup>1</sup>Biological and Environmental Sciences, University of Gothenburg,  
SE-405 30 Gothenburg, Sweden

<sup>2</sup>Chemical and Biological Engineering - Industrial Biotechnology,  
Chalmers University of Technology, 412 96 Gothenburg, Sweden

Received 12 March 2013; Accepted 05 August 2013

**Abstract:** The microalga *Porphyridium cruentum* (Rhodophyta) has several industrial and pharmaceutical uses, especially for its polysaccharide production. This study aimed to investigate the influence of nitrogen levels as reflected by altered N:P ratios on the production and content of biomass and carbohydrate. N:P molar ratios were altered in batch cultures to range from 1.6 to 50 using the Redfield ratio of 1:16 as reference. Algal growth (estimated as final cell number, biomass concentration and maximum specific growth rate) was negatively affected at low N:P ratios. The optimal N:P ratio for growth was identified at 35-50, with specific growth rates of 0.19 day<sup>-1</sup> and maximum cell concentrations of 59·10<sup>8</sup> cells L<sup>-1</sup> and 1.2 g dry weight of biomass L<sup>-1</sup>. In addition, variation in cell size was seen. Cells with larger diameters were at higher N:P ratios and smaller cells at lower ratios. The cellular carbohydrate content increased under reduced nitrogen availability. However, because accumulation was moderate at the lowest N:P ratio, 0.4 g per g dry weight biomass compared to 0.24 at the Redfield ratio of 16:1, conditions for increased total carbohydrate formation were identified at the N:P ratios optimal for growth. Additionally, carbohydrates were largely accumulated in late exponential to stationary phase.

**Keywords:** Rhodophyta • Red algae • Redfield ratio • Nitrogen-to-phosphorous ratio

© Versita Sp. z o.o.

## 1. Introduction

In a bio-based society, sustainable production processes should be based on renewable raw materials. Starch-based biomass of phototrophic organisms helps to fulfill this criterion. In this respect, studies on how culture conditions, such as nutrient availability, that would enhance cellular carbohydrate content are important. The microalga *Porphyridium* sp. (Rhodophyta) is a potential source for several products, such as fatty acids and lipids [1,2], pigments [3], and cell-wall polysaccharides [4]. These polysaccharides are sulphated and their structure gives rise to some unique properties that could lead to a broad range of industrial and pharmaceutical applications [5]. Polysaccharides surround the algal cell as an amorphous capsule. In the marine species *P. cruentum*, this capsule consists of glucose, galactose, xylose, glucuronic acid and methyl-glucuronic acid as sugar monomers [6]. The

outer part of the capsule can be partly excreted to the surroundings, i.e. exopolysaccharides, thereby increasing the viscosity of the medium [7]. Additionally, *P. cruentum* biomass contains starch [7] and cellular contents of carbohydrates of up to 57 % have been reported [8]. Thus, the combined amount of carbohydrates in biomass and exopolysaccharides of this microalga could potentially provide the carbon source for fermentation processes, such as bioethanol production [9].

Culture conditions affect carbohydrate content in microalgae. Limited levels of macro-nutrients, such as nitrogen and phosphorous, in the growth medium have been shown to affect formation of carbohydrates in several algae species. For instance, nitrogen limitation increased carbohydrate formation in *Chlamydomonas mexicana* [10] and *Chlorella* spp. [11] and increased starch production in *Tetraselmis subcordiformis* [12]. Phosphorus limitation induced an increase of carbohydrate formation

\* E-mail: albers@chalmers.se

in *Ankistrodesmus falcatus* [13]. Thus, as a mechanism of nutrient depletion, increased accumulation of carbohydrates is expected in the stationary growth phase. This has been documented for, e.g. *Chlamydomonas mexicana* [10], *Tetraselmis gracilis* [14] and *Porphyridium aeruginum* [15]. Another effect of variable nitrogen concentration is that cell-wall polysaccharides become increasingly soluble in nitrogen deficient media, and the majority of the cells-wall polysaccharides were found as exopolysaccharides [16]. Thus, to achieve optimal carbohydrate production, nitrogen availability needs to be balanced to give a simultaneously high carbohydrate accumulation and a reasonable growth rate and also to relate the concentration of nitrogen to the available phosphorus, *i.e.* the N:P ratio.

Nitrogen availability largely affects algae growth as indicated by the fact that nitrogen effluents intensify algal blooms [17]. Accordingly, in cultures of *Porphyridium* sp., high nitrogen concentrations were favorable for growth, whereas nitrogen deficiency limited growth [16,18]. For example in *P. purpureum* (syn. *P. cruentum*), nitrogen limitation led to cessation of growth with significantly reduced chlorophyll and phycoerythrin contents [19]. In this respect, the Redfield ratio (C:N:P 106:16:1 molar composition) [20] has been proposed as a general optimal ratio for algal growth. However, recently the oversimplification of this ratio has been questioned, as the optimal N:P ratios seem to be species specific and in the range of 8.2-45.0 [21].

Algal cellular content may vary in response to light intensity and the type of metabolism, *i.e.* photosynthetic or respiratory, and therefore also varies in response to light:dark cycles [22]. In *Porphyridium*, high light intensity stimulated starch and exopolysaccharide formation [7], whereas continuous illumination only enhanced exopolysaccharide formation [23]. For cost-effective production, the algal cultures are preferably illuminated by natural sunlight, which provides illumination with light:dark cycles. *P. cruentum*, grown outdoors in a semi-turbidostat culture, reached a cellular carbohydrate content of 31-58% (on ash free biomass basis) [24].

Given the potential to develop this species as a resource, an extended survey of the effects of N:P ratios is needed. Previous studies of *P. cruentum* and the effect of varying N:P ratio was limited to three ratios in the range of 5-20 [18]. The aim of this study was to compare a broader range of N:P ratios to determine the effect of nutrient availability on the growth properties and carbohydrate content and production, and to identify the optimal molar N:P ratio for *P. cruentum* growth. This was done using batch cultures illuminated in dark/light cycles with N:P ratios ranging from 2-50, using the Redfield ratio as a reference.

## 2. Experimental Procedures

### 2.1 Strain, medium and cultivation conditions

*Porphyridium cruentum* (strain GUMACC 25, UTEX 161) was obtained from Gothenburg University's Microalgae Algal Culture Collection, Sweden. The medium, f/2 [25], was prepared from filtered, autoclaved natural seawater with five times higher concentrations of sodium phosphate, trace metals, and vitamins compared to the original formula to promote good algal growth and ensure other nutrients did not become limited. Altered N:P ratios (1.6-50) were achieved by changing the nitrate concentration using 0.29-9.1 mmol L<sup>-1</sup> of NaNO<sub>3</sub>, while keeping the phosphate concentration constant at 0.18 mmol L<sup>-1</sup>, giving a range below (denoted as nitrogen limited) and above (denoted as nitrogen excess) the Redfield ratio of 16:1 mol N per mol P (used as the reference). The standard f/2 medium has a N:P ratio of 22. Cells were grown in batch cultures using culture flasks with ventilation caps (Sarstedt) at 23±1°C and 150 rpm agitation. The cultures were illuminated by fluorescent cool-white lamps (color code 840) at an irradiance of 98 µE m<sup>-2</sup> s<sup>-1</sup> (7000 lux) with an 18:6 hour light:dark cycle. Samples for total carbohydrates and cell concentration analyses were collected once a day during the dark phase. Two sets of experiments were conducted, one using cultures of 40 mL medium in 50 mL flasks, and the other using 150 mL medium in 250 mL flasks. This gave a volume to illuminated surface area of 1.6 and 0.86 mL cm<sup>-2</sup>, respectively. These experiments continued until the stationary phase was reached at days 14 and 12, respectively.

### 2.2 Analyses of cells and total carbohydrates

Cell concentration was estimated during cultivation using *in vivo* fluorescence of chlorophylls in 100 µL of cell suspension using black 96 well microtiter plates (FluoroNunc, Nunc Inc.). Fluorescence from chlorophylls was measured using a FLUOstar Omega plate reader (BMG LABTECH) held at a constant temperature of 25°C, equipped with optical filters for excitation at 400-480 nm and emission at 600-680 nm. Measurements on medium only were made in each run and used to correct for background. Maximum specific growth rates were calculated using linear regression of the natural logarithm of the *in vivo* fluorescence data against time during the exponential growth phase [26].

Accurate cell numbers (cells L<sup>-1</sup>) at the stationary phase were determined by counting cells in a Bürker chamber (C-Chip disposable hemocytometer DHC-B01, Digital Bio).

Biomass concentration ( $\text{g L}^{-1}$ ) was obtained using the dry weight of cells collected from 5 mL of culture broth on a  $0.45 \mu\text{m}$  hydrophilic polyethersulfonate filter (Sartorius Stedim Biotech, Germany). The filter was dried for 15 minutes in a microwave oven at 125 W, equilibrated in a desiccator overnight, and then weighed. An equivalent volume of media was filtered and the readings were used as background [27].

Cell morphology was studied using an inverted microscope (Leica DMI4000B) equipped with a CCD camera (Leica DFC360 FX) at  $100\times$  magnification (objective Leica HCX PL APO  $100\times$  NA 1.40). Average cell size for each N:P-ratio was estimated by determining the diameter of 90 cells from all replicates using the size tool of the accompanying software (Leica AF6000 E Software).

Total carbohydrates were analyzed using the phenol-sulfuric acid method [28] for the entire culture broth in triplicate. Glucose was used as the standard, and measurements on medium in each run were used as background. The assay was performed in microtiter plates. Absorbance was measured at 488 nm in a FLUOstar Omega plate reader (BMG LABTECH).


## 2.3 Statistics

A two tailed Student's *t*-test was used to test the null hypothesis that there was no difference between the different N:P ratios. Data was tested for significance against the reference (N:P ratio at 16). For deviant comparisons, statements are given with the data. The average standard deviation was calculated from the mean of variances of replicates within each set of experiments.

## 3. Results

### 3.1 Batch growth of *P. cruentum* at different N:P ratios

The effects of altered nutrient levels on *P. cruentum* growth and accumulation of carbohydrates in biomass was studied in a first set of cultivation using media with a large range of molar N:P ratios. The growth curves displayed the expected characteristics with a lag phase during the first one or two days, an exponential growth phase, a stationary phase when growth ceased, and finally a decay phase when cells were dying during the last few days (Figure 1A). The exponential growth phase ranged from the first to seventh day for cultures at N:P ratios of 16 and higher, whereas at nitrogen limitation the lag phase was longer (Figure 1A). The maximum cell concentration was lower at nitrogen limitation (N:P of 1.6 and 4.9) and higher at nitrogen excess (N:P of



**Figure 1.** *Porphyridium cruentum* cells in batch cultures (50 mL flasks) at different molar N:P ratios comparing nitrogen limitation at 1.6 (●), 4.9 (▲), and 11 (■); nitrogen excess at 24 (○), 35 (△), and 50 (□); and the Redfield ratio of 16:1 (X). A. Cell concentration estimated by *in vivo* fluorescence (f.u. = fluorescence units) on a logarithmic scale. B. Total carbohydrates ( $\text{g L}^{-1}$  broth). C. Carbohydrate content ( $\mu\text{g}$  glucose equivalents [glc eq.] per fluorescence units [f.u.]); representative data are shown (N:P of 11, 24, 35 and 50 are similar to the reference). Means for the triplicate cultivations at each time point are shown with  $\pm$  standard deviation as error bars.

35 and 50) when compared to the reference condition of 16:1.

Similarly, the total carbohydrate concentration was lower at nitrogen limitation and higher at nitrogen excess (Figure 1B). Cellular carbohydrate content increased slowly during the initial exponential growth phase, while a rapid increase in cellular carbohydrate

content was seen during the late exponential phase at day 4–5 (Figure 1C). Once the stationary phase was reached, carbohydrate content remained constant until the end of the experiment. In cultures with N:P ratios of 1.6 and 4.9, carbohydrate accumulation started later at day 6 marking the beginning of the stationary phase (Figure 1C). However, all conditions reached a similar maximum carbohydrate content of approximately  $26 \mu\text{g f.u.}^{-1}$ , except for the N:P ratio of 1.6, which was significantly lower at  $15 \mu\text{g f.u.}^{-1}$  ( $P < 0.01$ ,  $df = 19$ ).

The low cell concentrations observed at the lowest nitrogen availability of 1.6 and 4.9 were associated with low maximum specific growth rates (Figure 2A). Correspondingly, nitrogen excess at N:P ratios of 35 and 50 promoted a maximum specific growth rate nearly two times faster compared to the specific growth rate of  $0.12 \text{ day}^{-1}$  at the Redfield ratio.

At the end of the experiment for the first set of cultures (day 14), cells at the two highest N:P ratios were significantly heavier, and at the two lowest N:P ratios were significantly lighter (Figure 2B). Furthermore, there was a non-significant tendency for the cell diameters to be larger at nitrogen excess for N:P ratios of 35 and 50, and smaller at nitrogen limitation for N:P ratios of 1.6 and 4.9 (Figure 2C).


### 3.2 Cellular properties of *P. cruentum* at the stationary phase after growth at different N:P ratios

In the second set of cultures using larger volumes, limited nitrogen availability at an N:P ratio of 1.6 significantly reduced cell density. Similar to the first set of cultures, both cell number and biomass were significantly higher at nitrogen excess (N:P ratios of 35 and 50) compared to the reference of 16:1 (Table 1).

Total carbohydrate concentration followed a similar pattern to cell densities. However, only under nitrogen limiting conditions, at an N:P ratio of 4.9, a higher carbohydrate accumulation were strongly indicated at  $0.32 \text{ g g}^{-1}$  dry weight compared to the reference of 16:1 (Table 1). In spite of this, the carbohydrate content for the 1.6:1 ratio was even higher ( $0.40 \text{ g g}^{-1}$  dry weight) but yielded a non-significant comparison to the reference of 16:1, as the standard deviation was very large due to low cell densities. By contrast, no differences were found for the cellular carbohydrate content at the different N:P ratios (data not shown). Average content was  $42 \pm 6 \text{ pg glc eq. cell}^{-1}$ .

## 4. Discussion

Culture conditions that provide fast growth and high carbohydrate production for the red microalga



**Figure 2.** Cellular properties of *Porphyridium cruentum* cells in batch culture (50 mL flasks) at different molar N:P ratios, comparing nitrogen limitation at 1.6, 4.9, and 11; nitrogen excess at 24, 35, and 50; and the Redfield ratio at 16:1. Means for the triplicate cultures are shown. A. Maximum specific growth rate (days<sup>-1</sup>) at the exponential phase calculated from *in vivo* fluorescence data. Error bars show the average standard deviation (9.44%) estimated using all data, with 19 degrees of freedom. B. Cellular dry weight (μg f.u.<sup>-1</sup>) at the end of the experiment (f.u. = fluorescence units of *in vivo* fluorescence). Error bars show the average standard deviation (8.05%) estimated using all data giving 19 degrees of freedom. C. Cell diameter (μm) at the end of the experiment. The means of 90 cells from the triplicate cultures are shown, with standard deviation as error bars. Micrographs showing representative cells from N:P ratios 4.9, 16 and 35 are included with a scale bar representing 2 μm. Data significant at \*\*\* $P < 0.01$  and \*\* $P < 0.05$  and with strong indications at \* $P < 0.1$  compared to N:P ratio of 16.

*P. cruentum* were studied and described in this report. For optimal production of a cellular component, both

N:P ratio (N-mol P-mol <sup>-1</sup> )	Number of cells (x10 <sup>8</sup> cells L <sup>-1</sup> )	Biomass concentration (g dw L <sup>-1</sup> )	Total carbohydrate concentration (g glc eq. L <sup>-1</sup> )	Carbohydrate content (g glc eq. g dw cells <sup>-1</sup> )
1.6	2.8±0.8***	0.04±0.01***	0.02±0.004***	0.40±0.14
4.9	32.7±5.5	0.47±0.10	0.15±0.02	0.32±0.03*
11	33.7±3.8	0.48±0.02	0.15±0.02	0.29±0.03
16	29.9±3.9	0.51±0.06	0.13±0.01	0.24±0.03
24	32.2±2.8	0.54±0.03	0.13±0.01	0.25±0.02
35	59.4±3.3***	1.18±0.08***	0.22±0.01***	0.20±0.02
50	58.7±1.7***	1.22±0.15*	0.23±0.01***	0.20±0.03

**Table 1.** Cellular characteristics of *Porphyridium cruentum* at stationary phase (250 ml culture flasks). Data representative of the stationary phase are given - maximum number of cells (x10<sup>8</sup> cells L<sup>-1</sup>), biomass concentration (g dry weight [dw] L<sup>-1</sup>) and total carbohydrates (g glucose equivalents [glc eq.] L<sup>-1</sup>) at different molar N:P ratios of 1.6, 4.9, and 11 (nitrogen limitation), 24, 35, and 50 (nitrogen excess), and the Redfield ratio at 16:1. Means for the triplicate cultures and all days during the duration of the stationary phase (day 5-10) are shown with ± standard deviation.

Data significant (df = 10) at \*\*\*  $P < 0.01$  and with strong indications at \* $P < 0.1$  compared to N:P ratio of 16.

cellular abundance and total biomass formed are important. In our study, the maximum carbohydrate per cell was observed under nitrogen limited conditions with low overall cell density. Thus, the total amount of carbohydrate per volume of culture is a better variable to examine when identifying optimal culture conditions. While the carbohydrate content of cells only varied to a certain extent, the increase in content was severely counteracted by the fact that less biomass was formed; thus leading to an overall lower formation of carbohydrate at nitrogen limitation. Optimal production was thus identified during nitrogen excess.

Similar to other studies on *Porphyridium* strains [18,23], we found that high nitrogen availability, as reflected by the high N:P ratios, was very beneficial for growth. However, increasing the level of nitrogen too much causes a limitation of other nutrients, such as phosphorous [29], trace metals and vitamins. Under such conditions, growth will be negatively affected. Other factors that might limit phototrophic based growth are reduced light availability due to shading, agglomeration, and poor gas exchange rates with limited CO<sub>2</sub> availability.

The influence of nutrient limitation has been previously observed on accumulation of both carbohydrates and lipids [11,30]. Nutrient limitation also imposes other physiological changes in algal cells. By limiting nitrogen, cell volume of the green alga, *Ankistrodesmus falcatus*, decreased and displayed lower density. By contrast, limiting the availability of phosphorous resulted in the opposite effect [13]. Here, we observed effects on cell size in response to nutrient availability, with the smallest cell weight and diameters seen at reduced nitrogen availability, and the largest at increased availability. To some extent, this can be

explained by altered macromolecular composition and also by the fact that nitrogen limitation implies a decline in photosynthetic performance and remodeling of the photosynthetic machinery [31]. Additionally, nitrogen deprivation inhibits growth; only committed cells divide and the progeny is smaller.

Another explanation, possibly valid for *Porphyridium* spp., is that at conditions with small cells *i.e.* under nitrogen limitation, cell wall polysaccharides become increasingly soluble [18,23]. The expulsion of polysaccharides into the fluid could cause the cell size reduction and simultaneously increase the viscosity of the media. Though media viscosity was not measured, an increase in viscosity was previously observed during the ageing of the culture [7]. The composition of the exopolysaccharides is species dependent and consists of monosaccharides, methylated monosaccharides and sugar acids [32]. To the best of our knowledge no studies have been performed on how nitrogen availability alters the composition of exopolysaccharides. Such a study would be important in understanding how nitrogen limitation alters the proportion of exopolysaccharide to intracellular carbohydrate content.

By contrast, phosphorus abatement at higher N:P ratios blocks the cell cycle preventing the synthesis of nucleic acids and cell division. Consequently, cell size will be larger [29,33]. Additionally, at higher nitrogen concentrations, the less soluble cell wall polysaccharides add to an increased cell size. At higher N:P ratios the increased cell size allows for greater accumulation of carbohydrates in the cell. This is indicated by the similar cellular carbohydrate content of 42 pg glc eq. cell<sup>-1</sup> found for all N:P ratios.

The effect of nutrient limitation occurs when cells enter the stationary phase and accumulation



of carbohydrates is induced. This phenomenon was observed here and manifested by a 10-15 fold increase in the carbohydrate content independent of the N:P ratio when entering the stationary phase, as compared to the content for cells in the exponential growth phase (Figure 1C). However, as previously reported [16], an additional accumulation of carbohydrates during the stationary phase is expected. However, in our cultures only a small increase in cellular carbohydrates could be seen under some of the conditions. The carbohydrate concentration found in our study was lower (about ten times) than previous studies with the same species [7,18]. This difference is explained by the fact that the previous study cultivation was performed using 2 - 3% CO<sub>2</sub> in the inlet air which result in higher biomass formation (2-7 g dry weight L<sup>-1</sup> and 3-7.5·10<sup>7</sup> cells ml<sup>-1</sup>) and hence higher carbohydrate levels compared to our study that used only atmospheric air. On the other hand, the carbohydrate content found in our study, around 0.3 g g<sup>-1</sup> dry weight biomass or 42 pg cell<sup>-1</sup>, is comparable to the results of the aforementioned studies (25-44% and 40-60 pg cell<sup>-1</sup>). Thus, limitation in carbon supply reduced the possible carbohydrate production in our batch cultures and hence usage of cultivation aerated with elevated levels of carbon dioxide is desired for optimal production.

## References

- [1] Ahern T.J., Katoh S., Sada E., Arachidonic acid production by the red alga *Porphyridium cruentum*, *Biotechnol. Bioeng.*, 1983, 25, 1057-1070
- [2] Oh S.H., Han J.G., Kim Y., Ha J.H., Kim S.S., Jeong M.H., et al., Lipid production in *Porphyridium cruentum* grown under different culture conditions, *J. Biosci. Bioeng.*, 2009, 108, 429-434
- [3] Kathiresan S., Sarada R., Bhattacharya S., Ravishankar G.A., Culture media optimization for growth and phycoerythrin production from *Porphyridium purpureum*, *Biotechnol. Bioeng.*, 2007, 96, 456-463
- [4] Arad S.M., Levy-Ontman O., Red microalgal cell-wall polysaccharides: biotechnological aspects, *Curr. Opin. Biotechnol.*, 2010, 21, 358-364
- [5] Patel A.K., Laroche C., Marcati A., Ursu A.V., Jubeau S., Marchal L., et al., Separation and fractionation of exopolysaccharides from *Porphyridium cruentum*, *Bioresour. Technol.*, 2012, In Press, doi: 10.1016/j.biortech.2012.1012.1038
- [6] Heaney-Kieras J., Chapman D.J., Structural studies on the extracellular polysaccharide of the red alga, *Porphyridium cruentum*, *Carbohydr. Res.*, 1976, 52, 169-177
- [7] Arad S., Adda M., Cohen E., The potential production of sulfated polysaccharides from *Porphyridium*, *Plant Soil*, 1985, 89, 117-127
- [8] Becker E.W., *Microalgae: biotechnology and microbiology*, Cambridge University Press, Cambridge, 1994
- [9] John R.P., Anisha G.S., Nampoothiri K.M., Pandey A., Micro and macroalgal biomass: a renewable source for bioethanol, *Bioresour. Technol.*, 2011, 102, 186-193
- [10] Kroen W.K., Raynburn W.R., Influence of growth status and nutrients on extracellular polysaccharide synthesis by the soil agla *Chlamydomonas mexicana* (Chlorophyceae), *J. Phycol.*, 1984, 20, 253-257
- [11] Brányiková I., Marsalková B., Doucha J., Brányik T., Bisová K., Zachleder V., et al., Microalgae-novel highly efficient starch producers, *Biotechnol. Bioeng.*, 2011, 108, 766-776
- [12] Yao C., Ai J., Cao X., Xue S., Zhang W., Enhancing starch production of a marine green microalga *Tetraselmis subcordiformis* through nutrient limitation, *Bioresour. Technol.*, 2012, 118, 438-444
- [13] Kilham S.S., Kreeger D.A., Goulden C.E., Lynn S.G., Effect of nutrient limitation on biochemical

## Acknowledgements

We acknowledge the Olle Engkvist Byggmästare foundation (Stockholm, Sweden) for financial support.

- constituents of *Ankistrodesmus falcatus*, *Freshwater. Biol.*, 1997, 38, 591-596
- [14] Lourenco S.O., Lanfer Marquez U.M., Mancini-Filho J., Barbarino E., Aider E., Changes in biochemical profile of *Tetraselmis gracilis* I. Comparison of two culture media, *Aquaculture*, 1997, 148, 153-168
- [15] Ramus J., The production of extracellular polysaccharide by unicellular red alga *Porphyridium aerugineum*, *J. Phycol.*, 1972, 8, 97-111
- [16] Arad S.M., Friedman O.D., Rotem A., Effect of nitrogen on polysaccharide production in a *Porphyridium* sp., *Appl. Environ. Microbiol.*, 1988, 54, 2411-2414
- [17] Carstensen J., Henriksen P., Heiskanen A.S., Summer algal blooms in shallow estuaries: Definition, mechanisms, and link to eutrophication, *Limnol. Oceanogr.*, 2007, 52, 370-384
- [18] Adda M., Merchuk J.C., Arad S., Effect of nitrate on growth and production of cell-wall polysaccharide by the unicellular red alga *Porphyridium*, *Biomass*, 1986, 10, 131-140
- [19] Levy I., Gantt E., Development of photosynthetic activity in *Porphyridium purpureum* (Rhodophyta) following nitrogen starvation, *J. Phycol.*, 1990, 26, 62-68
- [20] Redfield A.C., The biological control of chemical factors in the environment, *Am. Sci.*, 1958, 46, 205-221
- [21] Klausmeier C.A., Litchman E., Daufresne T., Levin S.A., Optimal nitrogen-to-phosphorus stoichiometry of phytoplankton, *Nature*, 2004, 429, 171-174
- [22] MacIntyre H.L., Cullen J.J., Using cultures to investigate the physiological ecology of microalgae, In: Andersen R.A., Ed., *Algal culturing techniques*. Elsevier Academic Press, London, UK, 2005, 287-326
- [23] Thepenier C., Gudín C., Studies on optimal conditions for polysaccharide production by *Porphyridium cruentum*, *World J. Microbiol. Biotechnol.*, 1985, 1, 257-268
- [24] Vonshak A., Cohen Z., Richmond A., The feasibility of mass cultivation of *Porphyridium*, *Biomass*, 1985, 8, 13-25
- [25] Andersen R.A., Ed. *Algal culturing techniques*. Elsevier Academic Press, London, UK, 2005
- [26] Tunzi M.G., Chu M.Y., Bain R.C., In vivo fluorescence, extracted fluorescence, and chlorophyll concentrations in algal mass measurements, *Water Res.*, 1974, 8, 623-635
- [27] Lavens P., Sorgeloos P., Manual on the production and use of live food for aquaculture, *FAO Fisheries Technical Papers T361*, FAO, Rome, 1996, <ftp://ftp.fao.org/docrep/fao/003/w3732e/w3732e00.pdf>
- [28] Herbert D., Phipps P.J., Strange R.E., Chemical analysis of microbial cells, In: Norris J.R., Ribbons D.W., Eds., *Methods in microbiology*. Academic Press, London, 1971, 209-344
- [29] Lien T., Knutsen G., Phosphate as a control factor in cell division of *Chlamydomonas reinhardtii*, studied in synchronous culture, *Exp. Cell. Res.*, 1973, 78, 79-88
- [30] Roessler P.G., Environmental control of glycerolipid metabolism in microalgae: Commercial implications and future research directions, *J. Phycol.*, 1990, 26, 393-399
- [31] Young E.B., Beardall J., Photosynthetic function in *Dunaliella tertiolecta* (Chlorophyta) during a nitrogen starvation and recovery cycle, *J. Phycol.*, 2003, 39, 897-905
- [32] Percival E., Foyle R.A.J., Extracellular polysaccharides of *Porphyridium cruentum* and *Porphyridium aerugineum*, *Carbohydr. Res.*, 1979, 72, 165-176
- [33] Lien T., Knutsen G., Synchronous cultures of *Chlamydomonas reinhardtii*. Synthesis of repressed and derepressed phosphatase during the life cycle, *Biochim. Biophys. Acta.*, 1972, 287, 154-163
- [34] Arrigo K.R., Marine microorganisms and global nutrient cycles, *Nature*, 2005, 437, 349-355