

Glucose repression in Saccharomyces cerevisiae

Downloaded from: https://research.chalmers.se, 2024-04-28 04:31 UTC

Citation for the original published paper (version of record):

Kayikci, Ö., Nielsen, J. (2015). Glucose repression in Saccharomyces cerevisiae. FEMS Yeast Research, 15(6). http://dx.doi.org/10.1093/femsyr/fov068

N.B. When citing this work, cite the original published paper.

research.chalmers.se offers the possibility of retrieving research publications produced at Chalmers University of Technology. It covers all kind of research output: articles, dissertations, conference papers, reports etc. since 2004. research.chalmers.se is administrated and maintained by Chalmers Library



doi: 10.1093/femsyr/fov068 Advance Access Publication Date: 23 July 2015 Minireview

MINIREVIEW

Glucose repression in Saccharomyces cerevisiae

Ömur Kayikci^{1,2} and Jens Nielsen^{1,2,3,*}

¹Department of Biology and Biological Engineering, Kemivägen 10, Chalmers University of Technology, SE41296 Gothenburg, Sweden, ²Novo Nordisk Foundation Center for Biosustainability, Chalmers University of Technology, SE41296 Gothenburg, Sweden and ³Novo Nordisk Foundation Center for Biosustainability, Technical University of Denmark, DK2970 Hørsholm, Denmark

We dedicate this paper the late Professor Jure Piskur, a friend and colleague.

One sentence summary: The role of Snf1 signaling in glucose repression and carbon metabolism in Saccharomyces cerevisae.

Editor: Monique Bolotin-Fukuhara

ABSTRACT

Glucose is the primary source of energy for the budding yeast *Saccharomyces cerevisiae*. Although yeast cells can utilize a wide range of carbon sources, presence of glucose suppresses molecular activities involved in the use of alternate carbon sources as well as it represses respiration and gluconeogenesis. This dominant effect of glucose on yeast carbon metabolism is coordinated by several signaling and metabolic interactions that mainly regulate transcriptional activity but are also effective at post-transcriptional and post-translational levels. This review describes effects of glucose repression on yeast carbon metabolism with a focus on roles of the Snf3/Rgt2 glucose-sensing pathway and Snf1 signal transduction in establishment and relief of glucose repression.

Keywords: carbon metabolism; Snf1 signaling; carbon catabolite repression

INTRODUCTION

When glucose is accessible the yeast Saccharomyces cerevisiae prefers a fermentative metabolism despite presence of oxygen, and represses respiration, use of alternative carbon sources as well as gluconeogenesis (Klein, Olsson and Nielsen 1998; Rolland, Winderickx and Thevelein 2002). This repressive effect of glucose is transmitted to the cellular machinery by interlinked regulatory interactions and signaling pathways. These coordinative molecular activities mainly exert their effects at the transcriptional level, but they are also operative at posttranscriptional and post-translational levels.

Since S. cerevisiae primarily prefers glucose as a carbon source, sensing of extracellular and metabolized glucose levels is important for coordination of yeast carbon metabolism. Yeast cells adjust diverse cellular activities according to glucose levels detected extra- and intracellularly. The Snf3/ Rgt2 signaling pathway is a sensory cascades present in yeast for detecting extracellular glucose levels (Kaniak et al. 2004). Through this pathway, the cell can sense extracellular glucose levels and use this to regulate glucose uptake, and hereby trigger glucose repression. The Snf1 protein kinase signaling is central to functionality of glucose repression and to balance cellular energy levels. Snf1 kinase has a dual role in glucose repression, both as an activator and as a repressor. High glucose concentrations render Snf1 inactive which leaves the transcription factor Mig1 non-phosphorylated and hence being present in the nucleus where it exerts repression, together with Ssn6/ Tup1 complex, of genes involved in the utilization of alternative carbon sources (Gancedo 1998; Hedbacker and Carlson 2008). On the other hand, when glucose becomes limited Snf1 is active and phosphorylates Mig1 allowing release of glucose repression and expression of glucose-repressed genes. Beside its central role in regulating expression of glucose-repressed genes, Snf1 kinase has direct interaction with the transcriptional apparatus, and it has

^{*}Corresponding author: Department of Biology and Biological Engineering, Kemivägen 10, Chalmers University of Technology, SE41296 Gothenburg, Sweden. Tel: +46 (0) 31 772 3804; E-mail: nielsenj@chalmers.se

Received: 12 March 2015; Accepted: 20 July 2015

[©] FEMS 2015. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0/), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.



Figure 1. Snf3/ Rgt2 glucose-sensing pathway. Snf3/ Rgt2 glucose-sensing pathway dictates expression levels of hexose transporter (HXT) genes for optimal uptake of glucose at various concentrations.

been implicated in chromatin modification (Kuchin, Treich and Carlson 2000). Moreover, as a part of its role in energy homeostasis Snf1 regulates metabolic enzymes involved in fatty acid metabolism and carbohydrate storage as well as a role affecting GCN4 translation and hence amino acid biosynthesis (Usaite et al. 2009; Hedbacker and Carlson 2010; Zhang et al. 2011).

Snf1 is the yeast AMP-activated kinase (AMPK), a regulatory kinase that is highly conserved in eukaryal cells. Understanding its function and regulation is therefore of broad and general interest, and much knowledge about AMPK has been acquired through the use of *S. cerevisiae* as a model organism (Petranovic and Nielsen 2008; Petranovic *et al.* 2010). Understanding glucose repression is, however, also of utmost importance for the development of biotechnological processes where it is desirable to redirect the carbon fluxes towards products of interests (Chen and Nielsen 2013; Dai *et al.* 2015; Li and Borodina 2015). Due to the high relevance of glucose repression in yeast, we therefore here review key pathways involved in this complex process in yeast.

Snf3/Rgt2 signaling

Snf3/Rgt2 signaling pathway mainly senses varying concentrations of available glucose in the environment. The Snf3 and Rgt2 sensors belong to the HXT (HeXose transporters) gene family together with the Hxt1–17 and Gal2 proteins that all but Hxt12 can transport glucose (and other hexoses) but each with a different affinity for glucose (Wieczorke *et al.* 1999). Although structurally similar to hexose transporters, Snf3 and Rgt2 cannot transport glucose (Özcan, Dover and Johnston 1998). Snf3 senses low levels of extracellular glucose, while Rgt2 detects high levels of the sugar (Ozcan and Johnston 1999; Ozcan 2002). Intracellular signals generated upon detecting accessible amounts of glucose coordinate transcriptional regulation and expression of Hxt

proteins (Fig. 1). Low-affinity transporters, such as Hxt1, are expressed and activated when glucose is abundant while under such conditions expression of high-affinity transporters, such as Hxt7, are repressed (Ozcan 2002). Snf3 and Rgt2 are also likely to sense internal-to-external ratio of glucose concentrations to adjust glucose uptake and maintain intracellular glucose homeostasis (Karhumaa, Wu and Kielland-Brandt 2010). Regulation of transcriptional activity through the Snf3/Rgt2 pathway allows S. cerevisiae to finely coordinate glucose uptake in response to environmental availability of this hexose sugar. When extracellular glucose is sensed by Snf3 and Rgt2 transmembrane proteins, the membrane-attached type I casein kinases Yck1 and Yck2 are activated. These active kinases are required for degradation of Mth1 and Std1, two paralogous regulatory proteins recruited to the plasma membrane (Schmidt et al. 1999; Moriya and Johnston 2004). Although Mth1 and Std1 were thought to be directly phosphorylated by Yck1/2 kinases for degradation, more recent data show that Mth1 is degraded in the nucleus independent of Yck1/2 localization (Pasula et al. 2010). So, according to this finding the glucose sensors transmit the glucose signal to a yet unidentified protein to promote phosphorylation and degradation of Mth1. Upon phosphorylation, Mth1 and Std1 are targeted for ubiquitination and proteosome degradation (Spielewoy et al. 2004). Furthermore, when glucose is abundant, Mig1 represses MTH1 expression to maintain glucose repression of the HXT genes. In contrast, as Std1 is being degraded its expression is increased to ensure efficient expression of the HXT genes when glucose is exhausted (Kim and Johnston 2006). Upon degradation of Mth1 and Std1, protein kinase A hyperphophorylates and dislodges Rgt1, transcriptional repressor of glucoseinduced genes, from DNA (Palomino, Herrero and Moreno 2006). Glucose-induced translocation of PKA to the nucleus allows hyperphosphorylation of Rgt1 (Griffioen et al. 2000; Kim and Johnston 2006; Roy et al. 2013, 2014). This allows for expression of



Figure 2. The Snf1 protein kinase is central player in glucose repression pathway. The Snf1 protein kinase regulates glucose repression at transcriptional level by activating or inactivating expression of gluconeogenic genes and genes involved in respiration.

HXT genes and optimal uptake of glucose. Depletion of glucose renders Mth1 and Std1 available for Rgt1 interaction, which conceals PKA phosphorylation sites on Rgt1 and as a result the repressor remains bound to promoters suppressing expression of HXT genes when glucose is unavailable (Flick *et al.* 2003). Snf1 phosphorylation of Rgt1 triggers the repressor activity of Rgt1 and its propensity to bind DNA (Palomino, Herrero and Moreno 2006). This interaction between Rgt1 and Snf1 kinase is critical for graded derepression of HXT expression and plays an important role in overall glucose repression.

SNF1 signaling

When glucose in the environment is exhausted, yeast cells switch their metabolism from fermentation to respiration, and activate mechanisms and components for utilization of alternative carbon sources. This usually occurs in the late exponential phase of a batch culture, where the cells are often referred to as being 'derepressed'. Snf1 plays a central role in this metabolic shift by regulating a range of repressors and activators. Availability of glucose regulates activity of this central player in glucose repression. During growth on optimal glucose levels, Snf1 is inactive and excluded from the nucleus. This allows for a major downstream target of Snf1 the transcription repressor Mig1 to be non-phosphorylated and in the nucleus. Conversely, a drop in the glucose concentration (to about 0.2%) activates Snf1 which in turn phosphorylates and deactivates Mig1 relieving glucose repression (Fig. 2) (Piskur and Compagno 2014).

The yeast Snf1 is a widely conserved serine/theronine kinase in eukaryotes required for cellular energy homeostasis. Like the mammalian AMPK, Snf1 kinase complex in S. *cerevisiae* has a heterotrimeric structure with an alpha catalytic subunit (Snf1), a gamma regulatory subunit (Snf4) and one of three beta subunits (Sip1, Sip2, Gal83) (Carlson, Osmond and Botstein 1981; Amodeo, Rudolph and Tong 2007). When glucose levels are high, Snf1 kinase complex is inactive due to autoinhibition as a result of interaction between the N-terminal catalytic domain and C-terminal regulatory domains of Snf1 (Celenza and Carlson 1989; Jiang and Carlson 1996; Ludin, Jiang and Carlson 1998; Leech *et al.* 2003). Low glucose concentrations remove this autoinhibition and promote interactions between Snf4 regulatory subunit and Snf1 catalytic subunits. Additionally, phosphorylation of the conserved residue at Thr²¹⁰ is also required to activate Snf1 (McCartney and Schmidt 2001). In contrary to AMPK, Snf1 is not regulated by AMP but by ADP, which ensures a direct link between energy metabolism and this key regulator (Mayer *et al.* 2011).

Sak1, Elm1 and Tos3 are three protein kinases known to phosphorylate Thr210 in the Snf1 activation loop. These upstream kinases have overlapping functions since only deletion of all three abolishes Snf1 activation (Hong et al. 2003; Nath, McCartney and Schmidt 2003; Sutherland et al. 2003). So far, there has not been a report on how the glucose signal regulates the activating kinases of Snf1. Although each activating kinase contributes differently under different carbon source availability, Sak1 has the most stable interaction with Snf1; it is suggested to be the major activating kinase at conditions of growth on alternative carbon sources. The beta subunits also provide specificity for a particular upstream activating kinase under different conditions (McCartney, Rubenstein and Schmidt 2005). The three scaffolding (beta) subunits also regulate localization of Snf1 kinase. Sip1 directs Snf1 to vacuoles, while Sip2 keeps the enzyme in the cytoplasm and Gal83 play a role in nuclear localization of Snf1 (Vincent et al. 2001; Hedbacker, Hong and Carlson 2004). Each subunit has been indicated to have a distinct yet overlapping role in regulation of transcription and cellular metabolism. For example, Gal83 and Sip2 both play a major role in activating gluconeogenic genes and genes involved with the glyoxylate cycle while Sip1 is mainly involved in the regulation of genes associated with nitrogen metabolism (Zhang, Olsson and Nielsen 2010).

Another level of control is dephoshorylation of Snf1, which is carried out by Glc7/ Reg1 phosohotases (PP1) (Feng et al. 1991; Tu and Carlson 1995). This level of control has been hypothesized to be the main regulator of Snf1 activity since phosphorylation of Snf1 by upstream kinases (Sak1, Elm1 and Tos3) has not been

affected by changes in glucose concentrations (Rubenstein et al. 2008). On the other hand, dephosphorylation of the kinase has been shown to be correlated with glucose availability in the environment (Sanz et al. 2000). Although activity of the Reg1/Glc7 phosphatase remained unchanged in response to glucose fluctuation in the environment, it has been hypothesized that the accessibility of Snf1 for dephosphorylation by the phosphatases changes (Rubenstein et al. 2008). Subsequent work, however, revealed that addition of glucose causes a rapid increase in Reg1 activity leading to inactivation of Snf1 (Castermans et al. 2012). This process could be part of the mechanism that transmits the information about the glucose level to Snf1 kinase. In addition to detecting the shift in glucose availability to establish glucose repression, Snf1 has also been suggested to monitor absolute glucose levels (Bendrioua et al. 2014). With the activation of Snf1 by ADP, there is, however, also a direct coupling of Snf1 activity with the energy status of the cell, and this is clearly related to the glucose metabolism.

Transcriptional effects on carbon metabolism

Upon activation, Snf1 kinase interacts with a number of transcription factors, activating some while suppressing others. Mig1 transcriptional repressor is a major downstream target of Snf1 phosphorylation. When phosphorylated by Snf1, this transcription repressor is deactivated and released from DNA allowing expression of glucose-repressed genes, mainly genes for utilization of alternative carbon sources. Mig1 mediates suppression of glucose-repressed genes together with the general repressor complex Ssn6/Tup1 (Treitel and Carlson 1995; Lutfiyya et al. 1998). Mig1 also interacts with Hxk2 to suppress glucose-induced genes. Besides its metabolic role as glucose, kinase Hxk2 also affects transcriptional regulation of glucoserepressible genes. When glucose is abundant Hxk2 interacts with Mig1 at Ser³¹¹, a site that is also targeted by Snf1 for phosphorylation (Ahuatzi et al. 2007; Peláez, Herrero and Moreno 2010). By occupying this site, Hxk2 prevents Snf1 phosphorylation and hence removal of Mig1 from the nucleus. Furthermore, subsequent data suggest that Hxk2 is phosphorylated and dephosphorylated at Ser14 by Snf1 and Reg1/Glc7, respectively. Phosphorylation of Hxk2 prevents its nuclear localization and hence its interaction with transcription factors (Fernández-García et al. 2012).

Cat8 is another transcription factor regulated by Snf1 activation and required for gluconeogenesis and survival on alternative carbon sources (Fig. 2). Snf1 controls activity of this transcription factor at two levels. While removal of Mig1 repression by Snf1 allows for upregulation of CAT8 expression, Snf1 phosphorylation of Cat8 triggers its activation (Hedges, Proft and Entian 1995; Randez-Gil et al. 1997). Cat8 derepresses target genes by binding to carbon source-responsive elements (CSREs) in upstream regions of these genes (Young et al. 2003; Roth, Kumme and Schüller 2004). Key gluconeogenic genes as well as genes involved in the glyoxylate cycle and utilization of non-fermentable carbon sources, including FBP1, MLS1 and ICL1, depend on Cat8 for their transcriptional regulation (Randez-Gil et al. 1997; Tachibana et al. 2007; Biddick, Law and Young 2008; Weinhandl et al. 2014). Recently, Znf1 was identified as a transcription factor also involved in regulation of gluconeogenesis and the glyoxylate cycle, but it is not known if Snf1 regulates this transcription factor (Tangsombatvichit et al. 2015).

Another major yet indirect target of Snf1 is the Adr1 transcriptional activator. Under glucose-depleted conditions, Adr1 is important for ethanol utilization and fatty acid metabolism (Fig. 2). Adr1 activity is triggered by Snf1-mediated dephosphorylation and also Snf1 regulates chromatin binding of Adr1 when glucose is scarce (Young, Kacherovsky and Van Riper 2002; Schüller 2003; Ratnakumar et al. 2009; Turcotte et al. 2010). Under repressing conditions though Reg1/Glc7 phosphatases inhibit chromatin binding of Adr1 (Dombek, Kacherovsky and Young 2004). Moreover, Adr1 affects DNA binding of Cat8 (Tachibana et al. 2005; Biddick, Law and Young 2008). Like Cat8, Adr1 also binds to CSREs, and in fact they both target key genes for derepression. For example, Adr1 and Cat8 trigger ADH2 transcription synergistically by binding at the upstream activation sites (Verdone et al. 2002; Tachibana et al. 2005). This binding is required for maximal expression of the ADH2 gene product of which, alcohol dehydrogenase 2, is glucose repressed and required for ethanol catabolism (Walther and Schüller 2001). ACS1, acetyl CoA synthase, is another locus which complete derepression, requires Adr1 and Cat8 working synergistically (Kratzer and Schüller 1997). Adr1 coordinates metabolic activities mainly important for acetyl-CoA and NADH generation from alternative carbon sources, such as lipids (Young et al. 2003). Transcription of FOX2, a multifunctional enzyme involved in fatty acid degradation, also depends on Adr1 (Ratnakumar and Young 2010; Turcotte et al. 2010). Besides coordination of transcriptional factors, Snf1 regulates gene expression by chromatin remodeling. Upon glucose depletion, Snf1 kinase activation of Cat8 and Adr1 plays a role in chromatin remodeling for proper expression of glucoserepressed genes (Agricola et al. 2004; Tachibana et al. 2005). For example, Adr1 restructures the promoter of the ADH2 gene to ensure proper transcriptional activity (Di Mauro 2000; Verdone et al. 2002). Moreover, Snf1 mediates chromatin restructuring also directly via Gcn5 acetyltransferase, as in the case of ADY2 glucose-repressed gene (Abate et al. 2012). Snf1 directly interacts with and phosphorylates the histone acetyltranferase Gcn5 and triggers its histone acetyl transferase activity (Liu, Xu and Kuo 2010). Gcn5-mediated acetylation is critical for transcriptional activation of many stress-responsive genes. For example, binding of Adr1 activation sites depends on the acetylation state of nucleosomes (Verdone et al. 2002). Another way Snf1 exerts its effects on transcriptional activity is by directly interacting with RNAII-pol holoenzyme. In response to glucose limitation, Snf1 kinase physically interacts and phosphorylates the holoenzyme and hereby triggers the transcription process (Kuchin, Treich and Carlson 2000; Young et al. 2012).

Post-transcriptional and -translational effects on metabolism

Dynamic interactions of Snf1 at post-transcriptional and posttranslational levels are important for Snf1's role in balancing cellular energy levels when conditions are not favored. Snf1 achieves energy recalibration by inactivating energetically expensive cellular processes, such as amino acid and lipid biosynthesis, and meanwhile by upregulating programs, such as fatty acid oxidation, that generate energy. It is important to ensure coordinated regulation of carbon and nitrogen metabolism (Rødkær and Færgeman 2014), and under glucose starvation conditions Snf1 downregulates amino acid biosynthesis by inhibiting transcription and translation of GCN4 (Fig. 3). Deletion of SNF1 or inactivation of its kinase activity interestingly reveals significant induction in expression of genes mainly controlled by Gcn4 and required for amino acid generation (Shirra et al. 2008; Zaman et al. 2009). On the other hand, however, Snf1 has also been implicated in promoting translation initiation and activation of Gcn4 by two different mechanisms depending on



Figure 3. The Snf1 protein kinase regulates glucose repression at posttranscriptional and post-translational level. Prominent components of metabolic activity are direct targets of Snf1 kinase activity.

the limiting conditions. When amino acid levels are limited Snf1 interacts with and activates Gcn2, which leads to increased phosphorylation of eIF2 α for promoting Gcn4 translation initiation. Gcn2 and eukaryotic initiation factor 2α (eIF2 α) are two regulators of Gcn4 translation initiation. Phosphorylation of eIF2 α is critical for Gcn4 translation initiation. When Snf1 is active, it inhibits Sit4 and Glc7 phosphatases and hereby promotes the phosphorylation status of eIF2 α resulting in Gcn4 translation initiation (Cherkasova, Qiu and Hinnebusch 2010). The differential effects Snf1 has on GCN4 transcription and translation initiation under both carbon and nitrogen limitations emphasizes the interplay that Snf1 orchestrates between amino acid synthesis, carbon metabolism and cellular energy balance.

Snf1 coordinates carbon availability and cellular energy also through its effects on lipid metabolism. Fatty acid synthesis is energy demanding that has to be minimized during carbon source limitation (Klug and Daum 2014). Snf1 directly contributes to inhibition of this process by phosphorylating and inactivating acetyl carboxylase (Acc1) during energy deficiency (Fig. 3) (Woods et al. 1994; Shirra et al. 2001). This enzyme catalyzes the first rate-limiting step for de novo fatty acid biosynthesis, which is the formation of malonyl-Co A by carboxylation of acetyl CoA and removal of phosphorylation sites has been shown to result in an increased flux towards malonyl-CoA (Shi et al. 2014). In addition to inhibition of fatty acid synthesis, Snf1 promotes energy generation by stimulating degradation of fatty acids by peroxisome proliferation and β -oxidation. Fox2 is an additional factor associated with Snf1 in lipid degradation for energy synthesis (Usaite et al. 2009). This protein is a homodimeric multifunctional enzyme with 3-hydroxyacyl-CoA dehydrogenase and enoyl-CoA hydratese activities that allows Fox2 to have a wide substrate diversity for peroxisomal β -oxidation in yeast (Hiltunen et al. 2003). In silico studies have also identified Fox2 as a Snf1 phosphorylation target in lipid degradation (Ptacek et al. 2005; Usaite et al. 2009). Besides minimizing energy consumption and upregulating energy supply pathways, under prolonged glucose limitation Snf1 phoshorylates the general stress-responsive transcription factor Msn2 and hereby contributes to an adaptive response to poor glucose levels (Fig. 2) (De Wever et al. 2005).

Genetic and biochemical evidence suggest that Snf1 also plays a role in post-transcriptional control of gene regulation. mRNA decay is an important regulatory mechanism that together with nutrient-responsive transcriptional programs allows for a swift adaptation to unpredictable changes in environmental conditions. Inhibition of Snf1 activity leads to a rapid degradation of Snf1-dependent transcripts, whereas constitutive activation of Snf1 increases the stability of the same transcripts even in the presence of glucose (Young et al. 2012). Ccr4, Dhh1 and Xrn1 are three Snf1-targeted proteins involved in glucose-induced mRNA decay. Deletion of any of these proteins causes increased stability of Snf1-dependent transcripts in glucose-rich media (Young et al. 2012). Xrn1 has been suggested to be particularly important since an xrn1 mutant is not viable under conditions in which Snf1 is highly active (Haimovich et al. 2013; Braun and Young 2014). Moreover, an xrn1 reg1 doubledeletion mutant shows synthetic lethality implying how vital Xrn1 is under poor nutrient conditions when Snf1 is active. Xrn1 is a conserved 5' -> 3' exoribonuclease that is critical for transcription-coupled mRNA decay, a process that balances transcription rate and mRNA degradation rate. Xrn1 modulates synthesis and stability of transcripts for genes required at glucose-poor conditions in a Snf1-dependent manner. Snf1 phosphorylates Xrn1 on sites located at the C-terminus and phosphorylated Xrn1 increases the stability of mRNA synthesized from glucose-repressed genes (Fig. 3). Furthermore, Snf1 phosphorylation of Xrn1 is necessary for glucose-induced decay of the same transcripts when glucose is replenished (Braun and Young 2014). The direct effects of Snf1 on mRNA metabolism in response to carbon source shift highlight its regulatory role in modulation of glucose repression.

CONCLUSIONS

Glucose repression involves regulation of a multitude of genes and proteins involved in carbon source utilization and energy generation, and Snf3/Rgt2 sensors and the Snf1 kinase are important for this mechanism. Since yeast cells adjust molecular and cellular activities in response to levels of available glucose, integrating the glucose signal to carbon metabolism is highly regulated. Optimal uptake of glucose starts with the induction of the Snf3/Rgt2 pathway that upregulates expression of hexose transporters best suited for the concentration of glucose in the environment. Snf1 plays a pivotal role in orchestrating the effects of glucose on carbon metabolism. Inactive Snf1, in presence of glucose, mediates transcriptional repression of a multitude of genes including gluconeogenic genes, genes involved in respiration and utilization of non-fermentable carbon sources. Conversely, once activated upon glucose limitation Snf1 phosphorylates a number of proteins to mediate glucose derepression and activation of mechanisms required for energy generation from alternative carbon sources.

Although studies done so far indicate highly interactive and intricate roles Snf1 plays in regulating glucose repression and its effect on carbon metabolism, still much remains to be elucidated; for example, it is not known how Snf1 is inactivated in response to presence of excess glucose, and how Snf1 kinase interacts with other (known and unknown) regulators to control glucose repression.

Conflict of interest. None declared.

REFERENCES

Abate G, Bastonini E, Braun KA, et al. Snf1/AMPK regulates Gcn5 occupancy, H3 acetylation and chromatin remodelling at S. cerevisiae ADY2 promoter. BBA—Gene Regul Mech 2012;1819:419–27.

- Agricola E, Verdone L, Xella B, et al. Common chromatin Architecture, common chromatin remodeling, and common transcription kinetics of Adr1-dependent genes in *Saccharomyces cerevisiae*. Biochemistry 2004;**43**:8878–84.
- Ahuatzi D, Riera A, Peláez R, et al. Hxk2 regulates the phosphorylation state of Mig1 and therefore its nucleocytoplasmic distribution. J Biol Chem 2007;**282**:4485–93.
- Amodeo GA, Rudolph MJ, Tong L. Crystal structure of the heterotrimer core of Saccharomyces cerevisiae AMPK homologue SNF1. Nature 2007;449:492–5.
- Bendrioua L, Smedh M, Almquist J, et al. Yeast AMP-activated protein kinase monitors glucose concentration changes and absolute glucose levels. J Biol Chem 2014;**289**:12863–75.
- Biddick RK, Law GL, Young ET. Adr1 and Cat8 mediate coactivator recruitment and chromatin remodeling at glucoseregulated genes. PLoS One 2008;3:e1436.
- Braun KA, Young ET. Coupling mRNA synthesis and decay. Mol Cell Biol 2014;**34**:4078–87.
- Carlson M, Osmond BC, Botstein D. Mutants of yeast defective in sucrose utilization. *Genetics* 1981;**98**:25–40.
- Castermans D, Somers I, Kriel J, et al. Glucose-induced posttranslational activation of protein phosphatases PP2A and PP1 in yeast. Cell Res 2012;22:1058–77.
- Celenza JL, Carlson M. Mutational analysis of the Saccharomyces cerevisiae SNF1 protein kinase and evidence for functional interaction with the SNF4 protein. Mol Cell Biol 1989;9:5034–44.
- Chen Y, Nielsen J. Advances in metabolic pathway and strain engineering paving the way for sustainable production of chemical building blocks. Curr Opin Biotechnol 2013;24:965–72.
- Cherkasova V, Qiu H, Hinnebusch AG. Snf1 promotes phosphorylation of the subunit of eukaryotic translation initiation factor 2 by activating Gcn2 and inhibiting phosphatases Glc7 and Sit4. Mol Cell Biol 2010;**30**:2862–73.
- Dai Z, Liu Y, Guo J, et al. Yeast synthetic biology for high-value metabolites. FEMS Yeast Res 2015;15:1–11.
- De Wever V, Reiter W, Ballarini A, *et al*. A dual role for PP1 in shaping the Msn2-dependent transcriptional response to glucose starvation. *EMBO J* 2005;**24**:4115–23.
- Di Mauro E. Two distinct nucleosome alterations characterize chromatin remodeling at the Saccharomyces cerevisiae ADH2 promoter. J Biol Chem 2000;**275**:7612–8.
- Dombek KM, Kacherovsky N, Young ET. The Reg1-interacting proteins, Bmh1, Bmh2, Ssb1, and Ssb2, have roles in maintaining glucose repression in Saccharomyces cerevisiae. J Biol Chem 2004;**279**:39165–74.
- Feng ZH, Wilson SE, Peng ZY, et al. The yeast GLC7 gene required for glycogen accumulation encodes a type 1 protein phosphatase. J Biol Chem 1991;**266**:23796–801.
- Fernández-García P, Peláez R, Herrero P, et al. Phosphorylation of yeast hexokinase 2 regulates its nucleocytoplasmic shuttling. J Biol Chem 2012;**287**:42151–64.
- Flick KM, Spielewoy N, Kalashnikova TI, et al. Grr1-dependent inactivation of Mth1 mediates glucose-induced dissociation of Rgt1 from HXT gene promoters. Mol Biol Cell 2003;14: 3230–41.
- Gancedo JM. Yeast carbon catabolite repression. Microbiol Mol Biol R 1998;62:334–61.
- Griffioen G, Anghileri P, Imre E, et al. Nutritional control of nucleocytoplasmic localization of cAMP-dependent protein kinase catalytic and regulatory subunits in *Saccharomyces cerevisiae*. *J Biol Chem* 2000;**275**:1449–56.
- Haimovich G, Medina DA, Causse SZ, et al. Gene expression is circular: factors for mRNA degradation also foster mRNA synthesis. Cell 2013;**153**:1000–11.

- Hedbacker K, Carlson M. SNF1/AMPK pathways in yeast. Front Biosci 2008;13:2408–20.
- Hedbacker K, Carlson M. SNF1/AMPK pathway in yeast. Front Biosci 2010;1–3.
- Hedbacker K, Hong S-P, Carlson M. Pak1 protein kinase regulates activation and nuclear localization of Snf1-Gal83 protein kinase. Mol Cell Biol 2004;**24**:8255–63.
- Hedges D, Proft M, Entian KD. CAT8, a new zinc clusterencoding gene necessary for derepression of gluconeogenic enzymes in the yeast Saccharomyces cerevisiae. Mol Cell Biol 1995;15:1915–22.
- Hiltunen JK, Mursula AM, Rottensteiner H, et al. The biochemistry of peroxisomal beta-oxidation in the yeast Saccharomyces cerevisiae. FEMS Microbiol Rev 2003;27:35–64.
- Hong S-P, Leiper FC, Woods A, et al. Activation of yeast Snf1 and mammalian AMP-activated protein kinase by upstream kinases. P Natl Acad Sci USA 2003;**100**:8839–43.
- Jiang R, Carlson M. Glucose regulates protein interactions within the yeast SNF1 protein kinase complex. Gene Dev 1996;10:3105–15.
- Kaniak A, Xue Z, Macool D, et al. Regulatory network Connecting two glucose signal transduction pathways in Saccharomyces cerevisiae. Eukaryot Cell 2004;3:221–31.
- Karhumaa K, Wu B, Kielland-Brandt MC. Conditions with high intracellular glucose inhibit sensing through glucose sensor Snf3 in Saccharomyces cerevisiae. J Cell Biochem 2010;110:920–5.
- Kim J-H, Johnston M. Two glucose-sensing pathways converge on Rgt1 to regulate expression of glucose transporter genes in Saccharomyces cerevisiae. J Biol Chem 2006;281:26144–9.
- Klein CJL, Olsson L, Nielsen J. Glucose control in Saccharomyces cerevisiae: the role of MIG1 in metabolic functions. Microbiology 1998;144:13–24.
- Klug L, Daum G. Yeast lipid metabolism at a glance. FEMS Yeast Res 2014;**14**:369–88.
- Kratzer S, Schüller HJ. Transcriptional control of the yeast acetyl-CoA synthetase gene, ACS1, by the positive regulators CAT8 and ADR1 and the pleiotropic repressor UME6. Mol Microbiol 1997;26:631–41.
- Kuchin S, Treich I, Carlson M. A regulatory shortcut between the Snf1 protein kinase and RNA polymerase II holoenzyme. P Natl Acad Sci USA 2000;97:7916–20.
- Leech A, Nath N, McCartney RR, et al. Isolation of mutations in the catalytic domain of the snf1 kinase that render its activity independent of the snf4 subunit. *Eukaryot Cell* 2003;2: 265–73.
- Li M, Borodina I. Application of synthetic biology for production of chemicals in yeast Saccharomyces cerevisiae. FEMS Yeast Res 2015;**15**:1–12.
- Liu Y, Xu X, Kuo M-H. Snf1p regulates Gcn5p transcriptional activity by antagonizing Spt3p. Genetics 2010;184: 91–105.
- Ludin K, Jiang R, Carlson M Glucose-regulated interaction of a regulatory subunit of protein phosphatase 1 with the Snf1 protein kinase in Saccharomyces cerevisiae. P Natl Acad Sci USA 1998;95:6245–50.
- Lutfiyya LL, Iyer VR, DeRisi J, et al. Characterization of three related glucose repressors and genes they regulate in Saccharomyces cerevisiae. Genetics 1998;**150**:1377–91.
- McCartney RR, Rubenstein EM, Schmidt MC. Snf1 kinase complexes with different beta subunits display stress-dependent preferences for the three Snf1-activating kinases. *Curr Genet* 2005;47:335–44.
- McCartney RR, Schmidt MC. Regulation of Snf1 kinase. Activation requires phosphorylation of threonine 210 by an

upstream kinase as well as a distinct step mediated by the Snf4 subunit. J Biol Chem 2001;**276**:36460–6.

- Mayer FV, Heath R, Underwood E, et al. ADP regulates SNF1, the Saccharomyces cerevisiae homolog of AMP-activated protein kinase. Cell Metab 2011;**14**:707–14.
- Moriya H, Johnston M. Glucose sensing and signaling in Saccharomyces cerevisiae through the Rgt2 glucose sensor and casein kinase I. P Natl Acad Sci USA 2004;**101**:1572–7.
- Nath N, McCartney RR, Schmidt MC. Yeast Pak1 kinase associates with and activates Snf1. Mol Cell Biol 2003;23:3909–17.
- Ozcan S. Two different signals regulate repression and induction of gene expression by glucose. J Biol Chem 2002;277:46993–7.
- Özcan S, Dover J, Johnston M. Glucose sensing and signaling by two glucose receptors in the yeast *Saccharomyces cerevisiae*. EMBO J 1998;**17**:2566–73.
- Ozcan S, Johnston M. Function and regulation of yeast hexose transporters. Microbiol Mol Biol R 1999;63:554–69.
- Palomino A, Herrero P, Moreno F. Tpk3 and Snf1 protein kinases regulate Rgt1 association with *Saccharomyces cerevisiae* HXK2 promoter. *Nucleic Acids Res* 2006;**34**:1427–38.
- Pasula S, Chakraborty S, Choi JH, et al. Role of casein kinase 1 in the glucose sensor-mediated signaling pathway in yeast. BMC Cell Biol 2010;11:17.
- Peláez R, Herrero P, Moreno F. Functional domains of yeast hexokinase 2. Biochem J 2010;**432**:181–90.
- Petranovic D, Nielsen J. Can yeast systems biology contribute to the understanding of human disease? Trends Biotechnol 2008;26:584–90.
- Petranovic D, Tyo K, Vemuri GN, et al. Prospects of yeast systems biology for human health: integrating lipid, protein and energy metabolism. FEMS Yeast Res 2010;**10**:1046–59.
- Piskur J, Compagno C (eds). Molecular Mechanisms in Yeast Carbon Metabolism. Heidelberg: Springer, 2014.
- Ptacek J, Devgan G, Michaud G, et al. Global analysis of protein phosphorylation in yeast. *Nature* 2005;**438**:679–84.
- Randez-Gil F, Bojunga N, Proft M, et al. Glucose derepression of gluconeogenic enzymes in Saccharomyces cerevisiae correlates with phosphorylation of the gene activator Cat8p. Mol Cell Biol 1997;17:2502–10.
- Ratnakumar S, Kacherovsky N, Arms E, et al. Snf1 controls the activity of Adr1 through dephosphorylation of Ser230. *Genet*ics 2009;**182**:735–45.
- Ratnakumar S, Young ET. Snf1 dependence of peroxisomal gene expression is mediated by Adr1. J Biol Chem 2010;285: 10703–14.
- Rødkær SV, Færgeman NJ. Glucose- and nitrogen sensing and regulatory mechanisms in Saccharomyces cerevisiae. FEMS Yeast Res 2014;**14**:683–96.
- Rolland F, Winderickx J, Thevelein JM. Glucose-sensing and -signalling mechanisms in yeast. FEMS Yeast Res 2002;2: 183–201.
- Roth S, Kumme J, Schüller H-J. Transcriptional activators Cat8 and Sip4 discriminate between sequence variants of the carbon source-responsive promoter element in the yeast Saccharomyces cerevisiae. Curr Genet 2004;45:121–8.
- Roy A, Jouandot II D, Cho KH, et al. Understanding the mechanism of glucose-induced relief of Rgt1-mediated repression in yeast. Elsevier 2014;4:105–11.
- Roy A, Shin YJ, Cho KH, et al. Mth1 regulates the interaction between the Rgt1 repressor and the Ssn6-Tup1 corepressor complex by modulating PKA-dependent phosphorylation of Rgt1. Mol Biol Cell 2013;24:1493–503.
- Rubenstein EM, McCartney RR, Zhang C, et al. Access denied: Snf1 activation loop phosphorylation is controlled by

availability of the phosphorylated threonine 210 to the PP1 phosphatase. J Biol Chem 2008;**283**:222–30.

- Sanz P, Alms GR, Haystead TA, et al. Regulatory interactions between the Reg1-Glc7 protein phosphatase and the Snf1 protein kinase. Mol Cell Biol 2000;**20**:1321–8.
- Schmidt MC, McCartney RR, Zhang X, et al. Std1 and Mth1 proteins interact with the glucose sensors to control glucoseregulated gene expression in Saccharomyces cerevisiae. Mol Cell Biol 1999;19:4561–71.
- Schüller H-J. Transcriptional control of nonfermentative metabolism in the yeast Saccharomyces cerevisiae. Curr Genet 2003;43:139–60.
- Shi S, Chen Y, Siewers V, et al. Improving production of malonyl coenzyme A-derived metabolites by abolishing Snf1dependent regulation of Acc1. mBio 2014;5:e01130–14.
- Shirra MK, McCartney RR, Zhang C, et al. A chemical genomics study identifies Snf1 as a repressor of GCN4 translation. *J Biol Chem* 2008;**283**:35889–98.
- Shirra MK, Patton-Vogt J, Ulrich A, et al. Inhibition of acetyl coenzyme A carboxylase activity restores expression of the INO1 gene in a snf1 mutant strain of Saccharomyces cerevisiae. Mol Cell Biol 2001;21:5710–22.
- Spielewoy N, Flick K, Kalashnikova TI, et al. Regulation and recognition of SCFGrr1 targets in the glucose and amino acid signaling pathways. Mol Cell Biol 2004;24:8994–9005.
- Sutherland CM, Hawley SA, McCartney RR, et al. Elm1p is one of three upstream kinases for the Saccharomyces cerevisiae SNF1 complex. Curr Biol 2003;13:1299–305.
- Tachibana C, Biddick R, Law GL, *et al*. A poised initiation complex is activated by SNF1. *J Biol Chem* 2007;**282**:37308–15.
- Tachibana C, Yoo JY, Tagne JB, et al. Combined global localization analysis and transcriptome data Identify genes that are directly coregulated by Adr1 and Cat8. Mol Cell Biol 2005;25:2138–46.
- Tangsombatvichit P, Semkiv MV, Sibirny AA, et al. Zinc cluster protein Znf1, a novel transcription factor of nonfermentative metabolism in Saccharomyces cerevisiae. FEMS Yeast Res 2015;15:1–16.
- Treitel MA, Carlson M. Repression by SSN6-TUP1 is directed by MIG1, a repressor/activator protein. P Natl Acad Sci USA 1995;**92**:3132–6.
- Tu J, Carlson M. REG1 binds to protein phosphatase type 1 and regulates glucose repression in Saccharomyces cerevisiae. EMBO J 1995;14:5939–46.
- Turcotte B, Liang XB, Robert F, *et al*. Transcriptional regulation of nonfermentable carbon utilization in budding yeast. FEMS Yeast Res 2010;**10**:2–13.
- Usaite R, Jewett MC, Oliveira AP, *et al*. Reconstruction of the yeast Snf1 kinase regulatory network reveals its role as a global energy regulator. Mol Syst Biol 2009;**5**:1–12.
- Verdone L, Wu J, Van Riper K, et al. Hyperacetylation of chromatin at the ADH2 promoter allows Adr1 to bind in repressed conditions. EMBO J 2002;**21**:1101–11.
- Vincent O, Townley R, Kuchin S, et al. Subcellular localization of the Snf1 kinase is regulated by specific β subunits and a novel glucose signaling mechanism. *Gene Dev* 2001;**15**:1104–14.
- Walther K, Schüller HJ. Adr1 and Cat8 synergistically activate the glucose-regulated alcohol dehydrogenase gene ADH2 of the yeast Saccharomyces cerevisiae. *Microbiol Read Engl* 2001;**147**:2037–44.
- Weinhandl K, Winkler M, Glieder A, et al. Carbon source dependent promoters in yeasts. Microb Cell Fact 2014;**13**:5.
- Wieczorke R, Krampe S, Weierstall T, et al. Concurrent knockout of at least 20 transporter genes is required to block

uptake of hexoses in Saccharomyces cerevisiae. FEBS Lett 1999;464:123–8.

- Woods A, Munday MR, Scott J, et al. Yeast SNF1 is functionally related to mammalian AMP-activated protein kinase and regulates acetyl-CoA carboxylase in vivo. J Biol Chem 1994;269:19509–15.
- Young ET, Dombek KM, Tachibana C, et al. Multiple pathways are co-regulated by the protein kinase Snf1 and the transcription factors Adr1 and Cat8. J Biol Chem 2003;**278**:26146–58.
- Young ET, Kacherovsky N, Van Riper K. Snf1 protein kinase regulates Adr1 binding to chromatin but not transcription activation. J Biol Chem 2002;**277**:38095–103.
- Young ET, Zhang C, Shokat KM, et al. The AMP-activated protein kinase Snf1 regulates transcription factor binding, RNA poly-

merase II activity, and mRNA stability of glucose-repressed genes in Saccharomyces cerevisiae. J Biol Chem 2012;**287**: 29021–34.

- Zaman S, Lippman SI, Schneper L, et al. Glucose regulates transcription in yeast through a network of signaling pathways. Mol Syst Biol 2009;**5**:245.
- Zhang J, Olsson L, Nielsen J. The β -subunits of the Snf1 kinase in Saccharomyces cerevisiae, Gal83 and Sip2, but not Sip1, are redundant in glucose derepression and regulation of sterol biosynthesis. Mol Microbiol 2010;77: 371–83.
- Zhang J, Vaga S, Chumnanpuen P, et al. Mapping the interaction of Snf1 with TORC1 in Saccharomyces cerevisiae. Mol Syst Biol 2011;7:545.