



## **Corrigendum to: "Plasma concentration of the neurofilament light protein (NFL) is a biomarker of CNS injury in HIV infection: A cross-sectional**

Downloaded from: <https://research.chalmers.se>, 2025-12-04 23:21 UTC

Citation for the original published paper (version of record):

Gisslén, M., Price, R., Andreasson, U. et al (2016). Corrigendum to: "Plasma concentration of the neurofilament light protein (NFL) is a biomarker of CNS injury in HIV infection: A cross-sectional study" [EBioMedicine 3 (216), 2016, 135-140], DOI: 10.1016/j.ebiom.2015.11.036. EBioMedicine, 7: 287-288.  
<http://dx.doi.org/10.1016/j.ebiom.2016.04.021>

N.B. When citing this work, cite the original published paper.



## Corrigendum

Corrigendum to: “Plasma concentration of the neurofilament light protein (NFL) is a biomarker of CNS injury in HIV infection: A cross-sectional study”  
[EBioMedicine 3 (216) 135–140]



Magnus Gisslén, Richard W. Price, Ulf Andreasson, Niklas Norgren, Staffan Nilsson, Lars Hagberg, Dietmar Fuchs, Serena Spudich, Kaj Blennow, Henrik Zetterberg

The authors wish to point out that the units of CSF and plasma NFL in Table 2 and Figs. 1 and 2 should be corrected to ng/L. Corrected files are republished here.

**Table 2**


Median (IQR) plasma NFL (left columns) and CSF NFL (right columns) concentrations in different groups.

Groups	N	Plasma NFL	CSF NFL
		Median ng/L (IQR)	Median ng/L (IQR)
HIV negative (HIV-neg)	19	9.3 (5.9–13.1)	363 (264–487)
Primary HIV infection (PHI)	13	11.2 (8.2–16.6)	732 (456–1288)
Neuroasymptomatic HIV (NA)			
CD4 > 350	19	9.0 (6.5–14.3)	488 (374–694)
CD4 200–349	17	9.7 (7.6–12.2)	573 (464–764)
CD4 50–199	19	12.1 (7.8–44.7)	640 (490–1530)
CD4 < 50	20	39.6 (12.0–103)	2415 (690–4420)
HIV-associated dementia (HAD)	11	114 (46.0–235)	16,185 (1513–43,010)
HIV, treated-suppressed (ART)	22	11.1 (8.6–12.8)	582 (322–706)


DOI of original article: <http://dx.doi.org/10.1016/j.ebiom.2015.11.036>.

<http://dx.doi.org/10.1016/j.ebiom.2016.04.021>

2352–3964/© 2016 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).



**Fig. 1.** Association between CSF and plasma NFL and concentrations of biomarkers in 8 subject groups. Panels A–B and D–F, plot concentrations of different markers for the 8 subject groups, HIV-negative controls (HIV-neg); primary HIV infection (PHI); untreated neuroasymptomatic subjects in different CD4 cell strata (Asympt); HIV-associated dementia (HAD); and subjects on suppressive antiretroviral treatment (ART). Boxes depict median and IQR, whiskers show 10–90 percentiles and ‘+’ designate the means. Dotted horizontal lines in panels D–E show the limit of detection for HIV RNA (40 copies/mL), and in panel F the upper limit of normal for CSF neopterin (5.8 nmol/L). Panel C plot the correlation between log CSF and log plasma NFL.



**Fig. 2.** Association between NFL and age. Panel A displays the regression lines for age to log plasma and log CSF NFL in HIV-negative controls. Age was closely correlated with plasma NFL ( $r = 0.792, p < 0.0001$ ) and CSF NFL ( $r = 0.673, p = 0.002$ ) in HIV-negative controls. Panel B shows plasma NFL and panel C CSF NFL plotted against age in four HIV-infected groups: HIV-associated dementia (HAD); untreated neuroasymptomatic subjects (NA); subjects on suppressive antiretroviral treatment (ART); primary HIV-infection (PHI), and HIV-negative controls (HIV-neg).