



**CHALMERS**  
UNIVERSITY OF TECHNOLOGY

## **Metabolic engineering of *Saccharomyces cerevisiae* for overproduction of triacylglycerols**

Downloaded from: <https://research.chalmers.se>, 2019-09-22 10:47 UTC

Citation for the original published paper (version of record):


Ferreira, R., Teixeira, P., Gossing, M. et al (2018)

Metabolic engineering of *Saccharomyces cerevisiae* for overproduction of triacylglycerols

Metabolic Engineering Communications, 6: 22-27

<http://dx.doi.org/10.1016/j.meteno.2018.01.002>

N.B. When citing this work, cite the original published paper.



## Metabolic engineering of *Saccharomyces cerevisiae* for overproduction of triacylglycerols

Raphael Ferreira<sup>a,b</sup>, Paulo Gonçalves Teixeira<sup>a,b</sup>, Michael Gossing<sup>a,b</sup>, Florian David<sup>a,b</sup>, Verena Siewers<sup>a,b</sup>, Jens Nielsen<sup>a,b,c,\*</sup>

<sup>a</sup> Department of Biology and Biological Engineering, Chalmers University of Technology, SE412 96 Gothenburg, Sweden

<sup>b</sup> Novo Nordisk Foundation Center for Biosustainability, Chalmers University of Technology, SE412 96 Gothenburg, Sweden

<sup>c</sup> Novo Nordisk Foundation Center for Biosustainability, Technical University of Denmark, DK2800 Kgs Lyngby, Denmark



### A B S T R A C T

Triacylglycerols (TAGs) are valuable versatile compounds that can be used as metabolites for nutrition and health, as well as feedstocks for biofuel production. Although *Saccharomyces cerevisiae* is the favored microbial cell factory for industrial production of biochemicals, it does not produce large amounts of lipids and TAGs comprise only ~1% of its cell dry weight. Here, we engineered *S. cerevisiae* to reorient its metabolism for overproduction of TAGs, by regulating lipid droplet associated-proteins involved in TAG synthesis and hydrolysis. We implemented a push-and-pull strategy by overexpressing genes encoding a deregulated acetyl-CoA carboxylase, *ACC1*<sup>S659A/S1157A</sup> (*ACC1\*\**), as well as the last two steps of TAG formation: phosphatidic phosphatase (*PAH1*) and diacylglycerol acyltransferase (*DGA1*), ultimately leading to 129 mg·gCDW<sup>-1</sup> of TAGs. Disruption of TAG lipase genes *TGL3*, *TGL4*, *TGL5* and sterol acyltransferase gene *ARE1* increased the TAG content to 218 mg·gCDW<sup>-1</sup>. Further disruption of the beta-oxidation by deletion of *POX1*, as well as glycerol-3-phosphate utilization through deletion of *GUT2*, did not affect TAGs levels. Finally, disruption of the peroxisomal fatty acyl-CoA transporter *PXA1* led to accumulation of 254 mg·gCDW<sup>-1</sup>. The TAG levels achieved here are the highest titer reported in *S. cerevisiae*, reaching 27.4% of the maximum theoretical yield in minimal medium with 2% glucose. This work shows the potential of using an industrially established and robust yeast species for high level lipid production.

### 1. Introduction

In recent years, there has been an increasing demand for oils with specialized properties. Most oils found in nature are mainly composed of triacylglycerols (TAGs). Natural producers with specific TAG compositions, such as exotic plants, are usually poorly understood at the metabolic level, and lack available tools for introducing genetic modification, which ultimately makes them difficult to engineer. Consequently, there is an important need for alternative organisms that can accumulate high lipid amounts and be easily engineered for tailored oil production.

Although certain microorganisms are known to accumulate high levels of lipids, *Saccharomyces cerevisiae* remains the preferred cell factory for production of biochemicals (Nielsen and Keasling, 2016; Krivoruchko and Nielsen, 2015). It is a Generally Regarded As Safe (GRAS) organism with high tolerance to low pH and fermentation inhibitors. One of the major advantages of *S. cerevisiae* is the vast

knowledge available about its physiology and metabolism, providing a solid background for engineering approaches (Julleson et al., 2015). It is the most tractable organism, with several available tools for genome engineering, such as the CRISPR-Cas system, which has proven to be highly efficient (Dicarlo et al., 2013; Jakočiūnas et al., 2015; Zalatan et al., 2015).

TAGs are composed of three fatty acid chains esterified with a glycerol backbone. As such, the first key step towards increasing the TAG production potential in *S. cerevisiae* is the upregulation of the fatty acid biosynthesis. Fatty acids are synthesized as acyl-CoAs by the fatty acid synthetase complex, encoded by the genes *FAS1* and *FAS2* (Schweizer et al., 1984). Malonyl-CoA is used in this process as a substrate for iterative elongation of acyl chains, and is synthesized from acetyl-CoA by the enzyme acetyl-CoA carboxylase (Acc1). This enzyme is tightly regulated at the transcriptional and protein-level, rendering it a major regulation step in fatty acid biosynthesis (Hasslacher et al., 1993; Blank et al., 2017; Bozaquel-Morais et al., 2017). It is inhibited by

\* Corresponding author at: Department of Biology and Biological Engineering, Chalmers University of Technology, SE412 96 Gothenburg, Sweden.  
E-mail address: [nielsenj@chalmers.se](mailto:nielsenj@chalmers.se) (J. Nielsen).

Snf1 phosphorylation, and the expression of a double mutant *Acc1*<sup>S659A/S1157A</sup> with two removed phosphorylation sites, here shortened *Acc1*\*\* has been shown to increase production of malonyl-CoA and fatty acid-derived products up to 3-fold compared to the wild-type (wt) strain (Shi et al., 2014).

The second point of focus is increasing the flux of acyl chains towards TAG formation. TAG assembly is initiated with glycerol-3-phosphate (Gro-3-P), which is consecutively acylated with two acyl-chains, by GPAT and LPAT genes, to form phosphatidic acid (PA). PA is then dephosphorylated by the phosphatidic acid phosphohydrolase Pah1 to diacylglycerol (DAG), which is then used as a substrate by the diacylglycerol acyltransferase Dga1 or the phospholipid: diacylglycerol acyltransferase Lro1 to form TAG. Previous studies in *Yarrowia lipolytica* have shown that increasing levels of Gro-3-P by removing its utilization through the glycerol-3-phosphate dehydrogenase (Gut2) improved the TAG content 3-fold compared to the wt (Beopoulos et al., 2008). In *S. cerevisiae*, individual overexpression of either *PAH1* (Hung et al., 2016) or *DGA1* (Jensen et al., 2017, Runguphan and Keasling, 2014) resulted in a significant increase in TAG levels.

TAGs are stored in form of lipid droplets (LDs) together with sterol esters (SEs) (Kohlwein et al., 2013). When required by the cell, the acyl chains in these storage lipids can be cleaved by TAG lipases and SE hydrolases. Tgl3, Tgl4, and Tgl5 are members of the conserved patatin-domain-containing family of hydrolases (Athenstaedt and Daum, 2005) and simultaneous deletion of the respective genes has been proven to lead to an increase of TAG levels (Ploier et al., 2013). These studies indicate that TAG degradation mechanisms might counteract accumulation of high levels of these compounds, highlighting the importance of removing lipase activity and further fatty acid degradation such as fatty acid beta-oxidation.

The above-mentioned individual engineering strategies have been successful at increasing the TAG levels in *S. cerevisiae* and *Y. lipolytica*. However, there has not been any extensive study on exploring a combinatorial approach of all these strategies in a single strain. In this work, we developed a *S. cerevisiae* platform strain with high-level production and accumulation of TAGs. We therefore demonstrate the potential for *S. cerevisiae* to accumulate up to 25% of dry cell mass as TAGs without resorting to heterologous proteins.

## 2. Methods

### 2.1. Plasmid and strain construction

Strains, plasmids, links to plasmid maps, gRNA sequences and primers used in this study are listed in Table 1 and Additional File 1. Strain IMX581, and plasmids pROS10, pROS10, pMEL13 and pMEL10 were obtained from EUROSCARF (Frankfurt, Germany). Diagnostic primers and repair fragments were designed using the Yeastriction webtool (<http://yeastriction.tnw.tudelft.nl/>). Gene deletions were performed and verified as described in (Mans et al., 2015). DNA fragments for *DGA1* and *PAH1* were amplified from genomic DNA of CEN.PK 113–7D (provided by P. Kötter, University of Frankfurt, Germany). Sequences for *PGK1p* and *TEF1p* were amplified from pSP-G1 (Partow et al., 2010). Integrative plasmids were based on pCfB389 (XI-2-*loxP-KIURA3*) and

pCfB2055 (X-2-*loxP-KanMX4*), a derivative of pCfB353 (Jensen et al., 2014). The *KanMX4* cassette from pCfB2055 was subcloned with *XhoI-SalI* into pCfB389 to generate pMG95 (XI-2-*loxP-KanMX4*). The sequences for *PGK1p*, *TEF1p*, *PAH1* and *DGA1* were assembled into an integrative vector using pMG95 as backbone, generating pMG97. Sequences were assembled into integrative plasmids using CPEC (Quan and Tian, 2011). Construction of *HXT7p-ACC1*\*\* has been described elsewhere (Bergenholtz et al., submitted). For integration, generated integrative vectors were linearized with *NotI* and transformed directly using a PEG/LiOAc technique (Gietz et al., 2007). Correct integration was confirmed using primers located up- and downstream of the integration sites. For construction of *HXT7p-ACC1*\*\* *PAH1* + *DGA1*, the *KanMX4* marker was consecutively removed by expressing the *cre* recombinase gene from pSH65 as described (Jensen et al., 2014).

### 2.2. Growth medium

*S. cerevisiae* strains with auxotrophies were grown on YPD plates containing 20 g·L<sup>-1</sup> glucose, 10 g·L<sup>-1</sup> yeast extract, 20 g·L<sup>-1</sup> peptone from casein and 20 g·L<sup>-1</sup> agar. *URA3* plasmid carrying strains were grown on selective growth medium containing 6.9 g·L<sup>-1</sup> yeast nitrogen base without amino acids (Formedium, Hunstanton, UK), 0.77 g·L<sup>-1</sup> complete supplement mixture without uracil (Formedium), 20 g·L<sup>-1</sup> glucose and 20 g·L<sup>-1</sup> agar. *KanMX4* plasmid carrying strains were grown on YPD plates supplemented with 200 mg·L<sup>-1</sup> Geneticin®. Shake flask cultivations were performed in minimal medium containing 20 g·L<sup>-1</sup> glucose, 5 g·L<sup>-1</sup> (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 14.4 g·L<sup>-1</sup> KH<sub>2</sub>PO<sub>4</sub>, 0.5 g·L<sup>-1</sup> MgSO<sub>4</sub>·7H<sub>2</sub>O adjusted to pH 6. After sterilization, 2 mL·L<sup>-1</sup> trace element solution and 1 mL·L<sup>-1</sup> of vitamin solution were added. The composition of the trace element and vitamin solution has been reported earlier (Verduyn et al., 1992).

### 2.3. Shake flask cultivations

Biological triplicates were pre-cultivated in 5 mL minimal medium at 200 rpm and 30 °C for 18 h. Subsequently, the pre-culture was diluted into 15 mL minimal medium in a 100 mL shake flask to an OD<sub>600</sub> of 0.1. Shake flasks were incubated at 200 rpm and 30 °C for 72 h. A spectrophotometer (Genesis 20, Thermo Fisher Scientific, Waltham, MA, USA) was used to measure cell density at the end of the shake flask cultivations.

### 2.4. Lipid quantification

Samples for lipid analysis were taken at the end of the shake flask cultivations. The samples were centrifuged at 3000g for 5 min and the supernatant was discarded. Cell pellets were then washed with 50 mL deionized water, centrifuged at 3000g for 5 min and the supernatant was discarded. The pellets were kept at -20 °C for 10 min and then freeze-dried using a Christ alpha 2–4 LSC (Christ Gefrier Trocknungsanlagen, Osterode, Germany). Lipids were extracted using microwave extraction and analyzed by HPLC-CAD using 10 mg of dried cell biomass as previously described (Khoormung et al., 2013).

**Table 1**  
Strains used for this study.

Strain	Genotype	Parental strain	Reference
IMX581	<i>MATa ura3–52 can1::cas9-natNT2</i>	CEN.PK113–5D	Mans et al. (2015)
ADP	<i>MATa ura3–52 can1::cas9-natNT2 HXT7p-ACC1** PGK1p-PAH1 TEF1p-DGA1</i>	IMX581	This study
RF08	<i>MATa ura3–52 can1::cas9-natNT2 HXT7p-ACC1** PGK1p-PAH1 TEF1p-DGA1 tgl3/4/5Δ are1Δ</i>	ADP	This study
RF09	<i>MATa ura3–52 can1::cas9-natNT2 HXT7p-ACC1** PGK1p-PAH1 TEF1p-DGA1 tgl3/4/5Δ are1Δ pox1Δ</i>	RF08	This study
RF10	<i>MATa ura3–52 can1::cas9-natNT2 HXT7p-ACC1** PGK1p-PAH1 TEF1p-DGA1 tgl3/4/5Δ are1Δ pox1Δ gut2Δ</i>	RF09	This study
RF11	<i>MATa ura3–52 can1::cas9-natNT2 HXT7p-ACC1** PGK1p-PAH1 TEF1p-DGA1 tgl3/4/5Δ are1Δ pox1Δ gut2Δ pxa1Δ</i>	RF10	This study
RF12	<i>MATa ura3–52 can1::cas9-natNT2 HXT7p-ACC1** PGK1p-PAH1 TEF1p-DGA1 tgl3/4/5Δ are1Δ pox1Δ gut2Δ pxa1Δ faa2Δ</i>	RF11	This study

## 2.5. Lipid droplet staining

After 72 h of shake flask cultivation, 100  $\mu$ L of cell culture were transferred into a 1.5 mL Eppendorf tube, centrifuged and the pellet washed with 1 mL of deionized water. Cells were then centrifuged at 3000g for 5 min and resuspended with 100  $\mu$ L phosphate-buffered saline (PBS). Resuspended cells were treated with 1  $\mu$ L of BODIPY 493/503 (Thermo Fisher Scientific, Waltham, MA, USA) solution (1 mg·mL<sup>-1</sup> in ethanol) and kept at 4 °C in the dark for 10 min. Fluorescent microscope pictures were analyzed using a Leica AF 6000 inverted microscope using GFP filter settings (Leica Microsystems, Wetzlar, Germany) and processed with the Leica Application Suite (LAS) software.


## 3. Results

### 3.1. Push-and-pull strategy: Overexpression of ACC1\*\*, PAH1 and DGA1 leads to high accumulation of TAGs


We first co-overexpressed genes responsible for the first and last steps of TAG biogenesis, termed push-and-pull strategy. In that regard, we genomically integrated in chromosome X one copy of the double mutant version of *ACC1*, *ACC1*<sup>S659A/S1157A</sup> (*ACC1*\*\*), under control of the strong promoter *HXT7p* induced at low glucose concentrations (Jensen et al., 2014; Reifenberger et al., 1997), as the majority of fatty acids are produced during the ethanol phase (Teixeira et al., 2017, Bergenholm et al., submitted) (Fig. 1). We further engineered the strain by genomically integrating in chromosome XI one additional copy of *PAH1* and *DGA1* each, expressed under control of the strong constitutive promoters *PGK1p* and *TEF1p*, respectively, resulting in the ADP strain (Table 1) (Jensen et al., 2014). This combination of overexpressed genes led to 129 mg·gCDW<sup>-1</sup> of TAGs after 72 h shake flask cultures in 2% glucose minimal medium, a higher than 10-fold increase in comparison to the reference strain (Fig. 2). The accumulation of TAGs also led to an increase of the number of lipid droplets (Fig. S1). Notably, the integration of these genes did not alter the final OD of the resulting strain compared to the reference strain (Fig. S2).

### 3.2. Removal of the major TAG lipases and sterol acyltransferase increases TAG levels

In *S. cerevisiae*, three TAG lipases, namely Tgl3, Tgl4, and Tgl5, are responsible for most of the TAG lipase activity, and their deletion has been shown to lead to an increased level of TAGs (Athenstaedt and Daum, 2003).



**Fig. 1. Simplified schematic representation of the key metabolic fluxes targeted in this study.** Highlighted in green are genes overexpressed, *ACC1*\*\* (acetyl-CoA carboxylase), *DGA1* (diacylglycerol acyltransferase) and *PAH1* (phosphatidic acid phosphohydrolase). Highlighted in red are deleted genes, *GUT2* (glycerol-3-phosphate dehydrogenase), *ARE1* (acyl-CoA: sterol acyltransferase), *TGL3/4/5* (triacylglycerol lipases), *PXA1* (subunit of peroxisomal ABC-transporter), *POX1* (fatty acyl-CoA oxidase), and *FAA2* (fatty acyl-CoA synthetase). Marked in black are unmodified genes that are also involved in lipid metabolism: *FAS1/2* (fatty acid synthetase); *GPT2* and *SCT1* (GPAT genes); *LOA1*, *SLC1* and *ALE1* (LPAT genes); and *ARE2* (acyl-CoA: sterol acyltransferase). For the metabolic compounds: G3P: glyceraldehyde-3-phosphate; DHAP: dihydroxyacetone phosphate, Gro-3-P: glycerol-3-phosphate; LPA: Lysophosphatidic acid; PA: Phosphatidic acid; DAG: diacylglycerol; SE: sterol ester; TAG: triacylglycerol; FFA: free fatty acid. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)



**Fig. 2. TAG quantification of the different engineered strains.** TAGs were quantified for strains with the implemented push-and-pull approach (*ACC1*\*\*, *DGA1* and *PAH1* overexpression) and lacking TAG lipases (*TGL3/4/5* deletion), sterol acyltransferases (*ARE1* deletion),  $\beta$ -oxidation (*POX1* deletion), glycerol utilization (*GUT2* deletion), the acyl-CoA peroxisomal transporter (*PXA1* deletion) and fatty acyl-CoA synthetase (*FAA2* deletion). Strains were grown for 72 h in minimal medium containing 2% glucose and TAGs were quantified from biological quadruplicates. \* *p* value < .05 (Student's T-Test: two-tailed, two-sample equal variance).

The neutral lipids stored in lipid droplets are normally composed of 50% TAGs and 50% steryl esters (SEs) (Kohlwein et al., 2013). An SE results from the esterification of an acyl-CoA with a hydroxyl group of a sterol, catalyzed by acyl-CoA: sterol acyltransferases encoded by the *S. cerevisiae* genes *ARE1* and *ARE2* (Yang et al., 1996). We proposed to remove sterol ester formation through deletion of *ARE1* and *ARE2* in order to obtain a larger fraction of the neutral lipids in form of triacylglycerols. Removal of the major lipase genes *TGL3*, *TGL4* and *TGL5* and deletion of *ARE1* resulted in the new strain RF08. Although *ARE2* has been reported to be the major SE synthase (Sandager et al., 2002), we faced major technical difficulties in removing the gene during several attempts and therefore this deletion was not included in the following strain construction. RF08 showed a 68% increase in TAGs levels compared to strain ADP, with a final concentration of 218 mg-gCDW<sup>-1</sup> (Fig. 2). The resulting strain did not show any change in the final OD in comparison to the reference strain (Fig. S2).

### 3.3. Removal of the fatty acyl-CoA peroxisomal transporter further increases TAG levels

In *S. cerevisiae*, acyl-CoA is degraded back to acetyl-CoA by the peroxisomal  $\beta$ -oxidation pathway, and inactivation of this pathway leads to improved production of acyl-CoA derived chemicals (Valle-Rodríguez et al., 2014). Removal of the first step of the  $\beta$ -oxidation encoded by *POX1*, resulted in the strain RF09 which accumulated 204 mg-gCDW<sup>-1</sup> of TAGs, a 9% decrease ( $p > .05$ ) in comparison to RF08 (Fig. 2).

Next, we targeted the Gro-3-P utilization pathway by deleting the glycerol-3-phosphate dehydrogenase encoding gene *GUT2*. Gut2 catalyzes the conversion of Gro-3-P to dihydroxyacetone phosphate (DHAP) (Rønnow et al., 1993). Removal of *GUT2* resulted in the strain RF10, which accumulates TAGs at a level of 222 mg-gCDW<sup>-1</sup>, similar to the level of RF08 (Fig. 2).

We deleted *PXA1* encoding a subunit of the peroxisomal transporter involved in the import of acyl-CoA (Shani et al., 1995). The resulting strain RF11 reached a TAG level of 254 mg-gCDW<sup>-1</sup>, a 96% increase in comparison to the ADP strain ( $p$  value < .05), and a 14% increase in comparison to RF10 ( $p$  value < .05) (Fig. 2). The TAG titers produced by RF11 of 1.76 g-L<sup>-1</sup> from 20 g-L<sup>-1</sup> glucose translates to 27.4% of the maximum theoretical yield when considering the theoretical yield to be 0.32 gTAG-gGlucose<sup>-1</sup> (Table 2) (Caspeta and Nielsen, 2013). Notably, the titer achieved in RF11 is the highest reported in *S. cerevisiae* (Table S1). Finally, we deleted a gene encoding a peroxisomal fatty acid activating enzyme, *FAA2*, leading to strain RF12 strain. The final strain accumulated a similar level of TAG (248 mg-gCDW<sup>-1</sup>) compared to RF11.

## 4. Discussion

Lipid metabolism in eukaryotes is complex due to its crosstalk with essential cellular processes, such as synthesis of signaling molecules and cellular membranes. For this reason, it is tightly regulated at several levels, often making it a difficult target for metabolic engineering.

**Table 2**  
Yield calculations.

Strain	mgTAG/gDCW	Biomass (gDCW/L)	Titer (gTAG/L)	Yield (gTAG/gGLuc)	%Theoretical Yield*
IMX581	7.45 ± 1.51	7.30 ± 0.07	0.05 ± 0.01	0.003 ± 0.001	0.8% ± 0.2
ADP	129.30 ± 10.05	7.90 ± 0.26	1.02 ± 0.10	0.051 ± 0.005	16.0% ± 1.6
RF08	217.78 ± 12.43	7.95 ± 0.18	1.73 ± 0.10	0.087 ± 0.005	27.0% ± 1.6
RF09	201.23 ± 10.18	8.00 ± 0.22	1.61 ± 0.05	0.080 ± 0.003	25.1% ± 0.8
RF10	221.86 ± 20.77	6.74 ± 0.17	1.50 ± 0.17	0.075 ± 0.009	23.4% ± 2.7
RF11	254.02 ± 19.87	6.91 ± 0.04	1.76 ± 0.15	0.088 ± 0.007	27.4% ± 2.3
RF12	247.87 ± 10.13	6.73 ± 0.27	1.67 ± 0.12	0.083 ± 0.006	26.1% ± 1.8

\* Assuming 0.32 g/g as theoretical maximum yield.

Here, we started by implementing a typical push-and-pull strategy towards TAG accumulation. For that, engineering of *Acc1* at the transcriptional and phosphorylation level was a fundamental step due to the importance of this enzyme in controlling fatty acid metabolism. The initial approach of overexpressing genes encoding an *Acc1* with two removed phosphorylation sites, together with *Pah1* and *Dga1* resulted in TAG accumulation levels up to 12% of dry cell biomass, more than 10-fold above the levels in the reference strain. A similar strategy of overexpressing *ACC1* and *DGA1* in *Y. lipolytica* reached to a lipid content of 41.4% (Tai and Stephanopoulos, 2013). Previous studies based on individual expression of these three genes have reported similar titer improvements, with more pronounced effect for *Acc1* (Hofbauer et al., 2014; Hung et al., 2016; Runguphan and Keasling, 2014). Acetyl- and malonyl-CoA constitute the building blocks of fatty-acid biosynthesis, and these results show the importance of the *Acc1* level and activity being a central metabolic knot of the lipid metabolism in *S. cerevisiae*.

Next we removed most lipase activity and part of the sterol ester synthesis activity and observed an TAG increase of up to 80%, reaching close to 22% of the dry cell weight. These results strongly indicate that in the ADP strain, lipid homeostasis mechanisms are counterbalancing TAG accumulation with TAG hydrolysis, therefore limiting further production. Supporting this, previous reports have shown that lipase activity at transcriptional and protein level is affected by neutral lipid levels (Klein et al., 2016). Based on similar experiments realized in *S. cerevisiae*, we speculate that the removal of the lipases has had the major role in increasing TAG levels (Sandager et al., 2002; Athenstaedt and Daum, 2003).

In contrast with previous results obtained in *Y. lipolytica*, the disruption of the glycerol consumption gene, *GUT2*, did not lead to a significant increase in TAGs (Dulermo and Nicaud, 2011). Here, we hypothesize that the flux towards TAG biosynthesis is not constrained by the availability of Gro-3-P. This diverges from studies made in *Y. lipolytica*, where TAGs can accumulate to 80% of the cells' dry weight. This high lipid content in *Y. lipolytica* might indicate that fatty acid synthesis exceeds Gro-3-P availability in this organism, making Gro-3-P a limiting factor in TAG synthesis, which does not seem to be the case in *S. cerevisiae*. While not tackled in this study, recent studies have reported an improvement of TAG levels when overexpressing heterologous GPAT and LPAT genes, which suggests a potential additional engineering strategy reach higher TAG titers (Wei et al., 2017). Overexpression of the endogenous GPAT and LPAT genes could be considered as well (Fig. 1).

Additionally, neither the disruption of the first reaction of the  $\beta$ -oxidation, catalyzed by *Pox1*, nor removal of the import activation of FFAs into the peroxisomes catalyzed by *Faa2*, affected the TAG levels. In *S. cerevisiae*, in normal condition, a wild-type strain only produces very low levels of FFAs, as they are directly and quickly re-activated by fatty acyl-CoA synthetases to fatty acyl-CoAs (Zhou et al., 2016). Although previous studies have shown that *POX1* disruption increased TAGs in *S. cerevisiae* and *Y. lipolytica* (Valle-Rodríguez et al., 2014; Dulermo and Nicaud, 2011), the removal of the major TAG lipases further decreased the supply of FFAs, which could explain the null effect of *POX1* disruption. Additionally, the null effect from removing

FAA2 further confirms this hypothesis. Presumably, FAA2 disruption would have allowed more cytosolic FFA substrates to be converted to fatty acyl-CoA, ultimately resulting in more TAGs.

Finally, disruption of the peroxisomal fatty acyl-CoA transporter gene, *PXA1*, resulted in a 14% ( $p < .05$ ) increase of the TAG level. Pxa1 transports fatty acyl-CoA to the peroxisome which ultimately decreases the levels of available substrate for TAG biosynthesis. Altogether, these results highlight a potential limitation in cytosolic acyl-CoA availability for TAG biosynthesis.

We envision that other aspects of lipid metabolism can be engineered for a higher level of TAG accumulation. These include improvements in fatty acid biosynthesis, such as improving FAS activity, further enhancing the supply of acetyl-CoA and malonyl-CoA, or NADPH cofactor balancing (Chen et al., 2013, Yu et al., 2017; Qiao et al., 2017, Hung et al., 2016).

In conclusion, the strains generated in this study provide good platforms on which further development can be implemented to generate efficient TAG-producing yeast cell factories. Furthermore, through generation of these strains we gained new insight into the regulation of TAG biosynthesis in the important model organism *S. cerevisiae*.

## Acknowledgements

We thank the Chalmers Mass Spectrometry Infrastructure team for their help with the HPLC-CAD analysis. We also thank David Bergenholm for helpful feedback.

## Funding

This work was funded by the Novo Nordisk Foundation, Swedish Foundation for Strategic Research, the Knut and Alice Wallenberg foundation, and the U.S. Department of Energy, Office of Science, Office of Biological and Environmental Research, Genomic Science program, under Award Number DE-SC0008744.

## Author's contribution

R.F, V.S, F.D and J.N conceived and designed the project. R.F, P.G.T and M.G performed the experiments. R.F. and P.G.T wrote the manuscript. All authors read and approved the manuscript.

## Appendix A. Supporting information

Supplementary data associated with this article can be found in the online version at <http://dx.doi.org/10.1016/j.meteno.2018.01.002>.

## References

- Athenstaedt, K., Daum, G., 2003. *YMR313c/TGL3* encodes a novel triacylglycerol lipase located in lipid particles of *Saccharomyces cerevisiae*. *J. Biol. Chem.*
- Athenstaedt, K., Daum, G., 2005. Tgl4p and Tgl5p, two triacylglycerol lipases of the yeast *Saccharomyces cerevisiae* are localized to lipid particles. *J. Biol. Chem.*
- Beopoulos, A., Mrozova, Z., Thevenieau, F., Le Dall, M.T., Hapala, I., Papanikolaou, S., Chardot, T., Nicaud, J.M., 2008. Control of lipid accumulation in the yeast *Yarrowia lipolytica*. *Appl. Environ. Microbiol.*
- Blank, H.M., Perez, R., He, C., Maitra, N., Metz, R., Hill, J., Lin, Y., Johnson, C.D., Bankaitis, V.A., Kennedy, B.K., Aramayo, R., Polymenis, M., 2017. Translational control of lipogenic enzymes in the cell cycle of synchronous, growing yeast cells. *EMBO J.* 36, 487–502. <http://dx.doi.org/10.15252/embj.201695050>.
- Bozaquel-Morais, B.L., Madeira, J.B., Venâncio, T.M., Pacheco-Rosa, T., Masuda, C.A., Montero-Lomeli, M., 2017. A chemogenomic screen reveals novel Snf1p/AMPK independent regulators of acetyl-CoA carboxylase. *PLoS One* 12, 1–20. <http://dx.doi.org/10.1371/journal.pone.0169682>.
- Caspetta, L., Nielsen, J., 2013. Economic and environmental impacts of microbial bio-diesel. *Nat. Biotechnol.* 31 (9), 789–793 (Sep 1).
- Chen, Y., Daviet, L., Schalk, M., Siewers, V., Nielsen, J., 2013. Establishing a platform cell factory through engineering of yeast acetyl-CoA metabolism. *Metab. Eng.*
- Dicarlo, J.E., Norville, J.E., Rios, X., Aach, J., Church, G.M., 2013. Genome engineering in *Saccharomyces Cerevisiae* using CRISPR-Cas Systems, 41, pp. 4336–4343. doi:10.1093/nar/gkt135.
- Dulermo, T., Nicaud, J.M., 2011. Involvement of the G3P shuttle and  $\beta$ -oxidation

- pathway in the control of tag synthesis and lipid accumulation in *Yarrowia lipolytica*. *Metab. Eng.*
- Gietz, R.D., Schiestl, R.H., 2007. High-efficiency yeast transformation using the LiAc/SS carrier DNA/PEG method. *Nature protocols* 2 (1), 31–34.
- Hasslachner, M., Ivessa, A.S., Paltauf, F., Kohlwein, S.D., 1993. Acetyl-CoA carboxylase from yeast is an essential enzyme and is regulated by factors that control phospholipid metabolism. *J. Biol. Chem.*
- Hofbauer, H.F., Schopf, F.H., Schleifer, H., Knittelfelder, O.L., Pieber, B., Rechberger, G.N., Wolinski, H., Gaspar, M.L., Kappe, C.O., Stadlmann, J., Mechtler, K., 2014. Regulation of gene expression through a transcriptional repressor that senses acyl-chain length in membrane phospholipids. *Dev. Cell.*
- Hung, C.H., Kanehara, K., Nakamura, Y., 2016. In vivo reconstitution of algal triacylglycerol production in *Saccharomyces cerevisiae*. *Front. Microbiol.* 7.
- Jakočiūnas, T., Bonde, I., Herrgård, M., Harrison, S.J., Kristensen, M., Pedersen, L.E., Jensen, M.K., Keasling, J.D., 2015. Multiplex Metabolic Pathway Engineering using CRISPR / Cas9 in *Saccharomyces Cerevisiae*, 28, pp. 213–222. doi:10.1016/j.ymben.2015.01.008.
- Jensen, E.D., Ferreira, R., Jakočiūnas, T., Arsovska, D., Zhang, J., Ding, L., Smith, J.D., David, F., Nielsen, J., Jensen, M.K., Keasling, J.D., 2017. Transcriptional reprogramming in yeast using dCas9 and combinatorial gRNA strategies. *Microb. Cell Fact.*
- Jensen, N.B., Strucko, T., Kildegaard, K.R., David, F., Maury, J., Mortensen, U.H., Forster, J., Nielsen, J., Borodina, I., 2014. EasyClone: method for iterative chromosomal integration of multiple genes *Saccharomyces cerevisiae*. *FEMS Yeast Res.*
- Jullesson, D., David, F., Pfeleger, B., Nielsen, J., 2015. Impact of synthetic biology and metabolic engineering on industrial production of fine chemicals. *Biotechnol. Adv.* 33, 1395–1402. <http://dx.doi.org/10.1016/j.biotechadv.2015.02.011>.
- Khoomrung, S., Chumnanpuen, P., Jansa-Ard, S., Ståhlman, M., Nookaew, I., Borén, J., Nielsen, J., 2013. Rapid quantification of yeast lipid using microwave-assisted total lipid extraction and HPLC-CAD. *Anal. Chem.*
- Klein, I., Klug, L., Schmidt, C., Martina, Z., Korber, M., Daum, G., Athenstaedt, K., 2016. Regulation of the yeast triacylglycerol lipases Tgl4p and Tgl5p by the presence/absence of nonpolar lipids. *Mol. Biol. Cell* 27 (13), 2014–2024 (Jul 1).
- Kohlwein, Sepp D., Veenhuis, Marten, van der Klei, Ida J., 2013. "Lipid droplets and peroxisomes: key players in cellular lipid homeostasis or a matter of fat—store'em up or burn'em down. *Genetics* 193.1, 1–50.
- Krivoruchko, A., Nielsen, J., 2015. Production of natural products through metabolic engineering of *Saccharomyces cerevisiae*. *Curr. Opin. Biotechnol.* 35, 7–15. <http://dx.doi.org/10.1016/j.copbio.2014.12.004>.
- Mans, Robert, et al., 2015. CRISPR/Cas9: a molecular Swiss army knife for simultaneous introduction of multiple genetic modifications in *Saccharomyces cerevisiae*. *FEMS Yeast Res.* 15, 2.
- Nielsen, J., Keasling, J.D., 2016. *Eng. Cell. Metab.* Cell 164, 1185–1197. <http://dx.doi.org/10.1016/j.cell.2016.02.004>.
- Partow, S., Siewers, V., Bjørn, S., Nielsen, J., Maury, J., 2010. Characterization of different promoters for designing a new expression vector in *Saccharomyces cerevisiae*. *Yeast.*
- Ploier, B., Scharwey, M., Koch, B., Schmidt, C., Schatte, J., Rechberger, G., Kollroser, M., Hermetter, A., Daum, G., 2013. Screening for hydrolytic enzymes reveals Ayr1p as a novel triacylglycerol lipase in *Saccharomyces cerevisiae*. *J. Biol. Chem.*
- Qiao, K., Wasylenko, T.M., Zhou, K., Xu, P., Stephanopoulos, G., 2017. Lipid production in *Yarrowia lipolytica* is maximized by engineering cytosolic redox metabolism. *Nat. Biotechnol.*
- Quan, J., Tian, J., 2011. Circular polymerase extension cloning for high-throughput cloning of complex and combinatorial DNA libraries. *Nat. Protoc.* 6 (2), 242 (Feb 1).
- Reifenberger, E., Boles, E., Ciriacy, M., 1997. Kinetic characterization of individual hexose transporters of *Saccharomyces cerevisiae* and their relation to the triggering mechanisms of glucose repression. *FEBS J.*
- Runguphan, W., Keasling, J.D., 2014. Metabolic engineering of *Saccharomyces cerevisiae* for production of fatty acid-derived biofuels and chemicals. *Metab. Eng.*
- Rønnow, B., Kielland-Brandt, M.C., 1993. *GUT2*, a gene for mitochondrial glycerol 3-phosphate dehydrogenase of *Saccharomyces cerevisiae*. *Yeast* 9, 1121–1130. <http://dx.doi.org/10.1002/yea.320091013>.
- Sandager, L., Gustavsson, M.H., Ståhl, U., Dahlqvist, A., Wiberg, E., Banas, A., Lenman, M., Ronne, H., Stymne, S., 2002. Storage lipid synthesis is non-essential in yeast. *J. Biol. Chem.*
- Schweizer, M., Lebert, C., Höltke, J., Roberts, L.M., Schweizer, E., 1984. Molecular cloning of the yeast fatty acid synthetase genes, *FAS1* and *FAS2*: illustrating the structure of the *FAS1* cluster gene by transcript mapping and transformation studies. *MGG Mol. Gen. Genet.* 194, 457–465. <http://dx.doi.org/10.1007/BF00425558>.
- Shani, N., Watkins, P.A., Valle, D., 1995. *PXA1*, a possible *Saccharomyces cerevisiae* ortholog of the human adrenoleukodystrophy gene. *Proc. Natl. Acad. Sci.* 92 (13), 6012–6016 (Jun 20).
- Shi, S., Chen, Y., Siewers, V., Nielsen, J., 2014. Improving production of malonyl coenzyme A-derived metabolites by abolishing Snf1-dependent regulation of Accl1. *MBio* 5, 1–8. <http://dx.doi.org/10.1128/mBio.01130-14>.
- Tai, M., Stephanopoulos, G., 2013. Engineering the push and pull of lipid biosynthesis in oleaginous yeast *Yarrowia lipolytica* for biofuel production. *Metab. Eng.* 15, 1–9 (Jan 31).
- Teixeira, P.G., Ferreira, R., Zhou, Y.J., Siewers, V., Nielsen, J., 2017. Dynamic regulation of fatty acid pools for improved production of fatty alcohols in *Saccharomyces cerevisiae*. *Microb. Cell Fact.* 16, 45. <http://dx.doi.org/10.1186/s12934-017-0663-3>.
- Valle-Rodríguez, J.O., Shi, S., Siewers, V., Nielsen, J., 2014. Metabolic engineering of *Saccharomyces cerevisiae* for production of fatty acid ethyl esters, an advanced biofuel, by eliminating non-essential fatty acid utilization pathways. *Appl. Energy.*
- Verduyn, C., Postma, E., Scheffers, W.A., Van Dijken, J.P., 1992. Effect of benzoic acid on

- metabolic fluxes in yeasts: a continuous-culture study on the regulation of respiration and alcoholic fermentation. *Yeast* 8 (7), 501–517.
- Wei, Y., Gossing, M., Bergenholm, D., Siewers, V., Nielsen, J., 2017. Increasing cocoa butter-like lipid production of *Saccharomyces cerevisiae* by expression of selected cocoa genes. *AMB Express* 7 (1), 34 (Dec 1).
- Yu, T., Zhou, Y.J., Wenning, L., Liu, Q., Krivoruchko, A., Siewers, V., Nielsen, J., David, F., 2017. Metabolic engineering of *Saccharomyces cerevisiae* for production of very long chain fatty acid-derived chemicals. *Nat. Commun.*
- Zalatan, J.G., Lee, M.E., Almeida, R., Gilbert, L. a., Whitehead, E.H., La Russa, M., Tsai, J.C., Weissman, J.S., Dueber, J.E., Qi, L.S., Lim, W. a., 2015. Engineering complex synthetic transcriptional programs with CRISPR RNA scaffolds. *Cell* 9, 1–12. <http://dx.doi.org/10.1016/j.cell.2014.11.052>.
- Zhou, Y.J., Buijs, N.A., Zhu, Z., Qin, J., Siewers, V., Nielsen, J., 2016. Production of fatty acid-derived oleochemicals and biofuels by synthetic yeast cell factories. *Nat. Commun.* 7, 11709. <http://dx.doi.org/10.1038/ncomms11709>.