



QPEL: Quantum Program and Effect Language

Downloaded from: <https://research.chalmers.se>, 2019-03-24 18:17 UTC

Citation for the original published paper (version of record):

Adams, R. (2014)

QPEL: Quantum Program and Effect Language

Electronic Proceedings in Theoretical Computer Science, 172: 133-153

<http://dx.doi.org/10.4204/EPTCS.172.10>

N.B. When citing this work, cite the original published paper.

QPEL: Quantum Program and Effect Language

Robin Adams

Radboud University Nijmegen

r.adams@cs.ru.nl

We present the syntax and rules of deduction of QPEL (Quantum Program and Effect Language), a language for describing both quantum programs, and properties of quantum programs — *effects* on the appropriate Hilbert space. We show how semantics may be given in terms of *state-and-effect triangles*, a categorical setting that allows semantics in terms of Hilbert spaces, C^* -algebras, and other categories. We prove soundness and completeness results that show the derivable judgements are exactly those provable in all state-and-effect triangles.

1 Introduction

There is a growing number of quantum programming languages, and there is a need for a syntactic method of reasoning about these quantum programs: both in the hope of making automated tools for proving the correctness of programs, and because experience in other fields shows that many problems that are difficult when treated semantically

We present QPEL, a syntax for both describing quantum programs, and properties of quantum programs (quantum predicates, or effects). This system should be useful for reasoning about quantum programs and proving their correctness, as well as showing more generally how a language for quantum effects may be added on top of any quantum programming language. The part of the system that describes quantum programs is loosely based on Selinger’s Quantum Programming Language (QPL) [16].

The part of the system that describes quantum programs is a *linear* type theory (see [6, 7]): we are not able to duplicate data. Duplication of quantum data would violate the no-cloning theorem. We do allow deletion of data (which corresponds to e.g. measuring a qubit then throwing away the outcome of the measurement).

The part of the system that describes quantum predicates is based on the fact that the effects on a Hilbert state or C^* -algebra form an *effect algebra* - in fact, an *effect module* over the appropriate effect monoid [14].

There is a categorical structure called the *state-and-effect triangle* that has been shown to generalise several different ways of giving semantics to quantum computing, including Hilbert spaces and C^* -algebras. The first version of QPEL we present captures all and only the structure of a state-and-effect triangle. We show how to give semantics in an arbitrary triangle, and prove a Soundness and Completeness Theorem. We proceed to discuss what would need to be added to the system to represent other features of a quantum programming language, particularly qubits.

The language QPEL has a homepage at www.cs.ru.nl/~robina/QPEL

2 Preliminaries

2.1 Notation

If E and F are expressions involving partial functions, we write:

- $E = F$ to denote: E and F are both defined, and their values are equal;
- $E \simeq F$ to denote: E is defined if and only if F is defined, in which case their values are equal (this is sometimes known as *Kleene equality*);
- $E \xrightarrow{\sim} F$ to denote: if E is defined, then F is defined and their values are equal (this is sometimes known as *directed equality*).

2.2 Effect Algebras and Effect Monoids

We represent the effects on a quantum system by the elements of an *effect module* over an *effect monoid* M , whose elements we call *scalars*. The canonical example is the effects on a Hilbert space or C^* -algebra, which form an effect module over $[0, 1]$, with the scalars being probabilities. These concepts were introduced in [12].

Definition 1 (Partial Commutative Monoid). A *partial commutative monoid* consists of a set M ; an element $0 \in M$, the *zero*; and a partial binary operation $\odot : M^2 \rightarrow M$, the (*partial*) *sum*; such that:

- $x \odot y \simeq y \odot x$
- $x \odot (y \odot z) \simeq (x \odot y) \odot z$
- $x \odot 0 = x$

for all $x, y, z \in M$.

We write $x \perp y$, x is *orthogonal* to y , iff $x \odot y$ is defined.

Definition 2 (Effect Algebra). An *effect algebra* is a partial commutative monoid E with a (total) function $(-)^{\perp} : E \rightarrow E$, the *orthosupplement*, such that

- $x \odot y = 0^{\perp}$ iff $y = x^{\perp}$.
- If $x \perp 0^{\perp}$ then $x = 0$.

We write 1 for 0^{\perp} .

Definition 3 (Effect Algebra Homomorphism). Let E and F be effect algebras. An *effect algebra homomorphism* $\phi : E \rightarrow F$ is a function such that, for all $x, y \in E$:

$$\begin{aligned}\phi(x \odot y) &\xrightarrow{\sim} \phi(x) \odot \phi(y) \\ \phi(x^{\perp}) &= \phi(x)^{\perp}\end{aligned}$$

Lemma 4. For any effect algebra homomorphism ϕ , we have $\phi(0) = 0$.

Proof.

$$\begin{aligned}\phi(0 \odot 0) &= \phi(0) \\ \therefore \phi(0) \odot \phi(0) &= \phi(0) \\ &= \phi(0) \odot 0 \\ \therefore \phi(0) &= 0\end{aligned}$$

since any effect algebra satisfies cancellation [10]. □

Definition 5 (Effect Monoid). An *effect monoid* is an effect algebra E with a binary operation $\cdot : E^2 \rightarrow E$, the *multiplication*, such that

- $(x \otimes y) \cdot z \xrightarrow{\sim} (x \cdot z) \otimes (y \cdot z)$
- $x \cdot (y \otimes z) \xrightarrow{\sim} (x \cdot y) \otimes (x \cdot z)$
- $1 \cdot x = x \cdot 1 = x$
- $x \cdot (y \cdot z) = (x \cdot y) \cdot z$

The effect monoid is *commutative* iff $x \cdot y = y \cdot x$ for all x, y .

An effect monoid is a monoid in the category of effect algebras [12].

Lemma 6. In any effect monoid, $x \cdot 0 = 0 \cdot x = 0$.

Proof. We have

$$\begin{aligned} x \cdot (0 \otimes 0) &= x \cdot 0 \\ \therefore x \cdot 0 \otimes x \cdot 0 &= x \cdot 0 \\ &= x \cdot 0 \otimes 0 \\ \therefore x \cdot 0 &= 0 \end{aligned}$$

by cancellation. Similarly for $0 \cdot x$. □

Definition 7 (Effect Module). An *effect module* over an effect monoid E is an effect algebra A with a binary operation $\cdot : E \times A \rightarrow A$ called *scalar multiplication* such that, for all $x, y, z \in E$:

- $r \cdot (x \otimes y) \xrightarrow{\sim} (r \cdot x) \otimes (r \cdot y)$
- $(r \otimes s) \cdot x \xrightarrow{\sim} (r \cdot x) \otimes (s \cdot x)$
- $(r \cdot s) \cdot x = r \cdot (s \cdot x)$
- $1 \cdot x = x$

Definition 8 (Effect Module Homomorphism). Let A and B be effect modules over E . An *effect module homomorphism* $\phi : A \rightarrow B$ is an effect algebra homomorphism such that, for all $r \in E$ and $x \in A$,

$$\phi(r \cdot x) = r \cdot \phi(x) .$$

2.2.1 Examples

1. For any Hilbert space H , the set of effects over H forms an effect module over the effect monoid $[0, 1]$, with $F \otimes G = F + G$ iff $F + G$ is an effect [14].
2. Given a C^* -algebra A , the set of *effects* in A (positive elements below the unit) form an effect module over the real numbers $[0, 1]$.

2.3 Convex Sets

We describe the category of *convex sets* over any effect monoid. The *states* of a quantum system will form a convex set over the effect monoid of probabilities.

Definition 9. Given an effect monoid E , the *distribution monad* $\mathcal{D}_E : \mathbf{Set} \rightarrow \mathbf{Set}$ is defined as follows.

$$\mathcal{D}_E X = \{ \phi : X \rightarrow E : \text{supp } \phi \text{ is finite, } \sum_{x \in X} \phi(x) \text{ exists and is equal to } 1 \}$$

where $\text{supp } \phi = \{x \in X : \phi(x) \neq 0\}$.

For $f : X \rightarrow Y$,

$$\mathcal{D}_E f(\phi)(y) = \sum_{f(x)=y} \phi(x) \quad (\phi \in \mathcal{D}_E X, y \in Y) .$$

The unit $\eta_A : A \rightarrow \mathcal{D}_E A$ is defined by

$$\eta_A(a)(a') = \begin{cases} 1 & \text{if } a = a' \\ 0 & \text{if } a \neq a' \end{cases}$$

The multiplication $\mu_A : \mathcal{D}_E^2 A \rightarrow \mathcal{D}_E A$ is defined by

$$\mu_A(\Phi)(a) = \sum_{\phi \in \mathcal{D}_E A} (\Phi(\phi) \cdot \phi(a)) .$$

The category \mathbf{Conv}_E of *convex sets* and *affine functions* over E is the Eilenberg-Moore category of \mathcal{D}_E . A convex set may thus be thought of as a set X together with a function mapping any finite tuple $\langle r_1, \dots, r_n \rangle$ of elements of M that sum to 1, and any tuple $\langle x_1, \dots, x_n \rangle$ of elements of X , to an element $r_1 x_1 + \dots + r_n x_n$ of X .

Theorem 10. *The distribution monad is a strong monad. It is a commutative monad iff E is commutative.*

Proof. The tensorial strength $t_{AB} : A \times \mathcal{D}_E B \rightarrow \mathcal{D}_E(A \times B)$ is given by

$$t_{AB}(a, \phi)(a', b) = \begin{cases} \phi(b) & \text{if } a = a' \\ 0 & \text{if } a \neq a' \end{cases}$$

□

Corollary 10.1. *If E is commutative, then \mathbf{Conv}_E is a symmetric monoidal category.*

Proof. See [15].

□

The convex set $A \otimes B$ consists of all sums $r_1(a_1, b_1) + \dots + r_n(a_n, b_n)$ ($r_1 \otimes \dots \otimes r_n = 1, a_i \in A, b_i \in B$), quotiented by the appropriate equivalence relation. An affine function $f : A \otimes B \rightarrow C$ in \mathbf{Conv}_M is determined by the values $f(a, b)$ for $a \in A$ and $b \in B$

Theorem 11. *The hom-functors $\mathbf{Conv}_E[-, E] \dashv \mathbf{EMod}_E[-, E] : \mathbf{Conv}_E \rightleftarrows \mathbf{EMod}_E^{\text{op}}$ form an adjunction.*

Proof. To appear in [13]. The special case $E = [0, 1]$ was proved in [11].

□

3 Syntax and Rules of Deduction

We begin with a system that represents a symmetric monoidal closed category with distributive coproducts, with an effect module of predicates over each object.

Type	$A ::= A \otimes A \mid I \mid A + B$
Term	$M ::= x \mid M \otimes M \mid \text{let } x \otimes x = M \text{ in } M \mid \langle \rangle \mid$ $\text{inl}(M) \mid \text{inr}(M) \mid$ $(\text{case } M \text{ of } \text{inl}(x) \mapsto M \mid \text{inr}(x) \mapsto M) \mid$ $(\text{measure } \phi \mapsto M \mid \dots \mid \phi \mapsto M)$
Effect	$\phi ::= 0 \mid \phi \otimes \phi \mid \phi^\perp \mid \phi \cdot \psi \mid (\text{case } M \text{ of } \text{inl}(x) \mapsto \phi \mid \text{inr}(x) \mapsto \phi)$
Context	$\Gamma ::= \langle \rangle \mid \Gamma, x : A$
Judgement	$\mathcal{J} ::= \Gamma \vdash M : A \mid \Gamma \vdash M = N : A \mid \Gamma \vdash \phi \text{ eff} \mid \Gamma \vdash \phi \leq \psi$

The intuition is as follows:

- Each type represents a state space for a quantum computer at some stage of a calculation. For example, the type $(\mathbf{qbit} \otimes \mathbf{qbit}) + (\mathbf{qbit} \otimes \mathbf{qbit} \otimes \mathbf{qbit})$ represents a computer that has either two or three qubits in memory (depending on decisions earlier in the program). (The type \mathbf{qbit} will be introduced in Section 5.)

The type I represents a singleton data type. A term of type $A \otimes B$ is a pair consisting of a term of type A and a term of type B (possibly entangled). A term of type $A + B$ is either a term of type A or a term of type B (with 'or' understood here classically).

- A term M such that $\Gamma \vdash M : A$ represents a quantum algorithm that takes inputs as given by the context Γ , and returns an output of type A .

If the judgement $\Gamma \vdash M = N : A$ is derivable, then the algorithms M and N always produce the same output state given the same input state.

- An effect in context Γ represents an observable measurement that may be performed on the system denoted by Γ .

The effect 0 is the always false effect. The effect $\phi \otimes \psi$ is the sum of ϕ and ψ , which may only be formed if ϕ and ψ are orthogonal. The effect ϕ^\perp is the orthocomplement of ϕ .

We write

$$\begin{aligned}
 & 1 \text{ for } 0^\perp \\
 & \bigotimes_{i=1}^n \phi_i \text{ for } ((\dots(\phi_1 \otimes \phi_2) \otimes \dots) \otimes \phi_n) \\
 & \text{measure } \bigotimes_{i=1}^n \phi_i \mapsto M_i \text{ for } \text{measure } \phi_1 \mapsto M_1 \mid \dots \mid \phi_n \mapsto M_n \\
 & \phi \perp \psi \text{ for } \phi \leq \psi^\perp
 \end{aligned}$$

We write $\Gamma \vdash \phi \equiv \psi$ for the two judgements $\Gamma \vdash \phi \leq \psi$ and $\Gamma \vdash \psi \leq \phi$.

The rules of deduction are as follows.

Note Note in particular the rule (\otimes) . For $\phi \otimes \psi$ to be a well-formed effect in context Γ , we must first have a derivation of $\Gamma \vdash \phi \perp \psi$, i.e. $\Gamma \vdash \phi \leq \psi^\perp$.

Structural Rule

$$\text{(exch)} \frac{\Gamma, x : A, y : B, \Delta \vdash \mathcal{J}}{\Gamma, y : B, x : A, \Delta \vdash \mathcal{J}}$$

Term Formation

$$\begin{aligned} & \text{(var)} \frac{}{\Gamma \vdash x : A} (x : A \in \Gamma) \\ & (\otimes) \frac{\Gamma \vdash M : A \quad \Delta \vdash N : B}{\Gamma, \Delta \vdash M \otimes N : A \otimes B} \\ & \text{(let)} \frac{\Gamma \vdash M : A \otimes B \quad \Delta, x : A, y : B \vdash N : C}{\Gamma, \Delta \vdash \text{let } x \otimes y = M \text{ in } N : C} \\ & (\langle \rangle) \frac{}{\Gamma \vdash \langle \rangle : I} \\ & \text{(inl)} \frac{\Gamma \vdash M : A}{\Gamma \vdash \text{inl}(M) : A + B} \quad \text{(inr)} \frac{\Gamma \vdash M : B}{\Gamma \vdash \text{inr}(M) : A + B} \\ & \text{(case)} \frac{\Gamma \vdash M : A + B \quad \Delta, x : A \vdash N : C \quad \Delta, y : B \vdash P : C}{\Gamma, \Delta \vdash \text{case } M \text{ of inl}(x) \mapsto N \mid \text{inr}(y) \mapsto P : C} \\ & \text{(measure)} \frac{\Gamma \vdash 1 \leq \bigotimes_{i=1}^n \phi_i \quad \Delta \vdash M_i : A \quad (1 \leq i \leq n)}{\Gamma, \Delta \vdash \text{measure } \bigotimes_{i=1}^n \phi_i \mapsto M_i : A} \end{aligned}$$

Equality of Terms

$$\begin{aligned} & \text{(ref)} \frac{\Gamma \vdash M : A}{\Gamma \vdash M = M : A} \quad \text{(sym)} \frac{\Gamma \vdash M = N : A}{\Gamma \vdash N = M : A} \\ & \text{(trans)} \frac{\Gamma \vdash M = N : A \quad \Gamma \vdash N = P : A}{\Gamma \vdash M = P : A} \end{aligned}$$

Congruences

$$\begin{aligned} & (\otimes\text{-eq}) \frac{\Gamma \vdash M = M' : A \quad \Delta \vdash N = N' : B}{\Gamma, \Delta \vdash M \otimes N = M' \otimes N' : A \otimes B} \\ & \text{(let-eq)} \frac{\Gamma \vdash M = M' : A \otimes B \quad \Delta, x : A, y : B \vdash N = N' : C}{\Gamma, \Delta \vdash (\text{let } x \otimes y = M \text{ in } N) = (\text{let } x \otimes y = M' \text{ in } N') : C} \\ & \text{(inl-eq)} \frac{\Gamma \vdash M = N : A}{\Gamma \vdash \text{inl}(M) = \text{inl}(N) : A + B} \quad \text{(inr-eq)} \frac{\Gamma \vdash M = N : B}{\Gamma \vdash \text{inr}(M) = \text{inr}(N) : A + B} \\ & \text{(case-eq)} \frac{\Gamma \vdash M = M' : A + B \quad \Delta, x : A \vdash N = N' : C \quad \Delta, y : B \vdash P = P' : C}{\Gamma, \Delta \vdash (\text{case } M \text{ of inl}(x) \mapsto N \mid \text{inr}(y) \mapsto P) = (\text{case } M' \text{ of inl}(x) \mapsto N' \mid \text{inr}(y) \mapsto P') : C} \\ & \text{(measure-eq)} \frac{\Gamma \vdash 1 \leq \bigotimes_{i=1}^n \phi_i \quad \Gamma \vdash \phi_i \equiv \psi_i \quad (1 \leq i \leq n) \quad \Delta \vdash M_i = N_i : A \quad (1 \leq i \leq n)}{\Gamma, \Delta \vdash (\text{measure } \bigotimes_{i=1}^n \phi_i \mapsto M_i) = (\text{measure } \bigotimes_{i=1}^n \psi_i \mapsto N_i) : A} \end{aligned}$$

β -conversions

$$(\beta_{\otimes}) \frac{\Gamma \vdash M : A \quad \Delta \vdash N : B \quad \Theta, x : A, y : B \vdash P : C}{\Gamma, \Delta, \Theta \vdash (\text{let } x \otimes y = M \otimes N \text{ in } P) = [M/x, N/y]P : C}$$

$$(\beta_{+1}) \frac{\Gamma \vdash M : A \quad \Delta, x : A \vdash N : C \quad \Delta, y : B \vdash P : C}{\Gamma, \Delta \vdash \text{case inl}(M) \text{ of inl}(x) \mapsto N \mid \text{inr}(y) \mapsto P = [M/x]N : C}$$

$$+ (\beta_{+2}) \frac{\Gamma \vdash M : B \quad \Delta, x : A \vdash N : C \quad \Delta, y : B \vdash P : C}{\Gamma, \Delta \vdash \text{case inr}(M) \text{ of inl}(x) \mapsto N \mid \text{inr}(y) \mapsto P = [M/y]P : C}$$

 η -conversions

$$(\eta_{\otimes}) \frac{\Gamma \vdash M : A \otimes B}{\Gamma \vdash M = \text{let } x \otimes y = M \text{ in } x \otimes y : A \otimes B} \quad (\eta_I) \frac{\Gamma \vdash M : I}{\Gamma \vdash M = \langle \rangle : I}$$

$$(\eta_{+}) \frac{\Gamma \vdash M : A + B}{\Gamma \vdash M = \text{case } M \text{ of inl}(x) \mapsto \text{inl}(x) \mid \text{inr}(y) \mapsto \text{inr}(y) : A + B}$$

Commuting Conversions

$$(\text{let-commute}) \frac{\Gamma \vdash M : A \otimes B \quad \Delta, x : A, y : B \vdash N : C \otimes D \quad \Theta, t : C, u : D \vdash P : E}{\Gamma, \Delta, \Theta \vdash (\text{let } x \otimes y = M \text{ in let } t \otimes u = N \text{ in } P) = (\text{let } t \otimes u = \text{let } x \otimes y = M \text{ in } N \text{ in } P) : E}$$

$$(\text{let-case}) \frac{\Gamma \vdash M : A + B \quad \Delta, x : A \vdash N : C \otimes D \quad \Delta, y : B \vdash P : C \otimes D \quad \Theta, z : C, t : D \vdash Q : E}{\Gamma, \Delta, \Theta \vdash (\text{let } z \otimes t = \text{case } M \text{ of inl}(x) \mapsto N \mid \text{inr}(y) \mapsto P \text{ in } Q) = \text{case } M \text{ of inl}(x) \mapsto \text{let } z \otimes t = N \text{ in } Q \mid \text{inr}(y) \mapsto \text{let } z \otimes t = P \text{ in } Q : E}$$

$$(\text{let-}\otimes) \frac{\Gamma \vdash M : A \otimes B \quad \Delta, x : A, y : B \vdash N : C \quad \Theta \vdash P : D}{\Gamma, \Delta, \Theta \vdash (\text{let } x \otimes y = M \text{ in } N) \otimes P = \text{let } x \otimes y = M \text{ in } (N \otimes P)}$$

$$(\text{case-commute}) \frac{\Gamma \vdash M : A + B \quad \Delta, x : A \vdash N : C + D \quad \Delta, y : B \vdash P : C + D \quad \Theta, z : C \vdash Q : E \quad \Theta, t : D \vdash R : E}{\Gamma, \Delta, \Theta \vdash \text{case } M \text{ of inl}(x) \mapsto (\text{case } N \text{ of inl}(z) \mapsto Q \mid \text{inr}(t) \mapsto R) \mid \text{inr}(y) \mapsto (\text{case } P \text{ of inl}(z) \mapsto Q \mid \text{inr}(t) \mapsto R) = \text{case } (\text{case } M \text{ of inl}(x) \mapsto N \mid \text{inr}(y) \mapsto P) \text{ of inl}(z) \mapsto Q \mid \text{inr}(t) \mapsto R : E}$$

$$(\text{case-}\otimes) \frac{\Gamma \vdash Q : A + B \quad \Delta, a : A \vdash M : C \quad \Delta, b : B \vdash N : C \quad \Theta \vdash P : D}{\Gamma, \Delta, \Theta \vdash (\text{case } Q \text{ of inl}(a) \mapsto M \mid \text{inr}(b) \mapsto N) \otimes P = \text{case } Q \text{ of inl}(a) \mapsto M \otimes P \mid \text{inr}(b) \mapsto N \otimes P : C \otimes D}$$

Rules for Measurement

$$\begin{array}{c}
\text{(measure-perm)} \frac{\Gamma \vdash 1 \leq \bigotimes_{i=1}^n \phi_i \quad \Delta \vdash M_x : A \quad (1 \leq i \leq n)}{\Gamma, \Delta \vdash (\text{measure } \bigotimes_{i=1}^n \phi_i \mapsto M_i)} \quad (p \text{ a permutation of } \{1, \dots, n\}) \\
= (\text{measure } \bigotimes_{i=1}^n \phi_{p(i)} \mapsto M_{p(i)}) : A \\
\\
\text{(measure-0)} \frac{\Gamma \vdash 1 \leq \bigotimes_{i=1}^n \phi_i \quad \Delta \vdash M_i : A \quad (1 \leq i \leq n+1)}{\Gamma, \Delta \vdash (\text{measure } \phi_1 \mapsto M_1 \mid \dots \mid \phi_n \mapsto M_n \mid 0 \mapsto M_{n+1})} \\
= \text{measure } \phi_1 \mapsto M_1 \mid \dots \mid \phi_n \mapsto M_n : A \\
\\
\text{(measure-1)} \frac{\Gamma \vdash M : A}{\Gamma \vdash (\text{measure } 1 \mapsto M) = M : A} \\
\\
\text{(measure-plus)} \frac{\vdash 1 \leq \phi \otimes \psi \otimes \chi_1 \otimes \dots \otimes \chi_n \quad \Gamma \vdash M : A \quad \Gamma \vdash P_1 : A \quad \dots \quad \Gamma \vdash P_n : A}{\Gamma \vdash (\text{measure } \phi \otimes \psi \mapsto M \mid \chi_1 \mapsto P_1 \mid \dots \mid \chi_n \mapsto P_n)} \\
= (\text{measure } \phi \mapsto M \mid \psi \mapsto M \mid \chi_1 \mapsto P_1 \mid \dots \mid \chi_n \mapsto P_n) \\
\\
\text{(measure-case)} \frac{\Gamma, x : A \vdash 1 \leq \bigotimes_{i=1}^n \phi_i \quad \Gamma, y : B \vdash 1 \leq \bigotimes_{i=1}^n \psi_i \quad \Delta \vdash M : A + B \quad \Theta \vdash N_i : C \quad (1 \leq i \leq n)}{\Gamma, \Delta, \Theta \vdash \text{measure } \bigotimes_{i=1}^n (\text{case } M \text{ of } \text{inl}(x) \mapsto \phi_i \mid \text{inr}(y) \mapsto \psi_i) \mapsto N_i} \\
= \text{case } M \text{ of } \text{inl}(x) \mapsto (\text{measure } \bigotimes_{i=1}^n \phi_i \mapsto N_i) \mid \\
\text{inr}(y) \mapsto (\text{measure } \bigotimes_{i=1}^n \psi_i \mapsto N_i)
\end{array}$$

Effect Formation

$$\begin{array}{c}
(0) \frac{}{\Gamma \vdash 0 \text{ eff}} \quad (\perp) \frac{\Gamma \vdash \phi \text{ eff}}{\Gamma \vdash \phi^\perp \text{ eff}} \quad (\otimes) \frac{\Gamma \vdash \phi \perp \psi}{\Gamma \vdash \phi \otimes \psi \text{ eff}} \\
\\
(\text{mult}) \frac{\vdash \phi \text{ eff} \quad \Gamma \vdash \psi \text{ eff}}{\Gamma \vdash \phi \cdot \psi \text{ eff}} \\
\\
(\text{case}) \frac{\Gamma, x : A \vdash \phi \text{ eff} \quad \Gamma, y : B \vdash \psi \text{ eff} \quad \Delta \vdash M : A + B}{\Gamma, \Delta \vdash \text{case } M \text{ of } \text{inl}(x) \mapsto \phi \mid \text{inr}(y) \mapsto \psi \text{ eff}}
\end{array}$$

Derivability

$$\begin{array}{c}
(\leq\text{-ref}) \frac{\Gamma \vdash \phi \text{ eff}}{\Gamma \vdash \phi \leq \phi} \quad (\leq\text{-trans}) \frac{\Gamma \vdash \phi \leq \psi \quad \Gamma \vdash \psi \leq \chi}{\Gamma \vdash \phi \leq \chi} \\
\\
(0\leq) \frac{\Gamma \vdash \phi \text{ eff}}{\Gamma \vdash 0 \leq \phi} \\
\\
(\perp\text{-antitone}) \frac{\Gamma \vdash \phi \leq \psi}{\Gamma \vdash \psi^\perp \leq \phi^\perp} \quad (\perp\perp_0) \frac{\Gamma \vdash \phi \text{ eff}}{\Gamma \vdash \phi \leq \phi^{\perp\perp}} \\
\\
(\leq\text{-}\otimes) \frac{\Gamma \vdash \phi \perp \psi}{\Gamma \vdash \phi \leq \phi \otimes \psi} \quad (\otimes\text{-mono}) \frac{\Gamma \vdash \phi \leq \psi \quad \Gamma \vdash \psi \leq \chi^\perp}{\Gamma \vdash \phi \otimes \chi \leq \psi \otimes \psi}
\end{array}$$

$$\begin{array}{c}
(\otimes\text{-comm}_{\leq}) \frac{\Gamma \vdash \phi \perp \psi}{\Gamma \vdash \phi \otimes \psi \leq \psi \otimes \phi} \quad (\perp\text{-rotate}) \frac{\Gamma \vdash \phi \otimes \psi \perp \chi}{\Gamma \vdash \psi \otimes \chi \perp \phi} \\
(\otimes\text{-assoc}_{\leq}) \frac{\Gamma \vdash \phi \otimes \psi \perp \chi}{\Gamma \vdash \phi \otimes (\psi \otimes \chi) \leq (\phi \otimes \psi) \otimes \chi} \quad (\otimes\text{-0}) \frac{\Gamma \vdash \phi \text{ eff}}{\Gamma \vdash \phi \otimes 0 \leq \phi} \\
(\text{ortho}_1) \frac{\Gamma \vdash 1 \leq \phi \otimes \psi}{\Gamma \vdash \psi^\perp \leq \phi} \quad (\text{ortho}_2) \frac{\Gamma \vdash \phi \text{ eff}}{\Gamma \vdash 1 \leq \phi \otimes \phi^\perp} \\
(\text{dist}_L) \frac{\vdash \phi \perp \psi \quad \Gamma \vdash \chi \text{ eff}}{\Gamma \vdash \phi \cdot \chi \perp \psi \cdot \chi} \quad (\text{dist}_R) \frac{\vdash \phi \text{ eff} \quad \Gamma \vdash \psi \perp \chi}{\Gamma \vdash \phi \cdot \psi \perp \phi \cdot \chi} \\
\Gamma \vdash (\phi \otimes \psi) \cdot \chi \equiv \phi \cdot \chi \otimes \psi \cdot \chi \quad \Gamma \vdash \phi \cdot (\psi \otimes \chi) \equiv \phi \cdot \psi \otimes \phi \cdot \chi \\
(\text{unit}_L) \frac{\Gamma \vdash \phi \text{ eff}}{\Gamma \vdash 1 \cdot \phi \equiv \phi} \quad (\text{unit}_R) \frac{\vdash \phi \text{ eff}}{\Gamma \vdash \phi \cdot 1 \equiv \phi} \quad (\text{assoc}) \frac{\vdash \phi \text{ eff} \quad \vdash \psi \text{ eff} \quad \Gamma \vdash \chi \text{ eff}}{\Gamma \vdash \phi \cdot (\psi \cdot \chi) \equiv (\phi \cdot \psi) \cdot \chi} \\
(\text{comm}) \frac{\vdash \phi \text{ eff} \quad \vdash \psi \text{ eff}}{\vdash \phi \cdot \psi \equiv \psi \cdot \phi} \\
(\text{case-cong}) \frac{\Gamma, x : A \vdash \phi \text{ eff} \quad \Gamma, y : B \vdash \psi \text{ eff} \quad \Delta \vdash M = N : A + B}{\Gamma, \Delta \vdash \text{case } M \text{ of } \text{inl}(x) \mapsto \phi \mid \text{inr}(y) \mapsto \psi \equiv \text{case } N \text{ of } \text{inl}(x) \mapsto \phi \mid \text{inr}(y) \mapsto \psi} \\
(\text{case-mono}) \frac{\Gamma, x : A \vdash \phi \leq \phi' \quad \Gamma, y : B \vdash \psi \leq \psi' \quad \Delta \vdash M : A + B}{\Gamma, \Delta \vdash \text{case } M \text{ of } \text{inl}(x) \mapsto \phi \mid \text{inr}(y) \mapsto \psi \leq \text{case } M \text{ of } \text{inl}(x) \mapsto \phi' \mid \text{inr}(y) \mapsto \psi'} \\
(\beta_{+1}\text{-eff}) \frac{\Gamma, x : A \vdash \phi \text{ eff} \quad \Gamma, y : B \vdash \psi \text{ eff} \quad \Delta \vdash M : A}{\Gamma, \Delta \vdash \text{case } \text{inl}(M) \text{ of } \text{inl}(x) \mapsto \phi \mid \text{inr}(y) \mapsto \psi \equiv [M/x]\phi} \\
(\beta_{+2}\text{-eff}) \frac{\Gamma, x : A \vdash \phi \text{ eff} \quad \Gamma, y : B \vdash \psi \text{ eff} \quad \Delta \vdash M : B}{\Gamma, \Delta \vdash \text{case } \text{inr}(M) \text{ of } \text{inl}(x) \mapsto \phi \mid \text{inr}(y) \mapsto \psi \equiv [M/x]\psi} \\
(\eta_{+}\text{-eff}) \frac{\Gamma, z : A + B \vdash \phi \text{ eff}}{\Gamma, z : A + B \vdash \phi \equiv \text{case } z \text{ of } \text{inl}(x) \mapsto [\text{inl}(x)/z]\phi \mid \text{inr}(y) \mapsto [\text{inr}(y)/z]\phi} \\
(\text{case-}\otimes) \frac{\Gamma, x : A \vdash \phi \perp \phi' \quad \Gamma, y : B \vdash \psi \perp \psi' \quad \Delta \vdash M : A + B}{\Gamma, \Delta \vdash \text{case } M \text{ of } \text{inl}(x) \mapsto (\phi \otimes \phi') \mid \text{inr}(y) \mapsto (\psi \otimes \psi')} \\
\equiv (\text{case } M \text{ of } \text{inl}(x) \mapsto \phi \mid \text{inr}(y) \mapsto \psi) \otimes (\text{case } M \text{ of } \text{inl}(x) \mapsto \phi' \mid \text{inr}(y) \mapsto \psi') \\
(\text{case-}\perp) \frac{\Gamma, x : A \vdash \phi \text{ eff} \quad \Gamma, y : B \vdash \psi \text{ eff} \quad \Delta \vdash M : A + B}{\Gamma, \Delta \vdash \text{case } M \text{ of } \text{inl}(x) \mapsto \phi^\perp \mid \text{inr}(y) \mapsto \psi^\perp \equiv (\text{case } M \text{ of } \text{inl}(x) \mapsto \phi \mid \text{inr}(y) \mapsto \psi)^\perp} \\
(\text{case-}\leq) \frac{\Gamma \vdash M : A + B \quad \Delta, x : A \vdash \phi \leq \chi \quad \Delta, y : B \vdash \psi \leq \chi}{\Gamma, \Delta \vdash (\text{case } M \text{ of } \text{inl}(x) \mapsto \phi \mid \text{inr}(y) \mapsto \psi) \leq \chi} \\
(\text{case-times}) \frac{\Gamma, x : A \vdash \phi \text{ eff} \quad \Gamma, y : B \vdash \psi \text{ eff} \quad \Delta \vdash M : A + B \quad \vdash \chi \text{ eff}}{\Gamma, \Delta \vdash (\text{case } M \text{ of } \text{inl}(x) \mapsto \chi \cdot \phi \mid \text{inr}(y) \mapsto \chi \cdot \psi) \equiv \chi \cdot \text{case } M \text{ of } \text{inl}(x) \mapsto \phi \mid \text{inr}(y) \mapsto \psi}
\end{array}$$

3.1 Metatheorems

We can prove the following properties, which show that the typing system is well behaved.

Lemma 12.

1. *Substitution* If $\Gamma \vdash M : A$ and $\Delta, x : A, \Delta' \vdash \mathcal{J}$ then $\Delta, \Gamma, \Delta' \vdash [M/x] \mathcal{J}$.
2. *Weakening* If $\Gamma \vdash M : A$ and $\Gamma \subseteq \Delta$ then $\Delta \vdash M : A$.

Proof. The proof is straightforward, by induction on derivations. □

Lemma 13 (Equation Validity).

1. If $\Gamma \vdash M = N : A$ then $\Gamma \vdash M : A$ and $\Gamma \vdash N : A$.
2. If $\Gamma \vdash \phi \leq \psi$ then $\Gamma \vdash \phi$ eff and $\Gamma \vdash \psi$ eff.

Proof. Let QPEL' be the system where the rule (\heartsuit) is replaced with

$$(\heartsuit') \frac{\Gamma \vdash \phi \perp \psi \quad \Gamma \vdash \phi \text{ eff} \quad \Gamma \vdash \psi \text{ eff}}{\Gamma \vdash \phi \heartsuit \psi \text{ eff}}$$

It is straightforward to prove that QPEL' satisfies Equation Validity. It follows that the derivable judgements of QPEL and QPEL' are the same, and hence that QPEL satisfies Equation Validity. □

Lemma 14 (Functionality).

1. If $\Gamma \vdash M = N : A$ and $\Delta, x : A \vdash P : B$ then $\Gamma, \Delta \vdash [M/x]P = [N/x]P : B$.
2. If $\Gamma \vdash M = N : A$ and $\Delta, x : A \vdash \phi$ eff then $\Gamma, \Delta \vdash [M/x]\phi \equiv [N/x]\phi$.

Proof. Let QPEL'' be the system where (measure) is replaced with the rule

$$(\text{measure}'') \frac{\Gamma \vdash 1 \leq \heartsuit_{i=1}^n \phi_i \quad \Gamma \vdash \heartsuit_{i=1}^n \phi_i \text{ eff} \quad \Delta \vdash M_i : A \quad (1 \leq i \leq n)}{\Gamma, \Delta \vdash \text{measure}_{i=1}^n \phi_i \mapsto M_i : A}$$

We can prove that QPEL'' satisfies Equation Validity, using the same proof technique as Lemma 13. It follows that QPEL and QPEL'' have the same derivable judgements. It is straightforward to prove that QPEL'' satisfies Functionality, and so it follows that QPEL satisfies Functionality. □

Lemma 15. 1. If $\Gamma, x : A, y : B \vdash M : C$, $\Delta \vdash N : A \otimes B$ and $\Theta, z : C \vdash P : D$, then

$$\Gamma, \Delta, \Theta \vdash [\text{let } x \otimes y = N \text{ in } M/z]P = (\text{let } x \otimes y = N \text{ in } [M/z]P) : D$$

2. If $\Gamma \vdash M : A + B$, $\Delta, x : A \vdash N : C$, $\Delta, y : B \vdash P : C$ and $\Theta, z : C \vdash Q : D$, then

$$\begin{aligned} \Gamma, \Delta, \Theta \vdash [\text{case } M \text{ of } \text{inl}(x) \mapsto N \mid \text{inr}(y) \mapsto P/z]Q \\ = \text{case } M \text{ of } \text{inl}(x) \mapsto [N/z]Q \mid \text{inr}(y) \mapsto [P/z]Q : D \end{aligned}$$

Proof. The proof of this lemma involves noting that local definitions can be defined from the rules for I and \otimes , which to the best of my knowledge is a new result about linear type theory.

If $\Gamma \vdash M : A$ and $\Delta, x : A \vdash N : B$, we define the term $\text{let } x = M \text{ in } N$ to be

$$\text{let } x \otimes y = M \otimes \langle \rangle \text{ in } N$$

so $\Gamma, \Delta \vdash \text{let } x = M \text{ in } N : B$ and

$$\Gamma, \Delta \vdash (\text{let } x = M \text{ in } N) = [M/x]N : B .$$

From the rules of derivation in QPEL, we can show that:

- If $\Gamma \vdash N : A \otimes B$ and $\Delta, x : A, y : B \vdash M : C$ and $\Theta, z : C \vdash P : D$ then

$$\Gamma, \Delta, \Theta \vdash (\text{let } z = \text{let } x \otimes y = N \text{ in } M \text{ in } P) = (\text{let } x \otimes y = N \text{ in let } z = M \text{ in } P) : D$$

- If $\Gamma \vdash M : A + B$, $\Delta, x : A \vdash N : C$, $\Delta, y : B \vdash P : C$, and $\Theta, z : C \vdash Q : D$, then

$$\begin{aligned} \Gamma, \Delta, \Theta \vdash \text{let } z = (\text{case } M \text{ of } \text{inl}(x) \mapsto N \mid \text{inr}(y) \mapsto P) \text{ in } Q \\ = \text{case } M \text{ of } \text{inl}(x) \mapsto \text{let } z = N \text{ in } Q \mid \text{inr}(y) \mapsto \text{let } z = P \text{ in } Q : D \end{aligned}$$

The result then follows. □

4 Semantics

4.1 State and Effect Triangles

Let E be a commutative effect monoid. Recall the adjunction $\mathbf{Conv}_E[-, E] \dashv \mathbf{EMod}_E[-, E] : \mathbf{Conv}_E \rightleftarrows \mathbf{EMod}_E^{\text{op}}$.

Definition 16 (State-and-Effect Triangle). A *state-and-effect triangle* consists of:

- a symmetric monoidal category \mathcal{V} with binary coproducts that distribute over the tensor, such that the tensor unit is terminal;
- an effect monoid E ;
- a functor $P : \mathcal{V} \rightarrow \mathbf{EMod}_E^{\text{op}}$ that preserves finite coproducts and the terminal object;
- a symmetric monoidal functor $S : \mathcal{V} \rightarrow \mathbf{Conv}_E$;
- given a finite set $r_1, \dots, r_n \in PA$ such that $r_1 \otimes \dots \otimes r_n = 1$, an arrow $\text{meas}_A(r_1, \dots, r_n) : A \rightarrow n \cdot I$ in \mathcal{V} ;
- a natural transformation $\alpha : P \rightarrow \mathbf{Conv}_E[S-, E]$;
- a natural transformation $\beta : S \rightarrow \mathbf{EMod}_E[P-, E]$;

such that

1. given a permutation p on $\{1, \dots, n\}$, we have

$$\text{meas}_A(r_{p(1)}, \dots, r_{p(n)}) = \pi_p \circ \text{meas}_A(p_1, \dots, p_n)$$

where $\pi_p : n \cdot I \rightarrow n \cdot I$ satisfies

$$\pi_p \circ \kappa_i = \kappa_{p(i)}$$


2. $\text{meas}_A(p_1, \dots, p_n, 0) = \kappa_1 \circ \text{meas}_A(p_1, \dots, p_n) : A \rightarrow n \cdot I \rightarrow (n+1) \cdot I$
3. $\text{meas}_A(p \otimes q, r_1, \dots, r_n) = [\kappa_1, \kappa_1, \kappa_2, \dots, \kappa_{n+1}] \circ \text{meas}_A(p, q, r_1, \dots, r_n)$
4. meas_A is natural in A ; i.e. given $f : A \rightarrow B$,

$$\text{meas}_A(r_1, \dots, r_n) \circ f = \text{meas}_B(Pf(r_1), \dots, Pf(r_n))$$

5. $\alpha_A(p)(x) = \beta_A(x)(p)$ for all A, x, p .

We think of the arrows in \mathcal{V} as *computations*, the arrows $SA \rightarrow SB$ as *state transformers*, and the arrows $PA \rightarrow PB$ as *predicate transformers*.

We refer to α and β as the *validity transformations*, since the intuition is that $\alpha_A(p)(x) = \beta_A(x)(p)$ is the probability of the statement 'Predicate p is valid at state x '.



Examples The following are all examples of state-and-effect triangles:

- Take \mathcal{V} to be the category $\mathbf{FdHilb}_{\text{Un}}$ of finite-dimensional Hilbert spaces with unitary maps, PH to be the set of *effects* on H (positive operators less than I), and SH to be the set of *density matrices* on H .
- Take \mathcal{V} to be $\mathbf{Kl}(\mathcal{D}_E)$, the Kleisli category of the distribution monad \mathcal{D}_E . S is the canonical functor from the Kleisli category to the Eilenberg-Moore category. For $X \in \mathbf{Set}$, PX is the set of all functions $X \rightarrow \mathcal{D}_E(2)$, equivalently the set of functions $X \rightarrow E$.
- Take \mathcal{V} to be $\mathbf{CStar}_{\text{PU}}^{\text{op}}$, the category of C^* -algebras and positive unital maps. PA is the set of all effects on a , $[0, 1]_A = \{a \in A : 0 \leq a \leq 1\}$. SA is the set of all positive unital maps $A \rightarrow \mathbb{C}$.
- Take \mathcal{V} to be \mathbf{Set} , PA the power set of A , and $SA = A$. The effect monoid in this case is $\{0, 1\}$.

- More generally, let $(\mathcal{V}, \otimes, I)$ be any symmetric monoidal category with finite coproducts $(0, +)$ such that:
 - diagrams of the following form are always pullbacks in \mathcal{V} :

$$\begin{array}{ccc} A + X & \xrightarrow{1+f} & A + Y \\ \downarrow g+1 & & \downarrow g+1 \\ B + X & \xrightarrow{1+f} & B + Y \end{array} \qquad \begin{array}{ccc} A & \xrightarrow{\kappa_1} & A \\ \downarrow & & \downarrow \kappa_2 \\ A + X & \xrightarrow{1+f} & A + Y \end{array}$$

- for each non-zero $n \in \mathbb{N}$, the family of maps

$$[\triangleright_i, \kappa_2] : n \cdot X + 1 \rightarrow X + 1$$

are jointly monic where where, for $1 \leq i \leq n$, the ‘partial projection’ $\triangleright_i : n \cdot X \rightarrow X + 1$ is such that

$$\triangleright_i \circ \kappa_j = \begin{cases} \kappa_1 & \text{if } i = j \\ \kappa_2 \circ ! & \text{if } i \neq j \end{cases}$$

Take $E = \mathcal{V}[1, 2]$. Define P to be the functor $\mathcal{V}[-, 2]$, and S to be the functor $\mathcal{V}[-, 1]$. Define α and β by

$$\alpha(p)(\omega) = \beta(\omega)(p) = p \circ \omega .$$

for $p : A \rightarrow 2$ and $\omega : 1 \rightarrow A$.

The arrow $\text{meas}_A(r_1, \dots, r_n)$ is the unique arrow such that $\triangleright_i \circ \text{meas}_A(r_1, \dots, r_n) = r_i$.

See [13] for a verification that these constructions are all well-defined and satisfy the axioms of a state-and-effect triangle. The previous examples are all special cases of this construction.

Remarks

1. We do not want S always to be a strong monoidal functor. Intuitively, $S(A) \otimes S(B)$ gives the mixtures of pure states of $A \otimes B$, while $S(A \otimes B)$ also includes entangled states, and these will not be isomorphic in general.
2. The condition $\alpha_A(p)(x) = \beta_A(x)(p)$ can also be written as $\alpha = G\beta \circ \eta P$ or as $\beta = F\alpha \circ \varepsilon S$, where $F = \mathbf{EMod}_E[-, E] : \mathbf{EMod}_E^{\text{op}} \rightarrow \mathbf{Conv}_E$ and $G = \mathbf{Conv}_E[-, E] : \mathbf{Conv}_E \rightarrow \mathbf{EMod}_E^{\text{op}}$.

4.2 Semantics

Definition 17. Given any state-and-effect triangle, we interpret the syntax as follows.

- We associate with every type A an object $\llbracket A \rrbracket$ of \mathcal{V} thus:

$$\begin{aligned} \llbracket I \rrbracket &= I \\ \llbracket A \otimes B \rrbracket &= \llbracket A \rrbracket \otimes \llbracket B \rrbracket \\ \llbracket A + B \rrbracket &= \llbracket A \rrbracket + \llbracket B \rrbracket \end{aligned}$$

- We associate with every context Γ an object $\llbracket \Gamma \rrbracket$ of \mathcal{V} as follows.

$$\begin{aligned} \llbracket \langle \rangle \rrbracket &= I \\ \llbracket \Gamma, x : A \rrbracket &= \llbracket \Gamma \rrbracket \otimes \llbracket A \rrbracket \end{aligned}$$

- We associate with every term $\Gamma \vdash M : A$ an arrow $\llbracket M \rrbracket = \llbracket \Gamma \vdash M : A \rrbracket : \llbracket \Gamma \rrbracket \rightarrow \llbracket A \rrbracket$ in \mathcal{V} as follows.
 - $\llbracket x_1 : A_1, \dots, x_n : A_n \vdash x_i : A_i \rrbracket$ is the arrow

$$\llbracket A_1 \rrbracket \otimes \dots \otimes \llbracket A_n \rrbracket \xrightarrow{! \otimes \dots \otimes ! \otimes 1 \otimes ! \otimes \dots \otimes !} I \otimes \dots \otimes I \otimes \llbracket A_i \rrbracket \otimes I \otimes \dots \otimes I \xrightarrow{\cong} \llbracket A_i \rrbracket$$

- $\llbracket M \otimes N \rrbracket = \llbracket M \rrbracket \otimes \llbracket N \rrbracket$
- $\llbracket \Gamma, \Delta \vdash \text{let } x \otimes y = M \text{ in } N : C \rrbracket$ is

$$\llbracket \Gamma \rrbracket \otimes \llbracket \Delta \rrbracket \xrightarrow{\llbracket M \rrbracket \otimes 1} \llbracket A \rrbracket \otimes \llbracket B \rrbracket \otimes \llbracket \Delta \rrbracket \xrightarrow{\cong} \llbracket \Delta \rrbracket \otimes \llbracket A \rrbracket \otimes \llbracket B \rrbracket \xrightarrow{\llbracket N \rrbracket} \llbracket C \rrbracket$$

- $\llbracket \Gamma \vdash \langle \rangle : I \rrbracket = ! : \llbracket \Gamma \rrbracket \rightarrow I$
- $\llbracket \text{inl}(M) \rrbracket = \kappa_1 \circ \llbracket M \rrbracket$
- $\llbracket \text{inr}(M) \rrbracket = \kappa_2 \circ \llbracket M \rrbracket$
- $\llbracket \Gamma, \Delta \vdash \text{case } M \text{ of } \text{inl}(x) \mapsto N \mid \text{inr}(y) \mapsto P : C \rrbracket$ is the arrow

$$\llbracket \Gamma \rrbracket \otimes \llbracket \Delta \rrbracket \xrightarrow{\llbracket M \rrbracket \otimes 1} (\llbracket A \rrbracket + \llbracket B \rrbracket) \otimes \llbracket \Delta \rrbracket \xrightarrow{\cong} (\llbracket \Delta \rrbracket \otimes A) + (\llbracket \Delta \rrbracket \otimes B) \xrightarrow{\llbracket N \rrbracket, \llbracket P \rrbracket} \llbracket C \rrbracket$$

- $\llbracket \Gamma, \Delta \vdash \text{measure } \phi_1 \mapsto M_1 \mid \dots \mid \phi_n \mapsto M_n : A \rrbracket$ is the arrow

$$\llbracket \Gamma \rrbracket \otimes \llbracket \Delta \rrbracket \xrightarrow{\text{meas}_{\llbracket \Gamma \rrbracket}(\llbracket \phi_1 \rrbracket, \dots, \llbracket \phi_n \rrbracket) \otimes 1} n \cdot I \otimes \llbracket \Delta \rrbracket \xrightarrow{\cong} n \cdot \llbracket \Delta \rrbracket \xrightarrow{\llbracket M_1 \rrbracket, \dots, \llbracket M_n \rrbracket} \llbracket A \rrbracket$$

- We associate with every proposition ϕ such that $\Gamma \vdash \phi$ eff, an element $\llbracket \phi \rrbracket \in P[\llbracket \Gamma \rrbracket]$ as follows.

$$\begin{aligned} \llbracket 0 \rrbracket &= 0 \\ \llbracket \phi^\perp \rrbracket &= \llbracket \phi \rrbracket^\perp \\ \llbracket \phi \otimes \psi \rrbracket &= \llbracket \phi \rrbracket \otimes \llbracket \psi \rrbracket \\ \llbracket \phi \cdot \psi \rrbracket &= \llbracket \phi \rrbracket \cdot \llbracket \psi \rrbracket \end{aligned}$$

In this last line, if $\Gamma \vdash \phi \cdot \psi$ eff then $\llbracket \phi \rrbracket \in PI$ and $\llbracket \psi \rrbracket \in P[\llbracket \Gamma \rrbracket]$. We use the fact that $E \cong PI$ (since P preserves the terminal object), so we may take $\llbracket \phi \rrbracket$ to be an element of E .

$\llbracket \Gamma, \Delta \vdash \text{case } M \text{ of } \text{inl}(x) \mapsto \phi \mid \text{inr}(y) \mapsto \psi \rrbracket$ is defined as follows. We have

$$1 \otimes \llbracket M \rrbracket : \llbracket \Gamma \rrbracket \otimes \llbracket \Delta \rrbracket \rightarrow \llbracket \Gamma \rrbracket \otimes (\llbracket A \rrbracket + \llbracket B \rrbracket) = (\llbracket \Gamma \rrbracket \otimes \llbracket A \rrbracket) + (\llbracket \Gamma \rrbracket \otimes \llbracket B \rrbracket)$$

and so

$$P(1 \otimes \llbracket M \rrbracket) : P(\llbracket \Gamma \rrbracket \otimes \llbracket A \rrbracket) \times P(\llbracket \Gamma \rrbracket \otimes \llbracket B \rrbracket) \rightarrow P(\llbracket \Gamma \rrbracket \otimes \llbracket \Delta \rrbracket)$$

(Recall that $P : \mathcal{V} \rightarrow \mathbf{EMod}_E^{\text{op}}$ preserves binary coproducts, and so $P(A + B)$ is the product of PA and PB in \mathbf{EMod}_E .)

We define $\llbracket \text{case } M \text{ of } \text{inl}(x) \mapsto \phi \mid \text{inr}(y) \mapsto \psi \rrbracket$ to be

$$P(1 \otimes \llbracket M \rrbracket)(\llbracket \phi \rrbracket, \llbracket \psi \rrbracket) .$$

Lemma 18. 1. If $\Gamma, x : A \vdash M : B$ and $\Delta \vdash N : A$, then $\llbracket \Gamma, \Delta \vdash [N/x]M : B \rrbracket$ is the arrow

$$\llbracket \Gamma \rrbracket \otimes \llbracket \Delta \rrbracket \xrightarrow{1 \otimes \llbracket N \rrbracket} \llbracket \Gamma \rrbracket \otimes \llbracket A \rrbracket \xrightarrow{\llbracket M \rrbracket} \llbracket B \rrbracket$$

2. If $\Gamma, x : A \vdash \phi \text{ eff}$ and $\Delta \vdash M : A$ then

$$\llbracket [M/x]\phi \rrbracket = P(1_{\llbracket \Gamma \rrbracket} \otimes \llbracket M \rrbracket)(\llbracket \phi \rrbracket)$$

Proof. The two parts are proved simultaneously, by induction on M and ϕ . All cases are straightforward. \square

Definition 19. In a state-and-effect triangle, a judgement $\Gamma \vdash M = N : A$ is *true* iff $\llbracket \Gamma \vdash M : A \rrbracket = \llbracket \Gamma \vdash N : A \rrbracket$. A judgement $\Gamma \vdash \phi \leq \psi$ is *true* iff $\llbracket \Gamma \vdash \phi \text{ eff} \rrbracket \leq \llbracket \Gamma \vdash \psi \text{ eff} \rrbracket$, in the order in the effect module $P\llbracket \Gamma \rrbracket$.

Theorem 20 (Soundness). *Any derivable judgement is true in any state-and-effect triangle.*

Proof. Straightforward induction on derivations. \square

Theorem 21 (Completeness). *Any judgement that is true in every state-and-effect triangle is derivable.*

Proof. Define a state-and-effect triangle as follows.

The category \mathcal{V} is the category with objects the types of QPEL, and arrows $A \rightarrow B$ the pairs (x, M) such that $x : A \vdash M : B$, quotiented by:

- $(x : A \vdash M : B) = (y : A \vdash [y/x]M : B)$ if $x \neq y$ and y does not occur in M ;
- If $x : A \vdash M = N : B$ is derivable, then $(x : A \vdash M : B) = (x : A \vdash N : B)$.

The identity on A is $x : A \vdash x : A$. The composite of $x : A \vdash M : B$ and $y : B \vdash N : C$ is $x : A \vdash [M/y]N : C$. This is well-defined by Substitution and Functionality.

We shall write an arrow $x : A \vdash M : B$ as $M[x] : A \rightarrow B$, and then write $M[N]$ for the term $[N/x]M$.

Tensor Product For types A and B , the tensor product is $A \otimes B$.

Given arrows $M[a] : A \rightarrow A'$ and $N[b] : B \rightarrow B'$, define $M \otimes N : A \otimes B \rightarrow A' \otimes B'$ by

$$(M \otimes N)[z] = \text{let } a \otimes b = z \text{ in } M[a] \otimes N[b] .$$

Coproducts For types A and B , the coproduct is $A + B$, with injections

$$x : A \vdash \text{inl}(x) : A + B, \quad y : B \vdash \text{inr}(y) : A + B .$$

Given $M[a] : A \rightarrow C$ and $N[b] : B \rightarrow C$, the mediating arrow $[M, N] : A + B \rightarrow C$ is defined by

$$[M, N][x] = \text{case } x \text{ of } \text{inl}(a) \mapsto M[a] \mid \text{inr}(b) \mapsto N[b] .$$

Effect Monoid The effect monoid E is the set of all propositions ϕ such that $\vdash \phi$ eff, quotiented by: $\phi = \psi$ iff $\vdash \phi \leq \psi$ and $\vdash \psi \leq \phi$.

We have that $\phi \otimes \psi$ is defined iff $\vdash \phi \otimes \psi$ eff (equivalently, iff $\vdash \phi \leq \psi^\perp$), in which case the partial sum is $\phi \otimes \psi$. The zero element is 0, and the orthocomplement of ϕ is ϕ^\perp . The product of ϕ and ψ is $\phi \cdot \psi$.

Predicate Functor The functor P is defined by: PA is the set of all pairs (x, ϕ) such that $x : A \vdash \phi$ eff, quotiented by:

- $(x, \phi) = (y, [y/x]\phi)$ if $x \neq y$ and y does not occur in ϕ ;
- $(x, \phi) = (x, \psi)$ if $x : A \vdash \phi \equiv \psi$.

This is an effect module under $0, \perp, \otimes, \cdot$.

Given $M[a] : A \rightarrow B$, then $PM : PB \rightarrow PA$ is defined by

$$PM(b, \phi) \equiv (a, [M[a]/b]\phi) .$$

State Functor The functor S is defined by: SA is the set of all terms M such that $\vdash M : A$, quotiented by: $M = N$ iff $\vdash M = N : A$.

We make this into a convex set by setting

$$\phi_1 M_1 + \cdots + \phi_n M_n = \text{measure } \phi_1 \mapsto M_1 \mid \cdots \mid \phi_n \mapsto M_n .$$

Given $M[a] : A \rightarrow B$, we define $SM : SA \rightarrow SB$ by

$$SM(N) \equiv M[N] .$$

We make S into a symmetric monoidal functor by setting

$$\begin{aligned} \phi_{AB} &: SA \otimes SB \rightarrow S(A \otimes B) \\ \phi_{AB}(M, N) &= M \otimes N \\ \phi &: \{*\} \rightarrow SI \\ \phi(*) &= \langle \rangle \end{aligned}$$

Measurement Morphisms We have $\text{meas}_A(\phi_1, \dots, \phi_n) = \text{measure } \phi_1 \mapsto \text{in}_1(\langle \rangle) \mid \cdots \mid \phi_n \mapsto \text{in}_n(\langle \rangle)$, where the terms $\text{in}_i(M)$ are the n canonical terms such that $x : A \vdash \text{in}_i(x) : \overbrace{A + \cdots + A}^n$.

Validity Transformations The transformation α is given by $\alpha_A(x : A \vdash \phi \text{ eff})(\vdash M : A) \equiv (\vdash [M/x]\phi \text{ eff})$, and so β is given by $\beta_A(\vdash M : A)(x : A \vdash \phi \text{ eff}) \equiv (\vdash [M/x]\phi \text{ eff})$.

Proof of Completeness We will prove that, if a judgement is true in this triangle, then it is derivable.

Let $\Gamma \equiv x_1 : A_1, \dots, x_n : A_n$. Then a straightforward induction shows that:

$$\begin{aligned} [[\Gamma \vdash M : B]] &= z : A_1 \otimes \cdots \otimes A_n \vdash \text{let } x_1 \otimes \cdots \otimes x_n = z \text{ in } M : B \\ [[\Gamma \vdash \phi \text{ eff}]] &= z : A_1 \otimes \cdots \otimes A_n \vdash \text{let } x_1 \otimes \cdots \otimes x_n = z \text{ in } \phi \text{ eff} \end{aligned}$$

where this last effect is defined inductively thus:

$$\begin{aligned}
& \text{let } x_1 \otimes \cdots \otimes x_n = z \text{ in } 0 \equiv 0 \\
& \text{let } x_1 \otimes \cdots \otimes x_n = z \text{ in } \phi^\perp \equiv (\text{let } x_1 \otimes \cdots \otimes x_n = z \text{ in } \phi)^\perp \\
& \text{let } x_1 \otimes \cdots \otimes x_n = z \text{ in } \phi \otimes \psi \equiv (\text{let } x_1 \otimes \cdots \otimes x_n = z \text{ in } \phi) \otimes (\text{let } x_1 \otimes \cdots \otimes x_n = z \text{ in } \psi) \\
& \text{case } M \text{ of inl}(x) \mapsto \phi \mid \text{inr}(y) \mapsto \psi \text{ in } \text{let } x_1 \otimes \cdots \otimes x_n = z \text{ in } M \equiv \begin{cases} \text{case } (\text{let } x_1 \otimes \cdots \otimes x_n = z \text{ in } M) \text{ of} \\ \quad \text{inl}(x) \mapsto \phi \mid \text{inr}(y) \mapsto \psi \\ \quad \text{if } z \text{ occurs in } M \\ \text{case } M \text{ of inl}(x) \mapsto \text{let } x_1 \otimes \cdots \otimes x_n = z \text{ in } \phi \mid \\ \quad \text{inr}(y) \mapsto \text{let } x_1 \otimes \cdots \otimes x_n = z \text{ in } \psi \\ \text{otherwise} \end{cases}
\end{aligned}$$

Suppose that the judgement $\Gamma \vdash M = N : A$ is true in this triangle. Then we have

$$z : A_1 \otimes \cdots \otimes A_n \vdash (\text{let } x_1 \otimes \cdots \otimes x_n = z \text{ in } M) = (\text{let } x_1 \otimes \cdots \otimes x_n = z \text{ in } N) : A$$

is derivable. By Substitution, we have

$$\Gamma \vdash (\text{let } x_1 \otimes \cdots \otimes x_n = x_1 \otimes \cdots \otimes x_n \text{ in } M) = (\text{let } x_1 \otimes \cdots \otimes x_n = x_1 \otimes \cdots \otimes x_n \text{ in } N) : A$$

is derivable, and hence $\Gamma \vdash M = N : A$ is derivable by $(\eta \otimes)$.

Suppose that $\Gamma \vdash \phi \leq \psi$ is true in this triangle. Then

$$z : A_1 \otimes \cdots \otimes A_n \vdash (\text{let } x_1 \otimes \cdots \otimes x_n = z \text{ in } \phi) \leq (\text{let } x_1 \otimes \cdots \otimes x_n = z \text{ in } \psi)$$

is derivable. By Substitution, we have

$$\Gamma \vdash (\text{let } x_1 \otimes \cdots \otimes x_n = x_1 \otimes \cdots \otimes x_n \text{ in } \phi) \leq (\text{let } x_1 \otimes \cdots \otimes x_n = x_1 \otimes \cdots \otimes x_n \text{ in } \psi) .$$

It is easy to show, by induction on ϕ , that

$$\Gamma \vdash ((\text{let } x_1 \otimes \cdots \otimes x_n = x_1 \otimes \cdots \otimes x_n \text{ in } \phi) \equiv \phi) .$$

It follows that $\Gamma \vdash \phi \leq \psi$ is derivable. □

5 Qubits

There are several ways in which the system may be extended to represent qubits. The details below are based on the Measurement Calculus [8].

We extend the system with:

$$\begin{array}{lll}
\text{Type} & A & ::= \cdots \mid \mathbf{qbit} \\
\text{Term} & M & ::= \cdots \mid |+\rangle \mid XM \mid ZM \mid EMM \\
\text{Effect} & \phi & ::= \cdots \mid M = |+\alpha\rangle
\end{array}$$

where α is a real number in $[0, 2\pi)$.

The intention is that a term of type **qbit** represents a qubit. The term $|+\rangle$ represents a qubit in the phase

$$|+\rangle = \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle) .$$

The terms XM and ZM denote the result of applying the Pauli-X and Z gates to the qubit M . The term EMN denotes the result of applying the controlled Z gate to the pair of qubits M and N . The effect $M = |+\alpha\rangle$ denotes the projector on

$$|+\alpha\rangle = \frac{1}{\sqrt{2}}(|0\rangle + e^{i\alpha}|1\rangle)$$

Its orthocomplement, $|+\alpha\rangle^\perp$, is the projector on

$$|-\alpha\rangle = \frac{1}{\sqrt{2}}(|0\rangle - e^{i\alpha}|1\rangle)$$

We write $|-\rangle$ for $Z|+\rangle$

We extend the system with the following rules of deduction.

$$\begin{array}{c} \frac{}{\vdash \text{new } |+\rangle : \mathbf{qbit}} \quad \frac{\Gamma \vdash M : \mathbf{qbit}}{\Gamma \vdash XM : \mathbf{qbit}} \quad \frac{\Gamma \vdash M : \mathbf{qbit}}{\Gamma \vdash ZM : \mathbf{qbit}} \\ \frac{\Gamma \vdash M : \mathbf{qbit} \quad \Gamma \vdash N : \mathbf{qbit}}{\Gamma \vdash EMN : \mathbf{qbit} \otimes \mathbf{qbit}} \quad \frac{\Gamma \vdash M : \mathbf{qbit}}{\Gamma \vdash M = |+\alpha\rangle \text{ eff}} \quad (0 \leq \alpha < 2\pi) \\ \frac{\Gamma \vdash M : \mathbf{qbit} \quad \Gamma \vdash N : \mathbf{qbit}}{\Gamma \vdash E(XM)N = \text{let } x \otimes y = EMN \text{ in } Xx \otimes Zy : \mathbf{qbit} \otimes \mathbf{qbit}} \\ \frac{\Gamma \vdash M : \mathbf{qbit} \quad \Gamma \vdash N : \mathbf{qbit}}{\Gamma \vdash E(ZM)N = \text{let } x \otimes y = EMN \text{ in } Zx \otimes y : \mathbf{qbit} \otimes \mathbf{qbit}} \\ \frac{\Gamma \vdash M : \mathbf{qbit}}{\Gamma \vdash (XM = |+\alpha\rangle) \equiv (M = |+\alpha\rangle)} \quad (0 \leq \alpha < 2\pi) \\ \frac{\Gamma \vdash M : \mathbf{qbit}}{\Gamma \vdash (ZM = |+\alpha\rangle) \equiv (M = |+\alpha-\pi\rangle)} \quad (0 \leq \alpha < 2\pi) \\ \frac{\Gamma \vdash M : \mathbf{qbit}}{\Gamma \vdash X(XM) = M : \mathbf{qbit}} \quad \frac{\Gamma \vdash M : \mathbf{qbit}}{\Gamma \vdash Z(ZM) = M : \mathbf{qbit}} \\ \frac{\Gamma \vdash M : \mathbf{qbit}}{(X(ZM) = |+\alpha\rangle) \equiv (Z(XM) = |+\alpha\rangle)} \quad (0 \leq \alpha < 2\pi) \end{array}$$

The metatheorems in Section 3.1 all still hold for the expanded system. The expanded system can be given semantics in $\mathbf{CStar}_{\text{PU}}^{\text{op}}$ straightforwardly. We will show in a forthcoming paper how these rules are sufficient to prove the correctness of several quantum algorithms, including superdense coding and gate-based teleportation.

6 Natural Isomorphisms

It is interesting to consider the question of when the natural transformations α and β are isomorphisms. In the $\mathbf{FdHilb}_{\text{Un}}$ example, α and β are both isomorphisms. [14]. In the $\text{Kl}(\mathcal{D})$ example, α is an isomorphism but β is not. In the $\mathbf{CStar}_{\text{PU}}^{\text{op}}$ example, β is an isomorphism but α is not.

We can extend the system so it captures the state-and-effect triangles in which β is an isomorphism as follows.

Theorem 22 (Completeness). *Add to the system the rule*

$$\frac{\vdash \phi \text{ eff} \quad \Gamma \vdash M : A \quad \Gamma \vdash N : A \quad \Delta, x : A \vdash \psi \text{ eff}}{\Gamma, \Delta \vdash [(\text{measure } \phi \mapsto M | \phi^\perp \mapsto N) / x] \psi \equiv (\phi \cdot [M/x] \psi) \odot (\phi^\perp \cdot [N/x] \psi)}$$

If a judgement is true in every state-and-effect triangle in which α and β are natural isomorphisms, then it is derivable in this system.

I do not yet have a system that captures the state-and-effect triangles in which α is a natural isomorphism.

The case where α is an isomorphism is particularly interesting, as it is this that allows *weakest preconditions* in d’Hondt-Panangaden’s sense to be defined.

Definition 23. Let P and Q be quantum predicates, and F a quantum program. Then P is a *precondition* for Q with respect to M , PFQ , iff for all density matrices ρ , $\text{tr}(P\rho) \leq \text{tr}(QF(\rho))$. P is the *weakest precondition* for Q with respect to M , $P = wp(F)(Q)$ iff P is the greatest precondition for Q w.r.t. M under the Löwner order.

The weakest precondition for Q w.r.t. F always exists and is unique [9].

Lemma 24. *In the $\mathbf{FdHilb}_{\text{Un}}$ state-and-effect triangle, the weakest precondition for $Q \in PH$ with respect to $F : SK \rightarrow SH$ is $\alpha^{-1}(F \circ \alpha(P))$. The operation $wp(F)$ is therefore the effect module homomorphism $\alpha^{-1} \circ \mathbf{Conv}_M[1, F] \circ \alpha : PH \rightarrow PK$. The operation wp is therefore the natural transformation*

$$wp_{PHK} = \alpha^{-1} \circ \mathbf{Conv}_M[1, -] \circ \alpha : \mathbf{Conv}_M[SK, SH] \rightarrow \mathbf{EMod}_M[PH, PK]$$

Lemma 25. *Given $\Gamma \vdash M : A$ and $x : A \vdash \phi \text{ eff}$, then in the $\mathbf{FdHilb}_{\text{Un}}$ semantics:*

$$wp(\llbracket \Gamma \vdash M : A \rrbracket)(\llbracket x : A \vdash \phi \text{ eff} \rrbracket) = \llbracket \Gamma \vdash [M/x] \phi \text{ eff} \rrbracket$$

7 Conclusion, Related Work and Future Work

We have presented QPEL, a syntactic system involving both terms and propositions that captures the categorical notion of ‘state-and-effect triangle’ which has proved to be a general setting for describing both quantum programs, and effects. It is therefore a promising candidate for a language that allows us to reason about and prove properties of quantum programs, and shows how such a logic for quantum effects might be added on top of any quantum programming language.

Baltag and Smets in a series of papers [2, 5, 4, 3, 1] describe the language QDL, Quantum Dynamic Logic. This is also a language for describing quantum programs and properties of quantum programs. Their work differs from mine because their term language is an underspecification language (as is Dynamic Logic’s), and their propositions can denote all propositions expressible in classical logic, not just those that correspond to quantum effects.

d’Hondt-Panangaden [9] and Ying [18] have investigated the notion of a *quantum predicate*. Ying has given a Floyd-Hoare style logic which, given a program F written in his syntax, allows the weakest precondition of a predicate with respect to F to be calculated. Their work differs from mine because they do not give a syntax for the predicates, instead using the effects on a Hilbert space as the predicates directly.

In the future, the most important tasks are to apply the system to prove the correctness of a simple quantum program (e.g. the quantum teleportation protocol or quantum broadcasting), and to look for ways to extend the system in order to represent looping and/or recursion.

I will present the system in a more modular fashion, giving subsystems that can be interpreted in other state-and-effect triangles, for example using complete lattices instead of effect modules. This may lead to a general notion of a (2-)category of state-and-effect triangles.

I will also try to capture the conditions that make α or β a natural isomorphism. I will investigate the conditions that a state-and-effect triangle needs to satisfy to represent the type of qubits correctly, possibly involving Selinger's notion of a Quantum Flowchart Category. I will investigate formal translations between this system and other quantum programming languages, such as the quantum lambda calculus [17]. I will investigate which of Ying's equations on weakest preconditions [18] can be derived within our system.

Acknowledgements Thanks to Sam Staton and Bart Jacobs for many helpful discussions.

References

- [1] Alexandru Baltag & Sonja Smets (2004): *The Logic of Quantum Programs*. In: *QPL 2004*, pp. 39–56.
- [2] Alexandru Baltag & Sonja Smets (2005): *Complete Axiomatizations for Quantum Actions*. *International Journal of Theoretical Physics* 44, doi:10.1007/s10773-005-8022-2.
- [3] Alexandru Baltag & Sonja Smets (2005): *LQP: The Dynamic Logic of Quantum Information*. *Mathematical Structures in Computer Science*, doi:10.1017/s0960129506005299.
- [4] Alexandru Baltag & Sonja Smets (2011): *Quantum Logic as a Dynamic Logic*. *Synthese* 179, pp. 285–306, doi:10.1007/s11229-010-9783-6.
- [5] Alexandru Baltag & Sonja Smets (2012): *The Dynamic Turn in Quantum Logic*. *Synthese* 186, pp. 753–773, doi:10.1007/s11229-011-9915-7.
- [6] Nick Benton, Gavin Bierman, Valeria De Paiva & Martin Hyland (1993): *A Term Calculus for Intuitionistic Linear Logic*. In: *TLCA, Lecture Notes in Computer Science* 664, Springer-Verlag, pp. 75–90, doi:10.1007/BFb0037099.
- [7] P. N. Benton (1995): *A Mixed Linear and Non-Linear Logic: Proofs, Terms and Models (Extended Abstract)*. In: *Selected Papers from the 8th International Workshop on Computer Science Logic, CSL '94*, Springer-Verlag, London, UK, UK, pp. 121–135, doi:10.1007/bfb0022251.
- [8] Vincent Danos, Elham Kashefi, Prakash Panangaden & Simon Perdrix (2009): *Extended Measurement Calculus*. In: *Semantic Techniques for Quantum Computation*, chapter 5, Cambridge University Press, doi:10.1017/CBO9781139193313.008.
- [9] Ellie d'Hondt & Prakash Panangaden (2006): *Quantum Weakest Preconditions*. *Math Struct in Comp Science* 16, pp. 429–451, doi:10.1017/S0960129506005251.
- [10] A. Dvurečenskij & S. Pulmannov á (2000): *New Trends in Quantum Structures*. Kluwer Academic Publishers, Dordrecht, doi:10.1007/978-94-017-2422-7.
- [11] Bart Jacobs (2010): *Convexity, duality, and effects*. In C. S. Clade & V. Sassone, editors: *IFIP Theoretical Computer Science 2010, IFIP Adv. in Inf. and Comm. Techn.* 82 (1), Springer, Boston, pp. 1–19, doi:10.1007/978-3-642-15240-5_1.
- [12] Bart Jacobs (2011): *Probabilities, Distribution Monads, and Convex Categories*. *Theor. Comput. Sci.* 412(28), pp. 3323–3336, doi:10.1016/j.tcs.2011.04.005.
- [13] Bart Jacobs (2014): *New Directions in Categorical Logic, for Classical, Probabilistic and Quantum Logic*. To appear in *Logical Methods in Computer Science*. <http://arxiv.org/abs/1205.3940>
- [14] Bart Jacobs & Jorik Mandemaker (2013): *Relating Operator Spaces via Adjunctions*. In J. Chubb Reimann, V. Harizanov & A. Eskandarian, editors: *Logic and Algebraic Structures in Quantum Computing and Information*, Lect. Notes in Logic, Camb.

- [15] Anders Kock (1972): *Strong functors and monoidal monads*. *Archiv der Mathematik* 23(1), pp. 113–120, doi:10.1007/BF01304852.
- [16] Peter Selinger (2004): *Towards a Quantum Programming Language*. *Math Struct in Comp Science* 14(4), pp. 527–586, doi:10.1017/S0960129504004256.
- [17] P. Selinger & B. Valiron (2010): *Quantum Lambda Calculus*. In S. Gay & I Mackie, editors: *Semantical Techniques in Quantum Computation*, Cambridge University Press, pp. 135–172, doi:10.1017/cbo9781139193313.005.
- [18] Mingsheng Ying (2011): *Floyd-Hoare logic for quantum programs*. *ACM Trans. Program. Lang. Syst.* 33(6), p. 19, doi:10.1145/2049706.2049708.