

Benchmarking semiclassical approaches to strong-field QED: Nonlinear Compton scattering in intense laser pulses

Downloaded from: https://research.chalmers.se, 2024-05-01 21:47 UTC

Citation for the original published paper (version of record): Blackburn, T., Seipt, D., Bulanov, S. et al (2018). Benchmarking semiclassical approaches to strong-field QED: Nonlinear Compton scattering in intense laser pulses. Physics of Plasmas, 25(8). http://dx.doi.org/10.1063/1.5037967

N.B. When citing this work, cite the original published paper.

research.chalmers.se offers the possibility of retrieving research publications produced at Chalmers University of Technology. It covers all kind of research output: articles, dissertations, conference papers, reports etc. since 2004. research.chalmers.se is administrated and maintained by Chalmers Library



Benchmarking semiclassical approaches to strong-field QED: Nonlinear Compton scattering in intense laser pulses

T. G. Blackburn,^{1,a)} D. Seipt,^{2,3} S. S. Bulanov,⁴ and M. Marklund¹ ¹Department of Physics, Chalmers University of Technology, SE-41296 Gothenburg, Sweden ²Physics Department, Lancaster University, Lancaster LA1 4YB, United Kingdom ³The Cockcroft Institute, Daresbury Laboratory, Warrington WA4 4AD, United Kingdom ⁴Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA

(Received 30 April 2018; accepted 20 July 2018; published online 8 August 2018)

The recoil associated with photon emission is key to the dynamics of ultrarelativistic electrons in strong electromagnetic fields, as found in high-intensity laser-matter interactions and astrophysical environments such as neutron star magnetospheres. When the energy of the photon becomes comparable to that of the electron, it is necessary to use quantum electrodynamics (QED) to describe the dynamics accurately. However, computing the appropriate scattering matrix element from strong-field QED is not generally possible due to multiparticle effects and the complex structure of the electromagnetic fields. Therefore, these interactions are treated semiclassically, coupling probabilistic emission events to classical electrodynamics using rates calculated in the locally constant field approximation. Here, we provide comprehensive benchmarking of this approach against the exact QED calculation for nonlinear Compton scattering of electrons in an intense laser pulse. We find agreement at the percentage level between the photon spectra, as well as between the models' predictions of absorption from the background field, for normalized amplitudes $a_0 > 5$. We discuss possible routes towards improved numerical methods and the implications of our results for the study of QED cascades. © 2018 Author(s). All article content, except where otherwise noted, is licensed under a Creative Commons Attribution (CC BY) license (http://creativecommons.org/ licenses/by/4.0/). https://doi.org/10.1063/1.5037967

I. INTRODUCTION

Petawatt and multipetawatt laser facilities that reach focussed intensities in excess of $\sim 10^{22}$ W cm⁻² (Refs. 1 and 2) hold great promise for the study of the interaction of charged particles with electromagnetic fields of unprecedented strength.^{3–6} In these environments, the recoil associated with emission of radiation, known as radiation reaction, can become so large that it dominates the particle dynamics.^{7,8} Furthermore, when the energy of individual photons of this radiation becomes comparable to that of the emitting particle, it becomes essential to incorporate quantum effects on this radiation reaction⁹ in our modelling of plasmas as sources of high-energy photons,¹⁰ electron-positron pairs,^{11,12} or as laboratory analogues of high-field astrophysical environments.^{13,14}

However, it is not currently possible to use the most general and accurate approach, the theory of strong-field quantum electrodynamics (QED), to model many scenarios of interest, for reasons we will shortly outline. Instead, a semiclassical approach has been widely adopted for use in numerical simulations of laser-plasma and laser-particlebeam interactions. Inherent to this model are a number of assumptions, making it essential that we benchmark its predictions against those from QED. In this work, we do so for photon emission in the collision of ultrarelativistic electrons with intense laser pulses. We focus on the classically nonlinear, moderately quantum regime, motivated not only by progress in the development of the next generation of highintensity laser facilities,^{15,16} but also by recent experimental work on radiation reaction in strong fields.^{17–19}

Strong-field QED is not used directly to model these kinds of interactions for a number of reasons. First, a scattering-matrix calculation connects asymptotic free states, thereby requiring ab initio complete knowledge of the spatiotemporal structure of the background electromagnetic field; exact analytical calculations have thus far proven to be possible only for certain field configurations that possess high symmetry,²⁰ such as plane electromagnetic waves²¹ and static magnetic fields.²² Furthermore, it is generally assumed that back-reaction effects may be neglected. This is especially important when considering QED cascades,^{11,23,24} in which the initial state contains a single electron, positron, or photon and the final state many of these, because we expect significant absorption of energy from the background.²⁵ (See Ref. 26 for analysis beyond this approximation.) Even in the absence of significant depletion, the multiplicity alone makes the calculation of a cascade within strong-field QED extremely challenging. State-of-the-art results are those in which the final state contains two additional particles, such as double Compton scattering²⁷⁻²⁹ and trident pair crea $tion^{30-33}$ from single electrons.

These conditions, namely, complex field structure, strong depletion due to back-reaction, and high multiplicity, are ubiquitous in the interaction of high-intensity lasers with particle beams or plasma targets. As such, considerable effort has been devoted to the development of numerical



a)tom.blackburn@chalmers.se

schemes that can model QED cascades as well as selfconsistent plasma dynamics.^{34,35} We characterize these as semiclassical because they factorize the cascade into a product of first-order processes that occur in vanishingly small regions linked by classically determined trajectories; the production rates and spectra are calculated for the equivalent QED process in constant, crossed fields.³⁶ This is possible because at high intensity (to be defined in Sec. II), the formation length over which a photon is emitted, or an electronpositron pair is created, is much smaller than the length scale over which the field varies.²¹ The approximation that emission occurs instantaneously, and therefore that the rate for a constant, crossed field may be employed, is called the *locally* constant field approximation (LCFA). Monte Carlo implementations of QED processes based on this have found widespread adoption in particle-in-cell (PIC) codes (see Ref. 37 for details). Depletion in these codes is therefore treated classically, as QED processes alter the plasma current density **j**, which alters the energy density of the self-consistent electric and magnetic fields **E** and **B** via the $\mathbf{j} \cdot \mathbf{E}$ term in Poynting's theorem; see Refs. 38–40 for examples of how this drives laser absorption.

Identifying the parameter regime where this semiclassical picture works, and why, has been the subject of much theoretical work.^{41–43} However, there has been limited direct benchmarking of numerical and analytical results in regimes of experimental interest. For nonlinear Compton scattering (photon emission by an electron), Harvey *et al.*⁴⁴ compared the frequency and angular spectra predicted by (1) integration of the QED probability rate for a monochromatic plane wave and (2) semiclassical simulation of a 100 fs pulsed plane wave with a super-Gaussian temporal profile, concluding that the neglect of interference effects in the latter caused the harmonic structure to be missed.

In this work, we present systematic comparisons of not only the longitudinal and transverse momentum spectra (Secs. III A and III B) but also the absorption of energy from the background field (Sec. III C). We introduce a normalization framework in Sec. II C that guarantees that we compare precisely the same physical scenario. This permits direct, quantitative benchmarking of semiclassical codes against analytical results from QED in the parameter regime relevant for recent and upcoming experiments.

II. METHODS

The interaction geometry is illustrated in Fig. 1. An electron with initial Lorentz factor γ_0 collides head-on with a circularly polarized laser pulse that has dimensionless amplitude a_0 , central frequency ω_0 , and invariant duration τ . Throughout this work, we set $\hbar = c = 1$ and denote the elementary charge by e and the electron mass by m. The pulse vector potential $eA^{\mu}(\phi) = ma_0g(\phi)(0, \sin \phi, \cos \phi, 0)$, where $a_0 = eE_0/(m\omega_0)$ for electric field strength E_0 (Ref. 45) and $g(\phi) = \cos^2[\phi/(4\tau)]$ for phases $|\phi| < 2\pi\tau$. In all the results presented here, $\gamma_0 = 1000$ and $\omega_0 = 1.55 \text{ eV}$ (equivalent to a wavelength of $\lambda = 0.8 \,\mu\text{m}$). We will consider dimensionless amplitudes in the range of $5 \le a_0 \le 30$, which covers the transition between the weakly and highly nonlinear classical



FIG. 1. An electron (blue) with initial Lorentz factor $\gamma_0 \gg 1$ collides headon with an intense, circularly polarized laser pulse (red) that has strength parameter a_0 , angular frequency ω_0 , and duration τ .

regimes, and restrict the laser duration to be $\tau = 2$ or 3 so that the expected number of photons is of order one. This is because our QED calculations are performed for single scattering only, and so that we can gather sufficient statistics in the semiclassical simulations (the fraction of collisions in which only one photon is emitted is exponentially suppressed with increasing a_0).

The quantum interaction of charged particles and photons with strong fields is characterized by the invariants $\chi_e = e\sqrt{-(F.p)^2}/m^3$ and $\chi_{\gamma} = e\sqrt{-(F.k')^2}/m^3$, where *F* is the electromagnetic field tensor and *p* and *k'* are the fourmomenta of the electron and photon, respectively.²¹ χ_e may be interpreted as a measure of the field strength in the rest frame of the electron relative to that of the critical field of QED $E_{\rm crit} = m^2/e$.^{46–48} It is often referred to as the "quantum nonlinearity parameter" by analogy with a_0 , which is the classical nonlinearity parameter.²¹ We have $\chi_e \sim 0.1$ for the interaction parameters under consideration here, and so, quantum effects are non-negligible.

A. QED

The strong-field QED scattering matrix (S-matrix) connects asymptotic free states, evolving the initial state from the distant past to the distant future. The calculation is performed to all orders in the coupling to the background field a_0 , i.e., non-perturbatively, as the number of photons absorbed and reemitted by an electron in an intense laser field is very large. For the lowest order process shown in Fig. 2, the emission of one photon or single nonlinear Compton scattering,^{49–52} it reads

$$S_1 = -ie(2\pi)^3 \delta_{\mathrm{lf}}(p'+k'-p) \sum_j \mathcal{T}_j \mathcal{C}_j.$$
(1)

The delta function ensures the conservation of momentum in the lightfront and transverse directions. By *lightfront*



FIG. 2. Feynman diagrams for the emission of photons with four-momenta $k'_{1,2,...}$ in the scattering of a laser-dressed electron from asymptotic fourmomentum *p* to *p'*. The double lines indicate that the interaction with the background field is calculated to all orders in the coupling a_0 .

momentum, we mean $p^+ \equiv k.p/\omega_0$, which is conserved in a plane wave with wavevector k in the absence of radiation reaction. Other features are the transition operators \mathcal{F}_j , which are sensitive to the electron spins and the photon polarization, and \mathcal{C}_j , which are integrals over the laser phase (see the study by Seipt *et al.*⁵² for details)

$$\mathscr{C}_{j} = \int \mathrm{d}\phi \, \mathscr{F}_{j}(\phi) e^{i \int \mathrm{d}\phi \frac{k' \cdot \pi(\phi)}{k \cdot p'}}.$$
(2)

Here, \mathscr{F}_j are functions of the vector potential $A(\phi)$ and $\pi(\phi)$ in the exponent is the classical kinematic four-momentum of the electron, a solution to the radiation-free Lorentz force equation.

The one-photon emission probability is

$$\frac{\mathrm{d}^{3}\mathbb{P}_{1}}{\mathrm{d}f\mathrm{d}^{2}\mathbf{r}_{\perp}} = \frac{\alpha m^{2}}{(4\pi\omega_{0}p^{+})^{2}} \frac{f}{1-f} \frac{1}{2} \sum_{\mathrm{spin,pol}} \left| \sum_{j} \mathscr{T}_{j} \mathscr{C}_{j} \right|^{2}, \quad (3)$$

where we have defined the lightfront momentum transfer fraction $f = k'^+/p^+$ and normalized transverse photon momentum $\mathbf{r}_{\perp} = \mathbf{k}'_{\perp}/(fm)$. The magnitude of the latter $r_{\perp} = (p^+/m) \tan(\theta/2)$, where θ is the polar angle of the emitted photon, becoming $r_{\perp} \simeq \gamma \theta$ if $\gamma \gg 1$ and $\theta \ll 1$.

If $a_0^3/\chi_e \gg 1$ and *f* is not too small, the phase interval which contributes to the emission of a single photon becomes much smaller than the wavelength of the back-ground field²¹ and interference between emission from different formation regions is suppressed.⁴¹ In this case, the field may be treated as constant over the photon formation region. As the photons are emitted into a narrow cone around the direction of the electron's instantaneous momentum, we can integrate over the transverse momenta \mathbf{r}_{\perp} to obtain the instantaneous probability rate per unit phase and lightfront momentum transfer

$$\frac{\mathrm{d}W}{\mathrm{d}f} = -\frac{\alpha m^2}{\omega_0 p^+} \left[\mathrm{Ai}_1(z) + \left(\frac{2}{z} + \chi_e f \sqrt{z}\right) \mathrm{Ai}'(z) \right], \quad (4)$$

where $z^{3/2} = f/[\chi_e(1-f)]$ and $\chi_e \equiv \chi_e(\phi)$ the local value of the quantum parameter.

B. Semiclassical

In the semiclassical interpretation of the collision process, the electron follows a (radiation-free) classical trajectory between point-like, probabilistically determined, QED events. These events are implemented using the standard Monte Carlo algorithm,^{36,37} with rates calculated in the LCFA, i.e., Eq. (4) (see also Refs. 21, 22, and 53). We use circe, a particle-tracking code that simulates photon and positron production by high-energy electrons (and photons) that collide with laser pulses that have $a_0 \gg 1$. In one spatial dimension, the external field is assumed to be a plane electromagnetic wave, i.e., the fields are determined by a single parameter $\phi = \omega_0 n x$, where ω_0 is the wave frequency, $n = (1, \hat{\mathbf{k}})$ for the direction of propagation $\hat{\mathbf{k}}$, and x is the four-position of the electron. Between emissions, the electron dynamics are determined by the Lorentz force alone. The spatial components of the four-momentum $p = (\gamma m, \mathbf{p})$ that are perpendicular to the wavevector are obtained by integrating

$$\frac{\mathrm{d}\mathbf{p}_{\perp}}{\mathrm{d}\phi} = -\frac{e\mathbf{E}_{\perp}(\phi)}{\omega_0},\tag{5}$$

where \mathbf{E}_{\perp} is the electric field at phase ϕ . The remaining components of p are determined by the two conditions p^+ = const and $p^2 = m^2$. The four-position is determined by $\frac{dv}{d\phi} = p/(\omega_0 p^+)$.

Photon emission is implemented as follows: Each electron is assigned an optical depth against emission $\tau = -\log(1 - R)$ for pseudorandom $0 \le R < 1$, which evolves as

$$\frac{\mathrm{d}\tau}{\mathrm{d}\phi} = -\int_0^1 \frac{\mathrm{d}W}{\mathrm{d}f} \,\mathrm{d}f,\tag{6}$$

where *W* is the instantaneous probability rate of emission given by Eq. (4), until the point where it falls below zero. Then, the lightfront momentum transfer $f = \chi_{\gamma}/\chi_e$ is pseudorandomly sampled from the differential rate and τ is reset. Assuming that emission occurs in the direction parallel to the initial momentum, as the electron emits into a narrow cone of opening angle $1/\gamma$, the momenta of the electron and the photon after the scattering are

$$p' = (m\sqrt{1 + (1 - f)^{2}(\gamma^{2} - 1)}, (1 - f)\mathbf{p}),$$

$$k' = (fm\sqrt{\gamma^{2} - 1}, f\mathbf{p}).$$
(7)

As discussed in Ref. 34, this leads to an error in energy conservation of

$$\Delta \mathcal{E} = \frac{m}{2\gamma} \frac{f}{1-f} + O(\gamma^{-3}), \tag{8}$$

which is small for ultrarelativistic particles.

C. Comparison basis

In this work, we present quantitative, as well as qualitative, comparisons of electron and photon spectra predicted by the exact QED and semiclassical methods. We discuss here how the normalization may be set consistently, but independently by the two methods.

The final result of the QED calculation is the probability \mathbb{P}_1 that a single photon is emitted in collision of an electron with a pulsed electromagnetic plane wave. However, even for the short pulses under consideration here, the fact that $a_0 > 1$ makes it possible for $\mathbb{P}_1 > 1$. Where this occurs, the probability is generally interpreted as the mean number of emitted photons, i.e., $\mathbb{P}_1 \rightarrow N_{\gamma,\text{QED}}$, as this quantity can certainly exceed unity.^{9,44,54} We emphasise that this interpretation is exact only in the classical limit, where recoil can be neglected. The true probability for single scattering is given by the renormalized quantity $\mathbb{P}_1/(1 + \sum_{n=1}^{\infty} \mathbb{P}_n)$. To determine this would require the calculation of the scattering probability to a state containing an arbitrary number of photons *n*. Efforts to characterize such multiphoton interactions analytically have been limited due to the complexity of the

calculations; at present, all the results in the literature are restricted to $n \le 2$. For these reasons, we will define the QED "number of photons" as

$$N_{\gamma,\text{QED}} \equiv \int \frac{\mathrm{d}^3 \mathbb{P}_1}{\mathrm{d}f \mathrm{d}^2 \mathbf{r}_\perp} \,\mathrm{d}f \mathrm{d}^2 \mathbf{r}_\perp. \tag{9}$$

In the semiclassical calculation, multiphoton emission is accounted for by factorisation of the multiphoton emission into a product of first-order processes. Localizing emission in this way allows us to determine the branching fraction to a final state containing an arbitrary number of particles, thereby guaranteeing that $\mathbb{P}_1 < 1$. In the classical limit (i.e., negligible recoil per photon), one emission event is independent of any other and the probability that n photons are emitted in a given collision $\mathbb{P}_n = \lambda^n e^{-\lambda}/n!$, where $\lambda \equiv N_{\gamma,sc}$, the mean number of photons in the semiclassical case. However, we consider here collisions where $\gamma \sim 0.1$ and recoil is not negligible. As the emission rate (at fixed field strength) decreases with increasing particle energy, emitting a photon and so losing energy make it more probable that further photons are emitted. As such, the numbers of photons emitted in two non-overlapping intervals are not independent and the probability \mathbb{P}_n ceases to be Poisson-distributed. In summary, radiation reaction, the recoil due to photon emission, affects the average number of photons because "the emission of each photon modifies the electron state and, consequently, the next emissions."9

Since it is not possible, as yet, to determine the renormalization factor by which the QED results should be scaled, we propose this alternative. The QED results from Eq. (9) are *not* scaled. Equivalent semiclassical spectra are obtained statistically, by generating a large set of collision data, accepting only those collisions in which exactly one photon is emitted, and rescaling such that the spectra have integral $N_{\gamma,sc}$. The mean number of photons $N_{\gamma,sc}$ is determined by considering the entire set of collision data, i.e.,

$$N_{\gamma,\rm sc} \equiv \frac{\sum_i i N_i}{\sum N_i} \frac{1}{N_1} \int \frac{\mathrm{d}^2 N_1}{\mathrm{d}f \,\mathrm{d}r_\perp} \,\mathrm{d}f \,\mathrm{d}r_\perp, \tag{10}$$

where N_i is the number of simulated collisions in which exactly *i* photons are emitted.

This definition ensures that only collisions with a single emission contribute to the shape of the spectrum and that its integral may be interpreted in a manner consistent with the QED result. From now on, all differential spectra will be given in terms of the "number of photons" defined by Eqs. (9) and (10). We note that while this post-facto selection criterion lets us compare the same physical scenario as the QED approach, multiphoton and recoil corrections are still present because $N_{\gamma,sc}$ is affected by radiation reaction. We do not necessarily expect the QED "probability" \mathbb{P}_1 to satisfy $\mathbb{P}_1 \rightarrow N_{\gamma,\text{QED}} = N_{\gamma,sc}$ for this reason.

III. RESULTS

A. Lightfront momentum

parametrization for the differential scattering probability because the conservation of momentum for (single) nonlinear Compton scattering reads $u_0^+ = u_e^+ + u_\gamma^+$, where $u_e^+ = p'^+/m$ and $u_\gamma^+ = k'^+/m$ are the normalized lightfront momenta of the scattered electron and photon, respectively. We will plot differential spectra in terms of the transfer fraction $f = u_\gamma^+/u_0^+$ $= 1 - u_e^+/u_0^+$. In the back-scattering limit, $f \simeq \omega'/(\gamma_0 m)$, where ω' is the energy of the scattered photon.

We compare the analytical and simulation predictions for the number of photons emitted in the head-on collision of an electron with initial energy $\gamma_0 = 1000$ and a two- or threecycle laser pulse in Fig. 3(a), with examples of the differential spectra in Fig. 3(c). The percentage difference between the results of the two methods is given for the total number of photons in Fig. 3(b). We find that the semiclassical method systematically overestimates the number of photons but that the fractional discrepancy diminishes with increasing a_0 , falling below 10% when $a_0 \ge 20$. N_{γ} scales approximately linearly with a_0 as expected in the strong-field regime; the growing discrepancy towards the lower end of the plotted range is an indication of the transition to the perturbative regime where $N_{\gamma} \propto a_0^2$ instead.

The origin of the discrepancy is shown in Fig. 3(c). While there is very good agreement for large f, i.e., high energy, the semiclassical method strongly overestimates the number of photons with vanishing f. This is because the underlying LCFA rate contains an integrable singularity $\propto f^{-2/3}$ absent in the exact QED calculation.⁵⁵ In the latter case, the probability tends to a finite value⁴³

$$\lim_{f \to 0} \frac{\mathrm{d}\mathbb{P}_1}{\mathrm{d}f} = \frac{\alpha a_0^3}{2\chi_e} \int \mathrm{d}\phi \, g^2(\phi),\tag{11}$$

where $g(\phi)$ is the envelope function given in Sec. II. It is not surprising that the semiclassical spectra do not reproduce this limit because the LCFA arises from an expansion in the parameter $\chi_e/a_0^3 \ll 1.^{41}$

The physical meaning of this parameter is that it compares the formation length of the photon to the length scale over which the field varies. If this is sufficiently small, we can assume that emission occurs instantaneously and thereby neglect interference effects. As discussed in the study by Harvey et al.,⁴⁴ this means that Monte Carlo implementations of localized rates cannot reproduce the harmonic structure that becomes visible in the emission spectrum at small f. A simple way to estimate the smallest f for which the LCFA should be valid is to recall that in a monochromatic plane wave, emission over a complete phase oscillation contributes to photons at the first nonlinear Compton edge, for which the transfer fraction $f_C \simeq 2\chi_e/a_0^3$. The requirement that the formation length be smaller than the laser wavelength is then equivalent to having $f \gtrsim f_C$. This limit is consistent with the results shown in Fig. 3(c) and with a more detailed calculation performed by Di Piazza et al.⁴³ In fact, if we cut off the QED and semiclassical spectra below $f = f_C$, the percentage discrepancy in the number of photons falls from 17% to 5% at $a_0 = 10$ and from 5% to 2% at $a_0 = 30$.

While it is important to capture the number spectrum accurately, the dynamically significant quantity is the



FIG. 3. (a) The number of photons emitted in the collision of an electron with $\gamma_0 = 1000$ and a laser pulse with given a_0 and duration τ , calculated analytically from QED (points) and from semiclassical simulation (lines). (b) The percentage difference between the number of photons as evaluated by the two methods. (c) The lightfront-momentum spectrum for the specified collision parameters: results from QED (solid, grey) and semiclassical simulation (dashed, coloured). Vertical, dotted lines indicate f_C , the first Compton edge of a monochromatic plane wave with the same a_0 .

spectrum weighted by u_{γ}^+ , as this gives the momentum change of the electron or radiation reaction.^{56,57} When we compare the total radiated lightfront momentum

$$\mathbb{I}^+ \equiv u_0^+ \int f \, \frac{\mathrm{d}N_\gamma}{\mathrm{d}f} \, \mathrm{d}f \tag{12}$$

in Fig. 4(a), we find much better agreement between the two methods, with a relative discrepancy below 4% even for $a_0 = 5$. This is because the contribution to the electron recoil from the low-energy tail, i.e., the part of the spectrum where the LCFA fails, is very small and diminishes with increasing a_0 . This can also be seen in Figs. 4(c) and 4(d) where we show the intensity spectra without log scaling on the vertical axis.

B. Perpendicular momentum

We parametrize the perpendicular momentum spectrum using the scaled quantity $r_{\perp} \equiv u_{\gamma}^{\perp}/f$. For $\gamma_0 \gg a_0$ and $\gamma_0 \gg 1$, as we have here, $r_{\perp} \simeq \gamma_0 \theta$, where θ is the photon scattering angle. The comparison between the analytical and simulation results, shown in Fig. 5, is for r_{\perp} scaled by the laser strength parameter a_0 for the following reason.

Analysis of nonlinear Compton scattering in a monochromatic, circularly polarized plane wave only in terms of the number of laser photons absorbed has been shown to reproduce the classical result that the photons are typically emitted along the direction of the instantaneous momentum of the electron in the electromagnetic field.²⁵ Assuming that the electron and the laser were initially counterpropagating and that both the electron Lorentz factor γ_0 and the laser amplitude $a_0 \gg 1$, we have that the most probable angle of emission is $\tan \theta = 4a_0\gamma_0/(4\gamma_0^2 - a_0^2) \simeq a_0/\gamma_0$.

In the semiclassical calculation, we capture the electron's transverse oscillation directly by solving the classical equations of motion and rely on relativistic beaming to justify setting the photon's emission direction to be parallel to the electron's instantaneous momentum. We may derive a scaling relation for the average r_{\perp} for the pulsed plane waves under consideration here within the framework of the LCFA. The instantaneous angle between the electron momentum and the laser axis is $\theta(\phi) \simeq a_0 g(\phi)/\gamma_0$ for $\gamma_0 \gg a_0 \gg 1$, where $g(\phi)$ is the pulse envelope described in Sec. II. Assuming that the photon is emitted parallel to the electron momentum, we have that the mean value of r_{\perp}

$$\langle r_{\perp} \rangle = \frac{\gamma_0 \int \theta(\phi) W(\phi) \, \mathrm{d}\phi}{\int W(\phi) \, \mathrm{d}\phi} \simeq \frac{3a_0}{4},\tag{13}$$

where $W(\phi)$ is the emission rate [Eq. (4)] integrated over all f, and we obtain an analytical result by working in the classical limit $\chi_e \ll 1$. In this expression, the mean r_{\perp} is normalized to the number of photons. Consequently, if χ is not too large, we expect $\langle r_{\perp} \rangle / a_0$ predicted by semiclassical simulation to be independent of a_0 and the pulse duration τ . This is indeed what is shown in Fig. 5(a). The exact QED results are generally larger but tend towards the semiclassical results as a_0 is increased. This is because photons are emitted into a broader range of angles in the QED calculation, which can be seen by the fact that the standard deviations of the spectra (normalized to the mean) shown in Fig. 5(c), i.e., the widths of the distribution, are larger in the analytical case.

These integral comparisons lead us to expect important qualitative differences between the r_{\perp} spectra predicted by QED and by semiclassical simulation. Indeed, Fig. 5(e) shows that, unlike the former, the latter exhibit a universal shape when plotted as a function of r_{\perp}/a_0 . Furthermore, they diverge as $r_{\perp} \rightarrow a_0$. No photons are emitted with $r_{\perp} > a_0$. The range of accessible r_{\perp} is much larger for the QED spectra although we note that as a_0 increases, the peak of the spectrum (i.e., the most probable r_{\perp}) tends towards a_0 and the spectra generally become narrower. As we have scaled r_{\perp} down by a_0 , this indicates that the characteristic width of the spectrum is approximately constant for all a_0 .

The cause of these differences is the assumption of collinear emission in the semiclassical simulations, i.e.,



FIG. 4. (a) The total lightfront momentum lost by an electron with $\gamma_0 = 1000$ in a collision with a laser pulse with given a_0 and duration τ , calculated analytically from QED (points) and from semiclassical simulation (lines). (b) The percentage difference between the two. (c) and (d) The lightfront-momentum intensity spectrum for the specified collision parameters: results from QED (solid, grey) and semiclassical simulation (dashed, coloured).

assigning the final momenta according to Eq. (7). Relativistic beaming means that the photon is emitted forward into a cone of half-angle $\sim 1/\gamma_0$, corresponding to a width in r_{\perp} of $\sigma_r \sim 1$. We have neglected this extra angular divergence, which is why the semiclassical results have a sharp edge at $r_{\perp} = a_0$. The range of angles at which a photon can be emitted is bound by the angle between the electron's instantaneous momentum and the laser wavevector, which is at most a_0/γ_0 for the circularly polarized pulses under consideration here.

A more subtle discrepancy may be seen for small r_{\perp} in Fig. 5(e). Whereas the analytical results tend to zero as r_{\perp} does, exhibiting a pronounced shoulder as they do so, all the semiclassical spectra tend to a finite value of approximately 0.022. (This is consistent with a classical calculation of the angular spectrum, which gives $dN_{\gamma}/dr_{\perp} \simeq 5\alpha/\sqrt{3} \simeq 0.021$ for $r_{\perp} = 0$.) The difference is a consequence of the LCFA: recall that in the semiclassical approach, photons are emitted parallel to the electron's instantaneous momentum. Therefore, photons with small r_{\perp} originate from the leading and trailing edges of the pulse, where the local field strength is small and so too is the angle between the electron trajectory and the laser wavevector. In these regions, the effective a_0 is small enough that interference effects become important, suppressing photon emission.

C. Absorption

Energy-momentum conservation demands that the emission of a photon by an electron in a strong background field be accompanied by the absorption of energy from that background field. As the background under consideration here is an electromagnetic wave, this can be interpreted as the absorption of a certain number ℓ of photons. Seipt *et al.*²⁵ have shown that in a circularly polarized, monochromatic plane wave with strength parameter a_0 , the emission of a photon with quantum parameter $f = \chi_{\gamma}/\chi_e$ is associated with the absorption of $\ell = s$ laser photons, where

$$s = \frac{a_0^3}{\chi_e} \frac{f}{1 - f}.$$
 (14)

For the short pulses in this work, the probability that ℓ photons are absorbed is determined numerically.

In the semiclassical method, the background field is treated entirely classically. Nevertheless, we may define an equivalent number of absorbed photons by dividing the classical work done on the moving charge by the laser frequency ω_0

$$\ell = -\frac{1}{\omega_0} \int e \mathbf{v} \cdot \mathbf{E} \, \mathrm{d}x^0. \tag{15}$$



FIG. 5. (a) The mean r_{\perp} of the photon emitted in the collision of an electron with $\gamma_0 = 1000$ and a laser pulse with given a_0 and duration τ , calculated analytically from QED (points) and from semiclassical simulation (lines). (b) The percentage difference between the two. (c) The standard deviation of r_{\perp} , normalized to the mean, and (d) the percentage difference between the QED and semiclassical results. (e) Differential r_{\perp} spectra for the specified collision parameters: results from QED (solid, coloured) and semiclassical simulation (dashed, grey).



FIG. 6. (a) The number of laser photons absorbed in the emission of a single γ photon when an electron with $\gamma_0 = 1000$ collides with a laser pulse with given a_0 and duration τ : results from QED (points) and semiclassical simulation (lines). (b) The percentage difference between the two. (c) The weighted probability that ℓ photons are absorbed, from QED (grey, solid) and semiclassical simulation (coloured, dashed).

Note that for a plane wave, the above integral would be identically zero in the absence of radiation. (The same result holds in the QED calculation: if no photon is emitted, $\ell = 0$.) circe computes Eq. (15) for each test electron, integrating the work done across the entire trajectory of the electron in the pulse.

A comparison between the total number of absorbed photons $N_L \equiv \int \ell \, dN_{\gamma}$, as computed by the two methods, is shown in Fig. 6(a). We find that the semiclassical method systematically underestimates the absorption but that this difference occurs at the level of a few percent and decreases with increasing a_0 . This is in contrast to what we found for the number of photons and the total radiated lightfront momentum, where the semiclassical result was generally larger than the QED result. In all three cases, we expect errors to arise due to the finite size of the formation length and the associated interference; however, here, our results imply that there is some "missing" absorption.

In their analysis of electron-positron pair creation by a photon in a strong laser field, Meuren et al.42 divided the absorbed energy into "classical" and "quantum" parts, with the former being the work done accelerating the daughter particles out of the laser pulse and the latter being the absorption of photons over the formation length. They showed that the classical part scales approximately like a_0^3/χ_e and the quantum part like a_0/χ_e , concluding that the classical absorption should be dominant at high intensity. This is consistent with the results presented here, in that we capture the acceleration of the electron post-emission but not any absorption over the formation length, which is assumed to be vanishingly small. Recall that in the semiclassical simulations, there is an error in energy conservation due to the assumption of collinear emission (see Sec. IIB). Equation (8) predicts that the magnitude of this error is 2.8% at a_0 = 10 and 0.3% at a_0 = 30. This is comparable to the discrepancy shown in Fig. 6(b) but for the largest a_0 , where recoil corrections to $N_{\gamma,sc}$ take effect.

Equation (14) indicates that the larger the lightfront momentum transfer f, the more the photons are absorbed

from the external field. Both the lightfront momentum transferred to an individual photon and the number of emissions increase with a_0 , and so, we expect the absorption to increase as well. Integrating Eq. (14) weighted by the emission rate Eq. (4) over all f, we find that $N_L \sim a_0^2 \mathbb{I}^+$. Here, \mathbb{I}^+ is the total radiated lightfront momentum given by Eq. (12), which scales like a_0^2 in the classical limit. Then, we expect $N_L \sim a_0^4$, which agrees reasonably well with a power-law fit to the data shown in Fig. 6(a); we find $N_L \propto a_0^{3.7}$ for both the QED and semiclassical results. The true scaling is weaker than a_0^4 because of quantum corrections that reduce the radiated power.²²

We showed in Sec. III A that the semiclassical method predicts the large-*f* part of the emission spectrum accurately even for $a_0 = 5$. As this part of the spectrum is associated with the largest ℓ , the agreement between the QED and semiclassical results should be best for $\ell \gg 1$. Four examples of the spectrum of probable ℓ are shown in Fig. 6(c). The agreement between the two is excellent for $\ell > 10$, but the semiclassical method fails to capture the small- ℓ part of the spectrum accurately. This is because it localizes emission, thereby neglecting interference effects; these suppress the emission probability for small ℓ and give rise to the harmonic structure that can be seen in the QED spectra.

D. Exemplary case

Finally, we present a comparison for a specific set of collision parameters, drawing on the systematic results we have so far, to discuss the role of multiple emissions. Figure 7 shows the full set of double- and single-differential photon spectra for lightfront momentum, perpendicular momentum, and absorption for a collision between an electron with $\gamma_0 = 1000$ and a laser pulse with $a_0 = 20$ and $\tau = 2$. The average number of photons is $N_{\gamma} = 2.36$ for these parameters, and therefore, multiphoton effects should be taken into account. However, as the QED calculation is performed only for single scattering, we filter the semiclassical collision data to ensure that the same physical scenario is being compared.



FIG. 7. (Upper row) Double differential photon spectra $d^2N_{\gamma}/(df dr_{\perp})$ (log₁₀-scaled) for a collision between an electron with $\gamma_0 = 1000$ and a laser pulse with $a_0 = 20$ and $\tau = 2$: (a) from QED, (b) from semiclassical simulation, including all emissions, and (c) from semiclassical simulation, filtered to single scattering only. (Lower row) Single differential spectra for (d) the lightfront momentum transfer fraction *f*, (e) the scaled perpendicular momentum r_{\perp} , and (f) the number of laser photons absorbed: results from QED (grey, solid) and from semiclassical simulation including all emissions (blue, dotted) and single scattering only (red, dashed).

Now, we can show the effect of this filtering on the semiclassical results.

The double differential spectrum obtained when all emissions are taken into account is shown in Fig. 7(b); when only single scattering events are binned, we obtain the spectrum shown in Fig. 7(c). As discussed in Sec. III B, in the latter case, we find a sharp cutoff at $r_{\perp} = a_0$ as this is the largest angle between the electron momentum and the laser wavevector, and we assume that photons are emitted in the collinear direction. Then, photons with $r_{\perp} > a_0$ can only come from secondary scattering. Notice that while the QED result [Fig. 7(a)] is generally broader in the vertical direction, the probability that $r_{\perp} > a_0$ diminishes with increasing f, which apparently justifies the assumption of collinear emission for $f \sim 1$. The QED result is also smoother as it is free from the numerical noise inevitable in Monte Carlo simulations. The single scattering spectrum appears noisier as it represents only 20% of the collision data.

Figure 7(d) shows that the effect of the filtering on the lightfront intensity spectrum is rather small. Nevertheless, the agreement is better when only single scattering is included. The spectrum in this case is slightly harder, as secondary photons tend to be emitted with smaller f. Taken in isolation, Fig. 7(e) appears to suggest that the agreement between the QED and semiclassical spectra is better if we do not filter the collision data. The shape of the peak, if not its maximum value, is actually captured better when all emissions are included. This is coincidental. Recalling that we have assumed collinear emission in the semiclassical approach, we would expect the QED result for double Compton scattering to be even broader in r_{\perp} . In fact, the most significant difference is found when we compare the probability that ℓ photons are absorbed from the laser pulse in Fig. 7(f). The two semiclassical spectra have the same integral (by construction, see Sec. II C), and therefore, small values of ℓ must be suppressed when multiple emissions are included. The probability that $\ell = 2 \times 10^4$, for example, is $6 \times$ larger when all emissions are accounted for. Without the selection procedure we have introduced, it would not be possible to compare these against the QED result in a consistent way.

IV. DISCUSSION

It is generally assumed that the semiclassical approach used in modelling high-intensity laser interactions is valid when $a_0 \gg 1$ and $a_0^3/\chi_e \gg 1$. The precise value of a_0 for which these conditions are satisfied depends, however, on the particular quantity that is being calculated. We have shown that for a_0 as low as 5, semiclassical codes accurately capture the part of the emission spectrum for which $f \sim 1$. On the other hand, there is still a significant discrepancy in the total number of photons even when $a_0 = 30$. The error is concentrated in the low-f part of the spectrum, i.e., photons with low energy and large angle; clearly, a semiclassical code should not be used to predict the result of an experimental measurement in the spectral region $f \leq 2\chi_e/a_0^3$. Improving these codes could be accomplished by calculating this threshold and replacing the LCFA rate for photons with smaller f (Ref. 43) although this does require the external field to be treated as a slowly varying plane electromagnetic wave.

Alternatively, it might be possible to bypass this problem by using a PIC code, in which a hybrid description of the electromagnetic field is employed. The Nyquist frequency associated with the finite spacing of the grid naturally separates radiation into two components: lower frequencies are resolved on the grid, i.e., classically, and higher frequencies are treated as "photons," just as we outlined in Sec. II B. This is why many implementations of QED processes in PIC codes include the option of a low-frequency cutoff below which macrophotons are not emitted.⁵⁸ It would be interesting to compare the QED results in this work with the predictions of a semiclassical code in which the radiation spectrum below a certain cutoff is obtained by Fourier analysis of the Liénard-Wiechart potentials in the far field. This would ensure that the formation length is resolved at low f, thereby capturing interference effects. It is reasonable to expect a classical description to be appropriate because both the quantum corrections and the electron recoil should be negligible for photons with $f \ll 1$.

While it is important to make these improvements at low f, this part of the spectrum contributes negligibly to the momentum change of the electron, which is dominated by photons with large f. As the agreement between the QED and semiclassical spectra is much better here, it is not surprising that we find the average lightfront momentum loss predicted semiclassically to be within a few percent of the QED value even at $a_0 = 5$. At $a_0 = 10$, for example, the error in the total number of photons is 16% for both $\tau = 2$ and 3, whereas in the total radiated momentum \mathbb{I}^+ , it is 1.5%, an order of magnitude better. For the experimental parameters in the study by Cole *et al.*¹⁷ ($a_0 \simeq 10$, $\chi_e \simeq 0.1$, and $\gamma_0 \simeq 1000$), the lower limit on f is equivalent to $\omega \leq 100 \text{ keV}$; this part of spectrum represents approximately 16% of the total number of photons but only 0.04% of the total radiated energy, using their parametrization of the spectrum and the measured critical energy of 30 MeV.

This is encouraging for semiclassical or PIC-based modelling of radiation reaction of an electron population. In laser-beam interactions, the particle number density is generally low enough that the radiation spectrum can be obtained by incoherent summation over all photons emitted by the individual particles; this allows the treatment of a beam with a spectrum of energies and non-zero divergence. Although these two effects will wash out, for example, detailed harmonic structure in the momentum spectrum,⁵⁹ the overestimate at low f will survive. Nevertheless, as radiation reaction is an intrinsically multiphoton process,⁹ proper benchmarking requires the calculation of the higher-order diagrams shown in Fig. 2. The selection and scaling scheme we have presented here could easily be extended to comparisons with QED calculations of double, triple, etc., nonlinear Compton scattering.

Perhaps more important for the case of multiple emissions are the comparisons we present for the perpendicular momentum spectra. These test the assumption of collinear emission, which is distinct from the LCFA. While the peak at $r_{\perp} \simeq a_0$ is common to both QED and semiclassical results, the width of the distribution around this point is not captured semiclassically. The angle at which the photon and the electron travel after the scattering affects their quantum parameter, and so, the rates at which secondary processes occur. Even if the change to the rates is small, the cumulative error could become large if the multiplicity is high. Accurate modelling of the angular spectrum is important because, for example, the transverse broadening of an electron beam has been proposed as a signature of quantum effects on radiation reaction.⁶⁰ This broadening would be in addition to that from the finite beaming of the radiation and any initial divergence of the beam (a few milliradians for the laser-wakefieldaccelerated electron beams reported in Refs. 17 and 19). It would also be important for the study of QED avalanches, in which even a single electron accelerated by counterpropagating lasers can seed the creation of a critically dense electron-positron pair plasma. One possible approach would be to implement an angularly resolved LCFA rate that includes the finite $1/\gamma$ beaming of the scattered photon (see Ref. 53 for example).

We have also found that the absorption of energy from the background is dominated by the "classical" component, i.e., $\mathbf{j} \cdot \mathbf{E}$ work done by the external field in accelerating the scattered electron. This is assumed to be the case in PIC modelling of QED avalanches and suggests that a classical treatment of backreaction is reasonably accurate at high intensity. (Recall that in these codes, the fields and currents are evolved self-consistently but classically.) There remains the question of the "missing" absorption we discussed in Sec. III C. On the one hand, the fraction of the total depletion diminishes with increasing a_0 ; however, if this error does arise on a "per-emission" basis, the increased multiplicity at high a_0 could mean that it becomes significant. It is reasonable to expect a causal relationship between the absorption discrepancy and the assumption of collinear emission. This is something we will consider in future work.

V. CONCLUSIONS

In this work, we have presented benchmarking of semiclassical simulations against exact QED results for nonlinear Compton scattering in an intense laser pulse, using a method that guarantees that we compare precisely the same physical scenarios.

The differential spectra agree both qualitatively and quantitatively in the dynamically important region $f \gtrsim 2\chi_e/a_0^3$ that dominates the electron recoil and absorption from the laser fields. We find that the lightfront momentum loss and the number of absorbed photons from semiclassical simulations are within a few percent of the exact QED results for $a_0 > 5$. However, improvements are clearly called for at low f, where the LCFA breaks down, and in the angular distribution, where the agreement is only qualitative due to the assumption of collinear emission in the simulations.

It remains to be seen whether improving these will lead to significant differences in the results of simulations of laser-plasma interactions. In deciding what is most important, we should be guided by further comparison with QED calculations that include multiple emissions. These will place more stringent limits on the validity of the approximations that underpin the semiclassical approach. The fact that experimental exploration of the strong-field, multiphoton regime ($a_0 \sim 10$, $\chi_e \sim 0.1$, $N_\gamma \sim 10$) is already underway makes this an urgent question.

ACKNOWLEDGMENTS

The authors thank T. Heinzl and A. Ilderton for useful discussions and acknowledge support from the Knut and Alice Wallenberg Foundation (T.G.B. and M.M.), the Science and Technology Facilities Council, Grant No. ST/G008248/1 (D.S.), the U.S. DOE under Contract No. DE-

AC02-05CH11231 (S.S.B.), and the Swedish Research Council, Grant Nos. 2013-4248 and 2016-03329 (M.M.).

¹S.-W. Bahk, P. Rousseau, T. A. Planchon, V. Chvykov, G. Kalintchenko, A. Maksimchuk, G. A. Mourou, and V. Yanovsky, "Generation and characterization of the highest laser intensities (10²² W/cm²)," Opt. Lett. **29**, 2837–2839 (2004).

²A. S. Pirozhkov, Y. Fukuda, M. Nishiuchi, H. Kiriyama, A. Sagisaka, K. Ogura, M. Mori, M. Kishimoto, H. Sakaki, N. P. Dover, K. Kondo, N. Nakanii, K. Huang, M. Kanasaki, K. Kondo, and M. Kando, "Approaching the diffraction-limited, bandwidth-limited petawatt," Opt. Express 25, 20486–20501 (2017).

³G. A. Mourou, T. Tajima, and S. V. Bulanov, "Optics in the relativistic regime," Rev. Mod. Phys. **78**, 309–371 (2006).

⁴M. Marklund and P. K. Shukla, "Nonlinear collective effects in photonphoton and photon-plasma interactions," Rev. Mod. Phys. **78**, 591–640 (2006).

- ⁵S. V. Bulanov, T. Z. Esirkepov, Y. Hayashi, M. Kando, H. Kiriyama, J. K. Koga, K. Kondo, H. Kotaki, A. S. Pirozhkov, S. S. Bulanov, A. G. Zhidkov, P. Chen, D. Neely, Y. Kato, N. B. Narozhny, and G. Korn, "On the design of experiments for the study of extreme field limits in the interaction of laser with ultrarelativistic electron beam," Nucl. Instrum. Methods Phys. Res. A 660, 31–42 (2011).
- ⁶A. Di Piazza, C. Müller, K. Z. Hatsagortsyan, and C. H. Keitel, "Extremely high-intensity laser interactions with fundamental quantum systems," Rev. Mod. Phys. **84**, 1177–1228 (2012).
- ⁷S. V. Bulanov, T. Z. Esirkepov, J. Koga, and T. Tajima, "Interaction of electromagnetic waves with plasma in the radiation-dominated regime," Plasma Phys. Rep. **30**, 196–213 (2004).
- ⁸Y. Hadad, L. Labun, J. Rafelski, N. Elkina, C. Klier, and H. Ruhl, "Effects of radiation reaction in relativistic laser acceleration," Phys. Rev. D 82, 096012 (2010).
- ⁹A. Di Piazza, K. Z. Hatsagortsyan, and C. H. Keitel, "Quantum radiation reaction effects in multiphoton Compton scattering," Phys. Rev. Lett. **105**, 220403 (2010).
- ¹⁰A. Gonoskov, A. Bashinov, S. Bastrakov, E. Efimenko, A. Ilderton, A. Kim, M. Marklund, I. Meyerov, A. Muraviev, and A. Sergeev, "Ultrabright GeV photon source via controlled electromagnetic cascades in laser-dipole waves," Phys. Rev. X 7, 041003 (2017).
- ¹¹A. R. Bell and J. G. Kirk, "Possibility of prolific pair production with high-power lasers," Phys. Rev. Lett. **101**, 200403 (2008).
- ¹²C. P. Ridgers, C. S. Brady, R. Duclous, J. G. Kirk, K. Bennett, T. D. Arber, A. P. L. Robinson, and A. R. Bell, "Dense electron-positron plasmas and ultraintense γ rays from laser-irradiated solids," Phys. Rev. Lett. **108**, 165006 (2012).
- ¹³A. K. Harding and D. Lai, "Physics of strongly magnetized neutron stars," Rep. Prog. Phys. 69, 2631 (2006).
- ¹⁴R. Ruffini, G. Vereshchagin, and S.-S. Xue, "Electron-positron pairs in physics and astrophysics: From heavy nuclei to black holes," Phys. Rep. 487, 1–140 (2010).
- ¹⁵ELI-Extreme Light Infrastructure Science and Technology with Ultra-Intense Lasers Whitebook, edited by G. A. Mourou, G. Korn, W. Sandner, and J. L. Collier (THOSS Media GmbH, Berlin, 2011).
- ¹⁶G. Chériaux, F. Giambruno, A. Fréneaux, F. Leconte, L. P. Ramirez, P. Georges, F. Druon, D. N. Papadopoulos, A. Pellegrina, C. Le Blanc, I. Doyen, L. Legat, J. M. Boudenne, G. Mennerat, P. Audebert, G. Mourou, F. Mathieu, and J. P. Chambaret, "Apollon-10p: Status and implementation," AIP Conf. Proc. 1462, 78–83 (2012).
- ¹⁷J. M. Cole, K. T. Behm, E. Gerstmayr, T. G. Blackburn, J. C. Wood, C. D. Baird, M. J. Duff, C. Harvey, A. Ilderton, A. S. Joglekar, K. Krushelnick, S. Kuschel, M. Marklund, P. McKenna, C. D. Murphy, K. Poder, C. P. Ridgers, G. M. Samarin, G. Sarri, D. R. Symes, A. G. R. Thomas, J. Warwick, M. Zepf, Z. Najmudin, and S. P. D. Mangles, "Experimental evidence of radiation reaction in the collision of a high-intensity laser pulse with a laser-wakefield accelerated electron beam," Phys. Rev. X 8, 011020 (2018).
- ¹⁸T. N. Wistisen, A. Di Piazza, H. V. Knudsen, and U. I. Uggerhøj, "Experimental evidence of quantum radiation reaction in aligned crystals," Nat. Commun. 9, 795 (2018).
- ¹⁹K. Poder, M. Tamburini, G. Sarri, A. Di Piazza, S. Kuschel, C. D. Baird, K. Behm, S. Bohlen, J. M. Cole, D. J. Corvan, M. Duff, E. Gerstmayr, C. H. Keitel, K. Krushelnick, S. P. D. Mangles, P. McKenna, C. D. Murphy,
- Z. Najmudin, C. P. Ridgers, G. M. Samarin, D. R. Symes, A. G. R.

Thomas, J. Warwick, and M. Zepf, "Experimental signatures of the quantum nature of radiation reaction in the field of an ultraintense laser," Phys. Rev. X 8, 031004 (2018).

- ²⁰T. Heinzl and A. Ilderton, "Exact classical and quantum dynamics in background electromagnetic fields," Phys. Rev. Lett. **118**, 113202 (2017).
- ²¹V. I. Ritus, "Quantum effects of the interaction of elementary particles with an intense electromagnetic field," J. Sov. Laser Res. 6, 497–617 (1985).
- ²²T. Erber, "High-energy electromagnetic conversion processes in intense magnetic fields," Rev. Mod. Phys. 38, 626–659 (1966).
- ²³A. M. Fedotov, N. B. Narozhny, G. Mourou, and G. Korn, "Limitations on the attainable intensity of high power lasers," Phys. Rev. Lett. **105**, 080402 (2010).
- ²⁴S. S. Bulanov, T. Z. Esirkepov, A. G. R. Thomas, J. K. Koga, and S. V. Bulanov, "Schwinger limit attainability with extreme power lasers," Phys. Rev. Lett. **105**, 220407 (2010).
- ²⁵D. Seipt, T. Heinzl, M. Marklund, and S. S. Bulanov, "Depletion of intense fields," Phys. Rev. Lett. **118**, 154803 (2017).
- ²⁶A. Ilderton and D. Seipt, "Backreaction on background fields: A coherent state approach," Phys. Rev. D 97, 016007 (2018).
- ²⁷D. Seipt and B. Kämpfer, "Two-photon Compton process in pulsed intense laser fields," Phys. Rev. D 85, 101701 (2012).
- ²⁸F. Mackenroth and A. Di Piazza, "Nonlinear double Compton scattering in the ultrarelativistic quantum regime," Phys. Rev. Lett. **110**, 070402 (2013).
- ²⁹B. King, "Double Compton scattering in a constant crossed field," Phys. Rev. A 91, 033415 (2015).
- ³⁰H. Hu, C. Müller, and C. H. Keitel, "Complete QED theory of multiphoton trident pair production in strong laser fields," Phys. Rev. Lett. **105**, 080401 (2010).
- ³¹A. Ilderton, "Trident pair production in strong laser pulses," Phys. Rev. Lett. 106, 020404 (2011).
- ³²B. King and H. Ruhl, "Trident pair production in a constant crossed field," Phys. Rev. D 88, 013005 (2013).
- ³³V. Dinu and G. Torgrimsson, "Trident pair production in plane waves: Coherence, exchange, and spacetime inhomogeneity," Phys. Rev. D 97, 036021 (2018).
- ³⁴R. Duclous, J. G. Kirk, and A. R. Bell, "Monte Carlo calculations of pair production in high-intensity laser–plasma interactions," Plasma Phys. Controlled Fusion 53, 015009 (2011).
- ³⁵N. V. Elkina, A. M. Fedotov, I. Y. Kostyukov, M. V. Legkov, N. B. Narozhny, E. N. Nerush, and H. Ruhl, "QED cascades induced by circularly polarized laser fields," Phys. Rev. Spec. Top. Accel. Beams 14, 054401 (2011).
- ³⁶C. P. Ridgers, J. G. Kirk, R. Duclous, T. G. Blackburn, C. S. Brady, K. Bennett, T. D. Arber, and A. R. Bell, "Modelling gamma-ray photon emission and pair production in high-intensity laser-matter interactions," J. Comp. Phys. 260, 273–285 (2014).
- ³⁷A. Gonoskov, S. Bastrakov, E. Efimenko, A. Ilderton, M. Marklund, I. Meyerov, A. Muraviev, A. Sergeev, I. Surmin, and E. Wallin, "Extended particle-in-cell schemes for physics in ultrastrong laser fields: Review and developments," Phys. Rev. E 92, 023305 (2015).
- ³⁸E. N. Nerush, I. Y. Kostyukov, A. M. Fedotov, N. B. Narozhny, N. V. Elkina, and H. Ruhl, "Laser field absorption in self-generated electron-positron pair plasma," Phys. Rev. Lett. **106**, 035001 (2011).
- ³⁹T. Grismayer, M. Vranic, J. L. Martins, R. A. Fonseca, and L. O. Silva, "Laser absorption via quantum electrodynamics cascades in counter propagating laser pulses," Phys. Plasmas 23, 056706 (2016).
- ⁴⁰D. Del Sorbo, D. R. Blackman, R. Capdessus, K. Smali, C. Slade-Lowther, W. Luo, M. J. Duff, A. P. L. Robinson, P. McKenna, Z.-M. Sheng, J. Pasley, and C. P. Ridgers, "Efficient ion acceleration and dense electronpositron plasma creation in ultra-high intensity laser-solid interactions," New J. Phys. 20, 033014 (2018).
- ⁴¹V. Dinu, C. Harvey, A. Ilderton, M. Marklund, and G. Torgrimsson, "Quantum radiation reaction: From interference to incoherence," Phys. Rev. Lett. **116**, 044801 (2016).
- ⁴²S. Meuren, C. H. Keitel, and A. Di Piazza, "Semiclassical picture for electron-positron photoproduction in strong laser fields," Phys. Rev. D 93, 085028 (2016).
- ⁴³A. Di Piazza, M. Tamburini, S. Meuren, and C. H. Keitel, "Implementing nonlinear Compton scattering beyond the local-constant-field approximation," Phys. Rev. A 98, 012134 (2018).
- ⁴⁴C. N. Harvey, A. Ilderton, and B. King, "Testing numerical implementations of strong-field electrodynamics," Phys. Rev. A 91, 013822 (2015).

- ⁴⁶F. Sauter, "Uber das Verhalten eines Elektrons im homogenen elektrischen Feld nach der relativistischen Theorie Diracs," Z. Phys. 69, 742 (1931).
- ⁴⁷W. Heisenberg and H. Euler, "Folgerungen aus der Diracschen Theorie des Positrons," Z. Phys. 98, 714 (1936).
- ⁴⁸J. Schwinger, "On gauge invariance and vacuum polarization," Phys. Rev. 82, 664–679 (1951).
- ⁴⁹N. B. Narozhnyi, A. I. Nikishov, and V. I. Ritus, "Quantum processes in the field of a circularly polarized electromagnetic wave," Sov. Phys. J. Exp. Theor. Phys. **20**, 622 (1965).
- ⁵⁰M. Boca and V. Florescu, "Nonlinear Compton scattering with a laser pulse," Phys. Rev. A 80, 053403 (2009).
- ⁵¹K. Krajewska and J. Z. Kamiński, "Compton process in intense short laser pulses," Phys. Rev. A 85, 062102 (2012).
- ⁵²D. Seipt, V. Kharin, S. Rykovanov, A. Surzhykov, and S. Fritzsche, "Analytical results for nonlinear Compton scattering in short intense laser pulses," J. Plasma Phys. 82, 655820203 (2016).
- ⁵³V. N. Baier, V. M. Katkov, and V. M. Strakhovenko, *Electromagnetic Processes at High Energies in Oriented Single Crystals* (World Scientific, 1998).

- ⁵⁴M. E. Peskin and D. V. Schroeder, An Introduction to Quantum Field Theory (Westview Press, 1995).
- ⁵⁵V. Dinu, T. Heinzl, and A. Ilderton, "Infrared divergences in plane wave backgrounds," Phys. Rev. D 86, 085037 (2012).
- ⁵⁶C. P. Ridgers, T. G. Blackburn, D. Del Sorbo, L. E. Bradley, C. Slade-Lowther, C. D. Baird, S. P. D. Mangles, P. McKenna, M. Marklund, C. D. Murphy, and A. G. R. Thomas, "Signatures of quantum effects on radiation reaction in laser-electron-beam collisions," J. Plasma Phys. 83, 715830502 (2017).
- ⁵⁷F. Niel, C. Riconda, F. Amiranoff, R. Duclous, and M. Grech, "From quantum to classical modeling of radiation reaction: A focus on stochasticity effects," Phys. Rev. E 97, 043209 (2018).
- ⁵⁸T. D. Arber, K. Bennett, C. S. Brady, A. Lawrence-Douglas, M. G. Ramsay, N. J. Sircombe, P. Gillies, R. G. Evans, H. Schmitz, A. R. Bell, and C. P. Ridgers, "Contemporary particle-in-cell approach to laser-plasma modelling," Plasma Phys. Controlled Fusion **57**, 113001 (2015).
- ⁵⁹T. Heinzl, D. Seipt, and B. Kämpfer, "Beam-shape effects in nonlinear Compton and Thomson scattering," Phys. Rev. A 81, 022125 (2010).
- ⁶⁰D. G. Green and C. N. Harvey, "Transverse spreading of electrons in highintensity laser fields," Phys. Rev. Lett. **112**, 164801 (2014).