



Dissipation and thermal noise in hybrid quantum systems in the ultrastrong-coupling regime

Downloaded from: <https://research.chalmers.se>, 2025-12-10 01:21 UTC

Citation for the original published paper (version of record):

Settineri, A., MacRí, V., Ridolfo, A. et al (2018). Dissipation and thermal noise in hybrid quantum systems in the ultrastrong-coupling regime. *Physical Review A*, 98(5).
<http://dx.doi.org/10.1103/PhysRevA.98.053834>

N.B. When citing this work, cite the original published paper.

This is the accepted manuscript made available via CHORUS. The article has been published as:

Dissipation and thermal noise in hybrid quantum systems in the ultrastrong-coupling regime

Alessio Settineri, Vincenzo Macrì, Alessandro Ridolfo, Omar Di Stefano, Anton Frisk
Kockum, Franco Nori, and Salvatore Savasta

Phys. Rev. A **98**, 053834 — Published 20 November 2018

DOI: [10.1103/PhysRevA.98.053834](https://doi.org/10.1103/PhysRevA.98.053834)

Dissipation and Thermal Noise in Hybrid Quantum Systems in the Ultrastrong Coupling Regime

Alessio Settinieri,¹ Vincenzo Macrì,² Alessandro Ridolfo,² Omar Di
Stefano,² Anton Frisk Kockum,² Franco Nori,^{2,3} and Salvatore Savasta^{1,2}

¹*Dipartimento di Scienze Matematiche e Informatiche,*

Scienze Fisiche e Scienze della Terra,

Università di Messina, I-98166 Messina, Italy

²*Theoretical Quantum Physics Laboratory,*

RIKEN Cluster for Pioneering Research,

Wako-shi, Saitama 351-0198, Japan

³*Physics Department, The University of Michigan,*

Ann Arbor, Michigan 48109-1040, USA

(Dated: October 22, 2018)

Abstract

The interaction among the components of a hybrid quantum system is often neglected when considering the coupling of these components to an environment. However, if the interaction strength is large, this approximation leads to unphysical predictions, as has been shown for cavity-QED and optomechanical systems in the ultrastrong-coupling regime. To deal with these cases, master equations with dissipators retaining the interaction between these components have been derived for the quantum Rabi model and for the standard optomechanical Hamiltonian. In this article, we go beyond these previous derivations and present a general master equation approach for arbitrary hybrid quantum systems interacting with thermal reservoirs. Specifically, our approach can be applied to describe the dynamics of open hybrid systems with harmonic, quasi-harmonic, and anharmonic transitions. We apply our approach to study the influence of temperature on multiphoton vacuum Rabi oscillations in circuit QED. We also analyze the influence of temperature on the conversion of mechanical energy into photon pairs in an optomechanical system, which has been recently described at zero temperature. We compare our results with previous approaches, finding that these sometimes overestimate decoherence rates and underestimate excited-state populations.

I. INTRODUCTION

According to quantum mechanics, a closed system always displays a reversible evolution. However, no quantum system is completely isolated from its environment; for example, control and readout of a quantum system requires some coupling to the outside world, which leads to dissipation and decoherence (see, e.g., [1–4]). Realistic quantum systems should thus be regarded as open, taking into account the coupling to their environments. However, using an exact microscopic approach to include the environment (or reservoir) with its many degrees of freedom is often not feasible. Hence, it is highly desirable to model open quantum systems using a small number of variables. An adequate description of the time evolution of an open quantum system can be provided by the equation of motion for its density matrix: a quantum master equation [5, 6]. Another useful approach is based on the Heisenberg Langevin equation (see, e.g., [7–9]). Microscopic derivations of master equations start from the Hamiltonian dynamics of the total density matrix (for the system plus the environment). Then, tracing out the reservoir degrees of freedom, and introducing some approximations, a master equation can be derived describing the time evolution of the reduced density matrix only for the system [10]. It turns out that the resulting evolution, in general, is no longer unitary, and the open quantum system evolves into mixed states (see, e.g., [11]).

A hybrid quantum system combines two or more physical components or subsystems [12–14], with the goal of exploiting the advantages and strengths of the different systems in order to explore new phenomena and potentially bring about new quantum technologies. An important requirement for the realization of a functional hybrid quantum system is the ability to transfer, with high fidelity, quantum states and properties between its different components. Specifically, the effective coupling rate between the subsystems must be large enough to allow quantum state transfers between them within the shortest coherence time of the two subsystems [14]. This interaction regime is usually called the strong-coupling regime [15]. Cavity quantum electrodynamics (QED) in the strong-coupling regime has demonstrated great capability and potential for the control and manipulation of quantum states [3, 13, 15]. Further increasing the coupling strength, a hybrid quantum system enters the ultrastrong-coupling (USC) regime when the interaction rate becomes comparable to the transition frequency of at least one of the subsystems [3, 16, 17].

It has been shown that USC can give rise to several new interesting physical effects [18–34]. Ultrastrong coupling has been achieved in a variety of cavity-QED and other hybrid condensed-matter systems, including semiconductor polaritons in quantum wells [35–39], superconducting quantum circuits [40–53], a terahertz metamaterial coupled to the cyclotron resonance of a two-dimensional electron gas (2DEG) [54–58], organic molecules [59–64], and in an optomechanical system where a plasmonic picocavity was coupled to vibrations in a molecule [65]. In particular, in the case of superconducting quantum circuits, it is possible to reach the USC regime with even just a single artificial atom coupling to an electromagnetic resonator [40, 41, 52, 53, 66, 67]. Recently, coupling rates exceeding the transition frequencies of the components (deep-strong-coupling regime [20]) have been obtained in both a circuit-QED setup [46, 50] and with a 2DEG [58].

Although the Hamiltonian of a coupled light-matter system contains the so-called counter-rotating terms, allowing the simultaneous creation or annihilation of an excitation in both the matter system and the cavity mode, these terms can be safely neglected for small coupling rates, **if the components interact resonantly or almost resonantly**. However, when the coupling strength becomes a significant fraction of the cavity frequency (or of the emitter’s transition frequency), this often-invoked rotating-wave approximation (RWA) is no longer applicable and the anti-resonant terms in the interaction Hamiltonian significantly change the standard cavity-QED physics [35]. For example, the number of excitations in the cavity-emitter system is no longer conserved [30], even in the absence of drives and dissipation, and the system states become dressed by the presence of virtual excitations [68]. **It has also been demonstrated [69] that counter-rotating terms can induce anomalous qubit transitions (which do not conserve the excitation number) in a superconducting qubit-resonator system detuned from resonance.**

When deriving the master equation for a hybrid quantum system, the interaction between the subsystems is usually neglected when considering their coupling to the environment [see Fig. 1(a)]. This results in the standard quantum-optical master equation [5, 6] (see Sec. II A). This procedure works well in the weak-coupling regime, and can also be safely applied in the strong-coupling regime, when the density of states of the reservoirs and the system-bath interaction strengths are approximately flat (frequency independent) on the scale of the energy-level splittings induced by the interaction between the subsystems. However, it has been shown that when the light-matter interaction increases up to the breakdown




Figure 1. (a) The master-equation approach valid in the weak- and strong-coupling regimes. The light-matter coupling is neglected while deriving the dissipators. (b) The master-equation approach considering the light-matter coupling. As the coupling strength between the two subsystems increases, it becomes necessary to treat dissipation effects including the coupling between the subsystems. This can be done by developing the system operators describing the coupling to the reservoirs in the eigenbasis of the coupled light-matter system.

of the RWA, this approach leads to unphysical predictions, e.g., excitations in the system even at zero temperature [21]. A closely related problem arising in the USC regime is the failure of standard input-output theory [22, 24, 70–72], which predicts an unphysical output of photons when the hybrid quantum system is in its ground state.

In order to overcome the problems in the description of dissipation of cavity-QED systems in the USC regime, a master equation taking into account the non-Markovian nature of the baths has been developed [73]. Furthermore, Ref. [21] showed that a master equation working properly in the USC regime of cavity QED can be obtained by including the light-matter coupling in the derivation of dissipative terms of the master equation [see Fig. 1(b)]. This approach does not require the introduction of non-Markovian baths. The decoherence rates entering the modified master equation instead depend on the bath noise spectrum evaluated at the dressed transition frequencies of the light-matter system. Since this modified master equation is obtained after a post-trace RWA, it can only be applied to nonlinear interacting quantum systems with anharmonicity larger than the transition linewidths. This prevents the application of this approach to cavity-QED systems in the USC dispersive regime (see

Sec. III A), and to other hybrid quantum systems displaying a coexistence of harmonic (or quasi-harmonic) and anharmonic transitions, e.g., optomechanical systems. In order to describe the losses through the mirror of a cavity embedding matters, a master equation of a non-Lindblad form was also derived [74]. For optomechanical systems in the USC regime, an analogous *dressed-state* master-equation approach has been developed [75], but it also has limitations (see Sec. III B). A zero-temperature master equation able to describe systems with both anharmonic and (quasi-) harmonic transitions has been introduced to study a cavity-QED system in the USC and dispersive regimes [76]. However, a finite-temperature master equation is an essential tool for a precise analysis of experimental results, which, to some degree, are always affected by thermal noise. **A master equation without the post-trace RWA has been derived to describe a general spin-boson problem mapped into a finite-temperature Rabi model in ultra-strong coupling in Ref. [77].**

The main purpose of this article is to provide a general approach for the description of dissipation in arbitrary hybrid quantum systems with arbitrary coupling strengths between its components. We do this by presenting a generalized master equation able to describe systems with both harmonic and anharmonic transitions, also valid for non-zero-temperature reservoirs. The only key assumption in our derivation is a weak system-bath interaction, such that the usual second-order Born approximation can be applied (recently, different approaches where this assumption can be relaxed, have been developed in Refs. [78–81]).

In particular, we decompose the system operators in terms of the dressed states of the hybrid quantum system and derive the master equation without performing the usual secular approximation. Finally, we take care of possible numerical instabilities due to the presence of fast oscillating terms.

The outline of this article is as follows: We begin in Sec. II by briefly reviewing the standard quantum-optical master equation (Sec. II A) and the dressed master equation for anharmonic systems (Sec. II B). Section II C is devoted to the presentation of a non-Lindblad generalized master equation, able to overcome the limitations of the dressed approach of Sec. II B and to take into account non-zero-temperature reservoirs. We also give a suitable solution for some numerical stability problems of our generalized master equation. In Secs. III A and III B, we apply this generalized master equation to calculate the dynamics of a circuit-QED system and an optomechanical system, respectively, at non-zero temperatures, comparing the obtained results with the standard approaches used previously. We conclude

in Sec. IV. In Appendix A, we present more details for the derivation of the generalized dressed master equation.

II. MASTER EQUATIONS

In this section, we introduce dissipation for hybrid quantum systems following three different approaches. We start with the standard master equation, generally used for the description of open systems in quantum optics. Then we introduce the dressed master equation [21]. Finally, we consider a generalized dressed approach, able to describe the dissipation of hybrid quantum systems with arbitrary coupling strength, valid for systems displaying harmonic, quasi-harmonic, and anharmonic transitions, while also considering non zero-temperature reservoirs.

We begin by considering a generic system consisting of N interacting components or subsystems. Each i th component is weakly coupled to an independent bath, modelled as a collection of quantum harmonic oscillators, described by the free Hamiltonian ($\hbar = 1$ throughout this article)

$$\hat{H}_B^{(i)} = \sum_l \nu_l \hat{b}_{i,l}^\dagger \hat{b}_{i,l}, \quad (1)$$

where $\hat{b}_{i,l}$ ($\hat{b}_{i,l}^\dagger$) are bosonic annihilation (creation) operators for the l th bath mode with frequency ν_l of the i th reservoir. The system-bath (denoted by the subscript SB) interaction Hamiltonian is given by

$$\hat{H}_{SB} = \sum_{i,l} \alpha_{i,l} (\hat{s}_i + \hat{s}_i^\dagger) (\hat{b}_{i,l} + \hat{b}_{i,l}^\dagger), \quad (2)$$

where \hat{s}_i (\hat{s}_i^\dagger) are annihilation (creation) operators of the i th subsystem, mediating the interaction with the reservoirs. We denoted the coupling strength of the i th subsystem to the bath mode l of the i th reservoir by $\alpha_{i,l}$. In the interaction picture, the system-bath interaction Hamiltonian takes the form

$$\hat{\tilde{H}}_{SB} = \sum_{i,l} \alpha_{i,l} e^{\imath \hat{H}_S t} (\hat{s}_i + \hat{s}_i^\dagger) e^{-\imath \hat{H}_S t} (\hat{b}_{i,l} e^{-\imath \nu_{i,l} t} + \hat{b}_{i,l}^\dagger e^{\imath \nu_{i,l} t}), \quad (3)$$

where \hat{H}_S is the system Hamiltonian and \imath is the imaginary unit.

A. Standard Master Equation

In the standard approach, the components or subsystems are assumed to be independent while obtaining the dissipation. The coupling between the components is afterwards introduced in the system Hamiltonian. This leads to the Schrödinger-picture standard master equation

$$\dot{\hat{\rho}} = -i[\hat{H}_S, \hat{\rho}] + \mathcal{L}_{\text{bare}}\hat{\rho}, \quad (4)$$

where $\hat{\rho}$ is the density matrix of the system and

$$\mathcal{L}_{\text{bare}}\hat{\rho} = \sum_i \left\{ \gamma_i [1 + n(\omega_i, T_i)] \mathcal{D}[\hat{s}_i] \hat{\rho} + \gamma_i n(\omega_i, T_i) \mathcal{D}[\hat{s}_i^\dagger] \hat{\rho} \right\}, \quad (5)$$

with the generic dissipator

$$\mathcal{D}[\hat{O}] \hat{\rho} = \frac{1}{2} (2\hat{O}\hat{\rho}\hat{O}^\dagger - \hat{\rho}\hat{O}^\dagger\hat{O} - \hat{O}^\dagger\hat{O}\hat{\rho}). \quad (6)$$

In Eq. (5), the γ_i 's describe the leakage rates and $n(\omega_i, T_i)$ is the average thermal population of the i th reservoir at temperature T_i and the frequency ω_i at which \hat{s}_i rotates in the interaction picture. Pure dephasing effects can be included by adding to Eq. (5) the additional term $(\gamma_{\phi_i}/2) \mathcal{D}[\hat{d}_i] \hat{\rho}$, where \hat{d}_i are system operators that do not change the energy of the system, and γ_{ϕ_i} are the pure dephasing rates.

The master equation provided in Eq. (4) can be used to describe many cavity- and circuit-QED experiments in the weak- and strong-coupling regimes [3, 5, 6]. However, it has been shown that when the coupling between the components or subsystems increases beyond the point where the RWA is applicable, this approach leads to *unphysical* predictions, e.g., production of excitations in the system even at zero temperature [21].

B. Master equations in the dressed picture

1. Master equation for anharmonic systems

In order to overcome the limitations of the standard approach, Ref. [21] developed a dressed master equation, taking into account the coupling between all the components of the system. They also considered that transitions in the hybrid system occur between dressed eigenstates, not between the eigenstates of the free Hamiltonians of the components. In the

following, we briefly show some key points of the dressed master equation derivation. We first express the system Hamiltonian in the dressed basis of its energy eigenstates. We then switch to the interaction picture, writing the system operators as

$$\hat{S}_i(t) = \sum_{j,k>j} C_{jk} |j\rangle\langle k| e^{i\Delta_{jk}t}, \quad (7)$$

with

$$C_{jk} = \langle j | \hat{s}_i + \hat{s}_i^\dagger | k \rangle, \quad (8)$$

$$\Delta_{jk} = E_j - E_k, \quad (9)$$

and the reservoir operators as

$$\hat{B}(t) = \sum_{i,l} \alpha_{i,l} \hat{b}_{i,l} e^{-i\omega_l t}, \quad (10)$$

In this way, the system operators \hat{s}_i are expressed as a sum over transition operators $|j\rangle\langle k|$, which cause transitions (with frequency Δ_{jk}) between eigenstates of the hybrid quantum system $\{|j\rangle, |k\rangle\}$. Note that “ \sim ” identifies the operators in the interaction picture. With these new dressed operators, Eq. (3) can be split into two parts; one each for the dressed system operators with positive and negative frequencies:

$$\hat{H}_{\text{SB}} = \sum_i \left\{ \hat{S}_i(t) \hat{B}_i^\dagger(t) + \hat{S}_i^\dagger(t) \hat{B}_i(t) \right\}. \quad (11)$$

Note that, as shown in Ref. [21], the fast oscillating terms $\hat{S}_i^\dagger(t) \hat{B}_i^\dagger(t)$ and $\hat{S}_i(t) \hat{B}_i(t)$ have been dropped by an initial RWA and the diagonal terms arising from degenerate transitions with $j = k$ are neglected considering a system displaying parity symmetry (in this case $C_{jj} = 0$). By following the standard procedure [5] (second-order Born approximation, Markov approximation, assuming reservoirs with a continuum of frequencies and performing the secular approximation), as shown in detail in Ref. [21], for this simplified version of Eq. (3), we obtain a dressed master equation that in the Schrödinger picture can be written as,

$$\dot{\hat{\rho}} = -i[\hat{H}_S, \hat{\rho}] + \mathcal{L}_{\text{dressed}} \hat{\rho}, \quad (12)$$

with the Lindbladian superoperator

$$\mathcal{L}_{\text{dressed}} \hat{\rho} = \sum_i \sum_{j,k>j} \left\{ \Gamma_i^{jk} n(\Delta_{jk}, T_i) \mathcal{D}[|j\rangle\langle k|] \hat{\rho} + \Gamma_i^{jk} [1 + n(\Delta_{jk}, T_i)] \mathcal{D}[|k\rangle\langle j|] \hat{\rho} \right\}, \quad (13)$$

where the thermal populations are ($k_B = 1$ throughout this article)

$$n(\Delta_{jk}, T_i) = [\exp \{\Delta_{jk}/T_i\} - 1]^{-1} \quad (14)$$

and the damping rates are

$$\Gamma_i^{jk} = 2\pi g_i(\Delta_{jk}) |\alpha_i(\Delta_{jk})|^2 |C_{jk}|^2. \quad (15)$$

with $g(\Delta_{jk})$ being the reservoir density of states and $\alpha(\Delta_{jk})$ the system-reservoir coupling strength.

As shown by several studies [21–24, 29, 71, 82–85], the Lindbladian in Eq. (13) can correctly describe the dynamics of anharmonic cavity-QED systems in the USC regime. At $T = 0$, rather than exciting the system, the dissipators give relaxation to the true dressed ground state. At $T \neq 0$, these dissipators correctly describe the relaxation to the thermal-equilibrium density matrix for the interacting system [71]. However, because of the secular approximation used in the derivation of Eq. (13), this standard approach is *not* able to describe dissipation or decoherence in open quantum systems with mixed harmonic-anharmonic or quasi-harmonic spectra [21], e.g, for cavity QED in the dispersive regime and cavity optomechanics.

C. Generalized master equation

1. Derivation

In this section, we extend the previous treatment in order to derive a generalized dressed master equation able to describe both harmonic and mixed harmonic-anharmonic systems coupled to non-zero-temperature reservoirs. Moreover, the present derivation is not limited to systems with parity symmetry.

We start expressing the system Hamiltonian in the dressed basis of its energy eigenstates. We then switch to the interaction picture, writing the system operators as

$$\hat{S}_i(t) = \sum_{\epsilon' - \epsilon = \omega} \hat{H}(\epsilon) (\hat{s}_i + \hat{s}_i^\dagger) \hat{H}(\epsilon') e^{-i\omega t} = \sum_{\epsilon' - \epsilon = \omega} \hat{S}_i(\omega) e^{-i\omega t}, \quad (16)$$

and the reservoir operators as in Eq. (10), labelling the eigenvalues of \hat{H}_S by ϵ and denoting the projectors onto the respective eigenspaces by $\hat{\Pi}(\epsilon) \equiv |\epsilon\rangle\langle\epsilon|$. Recall that the symbol

“ \sim ” identifies interaction-picture operators. In this way, the system operators \hat{s}_i are expressed as a sum over transition operators, which cause transitions (with transition frequency ω) between energy eigenstates of the hybrid quantum system. For $\omega > 0$, $\hat{S}_i(\omega)$ is a positive-frequency operator that takes the system from an eigenstate with higher energy to one with lower energy. Conversely, for $\omega < 0$, $\hat{S}_i(\omega)$ is a negative-frequency operator which produces a transition to a higher-energy eigenstate. In the following, to emphasize these properties, we introduce the notation

$$\begin{aligned}\hat{S}_i^{(+)}(\omega) &= \hat{S}_i(\omega) & \text{for } \omega > 0, \\ \hat{S}_i^{(-)}(\omega) &= \hat{S}_i(-\omega) & \text{for } \omega > 0, \\ \hat{S}_i^{(0)} &= \hat{S}_i(\omega) & \text{for } \omega = 0,\end{aligned}\tag{17}$$

With these new dressed operators, Eq. (3) can be re-written in a way that makes it easy to derive the Born-Markov master equation for the system:

$$\hat{H}_{\text{SB}} = \sum_i \hat{S}_i(t) \left[\hat{B}_i^\dagger(t) + \hat{B}_i(t) \right].\tag{18}$$

Following the standard procedure (see Appendix A) the generalized dressed master equation can be obtained evaluating the double integrals in Eq. (A4) of Appendix A without assuming parity symmetry, and evaluating the two integrals without introducing the secular approximation $\omega = \omega'$. In this case, we obtain a Liouvillian superoperator \mathcal{L} that, considering all the different subsystems, in the Schrödinger picture, can be written in the general form

$$\begin{aligned}\mathcal{L}_{\text{gme}}\hat{\rho} = \frac{1}{2} \sum_i \sum_{\omega, \omega'} \bigg\{ & \Gamma_i(-\omega') n(-\omega', T_i) \left[\hat{S}_i(\omega') \hat{\rho}(t) \hat{S}_i(\omega) - \hat{S}_i(\omega) \hat{S}_i(\omega') \hat{\rho}(t) \right] \\ & + \Gamma_i(\omega) n(\omega, T_i) \left[\hat{S}_i(\omega') \hat{\rho}(t) \hat{S}_i(\omega) - \hat{\rho}(t) \hat{S}_i(\omega) \hat{S}_i(\omega') \right] \\ & + \Gamma_i(\omega) [n(\omega, T_i) + 1] \left[\hat{S}_i(\omega) \hat{\rho}(t) \hat{S}_i(\omega') - \hat{S}_i(\omega') \hat{S}_i(\omega) \hat{\rho}(t) \right] \\ & + \Gamma_i(-\omega') [n(-\omega', T_i) + 1] \left[\hat{S}_i(\omega) \hat{\rho}(t) \hat{S}_i(\omega') - \hat{\rho}(t) \hat{S}_i(\omega') \hat{S}_i(\omega) \right] \bigg\}\end{aligned}\tag{19}$$

where

$$\Gamma_i(\omega) = 2\pi g_i(\omega) |\alpha_i(\omega)|^2,\tag{20}$$

and “gme” refers to generalized master equation.

Equation (19) contains several terms since both the transition frequencies ω and ω' can be positive, negative and zero, although both $\Gamma_i(\omega)$ and $n(\omega, T_i)$ are non-zero for positive

frequencies only. Moreover, only a few of these terms are relevant in order to correctly describe the system dynamics. Indeed, the terms with oscillation frequencies significantly larger than the damping rates Γ_i of the system provide negligible contributions when integrating the master equation. Equation (19) also contains terms with $\omega' = \omega = 0$, originating from diagonal transition operators or, more generally, operators describing zero-frequency transitions. These terms give rise to additional pure dephasing contributions. Note that these terms can be regarded as a generalization of those appearing in the master equation for optomechanical systems in the USC regime [75].

Expanding Eq. (19), we obtain terms oscillating at frequencies $\pm(\omega' \pm \omega)$ arising from products of $\hat{S}_i^{(-)}$ and $\hat{S}_i^{(+)}$. We also obtain terms oscillating at frequencies $-\omega'$, $+\omega$ arising from products of $\hat{S}_i^{(-)}$ or $\hat{S}_i^{(+)}$ with $\hat{S}_i^{(0)}$, and non-oscillating terms arising from products between zero-frequency operators $\hat{S}_i^{(0)}$. Moreover, considering a system with well separated energy levels ($\omega \gg \Gamma_i$), the terms oscillating at $\pm(\omega + \omega')$, $+\omega$ and $-\omega'$ can be considered as rapidly oscillating and can be neglected. Including only those terms providing non-negligible contributions to the dynamics, the Liouvillian in Eq. (19) can be written as

$$\begin{aligned} \mathcal{L}_{\text{gme}}\hat{\rho} = & \frac{1}{2} \sum_i \sum_{(\omega, \omega') > 0} \left\{ \Gamma_i(\omega') n(\omega', T_i) [\hat{S}_i^{(-)}(\omega') \hat{\rho}(t) \hat{S}_i^{(+)}(\omega) - \hat{S}_i^{(+)}(\omega) \hat{S}_i^{(-)}(\omega') \hat{\rho}(t)] \right. \\ & + \Gamma_i(\omega) n(\omega, T_i) [\hat{S}_i^{(-)}(\omega') \hat{\rho}(t) \hat{S}_i^{(+)}(\omega) - \hat{\rho}(t) \hat{S}_i^{(+)}(\omega) \hat{S}_i^{(-)}(\omega')] \\ & + \Gamma_i(\omega) [n(\omega, T_i) + 1] [\hat{S}_i^{(+)}(\omega) \hat{\rho}(t) \hat{S}_i^{(-)}(\omega') - \hat{S}_i^{(-)}(\omega') \hat{S}_i^{(+)}(\omega) \hat{\rho}(t)] \\ & + \Gamma_i(\omega') [n(\omega', T_i) + 1] [\hat{S}_i^{(+)}(\omega) \hat{\rho}(t) \hat{S}_i^{(-)}(\omega') - \hat{\rho}(t) \hat{S}_i^{(-)}(\omega') \hat{S}_i^{(+)}(\omega)] \\ & + \Omega_i^+(T_i) [\hat{S}_i^{(0)} \hat{\rho}(t) \hat{S}_i^{(0)} - \hat{S}_i^{(0)} \hat{S}_i^{(0)} \hat{\rho}(t)] \\ & + \Omega_i'^+(T_i) [\hat{S}_i^{(0)} \hat{\rho}(t) \hat{S}_i^{(0)} - \hat{\rho}(t) \hat{S}_i^{(0)}(\omega') \hat{S}_i^{(0)}] \\ & + \Omega_i^-(T_i) [\hat{S}_i^{(0)} \hat{\rho}(t) \hat{S}_i^{(0)} - \hat{\rho}(t) \hat{S}_i^{(0)} \hat{S}_i^{(0)}] \\ & \left. + \Omega_i'^-(T_i) [\hat{S}_i^{(0)} \hat{\rho}(t) \hat{S}_i^{(0)} - \hat{S}_i^{(0)} \hat{S}_i^{(0)} \hat{\rho}(t)] \right\} \end{aligned} \quad (21)$$

with

$$\Omega_i'^{\pm}(T_i) = \int_0^t d\tau \int_0^\infty d\nu g_i(\nu) |\alpha_i(\nu)|^2 [n(\nu, T_i) + 1] e^{\pm i\nu\tau}, \quad (22)$$

$$\Omega_i^{\pm}(T_i) = \int_0^t d\tau \int_0^\infty d\nu g_i(\nu) |\alpha_i(\nu)|^2 n(\nu, T_i) e^{\pm i\nu\tau}. \quad (23)$$

We also observe that, for the particular case of an Ohmic bath, where

$$g_i(\nu) |\alpha_i(\nu)|^2 = \frac{\gamma_i \nu}{2\pi f_i}, \quad (24)$$

with γ_i and f_i being, respectively, the damping and the frequency of the considered subsystem, we obtain

$$\Gamma_i(\omega) = \frac{\gamma_i \omega}{f_i}, \quad (25)$$

and all the pure dephasing rates give the same result

$$\Omega_i'^{\pm}(T_i) = \Omega_i^{\pm}(T_i) = \Omega(T_i), \quad (26)$$

$$\Omega(T_i) = \frac{\gamma_i}{4f_i} T_i. \quad (27)$$

In the next section, we apply this generalized dressed master equation to two hybrid quantum systems, comparing the obtained numerical results with previous approaches.

2. Stability problems

We observe that the dissipator in Eq. (21) is not in Lindblad form and, consequently, properties like the positivity of the density matrix and the conservation of the probability cannot be guaranteed. Furthermore, in this framework, some useful theorems [86] on the steady-state behaviour have not been proven yet.

Actually, a careful inspection of Eq. (21) shows that it can be regarded as approximately Lindblad-like. Specifically, if we consider the interaction picture, each term of Eq. (21) (except the last) oscillates at frequencies $\pm(\omega - \omega')$. If $(\omega - \omega')$ is significantly larger than the damping rates Γ_i of the system, these terms provide negligible contributions when integrating the master equation. Hence, $|\omega - \omega'|$ can be assumed to be of the order of the system linewidths. It is thus reasonable to assume for the thermal populations of the reservoirs $n(\omega, T_i) \simeq n(\omega', T_i)$ and for the dampings $\Gamma_i(\omega) \simeq \Gamma_i(\omega')$. This analysis shows that, within a very good approximation, the dissipator in Eq. (21) can be regarded to be in Lindblad form.

Although the fast oscillating terms arising in Eq. (21), produced from transitions with high frequency differences (not present after the post-trace RWA), should not provide a significant contribution for $|\omega - \omega'| > \Gamma_i$, they can strongly increase the computation time and lead to computational instabilities. In order to overcome these difficulties, we use numerical filtering with a step-like function that sets to zero all the dissipator terms involving frequency differences higher than a certain value Λ . More specifically, the filtered Liouvillian

takes the form

$$\mathcal{L}_{\text{gme}}^{\text{filt}} \hat{\rho} = \mathcal{L}_{\text{gme}} \hat{\rho} \times F(\omega, \omega'), \quad (28)$$

where the filter function $F(\omega, \omega')$ can be written in a generalized form as

$$F(\omega, \omega') = \Theta(|\omega - \omega'|) - \Theta(|\omega - \omega'| - \Lambda), \quad (29)$$

with Θ the Heaviside step function and Λ the bandwidth of the filter.

III. DISSIPATION IN THE USC REGIME

In this section, we apply the generalized master equation presented in the previous section to study the influence of temperature on the dynamics of two open hybrid quantum systems in the USC regime. Specifically, we re-examine the dynamics of the two systems presented in Refs. [27] and [87]. The first example is a circuit-QED system in the dispersive regime, displaying multiphoton quantum Rabi oscillations. For this setup, we also compare the results obtained with the generalized master equation to those obtained using the dressed approach for anharmonic systems [21].

The second example is an optomechanical system with coexisting harmonic and anharmonic spectra. Specifically, we consider an ultra-high-frequency mechanical oscillator ultrastrongly coupled to a microwave resonator. Very recently, considering zero-temperature reservoirs, it has been shown [88] that this system is promising for the observation of the dynamical Casimir effect (DCE), which converts mechanical energy into photon pairs [89]. Here we analyze the influence of temperature on this fundamental quantum effect. Moreover, in order to understand the impact of the generalized master equation on the dynamics of hybrid quantum systems, we compare the obtained numerical results with those obtained using a previously developed approach for USC optomechanics [75]. **Note that all the numerical results are displayed in the lab frame.**

A. Circuit QED beyond the RWA

In this circuit-QED example, we study a flux qubit coupled to a single-mode resonator [27]. The bare qubit Hamiltonian can be written as

$$\hat{H}_q = \omega_q \hat{\sigma}_z / 2, \quad (30)$$

where the qubit resonance frequency is $\omega_q = \sqrt{\Delta^2 + (2I_p\delta\Phi_x)^2}$, with Δ the qubit energy gap, I_p the persistent current corresponding to the minima of the qubit potential, and $\delta\Phi_x$ the flux offset. The bare resonator Hamiltonian is

$$\hat{H}_c = \omega_c \hat{a}^\dagger \hat{a}, \quad (31)$$

where ω_c is the frequency of the resonator mode and \hat{a} (\hat{a}^\dagger) is the bosonic annihilation (creation) operator for that mode. The total quantum system is described by the generalized quantum Rabi Hamiltonian

$$\hat{H}_S = \hat{H}_q + \hat{H}_c + g\hat{X}[\cos(\theta)\hat{\sigma}_x + \sin(\theta)\hat{\sigma}_z], \quad (32)$$

where the flux dependence is encoded in $\cos(\theta) = \Delta/\omega_q$, $\hat{X} = \hat{a} + \hat{a}^\dagger$, and $\hat{\sigma}_x$, $\hat{\sigma}_z$ are Pauli matrices.

As shown in Ref. [27], the lowest energy levels of this system display a well-known avoided level crossing arising for $\omega_q \simeq \omega_c$ (vacuum Rabi splitting). This avoided crossing is due to the coherent coupling of the states $|e, 0\rangle$ and $|g, 1\rangle$, where g (e) indicates the ground (excited) state of the qubit and the second entry in the kets represents the photon number. However, when the RWA breaks down, the counter-rotating terms in Eq. (32) must be taken into account, and the total number of excitations in the system is no longer conserved [41, 69]. As a consequence, the coherent coupling between states with different numbers of excitations, not allowed in the standard Jaynes-Cummings model [90, 91], becomes possible through virtual transitions mediated by the counter-rotating terms [31]. This generates several additional avoided level crossings between states with different excitation numbers, e.g., between $|e, 0\rangle$ and $|g, 2\rangle$ [27].

For our numerical calculations, we consider, as in Ref. [27], $\omega_c/2\pi = 4.0$ GHz and a resonator-qubit coupling strength $g/\omega_c = 0.157$. We focus on the avoided crossing arising at $\omega_q \simeq 2\omega_c$ between the states $|\psi_\pm\rangle \simeq \frac{1}{\sqrt{2}}(|e, 0\rangle \pm |g, 2\rangle)$. We set $\omega_q/2\pi = 7.97$ GHz (obtained using the qubit parameters $\Delta/h = 2.25$ GHz, $2I_p = 1.97$ nA, and $\delta\Phi_x = 3.88\Phi_0$); this is where the splitting reaches its minimum [27]. **The minimum splitting $2\Omega_{\text{eff}}$ provides a direct measurement of the effective resonant coupling Ω_{eff} between the states $|e, 0\rangle$ and $|g, 2\rangle$.**

In order to probe this avoided crossing, we consider, as in Ref. [27], the case where the qubit is directly excited by a Gaussian π -pulse,

$$\hat{H}_p = \mathcal{E}(t) \cos(\omega t) \hat{\sigma}_x, \quad (33)$$

where $\mathcal{E}(t) = \Omega \exp[-(t - t_0)^2/2\tau^2]/(\tau\sqrt{2\pi})$. Here, τ is the standard deviation and $\Omega/\omega_c = (\pi/3) \times 10^{-1}$ the amplitude of the pulse. The center frequency of the pulse corresponds to the middle of the avoided crossing considered here. Specifically, $\omega = (\omega_{3,0} + \omega_{2,0})/2$, with $\omega_{i,j} = \omega_i - \omega_j$, where we labelled the energy values and the eigenstates of the hybrid system as ω_l and $|l\rangle$, with $l = 0, 1, \dots$, such that $\omega_k > \omega_j$ for $k > j$.

The system dynamics is then evaluated using the generalized master equation (gme)

$$\dot{\hat{\rho}} = -i[\hat{H}_S + \hat{H}_p, \hat{\rho}] + \mathcal{L}_{\text{gme}}\hat{\rho}, \quad (34)$$

where, considering an Ohmic bath, the Liouvillian dissipator can be written as

$$\begin{aligned} \mathcal{L}_{\text{gme}}\hat{\rho} = & \sum_{(\omega, \omega') > 0} \frac{1}{2} \left\{ \frac{\gamma\omega'}{\omega_q} n(\omega', T_\gamma) [\hat{P}^{(-)}(\omega') \hat{\rho} \hat{P}^{(+)}(\omega) - \hat{P}^{(+)}(\omega) \hat{P}^{(-)}(\omega') \hat{\rho}] \right. \\ & + \frac{\gamma\omega}{\omega_q} [n(\omega, T_\gamma) + 1] [\hat{P}^{(+)}(\omega) \hat{\rho} \hat{P}^{(-)}(\omega') - \hat{P}^{(-)}(\omega') \hat{P}^{(+)}(\omega) \hat{\rho}] \\ & + \frac{\gamma\omega}{\omega_q} n(\omega, T_\gamma) [\hat{P}^{(-)}(\omega') \hat{\rho} \hat{P}^{(+)}(\omega) - \hat{\rho} \hat{P}^{(+)}(\omega) \hat{P}^{(-)}(\omega')] \\ & + \frac{\gamma\omega'}{\omega_q} [n(\omega', T_\gamma) + 1] [\hat{P}^{(+)}(\omega) \hat{\rho} \hat{P}^{(-)}(\omega') - \hat{\rho} \hat{P}^{(-)}(\omega') \hat{P}^{(+)}(\omega)] \\ & + \frac{\kappa\omega'}{\omega_c} n(\omega', T_\kappa) [\hat{A}^{(-)}(\omega') \hat{\rho} \hat{A}^{(+)}(\omega) - \hat{A}^{(+)}(\omega) \hat{A}^{(-)}(\omega') \hat{\rho}] \\ & + \frac{\kappa\omega}{\omega_c} [n(\omega, T_\kappa) + 1] [\hat{A}^{(+)}(\omega) \hat{\rho} \hat{A}^{(-)}(\omega') - \hat{A}^{(-)}(\omega') \hat{A}^{(+)}(\omega) \hat{\rho}] \\ & + \frac{\kappa\omega}{\omega_c} n(\omega, T_\kappa) [\hat{A}^{(-)}(\omega') \hat{\rho} \hat{A}^{(+)}(\omega) - \hat{\rho} \hat{A}^{(+)}(\omega) \hat{A}^{(-)}(\omega')] \\ & \left. + \frac{\kappa\omega'}{\omega_c} [n(\omega', T_\kappa) + 1] [\hat{A}^{(+)}(\omega) \hat{\rho} \hat{A}^{(-)}(\omega') - \hat{\rho} \hat{A}^{(-)}(\omega') \hat{A}^{(+)}(\omega)] \right\}. \quad (35) \end{aligned}$$

Here κ and γ are the qubit and cavity damping rates, respectively, $\hat{A}^{(+)}$ and $\hat{A}^{(-)}$ are the positive- and negative-frequency dressed cavity operators ($\hat{s}_i = \hat{a}$), and $\hat{P}^{(+)}$ and $\hat{P}^{(-)}$ are the positive- and negative-frequency dressed qubit operators ($\hat{s}_i = \hat{\sigma}_-$). We neglected the very small pure dephasing term in the dissipator [see Eq. (21)] **and we did not apply any filtering procedure.**

Figure 2 displays the dynamics of the mean cavity photon number $\langle \hat{A}^{(-)} \hat{A}^{(+)} \rangle$ (a) and of the zero-delay two-photon correlation function $G_A^{(2)}(t, t) = \langle \hat{A}^{(-)}(t) \hat{A}^{(-)}(t) \hat{A}^{(+)}(t) \hat{A}^{(+)}(t) \rangle$ (b) after the arrival of a Gaussian π -pulse, evaluated for different temperatures and starting the dynamics with the system in its ground state. We used $T_\gamma/\omega_c = T_\kappa/\omega_c$ and the decoherence rates $\gamma/\omega_c = \kappa/\omega_c = 3.75 \times 10^{-4}$. Note that the output photon flux is proportional to $\langle \hat{A}^{(-)} \hat{A}^{(+)} \rangle$. At $T = 0$ our approach reproduces the two-photon vacuum Rabi




Figure 2. **Dynamics of anomalous two-photon vacuum Rabi oscillations.** Results obtained using the generalized-master-equation approach, varying the temperature of both subsystems. (a) Time evolution of the mean cavity photon number $\langle \hat{A}^{(-)} \hat{A}^{(+)} \rangle$ after the arrival of a Gaussian π -pulse to the qubit. The system starts in the ground state. (b) Two-photon correlation function for the cavity, obtained with the same parameters and conditions. After the arrival of the pulse, independent of the temperature of the reservoirs, the system undergoes vacuum Rabi oscillations showing the reversible exchange of photon pairs between the qubit and the resonator. However, when raising the temperature, due to the increasing decoherence, the oscillations become more damped and the correlation function reaches higher stationary values due to larger incoherent, thermal contributions. Note that the second and the fifth dips are shallower because of some spurious effects generated by other transitions excited by the coherent pulse. All parameters for the simulations are given in the text. Ω_{eff} on the x axis indicates the effective resonant coupling.

oscillations shown in Ref. [27]. Here we study the influence of non-zero temperature on the this anomalous atom-cavity energy exchange. Increasing the temperature, the oscillations become more damped and the energy exchange becomes less effective. This effect is even more pronounced for the two-photon correlation $G_A^{(2)}(t, t)$, which displays a stronger thermal sensitivity. These results help to set a limit on the system temperature for the observation of two-photon vacuum Rabi oscillations.

In order to further show the impact of the generalized approach presented in this paper on the dissipative dynamics of cavity-QED systems in the USC regime, we compare the numerical results obtained with the generalized dressed master equation with those obtained using the standard dressed approach of Ref. [21].

Figure 3 shows the mean cavity photon number (a) and the two-photon correlation function (b) evaluated using the generalized dressed master equation (red solid curves) and the standard dressed master equation [21] (black dashed curves), calculated with the atom and cavity reservoirs at temperature $T/\omega_c = 0.75$. Both approaches show the system undergoing multiphoton Rabi oscillations and the signals reaching the same stationary values, corresponding to the equilibrium thermal populations.

We observe that the standard approach overestimates decoherence effects. In the dispersive regime of cavity-QED, pairs of photon-like transitions partially overlap, reducing decoherence effects during the time evolution. This effect is completely neglected in the standard dressed master equation. Further calculations, not shown here, indicate that these discrepancies increase with temperature. These effects lead to an overestimation of the coherence losses of the system, which also can be seen in the behaviour of the two-photon correlation function [Fig. 3(b)].

It is also important to note that the generalized master equation is able to overcome another limit of validity of the standard dressed approach. As reported in Ref. [21], the standard dressed master equation breaks down in the limit of high excitation numbers, where more transitions might accidentally have the same frequency. The generalized master equation can handle such degenerate transitions well.




Figure 3. Comparison between the results obtained using the generalized-master-equation approach (red solid curves) and the standard dressed master equation (black dashed curves). (a) Time evolution of the mean cavity photon number $\langle \hat{A}^{(-)} \hat{A}^{(+)} \rangle$ at temperature $T/\omega_c = 0.75$ with all other parameters the same as in Fig. 2. (b) Two-photon correlation functions, obtained with the same parameters and conditions. After the arrival of the pulse, both approaches show the system undergoing two-photon Rabi oscillations and relaxing to thermal equilibrium. However, with the standard dressed master equation, the coherence losses are slightly overestimated (because of the post-trace RWA), so the oscillations are more damped and the stationary value is reached sooner.

B. Cavity optomechanics beyond the RWA

1. The full optomechanical Hamiltonian

In Sec. III A, we demonstrated that our generalized approach is able to correctly describe systems with quasi-harmonic spectra. In this section, we explore a mixed harmonic-anharmonic behaviour, considering a simple optomechanical system [87], where a single cavity mode of frequency ω_c is coupled by radiation pressure to a single mechanical mode of a mirror vibrating at frequency ω_m .

Denoting the mechanical bosonic operators \hat{b} , \hat{b}^\dagger and the cavity bosonic operators \hat{a} , \hat{a}^\dagger , the system Hamiltonian can be written as [92]

$$\hat{H}_S = \hat{H}_0 + \hat{V}_{\text{om}} + \hat{V}_{\text{DCE}}, \quad (36)$$

where

$$\hat{H}_0 = \omega_c \hat{a}^\dagger \hat{a} + \omega_m \hat{b}^\dagger \hat{b} \quad (37)$$

is the unperturbed Hamiltonian,

$$\hat{V}_{\text{om}} = g \hat{a}^\dagger \hat{a} (\hat{b} + \hat{b}^\dagger) \quad (38)$$

is the standard optomechanical interaction Hamiltonian, and

$$\hat{V}_{\text{DCE}} = \frac{g}{2} (\hat{a}^2 + \hat{a}^{\dagger 2}) (\hat{b} + \hat{b}^\dagger) \quad (39)$$

describes the emission of photon pairs induced by the mechanical motion predicted by the DCE [89, 93, 94]. When treating most optomechanics experiments until now, \hat{V}_{DCE} has been neglected. This is a very good approximation when the mechanical frequency is much smaller than the cavity frequency (which is the most common experimental situation), because \hat{V}_{DCE} connects bare states with an energy difference $2\omega_c \pm \omega_m$ which then is much larger than the coupling strength g . With this approximation, the resulting Hamiltonian, $\hat{H}_0 + \hat{V}_{\text{om}}$, conserves the number of photons and can be analytically diagonalized. However, when considering ultra-high-frequency mechanical oscillators, with resonance frequencies in the GHz spectral range, coupled to a microwave resonator, \hat{V}_{DCE} , **which does not conserve the photon number**, cannot be neglected any more [87].

As shown in Ref. [87], such a system displays an energy level spectrum with a ladder of avoided level crossings arising from the coherent coupling induced by \hat{V}_{DCE} between the

states $|n, k_n\rangle$ and $|n + 2, (k - q)_{n+2}\rangle$, occurring when the energies of the initial and final states coincide ($2\omega_c \simeq q\omega_m$). Here the first number in the ket denotes photon number and the second denotes phonon number (with the photon number as a subscript since the photons displace the mechanical Fock state). For example, with $q = 1$, we have the standard resonance condition for the DCE ($2\omega_c \simeq \omega_m$ [95]), in which case \hat{V}_{DCE} gives rise to a resonant coupling between the states $|0, k\rangle$ and $|2, (k - 1)_2\rangle$ with $k \geq 1$, converting a phonon into a photon pair.

When \hat{V}_{DCE} is taken into account, the system Hamiltonian does not conserve the number of photons (the phonon number is not conserved even in the standard optomechanical Hamiltonian). For example, the ground state of \hat{H}_S contains photons, i.e., $\langle E_0 | \hat{a}^\dagger \hat{a} | E_0 \rangle \neq 0$. Therefore, in analogy to USC cavity QED, a careful treatment of dissipation and input-output theory is required. If the standard photon and phonon operators were used to describe the interaction with the outside world, unphysical effects would arise.

2. The impact of temperature on the dynamical Casimir effect

It has been shown [87] that this system can be used to demonstrate the conversion of mechanical energy into photon pairs (DCE). The calculations in Ref. [87] were performed using a dressed master equation without the post-trace RWA, developed only for the case of zero-temperature reservoirs. Here we instead apply the generalized master equation presented in Sec. IIC, in order to study the influence of temperature on the energy conversion from phonons to photons.

For our numerical calculation we consider a normalized optomechanical coupling $g/\omega_m = 0.1$, a mechanical damping rate $\gamma/\omega_m = 0.05$, and a cavity damping rate $\kappa = \gamma/2$. We focus on the avoided level crossing between the states $|0, 2\rangle$ and $|2, 0_2\rangle$ at $\omega_m \simeq \omega_c$. We consider the resonant condition, corresponding to the minimum level splitting: $\omega_c/\omega_m = 1.016$.

As in Ref. [87], we consider a continuous coherent drive of the mechanical oscillator,

$$\hat{H}_d = \Omega (\hat{b} e^{-i\omega_m t} + \hat{b}^\dagger e^{i\omega_m t}), \quad (40)$$

with frequency resonant with the oscillating mirror and amplitude $\Omega = \gamma/2$. The dynamics giving rise to the DCE is then described by the filtered generalized master equation ($\Lambda = 10\gamma$)

$$\dot{\hat{\rho}} = -i [\hat{H}_S + \hat{H}_d, \hat{\rho}] + \mathcal{L}_{\text{gme}}^{\text{filt}} \hat{\rho}, \quad (41)$$

where the Liouvillian superoperator can be written as

$$\begin{aligned}
\mathcal{L}_{\text{gme}}^{\text{filter}} \hat{\rho} = & \sum_{(\omega, \omega') > 0} \frac{1}{2} \left\{ \gamma n(\omega', T_\gamma) [\hat{B}^{(-)}(\omega') \hat{\rho} \hat{B}^{(+)}(\omega) - \hat{B}^{(+)}(\omega) \hat{B}^{(-)}(\omega') \hat{\rho}] \right. \\
& + \gamma [n(\omega, T_\gamma) + 1] [\hat{B}^{(+)}(\omega) \hat{\rho} \hat{B}^{(-)}(\omega') - \hat{B}^{(-)}(\omega') \hat{B}^{(+)}(\omega) \hat{\rho}] \\
& + \gamma n(\omega, T_\gamma) [\hat{B}^{(-)}(\omega') \hat{\rho} \hat{B}^{(+)}(\omega) - \hat{\rho} \hat{B}^{(+)}(\omega) \hat{B}^{(-)}(\omega')] \\
& + \gamma [n(\omega', T_\gamma) + 1] [\hat{B}^{(+)}(\omega) \hat{\rho} \hat{B}^{(-)}(\omega') - \hat{\rho} \hat{B}^{(-)}(\omega') \hat{B}^{(+)}(\omega)] \\
& + \kappa n(\omega', T_\kappa) [\hat{A}^{(-)}(\omega') \hat{\rho} \hat{A}^{(+)}(\omega) - \hat{A}^{(+)}(\omega) \hat{A}^{(-)}(\omega') \hat{\rho}] \\
& + \kappa [n(\omega, T_\kappa) + 1] [\hat{A}^{(+)}(\omega) \hat{\rho} \hat{A}^{(-)}(\omega') - \hat{A}^{(-)}(\omega') \hat{A}^{(+)}(\omega) \hat{\rho}] \\
& + \kappa n(\omega, T_\kappa) [\hat{A}^{(-)}(\omega') \hat{\rho} \hat{A}^{(+)}(\omega) - \hat{\rho} \hat{A}^{(+)}(\omega) \hat{A}^{(-)}(\omega')] \\
& \left. + \kappa [n(\omega', T_\kappa) + 1] [\hat{A}^{(+)}(\omega) \hat{\rho} \hat{A}^{(-)}(\omega') - \hat{\rho} \hat{A}^{(-)}(\omega') \hat{A}^{(+)}(\omega)] \right\} \times F(\omega, \omega'), \quad (42)
\end{aligned}$$

where $\hat{A}^{(+)}$ and $\hat{A}^{(-)}$ are the positive- and negative-frequency dressed cavity operators ($\hat{s}_i = \hat{a}$), and $\hat{B}^{(+)}$ and $\hat{B}^{(-)}$ are the positive- and negative-frequency dressed mechanical operators ($\hat{s}_i = \hat{b}$).

In Fig. 4, we show the photonic and phononic populations, $\langle \hat{A}^{(-)} \hat{A}^{(+)} \rangle$ and $\langle \hat{B}^{(-)} \hat{B}^{(+)} \rangle$, and the relative two-photon and two-phonon correlation functions,

$$g_A^{(2)}(t, t) = \frac{\langle \hat{A}^{(-)}(t) \hat{A}^{(-)}(t) \hat{A}^{(+)}(t) \hat{A}^{(+)}(t) \rangle}{\langle \hat{A}^{(-)}(t) \hat{A}^{(+)}(t) \rangle^2}, \quad (43)$$

$$g_B^{(2)}(t, t) = \frac{\langle \hat{B}^{(-)}(t) \hat{B}^{(-)}(t) \hat{B}^{(+)}(t) \hat{B}^{(+)}(t) \rangle}{\langle \hat{B}^{(-)}(t) \hat{B}^{(+)}(t) \rangle^2}. \quad (44)$$

Figures 4(a, b) display the results of calculations done with zero-temperature reservoirs for both subsystems and starting the dynamics from the ground state. Figures 4(c, d) display the results of calculations for reservoirs with $T_\gamma/\omega_m = T_\kappa/\omega_m = 0.5$ and with the system initially in thermal equilibrium with those reservoirs.

At $T = 0$, with the system starting in its ground state, the photonic and phononic populations start from zero and, due to the coherent pumping, reach non-zero stationary values. The photonic correlation function $g_A^{(2)}(t, t)$ is initially much higher than two, suggesting photon-pair emission. As time goes on, $g_A^{(2)}(t, t)$ decreases significantly due to losses which affect the photon-photon correlations, and also due to the increase of the mean photon number (note that $g_A^{(2)}(t, t)$, owing to the squared denominator, is an intensity-dependent quantity). The mechanical correlation function $g_B^{(2)}(t, t)$, on the contrary, has an almost constant value ($g_B^{(2)}(t, t) \approx 1$), showing that the mechanical system is mainly in the coherent state produced by the pumping.




Figure 4. Results for the DCE at different temperatures, obtained using the generalized-master-equation approach. (a, b) System dynamics for $\omega_c \simeq \omega_m$, under coherent mechanical pumping, in perfect cooling conditions $T_\gamma = T_\kappa = 0$, starting the dynamics from the ground state. (c, d) The same, but with $T_\gamma/\omega_m = T_\kappa/\omega_m = 0.5$ and the initial state being the thermal state with $T/\omega_m = 0.5$. The blue dashed curves show the mean phonon number $\langle \hat{B}^{(-)} \hat{B}^{(+)} \rangle$ in (a, c) and the phonon-phonon correlation function $g_B^{(2)}(t, t)$ in (b, d). The red solid curves describe the mean cavity photon number $\langle \hat{A}^{(-)} \hat{A}^{(+)} \rangle$ in (a, c) and the zero-delay normalized photon-photon correlation function $g_A^{(2)}(t, t)$ in (b, d). All parameters for the simulations are given in the text.

For reservoirs with non-zero temperature, the phonon and photon populations, starting from their thermal-equilibrium values, equilibrate to lower steady-state values. This reduction of both populations originates from the increase of the decay rate of the coherent contributions with increasing temperature. We also note that the difference between the two steady-state values is reduced at higher temperatures, due to the thermal contributions. At $T \neq 0$, a fraction of the observed photons, as expected, does not come from the mechanical-to-optical energy conversion, but, trivially, from the photonic thermal reservoir.

This picture is confirmed by comparing the dynamics of the higher-order correlation functions [Figs. 4(c, d)]. Specifically, at higher temperature, we observe a strong decrease of

$g_A^{(2)}(t, t)$, showing that a reduced fraction of photons is emitted in pairs. However, the photon-photon correlation functions remains, even in the steady state, *higher* than the thermal value $g_A^{(2)}(t, t) = 2$. The phonon-phonon correlation starts from a value $\simeq 2$ corresponding to the initial incoherent thermal state and, as time goes on, decays to a stationary value higher than one due to the incoherent thermal excitations provided by the interaction with the thermal reservoirs.

Furthermore, in Fig. 4(d), the photon-photon and the phonon-phonon correlation functions do not start from the same initial value. This effect is due to the \hat{V}_{DCE} term which, owing to its non-bilinear form, modifies the thermodynamic equilibrium of the initial state of the system. The \hat{V}_{DCE} contribution leads to a separation of the correlation-function values with size proportional to the temperature. This separation thus vanishes trivially for $T = 0$, when the \hat{V}_{DCE} term becomes negligible.

The results obtained clearly show that the generalized dressed master equation provided here is able to describe dissipation in hybrid quantum systems with coexisting coherent phases (provided, e.g., by means of a continuous drive) and incoherent phases (provided, e.g., by thermal reservoirs or thermal-like pumping). The behaviour of the one- and two-photon correlation functions show that signatures of the DCE can be observed even in the presence of a non-negligible amount of thermal noise. It thus demonstrates that this effect can be observed in a real experimental set-up, where perfect cooling conditions cannot be reached. Although the number of Casimir photon pairs produced depends on the thermal noise injected into the system, our results here show that the DCE remains detectable even at relatively high temperatures.

3. Comparison to other approaches

As already mentioned in the introduction, and demonstrated in Ref. [21], the use of a master equation with a dissipator not taking into account the interaction between the subsystems can lead to unphysical results. Hu *et al.* derived [75] a dressed master equation specifically developed to describe dissipation in optomechanical systems characterized by the standard optomechanical Hamiltonian, $\hat{H}_S = \hat{H}_0 + \hat{V}_{\text{om}}$, in the USC regime. Here we show that this master equation *fails* when considering the complete optomechanical Hamiltonian $\hat{H}_S = \hat{H}_0 + \hat{V}_{\text{om}} + \hat{V}_{\text{DCE}}$.




Figure 5. State populations obtained using the master-equation approach of Ref. [75]. The red (blue) solid curve shows the time evolution of the population of the one-photon state $|1,0\rangle$ (the one-phonon state $|0,1\rangle$) labelled $\rho_{22}(t)$ [$\rho_{11}(t)$] under perfect cooling conditions $T_\gamma = T_\kappa = 0$ and without any pumping. The initial state is the ground state $|0,0\rangle$. All other parameters are the same as in Fig. 4. In these conditions, without any external driving or thermal excitations, the system is expected to remain in the ground state. However, the plot clearly shows a non-zero population in both the one-photon and one-phonon states. This indicates that this approach is *not* able to correctly describe optomechanical systems when the \hat{V}_{DCE} contribution no longer can be neglected.

In Fig. 5, we display results obtained describing the dynamics of our optomechanical system in perfect cooling conditions, without any pumping, with the master equation provided in Ref. [75], including the \hat{V}_{DCE} term as a perturbation in the dynamics. In these conditions, evaluating the dynamics with the system initially in the ground state (an eigenstate of the system), zero population is expected in the states with one photon, $|1,0\rangle$, and one phonon, $|0,1\rangle$. However, Fig. 5 clearly shows non-zero populations. This anomalous effect occurs because, due to the additional \hat{V}_{DCE} term, the number of photons is no longer conserved and consequently the eigenstates of the Hamiltonian changes. In this case, the master equation provided in Ref. [75] does not describe interactions between subsystems and reservoirs in terms of the correct eigenstates, which leads to an unphysical evolution of the initial ground

state. This result shows once more the importance of expressing the system operators in the basis of the system eigenstates when describing interactions with reservoirs to derive a correct master equation.

IV. CONCLUSIONS

We have presented a generalized dressed master equation, valid for arbitrary open hybrid quantum system interacting with thermal reservoirs and for arbitrary strength of the coupling between the components of the hybrid system. Our approach was derived within the Born-Markov approximation, including the pure dephasing terms and without performing the usual post-trace RWA. Therefore, our approach is able to handle dynamics in systems with both harmonic, quasi-harmonic, and anharmonic transitions. Moreover, this approach is not limited to systems displaying parity symmetry. Unfortunately the dissipator obtained includes rapidly oscillating terms that can cause numerical instabilities. In order to fix this problem, we introduced a filtering procedure which eliminates the fast-oscillating terms which do not contribute to the coarse-grained dynamics. This filtering has the added benefit of reducing computation times.

We applied our generalized approach to study the influence of temperature on multiphoton vacuum Rabi oscillation in a circuit-QED system in the dispersive regime. We compared our results with those obtained using the dressed master equation of Ref. [21]. We found that both approaches describe multiphoton Rabi oscillations and reach the same stationary state (the thermal equilibrium). However, the standard master equation overestimates decoherence effects since it does not take into account the partial overlap of photon-like transitions, which reduces the decoherence during the time evolution.

We also studied the influence of temperature on the conversion of mechanical energy into photon pairs (DCE) in an optomechanical system, recently described in Ref. [87] for zero-temperature reservoirs. In this case, we showed that the DCE can be observed also in the presence of a significant amount of thermal noise.

Finally, we demonstrated that the master-equation approach provided in Ref. [75] for optomechanical systems with ultrastrong coupling fails when considering the full optomechanical Hamiltonian including the \hat{V}_{DCE} term. Specifically, under these conditions, the master equation provided in Ref. [75] does not describe interactions between the components

and reservoirs correctly in terms of transitions between eigenstates of the hybrid system. Because of this shortcoming, that approach leads to an unphysical evolution of the initial ground state to excited states even at zero temperature and without any external pumping. This example clearly shows that the general master-equation approach provided here is necessary to describe dissipation of general open hybrid quantum systems interacting with thermal reservoirs.

V. ACKNOWLEDGEMENTS

F.N. is supported in part by the: MURI Center for Dynamic Magneto-Optics via the Air Force Office of Scientific Research (AFOSR) (FA9550-14-1-0040), Army Research Office (ARO) (Grant No. 73315PH), Asian Office of Aerospace Research and Development (AOARD) (Grant No. FA2386-18-1-4045), Japan Science and Technology Agency (JST) (the ImPACT program and CREST Grant No. JPMJCR1676), Japan Society for the Promotion of Science (JSPS) (JSPS-RFBR Grant No. 17-52-50023, and JSPS-FWO Grant No. VS.059.18N), RIKEN-AIST Challenge Research Fund, and the John Templeton Foundation.

Appendix A: Derivation of the dressed master equation

Starting from the system-bath Hamiltonian in Eq. (18), and following the standard procedure [5], i.e., performing the second-order Born approximation, the Markov approximation, and considering reservoirs with a continuum of frequencies, we obtain

$$\begin{aligned} \dot{\hat{\rho}}(t) = & \int_0^t dt' \left[\hat{S}_i(t') \hat{\rho}(t') \hat{S}_i(t) - \hat{S}_i(t) \hat{S}_i(t') \hat{\rho}(t') \right] \langle \hat{B}_i^\dagger(t) \hat{B}_i(t') \rangle \\ & + \int_0^t dt' \left[\hat{S}_i(t) \hat{\rho}(t') \hat{S}_i(t') - \hat{\rho}(t') \hat{S}_i(t') \hat{S}_i(t) \right] \langle \hat{B}_i^\dagger(t') \hat{B}_i(t) \rangle \\ & + \int_0^t dt' \left[\hat{S}_i(t') \hat{\rho}(t) \hat{S}_i(t) - \hat{S}_i(t) \hat{S}_i(t') \hat{\rho}(t) \right] \langle \hat{B}_i(t) \hat{B}_i^\dagger(t') \rangle \\ & + \int_0^t dt' \left[\hat{S}_i(t) \hat{\rho}(t') \hat{S}_i(t') - \hat{\rho}(t') \hat{S}_i(t') \hat{S}_i(t) \right] \langle \hat{B}_i(t') \hat{B}_i^\dagger(t) \rangle, \end{aligned} \quad (\text{A1})$$

where $\langle \hat{B}_i^\dagger(t) \hat{B}_i(t') \rangle$ and $\langle \hat{B}_i^\dagger(t') \hat{B}_i(t) \rangle$ are the reservoir correlation functions

$$\langle \hat{B}_i^\dagger(t) \hat{B}_i(t') \rangle = \int_0^\infty d\nu g_i(\nu) |\alpha_i(\nu)|^2 n(\nu, T_i) e^{\nu(t-t')}, \quad (\text{A2})$$

$$\langle \hat{B}_i(t) \hat{B}_i^\dagger(t') \rangle = \int_0^\infty d\nu g_i(\nu) |\alpha_i(\nu)|^2 [n(\nu, T_i) + 1] e^{-\nu(t-t')}, \quad (\text{A3})$$

with $g(\nu)$ being the reservoir density of states and $\alpha(\nu)$ the system-reservoir coupling strength. Substituting Eqs. (A2) and (A3) into Eq. (A1), and performing the change of variable $\tau = t - t'$, we obtain

$$\dot{\hat{\rho}}(t) = \sum_i \sum_{\omega, \omega'} \left[\hat{A}_{\omega, \omega'}^i(t) + \hat{B}_{\omega, \omega'}^i(t) + \hat{C}_{\omega, \omega'}^i(t) + \hat{D}_{\omega, \omega'}^i(t) \right], \quad (\text{A4})$$

where

$$\begin{aligned} \hat{A}_{\omega, \omega'}^i(t) &= \int_0^t d\tau e^{-i(\omega + \omega')t} e^{i\omega\tau} \left[\hat{S}_i(\omega') \hat{\rho}(t) \hat{S}_i(\omega) - \hat{S}_i(\omega) \hat{S}_i(\omega') \hat{\rho}(t) \right] \\ &\quad \times \int_0^\infty d\nu g_i(\nu) |\alpha_i(\nu)|^2 n(\nu, T_i) e^{i\nu\tau}, \\ \hat{B}_{\omega, \omega'}^i(t) &= \int_0^t d\tau e^{-i(\omega + \omega')t} e^{i\omega\tau} \left[\hat{S}_i(\omega') \hat{\rho}(t) \hat{S}_i(\omega) - \hat{\rho}(t) \hat{S}_i(\omega) \hat{S}_i(\omega') \right] \\ &\quad \times \int_0^\infty d\nu g_i(\nu) |\alpha_i(\nu)|^2 n(\nu, T_i) e^{-i\nu\tau}, \\ \hat{C}_{\omega, \omega'}^i(t) &= \int_0^t d\tau e^{-i(\omega + \omega')t} e^{i\omega\tau} \left[\hat{S}_i(\omega) \hat{\rho}(t) \hat{S}_i(\omega') - \hat{S}_i(\omega') \hat{S}_i(\omega) \hat{\rho}(t) \right] \\ &\quad \times \int_0^\infty d\nu g_i(\nu) |\alpha_i(\nu)|^2 [n(\nu, T_i) + 1] e^{-i\nu\tau}, \\ \hat{D}_{\omega, \omega'}^i(t) &= \int_0^t d\tau e^{-i(\omega + \omega')t} e^{i\omega\tau} \left[\hat{S}_i(\omega) \hat{\rho}(t) \hat{S}_i(\omega') - \hat{\rho}(t) \hat{S}_i(\omega') \hat{S}_i(\omega) \right] \\ &\quad \times \int_0^\infty d\nu g_i(\nu) |\alpha_i(\nu)|^2 [n(\nu, T_i) + 1] e^{i\nu\tau}, \end{aligned} \quad (\text{A5})$$

Assuming that the integrands decay on a much shorter time scale than that of the reservoir correlation functions, we can extend the τ integration to infinity. Evaluating both the integrals without performing any approximation except for the Born-Markov approximation, the master equation in the Schrödinger picture can be written

$$\dot{\hat{\rho}} = -i[\hat{H}_S, \hat{\rho}] + \mathcal{L}_{\text{gme}} \hat{\rho}, \quad (\text{A6})$$

with the Lindbladian superoperator that in the most general form can be written as,

$$\begin{aligned} \mathcal{L}_{\text{gme}} \hat{\rho} &= \frac{1}{2} \sum_i \sum_{\omega, \omega'} \left\{ \Gamma_i(-\omega') n(-\omega', T_i) \left[\hat{S}_i(\omega') \hat{\rho}(t) \hat{S}_i(\omega) - \hat{S}_i(\omega) \hat{S}_i(\omega') \hat{\rho}(t) \right] \right. \\ &\quad + \Gamma_i(\omega) n(\omega, T_i) \left[\hat{S}_i(\omega') \hat{\rho}(t) \hat{S}_i(\omega) - \hat{\rho}(t) \hat{S}_i(\omega) \hat{S}_i(\omega') \right] \\ &\quad + \Gamma_i(\omega) [n(\omega, T_i) + 1] \left[\hat{S}_i(\omega) \hat{\rho}(t) \hat{S}_i(\omega') - \hat{S}_i(\omega') \hat{S}_i(\omega) \hat{\rho}(t) \right] \\ &\quad \left. + \Gamma_i(-\omega') [n(-\omega', T_i) + 1] \left[\hat{S}_i(\omega) \hat{\rho}(t) \hat{S}_i(\omega') - \hat{\rho}(t) \hat{S}_i(\omega') \hat{S}_i(\omega) \right] \right\}. \end{aligned} \quad (\text{A7})$$

Both $\Gamma_i(\omega)$ and $n(\omega, T_i)$ are non zero only for $\omega > 0$ thus, using the definitions in Eq. (17), Eq. (A7) can be written as

$$\begin{aligned}
\mathcal{L}_{\text{gme}}\hat{\rho} = & \frac{1}{2} \sum_i \sum_{(\omega, \omega') > 0} \left\{ \Gamma_i(\omega') n(\omega', T_i) [\hat{S}_i^{(-)}(\omega') \hat{\rho}(t) \hat{S}_i^{(+)}(\omega) - \hat{S}_i^{(+)}(\omega) \hat{S}_i^{(-)}(\omega') \hat{\rho}(t)] \right. \\
& + \Gamma_i(\omega) n(\omega, T_i) [\hat{S}_i^{(-)}(\omega') \hat{\rho}(t) \hat{S}_i^{(+)}(\omega) - \hat{\rho}(t) \hat{S}_i^{(+)}(\omega) \hat{S}_i^{(-)}(\omega')] \\
& + \Gamma_i(\omega) [n(\omega, T_i) + 1] [\hat{S}_i^{(+)}(\omega) \hat{\rho}(t) \hat{S}_i^{(-)}(\omega') - \hat{S}_i^{(-)}(\omega') \hat{S}_i^{(+)}(\omega) \hat{\rho}(t)] \\
& + \Gamma_i(\omega') [n(\omega', T_i) + 1] [\hat{S}_i^{(+)}(\omega) \hat{\rho}(t) \hat{S}_i^{(-)}(\omega') - \hat{\rho}(t) \hat{S}_i^{(-)}(\omega') \hat{S}_i^{(+)}(\omega)] \\
& + \Gamma_i(\omega') n(\omega', T_i) [\hat{S}_i^{(-)}(\omega') \hat{\rho}(t) \hat{S}_i^{(-)}(\omega) - \hat{S}_i^{(-)}(\omega) \hat{S}_i^{(-)}(\omega') \hat{\rho}(t)] \\
& + \Gamma_i(\omega') [n(\omega', T_i) + 1] [\hat{S}_i^{(-)}(\omega) \hat{\rho}(t) \hat{S}_i^{(-)}(\omega') - \hat{\rho}(t) \hat{S}_i^{(-)}(\omega') \hat{S}_i^{(-)}(\omega)] \\
& + \Gamma_i(\omega) n(\omega, T_i) [\hat{S}_i^{(+)}(\omega') \hat{\rho}(t) \hat{S}_i^{(+)}(\omega) - \hat{\rho}(t) \hat{S}_i^{(+)}(\omega) \hat{S}_i^{(+)}(\omega')] \\
& + \Gamma_i(\omega) [n(\omega, T_i) + 1] [\hat{S}_i^{(+)}(\omega) \hat{\rho}(t) \hat{S}_i^{(+)}(\omega') - \hat{S}_i^{(+)}(\omega') \hat{S}_i^{(+)}(\omega) \hat{\rho}(t)] \\
& + \Gamma_i(\omega') n(\omega', T_i) [\hat{S}_i^{(-)}(\omega') \hat{\rho}(t) \hat{S}_i^{(0)} - \hat{S}_i^{(0)} \hat{S}_i^{(-)}(\omega') \hat{\rho}(t)] \\
& + \Gamma_i(\omega') [n(\omega', T_i) + 1] [\hat{S}_i^{(0)} \hat{\rho}(t) \hat{S}_i^{(-)}(\omega') - \hat{\rho}(t) \hat{S}_i^{(-)}(\omega') \hat{S}_i^{(0)}] \\
& + \Gamma_i(\omega) n(\omega, T_i) [\hat{S}_i^{(0)} \hat{\rho}(t) \hat{S}_i^{(+)}(\omega) - \hat{\rho}(t) \hat{S}_i^{(+)}(\omega) \hat{S}_i^{(0)}] \\
& + \Gamma_i(\omega) [n(\omega, T_i) + 1] [\hat{S}_i^{(+)}(\omega) \hat{\rho}(t) \hat{S}_i^{(0)} - \hat{S}_i^{(0)} \hat{S}_i^{(+)}(\omega) \hat{\rho}(t)] \\
& + \Omega_i^+(T_i) [\hat{S}_i^{(0)} \hat{\rho}(t) \hat{S}_i^{(0)} - \hat{S}_i^{(0)} \hat{S}_i^{(0)} \hat{\rho}(t)] \\
& + \Omega_i'^+(T_i) [\hat{S}_i^{(0)} \hat{\rho}(t) \hat{S}_i^{(0)} - \hat{\rho}(t) \hat{S}_i^{(0)}(\omega') \hat{S}_i^{(0)}] \\
& + \Omega_i^-(T_i) [\hat{S}_i^{(0)} \hat{\rho}(t) \hat{S}_i^{(0)} - \hat{\rho}(t) \hat{S}_i^{(0)} \hat{S}_i^{(0)}] \\
& + \Omega_i'^-(T_i) [\hat{S}_i^{(0)} \hat{\rho}(t) \hat{S}_i^{(0)} - \hat{S}_i^{(0)} \hat{S}_i^{(0)} \hat{\rho}(t)]
\end{aligned} \tag{A8}$$

with thermal populations

$$n(\omega, T_i) = [\exp \{\omega/T_i\} - 1]^{-1}, \tag{A9}$$

damping rates

$$\Gamma_i(\omega) = 2\pi g_i(\omega) |\alpha_i(\omega)|^2. \tag{A10}$$

and pure dephasing damping rates

$$\Omega_i^{\pm}(T_i) = \int_0^t d\tau \int_0^\infty d\nu g_i(\nu) |\alpha_i(\nu)|^2 [n(\nu, T_i) + 1] e^{\pm i\nu\tau}, \tag{A11}$$

$$\Omega_i^{\pm}(T_i) = \int_0^t d\tau \int_0^\infty d\nu g_i(\nu) |\alpha_i(\nu)|^2 n(\nu, T_i) e^{\pm i\nu\tau}. \tag{A12}$$

Specifically, the terms in the first four lines of Eq. (A8) oscillate at frequencies $\pm(\omega - \omega')$. If $(\omega - \omega')$ is significantly larger than the damping rates Γ_i of the system, these terms

provide negligible contributions when integrating the master equation. In the generalized approach, these terms are then eliminated by the numerical filtering. The terms in the next four lines of Eq. (A8) oscillate at $\pm(\omega' + \omega)$. These terms are clearly rapidly oscillating and thus provide negligible contributions. The terms in the following four lines, oscillating at $+\omega$, $-\omega'$, are fast-oscillating when considering systems displaying well-separated energy levels with $\omega \gg \Gamma_i$ and, in these cases, can be neglected. Finally, the term in the last four lines arise from degenerate transitions and describe pure dephasing. The contribution of these terms becomes negligible at very low temperatures in the particular case of Ohmic baths. Furthermore, it is important to note that, applying the post-trace RWA without considering any parity symmetry of the system, Eq. (A7) can be rewritten in a form equal to the standard dressed master equation as in Ref. [21], with a few additional terms provided by the zero-frequency operators $\hat{S}_i^{(0)} \neq 0$.

-
- [1] M. Brune, E. Hagley, J. Dreyer, X. Maître, A. Maali, C. Wunderlich, J. M. Raimond, and S. Haroche, “Observing the progressive decoherence of the “meter” in a quantum measurement,” *Phys. Rev. Lett.* **77**, 4887–4890 (1996).
 - [2] M. Hofheinz, H. Wang, M. Ansmann, R. C. Bialczak, E. Lucero, M. Neeley, A. D. O’Connell, D. Sank, J. Wenner, J. M. Martinis, and A. N. Cleland, “Synthesizing arbitrary quantum states in a superconducting resonator,” *Nature* **459**, 546 (2009).
 - [3] X. Gu, A. F. Kockum, A. Miranowicz, Y.-X. Liu, and F. Nori, “Microwave photonics with superconducting quantum circuits,” *Phys. Rep.* **718–719**, 1–102 (2017), [arXiv:1707.02046](#).
 - [4] S. N. A. Duffus, V. M. Dwyer, and M. J. Everitt, “Open quantum systems, effective Hamiltonians, and device characterization,” *Phys. Rev. B* **96**, 134520 (2017), [arXiv:1707.00593](#).
 - [5] H. P. Breuer and F. Petruccione, *The Theory of Open Quantum Systems* (Oxford University Press, 2002).
 - [6] C. W. Gardiner and P. Zoller, *Quantum Noise*, 3rd ed. (Springer, 2004).
 - [7] M. D. Kostin, “On the Schrödinger-Langevin equation,” *The Journal of Chemical Physics* **57**, 3589–3591 (1972).
 - [8] G. W. Ford and M. Kac, “On the quantum Langevin equation,” *Journal of Statistical Physics* **46**, 803–810 (1987).

- [9] S. Portolan, O. Di Stefano, S. Savasta, F. Rossi, and R. Girlanda, “Nonequilibrium Langevin approach to quantum optics in semiconductor microcavities,” [Physical Review B - Condensed Matter and Materials Physics](#) **77**, 035433 (2008), [arXiv:0611381 \[cond-mat\]](#).
- [10] F. Haake, “Statistical Treatment of Open Systems by Generalized Master Equations,” in [Springer Tracts in Modern Physics](#), edited by G. Höhler (Springer, Berlin, Heidelberg, 1973) pp. 98–168.
- [11] N. Shammah, S. Ahmed, N. Lambert, S. De Liberato, and F. Nori, “Open quantum systems with local and collective incoherent processes: Efficient numerical simulation using permutational invariance,” [arXiv:1805.05129](#), [Phys. Rev. A](#), in press (2018).
- [12] M. Wallquist, K. Hammerer, P. Rabl, M. Lukin, and P. Zoller, “Hybrid quantum devices and quantum engineering,” [Phys. Scr.](#) **T137**, 014001 (2009).
- [13] Z.-L. Xiang, S. Ashhab, J. Q. You, and F. Nori, “Hybrid quantum circuits: Superconducting circuits interacting with other quantum systems,” [Rev. Mod. Phys.](#) **85**, 623 (2013), [arXiv:1204.2137](#).
- [14] G. Kurizki, P. Bertet, Y. Kubo, K. Mølmer, D. Petrosyan, P. Rabl, and J. Schmiedmayer, “Quantum technologies with hybrid systems,” [Proc. Natl. Acad. Sci.](#) **112**, 3866 (2015), [arXiv:1504.00158](#).
- [15] S. Haroche, “Nobel lecture: Controlling photons in a box and exploring the quantum to classical boundary,” [Rev. Mod. Phys.](#) **85**, 1083–1102 (2013).
- [16] A. F. Kockum, A. Miranowicz, S. De Liberato, S. Savasta, and F. Nori, “Ultrastrong coupling between light and matter,” (2018), [arXiv:1807.11636](#).
- [17] P. Forn-Díaz, L. Lamata, E. Rico, J. Kono, and E. Solano, “Ultrastrong coupling regimes of light-matter interaction,” (2018), [arXiv:1804.09275](#).
- [18] S. De Liberato, C. Ciuti, and I. Carusotto, “Quantum Vacuum Radiation Spectra from a Semiconductor Microcavity with a Time-Modulated Vacuum Rabi Frequency,” [Phys. Rev. Lett.](#) **98**, 103602 (2007), [arXiv:0611282 \[cond-mat\]](#).
- [19] S. Ashhab and F. Nori, “Qubit-oscillator systems in the ultrastrong-coupling regime and their potential for preparing nonclassical states,” [Phys. Rev. A](#) **81**, 042311 (2010), [arXiv:0912.4888](#).
- [20] J. Casanova, G. Romero, I. Lizuain, J. J. García-Ripoll, and E. Solano, “Deep Strong Coupling Regime of the Jaynes-Cummings Model,” [Phys. Rev. Lett.](#) **105**, 263603 (2010), [arXiv:1008.1240](#).

- [21] F. Beaudoin, J. M. Gambetta, and A. Blais, “Dissipation and ultrastrong coupling in circuit QED,” *Phys. Rev. A* **84**, 043832 (2011), [arXiv:1107.3990](#).
- [22] A. Ridolfo, M. Leib, S. Savasta, and M. J. Hartmann, “Photon Blockade in the Ultrastrong Coupling Regime,” *Phys. Rev. Lett.* **109**, 193602 (2012), [arXiv:1206.0944](#).
- [23] R. Stassi, A. Ridolfo, O. Di Stefano, M. J. Hartmann, and S. Savasta, “Spontaneous Conversion from Virtual to Real Photons in the Ultrastrong-Coupling Regime,” *Phys. Rev. Lett.* **110**, 243601 (2013), [arXiv:1210.2367](#).
- [24] L. Garziano, A. Ridolfo, R. Stassi, O. Di Stefano, and S. Savasta, “Switching on and off of ultrastrong light-matter interaction: Photon statistics of quantum vacuum radiation,” *Phys. Rev. A* **88**, 063829 (2013).
- [25] S. De Liberato, “Light-Matter Decoupling in the Deep Strong Coupling Regime: The Breakdown of the Purcell Effect,” *Phys. Rev. Lett.* **112**, 016401 (2014), [arXiv:1308.2812](#).
- [26] E. Sanchez-Burillo, D. Zueco, J. J. Garcia-Ripoll, and L. Martin-Moreno, “Scattering in the Ultrastrong Regime: Nonlinear Optics with One Photon,” *Phys. Rev. Lett.* **113**, 263604 (2014), [arXiv:1406.5779](#).
- [27] L. Garziano, R. Stassi, V. Macrì, A. F. Kockum, S. Savasta, and F. Nori, “Multiphoton quantum Rabi oscillations in ultrastrong cavity QED,” *Phys. Rev. A* **92**, 063830 (2015), [arXiv:1509.06102](#).
- [28] L. Garziano, V. Macrì, R. Stassi, O. Di Stefano, F. Nori, and S. Savasta, “One Photon Can Simultaneously Excite Two or More Atoms,” *Phys. Rev. Lett.* **117**, 043601 (2016), [arXiv:1601.00886](#).
- [29] M. Cirio, S. De Liberato, N. Lambert, and F. Nori, “Ground State Electroluminescence,” *Phys. Rev. Lett.* **116**, 113601 (2016), [arXiv:1508.05849](#).
- [30] A. F. Kockum, A. Miranowicz, V. Macrì, S. Savasta, and F. Nori, “Deterministic quantum nonlinear optics with single atoms and virtual photons,” *Phys. Rev. A* **95**, 063849 (2017), [arXiv:1701.05038](#).
- [31] A. F. Kockum, V. Macrì, L. Garziano, S. Savasta, and F. Nori, “Frequency conversion in ultrastrong cavity QED,” *Sci. Rep.* **7**, 5313 (2017), [arXiv:1701.07973](#).
- [32] R. Stassi, V. Macrì, A. F. Kockum, O. Di Stefano, A. Miranowicz, S. Savasta, and F. Nori, “Quantum nonlinear optics without photons,” *Phys. Rev. A* **96**, 023818 (2017), [arXiv:1702.00660](#).

- [33] W. Qin, A. Miranowicz, P. Li, X. Lü, J. Q. You, and F. Nori, “Exponentially enhanced light-matter interaction, cooperativities, and steady-state entanglement using parametric amplification,” *Phys. Rev. Lett.* **120**, 093601 (2018), [arXiv:1709.09555](#).
- [34] A. Ridolfo, G. Falci, F. M. D. Pellegrino, and E. Paladino, “Photon pair production by STIRAP in ultrastrongly coupled matter-radiation systems,” (2018), [arXiv:1805.07079](#).
- [35] A. A. Anappara, S. De Liberato, A. Tredicucci, C. Ciuti, G. Biasiol, L. Sorba, and F. Beltram, “Signatures of the ultrastrong light-matter coupling regime,” *Phys. Rev. B* **79**, 201303 (2009), [arXiv:0808.3720](#).
- [36] G. Günter, A. A. Anappara, J. Hees, A. Sell, G. Biasiol, L. Sorba, S. De Liberato, C. Ciuti, A. Tredicucci, A. Leitenstorfer, and R. Huber, “Sub-cycle switch-on of ultrastrong light-matter interaction,” *Nature* **458**, 178 (2009).
- [37] Y. Todorov, A. M. Andrews, R. Colombelli, S. De Liberato, C. Ciuti, P. Klang, G. Strasser, and C. Sirtori, “Ultrastrong Light-Matter Coupling Regime with Polariton Dots,” *Phys. Rev. Lett.* **105**, 196402 (2010), [arXiv:1301.1297](#).
- [38] M. Geiser, F. Castellano, G. Scalari, M. Beck, L. Nevou, and J. Faist, “Ultrastrong Coupling Regime and Plasmon Polaritons in Parabolic Semiconductor Quantum Wells,” *Phys. Rev. Lett.* **108**, 106402 (2012), [arXiv:1111.7266](#).
- [39] B. Askenazi, A. Vasanelli, A. Delteil, Y. Todorov, L. C. Andreani, G. Beaudoin, I. Sagnes, and C. Sirtori, “Ultra-strong light-matter coupling for designer Reststrahlen band,” *New J. Phys.* **16**, 043029 (2014).
- [40] P. Forn-Díaz, J. Lisenfeld, D. Marcos, J. J. García-Ripoll, E. Solano, C. J. P. M. Harmans, and J. E. Mooij, “Observation of the Bloch-Siegert Shift in a Qubit-Oscillator System in the Ultrastrong Coupling Regime,” *Phys. Rev. Lett.* **105**, 237001 (2010), [arXiv:1005.1559](#).
- [41] T. Niemczyk, F. Deppe, H. Huebl, E. P. Menzel, F. Hocke, M. J. Schwarz, J. J. Garcia-Ripoll, D. Zueco, T. Hümmer, E. Solano, A. Marx, and R. Gross, “Circuit quantum electrodynamics in the ultrastrong-coupling regime,” *Nat. Phys.* **6**, 772 (2010), [arXiv:1003.2376](#).
- [42] A. Baust, E. Hoffmann, M. Haeberlein, M. J. Schwarz, P. Eder, J. Goetz, F. Wulschner, E. Xie, L. Zhong, F. Quijandría, D. Zueco, J.-J. García Ripoll, L. García-Álvarez, G. Romero, E. Solano, K. G. Fedorov, E. P. Menzel, F. Deppe, A. Marx, and R. Gross, “Ultrastrong coupling in two-resonator circuit QED,” *Phys. Rev. B* **93**, 214501 (2016), [arXiv:1412.7372](#).
- [43] P. Forn-Díaz, G. Romero, C. J. P. M. Harmans, E. Solano, and J. E. Mooij, “Broken selection

- rule in the quantum Rabi model,” *Sci. Rep.* **6**, 26720 (2016), [arXiv:1510.03379](#).
- [44] S. J. Bosman, M. F. Gely, V. Singh, A. Bruno, D. Bothner, and G. A. Steele, “Multi-mode ultra-strong coupling in circuit quantum electrodynamics,” *npj Quantum Inf.* **3**, 46 (2017), [arXiv:1704.06208](#).
 - [45] F. Yoshihara, T. Fuse, S. Ashhab, K. Kakuyanagi, S. Saito, and K. Semba, “Characteristic spectra of circuit quantum electrodynamics systems from the ultrastrong- to the deep-strong-coupling regime,” *Phys. Rev. A* **95**, 053824 (2017), [arXiv:1612.00121](#).
 - [46] F. Yoshihara, T. Fuse, S. Ashhab, K. Kakuyanagi, S. Saito, and K. Semba, “Superconducting qubit-oscillator circuit beyond the ultrastrong-coupling regime,” *Nat. Phys.* **13**, 44 (2017), [arXiv:1602.00415](#).
 - [47] N. K. Langford, R. Sagastizabal, M. Kounalakis, C. Dickel, A. Bruno, F. Luthi, D. J. Thoen, A. Endo, and L. DiCarlo, “Experimentally simulating the dynamics of quantum light and matter at deep-strong coupling,” *Nat. Commun.* **8**, 1715 (2017), [arXiv:1610.10065](#).
 - [48] J. Braumüller, M. Marthaler, A. Schneider, A. Stehli, H. Rotzinger, M. Weides, and A. V. Ustinov, “Analog quantum simulation of the Rabi model in the ultra-strong coupling regime,” *Nat. Commun.* **8**, 779 (2017), [arXiv:1611.08404](#).
 - [49] L. Magazzù, P. Forn-Díaz, R. Belyansky, J.-L. Orgiazzi, M. A. Yurtalan, M. R. Otto, A. Lupascu, C. M. Wilson, and M. Grifoni, “Probing the strongly driven spin-boson model in a superconducting quantum circuit,” *Nat. Commun.* **9**, 1403 (2018), [arXiv:1709.01157](#).
 - [50] F. Yoshihara, T. Fuse, Z. Ao, S. Ashhab, K. Kakuyanagi, S. Saito, T. Aoki, K. Koshino, and K. Semba, “Inversion of qubit energy levels in qubit-oscillator circuits in the deep-strong-coupling regime,” (2017), [arXiv:1712.05039](#).
 - [51] J. Puertas Martinez, S. Leger, N. Gheerart, R. Dassonneville, L. Planat, F. Foroughi, Y. Krupko, O. Buisson, C. Naud, W. Guichard, S. Florens, I. Snyman, and N. Roch, “Probing a transmon qubit via the ultra-strong coupling to a Josephson waveguide,” (2018), [arXiv:1802.00633](#).
 - [52] P. Forn-Díaz, J. J. García-Ripoll, B. Peropadre, J.-L. Orgiazzi, M. A. Yurtalan, R. Belyansky, C. M. Wilson, and A. Lupascu, “Ultrastrong coupling of a single artificial atom to an electromagnetic continuum in the nonperturbative regime,” *Nat. Phys.* **13**, 39 (2017), [arXiv:1602.00416](#).
 - [53] Z. Chen, Y. Wang, T. Li, L. Tian, Y. Qiu, K. Inomata, F. Yoshihara, S. Han, F. Nori,

- J. S. Tsai, and J. Q. You, “Single-photon-driven high-order sideband transitions in an ultra-strongly coupled circuit-quantum-electrodynamics system,” *Phys. Rev. A* **96**, 012325 (2017), [arXiv:1602.01584](#).
- [54] V. M. Muravev, I. V. Andreev, I. V. Kukushkin, S. Schmult, and W. Dietsche, “Observation of hybrid plasmon-photon modes in microwave transmission of coplanar microresonators,” *Phys. Rev. B* **83**, 075309 (2011).
- [55] G. Scalari, C. Maissen, D. Turcinkova, D. Hagenmuller, S. De Liberato, C. Ciuti, C. Reichl, D. Schuh, W. Wegscheider, M. Beck, and J. Faist, “Ultrastrong Coupling of the Cyclotron Transition of a 2D Electron Gas to a THz Metamaterial,” *Science* **335**, 1323 (2012), [arXiv:1111.2486](#).
- [56] C. Maissen, G. Scalari, F. Valmorra, M. Beck, J. Faist, S. Cibella, R. Leoni, C. Reichl, C. Charpentier, and W. Wegscheider, “Ultrastrong coupling in the near field of complementary split-ring resonators,” *Phys. Rev. B* **90**, 205309 (2014), [arXiv:1408.3547](#).
- [57] Q. Zhang, M. Lou, X. Li, J. L. Reno, W. Pan, J. D. Watson, M. J. Manfra, and J. Kono, “Collective non-perturbative coupling of 2D electrons with high-quality-factor terahertz cavity photons,” *Nat. Phys.* **12**, 1005 (2016), [arXiv:1604.08297](#).
- [58] A. Bayer, M. Pozimski, S. Schambeck, D. Schuh, R. Huber, D. Bougeard, and C. Lange, “Terahertz Light-Matter Interaction beyond Unity Coupling Strength,” *Nano Lett.* **17**, 6340 (2017), [arXiv:1807.00533](#).
- [59] T. Schwartz, J. A. Hutchison, C. Genet, and T. W. Ebbesen, “Reversible Switching of Ultrastrong Light-Molecule Coupling,” *Phys. Rev. Lett.* **106**, 196405 (2011).
- [60] S. Kéna-Cohen, S. A. Maier, and D. D. C. Bradley, “Ultrastrongly Coupled Exciton-Polaritons in Metal-Clad Organic Semiconductor Microcavities,” *Adv. Opt. Mater.* **1**, 827 (2013).
- [61] S. Gambino, M. Mazzeo, A. Genco, O. Di Stefano, S. Savasta, S. Patanè, D. Ballarini, F. Mangione, G. Lerario, D. Sanvitto, and G. Gigli, “Exploring Light-Matter Interaction Phenomena under Ultrastrong Coupling Regime,” *ACS Photonics* **1**, 1042 (2014).
- [62] C. R. Gubbin, S. A. Maier, and S. Kéna-Cohen, “Low-voltage polariton electroluminescence from an ultrastrongly coupled organic light-emitting diode,” *Appl. Phys. Lett.* **104**, 233302 (2014).
- [63] M. Mazzeo, A. Genco, S. Gambino, D. Ballarini, F. Mangione, O. Di Stefano, S. Patanè, S. Savasta, D. Sanvitto, and G. Gigli, “Ultrastrong light-matter coupling in electrically doped

- microcavity organic light emitting diodes,” *Appl. Phys. Lett.* **104**, 233303 (2014).
- [64] J. George, T. Chervy, A. Shalabney, E. Devaux, H. Hiura, C. Genet, and T. W. Ebbesen, “Multiple Rabi Splittings under Ultrastrong Vibrational Coupling,” *Phys. Rev. Lett.* **117**, 153601 (2016), [arXiv:1609.01520](#).
 - [65] F. Benz, M. K. Schmidt, A. Dreismann, R. Chikkaraddy, Y. Zhang, A. Demetriadou, C. Carnegie, H. Ohadi, B. de Nijs, R. Esteban, J. Aizpurua, and J. J. Baumberg, “Single-molecule optomechanics in “picocavities”,” *Science* **354**, 726 (2016).
 - [66] M. H. Devoret, S. Girvin, and R. Schoelkopf, “Circuit-QED: How strong can the coupling between a Josephson junction atom and a transmission line resonator be?” *Ann. Phys. (Leipzig)* **16**, 767 (2007).
 - [67] J. Bourassa, J. M. Gambetta, A. A. Abdumalikov, O. Astafiev, Y. Nakamura, and A. Blais, “Ultrastrong coupling regime of cavity QED with phase-biased flux qubits,” *Phys. Rev. A* **80**, 032109 (2009), [arXiv:0906.1383](#).
 - [68] O. Di Stefano, R. Stassi, L. Garziano, A. F. Kockum, S. Savasta, and F. Nori, “Feynman-diagrams approach to the quantum Rabi model for ultrastrong cavity QED: stimulated emission and reabsorption of virtual particles dressing a physical excitation,” *New J. Phys.* **19**, 053010 (2017), [arXiv:1603.04984](#).
 - [69] D. Sank, Z. Chen, M. Khezri, J. Kelly, R. Barends, B. Campbell, Y. Chen, B. Chiaro, A. Dunsworth, A. Fowler, E. Jeffrey, E. Lucero, A. Megrant, J. Mutus, M. Neeley, C. Neill, P. J. J. O’Malley, C. Quintana, P. Roushan, A. Vainsencher, T. White, J. Wenner, Alexander N. Korotkov, and J. M. Martinis, “Measurement-induced state transitions in a superconducting qubit: Beyond the rotating wave approximation,” *Phys. Rev. Lett.* **117**, 190503 (2016), [arXiv:1606.05721](#).
 - [70] C. Ciuti and I. Carusotto, “Input-output theory of cavities in the ultrastrong coupling regime: The case of time-independent cavity parameters,” *Phys. Rev. A* **74**, 033811 (2006), [arXiv:0606554 \[cond-mat\]](#).
 - [71] A. Ridolfo, S. Savasta, and M. J. Hartmann, “Nonclassical Radiation from Thermal Cavities in the Ultrastrong Coupling Regime,” *Phys. Rev. Lett.* **110**, 163601 (2013), [arXiv:1212.1280](#).
 - [72] L. Garziano, A. Ridolfo, S. De Liberato, and S. Savasta, “Cavity QED in the Ultrastrong Coupling Regime: Photon Bunching from the Emission of Individual Dressed Qubits,” *ACS Photonics* **4**, 2345 (2017), [arXiv:1702.00532](#).

- [73] S. De Liberato, D. Gerace, I. Carusotto, and C. Ciuti, “Extracavity quantum vacuum radiation from a single qubit,” *Phys. Rev. A* **80**, 053810 (2009), [arXiv:0906.2706](#).
- [74] M. Bamba and N. Imoto, “Maxwell boundary conditions imply non-lindblad master equation,” *Physical Review A* **94**, 033802 (2016), [arXiv:1603.00535](#).
- [75] D. Hu, S.-Y. Huang, J.-Q. Liao, L. Tian, and H.-S. Goan, “Quantum coherence in ultrastrong optomechanics,” *Phys. Rev. A* **91**, 013812 (2015), [arXiv:1410.2863](#).
- [76] K. K. W. Ma and C. K. Law, “Three-photon resonance and adiabatic passage in the large-detuning Rabi model,” *Phys. Rev. A* **92**, 023842 (2015).
- [77] J. Iles-Smith, N. Lambert, and A. Nazir, “Environmental dynamics, correlations, and the emergence of noncanonical equilibrium states in open quantum systems,” *Phys. Rev. A* **90**, 032114 (2014), [arXiv:1311.0016](#).
- [78] J. Iles-Smith, A. G. Dijkstra, N. Lambert, and A. Nazir, “Energy transfer in structured and unstructured environments: Master equations beyond the Born-Markov approximations,” *The Journal of chemical physics* **144**, 044110 (2016), [arXiv:1511.05181](#).
- [79] P. Strasberg, G. Schaller, N. Lambert, and Tobias Brandes, “Nonequilibrium thermodynamics in the strong coupling and non-markovian regime based on a reaction coordinate mapping,” *New Journal of Physics* **18**, 073007 (2016), [arXiv:1602.01340](#).
- [80] S. De Liberato, “Virtual photons in the ground state of a dissipative system,” *Nature Communications* **8**, 1465 (2017), [arXiv:1609.04249](#).
- [81] D. Zueco and J. García-Ripoll, “Ultrastrongly dissipative quantum Rabi model,” [arXiv:1806.01952](#) (2018).
- [82] F. Altintas and R. Eryigit, “Dissipative dynamics of quantum correlations in the strong-coupling regime,” *Phys. Rev. A* **87**, 022124 (2013).
- [83] D. Pagel, A. Alvermann, and H. Fehske, “Nonclassical light from few emitters in a cavity,” *Phys. Rev. A* **91**, 043814 (2015), [arXiv:1502.04529](#).
- [84] A. Le Boité, M.-J. Hwang, H. Nha, and M. B. Plenio, “Fate of photon blockade in the deep strong-coupling regime,” *Phys. Rev. A* **94**, 033827 (2016), [arXiv:1607.05062](#).
- [85] R. Stassi and F. Nori, “Long-lasting quantum memories: Extending the coherence time of superconducting artificial atoms in the ultrastrong-coupling regime,” *Phys. Rev. A* **97**, 033823 (2018), [arXiv:1703.08951](#).
- [86] B. Kraus, H. P. Büchler, S. Diehl, A. Kantian, A. Micheli, and P. Zoller, “Preparation of entan-

- gled states by quantum Markov processes,” *Phys. Rev. A* **78**, 042307 (2008), [arXiv:0803.1463](#).
- [87] V. Macrì, A. Ridolfo, O. Di Stefano, A. F. Kockum, F. Nori, and S. Savasta, “Nonperturbative Dynamical Casimir Effect in Optomechanical Systems: Vacuum Casimir-Rabi Splittings,” *Phys. Rev. X* **8**, 011031 (2018), [arXiv:1706.04134](#).
- [88] G. T. Moore, “Quantum Theory of the Electromagnetic Field in a Variable-Length One-Dimensional Cavity,” *J. Math. Phys.* **11**, 2679 (1970).
- [89] P. D. Nation, J. R. Johansson, M. P. Blencowe, and Franco Nori, “Colloquium: Stimulating uncertainty: Amplifying the quantum vacuum with superconducting circuits,” *Rev. Mod. Phys.* **84**, 1–24 (2012), [arXiv:1103.0835](#).
- [90] E. T. Jaynes and F. W. Cummings, “Comparison of quantum and semiclassical radiation theories with application to the beam maser,” *Proc. IEEE* **51**, 89 (1963).
- [91] B. W. Shore and P. L. Knight, “The Jaynes-Cummings Model,” *J. Mod. Opt.* **40**, 1195 (1993).
- [92] C. K. Law, “Interaction between a moving mirror and radiation pressure: A Hamiltonian formulation,” *Phys. Rev. A* **51**, 2537 (1995).
- [93] J. R. Johansson, G. Johansson, C. M. Wilson, and F. Nori, “Dynamical Casimir effect in superconducting microwave circuits,” *Phys. Rev. A* **82**, 52509 (2010), [arXiv:1007.1058](#).
- [94] C. M. Wilson, G. Johansson, A. Pourkabirian, M. Simoen, J. R. Johansson, T. Duty, F. Nori, and P. Delsing, “Observation of the dynamical Casimir effect in a superconducting circuit,” *Nature* **479**, 376 (2011), [arXiv:1105.4714](#).
- [95] A. Lambrecht, M.-T. Jaekel, and S. Reynaud, “Motion Induced Radiation from a Vibrating Cavity,” *Phys. Rev. Lett.* **77**, 615 (1996), [arXiv:9606029](#).