



## $L^\infty$ algebras for extended geometry

Downloaded from: <https://research.chalmers.se>, 2024-04-17 21:28 UTC

Citation for the original published paper (version of record):

Cederwall, M., Palmkvist, J. (2019).  $L^\infty$  algebras for extended geometry. Journal of Physics: Conference Series, 1194(1). <http://dx.doi.org/10.1088/1742-6596/1194/1/012021>

N.B. When citing this work, cite the original published paper.

PAPER • OPEN ACCESS

## $L_{\infty}$ algebras for extended geometry

To cite this article: Martin Cederwall and Jakob Palmkvist 2019 *J. Phys.: Conf. Ser.* **1194** 012021

View the [article online](#) for updates and enhancements.



**IOP | ebooks™**

Bringing you innovative digital publishing with leading voices to create your essential collection of books in STEM research.

Start exploring the **collection** - download the first chapter of every title for free.

# $L_\infty$ algebras for extended geometry

Martin Cederwall and Jakob Palmkvist

Division for Theoretical Physics, Dept. of Physics

Chalmers University of Technology

SE-412 96 Gothenburg, Sweden

E-mail: martin.cederwall@chalmers.se, jakob.palmkvist@chalmers.se

**Abstract.** Extended geometry provides a unified framework for double geometry, exceptional geometry, etc., *i.e.*, for the geometrisations of the string theory and M-theory dualities. In this talk, we will explain the structure of gauge transformations (generalised diffeomorphisms) in these models. They are generically infinitely reducible, and arise as derived brackets from an underlying Borchers superalgebra or tensor hierarchy algebra. The infinite reducibility gives rise to an  $L_\infty$  structure, the brackets of which have universal expressions in terms of the underlying superalgebra.

The results in this talk, presented by MC at Group32, Prague, July 9-13, 2018, are primarily based on refs. [1, 2].

The motivation for the investigation lies in the dualities appearing in string/M-theory, and the possibility to “geometrise” them. The plan of the talk is to briefly mention how the dualities arise, then display the basics of the general framework of extended geometry, and finally discuss the gauge structure, and the appearance of an  $L_\infty$  algebra.

When M-theory is compactified on an  $n$ -torus, the U-duality group is  $E_{n(n)}(\mathbb{Z})$ . The duality mixes momenta and brane windings. If this symmetry is to be geometrised, also diffeomorphisms and tensor gauge transformations need to be unified.

Membranes can wind on 2-cycles, 5-branes on 5-cycles. An example: Take a 6-dimensional torus  $T^6$ . There are  $\binom{6}{2} = 15$  membrane windings  $Z_{ij}$  and  $\binom{6}{5} = 6$  5-brane windings  $Z_{ijklm}$ . Collect them together with the momentum  $p^i$  into the generalised momentum  $P^M = (p^i, Z_{ij}, Z_{ijklm})$ . These generalised momenta span a 27-dimensional space, and transform under  $E_{6(6)}(\mathbb{Z})$ .

The discrete duality group contains the geometric mapping class group. It turns out that it can be geometrised, so that (roughly speaking) the duality group derives from an “extended geometry” like the mapping class group from geometry. Special cases are double geometry (for T-duality) [3–17], and exceptional geometry (for U-duality) [18–37]. Recently, a completely general framework has been formulated [1], where any Kac–Moody group can be used as structure group instead of the continuous version of a duality group, and where any integrable highest



weight representation can be used for the generalised momenta (at least for highest weights with vanishing Dynkin labels for short roots).

The gauge transformations in extended geometry — the generalised diffeomorphisms — unify diffeomorphisms and gauge transformations for tensor fields. Given a Kac–Moody algebra  $\mathfrak{g}$  and a lowest weight coordinate representation  $R(-\lambda)$  (we use conventions where extended tangent space vectors are in lowest weight modules and cotangent vectors in highest weight modules;  $R(-\lambda)$  denotes the lowest weight representation with lowest weight  $-\lambda$ , for dominant  $\lambda$ ), they are expressed in terms of a generalised Lie derivative as  $\delta_U \phi = \mathcal{L}_U \phi$ , for any field  $\phi$  transforming covariantly, where

$$\mathcal{L}_U V^M = U^N \partial_N V^M + Z_{PQ}{}^{MN} \partial_N U^P V^Q . \quad (1)$$

Here the invariant tensor  $Z$  has the universal expression [1, 36]

$$\sigma Z = -\eta_{\alpha\beta} t^\alpha \otimes t^\beta + (\lambda, \lambda) - 1 , \quad (2)$$

*i.e.*,  $Z_{PQ}{}^{MN} = -\eta_{\alpha\beta} (t^\alpha)_P{}^N (t^\beta)_Q{}^M + ((\lambda, \lambda) - 1) \delta_P^N \delta_Q^M$ , where  $(t^\alpha)_M{}^N$  are representation matrices in  $R(-\lambda)$ ,  $\sigma$  is the permutation operator and  $\eta_{\alpha\beta}$  the Killing metric.

In many cases (see below), the transformations form an “algebra”

$$[\mathcal{L}_U, \mathcal{L}_V]W = \mathcal{L}_{[U,V]}W , \quad (3)$$

where the “Courant bracket” is  $\llbracket U, V \rrbracket = \frac{1}{2}(\mathcal{L}_U V - \mathcal{L}_V U)$ , provided that the derivatives fulfil a *section constraint*. The section constraint ensures that fields locally depend only on an  $n$ -dimensional subspace of the coordinate space, on which a  $GL(n)$  subgroup acts. It reads

$$Y^{MN}{}_{PQ} \partial_M \otimes \partial_N = 0 , \quad (4)$$

where the two derivatives can act on any field (or parameter). The tensor  $Y$  is related to  $Z$  as  $Y = Z + 1$ , and can be seen (in eq. (1)) as the deviation of the generalised Lie derivative from a Lie derivative on the extended space. The section constraint means that any two momenta lie in a linear subspace of a minimal orbit of  $R(\lambda)$ .

The “algebra” is not a Lie algebra. The consequences will be examined below.

The closure of the generalised diffeomorphisms as in eq. (3) takes place only under certain conditions:

- The algebra  $\mathfrak{g}$  is finite-dimensional,
- $(\lambda, \theta) = 1$ , *i.e.*, the highest weight  $\lambda$  is a fundamental weight dual to a simple root with Coxeter label 1.

In other cases, so-called ancillary transformations occur. These are restricted local  $\mathfrak{g}$ -transformations, that remove degrees of freedom corresponding to mixed tensors (dual graviton, etc.)

The complete list of situations without ancillary transformations is:

- (i)  $\mathfrak{g}_r = A_r$ ,  $\lambda = \Lambda_p$ ,  $p = 1, \dots, r$  ( $p$ -form representations);



**Figure 1.** Dynkin diagram of  $\mathcal{B}(\mathfrak{g})$ . The grey node is a fermionic null node. The single line connecting to the Dynkin diagram of  $\mathfrak{g}$  can be replaced by multiple lines if  $\lambda$  is not a fundamental weight.

- (ii)  $\mathfrak{g}_r = B_r$ ,  $\lambda = \Lambda_1$  (the vector representation);
- (iii)  $\mathfrak{g}_r = C_r$ ,  $\lambda = \Lambda_r$  (the symplectic-traceless  $r$ -form representation);
- (iv)  $\mathfrak{g}_r = D_r$ ,  $\lambda = \Lambda_1, \Lambda_{r-1}, \Lambda_r$  (the vector and spinor representations);
- (v)  $\mathfrak{g}_r = E_6$ ,  $\lambda = \Lambda_1, \Lambda_5$  (the fundamental representations);
- (vi)  $\mathfrak{g}_r = E_7$ ,  $\lambda = \Lambda_1$  (the fundamental representation).

The Jacobi identity does not hold for the bracket  $[[\cdot, \cdot]]$ . Instead, one typically has

$$[[U, [V, W]]] + \text{cycl.} \sim d[[U, V, W]] , \quad (5)$$

where  $[[U, V, W]] \in R_2$  represents reducibility. The derivative  $d$  is linear in  $\partial_M$ .  $R_1, R_2, \dots$  are the positive levels (level = ghost number) of the Borchers superalgebra  $\mathcal{B}(\mathfrak{g})$  [25, 38] with Dynkin diagram given in Figure 1.

This is the beginning of the  $L_\infty$  structure.

What is an  $L_\infty$  algebra [39–42]? Consider a full set of ghosts, including ghosts for ghosts, etc. Let  $C = C_1 + C_2 + C_3 + \dots$ , where the subscript indicates ghost number. (We will later see this as an element in  $\mathcal{B}_+(\mathfrak{g})$ , the positive level subspace of  $\mathcal{B}(\mathfrak{g})$ .) The Batalin–Vilkovisky (BV) action [43], restricted to ghosts, can be expanded as

$$S(C, C^\star) = \sum_{n=1}^{\infty} \langle C^\star, [[C^N]] \rangle ; \quad (6)$$

where

$$[[C^m]] = \underbrace{[[C, C, \dots, C]]}_n \quad (7)$$

is the  $n$ -bracket.

The BV variation of  $C$  is

$$(S, C) = \sum_{n=1}^{\infty} [[C^m]] . \quad (8)$$

In order for it to be nilpotent,  $(S, (S, C)) = 0$ , the brackets must satisfy the generalised Jacobi identities

$$\sum_{i=0}^{n-1} (i+1) [[C^i, [[C^{m-i}]]] = 0 . \quad (9)$$



**Figure 2.** The Dynkin diagrams of  $\mathfrak{g}_{r+1}$ .



**Figure 3.** Equivalent Dynkin diagrams of  $\mathcal{B}(\mathfrak{g}_{r+1})$ .

The derivative is the 1-bracket.

$$R_1 \stackrel{\leftarrow}{\underset{d}{}} R_2 \stackrel{\leftarrow}{\underset{d}{}} R_3 \stackrel{\leftarrow}{\underset{d}{}} \dots \quad (10)$$

In order to be able to construct all the brackets, one needs to rely on some underlying structure. It would seem like the superalgebra  $\mathcal{B}(\mathfrak{g})$  provides it. However, there is no natural way of expressing the generalised Lie derivatives in terms of the (anti-)commutators of  $\mathcal{B}(\mathfrak{g})$ . For this one needs a further extension,  $\mathcal{B}(\mathfrak{g}_{r+1})$ . The bosonic extension  $\mathfrak{g}_{r+1}$  has the Dynkin diagram in Figure 2 (here, like for  $\mathcal{B}(\mathfrak{g})$ , the connection of the extending node to the Dynkin diagram of  $\mathfrak{g}$  can consist of multiple lines). Two Dynkin diagrams of  $\mathcal{B}(\mathfrak{g}_{r+1})$  are given in Figure 3. The leftmost line is always single. The equivalence between the two diagrams, expressing  $\mathcal{B}(\mathfrak{g}_{r+1})$  as an extension of  $\mathcal{B}(\mathfrak{g})$  or  $\mathfrak{g}_{r+1}$ , respectively, involves a fermionic Weyl reflection [44].


It was noted in ref. [45] (in that case for the exceptional series, but straightforwardly applied to the general situation) that the generalised diffeomorphisms have a natural expression in terms of the Lie super-brackets of  $\mathcal{B}(\mathfrak{g}_{r+1})$ . The actual expression is

$$\mathcal{L}_U V = [[U, F^{\flat M}], \partial_M V^\sharp] - [[\partial_M U^\sharp, F^{\flat M}], V], \quad (11)$$

The notation needs some explanation. We note that, in the double grading of Table 1, all  $\mathfrak{g}$ -modules at  $p \neq 0$  come in pairs. We use the nilpotent operations  $\sharp$  and  $\flat$  to raise and lower elements within the pairs.  $F^M$  are basis elements for  $R(\lambda)$  at  $p = -1$ ,  $q = 0$ , and their lowered counterpart  $F^{\flat M}$  span  $R(\lambda)$  at  $p = -1$ ,  $q = 0$ . The two terms reproduce the two terms in eq. (1).

The doubly extended algebra is needed also in order to harbour so called “ancillary” ghosts, which appear in most cases, including the exceptional series. At some level (ghost number) the derivative fails to be a derivation, and the generalised Lie derivative fails to be covariant. This gives rise to the extra ancillary ghosts. They are naturally encoded in the doubly extended algebra as elements in  $R_p$  at  $q = 1$ . They are restricted with respect to the section constraint in a certain sense, and can be constructed as a derivative together with an element in  $\tilde{R}_{p+1}$ . We refer to ref. [2] for details.

Using this Borchers superalgebra as an underlying structure, we are able to construct all brackets as derived brackets and check their identities [2].



**Table 1.** The general structure of the superalgebra  $\mathcal{B}(\mathfrak{g}_{r+1})$ . The blue lines are the  $L_\infty$ -levels, given by  $\ell = p + q$ . Red lines are the usual levels in the level decomposition of  $\mathcal{B}(\mathfrak{g}_{r+1})$ , and form  $\mathfrak{g}_{r+1}$  modules.

$$\begin{array}{ccccccc}
 & & & & K_{p_0} & \xleftarrow{d} & K_{p_0+1} & \xleftarrow{d} & K_{p_0+2} & \xleftarrow{d} & \dots \\
 & & & & \downarrow \mathfrak{b} & & \downarrow \mathfrak{b} & & \downarrow \mathfrak{b} & & \\
 0 & \xleftarrow{d} & C_1 & \xleftarrow{d} & \dots & \xleftarrow{d} & C_{p_0-1} & \xleftarrow{d} & C_{p_0} & \xleftarrow{d} & C_{p_0+1} & \xleftarrow{d} & C_{p_0+2} & \xleftarrow{d} & \dots
 \end{array}$$

**Table 2.** The typical structure of the action of the 1-bracket between the ghost modules, with ancillary ghosts appearing from level  $p_0 \geq 1$ .

The concrete expression for all brackets read:

$$\begin{aligned}
 \llbracket C \rrbracket &= dC, \\
 \llbracket K \rrbracket &= dK + K^{\mathfrak{b}}, \\
 \llbracket C^n \rrbracket &= k_n \left( (\text{ad } C)^{n-2} (\mathcal{L}_C C + X_C C) + \sum_{i=0}^{n-3} (\text{ad } C)^i R_C (\text{ad } C)^{n-i-3} \mathcal{L}_C C \right) \quad (12) \\
 \llbracket C^{n-1}, K \rrbracket &= \frac{k_n}{n} \left( (\text{ad } C)^{n-2} \mathcal{L}_C K + \sum_{i=0}^{n-3} (\text{ad } C)^i \text{ad } K (\text{ad } C)^{n-i-3} \mathcal{L}_C C \right),
 \end{aligned}$$

where the coefficients have the universal model-independent expression in terms of Bernoulli numbers

$$k_{n+1} = \frac{2^n B_n^+}{n!}, \quad n \geq 1. \quad (13)$$

All non-vanishing brackets except the 1-bracket contain at least one level 1 ghost  $C_1$ . No brackets contain more than one ancillary ghost. In the expressions for the brackets,  $\mathcal{L}_C = \mathcal{L}_{C_1}$ , i.e., only

the ghost number 1 part of  $C$  enters the generalised Lie derivative.  $X$  and  $R$  are ancillary contributions, and we refer to ref. [2] for their exact expressions.

The  $L_\infty$  algebra for double geometry was constructed in refs. [41, 46, 47]. Then, there are no ancillary ghosts, and the algebra stops at ghost number 2 and a 3-bracket. This is because the corresponding Borchers superalgebra is finite-dimensional.

The Borchers superalgebra is unable to handle situations where ancillary transformations appear in the commutator of two generalised Lie derivatives. The presence of ancillary ghost at ghost number  $n$  relies on the occurrence of a  $\mathfrak{g}$ -module  $\tilde{R}_p$  in  $\mathcal{B}(\mathfrak{g}_{r+1})$  (see Table 1). The Borchers superalgebra never contains  $\tilde{R}_1$ , which would be the signal of ancillary ghosts with ghost number 1, *i.e.*, of ancillary transformations  $\Sigma$  in the commutator of two generalised Lie derivatives,

$$[\mathcal{L}_U, \mathcal{L}_V]W = \mathcal{L}_{[U,V]}W + \Sigma_{U,V}W \quad (14)$$

The ancillary parameter  $\Sigma$  is a parameter for a restricted (with respect to the section constraint) local  $\mathfrak{g}$ -transformation, see refs. [34–36, 48]. Then one needs a *tensor hierarchy algebra* [49, 50], a generalisation of the Cartan-type superalgebras  $W(n)$  and  $S(n)$  in Kac’s classification [51]. Tensor hierarchy algebras are non-contragredient superalgebras, and therefore *a priori* not defined by standard Chevalley–Serre relations from a Dynkin diagram. In ref. [50] we presented a set of generators and relations for the (finite-dimensional) tensor hierarchy algebras  $W(r+1) = W(A_r)$  and  $S(r+1) = S(A_r)$ , based on the same Dynkin diagram as that of  $\mathcal{B}(A_r)$ . The straightforward generalisation of these relations seems to provide a good definition of  $W(\mathfrak{g})$  and  $S(\mathfrak{g})$  in general (see the talk by JP at the present meeting [52]). The superalgebra used for extended geometry with structure algebra  $\mathfrak{g}$  is  $S(\mathfrak{g}_{r+1})$ . These algebras agree with the Borchers superalgebras at positive levels, and turn out to “know” when ancillary transformations appear: they contain a module  $\tilde{R}_1$  exactly in these cases (*i.e.*, when  $\mathfrak{g}$  is infinite-dimensional and when  $\mathfrak{g}$  is finite-dimensional and  $(\lambda, \theta) > 1$ ). In the series of exceptional duality symmetries, this development is necessary starting from  $E_8$ . It also seems promising for incorporating dynamical fields (vielbein, torsion,...) in the present framework. Work is in progress concerning the rôle of tensor hierarchy algebras in extended geometry [53, 54], and we believe that they will encode the information needed for the gauge structure and dynamics also in cases with infinite-dimensional structure groups, such as  $E_9$  [36, 37],  $E_{10}$  and maybe  $E_{11}$  [55].

## References

- [1] Cederwall M and Palmkvist J 2018 Extended geometries *J. High Energy Phys.* JHEP02(2018)071 (*Preprint* arXiv:1711.07694).
- [2] Cederwall M and Palmkvist J 2018  $L_\infty$  algebras for extended geometry from Borchers superalgebras *Preprint* arXiv:1804.04377.
- [3] Tseytlin A A 1991 Duality symmetric closed string theory and interacting chiral scalars *Nucl. Phys. B* **350** 395.
- [4] Siegel W 1993 Two vierbein formalism for string inspired axionic gravity *Phys. Rev. D* **47**, 5453 (*Preprint* hep-th/9302036).
- [5] Siegel W 1993 Manifest duality in low-energy superstrings *International Conference on Strings 93 Berkeley* p 353 (*Preprint* hep-th/9308133).
- [6] Hitchin N 2010 Lectures on generalized geometry *Preprint* arXiv:1008.0973.
- [7] Hull C M 2005 A geometry for non-geometric string backgrounds *J. High Energy Phys.* JHEP10(2005)065 (*Preprint* hep-th/0406102).



- [8] Hull C M 2007 Doubled geometry and T-folds *J. High Energy Phys.* JHEP07(2007)080 (*Preprint* hep-th/0605149).
- [9] Hull C M and Zwiebach 2009 Double field theory *J. High Energy Phys.* JHEP09(2009)099 (*Preprint* arXiv:0904.4664).
- [10] Hohm O, Hull C M and Zwiebach W 2010 Background independent action for double field theory *J. High Energy Phys.* JHEP07(2010)016 (*Preprint* arXiv:1003.5027).
- [11] Hohm O, Hull C M and Zwiebach B 2010 Generalized metric formulation of double field theory *J. High Energy Phys.* JHEP08(2010)008 (*Preprint* arXiv:1006.4823).
- [12] Jeon I, Lee K, Park J-H and Suh Y 2013 Stringy unification of type IIA and IIB supergravities under  $N = 2$   $D = 10$  supersymmetric double field theory *Phys. Lett. B* **723** 245 (*Preprint* arXiv:1210.5078).
- [13] Park J-H 2013 Comments on double field theory and diffeomorphisms *J. High Energy Phys.* JHEP06(2013)098 (*Preprint* arXiv:1304.5946).
- [14] Berman D S, Cederwall M and Perry M J 2014 Global aspects of double geometry *J. High Energy Phys.* JHEP09(2014)066 (*Preprint* arXiv:1401.1311).
- [15] Cederwall M 2014 The geometry behind double geometry *J. High Energy Phys.* JHEP09(2014)070 (*Preprint* arXiv:1402.2513).
- [16] Cederwall M 2014 T-duality and non-geometric solutions from double geometry *Fortsch. Phys.* **62** 942 (*Preprint* arXiv:1409.4463).
- [17] Cederwall M 2016 Double supergeometry *J. High Energy Phys.* JHEP06(2016)155 (*Preprint* arXiv:1603.04684).
- [18] Hull C M 2007 Generalised geometry for M-theory *J. High Energy Phys.* JHEP07(2007)079 (*Preprint* hep-th/0701203).
- [19] Pacheco P P and Waldram D 2008 M-theory, exceptional generalised geometry and superpotentials *J. High Energy Phys.* JHEP09(2008)123 (*Preprint* arXiv:0804.1362).
- [20] Hillmann C 2008  $E_{7(7)}$  and  $d = 11$  supergravity *PhD thesis* (Humboldt-Universität zu Berlin) (*Preprint* arXiv:0902.1509).
- [21] Berman D S and Perry M J 2011 Generalized geometry and M theory *J. High Energy Phys.* JHEP06(2011)074 (*Preprint* arXiv:1008.1763).
- [22] Berman D S, Godazgar H and Perry M J 2011  $SO(5, 5)$  duality in M-theory and generalized geometry *Phys. Lett. B* **700** 65 (*Preprint* arXiv:1103.5733).
- [23] Coimbra A, Strickland-Constable C and Waldram D 2014  $E_{d(d)} \times \mathbb{R}^+$  generalised geometry, connections and M theory *J. High Energy Phys.* JHEP02(2014)054 (*Preprint* arXiv:1112.3989).
- [24] Coimbra A, Strickland-Constable C and Waldram D 2014 Supergravity as generalised geometry II:  $E_{d(d)} \times \mathbb{R}^+$  and M theory *J. High Energy Phys.* JHEP03(2014)019 (*Preprint* arXiv:1212.1586).
- [25] Berman D S, Cederwall M, Kleinschmidt A and Thompson D C 2013 The gauge structure of generalised diffeomorphisms *J. High Energy Phys.* JHEP01(2013)064 (*Preprint* arXiv:1208.5884).
- [26] Park J-H and Suh Y 2013 U-geometry :  $SL(5)$  *J. High Energy Phys.* JHEP04(2013)147 (*Preprint* arXiv:1302.1652).
- [27] Cederwall M, Edlund J and Karlsson A 2013 Exceptional geometry and tensor fields *J. High Energy Phys.* JHEP07(2013)028 (*Preprint* arXiv:1302.6736).
- [28] Cederwall M 2013 Non-gravitational exceptional supermultiplets *J. High Energy Phys.* JHEP07(2013)025 (*Preprint* arXiv:1302.6737).
- [29] Aldazabal G, Graña M, Marqués D and Rosabal J A 2013 Extended geometry and gauged maximal supergravity *J. High Energy Phys.* JHEP06(2013)046 (*Preprint* arXiv:1302.5419).
- [30] Hohm O and Samtleben H 2013 Exceptional form of  $D = 11$  supergravity *Phys. Rev. Lett.* **111** 231601 (*Preprint* arXiv:1308.1673).
- [31] Blair C D, Malek E and Park J-H 2014 M-theory and type IIB from a duality manifest action *J. High Energy Phys.* JHEP01(2014)172 (*Preprint* arXiv:1311.5109).
- [32] Hohm O and Samtleben H 2013 Exceptional field theory I:  $E_{6(6)}$  covariant form of M-theory and type IIB *Phys. Rev. D* **89** 066016 (*Preprint* arXiv:1312.0614).
- [33] Hohm O and Samtleben H 2014 Exceptional field theory II:  $E_{7(7)}$  *Phys. Rev. D* **89** 066017 (*Preprint* arXiv:1312.4542).
- [34] Hohm O and Samtleben H 2014 Exceptional field theory. III.  $E_{8(8)}$  *Phys. Rev. D* **90** 066002 (*Preprint* arXiv:1406.3348).
- [35] Cederwall M and Rosabal J A 2015  $E_8$  geometry *J. High Energy Phys.* JHEP07(2015)007 (*Preprint* arXiv:1504.04843).
- [36] Bossard G, Cederwall M, Kleinschmidt A, Palmkvist J and Samtleben H 2017 Generalized diffeomorphisms for  $E_9$  *Phys. Rev. D* **96** 106022 (*Preprint* arXiv:1708.08936).
- [37] Bossard G, Ciceri F, Inverso G, Kleinschmidt A and Samtleben H 2018  $E_9$  exceptional field theory I. The

- potential *Preprint* arXiv:1811.04088.
- [38] Cederwall M and Palmkvist J 2015 Superalgebras, constraints and partition functions *J. High Energy Phys.* JHEP08(2015)036 (*Preprint* arXiv:1503.06215).
  - [39] Lada T and Stasheff J 1993 Introduction to SH Lie algebras for physicists *Int. J. Theor. Phys.* **32** 1087 (*Preprint* hep-th/9209099).
  - [40] Zwiebach B 1993 Closed string field theory: Quantum action and the B-V master equation *Nucl. Phys. B* **390** 33 (*Preprint* hep-th/9206084).
  - [41] Hohm O and Zwiebach B 2017  $L_\infty$  algebras and field theory *Fortsch. Phys.* **65** 1700014 (*Preprint* arXiv:1701.08824).
  - [42] Roytenberg D and Weinstein A 1998 Courant algebroids and strongly homotopy Lie algebras *Preprint* math/9802118.
  - [43] Batalin I A and Vilkovisky G A 1981 Gauge algebra and quantization *Phys. Lett. B* **102** 27.
  - [44] Dobrev V K and Petkova V B 1987 Group theoretical approach to extended conformal supersymmetry: Function space realizations and invariant differential operators *Fortsch. Phys.* **35** 537.
  - [45] Palmkvist J 2015 Exceptional geometry and Borchers superalgebras *J. High Energy Phys.* JHEP11(2015)032 (*Preprint* arXiv:1507.08828).
  - [46] Deser A and Sämann C 2016 Extended Riemannian geometry I: Local double field theory (*Preprint* arXiv:1611.02772).
  - [47] Deser A and Sämann C 2017 Derived brackets and symmetries in generalized geometry and double field theory *17th Hellenic School and Workshops on Elementary Particle Physics and Gravity (CORFU2017)* (*Preprint* arXiv:1803.01659).
  - [48] Hohm O and Samtleben H 2013 U-duality covariant gravity *J. High Energy Phys.* JHEP09(2013)080 (*Preprint* arXiv:1307.0509).
  - [49] Palmkvist J 2014 The tensor hierarchy algebra *J. Math. Phys.* **55** 011701 (*Preprint* arXiv:1305.0018).
  - [50] Carbone L, Cederwall M and Palmkvist J 2018 Generators and relations for Lie superalgebras of Cartan type *Preprint* arXiv:1802.05767.
  - [51] Kac V G 1977 Lie superalgebras *Adv. Math.* **26** 8.
  - [52] Carbone L, Cederwall M and Palmkvist J 2018 Generators and relations for (generalised) Cartan superalgebras *talk by JP at Group32* (*Preprint* arXiv:1812.03068).
  - [53] Cederwall M and Palmkvist J 2019 Extended geometry and tensor hierarchy algebras I: Constructing the algebra from generators and relations, *to appear*.
  - [54] Cederwall M and Palmkvist J 2019 Extended geometry and tensor hierarchy algebras II: Gauge structure and dynamics, *to appear*.
  - [55] Bossard G, Kleinschmidt A, Palmkvist J, Pope C N and Sezgin E 2017 Beyond  $E_{11}$  *J. High Energy Phys.* JHEP05(2017)020 (*Preprint* arXiv:1703.01305).