



**CHALMERS**  
UNIVERSITY OF TECHNOLOGY

## **Bioconversion of pretreated wheat straw to ethanol by *monascus purpureus* CBS 109.07 and *fusarium venenatum* ATCC 20334 using simultaneous**

Downloaded from: <https://research.chalmers.se>, 2024-07-27 07:43 UTC

### Citation for the original published paper (version of record):

Yudianto, D., Nainggolan, E., Millati, R. et al (2019). Bioconversion of pretreated wheat straw to ethanol by *monascus purpureus* CBS 109.07 and *fusarium venenatum* ATCC 20334 using simultaneous saccharification and fermentation. *Biodiversitas*, 20(8): 2229-2235. <http://dx.doi.org/10.13057/biodiv/d200817>

N.B. When citing this work, cite the original published paper.

# Bioconversion of pretreated wheat straw to ethanol by *Monascus purpureus* CBS 109.07 and *Fusarium venenatum* ATCC 20334 using simultaneous saccharification and fermentation

DAVID YUDIANTO<sup>1</sup>, ELLYAS ALGA NAINGGOLAN<sup>2</sup>, RIA MILLATI<sup>3,\*</sup>, CHUSNUL HIDAYAT<sup>3</sup>, PATRIK LENNARTSSON<sup>4</sup>, MOHAMMAD J. TAHERZADEH<sup>4</sup>, CLAES NIKLASSON<sup>5</sup>

<sup>1</sup>Quality Assurance of Food Industry, Politeknik AKA Bogor, Jl. Pangeran Sogiri No. 283, Tanah Baru, Kota Bogor 16154, West Java, Indonesia. Tel./fax.: +62- 251-8650351, email: davidyudianto.se@gmail.com

<sup>2</sup>Department of Bioprocess Engineering, Faculty of Biotechnology, Institut Teknologi Del. Sitoluama, Toba Samosir 22381, North Sumatra, Indonesia

<sup>3</sup>Department of Food and Agricultural Product Technology, Faculty of Agricultural Technology, Universitas Gadjah Mada. Jl. Flora No. 1, Bulaksumur, Sleman 55281, Yogyakarta, Indonesia. Tel./fax.: +62-274-563062, \*email: ria\_millati@ugm.ac.id

<sup>4</sup>Swedish Center for Resource Recovery, University of Borås. 50190 Borås, Sweden

<sup>5</sup>Department of Chemistry and Chemical Engineering, Chalmers University of Technology. 41296 Gothenburg, Sweden

Manuscript received: 27 February 2019. Revision accepted: 21 June 2019.

**Abstract.** Yudianto D, Nainggolan EA, Millati R, Hidayat C, Lennartsson P, Taherzadeh MJ, Niklasson C. 2019. Bioconversion of pretreated wheat straw to ethanol by *Monascus purpureus* CBS 109.07 and *Fusarium venenatum* ATCC 20334 using simultaneous saccharification and fermentation. 20: 2229-2235. Fractions of sulfuric acid-pretreated wheat straw, i.e. solid, liquid, and a mixture of liquid and solid were used as substrates in simultaneous saccharification and fermentation (SSF) process to produce ethanol. The bioconversion was performed by *Monascus purpureus* CBS 109.07 and *Fusarium venenatum* ATCC 20334. The highest ethanol yields from solid, liquid and a mixture of solid and liquid fractions by *M. purpureus* CBS 109.07 were 0.36, 0.41, and 0.37 g/g glucose, respectively. The corresponding values by *F. venenatum* ATCC 20334 were 0.21, 0.54, 0.35 g/g glucose, respectively.

**Keywords:** Ethanol, *Fusarium venenatum* ATCC 20334, *Monascus purpureus* CBS 109.07, Simultaneous Saccharification and Fermentation, wheat straw

## INTRODUCTION

Wheat is one of the major crops in the United Kingdom (UK), and according to a report in 2009, the annual wheat straw yield was estimated to be between 8 and 10 million tons (Brander et al. 2009). Straws as lignocellulosic materials have been utilized for animal feed or making traditional objects (straw baskets or hats), but a major part of straw is discarded as waste (Hammond and Mansell 2018). One of potential usages of wheat straws is by converting it to more value-added products such as ethanol. The contents of glucan and xylan in wheat straw are approximately 36.3% and 19%, respectively, and have the potential to be converted to ethanol (Kootstra et al. 2009). Ethanol from lignocellulosic is a renewable fuel, which can reduce CO<sub>2</sub> emission and improve fuel combustion.

The basic steps of producing ethanol from lignocellulosic materials are pretreatment, hydrolysis, and fermentation. Cellulase complex can hydrolyze cellulose into monomeric sugars. These monomeric sugars can be converted to ethanol by simultaneous saccharification and fermentation (SSF) method, for example by *M. purpureus* CBS 109.07 and *F. venenatum* ATCC 20334. The previous study by Takeshita et al. (2016) showed that combination of *Monascus purpureus* NBRC 5965 and *Saccharomyces cerevisiae* K7 could produce ethanol in alcoholic beverages. The results showed that *M. purpureus* has the

capability as saccharification agent and ethanol producer. *Monascus purpureus* is not only used as a food colorant, flavoring agent, and preservative, but it is also widely applied in medical purposes to lower blood cholesterol, anti-diabetes, anti-inflammatory and to prevent osteoporosis (Arunachalam and Narmadhapriya 2011). *F. venenatum* has been cultured as a mycoprotein for a food source. Food products from *F. venenatum* have a fibrous texture and become rich source of high-quality protein, including essential amino acids (Hosseini and Khosravi-Darani 2010). *M. purpureus* CBS 109.07 and *F. venenatum* ATCC 20334 are still considered as new strains especially for ethanol production from lignocellulose.

The wheat straw used in this study had been pretreated using acid method to render the crystalline structure of the cellulose. The solid fraction of pretreated wheat straw contains high cellulose that can be used by fungi to produce ethanol in SSF, while the liquid fraction contains monomeric sugars. A mixture of solid and liquid fractions can be used as substrate for ethanol production in SSF. In SSF, glucose released during biomass saccharification strongly inhibits enzymes, particularly  $\beta$ -D-glucosidase that catalyzes hydrolysis of cellobiose to glucose. In SSF, glucose would be continuously converted to ethanol through a fungal fermentation process so that inhibition of enzymes by glucose can be avoided. To the best of our knowledge, there have been no studies on ethanol

production by *M. purpureus* CBS 109.07 and *F. venenatum* ATCC 20334 from wheat straw. Therefore, *M. purpureus* (CBS 109.07) together with *F. venenatum* ATCC 20334 were used to produce ethanol from wheat straw in this study. Therefore, the objective of this study was to evaluate ethanol production by *M. purpureus* CBS 109.07 and *F. venenatum* ATCC 20334 from a solid, liquid, mixed fraction (solid and liquid fractions) of pretreated wheat straw by SSF method.

## MATERIALS AND METHODS

### The Microorganisms

*Monascus purpureus* CBS 109.07 and *F. venenatum* ATCC 20334 used in this study were obtained from Swedish Center for Resource Recovery, University of Borås, Sweden. These fungi were maintained on Potato Dextrose Agar (PDA) at 4°C. The spore suspension was prepared by adding 20 ml of 0.05 M citrate buffer into agar plate. The spore suspension was dispersed with a disposable spreader.

### The Pretreated wheat straw

Pretreated wheat straw used in this study was obtained from the Swedish Center for Resource Recovery, University of Borås, Sweden. The chemical composition of the pretreated wheat straw is presented in Table 1. The wheat straw had been pretreated at SEKAB in Örnsköldsvik, Sweden. Pretreatment was conducted at 20 bar and pH 1.7 (adjusted with H<sub>2</sub>SO<sub>4</sub>) for 5-7 min. The pretreated wheat straw was stored at 4°C for further use.

Before fermentation process, the pretreated wheat straw was neutralized (detoxified) by sodium hydroxide 10 M because fungi could not grow on the pretreated wheat straw at very low pH condition. The pretreated wheat straw was diluted with deionized water (1:1) and added with 10 M sodium hydroxide until it reached pH 5.5. Filtration was carried out by filtering the pretreated wheat straw on the crucibles to separate the solid and the liquid fraction. The solid, the liquid, and the mixed fraction (a mixture of solid and liquid fractions) of pretreated wheat straw after detoxification process were individually used as substrates for *M. purpureus* CBS 109.07 and *F. venenatum* ATCC 20334 in the SSF process to produce ethanol. SSF of Avicel was also performed as reference (positive control).

**Table 1.** Chemical composition of pretreated wheat straw

Analysis	Concentration (% dry weight)
Total solid	19.74
Cellulose	41.13
Acid-insoluble lignin	29.10
Acid soluble lignin	0.71
Total lignin	29.82
Ash	6.45

### Enzymes activity

Two commercial enzymes, i.e. cellulase complex (NS22086) from Novozymes (Denmark) and  $\beta$ -glucosidase from almonds (Sigma-Aldrich, Germany) were used in all experiments as enzymatic complex. The activity of cellulase complex (NS22086) was 325 FPU/mL according to the method described by Yu et al. (2015). The activity of  $\beta$ -glucosidase was 2 units/mg solids.

### Cultivation of microorganisms

The media for culturing microorganisms contained (g/L): glucose 50; yeast extract 5; (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> 7.5; K<sub>2</sub>HPO<sub>4</sub> 3.5; MgSO<sub>4</sub>·7H<sub>2</sub>O 0.75; CaCl<sub>2</sub>·2H<sub>2</sub>O 1 and 0.05 M citrate buffer at pH 5.5. Fifty mL of medium in 250 mL cotton-plugged Erlenmeyer flasks were autoclaved at 121°C. After reaching room temperature; the medium was inoculated with the fungal microorganism. One milliliter of spore suspension, containing 1 x 10<sup>6</sup> spores/mL of *M. purpureus* CBS 109.07 or *F. venenatum* ATCC 20334, was added to each flask. The fungal culture was incubated for 30 hours at 30°C at 150 rpm. At the end of incubation, the content of Erlenmeyer flasks was aseptically centrifuged and was further used in SSF.

### Fermentation of pretreated wheat straw

SSF was performed under anaerobic conditions. The media contained (g/L): yeast extract 5; (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> 7.5; K<sub>2</sub>HPO<sub>4</sub> 3.5; MgSO<sub>4</sub>·7H<sub>2</sub>O 0.75; CaCl<sub>2</sub>·2H<sub>2</sub>O 1; either solid fraction of pretreated wheat straw 250 or a mixture of solid and liquid fraction of pretreated wheat straw 250 or pure cellulose (Avicel) 50; and 0.05 M buffer citrate. Meanwhile, the liquid fraction of pretreated wheat straw had a concentration of 100 mL/L media. The final volume in each flask was 100 mL. The pH of the media was adjusted to 5.5 with the addition of 10 M NaOH. All the media were autoclaved at 121°C for 15 min. The media were then inoculated with 1 g of the fungus (dry base). The required enzymes were also added into each flask aseptically. All the SSF experiments were performed at 30°C. The enzyme loading was 15 FPU/g cellulose and 30 IU/g total solid of pretreated wheat straw. The flasks were equipped with a loop-trap containing water and sterile plastic tubes as well as clamps for taking liquid samples Christia et al. (2016).

### Analysis of pretreated wheat straw

The pretreated wheat straw was analyzed for the ash content (Isroi et al. 2012), total solids (Isroi et al. 2012), total lignin (Sluiter et al. 2012) and cellulose content (Sluiter et al. 2012). The samples from SSF were analyzed by a High-Performance Liquid Chromatography (HPLC) (Isroi et al. 2012), which was equipped with UV/vis and RI detectors (Waters, UK). Cellobiose, glucose, xylose, and mannose were analyzed in the Biorad Aminex HPX-87P column (Bio-Rad, Richmond, CA, USA) equipped with the appropriate guard column with the HPLC conditions as follows: 25  $\mu$ L injection volume, 0.2  $\mu$ m filtered and degassed HPLC grade water mobile phase, 0.6 mL/minute flow rate, 85°C column temperature, 35 minutes run time. The analysis of ethanol, xylitol, glycerol, furfural, lactic,

acetic, and succinic acids were conducted according to the National Renewable Energy Laboratory (NREL) procedure that has been modified by Isroi et al. (2012). Those compounds were analyzed in the Biorad Aminex HPX-87H column (Bio-Rad, Richmond, CA, USA) equipped with the appropriate guard column with the HPLC conditions as follows: 25  $\mu$ L injection volume, 0.2  $\mu$ m filtered and degassed 0.005 M sulfuric acid mobile phase, 0.6 mL/minute flow rate, 60°C column temperature, 50 minutes run time.

## RESULTS AND DISCUSSION

### Composition of pretreated wheat straw

It can be seen in Table 1 that the percentages of total solids and cellulose were 19.74% and 41.13%, respectively. The percentage of cellulose was considerably high. Cellulose in wheat straw comprises of cellulose I allomorph with low crystallinity, and the stable cellulose I $\beta$  with polymorphic crystal structure (Volynets and Dahman 2011). Cellulose microfibrils are connected by amorphous serrated area in longitudinal formation in the epidermis, while in the parenchyma, they have a random arrangement. Cellulose microfibrils are entrapped in hemicellulose by hydrogen and covalent bond, which are stuck together by lignin. Yang et al. (2018) stated that the cellulose content in parenchyma is higher than in the epidermis. This phenomenon also occurs in every part in tissues, especially in vascular bundles and parenchyma tissue during plant growth. Furthermore, there is a similar crystallinity character of wheat straw's cellulose in different parts (Zhang et al. 2014). In epidermis, the cellulose is orientated along the growth direction, whereas in the parenchyma, there is no preferred orientation along the growth direction.

Total lignin of the pretreated wheat straw was 29.82%, which contains the acid-insoluble lignin (29.1%) and the soluble acid lignin (0.71%). A study by Qi et al. (2010) showed that the acid-insoluble lignin in pretreated wheat straw ranging from about 16% (pretreated with 0% sulfuric acid) to 33% (pretreated with 3% sulfuric acid). More severe pretreatment conditions resulted in higher acid-insoluble lignin in the pretreated solids (Qi et al. 2010). Total lignin in this study was quite high (29.82% of dry weight). This result indicates that there was no destruction of the lignin structure during pretreatment process. It means that only lignin bound to hemicelluloses that were damaged so that the lignin content in the substrate remains high. However, there is a possibility that chemical bonds between hemicellulose and cellulose may inhibit the cellulolytic enzyme activity. The presence of lignin and hemicellulose inhibits cellulolytic enzyme activity because of the structure of hemicellulose, which traps cellulose. The cellulolytic enzyme will have difficulty in hydrolyzing cellulose directly.

The ash content in the pretreated wheat straw was 6.45%. This value was slightly different from the ash content of raw wheat straw (7.9%) (Bakisgan et al. 2009). The differences in ash content might be caused by the acid pretreatment process carried out to wheat straw in this

study. Inorganic materials have association with lignocellulosic components especially hemicellulose in some biomass ashes (Vassilev et al., 2014). Acid pretreatment leads to degradation of lignocellulosic components, i.e. lignin and hemicellulose to its monomers. Whereas, untreated wheat straw still has a lot of lignocellulosic components associated with inorganic materials, which could be taken into account as ash during analysis.

### Simultaneous saccharification and fermentation by *M. purpureus* CBS 109.07


SSF of pretreated wheat straws (solid, liquid, and mixed solid and liquid fraction) were carried out under anaerobic condition. SSF with Avicel was also conducted as a reference/positive control. Cellulase and  $\beta$ -glucosidase enzymes produced glucose from the cellulose part of the pretreated wheat straws and *M. purpureus* CBS 109.07 simultaneously assimilated the glucose to ethanol. These enzymes help to increase the amount of ethanol yield during fermentation. *M. purpureus* CBS 109.07 is a cellobiose-utilizing microorganism, which can facilitate saccharification of cellulose. Fermentation of *M. purpureus* CBS 109.07 in the liquid fraction of pretreated wheat straw resulted in the highest maximum ethanol yield in SSF (0.41 g/g glucose) (Table 2), while the highest rate of ethanol production was obtained on the 2<sup>nd</sup> to the 3<sup>rd</sup> day of the fermentation process (Figure 1a). The ethanol concentration remained constant on the last 4 days of fermentation of the solid fraction, but it decreased slowly in the liquid fraction as well as in mixed (solid and liquid) fraction. Ethanol concentration increased in Avicel media. The increased ethanol production by *M. purpureus* CBS 109.07 in solid, liquid, and mixed fraction of pretreated wheat straw in SSF occurred on day-1 to day-3 of the fermentation process. The maximum ethanol concentration in Avicel medium was obtained on day-7 (Figure 1.A).

The concentration profiles of glucose, cellobiose, xylose, and mannose are presented in Figure 1.B-E. All of the simple sugars are intermediate products in the bioconversion chains from lignocellulosic biomass to ethanol. The concentrations of cellobiose, xylose, and mannose throughout all the experiments except glucose were less than 0.2, 0.8, and 0.51 g/L, respectively. The maximum glucose concentrations in a solid, liquid, a mixture of solid and liquid fractions of pretreated wheat straw, and Avicel were 17.94, 4.69, 8.81, 36.19 g/L, respectively. Each medium had different total solid content, except the liquid fraction of pretreated wheat straw. The total solid contents of the solid, a mixture of solid and liquid fractions of pretreated wheat straw, and Avicel were 4.93, 2.46, and 5 g, respectively. The cellulose contents of solid, a mixture of solid and liquid, and Avicel were 2, 1, and 5, respectively. The cellulose contents were calculated based on cellulose content of pretreated wheat straw (41.13% dry weight). Therefore, the maximum glucose concentrations in a solid, liquid, a mixture of solid and liquid fractions of pretreated wheat straw, and Avicel were different.


Furthermore, Kootstra et al (2009) stated that xylan was the dominant sugar in hemicelluloses of wheat straw (19%). Therefore, if the microorganisms can utilize both hexose and xylose that are present in the hydrolyzate, higher ethanol yield can be expected. There are several microorganisms that can produce ethanol from xylose. The results of the study by Qi et al. (2010) showed that the liquid fraction of pretreated wheat straw using 2% and 3% acid concentration yield the highest glucose content. Furthermore, study by Qi et al. (2010) also showed that xylose was the main monomeric sugar in all of the liquid fraction of pretreated wheat straw, and its yield was also significantly affected by acid concentration. Increasing acid concentration led to increased xylose yield. According to Qi et al. (2010), the highest xylose yield was 0.23 g

xylose/g wheat straw when the straw was pretreated with 2% sulfuric acid. Acid concentration >2% caused a slight decrease in xylose yield. Higher acid concentration causes further degradation of xylose to furfural.

Acetic acid was the major byproduct of SSF by *M. purpureus* CBS 109.07. The highest concentration of acetic acid was obtained from a liquid fraction of pretreated wheat straw (0.23 g/g biomass). The time courses of acetic acid production were similar to the time courses of ethanol. No further accumulation of acetic acid was obtained after 4 days of SSF. The yields of the other metabolites (xylitol, glycerol, furfural, succinic, and lactic acids) were negligible in all the SSF experiments by *M. purpureus* CBS 109.07, which were in the range of 0-0.08 g/g biomass (Table 2).



**Figure 1.** Concentration Profile of: (a). Ethanol, (b). Glucose, (c). Cellobiose, (d). Xylose, and (e). Mannose during SSF of Pretreated Wheat Straw and Avicel by *M. purpureus* CBS 109.07. (A) = solid fraction of pretreated wheat straw, (B) = liquid fraction of pretreated wheat straw, (C) = a mixture of the solid and liquid fraction of pretreated wheat straw



**Figure 2.** Concentration Profile of (a). Ethanol, (b). Glucose, (c). Cellobiose, (d). Xylose, and (e). Mannose during SSF of Pretreated Wheat Straw and Avicel by *F. venenatum* ATCC 20334. (A) = solid fraction of pretreated wheat straw, (B) = liquid fraction of pretreated wheat straw, (C) = a mixture of the solid and liquid fraction of pretreated wheat straw

### Simultaneous Saccharification and Fermentation by *F. venenatum* ATCC 20334

Ethanol production from pretreated wheat straw in SSF was also conducted using *F. venenatum* ATCC 20334. SSF of Avicel was also carried out as a control. The results are presented in Figure 2 and Table 2. *F. venenatum* ATCC 20334 is a filamentous fungus quite similar to *M. purpureus* CBS 109.07 in terms of formation of the metabolites, in which ethanol was the main metabolite and the main byproduct was acetic acid. The profile of ethanol formation (Figure 2.a) was quite similar to the *M. purpureus* CBS 109.07, where the major part of ethanol was produced in the first 2 to 3 days of fermentation. The concentration of ethanol remained constant within the last 4 days of SSF of the liquid fraction and a mixture of the solid and liquid fraction of pretreated wheat straw. However, it decreased slowly in the solid fraction of pretreated wheat

straw and increased in Avicel. The results showed that ethanol production by *F. venenatum* ATCC 20334 in all the substrates occurred on day-1 to day-4. However, it took about 7 days or more to reach the maximum ethanol concentration in Avicel (Figure 2a). The best ethanol yield by *F. venenatum* ATCC 20334 was 0.54 g/g glucose, which was obtained from the liquid fraction of pretreated wheat straw in SSF (Table 2). Enzymes act to enhance the overall ethanol yield from cellulose. Sakamoto et al. (2012) showed that addition of hemicellulolytic enzymes increased the production of monomeric sugars and ethanol from hemicellulosic materials of rice straw. The ethanol was produced not only from glucose but also from other monomeric sugars, such as cellobiose, xylose, and mannose. Therefore, if all those monomeric sugars could be assimilated, the theoretical ethanol yield could be higher than that of Table 2.

**Table 2.** Yield of ethanol and byproducts and ethanol concentration in simultaneous saccharification and fermentation of dilute-acid pretreated wheat straw with *M. purpureus* CBS 109.07 and *F. venenatum* ATCC 20334.

Strain	Substrate	Total solid (g/L)	Cellulose content (g/L)	Max ethanol concentration (g/L)	Max theoretical ethanol yield (%)	Max Yield (g/g glucose)						
						Ethanol	Xylitol	Succinic acid	Lactic acid	Glycerol	Acetic acid	Furfural
<i>M. purpureus</i> CBS 109.07	A	49.35	20.30	8.08	70.26 <sup>a</sup>	0.36	0.00	0.01	0.08	0.01	0.16	0.00
	B	-	4.69*	1.92	80.06 <sup>b</sup>	0.41	0.00	0.06	0.00	0.02	0.23	0.00
	C	24.68	10.15	4.19	72.86 <sup>a</sup>	0.37	0.00	0.01	0.02	0.01	0.11	0.00
	Avicel	50.00	50.00	9.07	32.11 <sup>a</sup>	0.16	0.00	0.00	0.04	0.00	0.01	0.00
<i>F. venenatum</i> ATCC 20334	A	49.35	20.30	4.79	41.65 <sup>a</sup>	0.21	0.00	0.00	0.11	0.01	0.02	0.01
	B	-	3.98*	2.16	78.62 <sup>b</sup>	0.54	0.02	0.05	0.05	0.02	0.32	0.01
	C	24.68	10.15	3.89	67.65 <sup>a</sup>	0.35	0.01	0.01	0.03	0.01	0.11	0.01
	Avicel	50.00	50.00	6.38	22.59 <sup>a</sup>	0.12	0.00	0.00	0.04	0.00	0.04	0.00

Note: \* Glucose content in Liquid Fraction of Pretreated Wheat Straw. <sup>a</sup> Max theoretical ethanol yield (solid, both of solid and liquid fraction, Avicel) = [(max produced ethanol (g/L))/[0.51×1.111×dry weight of biomass (g/L)×F]]×100. F = cellulose fraction in biomass = 0.997 for pure cellulose (Avicel) and 0.4113 for pretreated wheat straw. <sup>b</sup> Max theoretical ethanol yield (liquid fraction) = [(max produced ethanol (g/L))/[initial glucose concentration in liquid fraction of pretreated wheat straw (g/L)]]×100. <sup>c</sup> Max yield = [(max produced (g/L)-initial production (g/L))/[1.111×dry weight of biomass (g/L)×F]], where F = cellulose fraction in biomass = 0.997 for pure cellulose (Avicel) and 0.4113 for pretreated wheat straw. A = solid fraction of pretreated wheat straw. B = liquid fraction of pretreated wheat straw. C = a mixture of the solid and liquid fraction of pretreated wheat straw

In this study, ethanol yield in SSF of Avicel was lower than that of solid, liquid, and a mixture of solid and liquid fractions of pretreated wheat straw in all of the experiments (Table 2). This is similar to the results of the previous works, in which glucose and ethanol yields of pretreated sugar cane bagasse by cellulignin were higher than that of pure cellulose Avicel (Ferreira et al. 2010). Some other factors may also affect the ethanol yield. Physical properties and cellulose microstructure are among the potential factors that affect enzymatic hydrolysis (Ogeda et al. 2012). Furthermore, Avicel was purchased in dried form. It was reported that drying might damage the pores of the biomass irreversibly (Amit et al. 2018). Another reason of the lower yield in SSF of Avicel might be caused by the higher ethanol concentration in SSF of Avicel, which could reduce the activity of the enzyme (Márkus et al. 2017) and viability of microorganism (Chauhan et al. 2013).

The presence of residues of hemicellulose and lignin in the pretreated wheat straw could reduce the performance of the fungi during the bioconversion process of sugars to ethanol. It makes the ethanol yield in liquid fraction of pretreated wheat straw higher than that of in solid fraction and in a mixture of solid and liquid fraction of pretreated wheat straw.

The concentration profiles of glucose, cellobiose, xylose, and mannose are presented in Figure 2b-e. The concentrations of cellobiose, xylose, and mannose in SSF by *F. venenatum* remained under 0.22, 0.76, and 0.26 g/L, respectively throughout all the experiments except glucose. The maximum glucose concentrations resulted from a solid, liquid, mixture of solid and liquid fractions of pretreated wheat straw, and Avicel were 17.49, 3.98, 8.23, 39.93 g/L, respectively. The total solids and cellulose content of the solid, a mixture of solid and liquid fractions

of pretreated wheat straw, and Avicel were similar to SSF by *M. purpureus*. *F. venenatum* also produced acetic acid as the main byproduct and the highest yield of acetic acid was obtained from the fermentation of liquid fraction of pretreated wheat straw (0.32 g/g biomass), while the yields of acetic acid in solid, a mixture of solid and liquid fraction of pretreated wheat straw, and Avicel were 0.02 and 0.14 g/g biomass (Table 2). The yields of the other metabolites were negligible in all SSF experiments by *F. venenatum* ATCC 20334. The maximum yield of xylitol was in the range of 0-0.02, succinic acid was 0-0.05, lactic acid was 0.03-0.11, glycerol was 0-0.02, and furfural was 0-0.01 g/g biomass (Table 2).

Based on the results of this study, it can be concluded that *M. purpureus* CBS 109.07 and *F. venenatum* ATCC 20334 were able to produce ethanol from liquid and solid fraction of pretreated wheat straw. *M. purpureus* CBS 109.07 produced higher ethanol in the liquid and the solid fraction of pretreated wheat straw than *F. venenatum* ATCC 20334 did.

## ACKNOWLEDGEMENTS

This work was financially supported by The Swedish Council for Higher Education through Linnaeus-Palme Program.

## REFERENCES

- Amit K, Nakachew M, Yikal B, Mukesh Y. 2018. A review of factors affecting enzymatic hydrolysis of pretreated lignocellulosic biomass. Res J Chem Environ 22 (7): 62-67

- Arunachalam C, Narmadhapriya D. 2011. *Monascus* fermented rice and its beneficial aspects: a new review. *Asian J Pharm Clin Res* 4 (1): 29-31.
- Bakışgan C, Dumanli AG, Yürüm Y. 2009. Trace elements in Turkish biomass fuels: Ashes of wheat straw, olive bagasse and hazelnut shell. *Fuel* 88: 1842-1851.
- Brander M, Hutchison C, Sherrington C, Ballinger A, Beswick C, Baddeley A, et al. 2009. Methodology and evidence base on the indirect greenhouse gas effects of using wastes, residues, and by-products for biofuels and bioenergy—appendix 10 wheat straw case study—Data and Assumptions, In: Davis G, (editor). Report to RFA and DECC. Ecometrica.
- Chauhan NM, Shinde RB, Karuppaiyl SM. 2013. Effect of alcohols on filamentation, growth, viability and biofilm development in *Candida albicans*. *Braz J Microbiol* 44 (4): 1315-1320.
- Christia A, Setiowati AD, Millati R, Karimi K, Cahyanto MN, Niklasson C, Taherzadeh MJ. 2016. Ethanol production from alkali- pretreated oil palm empty fruit bunch by simultaneous saccharification and fermentation with *Mucor indicus*. *Intl J Green Energ* 13 (6): 566-572.
- Ferreira V, Faber MO, Mesquita SS, Pereira N. 2010. Simultaneous saccharification and fermentation process of different cellulosic substrates using a recombinant *Saccharomyces cerevisiae* harboring the  $\beta$ -glucosidase gene. *Electron J Biotechnol* 13 (2): 1-7.
- Hammond GP, Mansell RM. 2018. A comparative thermodynamic evaluation of bioethanol processing from wheat straw. *Appl Energ* 224: 136-146.
- Hosseini S, Khosravi-Darani K. 2010. Response surface methodology for mycoprotein production by *F. venenatum* ATCC 20334. *J Bioprocess Biotechnol* 1: 102. DOI: 10.4172/2155-9821.1000102
- Isroi, Ishola MM, Millati R, Syamsiah S, Cahyanto MN, Niklasson C, Taherzadeh MJ. 2012. Structural changes of oil palm empty fruit bunch (OPEFB) after fungal and phosphoric acid pretreatment. *Molecules* 17: 14995-15012.
- Kootstra AM J, Beftink H H, Scott EL, Sanders JPM. 2009. Comparison of dilute mineral and organic acid pretreatment for enzymatic hydrolysis of wheat straw. *Biochem Eng J* 46 (2): 126-131.
- Kootstra AMJ, Beftink HH, Scott EL, Sanders JPM. 2009. Optimization of the dilute maleic acid pretreatment of wheat straw. *Biotechnol Biofuels* 2: 31 DOI: 10.1186/1754-6834-2-31.
- Márkus Z, Bakó KB, Tóth G, Nemestóthy N, Gubicza L. 2017. Effect of chain length and order of alcohol on enzyme activity during enzymatic esterification in organic media. *Hung J Ind Chem* 45 (2): 35-39.
- Ogeda TL, Silva IB, Fidale LC, Seoud OAE, Petri DFS. 2012. Effect of cellulose physical characteristics, especially the water sorption value, on the efficiency of its hydrolysis catalyzed by free or immobilized cellulase. *J Biotechnol* 157: 246- 252.
- Qi B, Chen X, Wan Y. 2010. Pretreatment of wheat straw by nonionic surfactant-assisted dilute acid for enhancing enzymatic hydrolysis and ethanol production. *Bioresour Technol* 101 (13): 4875-4883.
- Sakamoto T, Hasunuma T, Hori Y, Yamada R, Kondo A. 2012. Direct ethanol production from hemicellulosic materials of rice straw by use of an engineered yeast strain codisplaying three types of hemicellulolytic enzymes on the surface of xylose-utilizing *Saccharomyces cerevisiae* cells. *J Biotechnol* 158 (4): 203-210.
- Sluiter A, Hames B, Ruiz R, Scarlata C, Sluiter J, Templeton D, Crocker D. 2012. Determination of Structural Carbohydrates and Lignin in Biomass. Laboratory Analytical Procedure (LAP), National Renewable Energy Laboratory (NREL). Colorado.
- Takeshita R, Saigusa N, Teramoto Y. 2016. Production and antioxidant activity of alcoholic beverages made from various cereal grains using *Monascus purpureus* NBRC 5965. *J Inst Brew* 122: 350-354.
- Vassilev SV, Vassileva CG, Baxter D. 2014. Trace element concentration and associations in some biomass ashes. *Fuel* 129: 292-313
- Volynets B, Dahman Y. 2011. Assessment of pretreatments and enzymatic hydrolysis of wheat straw as a sugar source for bioprocess industry. *Intl J Energ Environ* 2 (3): 427-446.
- Yang Z, Mei J, Liu Z, Huang G, Huang G, Han L. 2018. Visualization and semiquantitative study of the distribution of major components in wheat straw in mesoscopic scale using Fourier transform infrared microspectroscopic imaging. *Anal Chem* 90 (12): 7332-7340.
- Yu X, Liu Y, Cui Y, Cheng Q, Zhang Z, Lu JH, Meng Q, Teng L, Ren X. 2015. Measurement of filter paper activities of cellulose with microplate-based assay. *Saudi J Biol Sci* 23: S93-S98.
- Zhang H, Thygesen LG, Mortensen K, Kádár Z, Lindedam J, Jørgensen H, Felby C. 2014. Structure and enzymatic accessibility of leaf and stem from wheat straw before and after hydrothermal pretreatment. *Biotechnol Biofuels* 7: 74. DOI: 10.1186/1754-6834-7-74.