



CHALMERS
UNIVERSITY OF TECHNOLOGY

Symmetries of M-theory and free Lie superalgebras

Downloaded from: <https://research.chalmers.se>, 2026-06-09 02:39 UTC

Citation for the original published paper (version of record):

Gomis, J., Kleinschmidt, A., Palmkvist, J. (2019). Symmetries of M-theory and free Lie superalgebras. *Journal of High Energy Physics*, 2019(3). [http://dx.doi.org/10.1007/JHEP03\(2019\)160](http://dx.doi.org/10.1007/JHEP03(2019)160)

N.B. When citing this work, cite the original published paper.

Symmetries of M-theory and free Lie superalgebras

Joaquim Gomis,^a Axel Kleinschmidt^{b,c} and Jakob Palmkvist^d

^a*Departament de Física Quàntica i Astrofísica and Institut de Ciències del Cosmos (ICCUB),
Universitat de Barcelona,*

Martí i Franquès, ES-08028 Barcelona, Spain

^b*Max-Planck-Institut für Gravitationsphysik (Albert-Einstein-Institut),
Am Mühlenberg 1, DE-14476 Potsdam, Germany*

^c*International Solvay Institutes,
ULB-Campus Plaine CP231, BE-1050 Brussels, Belgium*

^d*Division for Theoretical Physics, Department of Physics, Chalmers University of Technology,
SE-412 96 Gothenburg, Sweden*

E-mail: joaquim.gomis@ub.edu, axel.kleinschmidt@aei.mpg.de,
jakob.palmkvist@chalmers.se

ABSTRACT: We study systematically various extensions of the Poincaré superalgebra. The most general structure starting from a set of spinorial supercharges Q_α is a free Lie superalgebra that we discuss in detail. We explain how this universal extension of the Poincaré superalgebra gives rise to many other algebras as quotients, some of which have appeared previously in various places in the literature. In particular, we show how some quotients can be very neatly related to Borcherds superalgebras. The ideas put forward also offer some new angles on exotic branes and extended symmetry structures in M-theory.

KEYWORDS: Global Symmetries, M-Theory, Space-Time Symmetries, Supersymmetric Gauge Theory

ARXIV EPRINT: [1809.09171](https://arxiv.org/abs/1809.09171)

Contents

1	Introduction	1
2	Review of Maxwell and free Lie (super)algebras	3
2.1	Maxwell algebra and free Lie algebras	3
2.2	Free Lie superalgebras	6
3	Instances of free Lie superalgebras	8
3.1	$D = 4$ and $\mathcal{N} = 1$ supersymmetry	8
3.1.1	Free Lie superalgebra	8
3.1.2	Comparison to Maxwell superalgebra in the literature	10
3.2	$D = 4$ and extended supersymmetry	11
3.2.1	$D = 4$ and $\mathcal{N} = 2$ supersymmetry	12
3.2.2	$D = 4$ and $\mathcal{N} > 2$ supersymmetry	12
3.3	$D = 11$ and $\mathcal{N} = 1$ supersymmetry	13
3.3.1	Free Lie superalgebra	13
3.3.2	Relation to E_{11} and ℓ_1 representation?	15
4	Free Lie superalgebras and Borchers superalgebras	18
4.1	$D = 4$ and $\mathcal{N} = 1$ supersymmetry	19
4.2	$D = 4$ and extended supersymmetry	24
4.3	$D = 11$ and $\mathcal{N} = 1$ supersymmetry	26
5	Conclusion	27
A	Gamma matrices in $D = 4$	28

1 Introduction

Translations P_a in the global Poincaré space-time symmetry algebra commute among themselves: $[P_a, P_b] = 0$. This is no longer true when one considers translations in the presence of a constant electro-magnetic background field [1–3] and the study of such deformations of the abelian algebra of translations has led to the investigation of so-called Maxwell algebras [3–6]. The most general deformation one can allow for (based on Lie algebra cohomology) is

$$[P_a, P_b] = Z_{ab}, \tag{1.1}$$

where the anti-symmetric generator $Z_{ab} = Z_{[ab]}$ is associated with the constant electro-magnetic background field and transforms as a tensor under Lorentz transformations. In

a next step one can analyse the commutator of Z_{ab} with P_a and iterate the procedure [6]. It was shown in [7] that the most general structure one obtains in this way is the infinite-dimensional *free* Lie algebra, called Maxwell_∞ there, generated by the translations P_a . Any free Lie algebra admits many Lie algebra ideals and associated quotient Lie algebras, some of which were studied for Maxwell_∞ in [7]. In particular [7] identified quotients that could be used in a particle model to describe motion in a general, non-constant, electro-magnetic field in an ‘unfolded’ treatment of the electro-magnetic field strength. This was achieved by introducing additional coordinates for the additional generators contained in the quotient of Maxwell_∞ . Associating the translations P_a with a coordinate x^a one in this way has an extended space with coordinates θ^{ab} for Z_{ab} and so on. The dynamical model based on the Maxwell_∞ algebra describes the dynamical object together with the background it moves in.

In the present article, we shall extend the picture of the free Lie algebra generalisation of the Poincaré algebra to a free Lie superalgebra generalisation of the Poincaré superalgebra. The basic building blocks in this construction are the (global) odd supersymmetry generators Q_α (with α an appropriate spinor index). The infinite-dimensional free Lie superalgebra generated by the Q_α has the most general generators in $\{Q_\alpha, Q_\beta\}$ and subsequent multi-(anti-)commutators. Again, it admits many non-trivial quotients that we shall discuss. Of particular interest to us will be quotients related to the eleven-dimensional supersymmetry algebra. As has been proved for example in [8, 9], the anti-commutator $\{Q_\alpha, Q_\beta\}$ should yield not only the translation generators P_a but also additional non-central terms that serve as sources for branes. These additional terms arise naturally in the free Lie superalgebra as we will show explicitly. Further extensions of the eleven-dimensional supersymmetry algebra were also studied in [10] (called ‘M-algebra’ there) where non-trivial multi-commutators arise. As we shall describe in more detail below, these again have a natural home in the free Lie superalgebra. The interpretation given in [10] of some of these generators as superpartners of the bosonic brane charges might give some idea of how to interpret the infinitely many new generators in the free Lie superalgebra. We will also discuss other possible connections to proposals of infinite-dimensional M-theory symmetries and extended space-times.

Studying the free Lie superalgebra in four space-time dimensions we are also able to make contact with the supersymmetric extension of the Maxwell algebra that was studied in [11] as it arises as a particular finite-dimensional quotient. In [11], a dynamical realisation of the supersymmetric Maxwell algebra was found in terms of a massless superparticle where again additional coordinates are introduced for the new generators and these have interpretations in terms of a Maxwell superfield background. Similar generalisations of the Maxwell algebra for extended supersymmetry have also been investigated [12] and we shall also exhibit how these \mathcal{N} -extended versions can be obtained from a free Lie superalgebra.

Our paper focusses almost exclusively on the algebraic aspects unifying the various algebras in terms of free Lie superalgebras. We shall not consider particle or string models realising these symmetries but it should be possible to construct these along the lines of [13].

An interesting observation, that also opens up connections to other algebraic structures investigated in the literature in connection with potential symmetries of M-theory, is that

many quotients of the free Lie superalgebra embed in Kac-Moody [14–16], Borchers [17–19] or tensor hierarchy algebras [20–23]. This is due to the definition of these algebras as being constructed as quotients of free algebras. A noteworthy difference is, however, that many of these conjectured algebraic structures in M-theory also have ‘negative level’ symmetry generators that are not present in our free Lie superalgebra constructions. This connection is discussed in detail in section 4.

The paper is structured as follows. We first review and motivate the free Lie algebra construction in relation to the bosonic Maxwell algebra and then introduce the free Lie superalgebra that is the main object of study in this paper. In section 3, we then discuss in detail various examples that are related to simple and extended supersymmetry in $D = 4$ and simple supersymmetry in $D = 11$. We also explain how our construction relates to various algebras in the literature and, potentially, to exotic branes. Section 4 contains a discussion of cases where quotients of the free Lie superalgebra can be re-expressed through Borchers superalgebras and how this in turn can be recast in terms of Kac-Moody algebras. Section 5 contains some concluding remarks and speculations.

2 Review of Maxwell and free Lie (super)algebras

In this section, we review some basic aspects of the conventional bosonic Maxwell algebra in order to motivate our generalisations first to free Lie algebras [7] and then to free Lie superalgebras.

2.1 Maxwell algebra and free Lie algebras

The starting point for all bosonic constructions here is the Poincaré algebra in D space-time dimensions,¹

$$[M_{ab}, M_{cd}] = 2\eta_{c[b}M_{a]d} - 2\eta_{d[b}M_{a]c}, \tag{2.1a}$$

$$[M_{ab}, P_c] = 2\eta_{c[b}P_{a]}, \tag{2.1b}$$

$$[P_a, P_b] = 0. \tag{2.1c}$$

The first line is just the Lorentz algebra $\mathfrak{so}(1, D - 1)$ of the anti-symmetric M_{ab} and the second line expresses that the translation generators P_a form a vector representation of the Lorentz algebra. The third line (2.1c) is the standard property of the Poincaré algebra that the translations commute. This is the relation that is deformed in the Maxwell algebra and its free Lie algebra generalisation.

As shown in [6] based on Lie algebra cohomology, the most general deformation of (2.1c) is

$$[P_a, P_b] = Z_{ab} \tag{2.2}$$

¹Our anti-symmetrisation convention is of strength one: $M_{[ab]} = \frac{1}{2}(M_{ab} - M_{ba})$. The fundamental Lorentz indices lie in the range $a, b \in \{0, \dots, D - 1\}$. We use the mostly plus convention for the flat Minkowski metric.

with a new generator $Z_{ab} = Z_{[ab]}$. This is also the commutator in the free Lie algebra based on the P_a by giving a name to $[P_a, P_b]$. The algebra spanned by $\{M_{ab}, P_a, Z_{ab}\}$ with trivial commutators $[P_a, Z_{bc}] = [Z_{ab}, Z_{cd}] = 0$ is the Maxwell algebra introduced in [3–6].

From the Lie algebra cohomology perspective, it is possible to deform also these trivial commutation relations. The next most general commutator is [6, 7]

$$[[P_a, P_b], P_c] = [Z_{ab}, P_c] =: Y_{ab,c}. \tag{2.3}$$

For this to form a consistent Lie algebra, the Jacobi identity has to be obeyed. This implies that the Lorentz tensor $Y_{ab,c}$ has the property $Y_{[ab,c]} = 0$. The comma notation for the indices reflects that the tensor also satisfies $Y_{[ab],c} = Y_{ab,c} = -Y_{ba,c}$. As a representation of the symmetric group it has the Young tableau $\begin{smallmatrix} \square & \square \\ \square \end{smallmatrix}$.²

This construction can be iterated to obtain all possible multiple commutators of the P_a up to the Jacobi identity (and anti-symmetry). This is by definition the *free Lie algebra* generated by the translation generators P_a and we denote it by $\mathfrak{f}(P)$. It has a natural graded structure given by how many times P_a appears in a multi-commutator. In this grading, we refer to the translation generators P_a as positive level $\ell = 1$, to their commutator $Z_{ab} = [P_a, P_b]$ as positive level $\ell = 2$ and so on.

The basic generators P_a on level $\ell = 1$ transform under the Lorentz algebra (as vectors) and consequently, we can make the generators at any level ℓ also transform under the Lorentz algebra. Thus, every level $\ell > 0$ of the free Lie algebra $\mathfrak{f}(P)$ consists of Lorentz tensors and we can form the semi-direct sum

$$\text{Maxwell}_\infty = \mathfrak{so}(1, D - 1) \oplus \mathfrak{f}(P) \tag{2.4}$$

with the $\mathfrak{so}(1, D - 1)$ Lorentz generators M_{ab} . This semi-direct sum is the infinite generalisation Maxwell_∞ of the Poincaré and Maxwell algebra introduced in [7]. We will refer to the algebra $\mathfrak{so}(1, D - 1)$ as level $\ell = 0$.

At this point it is important to remark that the construction of a free Lie algebra $\mathfrak{f}(P)$ only requires a set of D generators P_a , but that the transformation property of the P_a under level $\ell = 0$ is not something that is fixed by the free Lie algebra construction. In fact, we could also view the generators P_a as transforming in the fundamental of $\mathfrak{gl}(D)$ or any other Lie algebra \mathfrak{g}_0 with a D -dimensional representation. This representation need not even be irreducible. This level $\ell = 0$ algebra \mathfrak{g}_0 will then act on all the positive levels as well and we could arrange the higher levels in irreducible representations of \mathfrak{g}_0 . Choosing for instance $\mathfrak{g}_0 = \mathfrak{gl}(D)$, one obtains tensors of $\mathfrak{gl}(D)$ at higher levels. As $\mathfrak{gl}(D)$ does not have an invariant trace, the generator $Y_{ab,c}$ at $\ell = 3$ then is irreducible while it is reducible under $\mathfrak{so}(1, D - 1)$, see footnote 2. For yet different choices of \mathfrak{g}_0 acting on P_a each level ℓ arranges differently into irreducible representations of \mathfrak{g}_0 ; the total number of generators on each level of course does not change as this is determined by $\mathfrak{f}(P)$ which is defined independently of the choice of \mathfrak{g}_0 .³

²We note that this is not an irreducible representation of the Lorentz algebra as one can define the non-trivial trace $\eta^{bc} Y_{ab,c}$.

³Other interesting choices of \mathfrak{g}_0 relate to non-relativistic systems with a Galilean symmetry or very special relativity [24] where only the $\text{Sim}(D - 2)$ algebra with generators M_{+-} , M_{+i} and M_{ij} in a light-cone basis $\{x^\pm, x^i\}$ is kept [25].

The structure of a free Lie algebra $\mathfrak{f}(P)$ can be computed from a generating function [26] or explicitly using the Witt formula [27] that was reviewed in [7]. The generating function identity governing the free Lie algebra is

$$\bigotimes_{\ell=1}^{\infty} \left[\bigoplus_{k=0}^{\infty} (-1)^k t^{k\ell} \wedge^k T_{\ell} \right] = 1 - tT_1. \tag{2.5}$$

In this formula, t is a formal parameter and T_{ℓ} denotes the vector space of generators at level ℓ , such that $T_1 = \langle P_a \mid a = 0, \dots, D-1 \rangle$ and T_1 plays a special role since it fully determines $\mathfrak{f}(P)$. The identity (2.5) can be derived from the denominator formula for a Borcherds algebra. Expanding out the generating identity yields the following recursive identities at the first few powers of t :

$$T_1 = \langle P_a \rangle, \tag{2.6a}$$

$$T_2 = \wedge^2 T_1, \tag{2.6b}$$

$$T_3 = T_2 \otimes T_1 \ominus \wedge^3 T_1, \tag{2.6c}$$

$$T_4 = (T_3 \otimes T_1 \oplus \wedge^2 T_2) \ominus T_2 \otimes \wedge^2 T_1 \oplus \wedge^4 T_1, \tag{2.6d}$$

$$T_5 = (T_4 \otimes T_1 \oplus T_3 \otimes T_2) \ominus (T_3 \otimes \wedge^2 T_1 \oplus \wedge^2 T_2 \otimes T_1) \oplus T_2 \otimes \wedge^3 T_1 \ominus \wedge^5 T_1, \tag{2.6e}$$

...

The notation here is such that $\wedge^k R$ denotes the k -th anti-symmetric tensor power of a representation. With \ominus we denote the removal of representations from the module. The lowest levels are simple to understand as representing the anti-symmetry of the Lie bracket and the Jacobi identity of the free Lie algebra: T_2 consists of all commutators of the P_a in T_1 and therefore is the anti-symmetric product. The vector space T_3 is spanned by all commutators of P_a in T_1 with the elements in T_2 , as in (2.3), leading to the tensor product. The totally anti-symmetric Jacobi identity has to be satisfied by the triple commutators at $\ell = 3$ and so one has to remove $\wedge^3 T_1$ from the tensor product.

The tensor products here can be thought of as either just vector space operations or –and this will be more useful to us– as tensor products of representations of \mathfrak{g}_0 which is the level $\ell = 0$ algebra under which the P_a transform. The choice in (2.4) is $\mathfrak{g}_0 = \mathfrak{so}(1, D-1)$ and the tensor products would be those of Lorentz algebra representations.

Free Lie algebras admit many ideals and associated quotient Lie algebras. Several examples were discussed in [7] related to particle models in electro-magnetic backgrounds and previous examples in the literature. Any set of elements X in $\mathfrak{f}(P)$ generates an ideal, spanned by all multi-commutators obtained by acting on the elements X with the basic generators P_a , that is, the ideal is spanned by $[P, [P, \dots, [P, X]]]$. If the set X consists of only one element, then this is called a principal ideal. One can also consider ideals that are generated by a set which is a subspace of $\mathfrak{f}(P)$. In fact, any ideal generated by a set of elements X is the same as the ideal generated by the subspace $\langle X \rangle$ spanned by these elements.

The subspace $\langle X \rangle$ of $\mathfrak{f}(P)$ may be equal to T_{k+1} for some fixed finite level k . Then the ideal consists of all elements in $\mathfrak{f}(P)$ with more than k factors of P_a in the multi-commutator

T_ℓ	representation	generator
T_1	[2]	P_a
T_2	[2]	$Z_{ab} = \epsilon_{abc} Z^c$
T_3	$[2] \oplus [4]$	$Y_{ab,c} = \epsilon_{abd} \tilde{Y}^d{}_c$
T_4	$[2] \oplus [4] \oplus [6]$ [2]	$S_{ab,c,d}^1$ S_a^2
\vdots	\vdots	\vdots

Table 1. The free Lie algebra $\mathfrak{f}(P)$ in $D = 3$ space-time dimensions. Representations are labelled by the complexified Lorentz algebra of type A_1 while the generators represent the associated $\mathfrak{so}(1, 2)$ tensors. For mixed symmetry, these are reducible in the way described in footnote 2.

or, equivalently, the direct sum of all T_ℓ with $\ell > k$, and the quotient is a finite-dimensional truncation of $\mathfrak{f}(P)$ including only generators up to level k . For example, the truncation to $k = 1$ recovers the standard Poincaré algebra while the truncation to $k = 2$ gives the Maxwell algebra with P_a and Z_{ab} in the quotient.

More generally, the vector space $\langle X \rangle$ may be a subspace of T_{k+1} , and then the quotient may be infinite-dimensional. The Serre ideals appearing in the construction of Borcherds or Kac-Moody algebras discussed in section 4 are generated in this way. Even more generally, the vector space $\langle X \rangle$ may be decomposed into a direct sum $X = s_1 \oplus \dots \oplus s_n$ of finitely or infinitely many subspaces, or representations of \mathfrak{g}_0 . These representations s_i need not all be on the same level. Again, the quotient may be infinite-dimensional. An example of such an infinite-dimensional quotient was discussed in [7] where a realisation in terms of particles in general ‘unfolded’ electro-magnetic backgrounds was achieved.

As an example of a free Lie algebra that has not appeared in the literature before to the best of our knowledge, we consider the case of $D = 3$ translations P_a transforming under the Lorentz algebra $\mathfrak{g}_0 = \mathfrak{so}(1, 2)$ whose complexification is of Cartan type A_1 . The corresponding lowest levels are listed in table 1. Using standard mathematical Dynkin labels, the translation generators are denoted by [2].

The generators listed in table 1 can also be obtained by dimensional reduction from the general result in [7]. The three-dimensional case is of particular interest since one can find quotients of the Maxwell $_\infty$ algebra that admit an invariant and non-degenerate bilinear form that then can be used to construct three-dimensional Chern-Simons theory [28]. Since three-dimensional Chern-Simons theory is related to gravity, this leads to a Maxwell version of gravity [28–31] with a supersymmetric version studied in [32] and also the non-relativistic limit has been studied in this case [33]. A four-dimensional gravity system with Maxwell symmetry has been investigated in [34].

2.2 Free Lie superalgebras

The bosonic free Lie algebra $\mathfrak{f}(P)$ serves as a very universal object to construct kinematical algebras relevant in various contexts. It is natural to consider the corresponding Lie su-

peralgebra and in this section we give some general properties of its construction. Explicit examples will be discussed in the next section.

We shall consider the free Lie superalgebra generated by odd supertranslations Q_α where α labels the supersymmetry generators and we here think of them as spinorial generators. The exact range depends on the number of space-time dimensions and on whether one consider simple or extended supersymmetry. The general discussion of this section is independent of these details.

The free Lie superalgebra $\mathfrak{f}(Q)$ is given by all possible graded multi-commutators of the basic Q_α and subjected to the graded Jacobi identity

$$[\{Q_\alpha, Q_\beta\}, Q_\gamma] + [\{Q_\beta, Q_\gamma\}, Q_\alpha] + [\{Q_\gamma, Q_\alpha\}, Q_\beta] = 0. \quad (2.7)$$

‘Graded’ here refers to the \mathbb{Z}_2 -grading of superalgebras; the Q_α are by definition odd while their graded commutator $\{Q_\alpha, Q_\beta\}$ is even. One can again define a level ℓ on $\mathfrak{f}(Q)$ by counting the number of Q_α that occur in a general graded multi-commutator. For odd ℓ one obtains generators that are also odd in the \mathbb{Z}_2 sense while even ℓ gives \mathbb{Z}_2 even generators such that the two gradings are consistent.

Similar to the bosonic case one can describe the structure of the free Lie superalgebra in terms of a generating function. The generalisation of (2.5) is given by

$$\bigotimes_{\ell \text{ odd}} \left[\bigoplus_{k=0}^{\infty} (-1)^k t^{k\ell} \vee^k R_\ell \right] \otimes \bigotimes_{\ell \text{ even}} \left[\bigoplus_{k=0}^{\infty} (-1)^k t^{k\ell} \wedge^k R_\ell \right] = 1 - tR_1. \quad (2.8)$$

Here, t is again a formal parameter and the \mathbb{Z}_2 nature of the generators introduces some signs. $R_1 = \langle Q_\alpha \rangle$ denotes the generators on level $\ell = 1$ and we denote the generators on level ℓ by R_ℓ . Expanding out the generating identity one obtains for the first few levels

$$R_1 = \langle Q_\alpha \rangle, \quad (2.9a)$$

$$R_2 = \vee^2 R_1, \quad (2.9b)$$

$$R_3 = R_2 \otimes R_1 \ominus \vee^3 R_1, \quad (2.9c)$$

$$R_4 = (R_3 \otimes R_1 \oplus \wedge^2 R_2) \ominus R_2 \otimes \vee^2 R_1 \oplus \vee^4 R_1, \quad (2.9d)$$

$$R_5 = (R_4 \otimes R_1 \oplus R_3 \otimes R_2) \ominus (R_3 \otimes \vee^2 R_1 \oplus \wedge^2 R_2 \otimes R_1) \oplus R_2 \otimes \vee^3 R_1 \ominus \vee^5 R_1, \quad (2.9e)$$

...

Compared to (2.6), the anti-symmetric tensor powers \wedge^k are interchanged for symmetric ones, denoted \vee^k , in a few places. For instance R_2 is given by all *symmetric* products $\{Q_\alpha, Q_\beta\}$ of the Q_α and the Jacobi identity relevant for R_3 is totally symmetric as well.

We can again consider the case when the Q_α transform under some level $\ell = 0$ algebra \mathfrak{g}_0 . This could be the Lorentz algebra $\mathfrak{so}(1, D-1)$ in D dimensions or could also contain possible R-symmetries. We shall consider many examples below. Given a \mathfrak{g}_0 , the generators R_ℓ can be arranged into representations of \mathfrak{g}_0 by treating the tensor products as products of \mathfrak{g}_0 modules.

Just as free Lie algebras, free Lie superalgebras admit many quotients and one can again distinguish the case of finite-dimensional and infinite-dimensional quotients. Examples of

extensions of the standard supersymmetry algebra that have been discussed in the literature are finite-dimensional quotients and we shall illustrate this in examples below. Moreover, there are cases where these quotients can be described conveniently in terms of Borcherds superalgebras and Dynkin diagrams as we discuss in section 4.

3 Instances of free Lie superalgebras

In this section, we apply the algorithm (2.9) and construct the free Lie superalgebra generated by Q_α in various cases where \mathfrak{g}_0 is either the Lorentz algebra in D dimensions, or the direct sum of the Lorentz algebra and an R-symmetry algebra $\mathfrak{su}(\mathcal{N})$.

3.1 $D = 4$ and $\mathcal{N} = 1$ supersymmetry

We begin with the free Lie superalgebra generated by supersymmetry generators Q_α and then compare the results to various extensions of the Poincaré superalgebras that have appeared in the literature.

3.1.1 Free Lie superalgebra

Following the philosophy outlined in the previous section, we now construct the *free Lie superalgebra* $\mathfrak{f}(Q)$ generated by the Q_α in $D = 4$. The spinor index $\alpha = 1, \dots, 4$ here labels the four independent components of a Majorana spinor. Taking $\mathfrak{g}_0 = \mathfrak{so}(1, 3)$, the transformation of $R_1 = \langle Q_\alpha \rangle$ is given by

$$[M_{ab}, Q_\alpha] = \frac{1}{2} Q_\beta (\Gamma_{ab})^\beta{}_\alpha. \tag{3.1}$$

The real gamma matrices here satisfy $\{\Gamma_a, \Gamma_b\} = 2\eta_{ab}$. Gamma matrices with multiple indices are defined by $\Gamma^{ab} = \Gamma^{[a}\Gamma^{b]}$ etc. The gamma matrices that we use are given explicitly in appendix A. We use the signature $\eta = (-+++)$ and below we shall also encounter the symmetric combinations $(C\Gamma^a)^T = C\Gamma^a$ and $(C\Gamma^{ab})^T = C\Gamma^{ab}$ involving the charge conjugation matrix C .

The free Lie superalgebra $\mathfrak{f}(Q)$ is obtained by taking all possible (graded) commutators of the Q_α obeying only graded anti-symmetry and the Jacobi identity. Following the algorithm (2.9) this leads to table 2. In this table we have labelled the representations of the Lorentz algebra $\mathfrak{g}_0 = \mathfrak{so}(1, 3)$ in terms of Dynkin labels of the corresponding complex Lie algebra $A_1 \oplus A_1$, such that a Majorana spinor decomposes as $Q_\alpha \leftrightarrow [1, 0] \oplus [0, 1]$, the familiar decomposition into a left-handed and right-handed spinor in $D = 4$. Keeping this in mind, we can use the LiE software [35] to compute the tensor product decompositions of $A_1 \oplus A_1$. Generally, a complex representation of $A_1 \oplus A_1$ gives a real representation of $\mathfrak{so}(1, 3)$ if the representations of the two A_1 are balanced, i.e., either of the symmetric form $[p_1, p_2] \oplus [p_2, p_1]$ or directly $[p, p]$.

Let us also give the commutation relations for the free Lie superalgebra $\mathfrak{f}(Q)$ at lowest levels, using the names for the generators introduced in table 2. The first non-trivial commutator is that of the supercharges Q_α that we define to be

$$\{Q_\alpha, Q_\beta\} = (C\Gamma^a)_{\alpha\beta} P_a + \frac{1}{2} (C\Gamma^{ab})_{\alpha\beta} P_{ab}, \tag{3.2}$$

R_ℓ	representation	generator
R_1	$[1, 0] \oplus [0, 1]$	Q_α
R_2	$[1, 1]$	P_a
	$[2, 0] \oplus [0, 2]$	$P_{ab} = P_{[ab]}$
R_3	$[1, 0] \oplus [0, 1]$	Σ_α
	$[1, 0] \oplus [0, 1]$	$\tilde{\Sigma}_\alpha$
	$[1, 2] \oplus [2, 1]$	$\Sigma_{a\alpha}$
R_4	$[2, 0] \oplus [0, 2]$	$Z_{ab} = Z_{[ab]}$
	$[0, 0]$	B
	$3 \times [1, 1]$	vectorial
	$[3, 1] \oplus [1, 3] \oplus [1, 1]$	
	$[2, 2] \oplus [0, 2] \oplus [2, 0]$	
	$[2, 0] \oplus [0, 2]$	two-form
\vdots	\vdots	\vdots

Table 2. The free Lie superalgebra $\mathfrak{f}(Q)$ in $D = 4$ space-time dimensions for a single four-component spinor generator Q_α . We have grouped representations together that form a nice representation of $\mathfrak{so}(1, 3)$. All tensor-spinors are gamma traceless, but we retain the tensor traces, meaning that for example the first non-trivial Young tableau in R_4 represents a tensor structure $Y_{ab,c}$ where one can still take the Lorentz trace as discussed in footnote 2; this corresponds to the $[1, 1]$ representation listed there. Some representations occur with a non-trivial multiplicity and we have not given names to generators that do not appear elsewhere in this article.

where the term $P_{ab} = P_{[ab]}$ has already appeared in [36–38]. It does not appear in the standard $\mathcal{N} = 1$ supersymmetry algebra and it is not central as it does transform non-trivially under the Lorentz algebra. It can be interpreted as a source for the supersymmetric membrane in $D = 4$ [36] in a way similar to the complete $\{Q, Q\}$ commutators in eleven dimensions [9]. We shall come back to this interpretation below. We also note that (3.2) implies

$$P_a = \frac{1}{4}(\Gamma_a C^{-1})^{\alpha\beta} \{Q_\alpha, Q_\beta\}, \quad P_{ab} = -\frac{1}{4}(\Gamma_{ab} C^{-1})^{\alpha\beta} \{Q_\alpha, Q_\beta\}. \quad (3.3)$$

In the free Lie superalgebra, one can then form the triple commutators $[\{Q_\alpha, Q_\beta\}, Q_\gamma]$ at level $\ell = 3$ that have to be subjected to the Jacobi identity (2.7). The resulting most general expression can be split up into the commutators of Q_α with P_a and P_{ab} as

$$[Q_\alpha, P_a] = \Sigma_\beta (\Gamma_a)^\beta{}_\alpha + \Sigma_{a\alpha}, \quad (3.4a)$$

$$[Q_\alpha, P_{ab}] = \tilde{\Sigma}_\beta (\Gamma_{ab})^\beta{}_\alpha + \frac{4}{3} \Sigma_{[a\beta} (\Gamma_{b]}^\beta{}_\alpha. \quad (3.4b)$$

The anti-symmetrisation only refers to the vector indices a, b . As mentioned in the caption

of table 2, the tensor spinors we use are gamma traceless which means that

$$\Sigma_{a\alpha}(\Gamma^a)^\alpha{}_\beta = 0, \tag{3.5}$$

and accordingly they span a subspace of dimension $4 \times 4 - 4 = 12$. We also note that the Jacobi identities can imply that a naive counting of generators in commutation relations does not always work. For instance, in (3.4) the left-hand sides have superficially $4 \times 4 + 4 \times 6 = 40$ elements while the right-hand side has $4 + 12 + 4 = 20$ elements and the discrepancy is due to the Jacobi identity, meaning that there are 20 linear combinations of the 40 possible commutators that vanish due to the Jacobi identity. The consistency of the free Lie superalgebra commutation relations with the Jacobi identity requires the non-trivial gamma matrix identities (A.4). We shall see below how a quotient relates to the Maxwell superalgebra introduced in [11].

Continuing to level $\ell = 4$ we have to commute the level $\ell = 3$ generators with the basic Q_α on level $\ell = 1$. We shall not give the full commutations at this level but restrict ourselves to defining a part that is relevant to the comparison below,

$$\{Q_\alpha, \Sigma_\beta\} = \frac{1}{4}(C\Gamma^{ab})_{\alpha\beta}Z_{ab} + (C\Gamma_5)_{\alpha\beta}B + \dots \tag{3.6}$$

such that

$$[P_a, P_b] = Z_{ab} + \dots, \tag{3.7}$$

showing the relation between the free Lie superalgebra and the bosonic free Lie algebra. In the two equations above the dots denote additional generators that are present and crucial in the free Lie superalgebra. Their tensor type has been listed in table 2. For the comparison in the next section we do not require the precise form of these terms.

3.1.2 Comparison to Maxwell superalgebra in the literature

Let us now leave the free Lie superalgebra $\mathfrak{f}(Q)$ for the moment and go back to the Poincaré superalgebra in $D = 4$ (with $\mathcal{N} = 1$ supersymmetry):

$$[M_{ab}, P_c] = 2\eta_{c[b}P_{a]}, \tag{3.8a}$$

$$[M_{ab}, Q_\alpha] = \frac{1}{2}Q_\beta(\Gamma_{ab})^\beta{}_\alpha. \tag{3.8b}$$

$$\{Q_\alpha, Q_\beta\} = (C\Gamma^a)_{\alpha\beta}P_a, \tag{3.8c}$$

$$[P_a, Q_\alpha] = 0, \tag{3.8d}$$

$$[P_a, P_b] = 0. \tag{3.8e}$$

The Poincaré superalgebra has non-trivial cohomology and it can be extended as was studied in [11].⁴ It was shown there that one can in particular deform the commutators (3.8d) and (3.8e) through the introduction of a new (non-central) Majorana spinor

⁴We note that compared to [11] we have slightly changed conventions by removing some factors of i and rescaling some generators.

generator Σ_α by letting⁵

$$[Q_\alpha, P_a] = \Sigma_\beta (\Gamma_a)^\beta{}_\alpha. \quad (3.9)$$

In order to obtain a minimal $\mathcal{N} = 1$ supersymmetric extension of the Maxwell algebra that includes the generators $\{M_{ab}, P_a, Z_{ab}\}$ where according to (2.2)

$$[P_a, P_b] = Z_{ab} \quad (3.10)$$

one also has to impose by the Jacobi identity that

$$\{Q_\alpha, \Sigma_\beta\} = \frac{1}{4} (C\Gamma^{ab})_{\alpha\beta} Z_{ab} + (C\Gamma_5)_{\alpha\beta} B. \quad (3.11)$$

In the relation above we have also included a generator B that was introduced in [11] to pair with the bosonic Lorentz scalar chirality operator B_5 satisfying $[B_5, Q_\alpha] = Q_\beta (\Gamma_5)^\beta{}_\alpha$. From the commutation relations above one then deduces $[B_5, P_a] = 0$ and $[B_5, \Sigma_\alpha] = -\Sigma_\beta (\Gamma_5)^\beta{}_\alpha$. The total algebra considered in [11] then consists of

$$\underbrace{M_{ab}, B_5}_{\mathfrak{g}_0}, \underbrace{Q_\alpha}_{\ell=1}, \underbrace{P_a}_{\ell=2}, \underbrace{\Sigma_\alpha}_{\ell=3}, \underbrace{Z_{ab}, B}_{\ell=4} \quad (3.12)$$

We thus see that the generators of [11] form a subset of \mathfrak{g}_0 together with the free Lie superalgebra $\mathfrak{f}(Q)$ defined above and is consistent with the commutation relations. It is in fact a quotient of the free Lie superalgebra and the quotient can, moreover, be described in terms of a Borchers superalgebra as we shall explain in more detail in section 4. We also note that the algebra (3.12) admits an invariant Casimir of the form

$$C_2 = \frac{1}{2} P_a P^a + Q_\alpha C^{\alpha\beta} \Sigma_\beta - \frac{1}{2} M_{ab} Z^{ab} + B_5 B. \quad (3.13)$$

The quadratic Casimir (3.13) pairs generators on levels ℓ and $4 - \ell$; extending this beyond the generators in (3.12) would require generators on negative levels that pair with generators on level $\ell > 4$. Such a structure is provided by a tensor hierarchy algebra [20–23], yet another algebra that can be defined from the Poincaré superalgebra. In such an extension, the Casimir would also be made more manifestly graded symmetric by letting $B_5 B \rightarrow \frac{1}{2} (B_5 B + B B_5)$, $Q_\alpha C^{\alpha\beta} \Sigma_\beta \rightarrow \frac{1}{2} (Q_\alpha C^{\alpha\beta} \Sigma_\beta + \Sigma_\alpha C^{\alpha\beta} Q_\beta)$, etc.; in the algebra (3.12) this is not necessary as these generators commute up to central terms.

3.2 $D = 4$ and extended supersymmetry

The construction of a free Lie superalgebra $\mathfrak{f}(Q)$ described in section 2.2 allows for the Q to be odd generators transforming under some level $\ell = 0$ algebra \mathfrak{g}_0 . In the previous section, we considered the $D = 4$ Lorentz algebra $\mathfrak{g}_0 = \mathfrak{so}(1, 3)$ (possibly extended by a chirality operator). In this section, we shall consider the case of extended supersymmetry where there is a non-trivial R-symmetry acting on the supersymmetry generators. That is, we consider the case $\mathfrak{g}_0 = \mathfrak{so}(1, 3) \oplus \mathfrak{su}(\mathcal{N})$ for \mathcal{N} -extended supersymmetry. The corresponding basic generators will be denoted by Q_α^I where $\alpha = 1, \dots, 4$ is the Lorentz spinor index while $I = 1, \dots, \mathcal{N}$ is the R-symmetry index.

⁵A similar extension of the supersymmetry algebra in $D = 10$ dimensions had been introduced previously in order to write the Wess-Zumino term of the Green-Schwarz string as an invariant term [13]. The algebra of [13] has also been extended to study supersymmetric p -brane models and their κ -symmetry [39, 40].

R_ℓ	representation	generator
R_1	$[1, 0; 1] \oplus [0, 1; 1]$	Q_α^I
R_2	$[1, 1; 0]$	P_a
	$2 \times [0, 0; 0]$	P, P_5
	$[1, 1; 2]$	$P_a^{IJ} = P_a^{(IJ)}$
	$[2, 0; 2] \oplus [0, 2; 2]$	$P_{ab}^{IJ} = P_{ab}^{(IJ)} = P_{[ab]}^{IJ}$
\vdots	\vdots	\vdots

Table 3. The free Lie superalgebra $\mathfrak{f}(Q)$ in $D = 4$ space-time dimensions for extended $\mathcal{N} = 2$ supersymmetry. The $\mathfrak{su}(2)$ R-symmetry representation is given by the last Dynkin label while the first two specify the $A_1 \oplus A_1$ representation. The fundamental R-symmetry index is $I = 1, 2$.

3.2.1 $D = 4$ and $\mathcal{N} = 2$ supersymmetry

Complexifying the algebra as before we are therefore considering $A_1 \oplus A_1 \oplus A_1$ where each A_1 denotes a complexified $\mathfrak{su}(2)$. Then the level $\ell = 1$ generators Q_α^I have representation labels

$$R_1 = [1, 0; 1] \oplus [0, 1; 1], \tag{3.14}$$

so that the last label is the R-symmetry label, separated with a semicolon. The case $\mathcal{N} = 2$ is special as the R-symmetry representation $[1]$ can be treated as a real representation using ϵ^{IJ} as a complex structure. This is no longer the case for $\mathcal{N} > 2$ and we treat this case on its own below. Applying the algorithm (2.9) to the $\mathcal{N} = 2$ generator (3.14) leads to table 3, where we do not give the result for R_ℓ with $\ell > 2$ due to a proliferation of terms.

The commutation relation in the free Lie superalgebra representing the first two levels is

$$\begin{aligned} \{Q_\alpha^I, Q_\beta^J\} &= \epsilon^{IJ} (C\Gamma_5 \Gamma^a)_{\alpha\beta} P_a + \epsilon^{IJ} C_{\alpha\beta} P + \epsilon^{IJ} (C\Gamma_5)_{\alpha\beta} P_5 \\ &+ (C\Gamma^a)_{\alpha\beta} P_a^{IJ} + \frac{1}{2} (C\Gamma_5 \Gamma^{ab})_{\alpha\beta} P_{ab}^{IJ}. \end{aligned} \tag{3.15}$$

The first line corresponds to the central extension of the $\mathcal{N} = 2$ Poincaré superalgebra by an electric and magnetic charge. The second line introduces additional string and membrane charges [9, 37].

The calculation can be carried out to higher levels and will contain the hallmark relation $[P_a, P_b] = Z_{ab}$ of Maxwell algebras. An \mathcal{N} -extended supersymmetric version of the Maxwell algebra in $D = 4$ was introduced in [12] from a contraction of a superconformal algebra. Their algebra has an R-symmetry of \mathfrak{so} -type rather than the $\mathfrak{su}(\mathcal{N})$ that we have assumed here and therefore the resulting commutation relations are quite different, allowing in particular a raising and lowering of R-symmetry indices.

3.2.2 $D = 4$ and $\mathcal{N} > 2$ supersymmetry

One can similarly consider \mathcal{N} -extended supersymmetry in $D = 4$ dimension for $\mathcal{N} > 2$ by letting the generating elements of the free Lie superalgebra be Q_α^I with I a fundamental

index of $\mathfrak{su}(\mathcal{N})$. In order to write an anti-commutator of the elementary Q_α^I that yields standard translation generators P_a it is necessary to use 2-component Weyl spinors that we denote by Q_A^I and $\bar{Q}_{\dot{A} I}$ (with $A = 1, 2$ and $\dot{A} = \dot{1}, \dot{2}$). The important point here is that the two Weyl spinors transform in conjugate (fundamental) representations of $\mathfrak{su}(\mathcal{N})$. For $\mathcal{N} = 2$ we could use the invariant ϵ^{IJ} to relate the two representations, but for $\mathcal{N} > 2$ there is no corresponding invariant tensor to achieve this. The complexification of $\mathfrak{g}_0 = \mathfrak{so}(1, 3) \oplus \mathfrak{su}(\mathcal{N})$ is of type $A_1 \oplus A_1 \oplus A_{\mathcal{N}-1}$ and the corresponding complex representation is

$$R_1 = [1, 0; 1, 0, \dots, 0] \oplus [0, 1; 0, \dots, 0, 1], \tag{3.16}$$

illustrating that the two Weyl spinors transform in conjugate representations under $A_{\mathcal{N}-1}$. This space has complex dimension $\dim_{\mathbb{C}}(R_1) = 4\mathcal{N}$ which is twice the number of real supertranslations of \mathcal{N} -extended supersymmetry. A real slice through the complex representation is picked by a reality condition of the form

$$\bar{Q}_{\dot{A} I} = (Q_A^I)^*. \tag{3.17}$$

We note that the conjugation affects the R-symmetry index and spinor index at the same and the R-symmetry transformations in a real basis will therefore also affect the spinor index. This happens for instance for $\mathcal{N} = 8$ supergravity [41, 42]. Writing the $\mathfrak{su}(\mathcal{N})$ in terms of its real $\mathfrak{so}(\mathcal{N})$ subalgebra one can make the transition between the complex Weyl and a real Majorana basis manifest, but at the sake of giving up manifest $\mathfrak{su}(\mathcal{N})$ invariance. We shall not carry out this rewriting here but instead work with the (complex) Weyl spinors.

In terms of the Pauli matrices σ_{AA}^a one then obtains the (anti-)commutation relations

$$\{Q_A^I, \bar{Q}_{\dot{A} J}\} = 2\delta_J^I(\sigma^a)_{A\dot{A}}P_a + (\sigma^a)_{A\dot{A}}\tilde{P}_a^{IJ} \tag{3.18a}$$

$$\{Q_A^I, Q_B^J\} = P_{AB}^{IJ} + \epsilon_{AB}P^{IJ} \tag{3.18b}$$

$$\{\bar{Q}_{\dot{A} I}, \bar{Q}_{\dot{B} J}\} = \bar{P}_{\dot{A}\dot{B} IJ} + \epsilon_{\dot{A}\dot{B}}\bar{P}_{IJ}, \tag{3.18c}$$

where

$$\tilde{P}_a^{IJ} = 0, \quad P_{AB}^{IJ} = P_{(AB)}^{IJ} = P_{AB}^{(IJ)}, \quad P^{IJ} = P^{[IJ]}, \tag{3.19}$$

with similar relations for the conjugate generators. The generators P_{AB}^{IJ} and $\bar{P}_{\dot{A}\dot{B} IJ}$ with symmetric pairs of spinor indices combine into an antisymmetric two-form under the Lorentz group $\mathfrak{so}(1, 3)$ but this obscures their $\mathfrak{su}(\mathcal{N})$ properties in a way similar to (3.17).

3.3 $D = 11$ and $\mathcal{N} = 1$ supersymmetry

The last example we discuss in some detail is given by minimal supersymmetry in $D = 11$ space-time dimensions as this is the case relevant to M-theory.

3.3.1 Free Lie superalgebra

In this case the Q_α are real 32-component Majorana spinors that are irreducible under the Lorentz algebra $\mathfrak{so}(1, 10)$ and form the representation $R_1 = [0, 0, 0, 0, 1]$ of the complexified

R_ℓ	representation	generator
R_1	$[0, 0, 0, 0, 1]$	Q_α
R_2	$[1, 0, 0, 0, 0]$	P_a
	$[0, 1, 0, 0, 0]$	$P_{ab} = P_{[ab]}$
	$[0, 0, 0, 0, 2]$	$P_{a_1 \dots a_5} = P_{[a_1 \dots a_5]}$
R_3	$2 \times [0, 0, 0, 0, 1]$	spinors
	$2 \times [1, 0, 0, 0, 1]$	vector-spinors
	$[0, 1, 0, 0, 1]$	two-form spinor
	$[0, 0, 1, 0, 1]$	three-form spinor
	$[0, 0, 0, 1, 1]$	four-form spinor
R_4	$6 \times [0, 0, 0, 0, 2]$...
	$4 \times [0, 0, 0, 1, 0]$	
	$[0, 0, 0, 1, 2]$	
	$4 \times [0, 0, 1, 0, 0]$	
	$2 \times [0, 0, 1, 0, 2]$	
	$[0, 0, 1, 1, 0]$	
	$5 \times [0, 1, 0, 0, 0]$	
	$2 \times [0, 1, 0, 0, 2]$	
	$3 \times [0, 1, 0, 1, 0]$	
	$2 \times [0, 1, 1, 0, 0]$	
	$3 \times [1, 0, 0, 0, 0]$	
	$3 \times [1, 0, 0, 0, 2]$	
	$4 \times [1, 0, 0, 1, 0]$	
	$4 \times [1, 0, 1, 0, 0]$	
	$2 \times [1, 1, 0, 0, 0]$	
\vdots	\vdots	\vdots

Table 4. The free Lie superalgebra $\mathfrak{f}(Q)$ in $D = 11$ space-time dimensions for a single 32-component spinor generator Q_α . All $\mathfrak{so}(1,10)$ representations are labelled by their B_5 Dynkin labels in Bourbaki numbering. Tensor-spinors are gamma traceless but we retain tensor traces. Some representations occur with multiplicity as shown.

Lie algebra B_5 . The last label refers to the spinor node of the B_5 Dynkin diagram. Carrying out the algorithm (2.9) leads to table 4.

The commutation relations leading to level $\ell = 2$ in the free Lie superalgebra $\mathfrak{f}(Q)$ are

$$\{Q_\alpha, Q_\beta\} = (C\Gamma^a)_{\alpha\beta} P_a + \frac{1}{2}(C\Gamma^{ab})_{\alpha\beta} P_{ab} + \frac{1}{5!}(C\Gamma^{a_1 \dots a_5})_{\alpha\beta} P_{a_1 \dots a_5}. \quad (3.20)$$

The right-hand side contains the most general combination that can be written for two supersymmetry transformations in $D = 11$. This extension of the Poincaré superalgebra was first considered in [8], and is nowadays called the M-algebra (but should not be confused with the generalisation thereof constructed in [10]). In [9, 43, 44] it has been argued that the (non-central under Lorentz) terms P_{ab} and $P_{a_1\dots a_5}$ beyond the usual translation generator P_a are the ones that correspond to the M2-brane and M5-brane of eleven-dimensional supergravity. The even more general algebra constructed in [10] contained more brane sources and even fermionic ones that were obtained by treating the P_a , P_{ab} and $P_{a_1\dots a_5}$ as forms in superspace. The immediate interpretation of all these charges was not clear but some were used for studying the Wess-Zumino term of the M5-brane [10].

One possible quotient of the free Lie superalgebra $\mathfrak{f}(Q)$ is the standard Poincaré superalgebra in $D = 11$ (without P_{ab} and $P_{a_1\dots a_5}$) and another one is the truncation to $\ell = 2$ which then reproduces the M-algebra of [9]. Including also generators from higher levels one may construct quotients that agree with the algebra considered in [10].

We also note that very recently a supersymmetric extension of the $D = 11$ Maxwell algebra was proposed in [45, 46]. This algebra is a quotient of the free Lie superalgebra by retaining the following generators besides the Poincaré superalgebra: on $\ell = 2$ additionally P_{ab} but not $P_{a_1\dots a_5}$; on $\ell = 3$ only a single spinor Σ_α (similar to (3.9)) and on level $\ell = 4$ only a single two-form Z_{ab} . Other extensions of the $D = 11$ Poincaré superalgebra were studied in the context of free differential algebras in [47, 48].

3.3.2 Relation to E_{11} and ℓ_1 representation?

It is tempting to speculate on a relation between the free Lie superalgebra and exotic supersymmetric objects as they appear in supergravity and M-theory and are captured by approaches based on the Kac-Moody algebra E_{11} [15, 49]. Exotic branes in the sense of [50, 51] are extended objects of co-dimension at most two and they tend to be very non-perturbative when analysed in string theory, meaning their mass scales with the inverse string coupling to a power that is larger than 1. These exotic branes have been discovered originally using U-duality in low dimensions (see for instance [52, 53]). From the point of view of an electric space-time coupling they typically couple to *mixed symmetry* space-time potentials, meaning potentials that are not necessarily p -forms but represented by Young tableaux of more complicated type than a single column. Using this language all such supersymmetric branes have been classified and their relation to non-geometric backgrounds has been studied [54–57].

Even though exotic branes can typically be expressed as supersymmetric solutions of usual supergravity, meaning that they preserve some of the usual supersymmetries generated by the Q_α at $\ell = 1$ in $\mathfrak{f}(Q)$, their (electric) coupling to mixed symmetry potentials suggests to also investigate whether there can be any relation to higher levels $\ell > 1$ in the free Lie superalgebra. The motivation from this investigation comes from the fact that the adjoint representation of E_{11} captures all the relevant mixed symmetry potentials [55] and that the ℓ_1 representation of E_{11} seems to capture all the corresponding ‘electric’ charges [49, 58]. With electric charges we mean that while the M2-brane of $D = 11$ supergravity couples naturally to the 3-form potential, the M5-brane couples more naturally

‘electrically’ to its dual 6-form rather than ‘magnetically’ to the original 3-form. The electric coupling is always such that the corresponding Wess-Zumino term in the world-volume action is simply an integral over (the pull-back of) the corresponding potential. Roughly, for a supersymmetric p -brane the closed p -form $dX^{a_1} \wedge \dots \wedge dX^{a_p}$ on the world-volume Σ contributes to the supersymmetry algebra if the brane, embedded via the maps $X^a(\xi)$, wraps a topologically non-trivial cycle. This is due to the quasi-invariance of the Wess-Zumino coupling $\int_{\Sigma} C_{(p+1)}$ to a $(p+1)$ -form gauge field $C_{(p+1)}$ under supersymmetry [43]. The contribution to the supersymmetry algebra is of the form

$$\{Q_{\alpha}, Q_{\beta}\} = \frac{1}{p!} (C\Gamma_{a_1 \dots a_p})_{\alpha\beta} Z^{a_1 \dots a_p}, \quad \text{where } Z^{a_1 \dots a_p} \sim Q_p \int d^p \xi dX^{a_1} \wedge \dots \wedge dX^{a_p}, \quad (3.21)$$

with the integral over the non-trivial (spatial) cycle that the brane wraps and Q_p the charge of the brane. With this logic the 2-form P_{ab} in (3.20) is related to the M2-brane while the 5-form $P_{a_1 \dots a_5}$ is related to the M5-brane [9, 44]. This can be also understood through the gauge-field that the brane couples to as follows. The brane requires a $(p+2)$ -form flux in space-time $F_{(p+2)} = dC_{(p+1)} + \dots$ and in D space-time dimensions the transverse space has a sphere S^{D-p-2} at infinity over which the dual of the flux $F_{(p+2)}$ can be integrated to give the charge Q_p of the extended object. From this we see that the supersymmetry algebra for p -brane coupling to a form $C_{a_1 \dots a_{p+1}}$ should contain a (non-central) term $Z_{a_1 \dots a_p}$, i.e., a form with one index less.

As exotic branes couple to mixed symmetry potentials one might therefore by extension wonder whether there are mixed-symmetry ‘charges’ sitting somewhere in an extended supersymmetry algebra to which they couple. As there is no room in the standard $D = 11$ superalgebra (3.20) we look for them in the free Lie superalgebra $\mathfrak{f}(Q)$. In order to identify them we use the following tentative connection to E_{11} .

From the point of view of E_{11} , the generators P_a , P_{ab} and $P_{a_1 \dots a_5}$ appear naturally in the so-called ℓ_1 representation when it is decomposed under $\mathfrak{gl}(11) \subset E_{11}$ [49, 58]. The relation between space-time potentials in the adjoint of E_{11} and the ℓ_1 representation has also been discussed in these references. As E_{11} is conjectured to contain all the mixed symmetry space-time potentials that (supersymmetric) branes can couple to, it is an important observation that the ℓ_1 representation of E_{11} contains all the mixed symmetry ‘charges’ that can be obtained by removing one index from any of the E_{11} gauge potentials [49, 58, 59].

In order to compare the $\mathfrak{so}(1, 10)$ representations predicted by E_{11} and the content of the free Lie superalgebra $\mathfrak{f}(Q)$, we reproduce the lowest levels of the ℓ_1 representation of E_{11} in terms of $\mathfrak{gl}(11)$ representations in table 5. The table also lists their decomposition into $\mathfrak{so}(1, 10)$ representations. This should be compared to the *even* levels of the free Lie superalgebras listed in table 4.

The comparison has been carried out at the level of representations up to R_4 in the last column of table 5. As can be seen from that column, most of the mixed symmetry representations contained in the ℓ_1 representation arise already on the first two bosonic levels R_2 and R_4 of the free Lie superalgebra; it is possible that the question marks will be contained in the higher levels. As the charges in the ℓ_1 representation cover *all* branes, the analysis above includes standard and exotic branes.

Level	$\mathfrak{gl}(11)$ representation	$\mathfrak{so}(1, 10)$ representation	Occurrence in $\mathfrak{f}(Q)$
$\frac{3}{2}$	[1, 0, 0, 0, 0, 0, 0, 0, 0, 0]	[1, 0, 0, 0, 0]	R_2
$\frac{5}{2}$	[0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0]	[0, 1, 0, 0, 0]	R_2
$\frac{7}{2}$	[0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0]	[0, 0, 0, 0, 2]	R_2
$\frac{9}{2}$	[0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 1]	[1, 0, 0, 1, 0]	R_4
	[0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0]	[0, 0, 0, 0, 2]	R_4
$\frac{11}{2}$	[0, 0, 1, 0, 0, 0, 0, 1, 0, 0]	[0, 0, 2, 0, 0]	?
		[0, 1, 0, 1, 0]	R_4
		[1, 0, 0, 0, 2]	R_4
		[0, 0, 0, 0, 2]	R_4
	[0, 1, 0, 0, 0, 0, 0, 0, 0, 2]	[0, 1, 0, 0, 0]	R_4
		[1, 0, 1, 0, 0]	R_4
		[2, 1, 0, 0, 0]	?
	[0, 1, 0, 0, 0, 0, 0, 0, 1, 0]	[0, 0, 0, 1, 0]	R_4
		[0, 2, 0, 0, 0]	R_4
		[1, 0, 1, 0, 0]	R_4
$2 \times [1, 0, 0, 0, 0, 0, 0, 0, 0, 1]$	$2 \times [0, 1, 0, 0, 0]$	R_4	
	$2 \times [2, 0, 0, 0, 0]$?	
[0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0]	[0, 0, 0, 0, 0]	?	
\vdots	\vdots	\vdots	\vdots

Table 5. The first few levels of the ℓ_1 lowest weight representation of E_{11} . Level here refers to the natural grading under $\mathfrak{gl}(11) \subset E_{11}$. The last column lists a possible match with the even levels of the free Lie algebra in table 4.

We wish to stress that this comparison is speculative and we have only analysed the representations. The algebraic structure is quite different, however. In the E_{11} proposal of [49], all generators in ℓ_1 commute: they form a generalised abelian translation algebra of a generalised space-time in a way similar to the generalised diffeomorphisms of exceptional geometry [60–63]. One of the hallmarks of the Maxwell algebra and free Lie algebra approach pursued here is that the translations P_a no longer commute, see (2.2). This can be achieved in an E_{11} -covariant fashion by allowing non-trivial commutation relations for the elements of the ℓ_1 representation. One possibility would be an embedding in E_{12} along the lines of [58], see also [64]. This embedding embeds the semi-direct sum $E_{11} \oplus \ell_1$ as levels 0 and 1 of a graded decomposition of E_{12} under its obvious E_{11} subalgebra. We shall make more comments on such level decompositions in the next section. Another possibility is to consider the elements in the ℓ_1 representation as odd such that they satisfy non-trivial

anti-commutation relations rather than commutation relations. This structure can be embedded in the Borchers superalgebra denoted \mathcal{B}_{11} in [65], or the tensor hierarchy algebra considered in [23]. We refrain from speculating further in this direction, but we will in the next section apply the construction of a Borchers superalgebra to the case considered in section 3 where the basic objects are the supersymmetry generators Q_α .

4 Free Lie superalgebras and Borchers superalgebras

We have seen that a free Lie (super)algebra decomposes into a direct sum of subspaces at positive levels, and that we can add a Lie algebra \mathfrak{g}_0 at level zero, such that the positive level subspaces form representations of it. In order to obtain a simple Lie (super)algebra we need to continue to negative levels since the subspace spanned by all elements at levels $\ell > k$ for a given $k \geq 1$ otherwise constitutes an ideal, as described in the end of section 2.1. Any finite-dimensional complex semisimple Lie algebra has this structure, with the Cartan subalgebra at level zero and root vectors corresponding to positive and negative roots at positive and negative levels. The construction of the algebra from its Cartan matrix or Dynkin diagram can be generalised in different ways, leading to infinite-dimensional Kac-Moody algebras, the Borchers superalgebras that we will describe next, and most generally, to so-called contragredient Lie (super)algebras [66, 67]. As we will show, when \mathfrak{g}_0 is semisimple one can extend its Cartan matrix or Dynkin diagram and apply the generalised construction in order to obtain a corresponding extension of \mathfrak{g}_0 with two subalgebras at positive and negative levels, respectively, which are quotients of two (isomorphic) free Lie (super)algebras, and additional Cartan generators at level zero.

A Borchers superalgebra of rank r is defined from a symmetric Cartan matrix A_{ij} ($i, j = 1, 2, \dots, r$) with non-positive off-diagonal entries and a subset S of the index set $\{1, 2, \dots, r\}$ labelling the rows and columns. Here we will only consider the case when the Cartan matrix is non-degenerate, integer-valued and satisfies

$$i \notin S \Leftrightarrow A_{ii} = 2, \quad i \in S \Leftrightarrow A_{ii} = 0. \tag{4.1}$$

To the Cartan matrix we associate $3r$ generators e_i, h_i, f_i among which e_i, f_i are odd if and only if $i \in S$, and all h_i are even. The Borchers superalgebra is now defined as the Lie superalgebra generated by e_i, h_i, f_i modulo the relations

$$[h_i, e_j] = A_{ij}e_j, \quad [h_i, f_j] = -A_{ij}f_j, \quad \llbracket e_i, f_j \rrbracket = \delta_{ij}h_i, \quad [h_i, h_j] = 0, \tag{4.2a}$$

$$i \neq j \text{ and } i \notin S \Rightarrow (\text{ad } e_i)^{1-A_{ij}}e_j = 0, \quad (\text{ad } f_i)^{1-A_{ij}}f_j = 0, \tag{4.2b}$$

$$A_{ii} = 0 \Rightarrow \{e_i, e_i\} = 0, \quad \{f_i, f_i\} = 0. \tag{4.2c}$$

We have used the notation $\llbracket \cdot, \cdot \rrbracket$ to denote the graded commutator since the e_i and f_i can be odd or even. The off-diagonal entries of the Cartan matrix A_{ij} determine the Serre relations of (4.2b). The adjoint action there is also by the graded commutator.

The Dynkin diagram that we associate to the Cartan matrix consists of r nodes, where node i ($i = 1, 2, \dots, r$) is white (depicted with \circ) if $i \notin S \Leftrightarrow A_{ii} = 2$ and gray (depicted with \otimes) if $i \in S \Leftrightarrow A_{ii} = 0$, and where two nodes i and j are connected with $-A_{ij}$ lines.




Figure 1. Dynkin diagram of a Borchers Lie superalgebra that is related to a quotient of the free Lie superalgebra of $D = 4$ supersymmetry.

(We note that the relation between Cartan matrices and Dynkin diagrams is not one-to-one for superalgebras as their are odd Weyl reflections that can be applied [68].)

As noted above, Borchers superalgebras have a triangular decomposition: they consist of the Cartan subalgebra spanned by the h_i , a lower-triangular part generated by the f_i through multi-commutation and an upper-triangular part generated by the e_i through multi-commutation. Important for us is the observation that the upper-triangular part is a free Lie superalgebra modulo the Serre relations (4.2b) and (4.2c). In other words, the Serre relations define an ideal in a free Lie superalgebra that has to be quotiented out. We can refine the observation by allowing each generator e_i to have an arbitrary non-negative level v_i , and the corresponding f_i to have level $-v_i$. If $v_i = 1$ for all i , then we get the grading described above, with only the Cartan subalgebra at level zero. But we can also choose to have $v_i = 1$ only for some i , and $v_i = 0$ for the others. By setting $v_i = 1$ if $i \in S$ and $v_i = 0$ if $i \notin S$ we get a \mathbb{Z} -grading consistent with the \mathbb{Z}_2 -grading: odd elements appear at odd levels, and even elements at even levels. The level $\ell = 0$ subalgebra \mathfrak{g}_0 is then a semisimple Kac-Moody algebra with a Dynkin diagram given by removing the gray nodes from the Dynkin diagram of the Borchers superalgebra, centrally extended with the Cartan generators h_i of the removed nodes. In this level decomposition, the Serre relations (4.2b) determine the level $\ell = 0$ subalgebra \mathfrak{g}_0 and its representations at levels $\ell = \pm 1$ (which will be dual to each other), whereas the additional Serre relations (4.2c) define the ideal that is factored out from the free Lie superalgebras at positive and negative levels generated by the subspaces at level $\ell = 1$ and $\ell = -1$, respectively.

4.1 $D = 4$ and $\mathcal{N} = 1$ supersymmetry

We will now try to reconstruct an infinite-dimensional extension of the superalgebra considered in (3.12) as the subalgebra at non-negative levels of a Borchers superalgebra. In agreement with the discussion above we should have at level $\ell = 0$ the complexification $A_1 \oplus A_1$ of the Lorentz algebra $\mathfrak{so}(1, 3)$ in $D = 4$ and at level $\ell = 1$ the spinors $[0, 1] \oplus [1, 0]$. One way to arrange this is by taking the Cartan matrix

$$A = \begin{pmatrix} 2 & -1 & 0 & 0 \\ -1 & 0 & -2 & 0 \\ 0 & -2 & 0 & -1 \\ 0 & 0 & -1 & 2 \end{pmatrix}. \quad (4.3)$$

This can also be depicted in terms of a Dynkin diagram as in figure 1.

The two A_1 forming the Lorentz algebra in $D = 4$ correspond to nodes 1 and 4 of the Dynkin diagram and the gray nodes are fermionic and represent the two fundamental representations of the two A_1 as required by the fundamental spinor $Q_\alpha \leftrightarrow [1, 0] \oplus [0, 1]$.

Thus the subset S of the index set $\{1, 2, 3, 4\}$ is $S = \{2, 3\}$ here. The single lines connecting the gray nodes to nodes 1 and 4 imply that the corresponding Chevalley generators belong to a two-dimensional fundamental representation of the A_1 concerned.

Let us first see how to get the Lorentz algebra by removing the gray nodes. If we set

$$M_{02} = \frac{i}{2}(-e_1 + f_1 + e_4 - f_4), \quad M_{01} = \frac{1}{2}(-e_1 - f_1 - e_4 - f_4), \quad (4.4a)$$

$$M_{23} = \frac{i}{2}(e_1 + f_1 - e_4 - f_4), \quad M_{31} = \frac{1}{2}(-e_1 + f_1 - e_4 + f_4), \quad (4.4b)$$

$$M_{21} = \frac{i}{2}(h_1 - h_4), \quad M_{03} = \frac{1}{2}(h_1 + h_4), \quad (4.4c)$$

the commutation relations (2.1a) indeed follow from (4.2) in the case where the Cartan matrix A_{ij} is given by (4.3), or equivalently, by the Dynkin diagram in figure 1. Note that the imaginary unit i appears in (4.4) although we consider the real Lie algebra $\mathfrak{so}(1, 3)$ (no i appears in the commutation relations (2.1a)). This shows that $\mathfrak{so}(1, 3)$ is a non-split real form of $A_1 \oplus A_1$. If we instead just took the real span of $e_1, f_1, h_1, e_4, f_4, h_4$, then we would get the split real form $\mathfrak{sl}(2) \oplus \mathfrak{sl}(2)$ of $A_1 \oplus A_1$. Note also that $e_2, f_2, h_2, e_3, f_3, h_3$ do not appear in (4.4) since we have removed the gray nodes. However, the Cartan generators h_2 and h_3 do belong to level zero as well and by taking appropriate linear combinations of them, one can get the dilatation operator d and the chirality operator B_5 that also appears in the algebra (3.12).

We now put back the gray nodes and go from level $\ell = 0$ to $\ell = 1$. Looking for example at the e_2 Chevalley generator, the Serre relation implies

$$e_2 \neq 0, \quad [e_1, e_2] \neq 0, \quad \text{but} \quad [e_1, [e_1, e_2]] = 0, \quad (4.5)$$

so that there are two non-zero elements at level $\ell = 1$ associated with e_2 . This is the $[1, 0]$ representation of $A_1 \oplus A_1$. One can verify the Dynkin labels $[1, 0]$ by acting with the Cartan generators h_1 and h_4 on the highest weight vector $[e_1, e_2]$:

$$[h_1, [e_1, e_2]] = 1, \quad [h_4, [e_1, e_2]] = 0. \quad (4.6)$$

A similar reasoning for nodes 3 and 4 yields the $[0, 1]$ representation. The level ℓ here is given by the sum of the number of times e_2 or e_3 appear in a multi-commutator. Thus one of the two 2-dimensional Weyl spinors at level $\ell = 1$ is spanned (over the complex numbers) by e_2 and $[e_1, e_2]$ and the other one by e_4 and $[e_3, e_4]$. In order to get a Majorana spinor Q_α ($\alpha = 1, 2, 3, 4$) from them, we have to take the complex linear combinations

$$Q_1 = [e_1, e_2] - i[e_3, e_4], \quad (4.7a)$$

$$Q_2 = i[e_1, e_2] - [e_3, e_4], \quad (4.7b)$$

$$Q_3 = e_2 + ie_3, \quad (4.7c)$$

$$Q_4 = -ie_2 - e_3. \quad (4.7d)$$

Indeed, now the commutation relations (3.1), which state that Q_α transforms as a Majorana spinor, follow from (4.2), and the explicit form of the real gamma matrices given in A.

We proceed to level $\ell = 2$ and set

$$P_0 = i[e_1, \{e_2, [e_3, e_4]\}] - i\{e_2, e_3\}, \quad (4.8a)$$

$$P_1 = i[e_1, \{e_2, e_3\}] - i\{e_2, [e_3, e_4]\}, \quad (4.8b)$$

$$P_2 = -[e_1, \{e_2, e_3\}] - \{e_2, [e_3, e_4]\}, \quad (4.8c)$$

$$P_3 = i[e_1, \{e_2, [e_3, e_4]\}] + i\{e_2, e_3\}. \quad (4.8d)$$

Note that in each multi-commutator, e_2 and e_3 appear once each. A multi-commutator with two e_2 and no e_3 (or two e_3 and no e_2) would also belong to level $\ell = 2$, but any such element is zero because of the Serre relations

$$\{e_2, e_2\} = \{e_3, e_3\} = 0. \quad (4.9)$$

This means that P_{ab} , that appeared in the anti-commutation relation (3.2) of the free Lie superalgebra is not present in the Borchers superalgebra we consider here, since $\{e_2, e_2\}$ and $\{e_3, e_3\}$ are lowest weight vectors of the representation $[2, 0]$ and $[0, 2]$, respectively, which together form P_{ab} . Rather we get (3.8c) from (4.2). Also the relation (3.8a), which states that P_a transforms as vector under the Lorentz algebra, follow from (4.2).⁶

In drawing diagram 1 we have chosen a double line between nodes 2 and 3, corresponding to $A_{23} = A_{32} = -2$. For understanding the Serre relations drawing a single or multiple lines between the nodes does not matter, as any negative choice of A_{23} implies the same ideal. This is due to the Jacobi identity

$$[e_2, \{e_2, e_3\}] = [\{e_2, e_2\}, e_3] - [e_2, \{e_2, e_3\}]. \quad (4.10)$$

The first term on the right-hand side vanishes due to (4.2c) and the sign in front of the second term follows from the oddness of e_2 . This Jacobi identity therefore implies

$$[e_2, \{e_2, e_3\}] = 0 \quad (4.11)$$

independent of the value of A_{23} . The value of A_{23} does, however, lead to inequivalent Borchers superalgebras when one considers the action of the lower-triangular Chevalley generators f_i as well. As long as one is only interested in the upper triangular part, A_{23} does not matter. As we have just shown, (4.11) is a consequence of $\{e_2, e_2\} = 0$. In the free Lie superalgebra, this relation is not imposed. We thus conclude that the upper-triangular part of the Borchers superalgebra defined by (4.3) is a free Lie superalgebra on the $A_1 \oplus A_1$ representations $[1, 0] \oplus [0, 1]$ modulo the Serre relations (4.9) corresponding to an ideal generated by $[2, 0] \oplus [0, 2]$ at level $\ell = 2$.

⁶At this point, we note that there is the possibility of writing down a Borchers algebra whose positive levels agrees with the free Lie superalgebra including P_{ab} . In order to arrange for this, one has to consider more general Cartan matrices than the ones we have introduced in (4.1) by also allowing negative diagonal entries, e.g. $A_{ii} = -2$. The defining relations can for instance be found in [69] and the corresponding nodes are sometimes called ‘black’ nodes. For black nodes, the relation (4.2c) does not apply and therefore there are no Serre relations at positive levels associated with the black nodes, in other words they form a free Lie superalgebra.

B_ℓ	representation	generator
B_0	$[2, 0] \oplus [0, 2]$	M_{ab}
	$[0, 0]$	B_5
	$[0, 0]$	d
B_1	$[1, 0] \oplus [0, 1]$	Q_α
B_2	$[1, 1]$	P_a
B_3	$[1, 0] \oplus [0, 1]$	Σ_α
B_4	$[2, 0] \oplus [0, 2]$	$Z_{ab} = Z_{[ab]}$
	$[0, 0]$	B
B_5	$[1, 0] \oplus [0, 1]$	X_α
	$[1, 2] \oplus [2, 1]$	$X_{a\alpha}$
B_6	$[2, 0] \oplus [0, 2]$	
	$[1, 1]$	
	$[1, 1]$	
	$[1, 1]$	
	$[1, 3] \oplus [3, 1]$	
\vdots	\vdots	\vdots

Table 6. The non-negative levels of the Borcherds superalgebra defined by the Cartan matrix (4.3). This is a quotient of the free Lie superalgebra shown in table 2 and the notation is in that table. We have also included the Cartan generators from level $\ell = 0$. B_5 acts as chirality while d is a scaling operator. Up to level $\ell = 4$ this is the Maxwell superalgebra studied in [11].

The representation content of the positive levels of the Borcherds superalgebra can be constructed recursively using a similar algorithm as for the free Lie superalgebra. In the Borcherds superalgebra one has

$$B_1 = \langle Q_\alpha \rangle, \tag{4.12a}$$

$$B_2 = \vee^2 B_1 \ominus S_2, \tag{4.12b}$$

$$B_3 = B_2 \otimes B_1 \ominus \vee^3 B_1 \oplus S_3, \tag{4.12c}$$

$$B_4 = (B_3 \otimes B_1 \oplus \wedge^2 B_2) \ominus \vee^2 B_1 \otimes B_2 \oplus \vee^4 B_1 \ominus S_4, \tag{4.12d}$$

...

where the difference to the free Lie superalgebra (2.9) is the presence of a certain representation S_ℓ at each level that has to be removed. We have already seen that $S_2 = [2, 0] \oplus [0, 2]$ and an analysis similar to the one carried out in [26] for the case of only one gray node yields that in general $S_\ell = [\ell, 0] \oplus [0, \ell]$. This is displayed in table 6.

Alternatively, the representation content of the Borcherds superalgebra up to an arbitrary level k can be derived from the representation content of a corresponding Kac-Moody




Figure 2. Dynkin diagram of the Kac-Moody algebra of rank $k + 3$ corresponding to the Borchers superalgebra with the Dynkin diagram in figure 1 by the ‘Borchers-Kac-Moody correspondence’. The number of nodes in the lower line is $k - 1$, forming an A_{k-1} subdiagram. For $k = 3$, this Kac-Moody algebra is the exceptional Lie algebra E_6 , for $k = 4$ its affine extension E_6^+ , and for $k = 5$ its hyperbolic extension E_6^{++} .

algebra. This can for example be more useful than the method above when the representations S_ℓ are not known. This ‘Borchers-Kac-Moody correspondence’ was first explained in [18] in the context of E_{11} (see also [20]) and then generalised to other cases in [19, 70]. Here we will just briefly describe the consequence of the general procedure in the special case of the Borchers superalgebra with Cartan matrix (4.3) or with the Dynkin diagram in figure 1, and it reveals a remarkable connection between the Maxwell superalgebra and the exceptional Lie algebra E_6 . The method is only directly applicable if $A_{23} = A_{32} = -2$, and this is the reason why we have made this choice, although, as we have seen, any negative value of these entries in the Cartan matrix gives the same representation contents in the level decomposition.

The Dynkin diagram of the corresponding Kac-Moody algebra is obtained by replacing the gray nodes with ordinary white ones, removing the double line, and adding another $k - 1$ white nodes, each connected to the previous one with a single line, so that they form the Dynkin diagram of the Lie algebra A_{k-1} with split real form $\mathfrak{sl}(k)$. Furthermore, the new nodes 2 and 3 replacing the gray ones should be connected to the first node in this A_{k-1} subdiagram. The resulting Dynkin algebra is shown in figure 2. For $k = 3$, the corresponding Kac-Moody algebra is the exceptional Lie algebra E_6 , for $k = 4$ its affine extension E_6^+ , and for $k = 5$ its hyperbolic extension E_6^{++} . A level decomposition of this Kac-Moody algebra can be performed with respect to nodes 2 and 3 in the same way as for the original Borchers superalgebra, and since we now have a Lie algebra rather than a Lie superalgebra, the representations can be computed easily using the software `SimpLie` [71]. Furthermore, since we now at level $\ell = 0$ have not only $A_1 \oplus A_1$ and the Cartan generators h_2 and h_3 , but also $\mathfrak{sl}(k)$, each representation of $A_1 \oplus A_1$ comes together with a representation of $\mathfrak{sl}(k)$. It turns out that, at each level $\ell \leq k$, the $A_1 \oplus A_1$ representation that comes together with the k -th antisymmetrised tensor power of the fundamental k -dimensional representation of $\mathfrak{sl}(k)$ is precisely the representation B_ℓ appearing in the Borchers superalgebra. This can be verified for $k = 5$ by comparing tables 6 and 7. Using either tool we obtain the supersymmetric Maxwell algebra studied in [11] with an additional scaling operator d that can be chosen to have the level ℓ as its eigenvalue.

level ℓ	$A_1 \oplus A_1 \oplus A_4$ representation	generator
0	$[2, 0; 0, 0, 0, 0] \oplus [0, 2; 0, 0, 0, 0]$	M_{ab}
	$[0, 0; 0, 0, 0, 0]$	B_5
	$[0, 0; 0, 0, 0, 0]$	d
	$[0, 0; 1, 0, 0, 1]$	
1	$[1, 0; 1, 0, 0, 0] \oplus [0, 1; 1, 0, 0, 0]$	Q_α^I
2	$[1, 1; 0, 1, 0, 0]$	P_a^{IJ}
3	$[1, 0; 0, 0, 1, 0] \oplus [0, 1; 0, 0, 1, 0]$	Σ_α^{IJK}
4	$[2, 0; 0, 0, 0, 1] \oplus [0, 2; 0, 0, 0, 1]$	$Z_{ab}^{I_1 \dots I_4}$
	$[0, 0; 0, 0, 0, 1]$	$B^{I_1 \dots I_4}$
	$[0, 0; 1, 0, 1, 0]$	
5	$[1, 0; 0, 0, 0, 0] \oplus [0, 1; 0, 0, 0, 0]$	$X_\alpha^{I_1 \dots I_5}$
	$[1, 2; 0, 0, 0, 0] \oplus [2, 1; 0, 0, 0, 0]$	$X_{a\alpha}^{I_1 \dots I_5}$
	$[1, 0; 1, 0, 0, 1] \oplus [0, 1; 1, 0, 0, 1]$	
\vdots	\vdots	\vdots

Table 7. The non-negative levels of E_6^{++} (the hyperbolic extension of E_6) with Dynkin diagram in figure 2 for $k = 5$, decomposed with respect to the $A_1 \oplus A_1 \oplus A_4$ subalgebra obtained by removing node 2 and 3. The Dynkin labels of A_4 are separated from those of $A_1 \oplus A_1$ with a semicolon. Restricting to A_4 representations where the ℓ -th Dynkin label is equal to one and the others to zero (all zero for $\ell = 0$ and $\ell = 5$) we obtain the $A_1 \oplus A_1$ representations B_ℓ in table 2. We have illustrated this by putting ℓ fully antisymmetric indices $I, J, \dots = 1, \dots, 5$ on the generators in table 2 up to $\ell = 5$ (which coincide with those of the Maxwell superalgebra up to $\ell = 4$).

4.2 $D = 4$ and extended supersymmetry

The Borchers superalgebras considered in the preceding subsection can be extended in a way corresponding to extended supersymmetry. In order to get an R-symmetry algebra $\mathfrak{su}(\mathcal{N})$ at level $\ell = 0$ (commuting with the Lorentz algebra), and the two Weyl spinors at level $\ell = 1$ transforming in two \mathcal{N} -dimensional representations of $\mathfrak{su}(\mathcal{N})$, dual to each other (‘fundamental’ and ‘anti-fundamental’), we insert $\mathcal{N} - 1$ white nodes between the two gray nodes in the Dynkin diagram, as shown in figure 3. The generators e_2 and e_3 are then still lowest weight vectors of two representations of the subalgebra at level $\ell = 0$, but since this subalgebra now also contains $A_{\mathcal{N}-1}$ in addition to the (complexified) Lorentz algebra $A_1 \oplus A_1$, the representations now have $1 + 1 + (\mathcal{N} - 1)$ Dynkin labels. As in section 3.2 we have put the Dynkin labels corresponding to $A_{\mathcal{N}-1}$ after the two Dynkin labels corresponding to $A_1 \oplus A_1$, separated by a semicolon. The Dynkin labels of the two representations at level $\ell = 1$ are then $[1, 0; 1, 0, \dots, 0]$ and $[0, 1; 0, \dots, 0, 1]$, respectively, as can be verified by acting on e_2 and e_3 with the Cartan generators.




Figure 3. Dynkin diagram of the Borchers superalgebra for \mathcal{N} -extended supersymmetry. The middle part is the Dynkin diagram of type $A_{\mathcal{N}-1}$ corresponding to the complexified $\mathfrak{su}(\mathcal{N})$ R-symmetry. The outer nodes labelled 1 and 4 correspond to $A_1 \oplus A_1$ as before while the two gray nodes correspond to the two spinors.




Figure 4. Distinguished Dynkin diagram of the finite-dimensional contragredient Lie superalgebra $A(3, \mathcal{N} - 1)$, which has the superconformal algebra $\mathfrak{su}(2, 2|\mathcal{N})$ in $D = 4$ as a real form.

We also give the Cartan matrix corresponding to the simplest extended case $\mathcal{N} = 2$ for which there are five nodes in figure 3; it is

$$\begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & -1 & -1 \\ 0 & 0 & -1 & 2 & 0 \\ 0 & -1 & -1 & 0 & 2 \end{pmatrix}. \quad (4.13)$$

Row and column 5 correspond to the R-symmetry A_1 in the middle of the diagram.

At this point we note a certain similarity to the superconformal algebra $\mathfrak{su}(2, 2|\mathcal{N})$ in $D = 4$. This is a real form of the complex Lie superalgebra $A(3, \mathcal{N} - 1)$, which has a distinguished Dynkin diagram shown in figure 4. The fact that the Dynkin diagram is distinguished means that there is only one gray node, and this condition makes it unique. By relaxing this condition and performing so-called odd reflections [68] one can obtain equivalent non-distinguished Dynkin diagrams of the same Lie superalgebra, with more than one gray node. Following the rules for odd reflections (see for example [72]) it is straightforward to show that one of the possible diagrams look the same as the one in figure 3 [68, 73]. However, the meaning of gray nodes in this context is slightly different. The corresponding e and f generators are still odd elements, and the corresponding diagonal entries are still zero, but on one side of the gray node the corresponding entries in the Cartan matrix have opposite signs. For figure 4, this means that if the white nodes and lines on the left hand side of the gray node correspond to diagonal entries 2 and off-diagonal entries -1 , respectively, then those on the right hand side have diagonal entries -2 and off-diagonal entries 1, respectively. For figure 3, with $\mathcal{N} = 2$, this gives the corresponding Cartan matrix

$$\begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & -1 & 1 \\ 0 & 0 & -1 & 2 & 0 \\ 0 & 1 & 1 & 0 & -2 \end{pmatrix}, \quad (4.14)$$




Figure 5. Dynkin diagram of a Borchers Lie superalgebra that is related to a quotient of the free Lie superalgebra of $D = 11$ supersymmetry.

B_ℓ	B_5 representation	generator
B_0	$[0, 1, 0, 0, 0]$	M_{ab}
	$[0, 0, 0, 0, 0]$	d
B_1	$[0, 0, 0, 0, 1]$	Q_α
B_2	$[1, 0, 0, 0, 0]$	P_a
	$[0, 1, 0, 0, 0]$	P_{ab}
B_3	$[0, 0, 0, 0, 1]$	Σ_α
	$[1, 0, 0, 0, 1]$	$\Sigma_{a\alpha}$
B_4	$2 \times [0, 1, 0, 0, 0]$	$Z_{ab} = Z_{[ab]}$
	$[1, 0, 0, 0, 0]$	
	$[0, 0, 1, 0, 0]$	
	$[1, 1, 0, 0, 0]$	
	$[0, 0, 0, 0, 2]$	
	$[1, 0, 1, 0, 0]$	
$[1, 0, 0, 0, 0]$		
\vdots	\vdots	\vdots

Table 8. The non-negative levels of the Borchers superalgebra defined by the Dynkin diagram of figure 5. This is a quotient of the free Lie superalgebra shown in table 4 and the notation is as in that table.

of $A(3,1)$, where the entries in row or column 5 have opposite sign compared to (4.13). Accordingly, $A(3,1)$ is not a Borchers superalgebra but a contragredient Lie superalgebra [67], and this change of signs makes it finite-dimensional.

4.3 $D = 11$ and $\mathcal{N} = 1$ supersymmetry

We can also apply a Borchers construction to the eleven-dimensional case. The starting point for the Dynkin diagram is that of B_5 which is the complexification of $\mathfrak{so}(1,10)$. To obtain a spinor representation at level $\ell = 1$ one can attach a gray node to the ‘spinor node’ of the B_5 diagram with a single line. This leads to figure 5. The level decomposition of the Borchers algebra defined by this diagram can be computed using the methods described above and this leads to the $\mathfrak{so}(1,10)$ representation listed in table 8.

Table 8 should be compared to table 4 from which we see that the Borchers construction contains considerably fewer generators than the free Lie superalgebra. This is

of course expected since the positive levels of the Borcherds algebra are a quotient of the free Lie superalgebra by the Serre ideal. One noteworthy difference between the Borcherds algebra and the free Lie superalgebra is that the former does not contain the generator $P_{a_1\dots a_5}$ on level $\ell = 2$ associated with the M5-brane term in the superalgebra. Thus it agrees with the algebra given in [46] up to $\ell = 2$, but it extends it beyond that.⁷

5 Conclusion

In this paper, we have put forward the structure of a free Lie superalgebra for studying extensions of the Poincaré superalgebra. The various examples that we have presented comprise generalisations existing in the literature by quotienting and thus the free Lie superalgebra $\mathfrak{f}(Q)$ can be viewed as universal structure that encompasses all these proposals. We have focussed in this paper on the algebraic aspects and have not worked out any dynamical model realising the symmetry but this can be done for the point particle along the lines of [7, 11] and we expect a similar universal behaviour of a superparticle moving in a supersymmetric Maxwell background if a suitable ‘unfolded’ quotient is applied. An aspect that will deserve detailed study in this context is the κ -symmetry of the particle action. Extensions to string and brane models moving in backgrounds can also be constructed. A generalisation of the Polyakov-type term for the string invariant under the superalgebra can be written as

$$\int d^2\sigma \sqrt{-\gamma} \gamma^{ij} \left[L_i^a L_j^b \eta_{ab} + \frac{1}{2} L_i^{ab} L_j^{cd} \eta_{ac} \eta_{bd} + \dots \right], \tag{5.1}$$

where the $L_i^a = \partial_i x^a + \dots$ and $L_i^{ab} = \partial_i \theta^{ab} + \dots$ denotes the components of the Cartan-Maurer forms of a string moving on the group manifold of the superalgebra [7, 40].

Let us also point out a few other possible applications or connections to other work. As was emphasised in section 2 the construction of a free Lie (super-)algebra starts from a set of basic generators, e.g., the Q_α . Demanding that these basic generators transform in a representation of some level $\ell = 0$ algebra \mathfrak{g}_0 then leads to each level ℓ of the free Lie algebra $\mathfrak{f}(Q)$ transforming under \mathfrak{g}_0 . In the discussion of the free Lie superalgebra for $D = 11$ in section 3.3.1 we considered the case $\mathfrak{so}(1, 10)$, the eleven-dimensional Lorentz algebra. In view of possible relations to E_{11} it is worth noting that the Lorentz algebra is generalised to an algebra called $K(E_{11})$ (or sometimes $I_C(E_{11})$) in the context of West’s conjectures. This is an infinite-dimensional Lie algebra that is fixed by an involution acting on (the split real) E_{11} , and $K(E_{11})$ contains $\mathfrak{so}(1, 10)$. The general representation theory of this algebra has not been classified as it is not of standard type. However, it is known that $K(E_{11})$ can be made to act on the same 32 supertranslations Q_α that $\mathfrak{so}(1, 10)$ acts on [74, 75]. In this way, the whole free Lie superalgebra $\mathfrak{f}(Q)$ can be seen as a representation

⁷As in footnote 6, we note that it is in fact possible to get also $P_{a_1\dots a_5}$ at level $\ell = 2$ in a Borcherds superalgebra, but then one has to interchange the gray node 6 in figure 5 for a black node such that the corresponding diagonal entry in the Cartan matrix is not zero but negative. This removes the Serre relation $\{e_6, e_6\} = 0$ and gives rise to a 5-form since $\{e_6, e_6\}$ then is a lowest weight vector of the representation $[0, 0, 0, 0, 2]$.

of $K(E_{11})$ providing an infinite-dimensional realisation of $K(E_{11})$ albeit still of the same unfaithful type. Under $K(E_{11})$ one has for instance that $\ell = 2$ of $\mathfrak{f}(Q)$ forms an irreducible 528-dimensional representation. This perspective opens up the possibility of constructing particle or brane models with $K(E_{11})$ symmetry.

Yet another possibility of applying the free Lie superalgebra is in the context of so-called vector supersymmetry [76, 77]. In vector supersymmetry, the basic supersymmetry generators are not spinors Q_α but transform in a vector and a scalar representation of the Lorentz group. A particle realisation of this vector supersymmetry leads after quantisation to two uncoupled Dirac equations [76, 77].

In the physical realisations of Maxwell symmetries studied to date, the background field was non-dynamical. The proposals of [15, 78, 79] attempt to also derive the equations of a gravitational background from an algebraic formulation while [80] discusses the unfolding of the Maxwell equation for the gauge field itself. These constructions are based on algebras that are sometimes called Ogievetskii algebras that include translation generators and higher rank cousins of the type contained in free Lie algebras $\mathfrak{f}(P)$, however, the commutation relations appear to differ from those of the free Lie algebras. It is not clear to us how to reconcile these two pictures at the moment.

Acknowledgments

We would like to thank Carlo Meneghelli, Hermann Nicolai and Ergin Sezgin for helpful discussions. AK and JP thank the University of Barcelona for its hospitality. JG and AK would also like to thank the University of Brussels and the International Solvay Institutes for hospitality. JG has been supported in part by MINECO FPA2016-76005-C2-1-P and Consolider CPAN and by the Spanish government (MINECO/FEDER) under project MDM-2014-0369 of ICCUB (Unidad de Excelencia María de Maeztu). The work of JP is supported by the Swedish Research Council, project no. 2015-04268. It began at the Mitchell Institute for Fundamental Physics and Astronomy at the Texas A&M University and was then supported in part by NSF grant PHY-1214344.

A Gamma matrices in $D = 4$

An explicit real representation of the Γ^a matrices can be written as

$$\begin{aligned} \Gamma^0 &= \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{pmatrix} & \Gamma^1 &= \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix} \\ \Gamma^2 &= \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} & \Gamma^3 &= \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{pmatrix} \end{aligned}$$

$$C = \Gamma^0 = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{pmatrix} \quad \Gamma_5 = \Gamma^0 \Gamma^1 \Gamma^2 \Gamma^3 = \begin{pmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{pmatrix}. \quad (\text{A.1})$$

The matrix Γ^0 is anti-symmetric while the Γ^i (for $i = 1, 2, 3$) are symmetric. They satisfy $\{\Gamma^a, \Gamma^b\} = 2\eta^{ab}$ with $\eta^{ab} = \text{diag}(-1, 1, 1, 1)$. One also has $C^T = -C = C^{-1}$ and that

$$(C\Gamma^a)^T = C\Gamma^a, \quad (C\Gamma^{ab})^T = C\Gamma^{ab}, \quad (C\Gamma^{abc})^T = -C\Gamma^{abc}, \quad (C\Gamma^{abcd})^T = -C\Gamma^{abcd}, \quad (\text{A.2})$$

where $\Gamma^{ab} = \Gamma^{[a}\Gamma^b] = \frac{1}{2}(\Gamma^a\Gamma^b - \Gamma^b\Gamma^a)$ etc. The matrix Γ_5 anti-commutes with all the Γ^a . It squares to $(\Gamma_5)^2 = -\mathbb{1}$ and $(C\Gamma_5)^T = -C\Gamma_5$.

We use the following explicit spinor index notation on the gamma matrices, following the SW-NE contraction convention,

$$(\Gamma^a)^\alpha_\beta \quad \Rightarrow \quad (\Gamma^a\Gamma^b)^\alpha_\beta = (\Gamma^a)^\alpha_\gamma (\Gamma^b)^\gamma_\beta \quad (\text{A.3})$$

together with $C_{\alpha\beta}$ for the anti-symmetric charge conjugation matrix and $C^{\alpha\beta}$ for its inverse satisfying $C_{\alpha\gamma}C^{\gamma\beta} = \delta_\alpha^\beta$. The spinor index range is $\alpha = 1, \dots, 4$. The first relation in (A.2) reads in indices $(C\Gamma^a)_{\alpha\beta} = (C\Gamma^a)_{\beta\alpha}$, where $(C\Gamma^a)_{\alpha\beta} = C_{\alpha\gamma}(\Gamma^a)^\gamma_\beta$. We also note the cyclic identity that is satisfied by the gamma matrices in $D = 4$ are

$$(C\Gamma_a)_{(\alpha\beta}(C\Gamma^a)_{\gamma)\delta} = 0, \quad (\text{A.4a})$$

$$(C\Gamma_{ab})_{(\alpha\beta}(C\Gamma^{ab})_{\gamma)\delta} = 0, \quad (\text{A.4b})$$

$$(C\Gamma^a)_{(\alpha\beta}C_{\gamma)\delta} + (C\Gamma^{ab})_{(\alpha\beta}(C\Gamma_b)_{\gamma)\delta} = 0. \quad (\text{A.4c})$$

The identity (A.4a) and similar ones in $D = 3, 6, 10$ are important for the construction of supersymmetric Yang-Mills theories [81, 82], whereas (A.4b) is crucial for supersymmetric membranes [36]. In the context of free Lie superalgebras they appear in the Jacobi identity (2.7).

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] H. Bacry, P. Combe and J.L. Richard, *Group-theoretical analysis of elementary particles in an external electromagnetic field. 1. the relativistic particle in a constant and uniform field*, *Nuovo Cim. A* **67** (1970) 267 [[INSPIRE](#)].
- [2] H. Bacry, P. Combe and J.L. Richard, *Group-theoretical analysis of elementary particles in an external electromagnetic field. 2. the nonrelativistic particle in a constant and uniform field*, *Nuovo Cim. A* **70** (1970) 289 [[INSPIRE](#)].
- [3] R. Schrader, *The Maxwell group and the quantum theory of particles in classical homogeneous electromagnetic fields*, *Fortsch. Phys.* **20** (1972) 701 [[INSPIRE](#)].

- [4] J. Beckers and V. Hussin, *Minimal electromagnetic coupling schemes. II. Relativistic and nonrelativistic Maxwell Groups*, *J. Math. Phys.* **24** (1983) 1295 [INSPIRE].
- [5] D.V. Soroka and V.A. Soroka, *Tensor extension of the Poincaré algebra*, *Phys. Lett.* **B 607** (2005) 302 [hep-th/0410012] [INSPIRE].
- [6] S. Bonanos and J. Gomis, *Infinite Sequence of Poincaré Group Extensions: Structure and Dynamics*, *J. Phys.* **A 43** (2010) 015201 [arXiv:0812.4140] [INSPIRE].
- [7] J. Gomis and A. Kleinschmidt, *On free Lie algebras and particles in electro-magnetic fields*, *JHEP* **07** (2017) 085 [arXiv:1705.05854] [INSPIRE].
- [8] J.W. van Holten and A. Van Proeyen, *$N = 1$ Supersymmetry Algebras in $D = 2$, $D = 3$, $D = 4$ MOD-8*, *J. Phys.* **A 15** (1982) 3763 [INSPIRE].
- [9] P.K. Townsend, *P-brane democracy*, in *The World in eleven-dimensions: A Tribute to Oskar Klein*, M.J. Duff eds., CRC Press, Boca Raton U.S.A. (1995), pg. 375 [hep-th/9507048] [INSPIRE].
- [10] E. Sezgin, *The M algebra*, *Phys. Lett.* **B 392** (1997) 323 [hep-th/9609086] [INSPIRE].
- [11] S. Bonanos, J. Gomis, K. Kamimura and J. Lukierski, *Maxwell Superalgebra and Superparticle in Constant Gauge Backgrounds*, *Phys. Rev. Lett.* **104** (2010) 090401 [arXiv:0911.5072] [INSPIRE].
- [12] K. Kamimura and J. Lukierski, *Supersymmetrization Schemes of $D = 4$ Maxwell Algebra*, *Phys. Lett.* **B 707** (2012) 292 [arXiv:1111.3598] [INSPIRE].
- [13] M.B. Green, *Supertranslations, Superstrings and Chern-Simons Forms*, *Phys. Lett.* **B 223** (1989) 157 [INSPIRE].
- [14] V.G. Kac, *Infinite dimensional Lie algebras*, Cambridge University Press, Cambridge U.K. (1990).
- [15] P.C. West, *E_{11} and M-theory*, *Class. Quant. Grav.* **18** (2001) 4443 [hep-th/0104081] [INSPIRE].
- [16] T. Damour, M. Henneaux and H. Nicolai, *E_{10} and a 'small tension expansion' of M-theory*, *Phys. Rev. Lett.* **89** (2002) 221601 [hep-th/0207267] [INSPIRE].
- [17] R.E. Borcherds, *Generalized Kac-Moody algebras*, *J. Algebra* **155** (1988) 501.
- [18] M. Henneaux, B.L. Julia and J. Levie, *E_{11} , Borcherds algebras and maximal supergravity*, *JHEP* **04** (2012) 078 [arXiv:1007.5241] [INSPIRE].
- [19] J. Palmkvist, *Borcherds and Kac-Moody extensions of simple finite-dimensional Lie algebras*, *JHEP* **06** (2012) 003 [arXiv:1203.5107] [INSPIRE].
- [20] J. Palmkvist, *Tensor hierarchies, Borcherds algebras and E_{11}* , *JHEP* **02** (2012) 066 [arXiv:1110.4892] [INSPIRE].
- [21] J. Palmkvist, *The tensor hierarchy algebra*, *J. Math. Phys.* **55** (2014) 011701 [arXiv:1305.0018] [INSPIRE].
- [22] J. Greitz, P. Howe and J. Palmkvist, *The tensor hierarchy simplified*, *Class. Quant. Grav.* **31** (2014) 087001 [arXiv:1308.4972] [INSPIRE].
- [23] G. Bossard, A. Kleinschmidt, J. Palmkvist, C.N. Pope and E. Sezgin, *Beyond E_{11}* , *JHEP* **05** (2017) 020 [arXiv:1703.01305] [INSPIRE].

- [24] A.G. Cohen and S.L. Glashow, *Very special relativity*, *Phys. Rev. Lett.* **97** (2006) 021601 [[hep-ph/0601236](#)] [[INSPIRE](#)].
- [25] G.W. Gibbons, J. Gomis and C.N. Pope, *Deforming the Maxwell-Sim Algebra*, *Phys. Rev. D* **82** (2010) 065002 [[arXiv:0910.3220](#)] [[INSPIRE](#)].
- [26] M. Cederwall and J. Palmkvist, *Superalgebras, constraints and partition functions*, *JHEP* **08** (2015) 036 [[arXiv:1503.06215](#)] [[INSPIRE](#)].
- [27] E. Witt, *Treue Darstellung Liescher Ringe*, *J. Reine Angew. Math.* **177** (1937) 152.
- [28] P. Salgado, R.J. Szabo and O. Valdivia, *Topological gravity and transgression holography*, *Phys. Rev. D* **89** (2014) 084077 [[arXiv:1401.3653](#)] [[INSPIRE](#)].
- [29] P. Salgado and S. Salgado, $\mathfrak{so}(D-1, 1) \otimes \mathfrak{so}(D-1, 2)$ algebras and gravity, *Phys. Lett. B* **728** (2014) 5 [[INSPIRE](#)].
- [30] S. Hoseinzadeh and A. Rezaei-Aghdam, $(2+1)$ -dimensional gravity from Maxwell and semisimple extension of the Poincaré gauge symmetric models, *Phys. Rev. D* **90** (2014) 084008 [[arXiv:1402.0320](#)] [[INSPIRE](#)].
- [31] R. Caroca, P. Concha, O. Fierro, E. Rodríguez and P. Salgado-Rebolledo, *Generalized Chern-Simons higher-spin gravity theories in three dimensions*, *Nucl. Phys. B* **934** (2018) 240 [[arXiv:1712.09975](#)] [[INSPIRE](#)].
- [32] P. Concha, D.M. Peñafiel and E. Rodríguez, *On the Maxwell supergravity and flat limit in $2+1$ dimensions*, *Phys. Lett. B* **785** (2018) 247 [[arXiv:1807.00194](#)] [[INSPIRE](#)].
- [33] L. Avilés, E. Frodden, J. Gomis, D. Hidalgo and J. Zanelli, *Non-Relativistic Maxwell Chern-Simons Gravity*, *JHEP* **05** (2018) 047 [[arXiv:1802.08453](#)] [[INSPIRE](#)].
- [34] J.A. de Azcarraga, K. Kamimura and J. Lukierski, *Generalized cosmological term from Maxwell symmetries*, *Phys. Rev. D* **83** (2011) 124036 [[arXiv:1012.4402](#)] [[INSPIRE](#)].
- [35] M.A.A. van Leeuwen, A.M. Cohen and B. Lissner, *LiE, A Package for Lie Group Computations*, available from <http://www.mathlabo.univ-poitiers.fr/~maavl/LiE/>.
- [36] A. Achúcarro, J.P. Gauntlett, K. Itoh and P.K. Townsend, *World Volume Supersymmetry From Space-time Supersymmetry of the Four-dimensional Supermembrane*, *Nucl. Phys. B* **314** (1989) 129 [[INSPIRE](#)].
- [37] S. Ferrara and M. Porrati, *Central extensions of supersymmetry in four-dimensions and three-dimensions*, *Phys. Lett. B* **423** (1998) 255 [[hep-th/9711116](#)] [[INSPIRE](#)].
- [38] C. Chryssomalakos, J.A. de Azcarraga, J.M. Izquierdo and J.C. Perez Bueno, *The Geometry of branes and extended superspaces*, *Nucl. Phys. B* **567** (2000) 293 [[hep-th/9904137](#)] [[INSPIRE](#)].
- [39] E. Bergshoeff and E. Sezgin, *New Space-time Superalgebras and Their Kac-Moody Extension*, *Phys. Lett. B* **232** (1989) 96 [[INSPIRE](#)].
- [40] E. Bergshoeff and E. Sezgin, *Super p -brane theories and new space-time superalgebras*, *Phys. Lett. B* **354** (1995) 256 [[hep-th/9504140](#)] [[INSPIRE](#)].
- [41] E. Cremmer and B. Julia, *The SO(8) Supergravity*, *Nucl. Phys. B* **159** (1979) 141 [[INSPIRE](#)].
- [42] B. de Wit and H. Nicolai, *$N = 8$ Supergravity*, *Nucl. Phys. B* **208** (1982) 323 [[INSPIRE](#)].

- [43] J.A. de Azcarraga, J.P. Gauntlett, J.M. Izquierdo and P.K. Townsend, *Topological Extensions of the Supersymmetry Algebra for Extended Objects*, *Phys. Rev. Lett.* **63** (1989) 2443 [[INSPIRE](#)].
- [44] P.K. Townsend, *M theory from its superalgebra*, *NATO Sci. Ser. C* **520** (1999) 141 [[hep-th/9712004](#)] [[INSPIRE](#)].
- [45] D.M. Peñafiel and L. Ravera, *On the Hidden Maxwell Superalgebra underlying $D = 4$ Supergravity*, *Fortsch. Phys.* **65** (2017) 1700005 [[arXiv:1701.04234](#)] [[INSPIRE](#)].
- [46] L. Ravera, *Hidden role of Maxwell superalgebras in the free differential algebras of $D = 4$ and $D = 11$ supergravity*, *Eur. Phys. J. C* **78** (2018) 211 [[arXiv:1801.08860](#)] [[INSPIRE](#)].
- [47] L. Andrianopoli, R. D'Auria and L. Ravera, *Hidden Gauge Structure of Supersymmetric Free Differential Algebras*, *JHEP* **08** (2016) 095 [[arXiv:1606.07328](#)] [[INSPIRE](#)].
- [48] L. Andrianopoli, R. D'Auria and L. Ravera, *More on the Hidden Symmetries of 11D Supergravity*, *Phys. Lett. B* **772** (2017) 578 [[arXiv:1705.06251](#)] [[INSPIRE](#)].
- [49] P.C. West, *E_{11} , $SL(32)$ and central charges*, *Phys. Lett. B* **575** (2003) 333 [[hep-th/0307098](#)] [[INSPIRE](#)].
- [50] J. de Boer and M. Shigemori, *Exotic branes and non-geometric backgrounds*, *Phys. Rev. Lett.* **104** (2010) 251603 [[arXiv:1004.2521](#)] [[INSPIRE](#)].
- [51] J. de Boer and M. Shigemori, *Exotic Branes in String Theory*, *Phys. Rept.* **532** (2013) 65 [[arXiv:1209.6056](#)] [[INSPIRE](#)].
- [52] N.A. Obers and B. Pioline, *U duality and M-theory*, *Phys. Rept.* **318** (1999) 113 [[hep-th/9809039](#)] [[INSPIRE](#)].
- [53] B. de Wit and H. Nicolai, *Hidden symmetries, central charges and all that*, *Class. Quant. Grav.* **18** (2001) 3095 [[hep-th/0011239](#)] [[INSPIRE](#)].
- [54] E.A. Bergshoeff and F. Riccioni, *String Solitons and T-duality*, *JHEP* **05** (2011) 131 [[arXiv:1102.0934](#)] [[INSPIRE](#)].
- [55] A. Kleinschmidt, *Counting supersymmetric branes*, *JHEP* **10** (2011) 144 [[arXiv:1109.2025](#)] [[INSPIRE](#)].
- [56] E.A. Bergshoeff, F. Riccioni and L. Romano, *Branes, Weights and Central Charges*, *JHEP* **06** (2013) 019 [[arXiv:1303.0221](#)] [[INSPIRE](#)].
- [57] E.A. Bergshoeff, V.A. Penas, F. Riccioni and S. Risoli, *Non-geometric fluxes and mixed-symmetry potentials*, *JHEP* **11** (2015) 020 [[arXiv:1508.00780](#)] [[INSPIRE](#)].
- [58] A. Kleinschmidt and P.C. West, *Representations of G_{+++} and the role of space-time*, *JHEP* **02** (2004) 033 [[hep-th/0312247](#)] [[INSPIRE](#)].
- [59] M. Henneaux, A. Kleinschmidt and H. Nicolai, *Real forms of extended Kac-Moody symmetries and higher spin gauge theories*, *Gen. Rel. Grav.* **44** (2012) 1787 [[arXiv:1110.4460](#)] [[INSPIRE](#)].
- [60] D.S. Berman, H. Godazgar, M. Godazgar and M.J. Perry, *The Local symmetries of M-theory and their formulation in generalised geometry*, *JHEP* **01** (2012) 012 [[arXiv:1110.3930](#)] [[INSPIRE](#)].
- [61] D.S. Berman, H. Godazgar, M.J. Perry and P. West, *Duality Invariant Actions and Generalised Geometry*, *JHEP* **02** (2012) 108 [[arXiv:1111.0459](#)] [[INSPIRE](#)].

- [62] A. Coimbra, C. Strickland-Constable and D. Waldram, $E_{d(d)} \times \mathbb{R}^+$ generalised geometry, connections and M-theory, *JHEP* **02** (2014) 054 [[arXiv:1112.3989](#)] [[INSPIRE](#)].
- [63] D.S. Berman, M. Cederwall, A. Kleinschmidt and D.C. Thompson, The gauge structure of generalised diffeomorphisms, *JHEP* **01** (2013) 064 [[arXiv:1208.5884](#)] [[INSPIRE](#)].
- [64] A. Kleinschmidt, E_{11} as E_{10} representation at low levels, *Nucl. Phys. B* **677** (2004) 553 [[hep-th/0304246](#)] [[INSPIRE](#)].
- [65] A. Kleinschmidt and J. Palmkvist, Oxidizing Borchers symmetries, *JHEP* **03** (2013) 044 [[arXiv:1301.1346](#)] [[INSPIRE](#)].
- [66] V.G. Kac, Simple irreducible graded Lie algebras of finite growth, *Math. USSR Izv.* **2** (1968) 1271.
- [67] V.G. Kac, Lie Superalgebras, *Adv. Math.* **26** (1977) 8 [[INSPIRE](#)].
- [68] V.K. Dobrev and V.B. Petkova, Group Theoretical Approach to Extended Conformal Supersymmetry: Function Space Realizations and Invariant Differential Operators, *Fortsch. Phys.* **35** (1987) 537 [[INSPIRE](#)].
- [69] U. Ray, *Automorphic forms and Lie superalgebras*, Springer, Berlin Germany (2006).
- [70] P. Howe and J. Palmkvist, Forms and algebras in (half-)maximal supergravity theories, *JHEP* **05** (2015) 032 [[arXiv:1503.00015](#)] [[INSPIRE](#)].
- [71] E.A. Bergshoeff, I. De Baetselier and T.A. Nutma, E_{11} and the embedding tensor, *JHEP* **09** (2007) 047 [[arXiv:0705.1304](#)] [[INSPIRE](#)].
- [72] L. Frappat and A. Sciarrino and P. Sorba, *Dictionary on Lie algebras and Lie superalgebras*, Academic Press, New York U.S.A. (2000).
- [73] N. Beisert and M. Staudacher, The $N = 4$ SYM integrable super spin chain, *Nucl. Phys. B* **670** (2003) 439 [[hep-th/0307042](#)] [[INSPIRE](#)].
- [74] A. Kleinschmidt and H. Nicolai, IIA and IIB spinors from $K(E_{10})$, *Phys. Lett. B* **637** (2006) 107 [[hep-th/0603205](#)] [[INSPIRE](#)].
- [75] T. Damour, A. Kleinschmidt and H. Nicolai, $K(E_{10})$, Supergravity and Fermions, *JHEP* **08** (2006) 046 [[hep-th/0606105](#)] [[INSPIRE](#)].
- [76] R. Casalbuoni, J. Gomis, K. Kamimura and G. Longhi, Space-time Vector Supersymmetry and Massive Spinning Particle, *JHEP* **02** (2008) 094 [[arXiv:0801.2702](#)] [[INSPIRE](#)].
- [77] A. Barducci, R. Casalbuoni and J. Gomis, Confined dynamical systems with Carroll and Galilei symmetries, *Phys. Rev. D* **98** (2018) 085018 [[arXiv:1804.10495](#)] [[INSPIRE](#)].
- [78] A.B. Borisov and V.I. Ogievetsky, Theory of Dynamical Affine and Conformal Symmetries as Gravity Theory, *Theor. Math. Phys.* **21** (1975) 1179 [[INSPIRE](#)].
- [79] F. Riccioni and P. West, Local E_{11} , *JHEP* **04** (2009) 051 [[arXiv:0902.4678](#)] [[INSPIRE](#)].
- [80] N. Boulanger, P. Sundell and P. West, Gauge fields and infinite chains of dualities, *JHEP* **09** (2015) 192 [[arXiv:1502.07909](#)] [[INSPIRE](#)].
- [81] L. Brink, J.H. Schwarz and J. Scherk, Supersymmetric Yang-Mills Theories, *Nucl. Phys. B* **121** (1977) 77 [[INSPIRE](#)].
- [82] F. Gliozzi, J. Scherk and D.I. Olive, Supersymmetry, Supergravity Theories and the Dual Spinor Model, *Nucl. Phys. B* **122** (1977) 253 [[INSPIRE](#)].