



CHALMERS
UNIVERSITY OF TECHNOLOGY

Alginate and HM-pectin in sports-drink give rise to intra-gastric gelation in-vivo

Downloaded from: <https://research.chalmers.se>, 2025-03-23 10:39 UTC

Citation for the original published paper (version of record):

Marciani, L., Lopez-Sanchez, P., Pettersson, S. et al (2019). Alginate and HM-pectin in sports-drink give rise to intra-gastric gelation in-vivo. *Food and Function*, 10(12): 7892-7899 .
<http://dx.doi.org/10.1039/c9fo01617a>

N.B. When citing this work, cite the original published paper.

Cite this: *Food Funct.*, 2019, 10, 7892

Alginate and HM-pectin in sports-drink give rise to intra-gastric gelation *in vivo*

 Luca Marciani,^{a,b} Patricia Lopez-Sanchez,^{†c} Stefan Pettersson,^d Caroline Hoad,^{a,b} Nichola Abrehart,^{a,b} Martin Ahnoff,^c and Anna Ström^{*e,f}

The addition of gelling polysaccharides to sport-drinks may provide improved tolerability of drinks with high concentration of digestible carbohydrates (CHO), otherwise known to increase the risk of gastrointestinal complaints among athletes under prolonged exercise. The physico-chemical properties of a drink containing 14 wt% of digestible CHO (0.7:1 fructose and maltodextrin-ratio), 0.2 wt% of HM-pectin/alginate and 0.06 wt% sodium chloride were examined under *in vitro* gastric conditions using rheology and large deformation testing. The *in vivo* gelling behaviour of the drink was studied using magnetic resonance imaging of subjects at rest together with blood glucose measurements. The *in vivo* results confirm gelation of the test drink, with no gel remaining in the stomach at 60 min and blood glucose values were similar to control. The physico-chemical characterisation of the acidified test drink confirms the formation of a weak gel through which low M_w CHO can diffuse.

Received 22nd July 2019,
Accepted 19th November 2019

DOI: 10.1039/c9fo01617a

rsc.li/food-function

1. Introduction

Fuel substrate depletion (*i.e.* muscle and liver glycogen) and dehydration (>2% loss in body mass) have been identified as main factors decreasing performance during prolonged (>2 h) moderate to high-intensity exercise.^{1,2} To counteract dehydration and to sustain euglycemia and high carbohydrate (CHO) oxidation rates during competition and prolonged key training sessions, general recommendations encourage athletes to consume <8% glucose polymer and/or mono and disaccharide solutions including 20–50 mEq L⁻¹ sodium over water alone to enhance performance.^{3,4} However, if fluid needs are low (*e.g.* cooler conditions) and exercise duration exceeds 2.5 hours, it may be difficult for performance oriented athletes to provide carbohydrates at recommended rates (up to 1.5 g carbohydrates per min).⁵ Furthermore, excessive hypotonic fluid

consumption (*e.g.* traditional sports drink formulations or water) is a major mechanism involved in exercise-induced hyponatremia⁶ whereas a more concentrated CHO solution may provide a practical strategy to sustain exercise performance and health for both elite and slower recreational level athletes. However, hypertonic drinks have been suggested to increase water retention in the intestines that, together with malabsorption of residual CHO, might increase the risk of gastrointestinal (GI) discomfort.⁷

Attempts to change the basic formulation of CHO-rich products for sports nutrition involves the formation of a gel in various ways.^{8–10} Leiper *et al.* reported high gastric emptying rates for a drink containing a gel-forming high-molecular weight glucose polymer.⁸ Lopez-Sanchez *et al.* loaded alginate gel beads with low M_w CHO (60%). Low- M_w CHO was shown to diffuse unhindered through the beads under simulated gastric and intestinal conditions.⁹ Furthermore, a field study on elite long-distance runners reported high tolerability of an alginate containing drink with 30 wt% of CHO when used in individual training programs.¹¹ The effect of adding polysaccharides, such as alginates, to food or drinks, on uptake of CHO *in vivo* is not clear. While some studies report reduced gastric emptying rate, increased feelings of fullness^{12–16} and attenuated peak glucose and insulin response^{17,18} upon addition of polysaccharides to solid foods or drinks, others report absence of any effect of added fibers.^{19–22} The contradictory results are possibly related to variation in physico-chemical properties (such as viscosity and gel strength) of the consumed food and drinks as it has been suggested that food and/or gels above a certain strength (>0.65 N) may be retained in the stomach.^{13,16}

^aNottingham Digestive Diseases Centre and NIHR Nottingham Biomedical Research Centre, Nottingham University Hospitals NHS Trust and the University of Nottingham, Nottingham, UK

^bSir Peter Mansfield Imaging Centre, University of Nottingham, University Park, Nottingham, NG7 2RD, UK

^cMaurten AB, Biotech Center, Gothenburg, Sweden

^dCenter for Health and Performance, Department of Food and Nutrition, and Sport Science, University of Gothenburg, Sweden

^eApplied Chemistry, Chemistry and Chemical Engineering, Chalmers University of Technology, Gothenburg, Sweden

^fSuMo Biomaterials, VINN Excellence Center, Chalmers University of Technology, Gothenburg, Sweden

[†]Current address: Agrifood and Bioscience, RISE-Research Institutes of Sweden, Gothenburg, Sweden.



Attenuated or reduced CHO uptake is not wanted during prolonged exercise, where maintenance of blood glucose and increased rates of exogenous CHO is pivotal for performance.

The aim of this study was to test the hypotheses that a drink formulation containing low concentrations of HM-pectin and alginate together with a high concentration (14 wt%) of digestible CHO (fructose and maltodextrin) (1) is able to form a weak intra-gastric gel, and (2) has not a major effect on CHO uptake. For this investigation we carried out:

- An in depth *in vitro* characterization of the gels including rheology, microstructure and release of digestible CHO,
- In vivo* magnetic resonance imaging (MRI) of the intra-gastric behavior of the sports drink in healthy volunteers.

2. Materials and methods

Food grade sodium alginate of high guluronate content (Manugel DMB) was obtained from FMC Biopolymers and the pectin was a commercial citrus pectin (Genu Pectin Type B from CP Kelco, Denmark). The alginate had a guluronate content of 60–70% as defined by the supplier. The pectin had a degree of methylesterification (DM) > 50 as given by the supplier. Both alginate and pectin are anionic linear polymers, where the alginate is composed of (1,4)-linked β -D-mannuronic acid and α -L-guluronic acid residues and the pectin contains (1,4) linked α -D-galacturonate. Food grade maltodextrin (D.E. 16–19.9) was obtained from Cargill and food grade fructose (Fructopure 500) was obtained from Tate & Lyle. For simplicity, fructose and maltodextrin will from now on be referred to as digestible CHO. Glucono-delta-lactone (GDL) and NaCl used for *in vitro* experiments were obtained from Sigma-Aldrich, Sweden. For *in vivo* studies food grade NaCl (table salt) was used. Simulated gastric fluid without enzyme (pH 1.1–1.3, containing 0.7 M HCl and 0.1 M NaCl) and simulated intestinal fluid without enzyme (pH 6.5–6.6, containing $\sim 0.62 \text{ g L}^{-1}$ sodium hydroxide and $\sim 6.8 \text{ g L}^{-1}$ potassium phosphate monobasic) were obtained from Sigma Aldrich.

2.1 Preparation of samples

Preparation of test drink. The alginate, pectin, maltodextrin, fructose and NaCl were dry-mixed before adding to deionised water. For *in vivo* studies bottled water was used. The total polysaccharide concentration was 0.2% (in the dissolved drink) and the ratio of alginate to pectin was 60 : 40. The total digestible CHO (low molecular weight CHO, maltodextrin and fructose) concentration was 14 wt% and the ratio between maltodextrin and fructose was 1 : 0.7. The NaCl concentration was 0.06%. Osmolality of the drink was 490 mOsm kg^{-1} and pH 6.0. The details are summarised in Table 1.

Preparation of control drink. Maltodextrin, fructose and NaCl were used at the same concentrations and ratio as above, dry-mixed and added to water to yield a drink containing 14 wt% CHO and 0.06% NaCl. Osmolality of the drink was 485 mOsm kg^{-1} and pH 7.2. The details are summarised and compared to test drink in Table 1.

Table 1 Characteristics of test and control drinks

	Test drink	Control
<i>Contents per serving (g)</i>		
Total carbohydrates	31.7	31.7
Maltodextrin	18.1	18.1
Fructose	13.6	13.6
Sodium (Na ⁺)	0.20	0.20
Water	201	224
<i>Other ingredients</i>		
Alginate, pectin	—	—
pH	6.0	7.2
Osmolality ^a (mOsm kg ⁻¹ H ₂ O)	490	485

^aOsmolality was measured using a Type 13 Autocal osmometer (Roebbling Messentechnik, Bremen, Germany).

2.2 Characterisation of the gel

Rheology. Rotational rheology was used to determine the viscosity of the drink and oscillatory rheology to determine the pH of gelation and gel strength of the acidified drink. The rheometer used was stress controlled from Physica, Anton Paar, Germany, model MCR 300. A cone and plate geometry was used. The cone had a diameter of 50 mm and an angle of 1° (gap 50 μm). A shear sweep from 1 to 100 s^{-1} was selected to carry out viscosity measurements. GDL (0.75 g) was added to the drink (10 mL), quickly mixed until GLD was dispersed and loaded on the rheometer prior to gelation. To reduce evaporation a solvent trap was used. A small amplitude oscillatory shear test was carried out at a strain of 0.5% (chosen from the linear viscoelastic region) and frequency of 1 Hz. The change in pH over time was followed in parallel on a sample standing on the lab bench and measured using a pH meter. The measurements were performed at 37 °C controlled by a Peltier system.

Compression tests. GDL (2.25 g) was added to 30 mL of the drink while mixing until GDL was dispersed. The solution was poured into cylindrical moulds (10 mm diameter and 10 mm height) and the gels were let to cure for 48 hours at room temperature. After 48 hours the gels were gently removed from the moulds and their compression strength was measured using a texture analyser (HDi, Stable Micro Systems). Measurements were performed with a cylindrical probe of 20 mm diameter. Emery paper was glued to the probe and the bottom plate to reduce slippage. The compression speed was 0.1 mm s^{-1} . Average stress and strain at fracture of 8 gels were calculated. The final pH of the gels after 48 hours curing was 2.1.

Transmission electron microscopy. The microstructure of the alginate/HM pectin gels was studied by transmission electron microscopy (TEM). The gels were fixed in 2% glutaraldehyde solution. Dehydration was performed in a graded ethanol series starting at 30% ethanol, ending with propylene oxide prior to resin infiltration in TLV resin (TAAB Low Viscosity Resin). The samples were embedded in TLV resin and polymerized for 20 hours at 60 °C. Ultra-thin sections, approximately 100 nm thick, were prepared with a diamond knife using an ultramicrotome (PowerTome XL, RMC Products, Boeckeler Instruments, Tucson, AZ). The ultrathin sections were placed on 400-mesh gold grids and stained to visualise the polysac-



charides. The staining was done according to (Thiery, 1967) using periodic acid, thiosemicarbazide and silver proteinate. The thin-sectioned alginate gels were characterized with a TEM (LEO 706E, LEO Electron Microscopy, Oberkochen, Germany) at an accelerating voltage of 80 kV equipped with a very light sensitive CCD camera (Proscan).

Drink gelation in simulated gastric fluid. The test drink (20 mL) was gelled in simulated gastric fluid SGF (10 mL) in 7 different beakers. The formed gel was collected from each beaker after 0.5, 1, 2, 5, 10, 30 and 60 min, with the help of a metal sieve, and its weight measured. The CHO content in the remaining liquid was measured using a brixmeter (refractometer PAL-3, Atago, Tokyo). Measurements were done in duplicates. Results are shown as cumulative release as a function of time in gastric fluid:

$$\text{Cumulative release} = \frac{C_t}{C_\infty} \quad (1)$$

where C stands for the solute mass released in the medium at time (t) and infinite time (∞). C_∞ was set to equilibrium concentration, why a cumulative release of 1 would represent 0.093 g mL⁻¹ small M_w CHO. The results were corrected by the brix (%) in gastric fluid which was 0.4%. The experiment was repeated twice.

2.3 In vivo MRI study

Participants. The University of Nottingham Faculty of Medicine and Health Sciences Research Ethics Committee granted Ethics approval for this study and all participants gave written informed consent. Table 2 outlines the characteristics of the study participants showing the subjects having BMI between 15 and 23 kg m⁻², age 19 and 33, 2 males and 6 females.

Experimental design. This was a 2-way randomized, double-blind, crossover study in healthy adult volunteers. The participants attended in the morning after an overnight fast. Following a protocol similar to previous work with carbohydrate drinks,²³ they underwent a baseline fasted scan 45 min before receiving the test drink, provided in an opaque sports drink bottle. The test and control drinks were prepared and provided to the participants by a research fellow not involved in the data analysis, following a randomization blind code that was broken only after data analysis was completed. The participants ingested the control and/or test drink at a

volume of 500 ml after which they underwent a second MRI scan 15 min later, followed by another scan every hour for 5 h. At each MRI imaging time point the volunteers were asked to fill in an abdominal symptoms score questionnaire as previously used.²⁴ Capillary blood glucose levels were measured using the finger prick method. Single-use lancets (Unistix Owen Mumford, Oxfordshire, United Kingdom) and a hand-held blood glucose meter (Accu-check, Roche Diagnostics), were used.

MRI. MRI was carried out in the supine position on a 3 T Philips Achieva (Philips, Best, the Netherlands) scanner using a parallel imaging body coil wrapped around the abdomen. An axial HASTE (half-Fourier single shot turbo spin echo) sequence was acquired across the abdomen to measure gastric volumes and hence assess gastric emptying. Slice thickness was 10 mm with 30 axial slices acquired to cover the full stomach anatomy. This set of images was also used to select the axial imaging plane for quantitative T_2 mapping measurement of the transverse relaxation time of the gastric contents.²⁵ Each image set was acquired on a short breath hold.

Data analysis. Commercial software (Analyze 6, Biomedical Imaging Resources, Mayo Clinic, Rochester, MN) was used to trace manually around the region of interest (ROI) on each axial MRI image of the stomach contents. Text files containing the volumes or their signal intensity of each ROI for a given time point were extracted and the gastric volumes and T_2 values respectively calculated.

3. Results and discussion

The composition of the test drink, in terms of type and ratio of alginate and pectin was chosen so to form gels in the presence of acid.^{21,26} The contents of maltodextrin and fructose (multiple transporter CHO solutions) for the control and test drink were chosen based on previous research demonstrating increased intestinal CHO absorption and higher exogenous CHO oxidation rates for fructose-glucose/glucose polymer mixtures compared to isoenergetic glucose/glucose polymer intake only.²⁷

3.1 Physico-chemical characterisation of the test drink

The test drink, prepared as outlined in the Materials and methods section, is characterized by a Newtonian flow and with a shear viscosity of 6.5 ± 0.9 mPa s. The gelation of the drink was followed *in vitro* as a function of pH where pH was reduced using the slowly hydrolysed lactone, GDL. The GDL was dispersed into the test drink, added to the rheometer while still a fluid, and let to set on the rheometer prior measurements of storage (G') and loss (G'') modulus (Fig. 1A). A gel (here defined as $G' > G''$) is formed already at pH 3.4 (pK_a of both alginate and pectin being ~3.5), which strength increases with reduced pH (Fig. 1A), in agreement with previous studies.^{21,28} Ström and co-workers have further shown that the formation of a HM-pectin and alginate gel occurs within minutes²¹ once the pH is lowered close to the pK_a of alginate and pectin.

The gel will be subjected to forces such as shear and compression in the stomach, especially as it is pushed through the

Table 2 Characteristics such as gender, age, weight and height of MRI study participants

Magic ID	Gender	DOB	Age	Weight (kg)	Height (m)	BMI (kg m ⁻²)
1	M	1998-03-10	19	80	1.75	23
2	F	1987-01-02	30	58	1.58	18
3	F	1994-10-29	23	62	1.71	18
4	F	1995-09-13	22	53	1.64	16
5	F	1987-07-25	30	50	1.63	15
6	F	1995-10-23	22	61	1.63	19
7	F	1984-09-09	33	76	1.73	22
8	M	1997-04-30	20	74	1.77	21






Fig. 1 G' (triangle) and G'' (square) moduli of the test drink as a function of pH, determined at a strain of 0.5% and frequency of 1 Hz, all measurements performed at 37 °C (A) and a representative stress–strain curve for acidified test drink with a final pH of 2.1 (B).

antrum. The response of the gel to compression was therefore tested by forming cylindrical gels using a mould ($H = 10$ mm and $D = 10$ mm) in which the freshly prepared test drink plus GDL dispersion was poured and let to set for 24 hours. The cylinders were carefully removed after 48 hours and subjected to compression tests. The stress strain curves show a stress to fracture value of 7 ± 1 kPa, representing 0.5 N and a strain of fracture of $27 \pm 1.6\%$ (Fig. 1B). Such value of stress to fracture is just below the stated 10 kPa at force to fracture of gels previously shown to resist mechanical breakdown in the stomach,¹⁶ the formulation presented here should thus quickly pass on to the intestine.

The stability of the gel in gastric fluid and the release of low molecular weight (M_w) CHO is shown in Fig. 2. The test drink was added to a beaker containing simulated gastric juice, upon which a gel was formed instantaneously. The gel was stable *i.e.* no extensive shrinking or swelling occurred over the 60 minutes test in simulated gastric juice, contrary as was observed for calcium alginate beads.¹⁰ The release of CHO from the gel was fast, with CHO concentration outside the gel reaching 70% of C_∞ within ten minutes. In simulated intestinal juice the gel is expected to disintegrate, as pH of the gel increases to above the pK_a of the polysaccharides, thus deprotonating the polysaccharides leading to electrostatic repulsion and disintegration of the gel.

In general, the main driving forces for solute transport from gel matrices are related to the gradient in chemical potential, often expressed as the concentration difference of active solute between the gel matrix and the bulk according to Fick's law. Other factors that will impact the diffusion are the swelling or degradation and erosion of the matrix, which is not observed in simulated gastric juice.

The driving force for release of digestible CHO here is thus the gradient in chemical potential between the digestible CHO




Fig. 2 Cumulative release of low M_w CHO (filled circles) and average gel weight (open circles) as a function of time for the gel formed upon addition of the test drink to simulated gastric juice at $T = 37$ °C.

entrapped within the gel and the absence of digestible CHO in the simulated gastric juice.

Further, the voids and pores present in the HM pectin – alginate network are large (several 100 ds of nanometer), as observed using TEM (Fig. 3) and in agreement with previous studies on alginate HM-pectin gels²⁶ and calcium alginate.¹⁰ The polysaccharide network as visualised using TEM is corresponding to the black lines and dots. Keeping in mind that the size of the digestible CHO to be released, fructose with M_w of ~ 180 Da and maltodextrin with $M_w \sim 180$ –1500 Da, it is unlikely that the gel formed hinder the release of the CHO from the gel other than it is reducing coverage of the stomach wall as it is in its gelled state and not a solution.






Fig. 3 Transmission electron microscopy (TEM) images of gelled drink at two different magnifications (scale bar represents, from left to right, 1000 nm and 200 nm). White arrows indicate the presence of aggregates and thin strands.

It can be hypothesized from the physico-chemical characterisation of the formulation that upon ingestion of the test drink a gel will be formed in the acidic environment of the stomach, from which low- M_w CHO will be released *via* non hindered diffusion. The gel is however weak, suggesting little or no retention in the stomach. Once in the intestine, the increase of pH will force the gel to disintegrate owing to deprotonisation of the polysaccharides electrostatic repulsion.

3.2 Magnetic resonance imaging

The study was well tolerated by the participants and no adverse events were recorded. One of the subjects did not comply with the overnight fasting restrictions as their stomach showed the presence of food and liquid at the baseline scan. This participant was therefore excluded from the study.

It was possible to observe gelling of the test drink in the stomach of the seven remaining participants. The T_2 weighted images taken after ingestion of the test drink showed two dis-

tinct components in the stomach, one bright (consistent with a fluid component) and one darker (consistent with a gelled component). This is shown in Fig. 4 on the right hand panel. Conversely on the control drink study days the stomach contents were mostly bright (as seen in Fig. 4 on the left hand panel), with some artifacts probably due to flowing/moving fluid in the stomach. Looking at progressively longer echo times (*i.e.* images taken at different interval and collecting the signal later in time so that more of the signal from shorter time constants will have decayed) the fluid component remained brighter and did not change shape or appearance whilst the gelled component disappeared from the images, a clear sign that the darker component of the sports drink images had a much shorter T_2 than the brighter component. Intra-gastric gelling did not seem to be long-lived and in many subjects was not detectable by $T = 60$ min and beyond (Fig. 5).

Where apparent, separate regions of interest were drawn for the gel and fluid components visible in the stomach. If a




Fig. 4 T_2 weighted (TE = 300 ms) axial images of the stomach of one of the participants after they ingested the test drink on one study day (right hand panel) and the control drink on the other study day (left hand panel). The test drink image on the right shows two distinct components in the stomach, one bright (consistent with a fluid component) and one darker (consistent with a gelled component). Conversely on the left the stomach contents after the control drink are seen mostly bright, with some artifacts probably due to flowing/moving fluid in the stomach.






Fig. 5 The panel shows for corresponding axial images taken at different times (about 1 min apart from each other) from participant M7 on the test drink study day. Each image is from a similar location in the stomach but taken with progressively longer echo time TE, from 60 ms to 800 ms. At longer echo times most of the body organs and gel component have decayed (hence they appear black) leaving only bright form fluid water components.

single fluid component was visible, as in the case of the control drink, then one single region of interest was drawn. The signal decay sampled in the regions of interest was then fitted to a relaxation time model. Fluid values are more variable due to increased artifacts in the fluid regions, possibly due to the motion of the drink in the stomach reducing the signal intensity, so lower values (~ 0.5 s) could be underestimated. Most of the areas identified as gel seemed to have a T_2 around or below 0.2 s as shown in Fig. 6.

Measurement of blood glucose levels using the finger prick method over 120 min showed that control and test drinks gave rise to similar blood glucose levels (Fig. 7).

The MRI study confirmed the formation of a gel in the stomach 15 min after ingestion of the test drink, and the absence of gel in the stomach upon ingestion of the control drink. Furthermore, the study showed that no gel seemed to remain in the stomach at the second MRI scan (60 min later). It is worthwhile to note that none of the participants reported gastric distress or increased fullness upon ingestion of test drink in line with the hypothesis that the strength of the gel formed from the test drink used in this study is too weak to affect feeling of fullness or attenuate blood glucose levels, thus enabling efficient use of digestible CHO.

Considering that there seems to be a link between endurance performance, CHO ingestion rate and high exogenous CHO oxidation,^{29,30} sport drinks should be formulated to maximize CHO delivery without causing negative GI symptoms. The formulation tested here appears promising in this respect and randomized studies on exogenous CHO oxidation rates should be performed.




Fig. 6 Bar chart of the transverse relaxation time T_2 measured at the first imaging time point ($T = 0$) after ingestion in healthy participants who consumed 500 mL of test drink or control. Where apparent, separate regions of interest were drawn for the gel and fluid components visible in the stomach. If a single fluid component was visible as in the case of the control drink then one single region of interest was drawn. n indicates the number of subjects in whom a measurement was possible. The data are shown as mean \pm SD.




Fig. 7 Capillary blood glucose levels in participants of the double blind magnetic resonance imaging trial upon consumption of test and control drink.

4. Conclusions

We have shown that HM-pectin and alginate (0.2 wt%), in combination with digestible CHO (14 wt%), forms a weak gel under acid conditions, through which low- M_w CHO easily diffuses. MRI scanning confirms the presence of a gel *in vivo* in the stomach upon the first scan 15 minutes after ingestion of the test drink. Scanning the stomach 60 minutes after ingestion show that the gel is not retained, in line with the hypoth-



esis of the gel being weak enough to easily be emptied from the stomach. While a gel is present at early times in the stomach, the blood glucose level remains similar as for the control. No negative GI symptoms was observed for either of the test drink or the control despite their high content of digestible CHO. In order to gain more insight in the potential of polysaccharides to alleviate GI distress in conjunction with high-intensity exercise, further studies are needed, where conditions are more likely to provoke severe symptoms of gastric discomfort. Future studies should also involve a double-blind and randomized study on exogenous CHO oxidation.

Conflicts of interest

The study has been performed in collaboration with Maurten AB.

Acknowledgements

VINNOVA VINN Excellence Center, SuMo BIOMATERIALS, VINNMER and Innovationskontoret, Chalmers are acknowledged for financial support to A. S. and the study, Johan Bergenstahl for the use of the rheometer. Annika Altskär is gratefully acknowledged for TEM experiments.

References

- 1 S. N. Cheuvront, R. Carter 3rd and M. N. Sawka, Fluid balance and endurance exercise performance, *Curr. Sports Med. Rep.*, 2003, **2**, 202.
- 2 B. Essen, Intramuscular substrate utilization during prolonged exercise, *Ann. N. Y. Acad. Sci.*, 1977, **301**, 30.
- 3 M. N. Sawka, L. M. Burke, E. R. Eichner, R. J. Maughan, S. J. Montain and N. S. Stachenfeld, American College of Sports Medicine position stand: Exercise and fluid replacement, *Med. Sci. Sports Exercise*, 2007, **39**, 377.
- 4 X. Shi and C. V. Gisolfi, Fluid and carbohydrate replacement during intermittent exercise, *Sports Med.*, 1998, **25**, 157.
- 5 D. T. Thomas, K. A. Erdma and L. M. Burke, American college of sports medicine joint position statement. Nutrition and Athletic performance, *Med. Sci. Sports Exercise*, 2016, **48**, 543.
- 6 C. S. Almond, A. Y. Shin, E. B. Fortescue, R. C. Mannix, D. Wypij, B. A. Binstadt, C. N. Duncan, D. P. Olson, A. E. Salerno, J. W. Newburger and D. S. Greenes, Hyponatremia among runners in the Boston Marathon, *N. Engl. J. Med.*, 2005, **352**, 1550.
- 7 N. J. Rehrer, M. van Kemenade, W. Meester, F. Brouns and W. H. M. Saris, Gastrointestinal complaints in relation to dietary intake in triathletes, *Int. J. Sport Nutr.*, 1992, **2**, 48.
- 8 J. B. Leiper, K. P. Aulin and K. Soderlund, Improved gastric emptying rate in humans of a unique glucose polymer with gel-forming properties, *Scand. J. Gastroenterol.*, 2000, **35**, 1143.
- 9 P. Lopez-Sanchez, N. Fredriksson, A. Larsson, A. Altskär and A. Ström, High sugar content impacts microstructure, mechanics and release of calcium-alginate gels, *Food Hydrocolloids*, 2018, **84**, 26.
- 10 M. Ahnoff, O. Sköld and A. Ström, WO/2017/186940A1, 2017.
- 11 S. Sutehall, B. Muniz-Pardos, A. N. Bosch, A. Di Gianfrancesco and Y. P. Pitsiladis, Sports drinks on the edge of a new era, *Curr. Sports Med. Rep.*, 2018, **17**, 112.
- 12 L. Benini, G. Castellani, F. Brighenti, K. W. Heaton, M. T. Brentegani, M. C. Casiraghi, C. Sembenini, N. Pellegrini, A. Fioretta and G. Minniti, Gastric emptying of a solid meal is accelerated by the removal of dietary fibre naturally present in food, *Gut*, 1995, **36**, 825.
- 13 C. L. Hoad, P. Rayment, R. C. Spiller, L. Marciani, B. d. C. Alonso, C. Traynor, D. J. Mela, H. P. F. Peters and P. A. Gowland, In vivo imaging of intragastric gelation and its effect on satiety in humans, *J. Nutr.*, 2004, **134**, 2293.
- 14 M. Lyly, K.-H. Liukkonen, M. Salmenkallio-Marttila, L. Karhunen, K. Poutanen and L. Lähteenmäki, Fibre in beverages can enhance perceived satiety, *Eur. J. Nutr.*, 2009, **48**, 251.
- 15 K. R. Juvonen, A.-K. Purhonen, M. Salmenkallio-Marttila, L. Lähteenmäki, D. E. Laaksonen, K.-H. Herzig, M. I. J. Uusitupa, K. S. Poutanen and L. J. Karhunen, Viscosity of oat bran-enriched beverages influences gastrointestinal hormonal responses in healthy humans, *J. Nutr.*, 2009, **139**, 461.
- 16 L. Marciani, P. A. Gowland, A. Fillery-Travis, P. Manoj, J. Wright, A. Smith, P. Young, R. Moore and R. C. Spiller, Assessment of antral grinding of a model solid meal with echo-planar imaging, *Am. J. Physiol.: Gastrointest. Liver Physiol.*, 2001, **280**, G844.
- 17 I. Torsdottir, M. Alpsten, G. Holm, A. S. Sandberg and J. Tolli, A small dose of soluble alginate-fiber affects postprandial glycemia and gastric emptying in humans with diabetes, *J. Nutr.*, 1991, **121**, 795.
- 18 D. El Khoury, H. D. Goff, S. Berengut, R. Kubant and G. H. Anderson, Effect of sodium alginate addition to chocolate milk on glycemia, insulin, appetite and food intake in healthy adult men, *Eur. J. Clin. Nutr.*, 2014, **68**, 613.
- 19 R. D. Mattes, Effects of a combination fiber system on appetite and energy intake in overweight humans, *Physiol. Behav.*, 2007, **90**, 705.
- 20 N. C. Howarth, E. Saltzman and S. B. Roberts, Dietary fiber and weight regulation, *Nutr. Rev.*, 2001, **59**, 129.
- 21 A. Ström, R. Koppert, H. M. Boers, S. M. Melnikov, S. Wiseman and H. P. F. Peters, *Physico-chemical properties of polysaccharides determines their appetite effect in Gums and Stabilisers for the food industry 15*, Royal Society of Chemistry (RSC), 2010.
- 22 S. T. Odunsi, M. I. Vázquez-Roque, M. Camilleri, A. Papathanasopoulos, M. M. Clark, L. Wodrich, M. Lempke, S. McKinzie, M. Ryks, D. Burton and A. R. Zinsmeister, Effects of alginate on satiation, appetite,



- gastric function and selected gut satiety hormones in overweight and obesity, *Obesity*, 2010, **18**, 1579.
- 23 K. Murray, V. Wilkinson-Smith, C. Hoad, C. Costigan, E. Cox, C. Lam, L. Marciani, P. Gowland and R. C. Spiller, Differential effects of FODMAPs (fermentable oligo-, di-, mono-saccharides and polyols) on small and large intestinal contents in healthy subjects shown by MRI, *Am. J. Gastroenterol.*, 2014, **109**, 110.
- 24 G. Major, S. Pritchard, K. Murray, J. P. Alappadan, C. L. Hoad, L. Marciani, P. Gowland and R. Spiller, Colon hypersensitivity to distension, rather than excessive gas production, produces carbohydrate-related symptoms in individuals with irritable bowel syndrome, *Gastroenterology*, 2017, **152**, 124.
- 25 C. L. Hoad, E. F. Cox and P. A. Gowland, Quantification of T(2) in the abdomen at 3.0 T using a T(2)-prepared balanced turbo field echo sequence, *Magn. Reson. Med.*, 2010, **63**, 356.
- 26 P. Walkenström, S. Kidman, A.-M. Hermansson, P. B. Rasmussen and L. Hoegh, Microstructure and rheological behaviour of alginate/pectin mixed gels, *Food Hydrocolloids*, 2003, **17**, 593.
- 27 P. B. Wilson, Multiple transportable carbohydrates during exercise: Current limitations and directions for future research, *J. Strength Cond. Res.*, 2015, **29**, 2056.
- 28 V. J. Morris and G. R. Chilvers, Cold setting alginate-pectin mixed gels, *J. Sci. Food Agric.*, 1984, **35**, 1370.
- 29 K. Currell and A. E. Jeukendrup, Superior endurance performance with ingestion of multiple transportable carbohydrates, *Med. Sci. Sports Exercise*, 2008, **40**, 275.
- 30 D. Triplett, J. A. Doyle, J. C. Rupp and D. Benardot, An isocaloric glucose-fructose beverage's effect on simulated 100-km cycling performance compared with a glucose-only beverage, *Int. J. Sport Nutr. Exercise Metab.*, 2010, **20**, 122.

