



q-Independence of the Jimbo-Drinfeld Quantization

Downloaded from: <https://research.chalmers.se>, 2025-12-04 23:28 UTC

Citation for the original published paper (version of record):

Giselsson, O. (2020). q-Independence of the Jimbo-Drinfeld Quantization. *Communications in Mathematical Physics*, 376(3): 1737-1765. <http://dx.doi.org/10.1007/s00220-019-03660-9>

N.B. When citing this work, cite the original published paper.



q -Independence of the Jimbo–Drinfeld Quantization

Olof Giselsson 

Chalmers tvärgata 3, Göteborg, Sweden. E-mail: olofgi@chalmers.se

Received: 31 January 2019 / Accepted: 26 July 2019
Published online: 7 January 2020 – © The Author(s) 2020

Abstract: Let G be a connected semi-simple compact Lie group and for $0 < q < 1$, let $(\mathbb{C}[G]_q, \Delta_q)$ be the Jimbo–Drinfeld q -deformation of G . We show that the C^* -completions of $\mathbb{C}[G]_q$ are isomorphic for all values of q . Moreover, these isomorphisms are equivariant with respect to the right-actions of the maximal torus.

1. Introduction

The quantized universal enveloping algebra $U_q(\mathfrak{g})$ of a semi-simple Lie algebra \mathfrak{g} was introduced by Drinfeld and Jimbo in the mid-80's [3, 5]. In [4], Drinfeld also introduced their dual objects, deformations $\mathbb{C}[G]_q$ of the Hopf algebra of regular functions on a semi-simple Lie group G . Moreover, when G is compact, the algebra $\mathbb{C}[G]_q$ can be given the structure of a Hopf $*$ -algebra. In this case one can see that the enveloping C^* -algebra of $\mathbb{C}[G]_q$ exists, giving a natural q -analogue $C(G)_q$ of the algebra of continuous functions on G . The analytic approach to quantum groups was initially proposed by Woronowicz [14]. In the 90's Soibelman gave a complete classification of the irreducible $*$ -representations of $\mathbb{C}[G]_q$. These were shown to be in one-to-one correspondence with the symplectic leaves of G coming from the Poisson structure on $C(G)$ determined by the quantization when $q \rightarrow 1$. However, it was not clear how the C^* -algebraic structure of $C(G)_q$ was depending on the parameter q . In fact, several evidence pointed towards that the structure was actually independent of it. In the special case of SU_2 , it was observed (see [14]) that the C^* -algebras $C(SU_2)_q$, $q \in (0, 1)$ are all isomorphic. In the mid 90's, G. Nagy showed in [8] that the same holds for $C(SU_3)_q$. Moreover, it was also shown by Nagy (in [9]) that $C(SU_n)_q$ is KK-equivalent to $C(SU_n)_s$, for all $n \in \mathbb{N}$ and all $q, s \in (0, 1)$. This was extended by Neshveyev–Tuset in [10] to yield a KK-equivalence between $C(G)_q$ and $C(G)_s$ for any compact simply connected semi-simple Lie group G . In this paper, we show that some of the ideas that underpin Nagy's proof of the q -independence of $C(SU_3)_q$ can be extended to give the following result: for a fixed symplectic leaf $U \subseteq G$, with corresponding $*$ -representations π^q of $\mathbb{C}[G]_q$, we

have an isomorphism $\overline{\text{Im}\pi^q} \cong \overline{\text{Im}\pi^s}$ for all $q, s \in (0, 1)$. Using this, we prove that $C(G)_q \cong C(G)_s$, thus showing that these non-isomorphic compact quantum groups are all isomorphic as C^* -algebras. This confirms the conjecture made in [8].

The paper is organized as follows. We finish this section by giving some geometric intuitions underlying the proof and describing the idea of the proof using this geometric picture. We then present and prove the lifting theorem used in the proof of the main result. Section 2 goes through the formal definitions of $\mathbb{C}[G]_q$ as well as its representation theory. In Sect. 3, we prove some more specific results regarding representations and how these depend on the q parameter. In Sect. 4, we state and prove the main result.

1.1. Outline of the proof. To explain the main ideas of the proof, it is worth to start by considering the case of $G = \text{SU}_3$, previously covered by Nagy. There is an irreducible $*$ -representation $\pi^q : \mathbb{C}[\text{SU}_3]_q \rightarrow \mathcal{B}(\ell^2(\mathbb{Z}_+)^{\otimes 3})$ that, in Nagy's words, corresponds to the "big symplectic leaf" in SU_3 . However, there is an inherent problem when trying to determine if $\overline{\text{Im}\pi^q} \cong \overline{\text{Im}\pi^s}$ for different $q, s \in (0, 1)$. As π^q can be seen to vary (in a certain sense) continuously on q , one intuitive approach could be to let $q \rightarrow 0$, and then find some natural set of generators such that an isomorphic set of generators can be found in $\overline{\text{Im}\pi^q}$ for each $q \in (0, 1)$. This method works, for example, in the case of $C(\text{SU}_2)_q$. However, for $\mathbb{C}[\text{SU}_3]_q$, there seems to be no simple way of taking the limit $q \rightarrow 0$. As it sits, the image of π^q is simply too "twisted" to allow passing to any limit.

These problems are resolved in the following way: We observe that $\overline{\text{Im}\pi^q}$, for all $q \in (0, 1)$, contains the compact operators $\mathcal{K} \subseteq \mathcal{B}(\ell^2(\mathbb{Z}_+)^{\otimes 3})$, and moreover, the intersection $\text{Im}\pi^q \cap \mathcal{K}$ is non-trivial. The compact operators form a minimal ideal in $\overline{\text{Im}\pi^q}$, in the sense that it is contained in any other ideal. We now consider the composition

$$\mathbb{C}[\text{SU}_3]_q \xrightarrow{\pi^q} \mathcal{B}(\ell^2(\mathbb{Z}_+)^{\otimes 3}) \xrightarrow{p} \mathcal{B}(\ell^2(\mathbb{Z}_+)^{\otimes 3})/\mathcal{K} \cong \mathcal{Q}(\ell^2(\mathbb{Z}_+)^{\otimes 3}),$$

where p is the quotient map $x \mapsto x + \mathcal{K}$, and we then proceed by analyzing $p \circ \pi^q$. It is clear that π^q can not be a direct summand in $p \circ \pi^q$, as the elements mapped by π^q into $\text{Im}\pi^q \cap \mathcal{K}$ is now mapped into zero. It turns out that there are two Hilbert spaces H_1, H_2 , such that for every $q \in (0, 1)$, we have $*$ -representations

$$\mathbb{C}[\text{SU}_3]_q \xrightarrow{\Pi_1^q} \mathcal{B}(H_1), \mathbb{C}[\text{SU}_3]_q \xrightarrow{\Pi_2^q} \mathcal{B}(H_2)$$

and an isomorphism $\varphi_q : \overline{\text{Im}\pi^q}/\mathcal{K} \rightarrow \overline{\text{Im}(\Pi_1^q \oplus \Pi_2^q)}$, such that for every $a \in \mathbb{C}[\text{SU}_3]_q$, we have $\varphi_q(p \circ \pi^q(a)) = (\Pi_1^q \oplus \Pi_2^q)(a)$. Moreover, in this case, one can show that actually we have $\text{Im}(\Pi_1^q \oplus \Pi_2^q) = \overline{\text{Im}(\Pi_1^s \oplus \Pi_2^s)}$ for all $q, s \in (0, 1)$ as subspaces of $\mathcal{B}(H_1) \oplus \mathcal{B}(H_2)$. Thus, by quoting out the compact operators, we have successfully "untwisted" the $*$ -representation π^q . Letting $M = \overline{\text{Im}(\Pi_1^q \oplus \Pi_2^q)}$, one then shows that the injective homomorphisms $\varphi_q^{-1} : M \rightarrow \mathcal{Q}(\ell^2(\mathbb{Z}_+)^{\otimes 3})$ varies norm-continuously on q , meaning that, for a fixed $x \in M$, the map $q \in (0, 1) \rightarrow \varphi_q^{-1}(x)$ is a continuous function of q . We then get an isomorphism

$$p^{-1}(\varphi_q^{-1}(M)) \cong p^{-1}(\varphi_s^{-1}(M)), \quad q, s \in (0, 1) \quad (1)$$

using the lifting result (Lemma 1 below). As $\mathcal{K} \subseteq \overline{\text{Im}\pi^q}$, it follows that $p^{-1}(\varphi_q^{-1}(M)) = \overline{\text{Im}\pi^q}$ and hence we have established an isomorphism for different q . One needs further

argument to conclude that actually $C(\mathrm{SU}_3)_q \cong C(\mathrm{SU}_3)_s$, but by proving (1), the main effort is done. The proof below essentially systematizes this line of argument in a way that makes it also work for a general G , by quoting out ideals to "untwist" an irreducible $*$ -representation π^q of $C(G)_q$ further and further, until it is clear that the images of the resulting $*$ -representations are independent of q . Then one uses inductive arguments to check that also $\mathrm{Im}\pi^q \cong \mathrm{Im}\pi^s$ for $q, s \in (0, 1)$.

One can give a quite clear geometric heuristic of this, using the one-to-one correspondence between irreducible $*$ -representations of $\mathbb{C}[G]_q$ and symplectic leaves in G coming from the corresponding Poisson structure on $C(G)$ (see [7]). Recall that G can be decomposed into a disjoint union of symplectic leaves and that each leaf is an even-dimensional sub-manifold of G . Let U be a $2m$ -dimensional symplectic leaf of G , corresponding to a $*$ -representation π^q of $\mathbb{C}[G]_q$, and let $C_0(U)$ be the ideal of $C(\overline{U})$ of all continuous functions vanishing on $\overline{U} \setminus U$. Thus quoting out this ideal gives a homomorphism $C(\overline{U}) \rightarrow C(\overline{U} \setminus U)$. It turns out that $\overline{U} \setminus U$ can be written as a disjoint union $\cup_j U_j$, of symplectic leaves of dimension strictly less than $2m$, and that the leaves in this union of dimension $< 2m - 2$ are contained in the closures of the leaves of dimension $2m - 2$. In general, we can write \overline{U} as a disjoint union of symplectic leaves

$$\overline{U} = U \cup \left(\cup_j U_j^{(m-1)} \right) \cup \left(\cup_j U_j^{(m-2)} \right) \cup \dots \cup \left(\cup_j U_j^{(0)} \right)$$

such that each $U_j^{(k)}$ is a symplectic leaf of dimension $2k$ and

$$\cup_j \overline{U}_j^{(k)} = \left(\cup_j U_j^{(k)} \right) \cup \dots \cup \left(\cup_j U_j^{(0)} \right). \quad (2)$$

This shows that we can make a sequence of homomorphisms

$$C(\overline{U}) \longrightarrow \prod_j C(\overline{U}_j^{(m-1)}) \longrightarrow \dots \longrightarrow \prod_j C(\overline{U}_j^{(1)}) \longrightarrow \prod_j C(\overline{U}_j^{(0)}) \quad (3)$$

such that on each step, the homomorphism $\prod_j C(\overline{U}_j^{(k)}) \rightarrow \prod_j C(\overline{U}_j^{(k-1)})$ has kernel $\prod_j C_0(U_j^{(k)})$. Let us explain how a q -analogue of (3) is used.

Remark 1. For several reasons, the notations used here will differ somewhat from the ones used later in the text.

Let U and $U_j^{(k)}$ be as above. We can think of $\overline{\mathrm{Im}\pi^q}$ and the ideal $\mathcal{K} \subseteq \overline{\mathrm{Im}\pi^q}$ as q -analogs of $C(\overline{U})$ and $C_0(U)$, denoted by $C(\overline{U})_q$ and $C_0(U)_q$ respectively. There is then a sequence of homomorphisms

$$C(\overline{U})_q \xrightarrow{\partial_m^q} \prod_j C(\overline{U}_j^{(m-1)})_q \xrightarrow{\partial_{m-1}^q} \dots \xrightarrow{\partial_2^q} \prod_j C(\overline{U}_j^{(1)})_q \xrightarrow{\partial_1^q} \prod_j C(\overline{U}_j^{(0)})_q \quad (4)$$

such that on each step, the homomorphism $\partial_k : \prod_j C(\overline{U}_j^{(k)})_q \rightarrow \prod_j C(\overline{U}_j^{(k-1)})_q$ has kernel equal to $\prod_j C_0(U_j^{(k)})_q$. Let us denote by $C(\partial^{(k)}\overline{U})_q$, $k = 0, \dots, m-1$, the image of $C(\overline{U})_q$ in $\prod_j C(\overline{U}_j^{(k)})_q$ via the composition of homomorphisms in (4). The idea is to proceed by induction on the dimensions of the symplectic leaves. In the case of zero-dimensional leaves, the corresponding $*$ -representations are one-dimensional (maps to

©) and hence trivially q -independent. For higher dimensional leaves, we can use the induction hypothesis to connect the lower dimensional leaves for different $q, s \in (0, 1)$

$$\begin{array}{ccccccc}
 C(\overline{U})_q & \xrightarrow{\partial_m^q} & \prod_j C(\overline{U}_j^{(m-1)})_q & \xrightarrow{\partial_{m-1}^q} \cdots \xrightarrow{\partial_2^q} & \prod_j C(\overline{U}_j^{(1)})_q & \xrightarrow{\partial_1^q} & \prod_j C(\overline{U}_j^{(0)})_q \\
 \downarrow & & \downarrow \Gamma_{m-1}^{s,q} & & \downarrow \Gamma_1^{s,q} & & \downarrow \Gamma_0^{s,q} \\
 C(\overline{U})_s & \xrightarrow{\partial_m^s} & \prod_j C(\overline{U}_j^{(m-1)})_s & \xrightarrow{\partial_{m-1}^s} \cdots \xrightarrow{\partial_2^s} & \prod_j C(\overline{U}_j^{(1)})_s & \xrightarrow{\partial_1^s} & \prod_j C(\overline{U}_j^{(0)})_s
 \end{array} \quad (5)$$

Moreover, this can be done in a way such that the diagram (5) is commutative. The aim is then to construct a dotted arrow from $C(\overline{U})_q$ to $C(\overline{U})_s$ that makes the diagram commutative. The main obstacle to do this is to check that $C(\partial^{(m-1)}\overline{U})_q$ is mapped by $\Gamma_{m-1}^{s,q}$ to $C(\partial^{(m-1)}\overline{U})_s$. In order to prove this, one shows the following

- (i) for $k = 0, \dots, m-1$ the intersection of $C(\partial^{(k)}\overline{U})_q$ with $\prod C_0(U_j^{(k)})_q$ is mapped by $\Gamma_k^{s,q}$ to the intersection of $C(\partial^{(k)}\overline{U})_s$ with $\prod C_0(U_j^{(k)})_s$,
- (ii) the C^* -algebras $C(\partial^{(0)}\overline{U})_q$ are commutative and isomorphic for all $q \in (0, 1)$, via $\Gamma_0^{s,q}$,
- (iii) for $k = 1, \dots, m-1$, there is an approximate unit $\{u_{q,i}^{(k)}\}_{i=1}^\infty$ for $\prod_j C_0(U_j^{(k)})_q$ such that $\{u_{q,i}^{(k)}\}_{i=1}^\infty \subseteq C(\partial^{(k)}\overline{U})_q$ and $\Gamma_k^{s,q}(u_{q,i}^{(k)}) = u_{s,i}^{(k)}$ for all $i \in \mathbb{N}$.

Remark 2. In the actual proof below, we do not really use (iii), since by the way the arguments are constructed there, explicitly stating this point becomes unnecessary.

From the commutivity of the square

$$\begin{array}{ccc}
 \prod_j C(\overline{U}_j^{(k)})_q & \xrightarrow{\partial_k^q} & \prod_j C(\overline{U}_j^{(k-1)})_q \\
 \Gamma_k^{s,q} \downarrow & & \downarrow \Gamma_{k-1}^{s,q} \\
 \prod_j C(\overline{U}_j^{(k)})_s & \xrightarrow{\partial_k^s} & \prod_j C(\overline{U}_j^{(k-1)})_s
 \end{array} \quad (6)$$

it follows that if $\Gamma_{k-1}^{s,q}$ restricts to a $*$ -isomorphism from $C(\partial^{(k-1)}\overline{U})_q$ to $C(\partial^{(k-1)}\overline{U})_s$, then as for any $x \in C(\partial^{(k)}\overline{U})_q$, we have

$$\partial_k^s(\Gamma_k^{s,q}(x)) = \Gamma_{k-1}^{s,q}(\partial_k^q(x)) \in C(\partial^{(k-1)}\overline{U})_s,$$

it follows that $\Gamma_k^{s,q}(x) = y + c$ where $y \in C(\partial^{(k)}\overline{U})_s$ and $c \in \prod_j C_0(U_j^{(k)})_q$. By (iii), we have an approximate unit $\{u_{q,i}^{(k)}\}_{i=1}^\infty$ such that $xu_{q,i}^{(k)}$ is in the intersection of $C(\partial^{(k)}\overline{U})_q$ with $\prod_j C_0(U_j^{(k)})_q$. It now follows from (i) that for all $i \in \mathbb{N}$, we have

$$\begin{aligned}
 \Gamma_k^{s,q}(xu_{q,i}^{(k)}) &\in C(\partial^{(k)}\overline{U})_s, \\
 y\Gamma_k^{s,q}(u_{q,i}^{(k)}) &\in C(\partial^{(k)}\overline{U})_s
 \end{aligned}$$

and thus it follows that also $cu_{s,i}^{(k)} \in C(\partial^{(k)}\overline{U})_s$. Letting $i \rightarrow \infty$ now gives $c \in C(\partial^{(k)}\overline{U})_s$ and thus $\Gamma_k^{s,q}(x) \in C(\partial^{(k)}\overline{U})_s$. Using (ii), it now follows by induction that $C(\partial^{(m-1)}\overline{U})_q$ is mapped isomorphically onto $C(\partial^{(m-1)}\overline{U})_s$. As $\ker \partial_m^q = C_0(U)_q$, this is equivalent to

$$C(\overline{U})_q/C_0(U)_q \cong C(\overline{U})_s/C_0(U)_s, \quad q, s \in (0, 1).$$

After checking that these isomorphisms are varying norm-continuously as functions of q and s , the dotted arrow in (5) can then be constructed using Lemma 2 below.

1.2. Lifting results.

Lemma 1 (Lemma 2 in [8]). *Let H be a separable Hilbert space, let \mathcal{K} be the space of compact operators on H , let $\mathbb{Q}(H) = \mathcal{B}(H)/\mathcal{K}$ be the Calkin algebra and $p : \mathcal{B}(H) \rightarrow \mathbb{Q}(H)$ the quotient map. Suppose A is a fixed separable C^* -algebra of type I and $\phi_q : A \rightarrow \mathbb{Q}(H)$, $q \in [0, 1]$ is a point-norm continuous family of injective $*$ -homomorphisms. Denote*

$$\begin{aligned} \mathfrak{A}_q &:= \phi_q(A) : \\ M_q &:= p^{-1}(\mathfrak{A}_q). \end{aligned}$$

Then there exists a family of injective $$ -homomorphisms $\Phi_q : M_0 \rightarrow \mathcal{B}(H)$, $q \in [0, 1]$ with the following properties*

- (a) $\Phi_q(M_0) = M_q$ for $q \in [0, 1]$ and $\Phi_0 = \text{Id}_{M_0}$,
- (b) the family $\Phi_q : M_0 \rightarrow \mathcal{B}(H)$, $q \in [0, 1]$ is point-norm continuous,
- (c) for every $q \in [0, 1]$, the diagram

$$\begin{array}{ccc} M_0 & \xrightarrow{\Phi_q} & M_q \\ p \downarrow & & \downarrow p \\ \mathfrak{A}_0 & \xrightarrow{\phi_q \circ \phi_0^{-1}} & \mathfrak{A}_q \end{array} \quad (7)$$

is commutative.

We remind the reader that a C^* -algebra of type I, is one where the image of every irreducible $*$ -representation includes a non-zero compact operator. Here, we will use a modified version of Lemma 1.

Lemma 2. *Let H be a separable Hilbert space, let \mathcal{K} be the space of compact operators on H , let $\mathbb{Q}(H) = \mathcal{B}(H)/\mathcal{K}$ be the Calkin algebra and $p : \mathcal{B}(H) \rightarrow \mathbb{Q}(H)$ be the quotient map. For every $q \in (0, 1)$, suppose $A_q \subseteq \mathbb{Q}(H)$ is a separable C^* -algebra of type I and we have a family of $*$ -isomorphisms $\phi_{s,q} : A_q \rightarrow A_s$, $s, q \in (0, 1)$ which are continuous in the point-norm topology (i.e. for every fixed $q \in (0, 1)$ and $x \in A_q$, the map $s \in (0, 1) \mapsto \phi_{s,q}(x) \in \mathbb{Q}(H)$ is norm-continuous). Assume moreover that $\phi_{q,q} = \text{Id}_{A_q}$ and $\phi_{t,s} \circ \phi_{s,q} = \phi_{t,q}$ for all $t, s, q \in (0, 1)$. Denote*

$$B_q := p^{-1}(A_q).$$

Then there exists a family of inner $$ -isomorphisms $\Phi_{s,q} : \mathcal{B}(H) \rightarrow \mathcal{B}(H)$, $s, q \in (0, 1)$ (i.e. $\Phi_{s,q}(x) = U_{s,q}^* x U_{s,q}$ for some unitary $U_{s,q} \in \mathcal{B}(H)$) with the following properties*

- (i) $\Phi_{s,q}(\mathbf{B}_q) = \mathbf{B}_s$ for $s, q \in (0, 1)$,
- (ii) $\Phi_{q,q} = \text{Id}$ and $\Phi_{t,s} \circ \Phi_{s,q} = \Phi_{t,q}$ for all $t, s, q \in (0, 1)$,
- (iii) for fix q , the family $\Phi_{s,q} : \mathbf{B}_s \rightarrow \mathcal{B}(H)$, $s \in [0, 1]$ is point-norm continuous,
- (iv) for all $s, q \in (0, 1)$, the diagram

$$\begin{array}{ccc} \mathbf{B}_q & \xrightarrow{\Phi_{s,q}} & \mathbf{B}_s \\ p \downarrow & & \downarrow p \\ \mathbf{A}_q & \xrightarrow{\phi_{s,q}} & \mathbf{A}_s \end{array} \quad (8)$$

is commutative.

Proof. If $a < b$, then clearly the conclusion of Lemma 1 still holds if we change the interval to $[a, b]$. Let $a_k \in (0, 1)$, $k \in \mathbb{Z}$ be a strictly increasing sequence such that $a_k \rightarrow 1$ and $a_{-k} \rightarrow 0$ as $k \rightarrow \infty$. For $k \geq 0$, we apply Lemma 1 to the set of injective $*$ -homomorphisms

$$\tilde{\phi}_{q,k} := \phi_{q,a_k} : \mathbf{A}_{a_k} \rightarrow \mathcal{Q}(H), \quad q \in [a_k, a_{k+1}],$$

and let $M_{q,k} = p^{-1}(\tilde{\phi}_{q,k}(\mathbf{A}_{a_k})) = p^{-1}(\mathbf{A}_q)$. For $k < 0$, we instead apply Lemma 1 to

$$\tilde{\phi}_{q,k} := \phi_{q,a_{k+1}} : \mathbf{A}_{a_{k+1}} \rightarrow \mathcal{Q}(H), \quad q \in [a_k, a_{k+1}].$$

Let $\tilde{\Phi}_{q,k}$, $k \in \mathbb{Z}$, be the $*$ -isomorphisms aquired by applying Lemma 1. Note that $\tilde{\Phi}_{q,k}$ is an isomorphism from \mathbf{B}_{a_k} to \mathbf{B}_q for $q \in [a_k, a_{k+1}]$ and $k \geq 0$ and an isomorphism from $\mathbf{B}_{a_{k+1}}$ to \mathbf{B}_q if $k < 0$. Let us define a $*$ -isomorphism $\Phi_q : \mathbf{B}_{a_0} \rightarrow \mathbf{B}_q$ by the formula

$$\Phi_q = \begin{cases} \tilde{\Phi}_{q,k} \circ \tilde{\Phi}_{a_k,k-1} \circ \cdots \circ \tilde{\Phi}_{a_1,0}, & \text{if } q \in [a_k, a_{k+1}] \text{ and } k \geq 0 \\ \tilde{\Phi}_{q,k} \circ \tilde{\Phi}_{a_k,k+1} \circ \cdots \circ \tilde{\Phi}_{a_{-1},-1}, & \text{if } q \in [a_k, a_{k+1}] \text{ and } k < 0 \end{cases} \quad (9)$$

It follows from Lemma 1 and the construction of Φ_q that for $x \in \mathbf{B}_{a_0}$, the map $q \in (0, 1) \mapsto \Phi_q(x) \in \mathcal{B}(H)$ is norm-continuous. That $p \circ \Phi_q = \phi_{q,a_0} \circ p$ holds follows from $\phi_{t,s} \circ \phi_{s,q} = \phi_{t,q}$ and iteration of the commutative diagram

$$\begin{array}{ccccc} \mathbf{B}_{a_0} & \xrightarrow{\tilde{\Phi}_{a_1,0}} & \mathbf{B}_{a_1} & \xrightarrow{\tilde{\Phi}_{q,1}} & \mathbf{B}_q \\ p \downarrow & & p \downarrow & & \downarrow p \\ \mathbf{A}_{a_0} & \xrightarrow{\phi_{a_1,a_0}} & \mathbf{A}_{a_1} & \xrightarrow{\phi_{q,a_1}} & \mathbf{A}_q \end{array}$$

We then let $\Phi_{s,q} = \Phi_s \circ \Phi_q^{-1}$ for $s, q \in (0, 1)$. By (7), we have a commutative diagram

$$\begin{array}{ccccc} \mathbf{B}_q & \xrightarrow{\Phi_q^{-1}} & \mathbf{B}_{a_0} & \xrightarrow{\Phi_s} & \mathbf{B}_s \\ p \downarrow & & p \downarrow & & \downarrow p \\ \mathbf{A}_q & \xrightarrow{\phi_{q,a_0}^{-1}} & \mathbf{A}_{a_0} & \xrightarrow{\phi_{s,a_0}} & \mathbf{A}_s \end{array}$$

From this, we get the commutivity of (8), as

$$\phi_{s,a_0} \circ \phi_{q,a_0}^{-1} = \phi_{s,a_0} \circ \phi_{a_0,q} = \phi_{s,q}.$$

Hence, (i), (ii), (iii), and (iv) holds for $\Phi_{s,q}$ as a $*$ -isomorphism $B_q \rightarrow B_s$.

We now extend $\Phi_{s,q}$ as an inner automorphism of all of $\mathcal{B}(H)$. To do this, we first show that the restriction $\Phi_{s,q}|_{\mathcal{K}}$ is inner. We have $\mathcal{K} \subseteq B_q$, and we get from the diagram (8) that $\Phi_{s,q}(\mathcal{K}) \subseteq \mathcal{K}$, and as $\Phi_{s,q}^{-1} = \Phi_{q,s}$ it follows that actually

$$\Phi_{s,q}(\mathcal{K}) = \mathcal{K}. \quad (10)$$

So $\Phi_{s,q}$ is an irreducible representation of \mathcal{K} . It is known that any such is unitarily equivalent to the identity representation (e.g. see Corollary 1.10 in [2]). Hence, there exists a unitary $U_{s,q} \in \mathcal{B}(H)$ such that $\Phi_{s,q}(x) = U_{s,q}xU_{s,q}^*$ for all $x \in \mathcal{K}$. For arbitrary $y \in B_q$, we obtain

$$\Phi_{s,q}(x)\Phi_{s,q}(y) = \Phi_{s,q}(xy) = U_{s,q}xyU_{s,q}^* = \Phi_{s,q}(x)U_{s,q}yU_{s,q}^*$$

for all $x \in \mathcal{K}$. This gives $\Phi_{s,q}(y) = U_{s,q}yU_{s,q}^*$. \square

2. Preliminaries

In this section we recall some facts about the Hopf algebras $U_q(\mathfrak{g})$ and $\mathbb{C}[G]_q$. The presentation is mainly taken from [10]. A general reference for the technical claims made here is [7].

2.1. The quantum group $U_q(\mathfrak{g})$. Let G be a simply connected semisimple compact Lie group and let \mathfrak{g} denote its complexified Lie algebra. We write $U(\mathfrak{g})$ to denote the universal enveloping algebra of \mathfrak{g} equipped with a $*$ -involution induced by the real form derived from G . Moreover, we let $\mathfrak{h} \subseteq \mathfrak{g}$ be the Cartan sub-algebra coming from a maximal torus $T \subseteq G$. Let $\mathfrak{t} \subseteq \mathfrak{h}$ be the real subspace of skew-symmetric elements. Write Φ for the set of roots of \mathfrak{g} , Φ_+ for the set of positive roots and $\Omega = \{\alpha_1, \dots, \alpha_n\}$ for the set of simple roots.

We denote the Weyl group of \mathfrak{g} by W and identify its set of generators $s_i, i = 1, \dots, n$, (as a Coxeter group) with Ω by the identification $\alpha_i \mapsto s_{\alpha_i} =: s_i$. Moreover, we identify Φ_+ with the set $\{ws_iw^{-1} : s_i \in \Omega, w \in W\} \subseteq W$. In both instances, we write the identification as $\alpha \mapsto h_\alpha \in \mathfrak{t}$.

Let $q \in (0, 1)$. Let $(a_{ij})_{ij}$ be the Cartan matrix of \mathfrak{g} and $d_i = \frac{(\alpha_i, \alpha_i)}{2}$ for $i = 1, \dots, n$. Let $q_i = q^{d_i}$ for $i = 1, \dots, n$. The quantized universal enveloping algebra $U_q(\mathfrak{g})$ is the unital complex algebra generated by elements $E_i, F_i, K_i, K_i^{-1}, i = 1, \dots, n$ subject to the relations

$$K_i K_i^{-1} = K_i^{-1} K_i = 1, K_i K_j = K_j K_i, K_i E_j = q_i^{a_{ij}} E_j K_i, K_i F_j = q_i^{a_{ij}} F_j K_i,$$

$$E_i F_j - F_j E_i = \delta_{ij} \frac{K_i - K_i^{-1}}{q_i - q_i^{-1}}$$

$$\sum_{k=0}^{1-a_{ij}} (-1)^k \begin{bmatrix} 1-a_{ij} \\ k \end{bmatrix}_{q_i} E_i^k E_j E_i^{1-a_{ij}-k} = 0,$$

$$\sum_{k=0}^{1-a_{ij}} (-1)^k \begin{bmatrix} 1-a_{ij} \\ k \end{bmatrix}_{q_i} F_i^k F_j F_i^{1-a_{ij}-k} = 0$$

where $\begin{bmatrix} k \\ j \end{bmatrix}_{q_i} = \prod_{m=0}^{j-1} \frac{q_i^{-(k-m)} - q_i^{k-m}}{q_i^{-(m+1)} - q_i^{m+1}}$ is a q -analogue of the binomial coefficients.

$U_q(\mathfrak{g})$ becomes a Hopf $*$ -algebra when equipped with a co-associative co-product $\hat{\Delta}_q$, a co-unit $\hat{\epsilon}_q$, an antipode \hat{S}_q and a $*$ -involution given on generators as

$$\begin{aligned}\hat{\Delta}_q(K_i) &= K_i \otimes K_i, \quad \hat{\Delta}_q(E_i) = E_i \otimes 1 + K_i \otimes E_i, \quad \hat{\Delta}_q(F_i) = F_i \otimes K_i^{-1} + 1 \otimes F_i, \\ \hat{\epsilon}_q(E_i) &= \hat{\epsilon}_q(F_i) = 0, \quad \hat{\epsilon}_q(K_i) = 1, \\ \hat{S}_q(F_i) &= -F_i K_i, \quad \hat{S}_q(E_i) = -K_i^{-1} E_i, \quad \hat{S}_q(K_i) = K_i^{-1} \\ K_i^* &= K_i, \quad E_i^* = F_i K_i, \quad F_i^* = K_i^{-1} E_i.\end{aligned}$$

We denote the antipode of $U_q(\mathfrak{g})$ by \hat{S}_q . For $q = 1$, we let $U_1(\mathfrak{g}) = U(\mathfrak{g})$ be the ordinary universal enveloping algebra with the usual Hopf $*$ -algebra structure and denote the co-product, co-unit, antipode simply by $\hat{\Delta}$, $\hat{\epsilon}$, \hat{S} .

Let P be the set of weights for \mathfrak{g} . Let $P_+ \subseteq P$ be the set of dominant integral weights, i.e. $\lambda \in P$ such that $\langle \lambda, \alpha_i \rangle \geq 0$ for $i = 1, \dots, n$. Moreover, let $P_{++} \subseteq P_+ \subseteq P$ be the set of those dominant weights λ such that $\langle \lambda, \alpha_i \rangle > 0$ for $i = 1, \dots, n$.

The theory of $U_q(\mathfrak{g})$ -modules is very similar to the case $q = 1$ (see [7]). It is well known that for every $q \in (0, 1)$, the monoidal category $\mathcal{M}_q(\mathfrak{g})$ of admissible finite dimensional $U_q(\mathfrak{g})$ -modules are parameterized by $\lambda \in P_+$, with the same fusion rules as $\mathcal{M}(\mathfrak{g})$, the monoidal categories of finite dimensional $U(\mathfrak{g})$ -modules. Moreover, for any $\lambda \in P_+$, the vector spaces V_λ^q and V_λ have the same dimension. There is a similar decomposition into weight sub-spaces $V_\lambda^q(\gamma) \subseteq V_\lambda^q$, for $\gamma \in P$; these are the sub-spaces such that

$$K_i \cdot \eta = q_i^{\gamma(H_i)} \eta, \quad \eta \in V_\lambda^q(\gamma), \quad i = 1, \dots, n. \quad (11)$$

For each $\gamma \in P$, we also have a vector space isomorphism

$$V_\lambda^q(\gamma) \cong V_\lambda(\gamma).$$

In particular, the sub-space $V_\lambda^q(\lambda)$ is one-dimensional and is the highest weight-space of V_λ^q , in the sense that for $\xi \in V_\lambda^q(\lambda)$, we have

$$E_i \cdot \xi = 0, \quad \text{for } i = 1, \dots, n.$$

There is a non-degenerate inner product $\langle \cdot, \cdot \rangle$ on V_λ^q such that

$$\langle a \cdot \eta, \xi \rangle = \langle \eta, a^* \cdot \xi \rangle, \quad \forall a \in U_q(\mathfrak{g}), \quad \forall \eta, \xi \in V_\lambda^q.$$

Clearly, with respect to this inner product, different weight sub-spaces $V_\lambda^q(\gamma)$ are orthogonal for different γ . For any $w \in W$, the Weyl group of G , the weight sub-space $V_\lambda^q(w \cdot \lambda)$ is one dimensional. Thus we can choose a unit vector $\xi_{w \cdot \lambda} \in V_\lambda^q(w \cdot \lambda)$. Let us do this in such way that $\xi_{w \cdot \lambda} = \xi_{v \cdot \lambda}$ if $w \cdot \lambda = v \cdot \lambda$ and thus no ambiguity arises from this notation.

For $w \in W$, let $\ell(w) \in \mathbb{Z}_+$ denote the length of w . By definition, this is the smallest integer m such that w can be written as a product of m generators

$$w = s_{i_1} \cdots s_{i_m}. \quad (12)$$

For the identity element $e \in W$, we let $\ell(e) = 0$. If $\ell(w) = m$ in (12) then the product $w = s_{i_1} \cdots s_{i_m}$ is said to be *reduced*. Recall that the Bruhat order on W is the partial ordering determined by declaring that

$$v < w, \text{ if } \exists \alpha \in \Phi_+ \text{ such that } v\alpha = w \text{ and } \ell(v) = \ell(w) - 1. \quad (13)$$

Let $U_q(\mathfrak{b})$ be the sub-algebra of $U_q(\mathfrak{g})$ generated by K_i, K_i^{-1}, E_i for $i = 1, \dots, n$. We can concretely connect the Bruhat order of W with certain $U_q(\mathfrak{g})$ -modules in the following way:

Lemma 3 (Proposition 3.3 in [10]). *Let $v, w \in W$. The following are equivalent:*

- (i) *We have $v \leq w$ in the Bruhat order on W .*
- (ii) *For any $\lambda \in P_{++}$ we have $V_\lambda^q(v \cdot \lambda) \subset (U_q(\mathfrak{b}))V_\lambda^q(w \cdot \lambda)$.*

Both the subspaces $V_\lambda^q(v \cdot \lambda)$ and $(U_q(\mathfrak{b}))V_\lambda^q(w \cdot \lambda)$ are invariant under the actions of $K_i, i = 1, \dots, n$. Notice that these are commuting self-adjoint operators on V_λ^q that separate the weight-spaces. As the space $V_\lambda^q(v \cdot \lambda)$ is one-dimensional, it follows from (11) that if $v \not\leq w$ and hence $V_\lambda^q(v \cdot \lambda) \not\subset (U_q(\mathfrak{b}))V_\lambda^q(w \cdot \lambda)$, then we have the following corollary of Lemma 3.

Corollary 4. *If $\lambda \in P_{++}$ and $v \not\leq w$, then*

$$V_\lambda^q(v \cdot \lambda) \perp (U_q(\mathfrak{b}))V_\lambda^q(w \cdot \lambda).$$

2.2. The quantum group $\mathbb{C}[G]_q$. We define $\mathbb{C}[G]_q \subseteq (U_q(\mathfrak{g}))^*$ as the subspace of the dual generated by linear functionals of the form

$$\begin{aligned} C_{\eta, \xi}^\lambda(a) &:= \langle a \cdot \eta, \xi \rangle, \\ \text{for } a \in U_q(\mathfrak{g}), \eta, \xi \in V_\lambda^q, \lambda \in P_+. \end{aligned} \quad (14)$$

We let Δ_q, ϵ_q and S_q respectively be the dual of the product, co-product and antipode of $U_q(\mathfrak{g})$. Moreover, we define a $*$ -involution on $\mathbb{C}[G]_q$ by the formula

$$(C_{\eta, \xi}^\lambda)^*(a) = \overline{C_{\eta, \xi}^\lambda((\hat{S}_q(a))^*)}.$$

We let $\mathbb{C}[G] = \mathbb{C}[G]_1$ as well as $\Delta = \Delta_1, \epsilon = \epsilon_1$ and $S = S_1$ denote the $*$ -algebra of regular functions on G with the usual co-product, co-unit and antipode. For an irreducible module V_λ^q with an orthonormal basis $\xi_k, k = 1, \dots, m = \dim V_\lambda^q$, it follows from the definition of $\mathbb{C}[G]_q$ that for $i, k = 1, \dots, m$,

$$\Delta_q(C_{\xi_i, \xi_k}^\lambda) = \sum_{j=1}^m C_{\xi_i, \xi_j}^\lambda \otimes C_{\xi_j, \xi_k}^\lambda, \quad (15)$$

$$\epsilon_q(C_{\xi_i, \xi_k}^\lambda) = C_{\xi_i, \xi_k}^\lambda(I) = \langle \xi_i, \xi_k \rangle = \delta_{ik} 1. \quad (16)$$

$$\sum_{i=1}^m (C_{\xi_i, \xi_k}^\lambda)^* C_{\xi_i, \xi_j}^\lambda = \delta_{km} I, \quad k, j = 1, \dots, m. \quad (17)$$

To deduce (17), consider any $a \in U_q(\mathfrak{g})$, then it follows from the Hopf algebra axioms that (noticing that $C_{\eta,\xi}^\lambda(a) = C_{\xi,\eta}^\lambda(a^*)$ and hence $(C_{\eta,\xi}^\lambda)^*(a) = C_{\xi,\eta}^\lambda(\hat{S}_q(a))$)

$$\begin{aligned} \sum_{i=1}^m (C_{\xi_i,\xi_k}^\lambda)^* C_{\xi_i,\xi_j}^\lambda(a) &= \sum_{i=1}^m ((C_{\xi_i,\xi_k}^\lambda)^* \otimes C_{\xi_i,\xi_j}^\lambda)(\hat{\Delta}_q(a)) \\ &= \sum_{i=1}^m (C_{\xi_k,\xi_i}^\lambda \otimes C_{\xi_i,\xi_j}^\lambda)((\hat{S}_q \otimes \iota)\hat{\Delta}_q(a)) \\ &= C_{\xi_k,\xi_j}^\lambda(\sum \hat{S}_q(a_{(1)})a_{(2)}) = C_{\xi_k,\xi_j}^\lambda(\hat{e}_q(a)) = \delta_{kj}\hat{e}_q(a)I, \end{aligned}$$

where $\hat{\Delta}_q(a) = \sum a_{(1)} \otimes a_{(2)}$ in the Sweedler notation. We define inductively

$$\begin{aligned} \Delta_q^{(2)} &= \Delta_q. \\ \Delta_q^{(n)} &= \underbrace{(\Delta_q \otimes \iota \otimes \cdots \otimes \iota)}_{n-1 \text{ terms}} \circ \Delta_q^{(n-1)} : \mathbb{C}[G]_q \longrightarrow \underbrace{\mathbb{C}[G]_q \otimes \cdots \otimes \mathbb{C}[G]_q}_{n \text{ terms}}. \end{aligned} \quad (18)$$

Notice that by co-associativity $(\Delta_q \otimes \iota) \circ \Delta_q = (\iota \otimes \Delta_q) \circ \Delta_q$, it does not matter which tensor factor you apply Δ_q to in (18).

Let us denote by $C(G)_q$ the universal enveloping C^* -algebra of $\mathbb{C}[G]_q$. It is known from [7] that the universal enveloping C^* -algebra exists and that the natural homomorphism $\mathbb{C}[G]_q \hookrightarrow C(G)_q$ is injective. Hence we can identify $\mathbb{C}[G]_q$ with its inclusion $\mathbb{C}[G]_q \subseteq C(G)_q$. Moreover, the co-product can be extended to a $*$ -homomorphism $\Delta_q : C(G)_q \rightarrow C(G)_q \otimes C(G)_q$ (the minimal tensor product), giving a structure of $C(G)_q$ as a compact quantum group in the sense of Woronowicz [13]. We will use the same symbol for a $*$ -representation of $\mathbb{C}[G]_q$ as well as its extension to $C(G)_q$.

Recall the special case of SU_2 . Let V_λ^q be the unique 2-dimensional $U_q(\mathfrak{su}_2)$ -module, with basis $\xi_\lambda =: \xi_1, \xi_{-\lambda} =: \xi_2$ and let

$$t_{ij} = C_{\xi_i,\xi_j}^\lambda, \quad \text{for } i, j = 1, 2.$$

Then the elements t_{ij} generate $\mathbb{C}[SU_2]_q$ as an algebra, and they are subject to the relations

$$\begin{aligned} t_{11}t_{21} &= qt_{21}t_{11}, \quad t_{11}t_{12} = qt_{12}t_{11}, \quad t_{12}t_{21} = t_{21}t_{12}, \\ t_{22}t_{21} &= q^{-1}t_{21}t_{11}, \quad t_{22}t_{12} = q^{-1}t_{12}t_{22}, \\ t_{11}t_{22} - t_{22}t_{11} &= (q - q^{-1})t_{12}t_{21}, \quad t_{11}t_{22} - qt_{12}t_{21} = 1, \\ t_{11}^* &= t_{22}, \quad t_{12}^* = -qt_{21}. \end{aligned} \quad (19)$$

Moreover, the relations (19) determine $\mathbb{C}[SU_2]_q$, in the sense that this algebra is isomorphic to the universal $*$ -algebra with generators \hat{t}_{ij} , $i, j = 1, 2$, satisfying the relations (19).

Given two $*$ -representations π_1, π_2 , such that $\pi_i : C(G)_q \rightarrow \mathcal{B}(H_i)$ for $i = 1, 2$, we can define the tensor product using the co-multiplication as

$$\pi_1 \boxtimes \pi_2 := (\pi_1 \otimes \pi_2) \circ \Delta_q : C(G)_q \longrightarrow \mathcal{B}(H_1) \otimes \mathcal{B}(H_2) \subseteq \mathcal{B}(H_1 \otimes H_2) \quad (20)$$

where \otimes denotes the minimal tensor product between C^* -algebras. We will also use \otimes to denote the algebraic tensor product; it will always be clear from context which one we use (e.g. we will never take the algebraic tensor product between two C^* -algebras).

2.3. Representation theory of $\mathbb{C}[G]_q$. Recall a $*$ -representation Π_q of $\mathbb{C}[\mathrm{SU}_2]_q$ defined in the following way: let $C_q, S, d_q : \ell^2(\mathbb{Z}_+) \rightarrow \ell^2(\mathbb{Z}_+)$ be the operators defined on the natural orthonormal basis $\{e_j\}_{j \in \mathbb{Z}_+}$ as follows:

$$S e_n = e_{n+1}, \quad C_q e_n = \sqrt{1 - q^{2n}} e_n, \quad d_q e_n = q^n e_n. \quad (21)$$

Then the map

$$\Pi_q(t_{1,1}) = S^* C_q, \quad \Pi_q(t_{1,2}) = q d_q, \quad \Pi_q(t_{2,1}) = -d_q, \quad \Pi_q(t_{2,2}) = C_q S \quad (22)$$

extends to a $*$ -representation of $\mathbb{C}[\mathrm{SU}_2]_q$. Let $C^*(S) \subseteq \mathcal{B}(\ell^2(\mathbb{Z}_+))$ be the C^* -algebra generated by S (so that $C^*(S)$ is equal to the Toeplitz algebra). From the expressions (21) for C_q and d_q it is easy to see that $C_q, d_q \in C^*(S)$ and hence that

$$\Pi_q(\mathbb{C}[\mathrm{SU}_2]_q) \subseteq C^*(S).$$

We recall the $*$ -representation theory for $\mathbb{C}[G]_q$, $q \in (0, 1)$, due to Soibelman [7]: For each simple root $\alpha_i \in \Omega$ we get an injective Hopf $*$ -homomorphism $U_{q_i}(\mathfrak{su}_2) \rightarrow U_q(\mathfrak{g})$ that dualizes to a surjective Hopf $*$ -homomorphism

$$\varsigma_i^q : \mathbb{C}[G]_q \longrightarrow \mathbb{C}[\mathrm{SU}_2]_{q_i}, \quad i = 1, \dots, n$$

(this is true also for $q = 1$, we write $\varsigma_i = \varsigma_i^1$). For $i = 1, \dots, n$, we define $*$ -representations

$$\pi_i^q := \Pi_{q_i} \circ \varsigma_i^q : \mathbb{C}[G]_q \longrightarrow C^*(S) \subseteq \mathcal{B}(\ell^2(\mathbb{Z}_+)).$$

Let $w \in W$, with a reduced presentation $w = s_{j_1} \cdots s_{j_m}$ (and hence $m = \ell(w)$), and define

$$\pi_w^q := \pi_{j_1}^q \boxtimes \cdots \boxtimes \pi_{j_m}^q : \mathbb{C}[G]_q \longrightarrow C(S)^{\otimes \ell(w)} \subseteq \mathcal{B}(\ell^2(\mathbb{Z}_+)^{\otimes \ell(w)}) \quad (23)$$

when $e \in W$ is the identity element, we let $\pi_e = \epsilon_q$ (corresponding to the empty reduced presentation). From [7], we know that π_w^q does not depend on the reduced decomposition, in the sense that if we have two reduced presentations $w = s_{j_1} \cdots s_{j_m} = s_{j'_1} \cdots s_{j'_m}$, then the two corresponding $*$ -representations

$$\pi_{j_1}^q \boxtimes \cdots \boxtimes \pi_{j_m}^q, \quad \pi_{j'_1}^q \boxtimes \cdots \boxtimes \pi_{j'_m}^q$$

are unitarily equivalent. For simplicity, let us write

$$H_w := \ell^2(\mathbb{Z}_+)^{\otimes \ell(w)}.$$

Thus π_w^q is a $*$ -representation $\mathbb{C}[G]_q \rightarrow \mathcal{B}(H_w)$. We have a subgroup $T \subseteq G$ of a maximal torus, corresponding to the real sub-algebra $\mathfrak{t} \subseteq \mathfrak{g}$. Let ω_i , $i = 1, \dots, n$, be the fundamental weights for \mathfrak{g} . We have an isomorphism $T \cong \mathbb{T}^n$, given by

$$t = e^x \in T \mapsto (e^{\omega_1(x)}, \dots, e^{\omega_n(x)}) \in \mathbb{T}^n, \quad (24)$$

where $x \in \mathfrak{t}$. For every $t \in T$, we have mutually non-equivalent one-dimensional $*$ -representations $\chi_t : \mathbb{C}[G]_q \rightarrow \mathbb{C}$, such that, for $s, t \in T$, we have $\chi_s \boxtimes \chi_t = \chi_{st}$ and

$\chi_t \boxtimes \chi_{t^{-1}} = \chi_1 = \epsilon_q$. By [10], we can for every $t \in T$ associate a unitary operator $U_t \in \prod_{\lambda \in P_+} \mathcal{B}(V_\lambda^q)$, such that if $x \in \mathfrak{t}$ satisfies $t = e^x$, then for all $\lambda \in P_+$ and $\gamma \in P$

$$\langle U_t \eta, \xi \rangle = \chi_t(C_{\eta, \xi}^\lambda), \quad \eta, \xi \in V_\lambda^q, \quad (25)$$

$$U_t \eta = e^{\gamma(x)} \eta, \quad \eta \in V_\lambda^q(\gamma). \quad (26)$$

The following theorem characterizes all irreducible $*$ -representations of $\mathbb{C}[G]_q$ (up to unitary equivalence).

Theorem 5 (Theorem 6.2.7 in [7]).

- (i) For $w \in W$ and $t \in T$, the $*$ -representations $\pi_w^q \boxtimes \chi_t$ are irreducible and mutually non-equivalent, and
- (ii) every irreducible $*$ -representation π of $\mathbb{C}[G]_q$ is unitarily equivalent to some $\pi_w \boxtimes \chi_t$, $w \in W$, $t \in T$.

3. Basis Properties of $C(G)_q$ Under $*$ -Representations

3.1. Paths in the Weyl group and subsets of T . For elements $v, w \in W$, we write $v \triangleleft w$ to mean that

- (i) $v < w$ in the Bruhat order,
- (ii) there is no $r \in W$, such that $v < r < w$.

By Theorem 2.2.6 in [1], this means that there is a $\alpha \in \Phi_+$ such that $v\alpha = w$ and $l(v) = l(w) - 1$. Keeping with the established terminology, we also say that w covers v if $v \triangleleft w$. In general, we write $v \triangleleft^{(k)} w$ if we have elements $r_1, \dots, r_{k-1} \in W$ such that

$$v \triangleleft r_1 \triangleleft \dots \triangleleft r_{k-1} \triangleleft w.$$

It is a property of W that every chain

$$v \triangleleft r \triangleleft \dots \triangleleft w$$

must have the same length and hence that the relation $\triangleleft^{(k)}$ is actually well-defined for $k > 1$ (see Theorem 2.2.6 in [1]).

Let $v, w \in W$. If $v \triangleleft w$, we also write $v \xrightarrow{\gamma} w$, for a $\gamma \in \Phi_+$, if $v\gamma = w$. In general, if $v \leq w$, then we say that we have a path from v to w

$$v = v_1 \xrightarrow{\gamma_1} v_2 \xrightarrow{\gamma_2} \dots \xrightarrow{\gamma_{m-1}} v_m \xrightarrow{\gamma_m} v_{m+1} = w \quad (27)$$

if $v_j \triangleleft v_{j+1}$ and $v_j \gamma_{j+1} = v_{j+1}$ for $j = 1, \dots, m$. Clearly, every path from v to w has the same length $m = l(w) - l(v)$. We write (27) as the composition of paths

$$v \xrightarrow{\gamma} w, \text{ for } \gamma = \gamma_1 \circ \dots \circ \gamma_m \quad (28)$$

to indicate that we have a specific path between v and w .

For each path $v \xrightarrow{\gamma} w$ we associate a closed connected subgroup $T_\gamma \subseteq T$ by taking the exponential of the real subspace of \mathfrak{t} spanned by h_{γ_i} for $i = 1, \dots, m$. For the path $v \xrightarrow{\gamma_1} r \xrightarrow{\gamma_2} w$ and $\gamma = \gamma_2 \circ \gamma_1$, we have

$$T_{\gamma_2} T_{\gamma_1} := \{ts : t \in T_{\gamma_2}, s \in T_{\gamma_1}\} = T_\gamma. \quad (29)$$

Let T_v^w be the union of all the subgroups of T generated by paths from v to w . From (29), it follows that we have the following multiplicative property: If $v \leq r \leq w$ then

$$T_v^r T_r^w \subseteq T_v^w.$$

Clearly, it follows from (27) that for $v < w$, we have

$$T_v^w = \bigcup_{v < r \triangleleft w} T_v^r T_r^w, \quad (30)$$

where the union ranges over all $r \in W$ such that $v < r \triangleleft w$. For later use, we also note the following special case of (29): if $v \overset{\gamma}{\rightsquigarrow} r \triangleleft w$ with $r\alpha = w$ then

$$T_\gamma T_r^w = T_{\alpha \circ \gamma}. \quad (31)$$

3.2. Ideals and quotients. We will use the following results from [10] (though stated in a less general fashion in order to suit our purposes).

Theorem 6 (Theorem 4.1 (ii) in [10]). *Let $\sigma \in W$ and $Y \subseteq T$. For any $r \in W$ and $t \in T$, the kernel of the representation $\pi_r^q \boxtimes \chi_t$ contains the intersection of the kernels of the representations $\pi_\sigma^q \boxtimes \chi_s$, $s \in Y$ of $C(G)_q$ if and only if $r \leq \sigma$ and $t \in \bar{Y}T_r^\sigma$.*

Lemma 7 (Lemma 4.5 in [10]). *Let $t \in T$ and let $w \in W$. Assume $x \in C(G)_q$ is such that $(\pi_v^q \boxtimes \chi_s)(x) = 0$ for all $v \in W$ such that $v < w$ and $s \in tT_v^w$, then*

$$(\pi_w^q \boxtimes \chi_t)(x) \in \mathcal{K}_w. \quad (32)$$

Recall the definition of $C_{\eta, \xi}^\lambda \in \mathbb{C}[G]_q$, given by (14). To avoid multiple subscripts, let us write $w \cdot \lambda$ in place of $\xi_{w \cdot \lambda}$ in (14). Thus, for example, we write $C_{w \cdot \lambda, \lambda}^\lambda$ instead of $C_{\xi_{w \cdot \lambda}, \xi_\lambda}^\lambda$.

Lemma 8 (Lemma 2.3 in [10]). *Let $w \in W$ and $\lambda \in P_+$.*

- (i) $\pi_w^q(C_{w \cdot \lambda, \lambda}^\lambda)$ is a compact contractive diagonalizable operator with zero kernel, and the vector $e_0^{\otimes \ell(w)} \in H_w$ is its only eigenvector (up to scalar) with an eigenvalue of absolute value 1.
- (ii) If $\zeta \in V_\lambda^q$ is orthogonal to $(U_q(\mathfrak{b}))V_\lambda^q(w \cdot \lambda)$, then

$$\pi_w^q(C_{\zeta, \lambda}^\lambda) = 0.$$

When $q = 1$, we get a Hopf $*$ -algebra homomorphism $\tau_1 = \tau : \mathbb{C}[G] \rightarrow \mathbb{C}[T]$ by restriction to the subgroup $T \subseteq G$ (here $\mathbb{C}[T]$ is the Hopf $*$ -algebra of trigonometric polynomials on T). For general $q \in (0, 1)$, we define a surjective Hopf $*$ -algebra homomorphism $\tau_q : \mathbb{C}[G]_q \rightarrow \mathbb{C}[T]$, that extends to a homomorphism of compact quantum groups $\tau_q : C(G)_q \rightarrow C(T)$. We define τ_q in the following way: The compact operators $\mathcal{K} \subseteq \mathcal{B}(\ell^2(\mathbb{Z}_+))$ is a $*$ -ideal in $C^*(S)$, and it is well known that we have the isomorphism $p : C^*(S)/\mathcal{K} \xrightarrow{\sim} C(\mathbb{T})$ such that $S^* + \mathcal{K} \mapsto z$ (here $z \in C(\mathbb{T})$ is the coordinate function). Moreover, it is easy to see that we actually have a homomorphism of Hopf $*$ -algebras $\beta_q : \mathbb{C}[SU_2]_q \rightarrow \mathbb{C}[\mathbb{T}] \subseteq C(\mathbb{T})$ that factors as

$$\mathbb{C}[SU_2]_q \xrightarrow{\Pi_q} C^*(S) \xrightarrow{p} C^*(S)/\mathcal{K} \cong C(\mathbb{T}) \quad (33)$$

and such that $\beta_q(t_{11}^q) = z$. Consider now the $*$ -homomorphism

$$\tau_q : \mathbb{C}[G]_q \xrightarrow{\pi_1^q \boxtimes \dots \boxtimes \pi_n^q} C^*(S)^{\otimes n} \xrightarrow{p \otimes \dots \otimes p} C(\mathbb{T})^{\otimes n} \cong C(\mathbb{T}), \quad (34)$$

with the isomorphism $C(\mathbb{T})^{\otimes n} \cong C(\mathbb{T})$ induced by the isomorphism $\mathbb{T}^n \cong \mathbb{T}$ given by (24). By using (33), we can also factor τ_q as

$$\mathbb{C}[G]_q \xrightarrow{(\varsigma_1^q \otimes \dots \otimes \varsigma_n^q) \circ \Delta_q^{(n)}} \mathbb{C}[\mathrm{SU}_2]_{q_1} \otimes \dots \otimes \mathbb{C}[\mathrm{SU}_2]_{q_n} \xrightarrow{\beta_{q_1} \otimes \dots \otimes \beta_{q_n}} \mathbb{C}[\mathbb{T}]^{\otimes n} \cong \mathbb{C}[\mathbb{T}]. \quad (35)$$

If we consider $\mathbb{C}[\mathrm{SU}_2]_{q_1} \otimes \dots \otimes \mathbb{C}[\mathrm{SU}_2]_{q_n}$ as a tensor product of Hopf $*$ -algebras, hence also a Hopf $*$ -algebra, then it is easy to check that the $*$ -homomorphisms in (35) are actually Hopf $*$ -algebra morphisms. Thus, τ_q is a morphisms of Hopf $*$ -algebras.

Lemma 9. (i) *The $*$ -homomorphism $\tau_q : C(G)_q \rightarrow C(\mathbb{T})$ is surjective,*
(ii) *every χ_t , $t \in \mathbb{T}$, factors as*

$$\chi_t : C(G)_q \xrightarrow{\tau_q} C(\mathbb{T}) \xrightarrow{ev_t} \mathbb{C},$$

(iii) *for $\eta \in V_\lambda(\gamma_1)$, $\xi \in V_\lambda(\gamma_2)$, we have $\tau_q(C_{\eta, \xi}^\lambda) = 0$ unless $\gamma_1 = \gamma_2$ and $\langle \eta, \xi \rangle \neq 0$,*
(iv) *let ω_i , $i = 1, \dots, n$, be the fundamental weights, then the set $\tau_q(C_{\omega_i, \omega_i}^{\omega_i})$, $i = 1, \dots, n$, generates $C(\mathbb{T})$ as a C^* -algebra.*

Proof. Clearly, (i) follows from (iv) or (ii). We have that every one-dimensional $*$ -representation of $\mathbb{C}[G]_q$ is of the form χ_t , for some $t \in \mathbb{T}$, and different weight spaces are orthogonal. Hence we get (iii) from (25) and (26) by point evaluation. Note that the same argument shows that (iii) holds for *any* $*$ -homomorphism from $\mathbb{C}[G]_q$ to a commutative C^* -algebra.

By Lemma 2.2.1 in [7], the elements of the form $C_{\eta, \lambda}^\lambda$, $\lambda \in P_+$, $\eta \in V_\lambda^q$ generates $\mathbb{C}[G]_q$ as a $*$ -algebra. Thus it follows from (iii) that the elements in $C(\mathbb{T})$ of the form $\tau_q(C_{\lambda, \lambda}^\lambda)$, $\lambda \in P_+$ generate the image of τ_q . As every $\lambda \in P_+$ is a linear combination of the fundamental weights, it follows from (26) that the set $\{\tau_q(C_{\omega_i, \omega_i}^{\omega_i}) : i = 1, \dots, n\}$ generate the image of τ_q . Thus (ii) \Rightarrow (i) \Rightarrow (iv). To prove (ii), we establish a one-to-one correspondence between one-dimensional $*$ -representations and point evaluations of τ_q . By ([6], Theorem 14, Section 6.1.5), the set $F^i K^j E^k$, $j \in \mathbb{Z}$, $k \in \mathbb{Z}_+$, is a basis for $U_q(\mathfrak{su}_2)$. Using this, and keeping in mind that ξ_{ω_i} is a highest weight vector, we obtain

$$\varsigma_i^q(C_{\omega_j, \omega_j}^{\omega_j}) = \begin{cases} t_{11}^{q_i}, & \text{if } i = j \\ I, & \text{for } i \neq j. \end{cases} \quad (36)$$

It then follows that $(\beta_i \circ \varsigma_i^q)(C_{\omega_j, \omega_j}^{\omega_j}) = z$ for $i = j$ and $(\beta_i \circ \varsigma_i^q)(C_{\omega_j, \omega_j}^{\omega_j}) = I$ if $i \neq j$. If we extend $\xi_1 = \xi_{\omega_1}$ by ξ_2, \dots, ξ_m to an orthonormal basis for $V_{\omega_i}^q$, we then get from (15) and our comment after the proof of (iii), that if we let z_i be the i 'th coordinate function of \mathbb{T}^n , then

$$\tau_q(C_{\omega_i, \omega_i}^{\omega_i}) = z_i, \quad i = 1, \dots, n, \quad (37)$$

and from this it follows that the range of τ_q is dense in $C(T)$. By (26), if $t = e^x$, $x \in \mathfrak{t}$, then we have $\chi_t(C_{\omega_j, \omega_j}^{\omega_j}) = e^{\omega_j(x)}$ for $j = 1, \dots, n$. It thus follows that if we use the identification $T \cong \mathbb{T}^n$ given by (24), then $t = (e^{\omega_1(x)}, \dots, e^{\omega_n(x)})$ and hence

$$\chi_t(C_{\omega_j, \omega_j}^{\omega_j}) = e^{\omega_j(x)} = \text{ev}_t(z_j) = (\text{ev}_t \circ \tau_q)(C_{\omega_j, \omega_j}^{\omega_j}), \quad j = 1, \dots, n.$$

□

Let $\mathbb{C}[G]_q^{\text{inv}} \subseteq \mathbb{C}[G]_q$ denote the $*$ -subalgebra of elements invariant under the left-right-action of the maximal torus T . By definition, this is the subset $x \in \mathbb{C}[G]_q$, such that for every $t \in T$, we have $L_t(x) = x = R_t(x)$, where L_t and R_t are given by

$$L_t : \mathbb{C}[G]_q \xrightarrow{\Delta_q} \mathbb{C}[G]_q \otimes \mathbb{C}[G]_q \xrightarrow{\tau_q \otimes \text{id}} C(T) \otimes \mathbb{C}[G]_q \xrightarrow{\text{ev}_t \otimes \text{id}} \mathbb{C}[G]_q \text{ (Left-action).} \quad (38)$$

$$R_t : \mathbb{C}[G]_q \xrightarrow{\Delta_q} \mathbb{C}[G]_q \otimes \mathbb{C}[G]_q \xrightarrow{\text{id} \otimes \tau_q} \mathbb{C}[G]_q \otimes C(T) \xrightarrow{\text{id} \otimes \text{ev}_t} \mathbb{C}[G]_q \text{ (Right-action)} \quad (39)$$

Clearly $\mathbb{C}[G]_q^{\text{inv}}$ is a $*$ -subalgebra of $\mathbb{C}[G]_q$.

Lemma 10. *For every $w \in W$, there exists $\Upsilon_w \in \mathbb{C}[G]_q^{\text{inv}}$ such that*

- (i) $\pi_w^q(\Upsilon_w) \in \mathcal{B}(H_w)$ is a compact contractive positive operator with dense range,
- (ii) $e_0^{\otimes \ell(w)} \in H_w$ is the only eigenvector of $\pi_w^q(\Upsilon_w)$ (up to a scalar multiple) with eigenvalue 1,
- (iii) $\pi_v^q(\Upsilon_w) \neq 0$ if and only if $v \geq w$.

Proof. Take any $\lambda \in P_{++}$ and consider $C_{w \cdot \lambda, \lambda}^\lambda$. By combining Corollary 4 with Lemma 8 it follows that $\pi_v^q(C_{w \cdot \lambda, \lambda}^\lambda) = 0$ for any $v \not\geq w$. If $v \geq w$, then as $1 \in T_w^v$ (the identity of T) it follows from Theorem 6 that we have $\ker \pi_v^q \subseteq \ker \pi_w^q$. As $\pi_w^q(C_{w \cdot \lambda, \lambda}^\lambda) \neq 0$, it thus follows that also $\pi_v^q(C_{w \cdot \lambda, \lambda}^\lambda) \neq 0$. If we extend $\xi_\lambda, \xi_{w \cdot \lambda}$ to an orthonormal basis for V_λ^q , then we get from (15) and Lemma 9 (iii) that

$$L_t(C_{w \cdot \lambda, \lambda}^\lambda) = \chi_t(C_{w \cdot \lambda, w \cdot \lambda}^\lambda) \cdot C_{w \cdot \lambda, \lambda}^\lambda, \quad R_t(C_{w \cdot \lambda, \lambda}^\lambda) = \chi_t(C_{\lambda, \lambda}^\lambda) \cdot C_{w \cdot \lambda, \lambda}^\lambda, \quad t \in T. \quad (40)$$

Let us now define $\Upsilon_w := (C_{w \cdot \lambda, \lambda}^\lambda)^* C_{w \cdot \lambda, \lambda}^\lambda$. As L_t and R_t are $*$ -automorphisms for all $t \in T$, it follows from (40) that $\Upsilon_w \in \mathbb{C}[G]_q^{\text{inv}}$. Positivity follows from the definition of Υ_w , and the other claims in (i) and (iii) follow by Lemma 8 (i). To see (ii), note that

$$\langle \pi_w^q(\Upsilon_w) e_0^{\otimes \ell(w)}, e_0^{\otimes \ell(w)} \rangle = \|\pi_w^q(C_{w \cdot \lambda, \lambda}^\lambda) e_0^{\otimes \ell(w)}\|^2 = \|e_0^{\otimes \ell(w)}\|^2 = 1. \quad (41)$$

As $\pi_w^q(\Upsilon_w)$ is a positive contraction, it follows that $e_0^{\otimes \ell(w)}$ must be an eigenvector with eigenvalue 1. To see that $e_0^{\otimes \ell(w)}$ is the only eigenvector (up to a scalar multiple) of $\pi_w^q(\Upsilon_w)$ with eigenvalue 1, notice that (41) gives that $e_0^{\otimes \ell(w)}$ is also an eigenvector for $\pi_w^q(C_{w \cdot \lambda, \lambda}^\lambda)^*$ and thus the subspace generated by $e_0^{\otimes \ell(w)}$ is actually reducing $\pi_w^q(C_{w \cdot \lambda, \lambda}^\lambda)$. By (i) of Lemma 8, it follows that $\pi_w^q(C_{w \cdot \lambda, \lambda}^\lambda)$ must have norm strictly less than 1 when restricted to the orthogonal complement of $e_0^{\otimes \ell(w)}$ (otherwise there would be another eigenvector orthogonal to $e_0^{\otimes \ell(w)}$ with an eigenvalue of absolute value 1). The same then holds for $\pi_w^q(\Upsilon_w) = \pi_w^q(C_{w \cdot \lambda, \lambda}^\lambda)^* \pi_w^q(C_{w \cdot \lambda, \lambda}^\lambda)$. □

Definition 11. Let us call a $*$ -homomorphism of $\mathbb{C}[G]_q$ a *commutative $*$ -representation* if the image sits inside a commutative C^* -algebra.

Let $\chi : \mathbb{C}[G]_q \rightarrow C(X)$ be a commutative $*$ -representation. As every one-dimensional $*$ -representation of $\mathbb{C}[G]_q$ factors through the commutative C^* -algebra $C(T)$, it follows that the commutative $*$ -representation χ factors as

$$\begin{aligned}\chi &= \zeta \circ \tau_q \\ \mathbb{C}[G]_q &\xrightarrow{\tau_q} C(T) \xrightarrow{\zeta} C(X)\end{aligned}$$

for a unique $*$ -homomorphism ζ .

Definition 12. Let

$$\chi^q : \mathbb{C}[G]_q \rightarrow C(X), \quad q \in (0, 1)$$

be a family of $*$ -homomorphisms, where X is a fixed compact Hausdorff space. We say that the $*$ -homomorphisms $\{\chi^q\}_{q \in (0,1)}$ are *q -independent* if in the factorization

$$\begin{aligned}\chi^q &= \zeta^q \circ \tau_q, \\ \mathbb{C}[G]_q &\xrightarrow{\tau_q} C(T) \xrightarrow{\zeta^q} C(X),\end{aligned}$$

we have $\zeta^q = \zeta^s$ for all $q, s \in (0, 1)$.

Definition 13. (a) For $q \in (0, 1)$, let $B_w^q \subseteq \mathcal{B}(H_w)$ be the closure of the image $\pi_w : \mathbb{C}[G]_q \rightarrow \mathcal{B}(H_w)$.

(b) If $\chi : \mathbb{C}[G]_q \rightarrow C(X)$ is a commutative $*$ -representation, then let $B_{w,\chi}^q$ be the closure of the image of $\pi_w \boxtimes \chi : \mathbb{C}[G]_q \rightarrow B_w^q \otimes C(X)$.

Proposition 14. Let $w \in W$. Let $\mathcal{K}_w \subseteq \mathcal{B}(H_w)$ be the space of compact operators and let

$$p_w : \mathcal{B}(H_w) \longrightarrow \mathbb{Q}(H_w)$$

be the quotient map to the Calkin algebra. For every $v \triangleleft w$, there exists a subset $T_v^w \subseteq T$ and a commutative q -independent $*$ -representation $\chi_v^w : \mathbb{C}[G]_q \rightarrow C(T_v^w)$, such that the map

$$\eta_w^q : \pi_w^q(x) + \mathcal{K}_w \mapsto \bigoplus_{v \triangleleft w} (\pi_v^q \boxtimes \chi_v^w)(x), \text{ for } x \in \mathbb{C}[G]_q, \quad (42)$$

(where the sum ranges over all $v \in W$ covered by w) determines an isomorphism

$$\eta_w^q : B_w^q / \mathcal{K}_w \longrightarrow \overline{\bigoplus_{v \triangleleft w} (\pi_v^q \boxtimes \chi_v^w)(\mathbb{C}[G]_q)}. \quad (43)$$

We will postpone the proof of Proposition 14 until after Lemma 16.

Lemma 15. If $x \in \mathbb{C}[G]_q^{\text{inv}}$, then for any $w \in W$ and any commutative $*$ -representation χ , we have

$$(\pi_w \boxtimes \chi)(x) = \pi_w(x) \otimes I.$$

Proof. By (39) and (38), the left and right-actions by $t \in T$ on $\mathbb{C}[G]_q$ are respectively given as the compositions

$$\begin{aligned}\mathbb{C}[G]_q &\xrightarrow{(\tau_q \otimes \iota) \circ \Delta_q} C(T) \otimes \mathbb{C}[G]_q \xrightarrow{\text{ev}_T \otimes \iota} \mathbb{C}[G]_q, \\ \mathbb{C}[G]_q &\xrightarrow{(\iota \otimes \tau_q) \circ \Delta_q} \mathbb{C}[G]_q \otimes C(T) \xrightarrow{\iota \otimes \text{ev}_T} \mathbb{C}[G]_q.\end{aligned}$$

Let $x \in \mathbb{C}[G]_q^{\text{inv}}$. Clearly, it follows $L_t(x) = R_t(x) = x$ for all $t \in T$ if and only if

$$(\iota \otimes \tau_q) \circ \Delta_q(x) = x \otimes I, \quad (\tau_q \otimes \iota) \circ \Delta_q(x) = I \otimes x.$$

The statement now follows from the fact that any $*$ -homomorphism

$$\chi : C(G)_q \rightarrow C(X)$$

factors as $\chi = \zeta \circ \tau_q$ for a unique $*$ -representation $\zeta : C(T) \rightarrow C(X)$. \square

Lemma 16. *Let $\chi : C(G)_q \rightarrow C(X)$ be a $*$ -homomorphism such that we have $\chi(C(G)_q) = C(X)$. Then for any $w \in W$,*

$$\mathcal{K}_w \otimes C(X) \subset B_{w,\chi}^q. \quad (44)$$

Proof. If we, for any $\lambda \in P_{++}$, extend $\xi_\lambda, \xi_{w \cdot \lambda}$ to an orthonormal basis of V_λ^q , then we get from (15) and Lemma 9 (iii) that

$$(\pi_w^q \boxtimes \tau_q)(C_{w \cdot \lambda, \lambda}^\lambda) = \pi_w^q(C_{w \cdot \lambda, \lambda}^\lambda) \otimes \tau_q(C_{\lambda, \lambda}^\lambda).$$

By Lemma 10 and Lemma 15, if p_0 is the orthogonal projection onto $e_0^{\otimes \ell(w)}$, then $p_0 \otimes I \in B_{w, \tau_q}^q$. Thus

$$(p_0 \otimes I)((\pi_w^q \boxtimes \tau_q)(C_{w \cdot \lambda, \lambda}^\lambda))(p_0 \otimes I) \in B_{w, \tau_q}^q$$

and is by Lemma 8 a non-zero constant multiple of $p_0 \otimes \tau_q(C_{\lambda, \lambda}^\lambda)$. Since the functions

$$\tau_q(C_{\lambda, \lambda}^\lambda) \in C(T), \quad \lambda \in P_{++}$$

are generating $C(T)$ as a C^* -algebra, it follows that

$$p_0 \otimes C(T) \subseteq B_{w, \tau_q}^q. \quad (45)$$

By (Proposition 5.5 in [12]), the restriction of an irreducible $*$ -representation of $C(G)_q$ is still irreducible when restricted to

$$C(G/T)_q \stackrel{\text{def}}{=} \{a \in C(G)_q : (\iota \otimes \tau_q) \circ \Delta_q(a) = a \otimes I\}.$$

It follows that

$$\mathcal{K}_w \otimes I \subseteq (\pi_w^q \boxtimes \tau_q)(C(G/T)_q) \subseteq (\pi_w^q \boxtimes \tau_q)(C(G)_q),$$

and together with (45), this gives (44). \square

Proof of Proposition 14. Recall the definition of $T_v^w \subseteq T$. For $v \triangleleft w$, we let χ_v^w be the composition

$$\chi_v^w : C(G)_q \xrightarrow{\tau_q} C(T) \longrightarrow C(T_v^w),$$

where the second $*$ -homomorphism is the restriction of $f \in C(T)$ to $T_v^w \subseteq T$. By definition, these $*$ -representations are commutative and q -invariant. To prove the existence of the $*$ -homomorphism, we use Theorem 6 with $Y = \{1\}$ (the set containing only the identity of T). Thus for every $v \triangleleft w$ and $t \in T_v^w$ we have that $\ker \pi_w^q \subseteq \ker \pi_v^q \boxtimes \chi_t$ and hence we can define a $*$ -homomorphism

$$\begin{aligned} \varphi_v^w : B_w^q &\longrightarrow B_{v, \chi_v^w}^q \\ \pi_w^q(x) &\mapsto (\pi_v^q \boxtimes \chi_v^w)(x), \text{ for } x \in C(G)_q. \end{aligned}$$

By Lemma 10, we have $\pi_w^q(\gamma_w) \neq 0$ and $\pi_v^q(\gamma_w) = 0$ and by Lemma 15

$$(\pi_v^q \boxtimes \chi_v^w)(\gamma_w) = \pi_v^q(\gamma_w) \otimes I = 0.$$

Hence the kernel of φ_v^w is a non-trivial ideal of B_w^q . By Lemma 16, with $\chi = \epsilon_q$, we have $\mathcal{K}_w \subset B_w^q$. Thus any non-trivial ideal of B_w^q must contain \mathcal{K}_w and therefore $\mathcal{K}_w \subseteq \ker \varphi_v^w$. Now consider the $*$ -homomorphism

$$\bigoplus_{v \triangleleft w} \varphi_v^w : B_w^q \longrightarrow \prod_{v \triangleleft w} B_{v, \chi_v^w}^q.$$

By the definition of the φ_v^w 's, it follows that

$$\left(\bigoplus_{v \triangleleft w} \varphi_v^w \right) \circ \pi_w^q = \bigoplus_{v \triangleleft w} (\pi_v^q \boxtimes \chi_v^w). \quad (46)$$

As $\mathcal{K}_w \subset \ker \bigoplus_{v \triangleleft w} \varphi_v^w$ we can factor $\bigoplus_{v \triangleleft w} \varphi_v^w =: \eta_w^q \circ p_w$ for a $*$ -homomorphism

$$\eta_w^q : B_w^q / \mathcal{K}_w \longrightarrow \bigoplus_{v \triangleleft w} (\pi_v^q \boxtimes \chi_v^w)(C(G)_q).$$

From (46), it follows that (42) holds. Clearly, by (46) η_w^q is surjective. Hence we only need to show that the kernel of η_w^q is trivial. By definition, this is the same as showing

$$\ker \bigoplus_{v \triangleleft w} \varphi_v^w = \mathcal{K}_w. \quad (47)$$

In order to prove this, we are going to show that, for any $x \in C(G)_q$, if $(\pi_v^q \boxtimes \chi_t)(x) = 0$ for all $v \triangleleft w$ and $t \in T_v^w$, then also $(\pi_\sigma^q \boxtimes \chi_s)(x) = 0$ for all $\sigma < w$ and $s \in T_\sigma^w$. To see this, notice that by (30), there is a $\sigma \leq v \triangleleft w$ such that we have $s = s_1 s_2$ for $s_1 \in T_v^w$ and $s_2 \in T_\sigma^v$. As we have $(\pi_v^q \boxtimes \chi_{s_1})(x) = 0$ we get from Theorem 6 that also $(\pi_\sigma^q \boxtimes \chi_s)(x) = 0$. The equality (47) now follows from Lemma 7. \square

Proposition 17. Let $\mathcal{S} \subseteq W$ be a subset where all the elements have the same length i.e. $\ell(w) = \ell(v)$ for all $w, v \in \mathcal{S}$. Moreover, assume that for each $v \in \mathcal{S}$, we have a commutative $*$ -homomorphism $\chi_v : C(G)_q \rightarrow C(X_v)$ such that $\chi_v(C(G)_q) = C(X_v)$. Then

$$\prod_{v \in \mathcal{S}} \mathcal{K}_v \otimes C(X_v) \subseteq \bigoplus_{v \in \mathcal{S}} (\pi_v \boxtimes \chi_v)(C(G)_q) \subseteq \prod_{v \in \mathcal{S}} B_{v, \chi_v}^q. \quad (48)$$

Proof. As all the elements of \mathcal{S} have the same length, they must be mutually non-comparable in the partial ordering of W . It follows from Lemma 10 and Lemma 15 that for $v \in \mathcal{S}$, we have $(\pi_v^q \boxtimes \chi_v)(\Upsilon_v) = \pi_v^q(\Upsilon_v) \otimes I \neq 0$ and $(\pi_w^q \boxtimes \chi_w)(\Upsilon_v) = 0$ for any other $w \in \mathcal{S}$. As $\pi_v^q(\Upsilon_v)$ is a compact operator with dense range, it follows from Lemma 16 that

$$\begin{aligned} \overline{(\pi_v^q \boxtimes \chi_v)(\Upsilon_v(C(G)_q))} &= \mathcal{K}_v \otimes C(X_v) \\ \overline{(\pi_w^q \boxtimes \chi_w)(\Upsilon_v(C(G)_q))} &= \{0\}, \quad w \in \mathcal{S} \text{ such that } w \neq v. \end{aligned}$$

This gives (48). \square

3.3. Continuous deformations.

Lemma 18. *There are invertible co-algebra maps*

$$\theta^q : \mathbb{C}[G] \longrightarrow \mathbb{C}[G]_q, \quad q \in (0, 1)$$

such that for every $w \in W$, every q -independent commutative $$ -representation $\chi^q : \mathbb{C}[G]_q \rightarrow C(X)$ and any fixed $f \in \mathbb{C}[G]$, the map*

$$q \in (0, 1) \mapsto (\pi_w^q \boxtimes \chi^q)(\theta^q(f)) \in B_{w, \chi^q}^q \subseteq \mathcal{B}(H_w) \otimes C(X) \quad (49)$$

is continuous.

Proof. We will refer to the proof of Theorem 1.2 in [11]. We remark that our notation differs from theirs. Let t_{ij}^q , $i, j = 1, 2$ be the generators of $\mathbb{C}[\mathrm{SU}_2]_q$ for $q \in (0, 1]$. It follows from the proof that there are invertible co-algebra maps

$$\kappa_q : \mathbb{C}[\mathrm{SU}_2] \longrightarrow \mathbb{C}[\mathrm{SU}_2]_q, \quad q \in (0, 1], \quad \kappa_1 = \mathrm{Id},$$

such that $\kappa_q(t_{ij}^1) = t_{ij}^q$ and for every $f \in \mathbb{C}[\mathrm{SU}_2]$, the image $\kappa_q(f)$ is a non-commutative polynomial in t_{ij}^q , $i, j = 1, 2$, with coefficients continuous in q . Moreover, there exists an invertible co-algebra map $\vartheta_q : \mathbb{C}[G] \rightarrow \mathbb{C}[G]_q$, such that for every $i = 1, \dots, n$, there exists a continuous family of invertible co-algebra morphisms γ_i^q of $\mathbb{C}[G]$ that makes the following diagram commute

$$\begin{array}{ccccc} \mathbb{C}[\mathrm{SU}_2] & \xleftarrow{\quad \varsigma_i \quad} & \mathbb{C}[G] & \xrightarrow{\quad \gamma_i^q \quad} & \mathbb{C}[G] \\ \kappa_{qi} \downarrow & & & \swarrow \vartheta_q & \\ \mathbb{C}[\mathrm{SU}_2]_{qi} & \xleftarrow{\quad \varsigma_i^q \quad} & \mathbb{C}[G]_q & & \end{array} \quad (50)$$

This gives that

$$\kappa_{qi}^{-1} \circ \varsigma_i^q \circ \vartheta_q = \varsigma_i \circ (\gamma_i^q)^{-1} \quad (51)$$

varies continuously on $q \in (0, 1]$. The operators C_q and d_q , given by (21) varies continuously on $q \in (0, 1)$. Hence, by (22), it follows that the functions $q \in (0, 1) \mapsto$

$\Pi_q \circ \kappa_q(t_{ij}) = \Pi_q(t_{ij}^q) \subseteq \mathcal{B}(\ell^2(\mathbb{Z}_+))$, for $i, j = 1, 2$, are continuous. Thus, for any fixed $f \in \mathbb{C}[\text{SU}_2]$, we have a continuous function

$$q \in (0, 1) \mapsto \Pi_q \circ \kappa_q(f) \in \mathcal{B}(\ell^2(\mathbb{Z}_+)).$$

Composing this with (51), we get that for every $f \in \mathbb{C}[G]$, the function

$$q \in (0, 1) \mapsto \Pi_q \circ \varsigma_i^q \circ \vartheta_q(f) \in \mathcal{B}(\ell^2(\mathbb{Z}_+))$$

is continuous. As the maps in the diagram (50) are all (at least) co-algebra maps, so that $\Delta_q \circ \vartheta_q = (\vartheta_q \otimes \vartheta_q) \circ \Delta$, it follows for $f \in \mathbb{C}[G]$ and $w \in W$ with reduced presentation $w = s_{j_1} \cdots s_{j_m}$, that

$$\begin{aligned} \pi_w^q(\vartheta_q(f)) &= (\pi_{j_1}^q \otimes \cdots \otimes \pi_{j_m}^q) \circ \Delta_q^{(m)}(\vartheta_q(f)) \\ &= (\Pi_{q_{j_1}} \otimes \cdots \otimes \Pi_{q_{j_m}}) \circ ((\varsigma_{j_1}^q \circ \vartheta_q) \otimes \cdots \otimes (\varsigma_{j_m}^q \circ \vartheta_q)) \circ \Delta^{(m)}(f) \\ &= ((\Pi_{q_{j_1}} \circ \kappa_{q_{j_1}} \circ \varsigma_{j_1}) \otimes \cdots \otimes (\Pi_{q_{j_m}} \circ \kappa_{q_{j_m}} \circ \varsigma_{j_m})) \circ \Delta^{(m)}(f) \end{aligned}$$

and hence the function $q \in (0, 1) \mapsto \pi_w^q(\vartheta_q(f))$ is continuous. Combining this with (34), it follows that $q \in (0, 1) \mapsto \tau_q(\vartheta_q(f)) \in C(T)$ is also continuous. Thus

$$q \in (0, 1) \mapsto (\pi_w^q \boxtimes \tau_q)(\vartheta_q(f)) = ((\pi_w^q \circ \vartheta_q) \otimes (\tau_q \circ \vartheta_q)) \circ \Delta(f) \in \mathcal{B}(H_w) \otimes C(T)$$

is continuous. That (49) holds for all q -independent maps follows by the factorization $\chi^q = \zeta \circ \tau_q$. \square

Assume we have two subsets $T_1, T_2 \subseteq T$ and $T_3 = T_1 T_2$ (the point-wise multiplication). Let us denote by χ_i , for $i = 1, 2, 3$ the $*$ -homomorphism $\mathbb{C}[G]_q \rightarrow C(T_i)$, $i = 1, 2, 3$, given by restriction of τ_q to T_i . It follows that we have an identification

$$\chi_1 \boxtimes \chi_2 \sim \chi_3 \quad (52)$$

in the sense that χ_3 is the unique $*$ -homomorphism with the property that, using the isomorphism $C(T_1) \otimes C(T_2) \cong C(T_1 \times T_2)$, we have

$$(\chi_1 \boxtimes \chi_2)(a)(t_1, t_2) = \chi_3(a)(t_1 t_2), \quad t_1 \in T_1, t_2 \in T_2, a \in \mathbb{C}[G]_q. \quad (53)$$

The multiplication map $m : T_1 \times T_2 \rightarrow T_3$ gives an injective $*$ -homomorphism

$$C(T_3) \xrightarrow{m^*} C(T_1) \otimes C(T_2) \cong C(T_1 \times T_2)$$

and it follows from (53) that $\chi_1 \boxtimes \chi_2$ factors as

$$\mathbb{C}[G]_q \xrightarrow{\chi_3} C(T_3) \xrightarrow{m^*} C(T_1) \otimes C(T_2).$$

Furthermore, if for two subsets $T_1, T_2 \subseteq T$, we let $T_3 = T_1 \cup T_2$ and denote by χ_i , $i = 1, 2, 3$, the $*$ -homomorphisms $\mathbb{C}[G]_q \rightarrow C(T_i)$, $i = 1, 2, 3$, then we have an identification

$$\chi_1 \oplus \chi_2 \sim \chi_3 \quad (54)$$

via the injective $*$ -homomorphism $C(T_1 \cup T_2) \rightarrow C(T_1) \oplus C(T_2)$ determined by the two inclusions $T_i \subseteq T_3$ for $i = 1, 2$. Thus χ_3 satisfies

$$\chi_1(a) = \chi_3(a)|_{T_1}, \quad \chi_2(a) = \chi_3(a)|_{T_2}, \quad a \in \mathbb{C}[G]_q \quad (55)$$

where $\chi_3(a)|_{T_i}$, $i = 1, 2$ denotes the restriction of the function $\chi_3(a) \in C(T_3)$ to the subset $T_i \subseteq T_3$.

For a path $v \xrightarrow{\gamma} w$, let us denote by χ_γ the commutative q -independent $*$ -representation $\mathbb{C}[G]_q \rightarrow C(T_\gamma)$. If we have paths $v \xrightarrow{\gamma_1} r \xrightarrow{\gamma_2} w$, then it follows from (29) and (52) that if we have the composition of paths $\gamma = \gamma_1 \circ \gamma_2$, then

$$\chi_{\gamma_1} \boxtimes \chi_{\gamma_2} \sim \chi_\gamma. \quad (56)$$

4. The Main Result

Theorem 19. (i) For all $q, s \in (0, 1)$ and $w \in W$, we have an inner $*$ -automorphism $\Gamma_w^{s,q} : \mathcal{B}(H_w) \rightarrow \mathcal{B}(H_w)$ that restricts to a $*$ -isomorphism $B_w^q \rightarrow B_w^s$, such that $\Gamma_w^{s,q}(\mathcal{K}_w) = \mathcal{K}_w$ and we have

$$\begin{aligned} \Gamma_w^{s,t} \circ \Gamma_w^{t,q} &= \Gamma_w^{s,q}, \quad \text{for all } s, t, q \in (0, 1) \\ \Gamma_w^{q,q} &= \text{Id}, \quad \text{for all } q \in (0, 1). \end{aligned}$$

Moreover, for all $q, s \in (0, 1)$ and $w \in W$, the following diagram commutes

$$\begin{array}{ccc} B_w^q & \xrightarrow{\Gamma_w^{s,q}} & B_w^s \\ \eta_w^q \circ p_w \downarrow & & \downarrow \eta_w^s \circ p_w \\ \prod_{v \triangleleft w} B_{v, \chi_v^w}^q & \xrightarrow{\prod_{v \triangleleft w} (\Gamma_v^{s,q} \otimes \iota)} & \prod_{v \triangleleft w} B_{v, \chi_v^w}^s \end{array} \quad (57)$$

where η_w^q and χ_v^w are as in Proposition 14. The $*$ -isomorphisms $\Gamma_w^{s,q}$ are also continuous in the point-norm topology in the sense that, for fixed $q \in (0, 1)$ and $y \in B_w^q$, the function $s \in (0, 1) \mapsto \Gamma_w^{s,q}(y) \in \mathcal{B}(H_w)$ is continuous.

(ii) If $\chi^q : \mathbb{C}[G]_q \rightarrow C(X)$, $q \in (0, 1)$, are commutative q -independent $*$ -homomorphisms, then the $*$ -isomorphism $\Gamma_w^{s,q} \otimes \iota : B_w^q \otimes C(X) \rightarrow B_w^s \otimes C(X)$ restricts to a $*$ -isomorphism

$$\Gamma_w^{s,q} \otimes \iota : B_{w, \chi}^q \longrightarrow B_{w, \chi}^s.$$

Proof. We will prove (i) and (ii) simultaneously using induction on $k = \ell(w)$, starting at $k = 0$.

If $\ell(w) = 0$, then $w = e$. As $\pi_e = \epsilon_q$, we have $B_e = \mathbb{C}$. By the definition of a q -independent commutative $*$ -homomorphism $\chi^q = \zeta \circ \tau_q$, and hence

$$\chi^q(C(G)_q) = \zeta(\tau_q(C(G)_q)) = \zeta(C(T)).$$

So, (i) and (ii) hold in the case of $k = 0$ with $\Gamma_e^{s,q} = \text{Id}_{\mathbb{C}}$.

Assume now that (i) and (ii) hold for all $v \in W$ of length $\ell(v) < k$. By Proposition 14 we have a $*$ -homomorphism

$$\partial_1^q := \eta_w^q \circ p_w : B_w^q \longrightarrow B_w^q / \mathcal{K}_w \longrightarrow \prod_{v \triangleleft w} B_{v, \chi_v^w}^q \quad (58)$$

such that the image is isomorphic to $B_w^q / \mathcal{K}_w = p_w(B_w^q) \subseteq \mathbb{Q}(H_w)$. For $v \triangleleft w$, we have

$$((\eta_v^q \circ p_v) \otimes \iota) \circ (\pi_v^q \boxtimes \chi_v^w) = \prod_{\sigma \triangleleft v} (\pi_\sigma^q \boxtimes (\chi_\sigma^v \boxtimes \chi_v^w)) \sim \prod_{\sigma \triangleleft v} (\pi_\sigma \boxtimes \chi_\gamma) \quad (59)$$

where $\sigma \xrightarrow{\gamma} w$ is the path $\sigma \triangleleft v \triangleleft w$. Taking the product over all $v \triangleleft w$ we get from (59) a $*$ -homomorphism

$$\partial_2^q := \prod_{v \triangleleft w} (\eta_v^q \circ p_v) \otimes \iota : \prod_{v \triangleleft w} B_{v, \chi_v^w}^q \longrightarrow \prod_{\sigma \rightsquigarrow w}^{(2)} B_{v, \chi_\gamma}^q, \quad (60)$$

where the product $\prod_{\sigma \rightsquigarrow w}^{(2)}$ is indexed over all $\sigma \triangleleft^{(2)} w$ and all possible paths $\sigma \xrightarrow{\gamma} w$.

It follows from Lemma 16 that the kernel of ∂_2^q is equal to $\prod_{v \triangleleft w} \mathcal{K}_v \otimes C(T_v^w)$. If we iterate (60), then we get a sequence of $*$ -homomorphisms

$$B_w^q \xrightarrow{\partial_1^q} \prod_{v \triangleleft w} B_{v, \chi_v^w}^q \xrightarrow{\partial_2^q} \prod_{v \rightsquigarrow w}^{(2)} B_{v, \chi_\gamma}^q \xrightarrow{\partial_3^q} \dots \xrightarrow{\partial_{k-1}^q} \prod_{v \rightsquigarrow w}^{(k-1)} B_{v, \chi_\gamma}^q \xrightarrow{\partial_k^q} \prod_{e \rightsquigarrow w} C(T_\gamma), \quad (61)$$

where the product $\prod_{v \rightsquigarrow w}^{(i)}$ ranges over all elements $v \in W$ such that $v \triangleleft^{(i)} w$ and over all possible paths $v \xrightarrow{\gamma} w$. In the last product, $e \in W$ is the identity element and we suppress the upper index (k) as it is unnecessary in this case. In general, when we have a fixed element $v \in W$ such that $v \leq w$, then $v \rightsquigarrow w$ denotes the set of all possible paths $v \xrightarrow{\gamma} w$. As an example, for $v \in W$, we write $\prod_{v \rightsquigarrow w} B_{v, \chi_\gamma}$ to mean that the product ranges over all possible paths $v \xrightarrow{\gamma} w$. Similarly, we write $\prod_{v \triangleleft^{(i)} w}$ to mean that the product is over all $v \in W$ such that $v \triangleleft^{(i)} w$. Similar notations will also be used for direct sums, etc. Clearly, by Lemma 16, for every $i = 1, \dots, k$, we have

$$\ker \partial_{i+1}^q = \prod_{v \rightsquigarrow w}^{(i)} \mathcal{K}_v \otimes C(T_\gamma). \quad (62)$$

Moreover, the commutative C^* -algebra $\prod_{e \rightsquigarrow w} C(T_\gamma)$ does not depend on q . For $i = 1, \dots, k$, we let

$$\begin{aligned} \partial_i^q \circ \dots \circ \partial_1^q &=: \Psi_i^q : B_w^q \longrightarrow \prod_{v \rightsquigarrow w}^{(i)} B_{v, \chi_\gamma}^q \\ \partial_k^q \circ \dots \circ \partial_1^q &=: \Psi_k^q : B_w^q \longrightarrow \prod_{e \rightsquigarrow w} C(T_\gamma) \end{aligned}$$

be the composition of $*$ -homomorphisms in (61). By iteration of (42) and (54), we have for any $a \in \mathbb{C}[G]_q$

$$(\Psi_i^q \circ \pi_w^q)(a) = \bigoplus_{v \rightsquigarrow w}^{(i)} (\pi_v^q \boxtimes \chi_\gamma)(a), \quad i = 1, \dots, k. \quad (63)$$

By induction, for all $v \in W$ such that $l(v) < k$, we have a $*$ -isomorphism

$$\Gamma_v^{s,q} : B_v^q \longrightarrow B_v^s$$

such that $\Gamma_v^{s,q}(\mathcal{K}_v) = \mathcal{K}_v$ and the following diagram is commutative

$$\begin{array}{ccc} B_v^q & \xrightarrow{\Gamma_v^{s,q}} & B_v^s \\ \eta_v^q \circ p_v \downarrow & & \downarrow \eta_v^s \circ p_v \\ \prod_{\sigma \triangleleft v} B_{\sigma, \chi_\sigma^v}^q & \xrightarrow{\prod_{\sigma \triangleleft v} (\Gamma_\sigma^{s,q} \otimes \iota)} & \prod_{\sigma \triangleleft v} B_{\sigma, \chi_\sigma^v}^s \end{array}$$

It follows that for every $i = 1, \dots, k-1$, we have $*$ -isomorphisms

$$\prod_{v \rightsquigarrow w}^{(i)} (\Gamma_v^{s,q} \otimes \iota) : \prod_{v \rightsquigarrow w} B_{v, \chi_\gamma}^q \longrightarrow \prod_{v \rightsquigarrow w} B_{v, \chi_\gamma}^s \quad (64)$$

that maps $\prod_{v \rightsquigarrow w}^{(i)} \mathcal{K}_v \otimes C(T_\gamma)$ into itself and such that the following diagrams are commutative

$$\begin{array}{ccccc} B_w^q & \xrightarrow{\Psi_i^q} & \prod_{v \rightsquigarrow w}^{(i)} B_{v, \chi_\gamma}^q & \xrightarrow{\prod_{v \rightsquigarrow w}^{(i)} (\Gamma_v^{s,q} \otimes \iota)} & \prod_{v \rightsquigarrow w}^{(i)} B_{v, \chi_\gamma}^s \xleftarrow{\Psi_i^s} B_w^s \\ & \searrow \Psi_{i+1}^q & \downarrow \partial_{i+1}^q & & \downarrow \partial_{i+1}^s \\ & & \prod_{v \rightsquigarrow w}^{(i+1)} B_{v, \chi_\gamma}^q & \xrightarrow{\prod_{v \rightsquigarrow w}^{(i+1)} (\Gamma_v^{s,q} \otimes \iota)} & \prod_{v \rightsquigarrow w}^{(i+1)} B_{v, \chi_\gamma}^s \\ & & & & \nwarrow \Psi_{i+1}^s \end{array} \quad (65)$$

$$\begin{array}{ccccc} B_w^q & \xrightarrow{\Psi_{k-1}^q} & \prod_{v \rightsquigarrow w}^{(k-1)} B_{v, \chi_\gamma}^q & \xrightarrow{\prod_{v \rightsquigarrow w}^{(k-1)} (\Gamma_v^{s,q} \otimes \iota)} & \prod_{v \rightsquigarrow w}^{(k-1)} B_{v, \chi_\gamma}^s \xleftarrow{\Psi_{k-1}^s} B_w^s \\ & \searrow \Psi_k^q & \downarrow \partial_k^q & & \downarrow \partial_k^s \\ & & \prod_{e \rightsquigarrow w} C(T_\gamma) & \xrightarrow{\text{Id}} & \prod_{e \rightsquigarrow w} C(T_\gamma) \\ & & & & \nwarrow \Psi_k^s \end{array} \quad (66)$$

The idea is now to show that $\prod_{v \rightsquigarrow w}^{(i)} (\Gamma_v^{s,q} \otimes \iota)$ restricts to a $*$ -isomorphism between $\Psi_i^q(B_w^q)$ and $\Psi_i^s(B_w^s)$ for $i = 1, \dots, k$. We prove this by ‘climbing the ladder’ (65),

using induction on i , starting at $i = k$ (i.e the case (66)), and then we count down to $i = 1$.

The statement is clear at k , since by the q -independence of χ_γ and the fact that $\tau_q(C(G)_q) = C(T) = \tau_s(C(G)_s)$, we have

$$\Psi_k^q(B_w^q) = (\bigoplus_{e \rightsquigarrow w} \chi_\gamma)(C(G)_q) = (\bigoplus_{e \rightsquigarrow w} \chi_\gamma)(C(G)_s) = \Psi_k^s(B_w^s).$$

Assume now that the statement holds for $i + 1$. Consider $x \in \Psi_i^q(B_w^q)$. Then

$$\partial_{i+1}^q(x) \in \Psi_{i+1}^q(B_w^q),$$

and hence by induction

$$\left(\prod_{v \rightsquigarrow w}^{(i+1)} (\Gamma_v^{s,q} \otimes \iota) \right) (\partial_{i+1}^q(x)) \in \Psi_{i+1}^s(B_w^s).$$

But by the commutivity of the diagrams (65)–(66), this element is also equal to

$$\partial_{i+1}^s \left(\prod_{v \rightsquigarrow w}^{(i)} (\Gamma_v^{s,q} \otimes \iota)(x) \right) \in \Psi_{i+1}^s(B_w^s)$$

from which it follows, by (62), that

$$\prod_{v \rightsquigarrow w}^{(i)} (\Gamma_v^{s,q} \otimes \iota)(x) \in \Psi_i^s(B_w^s) + \prod_{v \rightsquigarrow w}^{(i)} \mathcal{H}_v \otimes C(T_\gamma) \quad (67)$$

and thus

$$\prod_{v \rightsquigarrow w}^{(i)} (\Gamma_v^{s,q} \otimes \iota)(x) = y + c, \quad y \in \Psi_i^s(B_w^s), \quad c \in \prod_{v \rightsquigarrow w}^{(i)} \mathcal{H}_v \otimes C(T_\gamma). \quad (68)$$

We show that actually $c \in \Psi_i^s(B_w^s)$. For a fixed $v \in W$ such that $v \triangleleft^{(i)} w$, we can embed

$$\mathcal{B}(H_v) \otimes C(T_v^w) \subseteq \prod_{v \rightsquigarrow w} \mathcal{B}(H_v) \otimes C(T_\gamma) \quad (69)$$

via the injective $*$ -homomorphisms $C(T_v^w) \rightarrow \prod_{v \rightsquigarrow w} C(T_\gamma)$ coming from the inclusions $T_\gamma \subseteq T_v^w$ and, by the definition of T_v^w , that $T_v^w = \cup_{v \rightsquigarrow w} T_\gamma$. Thus the embedding (69) is on simple tensors given by

$$x \otimes f \mapsto \prod_{v \rightsquigarrow w} (x \otimes f|_{T_\gamma}), \quad x \in \mathcal{B}(H_v), \quad f \in C(T_v^w),$$

where $f|_{T_\gamma}$ denoted the restriction of $f \in C(T_v^w)$ to the subset $T_\gamma \subseteq T_v^w$. Moreover, we have the embedding

$$\mathcal{H}_v \otimes C(T_v^w) \subseteq \prod_{v \rightsquigarrow w} \mathcal{B}(H_v) \otimes C(T_\gamma) \quad (70)$$

coming from (69). Using this embedding, we clearly have, for fixed $v \triangleleft^{(i)} w$, that

$$\bigoplus_{v \rightsquigarrow w} (\pi_v^q \boxtimes \chi_\gamma) : \mathbb{C}[G]_q \longrightarrow \mathcal{B}(H_v) \otimes C(T_v^w) \subseteq \prod_{v \rightsquigarrow w} \mathcal{B}(H_v) \otimes C(T_\gamma) \quad (71)$$

and that, as in (54), we can identify $\bigoplus_{v \rightsquigarrow w} (\pi_v^q \boxtimes \chi_\gamma) \sim \pi_v^q \boxtimes \chi_v^w$. It then follows from Lemma 16, that under the embeddings (69) and (70) we have

$$\begin{aligned} \mathcal{K}_v \otimes C(T_v^w) &\subseteq \overline{\bigoplus_{v \rightsquigarrow w} (\pi_v^q \boxtimes \chi_\gamma)(\mathbb{C}[G]_q)} \subseteq \mathcal{B}(H_v) \otimes C(T_v^w) \\ &\subseteq \prod_{v \rightsquigarrow w} \mathcal{B}(H_v) \otimes C(T_\gamma). \end{aligned} \quad (72)$$

Moreover, note that the left-hand sides (69) and (70) are clearly invariant under the homomorphism $\prod_{v \rightsquigarrow w} \Gamma_v^{s,q} \otimes \iota$. We can now use Proposition 17 and (63) to see that if we take the product of (72), ranging over all $v \triangleleft^{(i)} w$, then

$$\prod_{v \triangleleft^{(i)} w} \mathcal{K}_v \otimes C(T_v^w) \subseteq \Psi_i^q(B_w^q) \subseteq \prod_{v \triangleleft^{(i)} w} \mathcal{B}(H_v) \otimes C(T_v^w) \subseteq \prod_{v \rightsquigarrow w}^{(i)} \mathcal{B}(H_v) \otimes C(T_\gamma). \quad (73)$$

As $\prod_{v \rightsquigarrow w}^{(i)} (\Gamma_v^{s,q} \otimes \iota)$ clearly fixes the two sub-algebras on either side of $\Psi_i^q(B_w^q)$, it follows from (68) that

$$\prod_{v \rightsquigarrow w}^{(i)} (\Gamma_v^{s,q} \otimes \iota)(x) - y = c \in \prod_{v \triangleleft^{(i)} w} \mathcal{K}_v \otimes C(T_v^w) \subseteq \Psi_i^s(B_w^s).$$

From this, it follows that

$$\prod_{v \rightsquigarrow w}^{(i)} (\Gamma_v^{s,q} \otimes \iota)(\Psi_i^q(B_w^q)) \subseteq \Psi_i^s(B_w^s), \quad q, s \in (0, 1). \quad (74)$$

But as

$$\prod_{v \rightsquigarrow w}^{(i)} (\Gamma_v^{s,q} \otimes \iota) \circ \prod_{v \rightsquigarrow w}^{(i)} (\Gamma_v^{q,s} \otimes \iota) = \text{Id}, \quad q, s \in (0, 1)$$

we must have equality in (74).

Thus, we have an isomorphism

$$B_w^q / \mathcal{K}_w \cong B_w^s / \mathcal{K}_w, \quad q, s \in (0, 1)$$

via the $*$ -isomorphism

$$\mathcal{L}^{s,q} := (\eta_w^s)^{-1} \circ \left(\prod_{v \triangleleft w} (\Gamma_v^{s,q} \otimes \iota) \right) \circ \eta_w^q. \quad (75)$$

However, to be able to use Lemma 2 to conclude that the C^* -algebras B_w^q , $q \in (0, 1)$ are all isomorphic, we must also show that $\mathcal{L}^{s,q}$ are continuous in the point-norm topology, i.e. that for a fixed $q \in (0, 1)$ and an element $y \in B_w^q / \mathcal{K}_w$, we have a continuous function

$$s \in (0, 1) \rightarrow \mathbb{Q}(H_w), s \mapsto \mathcal{L}^{s,q}(y). \quad (76)$$

By a classical approximation argument, it is enough to prove this for the dense $*$ -subalgebra $(p_w \circ \pi_w^q)(\mathbb{C}[G]_q)$. By Lemma 18 we have invertible coalgebra morphisms $\theta^q : \mathbb{C}[G] \rightarrow \mathbb{C}[G]_q$ such that for fixed $f \in \mathbb{C}[G]$, the function $q \in (0, 1) \mapsto \pi_w^q(\theta^q(f)) \in \mathcal{B}(H_w)$ is continuous. Thus the function

$$q \in (0, 1) \mapsto (p_w \circ \pi_w^q)(\theta^q(f)) \in \mathcal{B}(H_w) / \mathcal{K}_w = \mathbb{Q}(H_w)$$

is also continuous. Let us write

$$F^q := (p_w \circ \pi_w^q)(\theta^q(f)) \in \mathbb{Q}(H_w).$$

By induction, the function

$$s \in (0, 1) \mapsto \left(\prod_{v \triangleleft w} (\Gamma_v^{s,q} \otimes \iota) \right) (\eta_w^q(F^q))$$

is continuous and $(\prod_{v \triangleleft w} (\Gamma_v^{q,q} \otimes \iota)) (\eta_w^q(F^q)) = \eta_w^q(F^q)$. Notice that by the definition of η_w^q and (46), we have

$$\eta_w^q(F^q) = \bigoplus_{v \triangleleft w} (\pi_v^q \boxtimes \chi_v^w)(\theta^q(f))$$

and thus by Lemma 18, the function

$$q \in (0, 1) \mapsto \eta_w^q(F^q) \in \prod_{v \triangleleft w} \mathcal{B}(H_v) \otimes C(T_v^w)$$

is continuous. It follows that for all $\epsilon > 0$ we have

$$\begin{aligned} & \left\| \left(\prod_{v \triangleleft w} (\Gamma_v^{s,q} \otimes \iota) \right) (\eta_w^q(F^q)) - (\eta_w^s(F^s)) \right\| \\ & \leq \left\| \left(\prod_{v \triangleleft w} (\Gamma_v^{s,q} \otimes \iota) \right) (\eta_w^q(F^q)) - \eta_w^q(F^q) \right\| + \left\| \eta_w^q(F^q) - \eta_w^s(F^s) \right\| < \epsilon \end{aligned}$$

for $|s - q| < \delta_1$, if $\delta_1 > 0$ is made small enough. If we apply the $*$ -isomorphism $(\eta_w^s)^{-1}$, we get

$$\|\mathcal{L}^{s,q}(F^q) - F^s\| < \epsilon, \text{ for } |s - q| < \delta_1 \quad (77)$$

and thus it follows that there is a $0 < \delta \leq \delta_1$, such that

$$\begin{aligned} & \|\mathcal{L}^{s,q}(F^q) - F^q\| \\ & \leq \|\mathcal{L}^{s,q}(F^q) - F^s\| + \|F^s - F^q\| < 2\epsilon \end{aligned}$$

when $|s - q| < \delta$. We can now apply Lemma 2 to get an inner $*$ -automorphism $\Gamma_w^{s,q} : \mathcal{B}(H_w) \rightarrow \mathcal{B}(H_w)$ that restricts to a $*$ -isomorphism $B_w^q \rightarrow B_w^s$, that is continuous in the point-set topology. That the diagram (57) commutes follows from the commutivity of (8) and the way $\mathcal{L}^{s,q}$ was defined. Clearly, the compact operators are invariant under $\Gamma_w^{s,q}$.

The case (ii). We prove it first for $\tau_q : \mathbb{C}[G]_q \rightarrow C(T)$ (see (34)). We combine the inclusion $B_{w,\tau_q}^q \hookrightarrow B_w^q \otimes C(T)$ with the sequence (61) by tensoring all the components with $C(T)$ in the following way

$$\begin{aligned} B_{w,\tau_q}^q &\hookrightarrow B_w^q \otimes C(T) \xrightarrow{\partial_1 \otimes \iota} \left(\prod_{v \triangleleft w} B_{v,\chi_v^w}^q \right) \otimes C(T) \xrightarrow{\partial_2 \otimes \iota} \left(\prod_{v \rightsquigarrow w}^{(2)} B_{v,\chi_v}^q \right) \otimes C(T) \\ &\xrightarrow{\partial_3 \otimes \iota} \dots \xrightarrow{\partial_{k-1} \otimes \iota} \left(\prod_{v \rightsquigarrow w}^{(k-1)} B_{v,\chi_v}^q \right) \otimes C(T) \xrightarrow{\partial_k \otimes \iota} \left(\prod_{e \rightsquigarrow w} C(T_\gamma) \right) \otimes C(T). \end{aligned} \quad (78)$$

If we define Ψ_i^q as before, then similar to (63), we have

$$(\Psi_i^q \otimes \iota) \circ (\pi_w^q \boxtimes \tau_q) = \left(\bigoplus_{v \rightsquigarrow w}^{(i)} (\pi_v^q \boxtimes \chi_\gamma) \right) \boxtimes \tau_q. \quad (79)$$

We can proceed exactly as before, using the commutative diagrams (65) and (66) (now tensored by $C(T)$), by induction on $i = 1, \dots, k$, starting at k . Clearly the images in $(\prod_{e \rightsquigarrow w} C(T_\gamma)) \otimes C(T)$ are the same, as the commutative $*$ -representation is q -independent. Assuming that

$$\left(\prod_{v \rightsquigarrow w}^{(i+1)} (\Gamma_v^{s,q} \otimes \iota) \right) \otimes \iota : (\Psi_{i+1}^q \otimes \iota) (B_{w,\tau_q}^q) \longrightarrow (\Psi_{i+1}^s \otimes \iota) (B_{w,\tau_s}^s)$$

is an $*$ -isomorphism gives for $x \in (\Psi_i^q \otimes \iota) (B_{w,\tau_q}^q)$, that

$$\left(\left(\prod_{v \rightsquigarrow w}^{(i)} (\Gamma_v^{s,q} \otimes \iota) \right) \otimes \iota \right) (x) \in (\Psi_i^s \otimes \iota) (B_{w,\tau_s}^s) + \left(\prod_{v \rightsquigarrow w}^{(i)} \mathcal{K}_v \otimes C(T_\gamma) \right) \otimes C(T). \quad (80)$$

The rest of the argument follows in a similar fashion as for B_w^q : we find an embedding

$$\prod_{v \triangleleft^{(i)} w} \mathcal{B}(H_v) \otimes C(T) \subseteq \left(\prod_{v \rightsquigarrow w}^{(i)} \mathcal{B}(H_v) \otimes C(T_\gamma) \right) \otimes C(T) \quad (81)$$

such that,

- (i) this subalgebra is invariant with respect to the map $(\prod_{v \rightsquigarrow w}^{(i)} (\Gamma_v^{s,q} \otimes \iota)) \otimes \iota$,
- (ii) we have

$$\prod_{v \triangleleft^{(i)} w} \mathcal{K}_v \otimes C(T) = \prod_{v \triangleleft^{(i)} w} \mathcal{B}(H_v) \otimes C(T) \cap \left(\prod_{v \rightsquigarrow w}^{(i)} \mathcal{K}_v \otimes C(T_\gamma) \right) \otimes C(T) \quad (82)$$

(iii) and the following inclusions holds

$$\prod_{v \triangleleft^{(i)} w} \mathcal{K}_v \otimes C(T) \subseteq (\Psi_i^s \otimes \iota)(B_{w, \tau_s}^s) \subseteq \prod_{v \triangleleft^{(i)} w} \mathcal{B}(H_v) \otimes C(T). \quad (83)$$

Clearly, this implies

$$\left(\left(\prod_{v \rightsquigarrow w}^{(i)} (\Gamma_v^{s, q} \otimes \iota) \right) \otimes \iota \right) (x) \in (\Psi_i^s \otimes \iota)(B_{w, \tau_s}^s).$$

To do this, we use the natural isomorphism

$$\left(\prod_{v \rightsquigarrow w}^{(i)} \mathcal{B}(H_v) \otimes C(T_\gamma) \right) \otimes C(T) \cong \prod_{v \rightsquigarrow w}^{(i)} \mathcal{B}(H_v) \otimes C(T_\gamma) \otimes C(T).$$

Notice that for every $v \rightsquigarrow w$, we have an embedding $C(T) \subseteq C(T_\gamma) \otimes C(T)$ determined by $C(T_\gamma) \otimes C(T) \cong C(T_\gamma \times T)$ and the multiplication map $T_\gamma \times T \rightarrow T$. As $\tau_q : \mathbb{C}[G]_q \rightarrow \mathbb{C}[T]$ is a morphism of Hopf $*$ -algebras (hence compatible with the multiplication in T), it follows that we have a $*$ -homomorphism

$$\pi_v^q \boxtimes \chi_\gamma \boxtimes \tau_q : C(G)_q \rightarrow \mathcal{B}(H_v) \otimes C(T) \subseteq \mathcal{B}(H_v) \otimes C(T_\gamma) \otimes C(T).$$

We can then, for fixed $v \triangleleft^{(i)} w$, embed diagonally

$$\mathcal{B}(H_v) \otimes C(T) \subseteq \prod_{v \rightsquigarrow w} \mathcal{B}(H_v) \otimes C(T_\gamma) \otimes C(T).$$

By taking the product over all $v \triangleleft^{(i)} w$, we get an embedding (81) such that (83) holds (the first inclusion follows from Proposition (17)). Clearly, we also have (82). Thus, it follows that $(\Gamma_w^{s, q} \otimes \iota)(B_{w, \tau_q}^q) = B_{w, \tau_q}^s$, since this is the case $i = 1$.

This implies the general case: let $\chi^q : C(G)_q \rightarrow C(X)$ be commutative q -independent $*$ -homomorphisms and $\zeta : C(T) \rightarrow C(X)$ the $*$ -homomorphism such that $\chi^q = \zeta \circ \tau_q$. Then $\iota \otimes \zeta$ is a surjective $*$ -homomorphism $B_{w, \tau_q}^q \rightarrow B_{v, \chi^q}^q$. As $\Gamma_w^{s, q} \otimes \iota$ is a $*$ -isomorphism $B_{w, \tau_q}^q \rightarrow B_{w, \tau_s}^s$, we have

$$\begin{aligned} (\Gamma_w^{s, q} \otimes \iota)(B_{w, \chi^q}^q) &= (\Gamma_w^{s, q} \otimes \iota) \circ (\iota \otimes \zeta)(B_{w, \tau_q}^q) = (\iota \otimes \zeta) \circ (\Gamma_w^{s, q} \otimes \iota)(B_{w, \tau_q}^q) \\ &= (\iota \otimes \zeta)(B_{w, \tau_s}^s) = B_{w, \chi^s}^s. \end{aligned}$$

□

Corollary 20. *The universal enveloping C^* -algebras of $\mathbb{C}[G]_q$ are isomorphic for all $q \in (0, 1)$. These isomorphisms are equivariant with respect to the right-action of T .*

Proof. If $\omega \in W$ is the unique element of longest length in the Weyl group and $\tau_q : C(G)_q \rightarrow C(T)$ the commutative q -independent $*$ -homomorphism coming from the embedding of the maximal torus $T \subseteq G$, then it follows from Theorem 6 that any irreducible $*$ -representation of $C(G)_q$ must factor through $\pi_\omega^q \boxtimes \tau_q$. Thus $B_{\omega, \tau_q}^q \cong C(G)_q$. The q -independence follows from Theorem 19(ii). For $x \in C(G)_q$ and $t \in T$ we have

$$\begin{aligned} (\pi_\omega^q \boxtimes \tau_q)(R_t(x)) \\ = (\iota \otimes \iota \otimes \text{ev}_t) \circ (\iota \otimes \Delta_T)((\pi_\omega^q \boxtimes \tau_q)(x)) \end{aligned}$$

and thus equivariance with respect to the right-action follows as the isomorphism $B_{\omega, \tau_q}^q \rightarrow B_{\omega, \tau_s}^s$ is of the form $\Gamma_{\omega}^{s,q} \otimes \iota$. \square

Acknowledgements. Open access funding provided by Chalmers University of Technology. The author is grateful to Mahmood Alagmandan and Lyudmila Turowska for helpful discussions and comments. The author was partially supported by GS Magnusons fond, Grant Number MG2018-0083, as well as Knut och Alice Wallenbergs Stiftelse.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

References

1. Bjorner, A., Brenti, F.: Combinatorics of Coxeter Groups. Graduate Texts in Mathematics, vol. 231. Springer, New York (2005)
2. Davidson, K.: C^* -algebras by Example. Fields Institute Monographs, vol. 6. American Mathematical Society, Providence, RI (1996)
3. Drinfeld, V.G.: Hopf algebras and the quantum Yang–Baxter equation (Russian). Dokl. Akad. Nauk SSSR **283**(5), 1060–1064 (1985)
4. Drinfeld, V.G.: Quantum groups. In: Proceedings of the International Congress of Mathematicians, vol. 1, 2, pp. 798–820. Berkeley, CA (1986)
5. Jimbo, M.: A q -analogue of $U(\mathfrak{g})$ and the Yang–Baxter equation. Lett. Math. Phys. **10**(1), 63–69 (1985)
6. Klimyk, A., Schmüdgen, K.: Quantum Groups and Their Representations. Texts and Monographs in Physics. Springer, Berlin (1997)
7. Korogodski, L.I., Soibelman, Y.S.: Algebra of Functions on Quantum Groups: Part 1. Mathematical Surveys and Monographs, vol. 56. American Mathematical Society, Providence, RI (1998)
8. Nagy, G.: A rigidity property for quantum $SU(3)$ groups. In: Brylinski, J.L., Brylinski, R., Nistor, V., Tsygan, B., Xu, P. (eds.) Advances in Geometry. Progress in Mathematics, vol. 172, pp. 297–336. Birkhäuser, Boston, MA (1999)
9. Nagy, G.: Deformation quantization and K-theory. In: Coburn, L.A., Rieffel, M.A. (eds.) Perspectives on Quantization (South Hadley, MA, 1996). Contemporary Mathematics, vol. 214, pp. 111–134. American Mathematical Society, Providence, RI (1998)
10. Neshveyev, S., Tuset, L.: Quantized algebras of functions on homogeneous spaces with Poisson stabilizers. Commun. Math. Phys. **312**(1), 223–250 (2012)
11. Neshveyev, S., Tuset, L.: K-homology class of the Dirac operator on a compact quantum group. Doc. Math. **16**, 767–780 (2011)
12. Stokman, J.V., Dijkhuizen, M.S.: Quantized flag manifolds and irreducible $*$ -representations. Commun. Math. Phys. **203**(2), 297–324 (1999)
13. Woronowicz, S.L.: Compact matrix pseudogroups. Commun. Math. Phys. **111**(4), 613–665 (1987)
14. Woronowicz, S.L.: Twisted $SU(2)$ group. An example of a noncommutative differential calculus. Publ. Res. Inst. Math. Sci. **23**(1), 117–181 (1987)

Communicated by Y. Kawahigashi