Thermovoltage in quantum dots with attractive interaction

Downloaded from: <https://research.chalmers.se>, 2023-04-26 12:47 UTC

Citation for the original published paper (version of record):

Schulenburg, J., Wegewijs, M., Splettstoesser, J. (2020). Thermovoltage in quantum dots with attractive interaction. *Applied Physics Letters*, 116(24). <http://dx.doi.org/10.1063/5.0008866>

N.B. When citing this work, cite the original published paper.

Thermovoltage in quantum dots with attractive interaction

Cite as: Appl. Phys. Lett. **116**, 243103 (2020); <https://doi.org/10.1063/5.0008866>

Submitted: 26 March 2020 . Accepted: 31 May 2020 . Published Online: 16 June 2020

Jens Schulenborg, Maarten R. Wegewijs, and Janine Splettstoesser

View Online

Export Citation

CrossMark

ARTICLES YOU MAY BE INTERESTED IN

[An elementary photo-thermoelectric transistor: Experimental demonstration](#)

Applied Physics Letters **116**, 243501 (2020); <https://doi.org/10.1063/5.0010264>

[Niobium Dayem nano-bridge Josephson gate-controlled transistors](#)

Applied Physics Letters **116**, 242601 (2020); <https://doi.org/10.1063/5.0011304>

[Spin-orbit torque field-effect transistor \(SOTFET\): Proposal for a magnetoelectric memory](#)

Applied Physics Letters **116**, 242405 (2020); <https://doi.org/10.1063/5.0002909>

David Daughton, PhD
Applications Scientist
Lake Shore Cryotronics

Houston Fortney
Development Engineer
Lake Shore Cryotronics

WEBINAR

A New Concept in Semiconductor
Material/Device Characterization

Combining DC and AC Sourcing and Measuring

Watch Now

Thermovoltage in quantum dots with attractive interaction

Cite as: Appl. Phys. Lett. **116**, 243103 (2020); doi: [10.1063/5.0008866](https://doi.org/10.1063/5.0008866)

Submitted: 26 March 2020 · Accepted: 31 May 2020 ·

Published Online: 18 June 2020

View Online

Export Citation

CrossMark

Jens Schulenburg,^{1,2,a)} Maarten R. Wegewijs,^{3,4} and Janine Splettstoesser²

AFFILIATIONS

¹Center for Quantum Devices, Niels Bohr Institute, University of Copenhagen, 2100 Copenhagen, Denmark

²Department of Microtechnology and Nanoscience, Chalmers University of Technology, 41296 Göteborg, Sweden

³Institute for Theory of Statistical Physics, RWTH Aachen, 52056 Aachen, Germany

⁴Peter Grünberg Institut and JARA, Forschungszentrum Jülich, 52425 Jülich, Germany

^{a)}Author to whom correspondence should be addressed: jens.schulenburg@nbi.ku.dk

ABSTRACT

We study the linear and nonlinear thermovoltage of a quantum dot with effective *attractive* electron–electron interaction and weak, energy-dependent tunnel coupling to electronic contacts. Remarkably, we find that the thermovoltage shows signatures of *repulsive* interaction, which can be rationalized. These thermovoltage characteristics are robust against large potential and temperature differences well into the nonlinear regime, which we expect can be demonstrated in current state-of-the-art experiments. Furthermore, under nonlinear operation, we find extended regions of large power production at efficiencies on the order of the Curzon–Ahlborn bound interrupted only by a characteristic sharp dip.

Published under license by AIP Publishing. <https://doi.org/10.1063/5.0008866>

Recently, different types of devices with an effectively attractive electron–electron interaction¹ have been experimentally investigated² and quantum dot structures with attractive onsite-interaction have also been realized.^{3,4} In these quantum dots, signatures of pair tunneling^{5–7} induced by the attractive onsite interaction could be identified in transport properties.

In the present letter, we predict surprising features in the thermovoltage of such quantum dots. We show that the linear-response thermovoltage—the Seebeck coefficient—shows signatures at quantum-dot level positions that are characteristic of Coulomb oscillations due to *repulsive* onsite interaction. We rationalize this fact and show that it can be exploited in an analysis of how these features are modified under various realistic experimental conditions. The discussed effects are highly relevant for the characterization of attractive systems, which has only recently started.

Simultaneously, there has been significant progress in investigating linear and nonlinear response thermoelectrics in quantum dot devices (see Refs. 8–20 and references therein). These are of interest for on-chip energy harvesting, and their Seebeck coefficient is a key parameter to characterize them. Our analysis instead reveals the thermoelectric properties of systems with *strong attractive* electron–electron interaction. We also explicitly address energy-dependent tunnel couplings between the dot and the environment as energy filters in addition to the quantum-dot

levels. Efficient nanoscale thermoelectrics, in particular three-terminal energy harvesters,^{10,21–24} crucially rely on this energy-dependent coupling. In this Letter, we characterize the performance of such quantum dots with attractive interaction as steady-state heat engines and find extended regions of large power production and efficiency.

Finally, the thermoelectric response of *repulsive* quantum dots^{16,19} has successfully been analyzed using a mapping based on a fermionic duality relation,²⁵ providing simple analytical formulas. Here, this relation enables us to explain the thermoelectric response of a quantum dot with attractive interaction in terms of the well understood physics of a repulsive dot. This simple description can serve as a guide for future experiments.

The quantum dot of interest is sketched in Fig. 1(a). It is modeled as a single spin-degenerate level, with an *attractive* electron–electron interaction. We assume the level spacing to be large compared to any other energy scale relevant for transport, such as voltage bias and temperatures; indeed, recent experimental realizations of quantum dots with attractive interactions have been well explained in terms of such a model.⁴ The isolated dot is then described by the Hamiltonian,

$$H = \bar{\epsilon}N - |U|N_{\uparrow}N_{\downarrow}, \quad (1)$$

with $\bar{\epsilon}$ being the single-level energy, $N = N_{\uparrow} + N_{\downarrow}$ the local particle-number operator with spin-resolved components $N_{\uparrow}$ and $N_{\downarrow}$, and

FIG. 1. (a) Sketch of a quantum dot with attractive interaction. (b) Seebeck coefficient S (blue solid) and its derivative (green dashed) and (c) equilibrium charge n and its dual n_d , as a function of dot level $\varepsilon = \tilde{\varepsilon} - \mu$. We take $T = |U|/10$ and $\Gamma_{L,R}$ that are energy-independent.

$|U| = -U$ the interaction strength. The quantum dot is tunnel-coupled to two electronic reservoirs, $\alpha = L, R$, with electrochemical potentials $\mu_L = \mu - V$ and $\mu_R = \mu$ and temperatures $T_L = T + \Delta T$ and $T_R = T$. We assume the experimentally relevant case of weak tunnel rates $\Gamma_\alpha(E) \ll T_\alpha$, implying that pair tunneling is enabled by thermal excitations.^{4,26,27} We allow these rates to strongly depend on the energy of the tunneling process. We set $k_B = \hbar = |e| = 1$.

We analyze the full thermoelectric response of the quantum dot using expressions obtained from a recently established, general *fermionic duality*.^{16,19,25,28} This purely dissipative symmetry, applied to a weak-coupling master-equation description, maps the transport dynamics of fermionic open nonequilibrium systems to those of *dual* systems with sign-inverted local energies, chemical potentials, and energy dependencies of the tunnel couplings, replacing $[\tilde{\varepsilon}, -|U|, \mu_\alpha, \Gamma_\alpha(E)] \rightarrow [-\tilde{\varepsilon}, |U|, -\mu_\alpha, \Gamma_\alpha(-E)]$. In the present case, the remarkable duality enables us to predict *nonequilibrium* effects in the thermoelectric response of the attractive dot of interest by relating them to the properties of the quantum dot with *equilibrium* parameters and attractive interaction as well as to its dual model with *repulsive* interaction.²⁹ The key quantities in this analysis are the equilibrium dot occupation for attractive interaction,

$$n(\varepsilon, -|U|) = \langle N \rangle = \frac{2f(\varepsilon)}{1 + f(\varepsilon) - f(\varepsilon - |U|)}, \quad (2)$$

with $\varepsilon = \tilde{\varepsilon} - \mu$ and $f(x) = [\exp(x/T) + 1]^{-1}$ the Fermi function, and most importantly, the *dual* occupation,¹⁹

$$n_d = n(-\varepsilon, |U|) = \frac{2[1 - f(\varepsilon)]}{1 - f(\varepsilon) + f(\varepsilon - |U|)}, \quad (3)$$

to which the duality assigns a repulsive interaction $|U|$. Figure 1(c) shows both n and n_d as functions of ε .

We start with the linear response of the thermovoltage, for small V and ΔT , and the linear Seebeck coefficient $S = V/\Delta T|_{I=0}$ at vanishing charge current $I = 0$ [see Eq. (10)] across the dot and first consider energy-independent tunnel couplings $\Gamma_\alpha(E) \rightarrow \Gamma_\alpha$. The explicit formula for S has a remarkably simple form¹⁶ in terms of the dual dot occupation (3),

$$S \times T = \varepsilon - |U|(2 - n_d)/2. \quad (4)$$

The consequences of the attractive interaction are shown in Fig. 1(b). On the one hand, we find a linear ε -dependence $S \times T \approx \varepsilon - |U|/2$ around the zero-crossing at $\varepsilon = |U|/2$. This is intuitively expected in analogy to the well-known case of repulsive quantum dots:^{16,30–32} as

reflected by $n(\varepsilon)$ in Fig. 1(c), the attractive dot effectively acts as a single resonance at $\varepsilon = |U|/2$. On the other hand, we find $S \times T \rightarrow \varepsilon - |U|$ for $\varepsilon < 0$ and $S \times T \rightarrow \varepsilon$ for $\varepsilon > |U|$. Here, the attractive interaction does not anymore favor thermally excited pair transitions over single-electron transitions. The resulting crossovers between all three identified regimes lead to surprising kinks in the Seebeck coefficient $S(\varepsilon)$ [blue line in Fig. 1(b)]. These kinks are even better visible in its derivative³³ [green, dashed line in Fig. 1(b)], measurable using lock-in techniques,

$$T \times \frac{\partial S}{\partial \varepsilon} = 1 + \delta n_d^2 \times \frac{|U|}{2T}. \quad (5)$$

Indeed, this derivative depends on the equilibrium charge fluctuations $\delta n_d^2 = \langle N^2 \rangle_d - n_d^2$ after the duality mapping. This implies features in $S(\varepsilon)$ at the Coulomb resonances $\varepsilon = 0$ and $\varepsilon = -U = |U|$ of a *repulsive* dot. The appearance of the dual occupation n_d in Eq. (4) is not expected from a brute-force “Fermi’s Golden rule” calculation and defies common physical intuition. This is typical for insights offered by fermionic duality.^{16,19,25}

The mapping to a repulsive system via n_d enables a further prediction for experiments. The peaks in $\partial S/\partial \varepsilon$ are shifted by approximately $\pm T \ln(2)$ away from the zero-temperature resonances $\varepsilon = 0, |U|$ of a repulsive dot. In the latter case, this shift is well understood as the entropy of the singly occupied state due to the spin degeneracy.^{34–36} In an attractive dot, single occupation is never a stable equilibrium state, but remarkably, our dual picture reveals that its spin degeneracy, nevertheless, affects the Seebeck coefficient.

A relevant question is how energy-dependent couplings $\Gamma_\alpha(E)$ affect the thermovoltage. In experiments, the environment density of states may sizably vary around the Fermi energy and, thereby, give rise to such an energy dependency. Moreover, an appropriately tuned energy dependence can be beneficial for efficient nanoscale energy harvesting.^{10,17,21–23}

We account for this by assuming arbitrary smoothly energy-dependent rates $\Gamma_{\alpha=L,R}(E) \ll T$ within the weak-coupling constraint. Following Ref. 19, S is then determined by

$$S \times T = \varepsilon - \frac{(1 + \Lambda)(2 - n_d)}{(1 - \Lambda)n_d + (1 + \Lambda)(2 - n_d)} |U|. \quad (6)$$

This introduces the energy asymmetry Λ of the coupling,

$$\Lambda = \frac{\Gamma_{UL}\Gamma_{UR}\Gamma_\varepsilon - \Gamma_{eL}\Gamma_{eR}\Gamma_U}{\Gamma_{UL}\Gamma_{UR}\Gamma_\varepsilon + \Gamma_{eL}\Gamma_{eR}\Gamma_U}, \quad (7)$$

with $\Gamma_{eL} = \Gamma_L(\varepsilon)$, $\Gamma_{eR} = \Gamma_R(\varepsilon)$, $\Gamma_{UL} = \Gamma_L(\varepsilon - |U|)$, $\Gamma_{UR} = \Gamma_R(\varepsilon - |U|)$, $\Gamma_\varepsilon = \Gamma_{eL} + \Gamma_{eR}$, and $\Gamma_U = \Gamma_{UL} + \Gamma_{UR}$. The result for S in the presence of energy-dependent tunnel coupling is shown in Fig. 2. Equation (6) enables us to systematically isolate how energy-dependent couplings $\Gamma_\alpha(E)$ influence the linear thermovoltage for different level positions ε at fixed $U = -|U|$ and T .^{37,47–50} In Fig. 2, we identify—as one main qualitative effect of energetic coupling asymmetry—a shift of the zero-crossing of S as a function of ε away from $\varepsilon_0 = |U|/2$. This shift, which we call $\Delta\varepsilon_0$, can be understood from Eq. (6). We exploit that well within $0 < \varepsilon < |U|$ and for $|U| \gg T$, repulsive Coulomb blockade induces a plateau at $n_d \approx 1$ in the dual occupation. This simplifies Eq. (6) to

$$S \times T \rightarrow \varepsilon - (1 + \Lambda)|U|/2 \quad \text{for} \quad 0 < \varepsilon < |U|, \quad (8)$$

FIG. 2. Seebeck coefficient [(a) and (b)] and its derivative at fixed Λ (c) as a function of dot level ε and coupling-asymmetry Λ . We take energy-dependent $\Gamma_L(E)$, $\Gamma_R(E)$, and $T = |U|/10$.

implying that an asymmetry $\Lambda > 0$ favors emission from a doubly occupied dot at addition energy $\varepsilon - |U|$ and $\Lambda < 0$ favors absorption into an empty dot at energy ε . This offset of $S(\varepsilon)$ in Eq. (8) involves a zero crossing ε_0 shifted away from $|U|/2$ by $\Delta\varepsilon_0 \rightarrow |U|/2 \times \Lambda$, predicting a pronounced effect for strong attractive interaction. The limit $\Delta\varepsilon_0 \rightarrow \pm|U|/2$ for $\Lambda \rightarrow \pm 1$ reflects that transport is reduced to a single resonance at $\varepsilon = 0$ or $\varepsilon = |U|$, annulling all two-electron features.

Next, Fig. 2(c) demonstrates how the peaks in $\partial S/\partial\varepsilon$, and hence the kinks in S visible in Fig. 1(b), change with energy-dependent couplings. For $\Lambda > 0$, the peak in $\partial S/\partial\varepsilon$ around $\varepsilon = 0$ shrinks with larger $|\Lambda|$, whereas the peak in $\partial S/\partial\varepsilon$ around $\varepsilon = |U|$ grows. At the same time, the latter also moves substantially further away from resonance—we refer to this shift from resonance as $\Delta\varepsilon_p$. For $\Lambda < 0$, the behavior is opposite with respect to the two peaks. In the single-resonance limit $|\Lambda| \rightarrow 1$, the slope $\partial S/\partial\varepsilon \times T \rightarrow 1$ becomes constant and both peaks disappear, as expected.

The change in the relative peak height follows from the offset of S within $0 < \varepsilon < |U|$ described in Eq. (8). For example, $\Lambda > 0$, the shift to smaller S in this ε -interval causes the step of $S(\varepsilon)$ around $\varepsilon = 0$ to be smaller than around $\varepsilon = |U|$, see Fig. 2(b). This leads to a smaller relative peak height in $\partial S/\partial\varepsilon$.

The growing shift $\Delta\varepsilon_p$ of the higher peak for increasing $|\Lambda|$ stems from the fact that the coupling asymmetry Λ not only affects S for $0 < \varepsilon < |U|$ and large $|U|/T$ as in Eq. (8) but, in general, also influences where the crossover between the single-particle, linear limits ($S(\varepsilon) \sim \varepsilon$ and $S(\varepsilon) \sim \varepsilon - |U|$) and the two-particle limit ($S(\varepsilon) \sim \varepsilon - |U|/2$) takes place [see Eq. (6)]. For both small $|\Lambda|$ and large $|\Lambda| \leq 1$, a useful analytical expression is [including the spin-degeneracy shift $T \ln(2)$]

$$\Delta\varepsilon_p \approx T \times \text{sgn}(\Lambda) \times \ln[2/(1 - |\Lambda|)]. \quad (9)$$

For example, $\Lambda = 0.8$ yields $\Delta\varepsilon_p \approx 3.3T \ln(2)$, which substantially deviates from the wideband limit result, where the shift away from resonance is given by $T \ln(2)$. In this case, Γ_U is large enough compared to Γ_ε , such that even for a considerable interval with $\varepsilon > 0$ and $\varepsilon - |U| > 0$, the physics of pair tunneling prevails. Namely, a thermally excited electron entering the dot at energy ε causes transport of further electrons at energy $\varepsilon - |U|$ before the dot is emptied again.

Next, we demonstrate how the Seebeck coefficient gets modified in the *nonlinear regime* due to large ΔT and V . This is also relevant below where we discuss the performance of the quantum dot as a thermoelectric device. The nonlinear thermovoltage $S_{nl} = V|_{I=0}/\Delta T$ quantifies the voltage $V = \mu - \mu_L$ required to suppress a charge current I induced by a large temperature difference $\Delta T = T_L - T$ across the junction.

We have previously shown¹⁶ the nonlinear current to take the compact form,

$$I = \frac{\gamma_L \gamma_R}{\gamma_L + \gamma_R} (n_L - n_R), \quad (10)$$

in the wideband limit ($\Gamma_{\varepsilon\alpha} = \Gamma_{U\alpha} = \Gamma_\alpha$). This depends on the difference between *equilibrium* occupations $n_R = n$ and $n_L = n|_{\mu, T \rightarrow \mu_L, T_L}$ and on the energy-level dependent charge relaxation rates^{38,39} $\gamma_{R/L} = \Gamma_{R/L}[1 + f_{R/L}(\varepsilon) - f_{R/L}(\varepsilon - |U|)]/2 > 0$. Both the occupations and relaxation rates can be understood as if the dot was coupled only to the right or left lead; the symbols $f_R(x) = f(x)$ and $f_L(x) = f(x)|_{\mu, T \rightarrow \mu_L, T_L}$ denote the corresponding Fermi functions. Setting $I = 0$ while keeping the potential $\mu_R = \mu$ and temperature $T_R = T$ fixed, Eq. (10) yields the helpful analytical result for the thermovoltage,¹⁶

$$S_{nl} \times T = \varepsilon - |U| - \frac{T + \Delta T}{\Delta T/T} \times \ln \left[\frac{1 - n_d + \sqrt{(1 - n_d)^2 + \exp\left[-\frac{|U|\Delta T}{T(T + \Delta T)}\right]} n_d (2 - n_d)}{2 - n_d} \right], \quad (11)$$

again expressed in terms of the dual occupation number n_d .

Figure 3 shows $S_{nl} \times T$ and its ε -derivative⁴⁰ as a function of level position ε and temperature difference $\Delta T > 0$. As expected, the zero crossing at the particle-hole symmetric point $\varepsilon = |U|/2$ persists. Importantly, the counter intuitive features at $\varepsilon = 0, |U|$ also continue to exist. This is indeed suggested by Eq. (11), in which the ε -dependence enters *entirely* through the dual occupation n_d determined by repulsive interaction. Specifically, in Fig. 3(c), an increasing ΔT transforms the steps at $\varepsilon = 0, |U|$ between the three regimes of $S_{nl}(\varepsilon) \times T$ with equal ε -slopes into temperature-broadened transitions between three regimes of *different* slopes. For $\varepsilon < 0$ and $\varepsilon > |U|$, $S_{nl}(\varepsilon) \times T$ still grows with a slope of 1 as function of ε , just as the Seebeck coefficient $S(\varepsilon) \times T$ [see Fig. 1(b)]. This again reflects that transport is effectively governed by single-particle physics (see above discussion on energy-dependent couplings). For levels $0 < \varepsilon < |U|$ at which two-particle effects are relevant, a linear ε -dependence of $S_{nl} \times T$ with a larger slope of ~ 2 emerges, as can be qualitatively understood from an analysis of the nonlinear charge current (10): a small $T \ll |U|$ and a large $\Delta T \geq |U|$ correspond to a sharp two-particle transition of $n_R = 2 \rightarrow 0$, yet a smooth behavior of n_L as a function of ε around

FIG. 3. Nonlinear thermovoltage [(a) and (c)] and its derivative (b) as a function of dot level ε and temperature gradient $\Delta T/|U|$. We take $T = |U|/10$ and Γ_L, Γ_R that are energy-independent.

$\varepsilon = |U|/2$. Consequently, fulfilling $n_L = n_R$ to achieve $I = 0$ for fixed ε and μ requires a relatively large shift of $V = \mu - \mu_L$. In particular, the slope of $2 \times S_{nl}(\varepsilon) \times T = \frac{\sqrt{T}}{\Delta T} \Big|_{I=0}$ in the limit $\Delta T/T \gg 1$ reflects the sharp change of n_R by 2 at $\varepsilon = |U|/2$ due to attractive interaction [see Fig. 1(c)].

Finally, let us consider the power output of the dot. As is well-known,^{41–44} a sharp spectral resonance of a conductor is beneficial for its thermoelectric performance. Hence, quantum dots have been studied as thermoelectric elements operated in the nonlinear regime of large temperature and voltage biases, both theoretically^{8,9,12,16} and experimentally.^{11,20} We now show that also in the presence of strong attractive interaction, finite power output is possible at high efficiencies.

We study the power output $P = I \times V$ with current I given by Eq. (10) as well as the efficiency $\eta = P/J$, where J is the heat current out of the left (hot) reservoir. Analytical expressions for J are derived in Ref. 16 for a generic onsite interaction. In Fig. 4(a), we show P and η as a function of V for $\Delta T = -|U|$, $\Gamma_L = \Gamma_R = \Gamma$ and for an $\varepsilon = 1.23|U|$ in the vicinity of the crossover between single- and two-particle regimes, optimized for maximal power output. The power clearly behaves non-monotonically with a peak at $|V| \approx \Delta T$. Efficiency η increases with voltage and assumes about 0.6 times the Carnot efficiency $\eta_C = 1 - T_R/T_L$ at maximum power. These efficiencies are sizable at finite power output, as can clearly be seen in Fig. 4(b), where η and P are shown for several temperature differences ΔT . These also show that efficiencies reach the Carnot limit when power is suppressed at large voltages.⁴⁵

Figures 4(c) and 4(d) show the power $P_{\max}$ maximized over V at otherwise fixed parameters and the efficiency $\eta_{P_{\max}}$ at $P_{\max}$. Extended regions of ε - and ΔT -values have sizable power $P_{\max} \gtrsim \Gamma T$. Interestingly, Fig. 4(c) shows that the maximum power is fully suppressed *only* at $\varepsilon = |U|/2$. This can be understood by the fact that the

nonlinear thermovoltage S_{nl} only disappears at this level position, as shown above. However, in the whole two-particle regime, $0 < \varepsilon < |U|$, the power is small compared to the single-particle regime, $\varepsilon < 0$ and $\varepsilon > |U|$. The reason is that the charge relaxation rate γ_R entering the current, Eq. (10), is suppressed in this regime by attractive Coulomb blockade.²⁵ This leads to low power production.

Important for the performance of the attractive quantum dot as a heat engine is our finding that the efficiency at maximum output power is on the order of the Curzon–Ahlborn bound,⁴⁶ $\eta_{CA} = 1 - \sqrt{T_R/T_L}$, in the whole range in which the output power is sizable. It even reaches this bound close to the *dual* resonances, $\varepsilon = 0, |U|$, namely, the level positions at which n_d obtained from the dual mapping changes by 1. It is also remarkable that for the finite power output, prominent features appear at the resonances of the dual repulsive model.

To conclude, we have analyzed the thermoelectric response of a weakly coupled, single-level quantum dot with attractive interaction together with its performance as a steady-state heat engine. The presented results are expected to be important for future experiments aiming to characterize systems with strong attractive onsite interaction. At the same time, they demonstrate that nanodevices based on quantum dots with attractive interaction can also efficiently convert heat into work.

The most relevant qualitative features that we found are (i) two stepped features in the Seebeck coefficient S instead of one, *unexpectedly* located at the positions for resonances of a repulsive dot (Fig. 1). (ii) Nonlinear Seebeck coefficient S_{nl} is constant up to sizeable thermal bias (Fig. 3). (iii) Sharp dip in the maximum power and efficiency at the position *expected* for the attractive dot (Fig. 4). We anticipate these effects in state-of-the-art experiments as, e.g., in quantum dots defined at an oxide interface⁴ where clear features of attractive interaction have already been seen in voltage-driven charge transport measurements. Here, applying a temperature bias would allow to verify our predictions, in particular, due to the available electrical control over the level position. In contrast, nanostructures such as in Ref. 3 allow the magnitude and sign of the *real* interaction to be altered. This would enable a direct comparison between attractive and *real* repulsive quantum dots. Predictions for such a comparison are outlined in the [supplementary material](#). However, these setups would need to be extended to allow for transport measurements.

The remarkable appearance of prominent features at level positions characteristic of a repulsive quantum dot was rationalized with a dual mapping emerging from a dissipative symmetry for master equations. For attractive quantum dots, the role of this dual mapping is particularly important: The dual features with respect to the original attractive system *do not* appear at special positions, where, e.g., particle-hole symmetry imposes restrictions. Their prominent role could hence not have been predicted straightforwardly in another way.

See the [supplementary material](#) for a comparison to a quantum dot with *real* repulsive interaction.

We acknowledge financial support from the Knut and Alice Wallenberg foundation and the Swedish VR (J.Sp., J.Sc.) and the Danish National Research Foundation (J.Sc.).

DATA AVAILABILITY

The data that support the findings of this study are available within this article.

FIG. 4. (a) Power P and efficiency η as a function of the voltage bias V at the fixed level position, $\varepsilon = 1.23|U|$, $\Delta T = |U|$, $T = |U|/10$. (b) Efficiency vs power at the fixed level position, $\varepsilon = 1.23|U|$ for different temperature biases ΔT . (c) Power $P_{\max}$ maximized over V . (d) Efficiency $\eta_{P_{\max}}$ at maximum power. In all panels, we set $\Gamma_L = \Gamma_R = \Gamma$.

REFERENCES

- ¹M. R. Butler, B. Movaghar, T. J. Marks, and M. A. Ratner, "Electron pairing in designer materials: A novel strategy for a negative effective Hubbard U," *Nano Lett.* **15**, 1597–1602 (2015).
- ²G. Cheng, M. Tomczyk, S. Lu, J. P. Veazey, M. Huang, P. Irvin, S. Ryu, H. Lee, C.-B. Eom, C. S. Hellberg, and J. Levy, "Electron pairing without superconductivity," *Nature* **521**, 196 (2015).
- ³A. Hamo, A. Benyamini, I. Shapir, I. Khivrich, J. Waissman, K. Kaasbjerg, Y. Oreg, F. von Oppen, and S. Ilani, "Electron attraction mediated by Coulomb repulsion," *Nature* **535**, 395 (2016).
- ⁴G. E. D. K. Prawiroatmodjo, M. Leijnse, F. Trier, Y. Chen, D. V. Christensen, M. von Soosten, N. Pryds, and T. S. Jespersen, "Transport and excitations in a negative-U quantum dot at the LaAlO₃/SrTiO₃ interface," *Nat. Commun.* **8**, 395 (2017).
- ⁵J. Koch, M. E. Raikh, and F. von Oppen, "Pair tunneling through single molecules," *Phys. Rev. Lett.* **96**, 056803 (2006).
- ⁶E. Sela, H.-S. Sim, Y. Oreg, M. E. Raikh, and F. von Oppen, "Electron-pair resonance in the Coulomb blockade," *Phys. Rev. Lett.* **100**, 056809 (2008).
- ⁷M. Leijnse, M. R. Wegewijs, and M. H. Hettler, "Pair tunneling resonance in the single-electron transport regime," *Phys. Rev. Lett.* **103**, 156803 (2009).
- ⁸M. Esposito, K. Lindenberg, and C. Van den Broeck, "Thermoelectric efficiency at maximum power in a quantum dot," *Europhys. Lett.* **85**, 60010 (2009).
- ⁹M. Leijnse, M. R. Wegewijs, and K. Flensberg, "Nonlinear thermoelectric properties of molecular junctions with vibrational coupling," *Phys. Rev. B* **82**, 045412 (2010).
- ¹⁰R. Sánchez and M. Büttiker, "Optimal energy quanta to current conversion," *Phys. Rev. B* **83**, 085428 (2011).
- ¹¹S. F. Svensson, E. A. Hoffmann, N. Nakpathomkun, P. M. Wu, H. Q. Xu, H. A. Nilsson, D. Sánchez, V. Kashcheyevs, and H. Linke, "Nonlinear thermovoltage and thermocurrent in quantum dots," *New J. Phys.* **15**, 105011 (2013).
- ¹²D. M. Kennes, D. Schuricht, and V. Meden, "Efficiency and power of a thermoelectric quantum dot device," *Europhys. Lett.* **102**, 57003 (2013).
- ¹³B. Sothmann, R. Sánchez, and A. N. Jordan, "Thermoelectric energy harvesting with quantum dots," *Nanotechnology* **26**, 032001 (2015).
- ¹⁴A. Svilans, A. M. Burke, S. F. Svensson, M. Leijnse, and H. Linke, "Nonlinear thermoelectric response due to energy-dependent transport properties of a quantum dot," *Physica E* **82**, 34–38 (2016).
- ¹⁵D. Sánchez and R. López, "Nonlinear phenomena in quantum thermoelectrics and heat," *C. R. Phys.* **17**, 1060–1071 (2016).
- ¹⁶J. Schulenburg, A. D. Marco, J. Vanherck, M. R. Wegewijs, and J. Splettstoesser, "Thermoelectrics of interacting nanosystems—Exploiting superselection instead of time-reversal symmetry," *Entropy* **19**, 668 (2017).
- ¹⁷N. Walldorf, A.-P. Jauho, and K. Kaasbjerg, "Thermoelectrics in Coulomb-coupled quantum dots: Cotunneling and energy-dependent lead couplings," *Phys. Rev. B* **96**, 115415 (2017).
- ¹⁸P. A. Erdman, F. Mazza, R. Bosisio, G. Benenti, R. Fazio, and F. Taddei, "Thermoelectric properties of an interacting quantum dot based heat engine," *Phys. Rev. B* **95**, 245432 (2017).
- ¹⁹J. Schulenburg, J. Splettstoesser, and M. R. Wegewijs, "Duality for open fermion systems: Energy-dependent weak coupling and quantum master equations," *Phys. Rev. B* **98**, 235405 (2018).
- ²⁰M. Josefsson, A. Svilans, A. M. Burke, E. A. Hoffmann, S. Fahlvik, C. Thelander, M. Leijnse, and H. Linke, "A quantum-dot heat engine operating close to the thermodynamic efficiency limits," *Nat. Nanotechnol.* **13**, 920–924 (2018).
- ²¹F. Hartmann, P. Pfeffer, S. Höfling, M. Kamp, and L. Worschech, "Voltage fluctuation to current converter with Coulomb-coupled quantum dots," *Phys. Rev. Lett.* **114**, 146805 (2015).
- ²²B. Roche, P. Rouleau, T. Jullien, Y. Jompol, I. Farrer, D. A. Ritchie, and D. C. Glattli, "Harvesting dissipated energy with a mesoscopic ratchet," *Nat. Commun.* **6**, 6738 (2015).
- ²³H. Thierschmann, R. Sánchez, B. Sothmann, F. Arnold, C. Heyn, W. Hansen, H. Buhmann, and L. W. Molenkamp, "Three-terminal energy harvester with coupled quantum dots," *Nat. Nanotechnol.* **10**, 854 (2015).
- ²⁴H. Thierschmann, R. Sánchez, B. Sothmann, H. Buhmann, and L. W. Molenkamp, "Thermoelectrics with Coulomb-coupled quantum dots," *C. R. Phys.* **17**, 1109–1122 (2016).
- ²⁵J. Schulenburg, R. B. Saptsov, F. Haupt, J. Splettstoesser, and M. R. Wegewijs, "Fermion-parity duality and energy relaxation in interacting open systems," *Phys. Rev. B* **93**, 081411 (2016).
- ²⁶E. Kleinherbers, P. Stegmann, and J. König, "Revealing attractive electron–electron interaction in a quantum dot by full counting statistics," *New J. Phys.* **20**, 073023 (2018).
- ²⁷B. A. Placke, T. Pluecker, J. Splettstoesser, and M. R. Wegewijs, "Attractive and driven interactions in quantum dots: Mechanisms for geometric pumping," *Phys. Rev. B* **98**, 085307 (2018).
- ²⁸V. Bruch, K. Nestmann, J. Schulenburg, and M. R. Wegewijs, "Fermionic duality: General symmetry of open systems with strong dissipation and memory" (to be published).
- ²⁹Up to linear order in the couplings Γ_z , the dual model is a physically valid model and can be interpreted physically. This breaks down in higher-than-linear orders in Γ_z , but the duality remains useful.²⁸
- ³⁰C. W. J. Beenakker and A. A. M. Staring, "Theory of the thermopower of a quantum dot," *Phys. Rev. B* **46**, 9667 (1992).
- ³¹A. A. M. Staring, L. W. Molenkamp, B. W. Alphenaar, H. van Houten, O. J. A. Buyk, M. A. A. Mabesoone, C. W. J. Beenakker, and C. T. Foxon, "Coulomb-blockade oscillations in the thermopower of a quantum dot," *Europhys. Lett.* **22**, 57 (1993).
- ³²A. S. Dzurak, C. G. Smith, M. Pepper, D. A. Ritchie, J. E. F. Frost, G. A. C. Jones, and D. G. Hasko, "Observation of Coulomb blockade oscillations in the thermopower of a quantum dot," *Solid State Commun.* **87**, 1145–1149 (1993).
- ³³The relation $\delta n_d^2 = T \partial n_d / \partial \varepsilon$ is proven in Ref. 16.
- ³⁴M. M. Deshmukh, E. Bonet, A. N. Pasupathy, and D. C. Ralph, "Equilibrium and nonequilibrium electron tunneling via discrete quantum states," *Phys. Rev. B* **65**, 073301 (2002).
- ³⁵E. Bonet, M. M. Deshmukh, and D. C. Ralph, "Solving rate equations for electron tunneling via discrete quantum states," *Phys. Rev. B* **65**, 045317 (2002).
- ³⁶S. Juergens, F. Haupt, M. Moskalets, and J. Splettstoesser, "Thermoelectric performance of a driven double quantum dot," *Phys. Rev. B* **87**, 245423 (2013).
- ³⁷ Λ actually depends on ε via $\Gamma_z(E)$ [Eq. (7)]. In Fig. 2, ε is swept while adjusting $\Gamma_z(E)$ to keep Λ constant. However, as Ref. 19 has shown, an ε -sweep that keeps Λ constant is achieved without adjusting the couplings for an exponential $\Gamma_z(E) \sim \exp((E - E_0)/D)$. Such an energy profile has proven to be a reasonable assumption in analyzing several previous experiments on few-electron transport from quantum dots.^{47–50}
- ³⁸J. Splettstoesser, M. Governale, J. König, and M. Büttiker, "Charge and spin dynamics in interacting quantum dots," *Phys. Rev. B* **81**, 165318 (2010).
- ³⁹J. Vanherck, J. Schulenburg, R. B. Saptsov, J. Splettstoesser, and M. R. Wegewijs, "Relaxation of quantum dots in a magnetic field at finite bias: Charge, spin, and heat currents," *Phys. Status Solidi B* **254**, 1600614 (2017).
- ⁴⁰The derivative can be straightforwardly analytically obtained from Eq. (11).
- ⁴¹G. D. Mahan and J. O. Sofo, "The best thermoelectric," *Proc. Natl. Acad. Sci. U. S. A.* **93**, 7436–7439 (1996).
- ⁴²L. D. Hicks and M. S. Dresselhaus, "Effect of quantum-well structures on the thermoelectric figure of merit," *Phys. Rev. B* **47**, 12727–12731 (1993).
- ⁴³L. D. Hicks and M. S. Dresselhaus, "Thermoelectric figure of merit of a one-dimensional conductor," *Phys. Rev. B* **47**, 16631–16634 (1993).
- ⁴⁴T. E. Humphrey and H. Linke, "Reversible thermoelectric nanomaterials," *Phys. Rev. Lett.* **94**, 096601 (2005).
- ⁴⁵The sharp feature at low power and high efficiencies stems from the exponential tails of $n_{L/R}$, prohibiting an exactly suppressed current. We expect this to be smoothed by here neglected higher-order tunneling effects.
- ⁴⁶F. L. Curzon and B. Ahlborn, "Efficiency of a Carnot engine at maximum power output," *Am. J. Phys.* **43**, 22 (1975).
- ⁴⁷K. A. Matveev and L. I. Glazman, "Coulomb blockade of activated conduction," *Phys. Rev. B* **54**, 10339–10341 (1996).
- ⁴⁸N. M. Zimmerman, E. Hourdakis, Y. Ono, A. Fujiwara, and Y. Takahashi, "Error mechanisms and rates in tunable-barrier single-electron turnstiles and charge-coupled devices," *J. Appl. Phys.* **96**, 5254–5266 (2004).
- ⁴⁹J. D. Fletcher, M. Kataoka, S. P. Giblin, S. Park, H.-S. Sim, P. See, D. A. Ritchie, J. P. Griffiths, G. A. C. Jones, H. E. Beere, and T. J. B. M. Janssen, "Stabilization of single-electron pumps by high magnetic fields," *Phys. Rev. B* **86**, 155311 (2012).
- ⁵⁰B. Kaestner and V. Kashcheyevs, "Non-adiabatic quantized charge pumping with tunable-barrier quantum dots: A review of current progress," *Rep. Prog. Phys.* **78**, 103901 (2015).