



A two-sided but significantly beamed jet in the supercritical accretion quasar IRAS F11119+3257

Downloaded from: <https://research.chalmers.se>, 2025-12-04 20:05 UTC

Citation for the original published paper (version of record):

Yang, J., Paragi, Z., An, T. et al (2020). A two-sided but significantly beamed jet in the supercritical accretion quasar IRAS F11119+3257. *Monthly Notices of the Royal Astronomical Society*, 494(2): 1744-1750. <http://dx.doi.org/10.1093/MNRAS/STAA836>

N.B. When citing this work, cite the original published paper.

A two-sided but significantly beamed jet in the supercritical accretion quasar IRAS F11119+3257

Jun Yang¹,¹★ Zsolt Paragi,² Tao An³, Willem A. Baan^{4,5}, Prashanth Mohan³ and Xiang Liu⁵

¹Department of Space, Earth and Environment, Chalmers University of Technology, Onsala Space Observatory, SE-439 92 Onsala, Sweden

²Joint Institute for VLBI ERIC (JIVE), Postbus 2, NL-7990 AA Dwingeloo, the Netherlands

³Shanghai Astronomical Observatory, Key Laboratory of Radio Astronomy, Chinese Academy of Sciences, 200030 Shanghai, China

⁴Netherlands Institute for Radio Astronomy ASTRON, NL-7991 PD Dwingeloo, the Netherlands

⁵Xinjiang Astronomical Observatory, Key Laboratory of Radio Astronomy, Chinese Academy of Sciences, 150 Science 1-Street, 830011 Urumqi, China

Accepted 2020 March 20. Received 2020 March 20; in original form 2020 February 7

ABSTRACT

Highly accreting quasars are quite luminous in the X-ray and optical regimes; while, they tend to become radio quiet and have optically thin radio spectra. Among the known quasars, IRAS F11119+3257 is a supercritical accretion source because it has a bolometric luminosity slightly above the Eddington limit and extremely powerful X-ray outflows. To probe its radio structure, we investigated its radio spectrum between 0.15 and 96.15 GHz and performed very-long-baseline interferometric (VLBI) observations with the European VLBI Network (EVN) at 1.66 and 4.93 GHz. The deep EVN image at 1.66 GHz shows a two-sided jet with a projected separation about 200 pc and a very high flux density ratio of about 290. Together with the best-fitting value of the integrated spectral index of -1.31 ± 0.02 in the optically thin part, we infer that the approaching jet has an intrinsic speed at least 0.57 times of the light speed. This is a new record among the known all kinds of super-Eddington accreting sources and unlikely accelerated by the radiation pressure in a certain models. We propose a scenario in which IRAS F11119+3257 is an unusual compact symmetric object with a small jet viewing angle and a radio spectrum peaking at 0.53 ± 0.06 GHz mainly due to the synchrotron self-absorption.

Key words: galaxies: active – galaxies: jets – quasars: individual: IRAS F11119+3257 – radio continuum: galaxies.

1 INTRODUCTION

Nearby active galactic nuclei (AGNs) hosting highly accreting (accretion rate comparable to or exceeding the Eddington rate) supermassive black holes (SMBHs) are useful targets for monitoring observations in several contexts. Some of these include studies of powerful X-ray absorption outflows and their impact on the host galaxy evolution (e.g. Kormendy & Ho 2013; Nardini et al. 2015; Tombesi et al. 2015; Tombesi 2016), and in exploring radio jet and outflow activity in an extreme accretion regime (e.g. Giroletti et al. 2017; Yang et al. 2018). These highly accreting AGNs are typically radio-quiet (e.g. Greene, Ho & Ulvestad 2006; Panessa et al. 2007; Sikora, Stawarz & Lasota 2007) with a steep radio spectrum (Laor, Baldi & Behar 2019) when their accretion rates approach or exceed the Eddington limit. The existence of young,

possibly episodic jets moving at mildly relativistic jet speeds in these AGNs (e.g. Panessa et al. 2019) thus remain debatable. Further, their high-redshift counterparts are probes of rapid SMBH growth (e.g. Volonteri, Silk & Dubus 2015), feedback interaction with the host galaxy through jet and radiative processes (e.g. Pacucci, Volonteri & Ferrara 2015), and are used as standard candles for determining cosmological luminosity distance (e.g. Wang et al. 2013; Marziani & Sulentic 2014).

The quasar IRAS F11119+3257 (J1114+3241, B1111+329) at a redshift $z = 0.189$ is a type 1 ultraluminous infrared galaxy (ULIRG) and hosts strong molecular outflows, with the emission dominated by the AGN component (Veilleux et al. 2013, 2017). Based on an SMBH mass of $M_{\text{bh}} \approx 2 \times 10^7 M_{\odot}$ calibrated for a sample of similar ULIRG sources (Kawakatu, Imanishi & Nagao 2007), the bolometric luminosity $L_{\text{b}} = 5 L_{\text{Edd}}$, where L_{Edd} is the Eddington luminosity (Tombesi et al. 2015). Using a correlation relation between infrared and radio luminosities for starburst galaxies, it is found that for IRAS F11119+3257 the AGN component far exceeds

* E-mail: jun.yang@chalmers.se

Table 1. List of the radio flux densities of IRAS F11119+3257. The used radio facilities are the Giant Metrewave Radio Telescope (GMRT), the Westerbork Synthesis Radio Telescope (WSRT), the Robert C. Byrd Green Bank Telescope (GBT), the Karl G. Jansky Very Large Array (VLA), and the Atacama Large Millimeter Array (ALMA).

Freq. (GHz)	Flux (mJy)	Array	Reference
0.15	154 ± 16	GMRT	Intema et al. (2017)
0.33	241 ± 24	WSRT	Rengelink et al. (1997)
1.40	105 ± 5	VLA	Becker, White & Helfand (1995)
1.40	110 ± 3	VLA	Condon et al. (1998)
1.43	106 ± 5	VLA	This paper
4.85	23 ± 4	GBT	Gregory et al. (1996)
5.23	21.0 ± 1.1	VLA	Berton et al. (2018)
8.44	11 ± 0.6	VLA	This paper
14.94	4.6 ± 0.3	VLA	This paper
96.15	0.5 ± 0.05	ALMA	Veilleux et al. (2017)

the starburst contribution (Komossa et al. 2006). In addition to the molecular outflows, it also hosts wide-aperture energetic radiation-driven X-ray emitting winds, suggesting a likely energy conserving quasar-mode feedback (Tombesi et al. 2015, 2017).

The quasar IRAS F11119+3257 has a relatively bright radio counterpart. The early survey of the Bologna Northern Cross Radio Telescope (BNCRT) at 408 MHz (Colla et al. 1970) found a half-Jy radio counterpart. Later, it was observed by more radio telescopes at multiple frequencies. The radio flux densities reported in literature over the past 30 yr are listed in Table 1. All these existing observations indicate a compact emission structure (e.g. Veilleux et al. 2017; Berton et al. 2018). Although it has an increasingly steep radio spectrum at frequencies $\gtrsim 1$ GHz, it may not be classified as radio-quiet owing to a high radio power ($\approx 10^{25}$ W Hz $^{-1}$) reminiscent of a radio-loud quasar (Komossa et al. 2006).

The radio jet, radiation pressure from the disc emission and disc winds can contribute to driving the powerful X-ray outflows in this source (e.g. Tombesi 2016). To probe this and to possibly resolve the compact but steep spectrum radio source, we conducted high-resolution very-long-baseline interferometric (VLBI) observations of the pc-scale region. This paper is compiled as follows. In Section 2, we describe the radio observations of IRAS F11119+3257 and the data reduction. In Section 3, we report the broad-band radio spectrum and the radio morphology. In Section 4, we discuss the radio core, constraints on the jet parameters, and the relation to young radio sources. In Section 5, we present the main conclusions from our study. Throughout the paper, a standard Λ CDM cosmological model with $H_0 = 71$ km s $^{-1}$ Mpc $^{-1}$, $\Omega_m = 0.27$, $\Omega_\Lambda = 0.73$ is adopted; the VLBI images then have a scale of 3.9 pc mas $^{-1}$.

2 OBSERVATIONS AND DATA REDUCTION

2.1 The EVN experiments at 1.66 and 4.93 GHz

We observed IRAS F11119+3257 with the European VLBI Network (EVN) at 1.66 and 4.93 GHz in 2016. The experiment set-ups are summarized in Table 2. The participating telescopes were Robledo (Ro), Sardinia (Sr), Hartesbeesthoek (Hh), Zelenchukskaya (Zc), Sveltoe (Sv), Torun (Tr), Urumqi (Ur), Tianma (T6), Onsala (O8), Medicina (Mc), Effelsberg (Ef), Westerbork (Wb, single dish), Jodrell Bank Lovell (Jb1), and Mk2 (Jb2). The correlation was done by the EVN software correlator (SFXC; Keimpema et al. 2015) at

JIVE (Joint Institute for VLBI, ERIC) using the typical correlation parameters.

The short EVN observations at 4.93 GHz were performed in the e-VLBI mode on 2016 November 16. The data rate was reduced to 1024 Mbps for T6 due to the network limitation, and Tr and Jb2 due to the VLBI backends. To calibrate the data and measure a precise position for our faint target IRAS F11119+3257, a bright compact source J1111+3252 (Helmholtz et al. 2007), about 41 arcmin apart from IRAS F11119+3257, was also observed periodically. The phase-referencing calibrator position is RA = 11^h11^m31^s.77219, Dec. = 32°52′55″.7847 (J2000, $\sigma_{ra} = \sigma_{dec} = 0.16$ mas) in the source catalogue¹ provided by L. Petrov from the Goddard Space Flight Centre VLBI group. The calibrator position has an offset of 1.4 mas with respect to the sub-mas-precision optical position in the second data release (DR2; Brown et al. 2018) of the *Gaia* mission (Prusti et al. 2016). The nodding observations used a cycle period of about six min (1.5 min for J1111+3252, 4 min for IRAS F11119+3257). All the telescopes had an elevation of ≥ 18 deg during the observations.

The full EVN observations at 1.66 GHz were carried out in the disc-recording mode on 2016 March 8. There were 13 stations participating in the 4-h observations. Because the target source is relatively bright at 1.66 GHz, the phase-referencing calibrator was not observed. This allowed us to reach an on-target time of about 200 min.

The visibility data were calibrated using the National Radio Astronomy Observatory (NRAO) software package Astronomical Image Processing System (AIPS; Greisen 2003). As the digital filters provided by the European digital VLBI backends at the most stations had a valid bandwidth of only about 75 per cent, the side channels were dropped out in loading the data into AIPS. With a reduction of the bandwidth, the task ACCOR was performed to correct the cross-correlation amplitude.

A priori amplitude calibration was performed with the system temperatures and the antenna gain curves. For cases where the telescope monitoring data were missing, nominal values of the system equivalent flux density in the EVN status table were used. The ionospheric dispersive delays were corrected according to a map of total electron content provided by Global Positioning System (GPS) satellite observations. Phase errors due to antenna parallactic angle variations were removed. After a manual phase calibration was carried out, the global fringe fitting and the bandpass calibration were performed.

At 4.93 GHz, we first imaged the phase-referencing calibrator J1111+3252. We iteratively ran model fitting with point sources and self-calibration in DIFMAP (version 2.5e; Shepherd, Pearson & Taylor 1994), fringe fitting, and self-calibration to remove its structure-dependent phase errors in AIPS. The calibrator has a single-side core-jet structure with a total flux density of 0.16 ± 0.02 Jy. Its radio core, i.e. the jet base, has a peak brightness of 0.09 ± 0.01 Jy beam $^{-1}$ and its position was used as the reference point in the phase-referencing calibration. We also ran amplitude self-calibrations on the calibrator data and transferred the solutions to the target data. Owing to the limited uv coverage of the short observations in particular on the long baselines, the deconvolution was performed by fitting the visibility data directly to some point source models in DIFMAP to minimize the potential deconvolution errors of CLEAN.

¹http://astrogeo.org/vlbi/solutions/rfc_2019d/

Table 2. The experiment set-ups of the used EVN observations. Columns give (1) date, (2) frequency, (3) total time, (4) data rate, (5) baseband filter, (6) project code, (7) observing mode, and (8) participating stations (see Section 2.1 for the explanation on the telescope codes).

Date	ν_{obs} (GHz)	Time (h)	Rate (Mbps)	Filter (MHz)	Project code	Observing mode	Participating stations
2016 Mar 08	1.66	4.0	1024	16	EY024B	Disc recording VLBI	Ro, Sr, Hh, Zc, Sv, Tr, Ur, T6, O8, Mc, Ef, Wb, Jb1
2016 Nov 16	4.93	2.5	2048	32	RSY04	Real-time e-VLBI	Ir, Hh, T6, Ys, O8, Nt, Mc, Ef, Wb, Jb2

We performed about 10 times iterations of the deconvolution and the self-calibration at 1.66 GHz. At each iteration, the new image model was used to solve for the residual systematic phase and amplitude errors in AIPS. When some significant ($\gtrsim 5\sigma$) positive features were found in the residual intensity map and the self-calibration failed to remove them, new model-fitting delta components were manually added to improve the fitting in DIFMAP. Totally, there were 35 point-source components used at 1.66-GHz. The reliability of each main feature was also further verified by removing it and seeing the beam pattern around it with proper data weighting and tapering. As a consequence of no observations of the phase-referencing calibrator at 1.66 GHz, the image peak positions were used to align the images at 1.66 and 4.93 GHz.

We also tried the standard CLEAN algorithm with carefully applying data weighting and tapering in DIFMAP. Using the final self-calibrated data, we could make a consistent map while with a relatively high (about 1.3 times) noise level mainly in the on-source region. This is a known limitation of the CLEAN algorithm because it generated a certain spurious structure in the form of spots or ridges as modulation on the broad features. Thus, we prefer the model-fitting technique to the CLEAN algorithm.

2.2 The VLA archive data

To study the compactness of IRAS F11119+3257 on the kpc scales, we downloaded the archive visibility data of project AK0311 observed by the VLA at 1.4 and 8.4 GHz on 1992 December 20. The data were calibrated and provided by the image retrieval tool of the NRAO science data archive. The flux densities were reported in Table 1.


We also downloaded the raw visibility data of the VLA project AN0104. The project was observed by Nagar et al. (2003) at 15 GHz on 2002 February 23. According to the standard calibration steps recommended by the AIPS cookbook, we calibrated the data manually with a short solution interval of 1 min. An image sensitivity of 0.16 mJy beam⁻¹ was achieved. The quasar IRAS F11119+3257 is detected with a signal-to-noise ratio (SNR) of 28, against the early report of non-detection (Nagar et al. 2003).

3 RESULTS

3.1 Broad-band radio spectrum

On the kpc scales, IRAS F11119+3257 is unresolved in the VLA images. With a natural weighing, the VLA image sensitivities reach 0.13 mJy beam⁻¹ at 1.43 GHz, 0.03 mJy beam⁻¹ at 8.44 GHz, and 0.16 mJy beam⁻¹ at 14.94 GHz. These VLA flux density measurements are also reported in Table 1.

There is no significant flux density variability observed so far. Our 1.43 GHz flux measurements are consistent with the results of the VLA surveys of the FIRST (Faint Images of the Radio Sky at Twenty Centimeters, Becker et al. 1995) and the NVSS (NRAO VLA Sky Survey, Condon et al. 1998). The new 5.23 GHz flux

**Figure 1.** The broad-band radio spectra of IRAS F11119+3257. The data are listed in Table 1. The grey curve shows the best-fitting results of equation (1).

measurement (Berton et al. 2018) has no significant difference from that observed by the GBT about 30 yr ago (Gregory et al. 1996).

The broad-band radio spectrum of IRAS F11119+3257 in Fig. 1 shows a spectral shape similar to GHz-peaked spectrum sources (e.g. O’Dea 1998) with a steep spectral slope at the high frequencies. To characterize the spectra, we also fit the data to a function similar to the synchrotron self-absorption model for the spherical homogeneous plasma. The function is

$$S_\nu(\nu) = \frac{S_t}{1 - \exp(-1)} \left(\frac{\nu}{\nu_t} \right)^{\alpha_r} \left\{ 1 - \exp \left[- \left(\frac{\nu}{\nu_t} \right)^{\alpha - \alpha_r} \right] \right\}, \quad (1)$$

where S_t is the flux density at the spectral turnover frequency ν_t , ν is the observing frequency, α is the spectral index in the optically thin region, and α_r is the spectral index of the rising region, and $\alpha_r = 2.5$ in the synchrotron self-absorption mode for the spherical homogeneous plasma. The least-square fitting gives $S_t = 262 \pm 31$ mJy, $\nu_t = 0.53 \pm 0.06$ GHz, $\alpha = -1.31 \pm 0.02$, $\alpha_r = 0.79 \pm 0.21$, and the reduced $\chi_r^2 = 1.01$.

3.2 Parsec-scale radio morphology

All the EVN imaging results of IRAS F11119+3257 are shown in Fig. 2. The parsec-scale radio morphology shows a strong dependence on the observing frequency, the image resolution, and sensitivity. Compared to the total flux densities predicted by our model, our VLBI images have restored nearly 100 percent at 1.66 GHz and about 50 percent at 4.93 GHz.




Figure 2. The two-sided but significantly beamed jet observed with the EVN in the highly accreting quasar IRAS F11119+3257. The yellow cross marks the optical centroid measured by *Gaia*. The first contours in all the images are at 3σ level. (a) The high-resolution 4.93-GHz image is made with natural weighting. The full width at half-maximum (FWHM) is $3.4 \text{ mas} \times 0.77 \text{ mas}$ at 12.0 deg . The contours are $0.07 \times (-1, 1, 2, 4, \dots, 32) \text{ mJy beam}^{-1}$. (b) The high-resolution 1.66-GHz image is made with uniform weighting. The FWHM is $5.4 \text{ mas} \times 2.0 \text{ mas}$ at 2.7 deg . The contours are $0.045 \times (-1, 1, 2, \dots, 512) \text{ mJy beam}^{-1}$. (c) The high-sensitivity 1.66-GHz image is made with a circular Gaussian beam. The FWHM is 7.5 mas . The contours are $0.02 \times (-1, 1, 2, 4, \dots, 2048) \text{ mJy beam}^{-1}$.

The 4.93-GHz image in the left-hand panel has the highest image resolution, up to 0.8 mas in the direction of East–west, because of the significant contribution of the Tianma 65-m radio telescope. Compared to the earlier imaging results from snapshot observations with the Very Long Baseline Array (VLBA) at 2.3 GHz (Petrov 2013), our image reveals the more details. Beside the peak component, there are a significant extension towards North and a few faint features with the decreasing brightness towards West. Because of the limited image sensitivity and uv coverage, the extension towards West is coincidentally similar to the typical one-sided core-jet structures observed in flat-spectrum blazars (e.g. Cheng et al. 2018). With respect to the phase-referencing calibrator, the image has quite precise position measurements. The optical centroid, reported by the *Gaia* DR2 (Brown et al. 2018), is marked as a yellow cross (J2000, RA = $11^{\text{h}}14^{\text{m}}38^{\text{s}}.89019$, Dec. = $32^{\circ}41'33''.4824$, $\sigma_{\text{ra}} = \sigma_{\text{dec}} = 0.1 \text{ mas}$, the astrometric excess noise: 0.4 mas). With respect to the optical centroid, the radio peak has an offset of about 18 mas .

There are more faint features recovered at 1.66 GHz in the middle and right-hand panels. As all the big antennas (Ef, Jb1, Ro, Sr, T6) are included, even a 4-h EVN observations still allows us to achieve an extremely high image sensitivity of $6.5 \mu\text{Jy beam}^{-1}$ (1σ) with natural weighting. To clearly reveal these faint features, we convolved the source model with a larger circular beam of 7.5 mas . The low-resolution image in the right-hand panel displays four relatively discrete components. According to their positions, they are marked as N, C, S, and W. In the middle panel, component S is resolved into S1 and S2, and component N is not seen because of the relatively high image resolution. Both components S1 and S2 have faint extensions in almost all the directions. There also exists significant continuous radio emission connecting components S and C. To quantitatively describe these components, we also fit them

with circular Gaussian models in DIFMAP 2.5e. The least-square fitting results including the formal 1σ uncertainties at the reduced $\chi_r^2 = 1$ are summarized in Table 3. The errors for the positions were truncated to 0.01 mas . The errors for the sizes were truncated to 0.1 mas . The empirical systematic uncertainties for S_{obs} and L_R are 5 per cent.

All the detected components in IRAS F11119+3257 have a brightness temperature of $\geq 1.0 \times 10^6 \text{ K}$ at 1.6 and 5 GHz . Component S1 has the highest brightness temperature reaching $3.2 \times 10^9 \text{ K}$ at 1.6 GHz . The next to last column in Table 3 reports T_b , estimated as (e.g. Condon et al. 1982)

$$T_b = 1.22 \times 10^9 \frac{S_{\text{obs}}}{\nu_{\text{obs}}^2 \theta_{\text{size}}^2} (1+z), \quad (2)$$

where S_{obs} is the total flux density in mJy, ν_{obs} is the observing frequency in GHz, θ_{size} is the FWHM of the circular Gaussian model in mas, and z is the redshift.

4 DISCUSSION

4.1 No evidence for the star formation activity

All these VLBI-detected components originate in the AGN activity instead of the star formation activity. Their radio emission is unlikely dominated by thermal emission of the star-forming activity because of their $T_B \gtrsim 10^6 \text{ K}$. It is also difficult to associate them with single young supernova or many supernovae remnants produced by the star formation activity. The radio luminosity $L_R = \nu L_\nu$ of each component is listed in the last column of Table 3. A rarely seen young supernova may reach a peak luminosity, $L_R \sim 5 \times 10^{38} \text{ erg s}^{-1}$ (e.g. Weiler et al. 2002), comparable to L_R of the two faint components N and W, while fail to explain their

Table 3. Circular Gaussian model-fitting results of the major components found in IRAS F11119+3257. Columns give (1) name, (2) observing frequency, (3) integrated flux density, (4–5) relative offsets in right ascension and declination with respect to the optical *Gaia* position, (6) size, (7) brightness temperature, and (8) radio luminosity.

Name	ν_{obs} (GHz)	S_{obs} (mJy)	$\Delta\alpha\cos\delta$ (mas)	$\Delta\delta$ (mas)	θ_{size} (mas)	T_{b} (K)	L_{R} (erg s ⁻¹)
N	1.66	0.26 ± 0.03	$+1.13 \pm 0.61$	$+36.55 \pm 0.56$	11.6 ± 1.3	1.0×10^6	4.3×10^{38}
C	1.66	1.28 ± 0.01	-1.75 ± 0.02	-1.73 ± 0.03	1.7 ± 0.1	2.4×10^8	2.1×10^{39}
S1	1.66	64.95 ± 0.01	-4.82 ± 0.01	-16.78 ± 0.01	3.3 ± 0.1	3.2×10^9	1.1×10^{41}
S2	1.66	11.43 ± 0.01	-11.14 ± 0.01	-16.00 ± 0.01	2.9 ± 0.1	3.9×10^8	1.9×10^{40}
W	1.66	0.29 ± 0.01	-19.83 ± 0.26	-4.54 ± 0.29	4.6 ± 0.6	7.2×10^6	4.8×10^{38}
S1	4.93	9.54 ± 0.07	-5.13 ± 0.02	-16.95 ± 0.01	2.7 ± 0.1	8.1×10^7	4.6×10^{40}
S2	4.93	1.67 ± 0.10	-11.71 ± 0.17	-14.30 ± 0.11	5.3 ± 0.1	3.5×10^6	8.1×10^{39}

extended structure. Additionally, because IRAS F11119+3257 is not a starburst galaxy (Komossa et al. 2006) like Arp 220 (e.g. Varenus et al. 2019), the two components cannot be composed of overlapping emission from supernova remnants in the nuclear region.

4.2 Non-detection of the flat-spectrum radio core

None of these VLBI-detected components can be identified as the flat-spectrum radio core of IRAS F11119+3257 because of their extended structure ($\theta_{\text{size}} \geq 1.7$ mas) and optically thin spectra. Component C is relatively close to the optical centroid. However, it has a steep spectrum: with flux densities of 1.28 mJy at 1.66 GHz, ≤ 0.07 mJy beam⁻¹ (3σ) at 4.93 GHz, $\alpha \leq -2.7$ ($S_{\nu} \propto \nu^{\alpha}$). The bright components S1 and S2 have a similar spectral index of $\alpha = -1.8$. The remaining components N and W are representative of the more extended structure and also have steep ($\alpha \leq -1.3$) spectra. Moreover, according to the map of the velocity gradient of the CO(1–0) line emission of the host galaxy (Veilleux et al. 2017), there exists a hint for a small offset between the zero-velocity component and the continuum source, i.e. component S.

The undetected radio core is most likely near component C. First, this allows us to naturally explain the extended conical structure formed by components C and S at 1.66 GHz. Secondly, this is consistent with the *Gaia* position. Generally, the *Gaia* positions are strongly dependent on the assumption of the point source structure. As a luminous quasar, IRAS F11119+3257 has a compact optical structure. This has been indirectly confirmed by the small astrometric excess noise ($1\sigma = 0.4$ mas) in the *Gaia* position. Finally, according to a statistical analysis of systematic differences in the positions of *Gaia* DR2 with respect to VLBI (Kovalev, Petrov & Plavin 2017), the separation ~ 18 mas between the radio peak and the optical centroid is too large to be explained as a potential systematic position error (≤ 10 mas).

The non-detection of the radio core allows us to set 3σ limits on its radio luminosity: $L_{\text{R}} \leq 3.4 \times 10^{38}$ erg s⁻¹ at 4.93 GHz and $L_{\text{R}} \leq 3.5 \times 10^{37}$ erg s⁻¹ at 1.66 GHz.

The radio core may be intrinsically faint. This is also in agreement with the extremely steep spectra between $\nu_{\text{t}} = 0.53 \pm 0.06$ and 96 GHz. At the low accretion rate state, there exists a correlation (e.g. Merloni, Heinz & Di Matteo 2003) among the radio core luminosity at 5 GHz, the X-ray luminosity (L_{X}) in the 2–10 keV, and the black hole mass (M_{BH})

$$\log L_{\text{R}} = (0.60_{-0.11}^{+0.11}) \log L_{\text{X}} + (0.78_{-0.09}^{+0.11}) \log M_{\text{BH}} + 7.33_{-4.07}^{+4.05} \quad (3)$$

According to the X-ray observations, $L_{\text{X}} = 3.8 \times 10^{42}$ erg s⁻¹ (Teng & Veilleux 2010) at the high accretion rate state. Since the high-state L_{X} is most likely an upper limit, it would predict $L_{\text{R}} \leq 10^{38.6 \pm 0.88}$ erg s⁻¹, comparable to our observational limits.

Another possible reason for the non-detection of the radio core may be its quenching after a transition from the low to high accretion rate states. By analogy with the unification evolution model for Galactic X-ray binaries (Fender, Belloni & Gallo 2004), and assuming that the jet radiative properties are scale free (e.g. Ruan et al. 2019), the radio cores in AGNs could be quenched when their accretion rates approach or exceed their Eddington accretion rates. Currently, the radio quiescence in the high state has been observed by Greene et al. (2006) in a sample of 19 low-mass galaxies with candidate massive black holes.

4.3 Constraints on the jet parameters

These components in IRAS F11119+3257 cannot be explained as the wide-angle outflows or winds. They have a ratio of $\frac{L_{\text{R}}}{L_{\text{X}}}$, at least one order of magnitude higher than the relation $\frac{L_{\text{R}}}{L_{\text{X}}} \sim 10^{-5}$ observed by Laor & Behar (2008) in the radio-quiet Palomar–Green quasar sample. Moreover, there is no diffuse biconical structure observed.

We can identify bright components C, S, and W as the approaching jet components with respect to the optical position. The non-detection of the radio core and the rapid drop of brightness and flux density in the inner component C indicates that they were from a relatively short-duration ejection event. Component N can be identified as a receding jet component or a relic jet because of its faintness.

The large flux density of component S may be mainly attributable to the Doppler boosting while the starkly lower flux density of component N to Doppler de-boosting thus resulting in the large flux density ratio. This involves approaching and receding components as opposed to them being expanding shocks. This is as there are no high brightness hot spots and edges, which can be brightened significantly by strong shocks, and, the linear polarization (indicative of an interaction at the interface between the shock and surrounding medium) is very low ($3\sigma \leq 1.3$ mJy) from the NVSS survey (Condon et al. 1998).

The apparent flux density ratio R_{flux} between approaching and receding jets emitting the same radio luminosity isotropically in their respective rest frames is (e.g. Böttcher, Harris & Krawczynski 2012)

$$R_{\text{flux}} = \frac{S_{\text{app}}}{S_{\text{rec}}} = \left(\frac{1 + \beta \cos \theta_{\text{v}}}{1 - \beta \cos \theta_{\text{v}}} \right)^{\eta}, \quad (4)$$




Figure 3. The relation between the flux density ratio $\frac{S_{\text{app}}}{S_{\text{rec}}}$ and the jet speed $\beta \cos \theta_v$ derived from equation (4). The cross marks the results of components S1 and N.

where S_{app} and S_{rec} are the flux densities of the approaching and receding components, respectively, β is the intrinsic jet speed in unit of the light speed c , θ_v is the jet viewing angle, $\eta = 3 - \alpha$ for a pair of discrete jet components and $\eta = 2 - \alpha$ for a continuous two-sided jet, and $\alpha = -1.31 \pm 0.02$ for our target. The relation is also plotted in Fig. 3. The arm length ratio R_{arm} between approaching and receding components is also affected by the Doppler beaming effect, while much weaker than the flux density ratio ($R_{\text{arm}}^\eta = R_{\text{flux}}$). Because the approaching jet might bend significantly and the radio core is not identified, the arm length ratio is not available to constrain the jet parameters.

We can constrain the jet parameters β and θ_v using equation (4). If component S1 and N are a pair of approaching and receding ejecta and $\eta = 3 - \alpha$, we can derive a ratio of 246 and a constraint of $\beta \cos \theta_v = 0.57c$. The flux density ratio is also marked as a cross in Fig. 3. The constraint gives $\beta \geq 0.57c$ and $\theta_v \leq 55$ deg. Owing to the observed high ratio, the constraint on $\beta \cos \theta_v$ is quite robust. It will not change significantly because the approaching jet may also include other much fainter components and $\eta = 2 - \alpha$. If component N is not the counter jet of component S1, the ratio will represent a lower limit and the limits on β and θ_v will remain valid. The small θ_v would allow us to simply explain the apparent wide opening angle (about 60 deg) and large change of the jet direction as a consequence of a projection effect.

Our limit corresponds to an accretion disc inclination of $\theta_i \leq 55$ deg assuming the jet is perpendicular to the accretion disc. As the lower limit of the jet speed is higher than that observed in the powerful X-ray outflows (Tombsi et al. 2015) and only a maximum jet speed of upto $0.5c$ is achievable for a radiative jet in a supercritical accretion disc (Sadowski & Narayan 2015), the jet components in IRAS F11119+3257 may not be driven by radiation pressure alone. These jet components are not subject to highly relativistic beaming since they have an extended morphology, relatively low T_b , and no high-energy γ -ray counterpart in the *Fermi* LAT 8-Year Point Source Catalogue (Abdollahi et al. 2020).

IRAS F11119+3257 has the highest jet speed among the known highly accreting objects. Super-Eddington accretions on to black holes are generally found in luminous AGNs, tidal disruption events (TDEs), and ultraluminous X-ray (ULX) sources. So far, all these supercritical accretion systems tend to have strong winds/outflows rather than relativistic jets. Among the TDEs, Swift J1644+57 had multiband powerful non-thermal emission but an absence of a collimated synchrotron jet (Yang et al. 2016a). The only known TDE jet, found by Mattila et al. (2018) in Arp 299-B, had a relatively low jet speed of about $0.2c$ on the pc-scale. Compared

to the TDEs, the highly accreting AGNs have a much more stable radio emission. Currently, there are only a few targets imaged by the VLBI observations (e.g. Giroletti et al. 2017; Yang et al. 2018). This is also as most of them are radio quiet (e.g. Panessa et al. 2007; Sikora et al. 2007) and have steep radio spectra (Laor et al. 2019), with a consequent lack of multi-epoch deep VLBI observations to constrain their jet speed. Additionally, extragalactic ULXs are rather faint for the VLBI observations to detect them (Yang et al. 2016b).

4.4 An unusual compact symmetric object

IRAS F11119+3257 may be identified as an uncommon compact symmetric object (CSO; e.g. Wilkinson et al. 1994; Readhead et al. 1996) from its two-sided jet morphology, compact size, and exceptionally high-flux density ratio. Most CSOs have faint radio cores and two-sided mini-lobes with projected sizes $\lesssim 1$ kpc and flux density ratios $\lesssim 10$ due to their jet speeds $\beta \lesssim 0.9c$ and jet viewing angles $\theta_v \gtrsim 45$ deg (e.g. Owsianik & Conway 1998; Polatidis & Conway 2003; An & Baan 2012). Compared to other CSOs (e.g. Sokolovsky et al. 2011; An et al. 2012), its jet has a lower T_b owing to its steep spectrum and relatively extended structure. Moreover, it cannot be taken as a representative CSO because of its especially high-flux density ratio.

According to the observed broad-band radio spectrum, IRAS F11119+3257 is a Gigahertz-peaked spectrum (GPS) source (e.g. Stanghellini et al. 1998) or a compact steep-spectrum source (CSS; e.g. O’Dea 1998). Among the GPS source, CSOs are frequently found (e.g. Xiang et al. 2005). According to the correlation relation between the linear size l_{jet} and the turnover frequency ν_t , $\log l_{\text{jet}} = -1.54 \log \nu_t - 0.32$, observed in GPS and CSS sources (O’Dea 1998), the observed turnover frequency allows us to set a 3σ upper limit of 10 kpc for its linear size. Using the lower limit of its jet speed ($0.57c$), we can derive an upper limit on its kinematic age, 6×10^5 yr. If these jets were launched at the early low accretion rate state, the age would set the first observational constraint on the time of the state transition of the accreting SMBH. Moreover, IRAS F11119+3257 is also a low-power or low-luminosity ($L_R \lesssim 10^{41}$ erg s $^{-1}$ at 1.4 GHz) compact radio source (Giroletti, Giovannini & Taylor 2005). It would require more energy injections for the compact source to become a large-scale Fanaroff–Riley type radio source (e.g. Kunert-Bajraszewska et al. 2010; An & Baan 2012).

The turnover in the broad-band radio spectrum of IRAS F11119+3257 might mainly result from synchrotron self-absorption. As the brightest component, component S1 has an average T_b from 8.1×10^7 K at 5 GHz to 3.2×10^9 K at 1.66 GHz. Assuming that $\log T_b$ and $\log \nu_{\text{obs}}$ are linearly related, component S would have an average T_b of 1.3×10^{11} K at $\nu_t = 0.53$ GHz. On the other hand, when a source is self-absorbed, the theoretical T_b in the emission rest frame is also $\sim 10^{11}$ K at the peak frequency (Readhead 1994). Thus, the inner part of component S1 might suffer significant synchrotron self-absorption.

5 CONCLUSIONS

As an optically luminous quasar, IRAS F11119+3257 hosts a supercritical accretion, the emission from which drives powerful X-ray outflows. Its radio spectrum between 0.15 and 96 GHz shows a peak at 0.53 ± 0.06 GHz and a steep slope of $\nu^{-1.31 \pm 0.02}$ in the optically thin part. From the EVN observations at 1.66 and 4.93 GHz, the quasar displays a two-sided jet with a projected separation of about 200 parsec. From the large flux density ratios

between the approaching and receding jet components, we inferred that the jet has an intrinsic speed of $\geq 0.57c$. This is higher than that observed in the X-ray winds and is thus unlikely to be driven by the radiation pressure alone. Among the known super-Eddington accretion systems, the jet in IRAS F11119+3257 has the highest speed. Moreover, we identified IRAS F11119+3257 as an unusual CSO with a jet viewing angle of $\leq 55^\circ$, a kinematic age of $\leq 6 \times 10^5$ yr, and a synchrotron self-absorbed radio spectrum.

ACKNOWLEDGEMENTS

This work was partly supported by the Square Kilometer Array (SKA) pre-research funding from the Ministry of Science and Technology of the People's Republic of China (2018YFA0404603) and the Chinese Academy of Sciences (CAS, No. 114231KYSB20170003). The European VLBI Network is a joint facility of independent European, African, Asian, and North American radio astronomy institutes. Scientific results from data presented in this publication are derived from the following EVN project codes: EY024 and RSY04. e-VLBI research infrastructure in Europe is supported by the European Union's Seventh Framework Programme (FP7/2007-2013) under grant agreement number RI-261525 NEXPreS. The National Radio Astronomy Observatory is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc. This work has made use of data from the European Space Agency (ESA) mission *Gaia* (<https://www.cosmos.esa.int/gaia>), processed by the *Gaia* Data Processing and Analysis Consortium (DPAC, <https://www.cosmos.esa.int/web/gaia/dpac/consortium>). Funding for the DPAC has been provided by national institutions, in particular the institutions participating in the *Gaia* Multilateral Agreement. We thank the staff of the GMRT that made these observations possible. The GMRT is run by the National Centre for Radio Astrophysics of the Tata Institute of Fundamental Research.

REFERENCES

- Abdollahi S. et al., 2020, *ApJS*, 247, 33
 An T., Baan W. A., 2012, *ApJ*, 760, 77
 An T. et al., 2012, *ApJS*, 198, 5
 Becker R. H., White R. L., Helfand D. J., 1995, *ApJ*, 450, 559
 Berton M. et al., 2018, *A&A*, 614, A87
 Böttcher M., Harris D. E., Krawczynski H., 2012, *Relativistic Jets from Active Galactic Nuclei*. Wiley, Weinheim
 Brown A. G. A. et al., 2018, *A&A*, 616, A1
 Cheng X.-P. et al., 2018, *ApJS*, 234, 17
 Colla G. et al., 1970, *A&AS*, 1, 281
 Condon J. J., Condon M. A., Gisler G., Puschell J. J., 1982, *ApJ*, 252, 102
 Condon J. J., Contton W. D., Greisen E. W., Yin Q. F., Perley R. A., Taylor G. B., Broderick J. J., 1998, *AJ*, 115, 1693
 Fender R. P., Belloni T. M., Gallo E., 2004, *MNRAS*, 355, 1105
 Giroletti M., Giovannini, Taylor G. B., 2005, *A&A*, 441, 89
 Giroletti M., Panessa F., Longinotti A. L., Krongold Y., Guainazzi M., Costantini E., Santos-Lleo M., 2017, *A&A*, 600, A87
 Greene J. E., Ho L. C., Ulvestad J. S., 2006, *ApJ*, 636, 56
 Gregory P. C., Scott W. K., Douglas K., Condon J. J., 1996, *ApJS*, 103, 427
 Greisen E. W., 2003, in Heck A., ed., *Astrophysics and Space Science Library*, Vol. 285, Information Handling in Astronomy: Historical Vistas. Kluwer, Dordrecht, p. 109
 Helmboldt J. F. et al., 2007, *ApJ*, 658, 203
 Intema H. T., Jagannathan P., Mooley K. P., Frail D. A., 2017, *A&A*, 598, A78
 Kawakatu N., Imanishi M., Nagao T., 2007, *ApJ*, 661, 660
 Keimpema A. et al., 2015, *Exp. Astron.*, 39, 259
 Komossa S., Voges W., Xu D., Mathur S., Adorf H.-M., Lemson G., Duschl W. J., Grupe D., 2006, *AJ*, 132, 531
 Kormendy J., Ho L. C., 2013, *ARA&A*, 51, 511
 Kovalev Y. Y., Petrov L., Plavin A. V., 2017, *A&A*, 598, L1
 Kunert-Bajraszewska M., Gawroński M. P., Labiano A., Siemiginowska A., 2010, *MNRAS*, 408, 2261
 Laor A., Behar E., 2008, *MNRAS*, 390, 847
 Laor A., Baldi R. D., Behar E., 2019, *MNRAS*, 482, 5513
 Marziani P., Sulentic J. W., 2014, *MNRAS*, 442, 1211
 Mattila S. et al., 2018, *Science*, 361, 482
 Merloni A., Heinz S., Di Matteo T., 2003, *MNRAS*, 345, 1057
 Nagar N. M., Wilson A. S., Falcke H., Veilleux S., Maiolino R., 2003, *A&A*, 409, 115
 Nardini E. et al., 2015, *Science*, 347, 860
 O'Dea C. P., 1998, *PASP*, 110, 493
 Owsianik I., Conway J. E., 1998, *A&A*, 337, 69
 Pacucci F., Volonteri M., Ferrara A., 2015, *MNRAS*, 452, 1922
 Panessa F., Barcons X., Bassani L., Cappi M., Carrera F. J., Ho L. C., Pellegrini S., 2007, *A&A*, 467, 519
 Panessa F., Baldi R. D., Laor A., Padovani P., Behar E., McHardy I., 2019, *Nat. Astron.*, 3, 387
 Petrov L., 2013, *AJ*, 146, 5
 Polatidis A. G., Conway J. E., 2003, *Publ. Astron. Soc. Aust.*, 20, 69
 Prusti T. et al., 2016, *A&A*, 595, A1
 Readhead A. C. S., 1994, *ApJ*, 426, 51
 Readhead A. C. S., Taylor G. B., Xu W., Pearson T. J., Wilkinson P. N., Polatidis A. G., 1996, *ApJ*, 460, 612
 Rengelink R. B., Tang Y., de Bruyn A. G., Miley G. K., Bremer M. N., Roettgering H. J. A., Bremer M. A. R., 1997, *A&AS*, 124, 259
 Ruan J. J., Anderson S. F., Eracleous M., Green P. J., Haggard D., MacLeod C. L., Runnøe J. C., Sobolewska M. A., 2019, *ApJ*, 883, 76
 Sadowski A., Narayan R., 2015, *MNRAS*, 453, 3213
 Shepherd M. C., Pearson T. J., Taylor G. B., 1994, *BAAS*, 26, 987
 Sikora M., Stawarz Ł., Lasota J.-P., 2007, *ApJ*, 658, 815
 Sokolovsky K. V., Kovalev Y. Y., Pushkarev A. B., Mimica P., Perucho M., 2011, *A&A*, 535, A24
 Stanghellini C., O'Dea C. P., Dallacasa D., Baum S. A., Fanti R., Fanti C., 1998, *A&AS*, 131, 303
 Teng S. H., Veilleux S., 2003, *ApJ*, 725, 1848
 Tombesi F., 2016, *Astron. Nachr.*, 337, 410
 Tombesi F., Meléndez M., Veilleux S., Reeves J. N., González-Alfonso E., Reynolds C. S., 2015, *Nature*, 519, 436
 Tombesi F., Veilleux S., Meléndez M., Lohfink A., Reeves J. N., Piconcelli E., Fiore F., Feruglio C., 2017, *ApJ*, 850, 151
 Varenus E. et al., 2019, *A&A*, 623, A173
 Veilleux S. et al., 2013, *ApJ*, 776, 27
 Veilleux S., Bolatto A., Tombesi F., Meléndez M., Sturm E., González-Alfonso E., Fischer J., Rupke D. S. N., 2017, *ApJ*, 843, 18
 Volonteri M., Silk J., Dubus G., 2015, *ApJ*, 804, 148
 Wang J.-M., Du P., Valls-Gabaud D., Hu C., Netzer H., 2013, *Phys. Rev. Lett.*, 110, 081301
 Weiler K. W., Panagia N., Montes M. J., Sramek R. A., 2002, *ARA&A*, 40, 387
 Wilkinson P. N., Polatidis A. G., Readhead A. C. S., Xu W., Pearson T. J., 1994, *MNRAS*, 269, 67
 Xiang L., Dallacasa D., Cassaro P., Jiang D., Reynolds C., 2005, *A&A*, 434, 123
 Yang J., Paragi Z., van der Horst A. J., Gurvits L. I., Campbell R. M., Giannios D., An T., Komossa S., 2016a, *MNRAS*, 462, L66
 Yang X. et al., 2016b, *MNRAS*, 464, L70
 Yang J., An T., Zheng F., Baan W. A., Paragi Z., Mohan P., Zhang Z., Liu X., 2018, *MNRAS*, 482, 1701

This paper has been typeset from a \LaTeX file prepared by the author.