



## Non-Markovian Steady States of a Driven Two-Level System

Downloaded from: <https://research.chalmers.se>, 2025-12-09 00:50 UTC

Citation for the original published paper (version of record):

Josefsson Ask, A., Johansson, G. (2022). Non-Markovian Steady States of a Driven Two-Level System. Physical Review Letters, 128(8). <http://dx.doi.org/10.1103/PhysRevLett.128.083603>

N.B. When citing this work, cite the original published paper.

# Non-Markovian Steady States of a Driven Two-Level System

Andreas Ask and Göran Johansson 

*Department of Microtechnology and Nanoscience (MC2), Chalmers University of Technology, SE-41296 Göteborg, Sweden*



(Received 11 March 2021; revised 20 December 2021; accepted 1 February 2022; published 23 February 2022)

We show that an open quantum system in a non-Markovian environment can reach steady states that it cannot reach in a Markovian environment. As these steady states are unique for the non-Markovian regime, they could offer a simple way of detecting non-Markovianity, as no information about the system's transient dynamics is necessary. In particular, we study a driven two-level system (TLS) in a semi-infinite waveguide. Once the waveguide has been traced out, the TLS sees an environment with a distinct memory time. The memory time enters the equations as a time delay that can be varied to compare a Markovian to a non-Markovian environment. We find that some non-Markovian states show exotic behaviors such as population inversion and steady-state coherence beyond  $1/\sqrt{8}$ , neither of which is possible for a driven TLS in the Markovian regime, where the time delay is neglected. Additionally, we show how the coherence of quantum interference is affected by time delays in a driven system by extracting the effective Purcell-modified decay rate of a TLS in front of a mirror.

DOI: [10.1103/PhysRevLett.128.083603](https://doi.org/10.1103/PhysRevLett.128.083603)

There are no truly closed quantum systems. In one way or another, a quantum system is always in contact with a noisy environment and will inevitably lose its quantum properties [1]. If the dynamics are Markovian in nature, the environment can be considered memoryless, and there is no backflow of information. Such systems are described by a quantum dynamical semigroup, whose generator governs a master equation in Lindblad form [2,3]. In many realistic systems, the requirements for a Markovian time evolution, such as weak interaction and short environment correlation times, are not satisfied, and long-time memory effects of the environment influence the system dynamics. In what ways such interaction affects the evolution of a quantum system is not only interesting from a fundamental perspective, but could also prove useful to probe properties of the environment [4,5], and ultimately lead to a better understanding of the decoherence mechanisms of quantum systems [6].

Although one cannot translate the classical definition of a Markov process directly to the quantum regime [7], several definitions and corresponding measures of non-Markovianity for open quantum systems have been introduced [8–13]. These measures are all constructed to detect deviations from Markovianity by characterizing the system's transient dynamics. To our knowledge, the question whether a non-Markovian environment can be distinguished from the

steady-state of a driven system has not been addressed. That would not only simplify the characterization of non-Markovianity in such cases, but it is also an interesting question in itself.

In this Letter, we show that a driven two-level quantum system coupled to a non-Markovian environment can reach a unique set of steady states that are out of reach for the system coupled to a Markovian environment. We call these states “non-Markovian steady states,” as they can be distinguished from *any* state in the Markovian regime. These states are different from the oscillatory non-Markovian steady states for undriven systems discussed, e.g., in Refs. [14] and [15]. To quantify these steady states' uniqueness, we propose a measure based on trace distance and distinguishability [16]. We note that our measure does not attempt to quantify the degree of non-Markovianity in these systems but rather gives a quantitative measure on how easily one can distinguish these states from the states in the Markovian regime.

To demonstrate when non-Markovian steady states can occur, we study a driven two-level system (TLS) in a semi-infinite waveguide (an atom in front of a mirror) [17–25], see Fig. 1. The drive amplitude and the system's coupling strength to the waveguide are taken as fixed parameters throughout the system evolution. Once the waveguide has been traced out, the distance to the mirror gives the environment seen by the TLS a distinct memory time. This memory time enters the equations for the system dynamics as a time delay, which can be set to zero for comparison between a Markovian and a non-Markovian environment. Thus, the physical origin of any non-Markovian effects in this system has an easy interpretation in terms of coherent quantum feedback. Additionally, an atom in front of a mirror has been realized with both artificial

*Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/) license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by [Bibsam](https://www.bibsam.se/).*




FIG. 1. (a) Schematic of an atom in front of a mirror. Photons emitted to the left are reflected from the mirror and interacts with the atom again, giving the atom's environment an effective memory time. (b) Representation of one time step in the evolution of the setup in (a) as a MPS. The waveguide is represented by time bins (gray), which moves in a conveyor belt fashion one time step,  $\Delta t$ , at a time, and interacts with the atom (turquoise) twice.

and natural atoms in a variety of systems already [26–32], so the physics discussed in this Letter could be further investigated experimentally immediately. We also note that a similar system to the atom in front of a mirror (in fact, they are fully mappable to each other in some parameter regimes) is the giant atom [15,33–36], which was recently realized in both the Markovian [37,38] and non-Markovian regime [39].

Despite being an archetypical quantum-optical model system for decades [17], the atom in front of a mirror has remained a hard system to simulate without resorting to substantial approximations. The propagation time delay between the atom and the mirror prohibits a treatment based on Markovian master equations, and earlier work on non-Markovianity has been limited to either few excitations [40,41] or short timescales [42,43]. It was only recently that Pichler *et al.* [23] proposed a method based on matrix product states (MPSs) which could allow the system to be integrated all the way to steady state, while still allowing for many excitations in the feedback loop and long delay times. There, it was shown that the time delay does in fact alter the steady state of the atom. However, that does not imply that the state is *unique* for the non-Markovian regime, i.e., that the same state cannot be reached using a different drive strength and neglecting the time delay. It could also happen that a non-Markovian environment reduces the purity of the steady state, in which case the effect of the non-Markovian environment could be captured by adding additional pure dephasing to a fully Markovian treatment. In fact, we find that the non-Markovian environment mostly produces steady states one cannot distinguish from those reachable in the Markovian environment. For some system parameters, however, we find non-Markovian steady states that are not only unique for the non-Markovian regime but also

show exotic behaviors such as population inversion in the TLS or steady-state coherence beyond  $1/\sqrt{8}$ , neither of which is possible in the Markovian regime.

**Definition of non-Markovian steady states.**—In the Markovian regime, the dynamics of the TLS is given by the Markovian master equation in Lindblad form [2,17,26],

$$\dot{\rho}_M = -i[H_{\text{TLS}}, \rho] + \frac{\gamma}{2}(2\sigma^- \rho \sigma^+ - \{\sigma^+ \sigma^-, \rho\}) + \gamma_\phi(2\sigma^+ \sigma^- \rho \sigma^+ \sigma^- - \{\sigma^+ \sigma^-, \rho\}), \quad (1)$$

where  $\gamma_\phi$  is a pure dephasing rate,  $\gamma$  is a renormalized decay rate due to the mirror  $\gamma = 2\gamma' \cos(\phi)$ , where  $\gamma'$  is the bare decay rate (without the mirror),  $\phi$  is a phase shift,  $H_{\text{TLS}} = \Delta \sigma^+ \sigma^- + (\Omega/2)(\sigma^+ + \sigma^-)$ , where  $\Delta = \omega_d - \omega_0$  is the detuning between the TLS transition frequency  $\omega_0$  and the drive frequency  $\omega_d$ ,  $\Omega$  is the amplitude of the driving field, and  $\sigma^+$  ( $\sigma^-$ ) creates (annihilates) an excitation in the TLS. The solutions to Eq. (1) yield an elliptical area in the Bloch sphere of possible steady states, whose outer boundary is given by  $\Omega/\gamma = [0, \infty]$  and  $\gamma_\phi = 0$ , see Fig. 2. The mirror is thus irrelevant for determining possible steady states; it only rescales the decay rate. Finite  $\Delta$  does not change the possible steady states and for that reason we always consider resonant driving throughout the Letter,  $\Delta = 0$ . We let  $\rho_M(\gamma, \gamma_\phi)$  denote the steady state solution to Eq. (1) for a fixed drive strength, and  $\rho$  denote the steady-state reduced density matrix of the TLS in a non-Markovian environment. Then, the ability to distinguish the non-Markovian regime from the Markovian regime can be captured by




FIG. 2. Markovian versus non-Markovian regimes of steady states for a driven TLS in front of a mirror. By neglecting the time delay, the system evolves according to a Markovian master equation and can only reach states lying either on the solid black line (for no pure dephasing) or its inside (with dephasing). If the time delay is taken into account the system can reach steady states which are, e.g., outside of the Markovian regime (green crosses), precisely on the boundary between the two regimes (magenta triangles), or well inside the Markovian regime (blue circles). In all cases the following parameters were used:  $\gamma = \gamma_L + \gamma_R = 1$ ,  $\gamma_{L/R} = \gamma/2$ , and  $\Omega/\gamma = [0.1, 4]$  ( $\Omega/\gamma = [0.1, 3.5]$  for  $\phi = \pi/2$ ).

$$\mathcal{N}_{ss} = \min_{\gamma, \gamma_\phi} T[\rho, \rho_M(\gamma, \gamma_\phi)], \quad (2)$$

where  $T[\rho, \rho_M] = \frac{1}{2} \text{Tr} \sqrt{(\rho - \rho_M)^\dagger (\rho - \rho_M)}$  is the trace distance between  $\rho$  and  $\rho_M$ . As the trace distance is closely related to the *distinguishability* of quantum states [16,44], we define a steady state as non-Markovian if  $\mathcal{N}_{ss} > 0$ . With this definition, a non-Markovian steady state is a state that is unique for the system in a non-Markovian environment. For the atom in front of a mirror, we show that most non-Markovian steady states would correctly be classified as belonging to a non-Markovian system according to the definition of non-Markovianity in Ref. [9]. However, we note that a large degree of non-Markovianity does not necessarily correspond to a large  $\mathcal{N}_{ss}$ .

*Model.*—To model a non-Markovian environment, we put the driven TLS in a semi-infinite waveguide, see Fig. 1(a). After the waveguide has been traced out, the distance to the mirror gives the environment of the TLS a distinct memory time,  $\tau$ . Emission towards the mirror enters a coherent feedback loop, in which it picks up a propagation phase,  $\phi$ , that in our calculation includes any extra phase shift imposed by the field's boundary condition at the mirror. In a frame rotating with the drive frequency,  $\omega_d$ , the total Hamiltonian has two parts in the interaction picture  $H(t) = H_{\text{TLS}} + H_{\text{int}}(t)$ , where the interaction Hamiltonian

$$H_{\text{int}}(t) = i(\sqrt{\gamma_L} b_L^\dagger(t) + \sqrt{\gamma_R} b_R^\dagger(t)) \sigma^- + \text{H.c.}, \quad (3)$$

can be rewritten in terms of a single bath operator,  $b(t)$ , since the mirror couples left- and right-going modes  $b_R(t) = b_L(t - \tau) e^{i\phi}$ ,

$$H_{\text{int}}(t) = i[\sqrt{\gamma_L} b^\dagger(t) + \sqrt{\gamma_R} b^\dagger(t - \tau) e^{i\phi}] \sigma^- + \text{H.c.}, \quad (4)$$

where  $\gamma_L$  and  $\gamma_R$  denotes the decay rate into left ( $L$ )- and right ( $R$ )-going modes in the waveguide, respectively,  $\phi$  is the phase shift acquired by a photon (or phonon) traveling to the mirror and back. The phase shift is in fact related to the drive frequency,  $\phi = \omega_d \tau$ , but we keep it as an independent variable in order to study the effect of the phase shift and delay time separately. The bosonic operator,  $b(t)$ , obeys the quantum white-noise commutation relation  $[b(t), b^\dagger(t')] = \delta(t - t')$ , and is defined as  $b(t) = \int_B d\omega b(\omega) \exp[-i(\omega - \omega_d)t]$ , where  $b^\dagger(\omega)$  [ $b(\omega)$ ] creates [annihilates] a photon at frequency  $\omega$ , satisfying the commutation relation  $[b(\omega), b^\dagger(\omega')] = \delta(\omega - \omega')$ . A full derivation of the interaction Hamiltonian in Eq. (4) can be found in Ref. [23]. The interpretation, however, is clear: the TLS interacts with a bosonic bath at time  $t$ ; after some time  $\tau$  the TLS interacts with the bath again; the state of the bath at this later time has to be the state of the bath at an earlier time  $t - \tau$ , taking into account the traveling phase acquired during this time.

The system dynamics is calculated by solving the quantum stochastic Schrödinger equation (QSSE)

$$i \frac{d}{dt} |\Psi(t)\rangle = H(t) |\Psi(t)\rangle, \quad (5)$$

using the MPS method formulated in Ref. [23]. Matrix product states have shown to be efficient representations of 1D many-body systems [45–51]. Solving the QSSE is turned into a many-body problem by discretizing time,  $t_n = n\Delta t$ , turning Eq. (5) into a dynamical map  $|\Psi(t_{n+1})\rangle = U_n |\Psi(t_n)\rangle$ . Throughout the Letter we use a time step much smaller than all other timescales involved:  $\Delta t \ll \{1/\gamma, 1/\Omega\}$ . The state of the field in the waveguide is represented by time bins, with associated bosonic noise increments  $\Delta B(t_n) = \int_{t_n}^{t_{n+1}} b(t) dt$ , which fulfills the commutation relation  $[\Delta B(t_n), \Delta B^\dagger(t_{n'})] = \Delta t \delta_{n,n'}$ . The operator  $\Delta B^\dagger(t_n)$  can thus be seen as a creation operator for time bin  $n$ , with a corresponding Fock state defined as  $|n\rangle_I \equiv [(\Delta B_I^\dagger)^n / \sqrt{n! \Delta t^n}] |\text{vac}\rangle_I$ . The unitary,  $U_n$ , is written in this time-bin formulation as

$$U_n = \exp[iH_{\text{TLS}} \Delta t + (\sqrt{\gamma_L} \Delta B^\dagger(t_n) \sigma^- + \sqrt{\gamma_R} \Delta B^\dagger(t_{n-k}) e^{i\phi} \sigma^- - \text{H.c.})], \quad (6)$$

here  $t_k = k\Delta t$  is the feedback time  $\tau$ . The total quantum state at time  $t_n$ , for both the bath and the TLS, is then written as

$$|\Psi(t_n)\rangle = \sum_{i_T, i_1, \dots, i_N} \Psi_{i_T, i_1, \dots, i_N}(t_n) |i_T\rangle \otimes |i_1\rangle \otimes \dots \otimes |i_N\rangle, \quad (7)$$

where  $t_N = N\Delta t$  is the total integration time,  $i_T$  denotes the state of the TLS, and  $i_j$  is the photon number in time bin  $j$ . The initial state is written as a MPS ansatz

$$\Psi_{i_S, i_1, \dots, i_N}(t_0) = M[S]^{i_T} M[1]^{i_1} \dots M[N]^{i_N}, \quad (8)$$

where  $M[j]^{i_j}$  is a matrix of dimension  $D_j \times D_{j+1}$ . The maximum matrix dimension  $D_{\text{max}}$  in the MPS chain is referred to as the bond dimension and sets an upper limit to the amount of entanglement in the system, which in our system depends on the length of the feedback loop [23]. We use a bond dimension of  $D_{\text{max}} = 32$  for the moderate time delays considered here. The state amplitudes are updated in each time step using standard MPS techniques [45,52].

*Non-Markovian regime.*—Two parameters are important for quantifying the significance of the feedback:  $\gamma\tau$ , where  $\gamma = \gamma_L + \gamma_R$ , and  $\Omega\tau$ . Only when both  $\gamma\tau \ll 1$  and  $\Omega\tau \ll 1$  does the system not have time to evolve during the feedback time, and a Markovian treatment is valid. The situation is thus different from the nondriven TLS in which the single parameter  $\gamma\tau$  determines the non-Markovian properties alone.






FIG. 3. Non-Markovianity of the steady state evaluated using the measure introduced in Eq. (2). A phase shift of  $\phi = 0$  was used unless it is stated otherwise in the figure. A large  $\mathcal{N}_{ss}$  means that the state is easily distinguishable from any state in the Markovian regime.

We first study the effect of the time delay, and set  $\phi = 0$ . The feedback is thus in phase with the drive and they interfere constructively. In all calculations that follow we use  $\gamma = 2\gamma_L = 2\gamma_R = 1$ . In Fig. 2 we plot the steady-state reduced density-matrix elements of the TLS for drive strengths in the range  $\Omega/\gamma = [0.1, 4]$  for  $\gamma\tau = 0.5$  (green crosses) and  $\gamma\tau = 3$  (magenta triangles). The trace distance to the closest Markovian steady state,  $\mathcal{N}_{ss}$  [Eq. (2)], is plotted for a greater variety of time delays in Fig. 3. From these two figures we make the following observations: (i) When the driving is weak, the steady state cannot be distinguished from a Markovian steady state, independent of time delay. (ii) The states start to deviate from the Markovian regime initially for  $\gamma\tau > 0$ , reach a maximum deviation at  $\gamma\tau \approx 0.5$ , and then start to approach the Markovian regime again. For sufficiently long time delays the states cannot be distinguished from a Markovian state anymore. (iii) For  $\gamma\tau = 0.5$  the TLS shows both population inversion,  $\rho_{ee} > 1/2$ , for sufficiently strong driving, and larger coherence than what is possible in the Markovian regime,  $|\rho_{eq}| > 1/\sqrt{8}$ . The oscillatory behavior that can be seen in both Figs. 2 and 3 is due to the additional phase shift,  $\Omega\tau$ , induced by the drive, primarily noticeable for the longer time delays. We also note that Fig. 3 could serve as a mean to extract the delay time experimentally by using  $\tau$  as a fitting parameter, which could be easier than, e.g., measuring two-time correlation functions.

When the phase shift deviates from 0 mod  $2\pi$ , the states either approach or fall well inside the Markovian regime (blue circles in Fig. 2). Inside the Markovian regime  $\mathcal{N}_{ss} = 0$  per definition.

Coherent quantum interference phenomena play an important role in many applications in waveguide QED. It is well known, e.g., that the mirror doubles the TLS' decay rate to the waveguide due to the Purcell effect in the Markovian regime (for  $\phi = 0$ ). Increasing the distance to the mirror reduces the coherence of the radiation that comes back to the TLS. The longer time it takes for the feedback to




FIG. 4. Effective decay rate for  $\phi = 0$  as a function of drive strength for various time delays.

come back, the higher chance for spontaneous emission to occur in the TLS. For long enough time delay, the TLS should behave as if it was positioned in an infinite waveguide instead, without the mirror present. This is precisely what we observe in Fig. 4. We extract the “effective” decay rate by calculating the ratio between the output time-bin population and the TLS population,  $\gamma_{\text{eff}} = \langle \Delta B_{\text{out}}^\dagger \Delta B_{\text{out}} \rangle_{ss} / \rho_{ee}$ . For weak driving we observe that the decay rate is not affected by the time delay, as the drive strength increases, however, it approaches the expected value of  $\gamma$ . We note that the effective decay rate could have been extracted from the master equation in Eq. (1) if negative dephasing rates were allowed. In fact, all the non-Markovian steady states seen in Fig. 2 could be described by the master equation with a negative dephasing rate. Since negative dephasing rates have been used to describe temporary increases in quantum coherence for non-Markovian systems [53,54], we find it interesting to note that such effects can persist all the way to the steady state.

*Non-Markovianity measure of the transient dynamics.*— Finally, we make the remark that maximizing  $\mathcal{N}_{ss}$  is not necessarily the same thing as maximizing the non-Markovianity of the system in a traditional sense. For this




FIG. 5. Non-Markovianity according to the measure in Eq. (9) as a function of time delay. Dashed lines are for the same drive strengths as written in the figure but for  $\phi = \pi/2$ . By comparing with Fig. 2 we conclude that a large non-Markovianity does not correspond to a steady state that can be distinguished from the Markovian regime.

argument, we compare  $\mathcal{N}_{ss}$  to the measure proposed in Ref. [9],

$$\mathcal{N} = \max_{\rho_{1,2}} \int_{\sigma>0} dt \sigma(t, \rho_{1,2}), \quad (9)$$

where  $\sigma(t, \rho_{1,2}) = (d/dt)T[\rho_1(t), \rho_2(t)]$ , and  $T[\rho_1, \rho_2]$  denotes the trace distance. The maximum is taken over all pairs of initial states. For our system, we can safely choose the ground and excited states as the two initial states [55]. We plot  $\mathcal{N}$  as a function of time delay for various drive strengths in Fig. 5 for both  $\phi = 0$  (solid lines) and  $\phi = \pi/2$  (dashed lines). In Fig. 3, we saw the largest  $\mathcal{N}_{ss}$  for  $\gamma\tau \approx 0.5-1$ , which is barely non-Markovian according to  $\mathcal{N}$ , whereas longer time delays and stronger driving increases  $\mathcal{N}$  significantly.

**Conclusion.**—We have introduced the concept of non-Markovian steady states as a set of steady states unique for open quantum systems in a non-Markovian environment. The system cannot reach these states if it is coupled to a Markovian environment and could thus offer a simple way of detecting non-Markovianity as only a steady state measurement is required. Moreover, we introduced an appropriate measure for these states' uniqueness based on the trace distance to the closest Markovian steady state. As an example, we show that non-Markovian steady states occur in a driven TLS in a semi-infinite waveguide. Among the non-Markovian steady states, we find states with population inversion in the TLS, or steady state coherence larger than  $1/\sqrt{8}$ , two impossible scenarios in the Markovian regime. Additionally, we showed that time delay could have a detrimental effect on coherent quantum interference in waveguides by extracting the effective Purcell-modified decay rate as a function of time delay and drive strength.

Further studies need to be made in order to generalize the findings of this work to more general non-Markovian environments which could be of practical importance, e.g., in the field of quantum optics. Here, we specially note the Redfield theory, which can be used to capture non-Markovian effects for time delays comparable to the system relaxation timescale [56,57], which is indeed where we find non-Markovian steady states.

We thank Arne L. Grimsmo for valuable discussions, and acknowledge funding from the Swedish Research Council (VR) through Grant No. 2017-04197. G.J. also acknowledges funding from the Knut and Alice Wallenberg Foundation (KAW) through the Wallenberg Centre for Quantum Technology (WACQT).

- 
- [1] H. P. Breuer and F. Petruccione, *The Theory of Open Quantum Systems* (OUP, Oxford, 2007).
  - [2] G. Lindblad, On the generators of quantum dynamical semigroups, *Commun. Math. Phys.* **48**, 119 (1976).

- [3] V. Gorini, A. Kossakowski, and E. C. G. Sudarshan, Completely positive dynamical semigroups of N-level systems, *J. Math. Phys. (N.Y.)* **17**, 821 (1976).
- [4] S. Cialdi, A. Smirne, M. G. A. Paris, S. Olivares, and B. Vacchini, Two-step procedure to discriminate discordant from classical correlated or factorized states, *Phys. Rev. A* **90**, 050301(R) (2014).
- [5] M. Gessner, M. Ramm, T. Pruttivarasin, A. Buchleitner, H.-P. Breuer, and H. Häffner, Local detection of quantum correlations with a single trapped ion, *Nat. Phys.* **10**, 105 (2014).
- [6] H.-P. Breuer, E.-M. Laine, J. Piilo, and B. Vacchini, Colloquium: Non-Markovian dynamics in open quantum systems, *Rev. Mod. Phys.* **88**, 021002 (2016).
- [7] B. Vacchini, A. Smirne, E.-M. Laine, J. Piilo, and H.-P. Breuer, Markovianity and non-Markovianity in quantum and classical systems, *New J. Phys.* **13**, 093004 (2011).
- [8] M. M. Wolf, J. Eisert, T. S. Cubitt, and J. I. Cirac, Assessing Non-Markovian Quantum Dynamics, *Phys. Rev. Lett.* **101**, 150402 (2008).
- [9] H.-P. Breuer, E.-M. Laine, and J. Piilo, Measure for the Degree of Non-Markovian Behavior of Quantum Processes in Open Systems, *Phys. Rev. Lett.* **103**, 210401 (2009).
- [10] A. Rivas, S. F. Huelga, and M. B. Plenio, Entanglement and Non-Markovianity of Quantum Evolutions, *Phys. Rev. Lett.* **105**, 050403 (2010).
- [11] S. C. Hou, X. X. Yi, S. X. Yu, and C. H. Oh, Alternative non-Markovianity measure by divisibility of dynamical maps, *Phys. Rev. A* **83**, 062115 (2011).
- [12] S. Luo, S. Fu, and H. Song, Quantifying non-Markovianity via correlations, *Phys. Rev. A* **86**, 044101 (2012).
- [13] R. Vasile, S. Maniscalco, M. G. A. Paris, H.-P. Breuer, and J. Piilo, Quantifying non-Markovianity of continuous-variable Gaussian dynamical maps, *Phys. Rev. A* **84**, 052118 (2011).
- [14] R. Finsterhölzl, M. Katzer, and A. Carmele, Nonequilibrium non-Markovian steady states in open quantum many-body systems: Persistent oscillations in Heisenberg quantum spin chains, *Phys. Rev. B* **102**, 174309 (2020).
- [15] L. Guo, A. F. Kockum, F. Marquardt, and G. Johansson, Oscillating bound states for a giant atom, *Phys. Rev. Research* **2**, 043014 (2020).
- [16] C. A. Fuchs and J. van de Graaf, Cryptographic distinguishability measures for quantum-mechanical states, *IEEE Trans. Inf. Theory* **45**, 1216 (1999).
- [17] U. Dorner and P. Zoller, Laser-driven atoms in half-cavities, *Phys. Rev. A* **66**, 023816 (2002).
- [18] T. Tufarelli, F. Ciccarello, and M. S. Kim, Dynamics of spontaneous emission in a single-end photonic waveguide, *Phys. Rev. A* **87**, 013820 (2013).
- [19] D. Meschede, W. Jhe, and E. A. Hinds, Radiative properties of atoms near a conducting plane: An old problem in a new light, *Phys. Rev. A* **41**, 1587 (1990).
- [20] A. Beige, J. Pachos, and H. Walther, Spontaneous emission of an atom in front of a mirror, *Phys. Rev. A* **66**, 063801 (2002).
- [21] K. Koshino and Y. Nakamura, Control of the radiative level shift and linewidth of a superconducting artificial atom through a variable boundary condition, *New J. Phys.* **14**, 043005 (2012).

- [22] Y. Wang, J. Minář, G. Hétet, and V. Scarani, Quantum memory with a single two-level atom in a half cavity, *Phys. Rev. A* **85**, 013823 (2012).
- [23] H. Pichler and P. Zoller, Photonic Circuits with Time Delays and Quantum Feedback, *Phys. Rev. Lett.* **116**, 093601 (2016).
- [24] H. Pichler, S. Choi, P. Zoller, and M. D. Lukin, Universal photonic quantum computation via time-delayed feedback, *Proc. Natl. Acad. Sci. U.S.A.* **114**, 11362 (2017).
- [25] E. Wiegand, B. Rousseaux, and G. Johansson, Semiclassical analysis of dark-state transient dynamics in waveguide circuit QED, *Phys. Rev. A* **101**, 033801 (2020).
- [26] J. Eschner, C. Raab, F. Schmidt-Kaler, and R. Blatt, Light interference from single atoms and their mirror images, *Nature (London)* **413**, 495 (2001).
- [27] M. A. Wilson, P. Bushev, J. Eschner, F. Schmidt-Kaler, C. Becher, R. Blatt, and U. Dörner, Vacuum-Field Level Shifts in a Single Trapped Ion Mediated by a Single Distant Mirror, *Phys. Rev. Lett.* **91**, 213602 (2003).
- [28] F. Dubin, D. Rotter, M. Mukherjee, C. Russo, J. Eschner, and R. Blatt, Photon Correlation versus Interference of Single-Atom Fluorescence in a Half-Cavity, *Phys. Rev. Lett.* **98**, 183003 (2007).
- [29] I. C. Hoi, A. F. Kockum, L. Tornberg, A. Pourkabirian, G. Johansson, P. Delsing, and C. M. Wilson, Probing the quantum vacuum with an artificial atom in front of a mirror, *Nat. Phys.* **11**, 1045 (2015).
- [30] P. Y. Wen, A. F. Kockum, H. Ian, J. C. Chen, F. Nori, and I.-C. Hoi, Reflective Amplification without Population Inversion from a Strongly Driven Superconducting Qubit, *Phys. Rev. Lett.* **120**, 063603 (2018).
- [31] P. Y. Wen, K.-T. Lin, A. F. Kockum, B. Suri, H. Ian, J. C. Chen, S. Y. Mao, C. C. Chiu, P. Delsing, F. Nori, G.-D. Lin, and I.-C. Hoi, Large Collective Lamb Shift of Two Distant Superconducting Artificial Atoms, *Phys. Rev. Lett.* **123**, 233602 (2019).
- [32] Y. Lu, A. Bengtsson, J. J. Burnett, E. Wiegand, B. Suri, P. Krantz, A. F. Roudsari, A. F. Kockum, S. Gasparinetti, G. Johansson, and P. Delsing, Characterizing decoherence rates of a superconducting qubit by direct microwave scattering, *npj Quantum Inf.* **7**, 35 (2021).
- [33] A. Frisk Kockum, P. Delsing, and G. Johansson, Designing frequency-dependent relaxation rates and lamb shifts for a giant artificial atom, *Phys. Rev. A* **90**, 013837 (2014).
- [34] L. Guo, A. Grimsmo, A. F. Kockum, M. Pletyukhov, and G. Johansson, Giant acoustic atom: A single quantum system with a deterministic time delay, *Phys. Rev. A* **95**, 053821 (2017).
- [35] A. F. Kockum, G. Johansson, and F. Nori, Decoherence-Free Interaction between Giant Atoms in Waveguide Quantum Electrodynamics, *Phys. Rev. Lett.* **120**, 140404 (2018).
- [36] A. Ask, M. Ekström, P. Delsing, and G. Johansson, Cavity-free vacuum-Rabi splitting in circuit quantum acoustodynamics, *Phys. Rev. A* **99**, 013840 (2019).
- [37] B. Kannan, M. J. Ruckriegel, D. L. Campbell, A. Frisk Kockum, J. Braumüller, D. K. Kim, M. Kjaergaard, P. Krantz, A. Melville, B. M. Niedzielski, A. Vepsäläinen, R. Winik, J. L. Yoder, F. Nori, T. P. Orlando, S. Gustavsson, and W. D. Oliver, Waveguide quantum electrodynamics with superconducting artificial giant atoms, *Nature (London)* **583**, 775 (2020).
- [38] A. M. Vadiraj, A. Ask, T. G. McConkey, I. Nsanzineza, C. W. Sandbo Chang, A. F. Kockum, and C. M. Wilson, Engineering the level structure of a giant artificial atom in waveguide quantum electrodynamics, *Phys. Rev. A* **103**, 023710 (2021).
- [39] G. Andersson, B. Suri, L. Guo, T. Aref, and P. Delsing, Non-exponential decay of a giant artificial atom, *Nat. Phys.* **15**, 1123 (2019).
- [40] T. Tufarelli, M. S. Kim, and F. Ciccarello, Non-markovianity of a quantum emitter in front of a mirror, *Phys. Rev. A* **90**, 012113 (2014).
- [41] Y.-L. L. Fang, F. Ciccarello, and H. U. Baranger, Non-markovian dynamics of a qubit due to single-photon scattering in a waveguide, *New J. Phys.* **20**, 043035 (2018).
- [42] P. Haikka and S. Maniscalco, Non-Markovian dynamics of a damped driven two-state system, *Phys. Rev. A* **81**, 052103 (2010).
- [43] P. Haikka, J. D. Cresser, and S. Maniscalco, Comparing different non-Markovianity measures in a driven qubit system, *Phys. Rev. A* **83**, 012112 (2011).
- [44] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information: 10th Anniversary Edition* (Cambridge University Press, Cambridge, England, 2010).
- [45] U. Schollwöck, The density-matrix renormalization group in the age of matrix product states, *Ann. Phys. (Amsterdam)* **326**, 96 (2011).
- [46] G. Vidal, Efficient Classical Simulation of Slightly Entangled Quantum Computations, *Phys. Rev. Lett.* **91**, 147902 (2003).
- [47] G. Vidal, Efficient Simulation of One-Dimensional Quantum Many-Body Systems, *Phys. Rev. Lett.* **93**, 040502 (2004).
- [48] F. Verstraete and J. I. Cirac, Matrix product states represent ground states faithfully, *Phys. Rev. B* **73**, 094423 (2006).
- [49] M. B. Hastings, An area law for one-dimensional quantum systems, *J. Stat. Mech.* (2007) P08024.
- [50] S. R. White, Density Matrix Formulation for Quantum Renormalization groups, *Phys. Rev. Lett.* **69**, 2863 (1992).
- [51] S. Östlund and S. Rommer, Thermodynamic Limit of Density Matrix Renormalization, *Phys. Rev. Lett.* **75**, 3537 (1995).
- [52] Y.-Y. Shi, L.-M. Duan, and G. Vidal, Classical simulation of quantum many-body systems with a tree tensor network, *Phys. Rev. A* **74**, 022320 (2006).
- [53] U. Shrikant, R. Srikanth, and S. Banerjee, Non-Markovian dephasing and depolarizing channels, *Phys. Rev. A* **98**, 032328 (2018).
- [54] X. Cai, Quantum dephasing induced by non-Markovian random telegraph noise, *Sci. Rep.* **10**, 88 (2020).
- [55] S. Wißmann, A. Karlsson, E.-M. Laine, J. Piilo, and H.-P. Breuer, Optimal state pairs for non-Markovian quantum dynamics, *Phys. Rev. A* **86**, 062108 (2012).
- [56] F. Damanet, A. J. Daley, and J. Keeling, Atom-only descriptions of the driven-dissipative dicke model, *Phys. Rev. A* **99**, 033845 (2019).
- [57] R. Hartmann and W. T. Strunz, Accuracy assessment of perturbative master equations: Embracing nonpositivity, *Phys. Rev. A* **101**, 012103 (2020).