



## ON THE INHOMOGENEOUS VINOGRADOV SYSTEM

Downloaded from: <https://research.chalmers.se>, 2025-02-13 05:36 UTC

Citation for the original published paper (version of record):

Brandes, J., Hughes, K. (2022). ON THE INHOMOGENEOUS VINOGRADOV SYSTEM. *Bulletin of the Australian Mathematical Society*, 106(3): 396-403.

<http://dx.doi.org/10.1017/S0004972722000284>

N.B. When citing this work, cite the original published paper.

## ON THE INHOMOGENEOUS VINOGRADOV SYSTEM

JULIA BRANDES  and KEVIN HUGHES 

(Received 15 January 2022; accepted 15 February 2022)

### Abstract

We show that the system of equations

$$\sum_{i=1}^s (x_i^j - y_i^j) = a_j \quad (1 \leq j \leq k)$$

has appreciably fewer solutions in the subcritical range  $s < \frac{1}{2}k(k+1)$  than its homogeneous counterpart, provided that  $a_\ell \neq 0$  for some  $\ell \leq k-1$ . Our methods use Vinogradov's mean value theorem in combination with a shifting argument.

2020 *Mathematics subject classification*: primary 11D45; secondary 11D85, 11L15, 11P05, 42B05.

*Keywords and phrases*: Diophantine equations, exponential sums.

### 1. Introduction

Sitting squarely at the interface of harmonic analysis and analytic number theory, systems of diagonal equations have long attracted a significant amount of attention. In particular, thanks to the resolution of the main conjecture associated with Vinogradov's mean value theorem by Wooley [10] and Bourgain *et al.* [2] (see also [11]), we now have essentially sharp upper bounds for the number of integer solutions of systems of equations of Vinogradov type.

For integral  $s, k$ , denote by  $J_{s,k}(X)$  the number of solutions  $\mathbf{x}, \mathbf{y} \in \mathbb{Z}^s \cap [-X, X]^s$  satisfying

$$\sum_{i=1}^s (x_i^j - y_i^j) = 0 \quad (1 \leq j \leq k). \quad (1.1)$$

We remark for future reference that by transitioning to exponential sums, the quantity  $J_{s,k}(X)$  may be defined also for noninteger positive  $s$  (see (2.4)). In this notation,

---

During the production of this manuscript, the first author was supported by the Swedish Science Foundation (Vetenskapsrådet) via Starting Grant no. 2017-05110.

© The Author(s), 2022. Published by Cambridge University Press on behalf of Australian Mathematical Publishing Association Inc. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

[2, Theorem 1.1] (see also [11, Corollary 1.3]) shows that for all positive numbers  $s$  and all  $k \in \mathbb{N}$ ,

$$J_{s,k}(X) \ll X^\varepsilon (X^s + X^{2s-k(k+1)/2}). \tag{1.2}$$

The second term in (1.2), dominant when  $s > \frac{1}{2}k(k + 1)$ , conforms with the circle method heuristic which states that a system of equations in  $n$  variables and of total degree  $K$  should have  $\sim cX^{n-K}$  solutions over the integers of size at most  $X$ , where the factor  $c$  encodes the density of solutions of the underlying system over all completions of  $\mathbb{Q}$ . Indeed, in the case of (1.1), such an asymptotic formula is readily established for  $s > \frac{1}{2}k(k + 1)$  by a classical application of the Hardy–Littlewood method (see [7, Ch. 7]). Moreover, these methods are robust with respect to small modifications to the system (1.1) such as nontrivial coefficients or a nonvanishing expression on the right-hand side of (1.1).

When  $s$  is smaller than  $\frac{1}{2}k(k + 1)$ , the circle method heuristic fails for (1.1), as we have to allow for the existence of diagonal solutions in which the variables  $(x_1, x_2, \dots, x_s)$  are a permutation of  $(y_1, y_2, \dots, y_s)$ . The contribution of these solutions amounts to  $s! X^s$ , so it is natural to expect an asymptotic formula of the shape

$$J_{s,k}(X) \sim s! X^s \tag{1.3}$$

to hold in the subcritical range. Adventurous souls might even go as far as to conjecture that the nondiagonal contribution to the mean value  $J_{s,k}(X)$  should be no larger than  $O(X^\varepsilon(1 + X^{2s-k(k+1)/2}))$ , based on an expectation that the circle method heuristic should continue to be valid in that setting. Somewhat frustratingly, however, we are able to establish (1.3) only in the range  $s \leq k + 1$  (see [8, Theorem 1]), which falls far short of the conjectured range  $s < \frac{1}{2}k(k + 1)$ .

The purpose of this note is to study an inhomogeneous version of the system (1.1) that does not allow for diagonal contributions. For  $s, k \in \mathbb{N}$  and  $a_1, \dots, a_k \in \mathbb{Z}$ , write  $J_{s,k}(X; \mathbf{a})$  for the number of  $\mathbf{x}, \mathbf{y} \in \mathbb{Z}^s \cap [-X, X]^s$  satisfying the system of equations

$$\sum_{i=1}^s (x_i^j - y_i^j) = a_j \quad (1 \leq j \leq k). \tag{1.4}$$

Systems of this or similar types have recently arisen in a range of different contexts (for example, [1, 4, 5]). It follows easily by a standard application from the triangle inequality that

$$J_{s,k}(X; \mathbf{a}) \leq J_{s,k}(X) \ll X^{s+\varepsilon} \tag{1.5}$$

in the subcritical range. We show that for most  $\mathbf{a}$ , we can do better.

**THEOREM 1.1.** *Suppose that  $k \geq 2$  and  $\mathbf{a} \in \mathbb{Z}^k \setminus \{\mathbf{0}\}$ . Let  $\ell \in \{1, \dots, k\}$  be the smallest integer for which  $a_\ell \neq 0$ . Then for any integer  $s < \frac{1}{2}k(k + 1)$  and any  $\varepsilon > 0$ ,*

$$J_{s,k}(X; \mathbf{a}) \ll X^{s-1/2+\varepsilon} + X^{s-\eta_{s,k}(\ell)+\varepsilon}, \tag{1.6}$$

where

$$\eta_{s,k}(\ell) = \frac{(k-\ell)(k-\ell+1)}{2} \left(1 - \frac{2s}{k(k+1)}\right).$$

By a brief computation, the first term in (1.6) dominates for  $s = \frac{1}{2}k(k+1) - 1$  as soon as  $a_\ell \neq 0$  for some  $\ell \leq k - \frac{1}{2}(\sqrt{2k^2 + 2k + 1} - 1)$ . Hence, we have the following simple consequence of Theorem 1.1.

**COROLLARY 1.2.** *Suppose that  $k \geq 2$  and assume that  $a_\ell \neq 0$  for some*

$$\ell \leq k - \frac{1}{2}(\sqrt{2k^2 + 2k + 1} - 1).$$

*Then for any integer  $s \leq \frac{1}{2}k(k+1) - 1$  and any  $\varepsilon > 0$ ,*

$$J_{s,k}(X; \mathbf{a}) \ll X^{s-1/2+\varepsilon}.$$

Here, one can compute

$$\begin{aligned} k - \frac{1}{2}(\sqrt{2k^2 + 2k + 1} - 1) &= \left(1 - \frac{1}{\sqrt{2}}\right)\left(k + \frac{1}{2}\right) + O(1/k) \\ &= 0.292\dots \cdot k + 0.146\dots + O(1/k). \end{aligned}$$

It may be worth pointing out that we do not expect to obtain power savings over the bound (1.5) when  $s$  is at or beyond the critical point and  $|a_j| \leq 2sX^j$  for  $1 \leq j \leq k$ , since at that point, the contribution from a major arcs analysis will be of size  $\gg X^{2s-k(k+1)}$  (see [7, Ch. 7] for details). In this sense, the range  $s \leq \frac{1}{2}k(k+1) - 1$  is optimal.

The proof of Theorem 1.1 relies crucially on the absence of translation invariance in the system (1.4). For this reason, our result becomes gradually weaker as the system acquires larger translation-invariant subsystems, up to the point when  $\mathbf{a} = (0, \dots, 0, a_k)^t$ , where we have  $\ell = k$  and  $\eta_{s,k}(k) = 0$ . At this point, the entire system becomes translation-invariant, so we fail to make any progress beyond the trivial bound (1.5).

Our strategy transfers to several other settings, but in the interest of a slick presentation, we have opted to focus our attention on the inhomogeneous Vinogradov system, since this seems to be by far the most relevant example. However, we discuss some possible extensions of our result as well as some of its limitations in the final section.

*Notation.* We denote the unit torus by  $\mathbb{T} = \mathbb{R}/\mathbb{Z}$ . Throughout, the letter  $\varepsilon$  will be used to denote an arbitrary positive number. We adopt the convention that whenever  $\varepsilon$  appears in a statement, the statement holds for all  $\varepsilon > 0$ . Finally, we take  $X$  to be a large positive number which, just like the implicit constants in the notations of Landau and Vinogradov, is permitted to depend at most on  $s$ ,  $k$  and  $\varepsilon$ .

**2. Proof of the main theorem**

In our proof, we employ a strategy inspired by work of Wooley [9] to understand systems with incomplete translation-invariant structure. Our first step is to bound  $J_{s,k}(X; \mathbf{a})$  in terms of a different quantity which will be easier to handle. Define polynomials  $p_j$  by setting

$$p_j(h) = \sum_{m=1}^j \binom{j}{m} a_m h^{j-m} \quad (1 \leq j \leq k). \tag{2.1}$$

Write  $H_{s,k}(X; \mathbf{a})$  for the number of  $\mathbf{z}, \mathbf{w} \in \mathbb{Z}^s \cap [-2X, 2X]^s$  and  $h \in \mathbb{Z} \cap [-X, X]$  satisfying the system of equations

$$\sum_{i=1}^s (z_i^j - w_i^j) = p_j(h) \quad (1 \leq j \leq k). \tag{2.2}$$

**LEMMA 2.1.** *We have*

$$J_{s,k}(X; \mathbf{a}) \ll X^{-1} H_{s,k}(X; \mathbf{a}).$$

**PROOF.** Suppose that  $(\mathbf{x}, \mathbf{y})$  is a solution to (1.4), and fix a parameter  $h \in \mathbb{Z}$ . It follows from the binomial theorem that  $(\mathbf{x}, \mathbf{y})$  is also a solution of the shifted system

$$\sum_{i=1}^s ((x_i + h)^j - (y_i + h)^j) = p_j(h) \quad (1 \leq j \leq k).$$

If now  $|h| \leq X$ , then the number of such solutions  $\mathbf{x}, \mathbf{y} \in \mathbb{Z}^s \cap [-X, X]^s$  is certainly no larger than the number of  $\mathbf{z}, \mathbf{w} \in \mathbb{Z}^s \cap [-2X, 2X]^s$  satisfying (2.2) with that particular value of  $h$ . The desired conclusion now follows upon summing over all  $|h| \leq X$ .  $\square$

Next, we write  $H_{s,k}(X; \mathbf{a})$  in terms of mean values over exponential sums. Set

$$f_k(\boldsymbol{\alpha}; X) = \sum_{|x| \leq X} e(\alpha_1 x + \dots + \alpha_k x^k)$$

and

$$g_k(\boldsymbol{\alpha}; X) = \sum_{|x| \leq X} e(\alpha_1 p_1(h) + \dots + \alpha_k p_k(h)), \tag{2.3}$$

and recall that in this notation, we have

$$J_{s,k}(X) = \int_{\mathbb{T}^k} |f_k(\boldsymbol{\alpha}; X)|^{2s} d\boldsymbol{\alpha}. \tag{2.4}$$

Similarly, it follows from standard orthogonality relations that we may write

$$H_{s,k}(X; \mathbf{a}) = \int_{\mathbb{T}^k} |f_k(\boldsymbol{\alpha}; 2X)|^{2s} g_k(-\boldsymbol{\alpha}; X) d\boldsymbol{\alpha}. \tag{2.5}$$

From the definition of  $p_j$  in (2.1), we see that  $g_k(\boldsymbol{\alpha}; X)$  is an exponential sum of degree at most  $k - 1$ ; however, its degree may be lower depending on the values of the coefficients  $a_1, a_2, \dots, a_k$ . In particular, recalling that  $\ell \in \{1, \dots, k\}$  denotes the

smallest integer for which  $a_\ell \neq 0$ , we discern from (2.1) that

$$\deg p_j = \max\{0, j - \ell\} \quad (1 \leq j \leq k). \quad (2.6)$$

Thus,  $g_k(\alpha; X)$  is an exponential sum containing  $k - \ell$  integer polynomials with degrees  $1, \dots, k - \ell$ .

We now apply Hölder's inequality to (2.5) and find that

$$H_{s,k}(X; \mathbf{a}) \leq \left( \int_{\mathbb{T}^k} |f_k(\alpha; 2X)|^{k(k+1)} d\alpha \right)^{2s/k(k+1)} \left( \int_{\mathbb{T}^k} |g_k(\alpha; X)|^{2\sigma} d\alpha \right)^{1/(2\sigma)}, \quad (2.7)$$

where

$$\sigma = \frac{k(k+1)}{2(k(k+1) - 2s)}. \quad (2.8)$$

Clearly, the first integral in (2.7) is  $J_{k(k+1)/2,k}(2X)$ . Moreover, from (2.3), (2.1) and (2.6), it follows via an integer change of variables on  $\alpha$  together with the periodicity of the exponential sums that

$$\int_{\mathbb{T}^k} |g_k(\alpha; X)|^{2\sigma} d\alpha = \int_{\mathbb{T}^{k-\ell}} |f_{k-\ell}(\gamma; X)|^{2\sigma} d\gamma = J_{\sigma, k-\ell}(X).$$

Consequently, upon inserting the bound (1.2) from [2, Theorem 1.1], we conclude that

$$\begin{aligned} H_{s,k}(X; \mathbf{a}) &\ll (J_{k(k+1)/2,k}(2X))^{2s/k(k+1)} (J_{\sigma, k-\ell}(X))^{1/(2\sigma)} \\ &\ll X^\varepsilon (X^{k(k+1)/2})^{2s/k(k+1)} (X^\sigma + X^{2\sigma - (k-\ell)(k-\ell+1)/2})^{1/(2\sigma)} \\ &\ll X^{s+\varepsilon} (X^{1/2} + X^{1-(k-\ell)(k-\ell+1)/(4\sigma)}). \end{aligned}$$

The proof of Theorem 1.1 is now complete upon recalling Lemma 2.1 and inserting the value of  $\sigma$  from (2.8).

### 3. Further discussion: Generalisations and limitations

It is natural to ask in what ways the proof of Theorem 1.1 may be generalised. As mentioned in the introduction, a close reading of our methods reveals that the translation-invariant structure of the system does indeed play a crucial role in our arguments. Consequently, there is little hope to extend our results to incomplete Vinogradov systems such as those considered in [3] without fundamentally different ideas.

However, the idea underpinning the proof of Theorem 1.1 is quite general, and can be used to bound the number of solutions of the inhomogeneous analogues of other translation-invariant systems, including multidimensional ones such as those considered in [6]. Indeed, beyond a sharp mean value estimate for such systems in the subcritical range, we only need a nontrivial bound for a derived mean value over a family of 'shifting polynomials' analogous to the polynomials  $p_j$  occurring in Lemma 2.1. Owing to its genesis via shifting the variables, this derived mean value can be seen to be of the same general shape as the original one (but with lower degree), and therefore a similar mean value estimate can usually be applied, leading to

a nontrivial bound as soon as at least one of the shifting polynomials is nonconstant. Alternatively, one could also use square-root cancellation at the second moment (Plancherel’s theorem) and interpolate with trivial bounds at  $L^\infty$  to obtain cancellation at all moments greater than 2. In this manner, our method shows that the numbers of solutions of most inhomogeneous translation-invariant systems of equations are measurably smaller than the corresponding bounds for their homogeneous analogues for all even moments in the subcritical range.

A different direction for generalisations is that in which the variables  $x_i, y_i$  are restricted to some subset  $\mathcal{X} \subseteq [-X, X] \cap \mathbb{Z}$ . Fixing such a set  $\mathcal{X}$  as well as a set of shifting variables  $\mathcal{H} \subset [-X, X] \cap \mathbb{Z}$ , put  $J_{s,k}(\mathcal{X}; \mathbf{a})$  for the number of  $\mathbf{x}, \mathbf{y} \in \mathcal{X}^s$  satisfying (1.4), and write  $H_{s,k}(\mathcal{X}, \mathcal{H}; \mathbf{a})$  for the number of  $\mathbf{x}, \mathbf{y} \in (\mathcal{X} + \mathcal{H})^s$  and  $h \in \mathcal{H}$  solving the system (2.2). Then an inspection of the proof of Lemma 2.1 shows that

$$J_{s,k}(\mathcal{X}; \mathbf{a}) \ll |\mathcal{H}|^{-1} H_{s,k}(\mathcal{X}, \mathcal{H}; \mathbf{a}).$$

Since it follows from [2, Theorem 1.2] and [11, Theorem 1.1] that

$$J_{s,k}(\mathcal{X}; \mathbf{0}) \ll X^\varepsilon (|\mathcal{X}|^s + |\mathcal{X}|^{2s-k(k+1)/2}),$$

the same argument as above leads *mutatis mutandis* to the following conclusion.

**THEOREM 3.1.** *Suppose that  $k \geq 2$  and  $\mathbf{a} \in \mathbb{Z}^k \setminus \{\mathbf{0}\}$ . Let  $\ell \in \{1, \dots, k\}$  be the smallest integer for which  $a_\ell \neq 0$ . Moreover, fix a set  $\mathcal{X} \subseteq [-X, X] \cap \mathbb{Z}$ . Then for any integer  $s < \frac{1}{2}k(k+1)$ , any set  $\mathcal{H} \subseteq [-X, X] \cap \mathbb{Z}$  and any  $\varepsilon > 0$ ,*

$$J_{s,k}(\mathcal{X}; \mathbf{a}) \ll X^\varepsilon |\mathcal{X} + \mathcal{H}|^s (|\mathcal{H}|^{-1/2} + |\mathcal{H}|^{-\eta_{s,k}(\ell)}),$$

where  $\eta_{s,k}(\ell)$  is as in Theorem 1.1.

From the point of view of harmonic analysis, it would be of interest to have an analogue of Theorem 1.1 involving systems of the type (1.4) in which each solution is counted by a complex weight. Such situations arise habitually in the context of restriction and extension operators. For a given complex-valued sequence  $(c_n)_{n \in \mathbb{Z}}$ , one defines the truncated extension operator along the moment curve  $(t, t^2, \dots, t^k) \subset \mathbb{R}^k$  via

$$E_X c(\boldsymbol{\alpha}) = \sum_{|x| \leq X} c_x e(\alpha_1 x + \dots + \alpha_k x^k),$$

so that  $f_k(\boldsymbol{\alpha}; X) = E_X \mathbb{1}(\boldsymbol{\alpha})$ . In this notation, the conclusion of [2, Theorem 1.2] and [11, Theorem 1.1] reads

$$\|E_X c\|_{L^{2s}(\mathbb{T}^k)} \ll X^\varepsilon (1 + X^{1/2-k(k+1)/(4s)}) \|c\|_{\ell^2([-X, X])}.$$

Consider the Fourier transform of  $|E_X c(\boldsymbol{\alpha})|^{2s}$ , which is given by

$$\Phi(\mathbf{n}) = \int_{\mathbb{T}^k} |E_X c(\boldsymbol{\alpha})|^{2s} e(-\boldsymbol{\alpha} \cdot \mathbf{n}) d\boldsymbol{\alpha},$$

so that in particular  $\|E_X c\|_{L^{2s}(\mathbb{T}^k)}^{2s} = \Phi(\mathbf{0})$ . Of course, it follows directly via the triangle inequality that  $|\Phi(\mathbf{n})| \leq \Phi(\mathbf{0})$  for all  $\mathbf{n} \in \mathbb{Z}^k$ , but it would be desirable to have a stronger

inequality and ideally one that exhibits a power saving over the trivial bound. In the special case when  $\mathfrak{c}$  is the indicator function on a set  $X \subseteq \mathbb{Z} \cap [-X, X]$ , the results of Theorems 1.1 and 3.1 achieve such a power saving for a large selection of  $\mathfrak{n}$  and for all  $s \in \mathbb{N}$  below the critical point. In fact, it seems plausible that a careful modification of our arguments should apply also to a setting where the sequence  $(c_x)_{x \in \mathbb{Z}}$  takes nonnegative real values. However, it is less obvious how the proof of Lemma 2.1 would carry over to the general situation in which the weights  $\mathfrak{c}$  may take complex values.

Finally, we remark that the saving of  $X^{1/2}$  in Theorem 1.1 is optimal within the methods. This can be seen by noting that the shifting and averaging operation creates one dummy variable, on which we can expect no more than square-root cancellation. The analytically inclined reader may also be interested to note that the results of [2, 11] require the underlying curve to have torsion and this is what enables the strong bounds on mean values of the exponential sum  $f_k(\alpha; X)$ . The curve underlying the exponential sum  $g_k(\alpha; X)$ , given by  $(p_1(t), \dots, p_k(t))$ , does not have torsion over  $\mathbb{R}^k$ , but it can be restricted to a  $(k - \ell)$ -dimensional subspace in which it has torsion. This gives a geometric explanation for why our results are weaker as  $\ell$  increases, and fail to give any improvement at all when  $\ell$  is equal to  $k$ .

### Acknowledgement

The authors are grateful to Igor Shparlinski for useful discussions surrounding the topic of this paper.

### References

- [1] R. C. Baker, M. Munsch and I. E. Shparlinski, ‘Additive energy and a large sieve inequality for sparse sequences’, *Mathematika*, to appear.
- [2] J. Bourgain, C. Demeter and L. Guth, ‘Proof of the main conjecture in Vinogradov’s mean value theorem for degrees higher than three’, *Ann. of Math. (2)* **184**(2) (2016), 633–682.
- [3] J. Brandes and T. D. Wooley, ‘Vinogradov systems with a slice off’, *Mathematika* **63**(3) (2017), 797–817.
- [4] C. Chen, B. Kerr, J. Maynard and I. E. Shparlinski, ‘Metric theory of Weyl sums’, *Math. Ann.*, to appear. <https://doi.org/10.1007/s00208-021-02352-x>.
- [5] S. Dendrinos, K. Hughes and M. Vitturi, ‘Some subcritical estimates for the  $\ell^p$ -improving problem for discrete curves’, *J. Fourier Anal. Appl.*, to appear.
- [6] S. Guo and P. Zorin-Kranich, ‘Decoupling for moment manifolds associated to Arkhipov–Chubarikov–Karatsuba systems’, *Adv. Math.* **360** (2020), Article no. 106889.
- [7] R. C. Vaughan, *The Hardy–Littlewood Method*, 2nd edn (Cambridge University Press, Cambridge, 1997).
- [8] R. C. Vaughan and T. D. Wooley, ‘A special case of Vinogradov’s mean value theorem’, *Acta Arith.* **79**(3) (1997), 193–204.
- [9] T. D. Wooley, ‘Rational solutions of pairs of diagonal equations, one cubic and one quadratic’, *Proc. Lond. Math. Soc. (3)* **110**(2) (2015), 325–356.
- [10] T. D. Wooley, ‘The cubic case of the main conjecture in Vinogradov’s mean value theorem’, *Adv. Math.* **294** (2016), 532–561.
- [11] T. D. Wooley, ‘Nested efficient congruencing and relatives of Vinogradov’s mean value theorem’, *Proc. Lond. Math. Soc. (3)* **118**(4) (2019), 942–1016.



JULIA BRANDES, Mathematical Sciences,  
University of Gothenburg and Chalmers Institute of Technology,  
412 96 Göteborg, Sweden  
e-mail: [brjulia@chalmers.se](mailto:brjulia@chalmers.se)

KEVIN HUGHES, School of Mathematics,  
University of Bristol, Fry Building, Woodland Road, Bristol BS8 1UG, UK  
and  
Heilbronn Institute for Mathematical Research  
e-mail: [khughes.math@gmail.com](mailto:khughes.math@gmail.com)