



CHALMERS
UNIVERSITY OF TECHNOLOGY

A Combinatorial Description of Certain Polynomials Related to the XYZ Spin Chain. II. The Polynomials $p(n)$

Downloaded from: <https://research.chalmers.se>, 2024-08-07 23:05 UTC

Citation for the original published paper (version of record):

Hietala, L. (2022). A Combinatorial Description of Certain Polynomials Related to the XYZ Spin Chain. II. The Polynomials $p(n)$. *Symmetry, Integrability and Geometry - Methods and Applications*, 18. <http://dx.doi.org/10.3842/SIGMA.2022.036>

N.B. When citing this work, cite the original published paper.

A Combinatorial Description of Certain Polynomials Related to the XYZ Spin Chain.

II. The Polynomials p_n

Linnea HIETALA ^{ab}

a) *Department of Mathematics, Uppsala University, Box 480, 751 06 Uppsala, Sweden*
E-mail: linnea.hietala@math.uu.se

b) *Department of Mathematical Sciences, Chalmers University of Technology and University of Gothenburg, 412 96 Gothenburg, Sweden*

Received August 06, 2021, in final form April 29, 2022; Published online May 15, 2022

<https://doi.org/10.3842/SIGMA.2022.036>

Abstract. By specializing the parameters in the partition function of the 8VSOS model with domain wall boundary conditions and diagonal reflecting end, we find connections between the three-color model and certain polynomials $p_n(z)$, which are conjectured to be equal to certain polynomials of Bazhanov and Mangazeev, appearing in the eigenvectors of the Hamiltonian of the supersymmetric XYZ spin chain. This article is a continuation of a previous paper where we investigated the related polynomials $q_n(z)$, also conjectured to be equal to polynomials of Bazhanov and Mangazeev, appearing in the eigenvectors of the supersymmetric XYZ spin chain.

Key words: eight-vertex SOS model; domain wall boundary conditions; reflecting end; three-color model; XYZ spin chain; polynomials; positive coefficients

2020 Mathematics Subject Classification: 82B23; 05A15; 33E17

1 Introduction

In 1967, Lieb [15] solved the first special cases of the six-vertex (6V) model on a square lattice, with periodic boundary conditions in both directions. Later the same year, Sutherland [25] solved the general case. A bijection from the states of the 6V model to three-colorings of a square lattice, where the color is fixed on one face and all adjacent faces have different colors, was found by Lenard [15, (note added in proof)]. The three-color model, where each color is given a weight, was introduced by Baxter [1].

Izergin [10, 11] obtained a determinant formula for the partition function of the 6V model with domain wall boundary conditions (DWBC) [12]. By specializing the parameters in this determinant formula, Kuperberg [13] gave an alternative proof of the alternating sign matrix (ASM) conjecture of Mills, Robbins and Rumsey [17], which gives a formula for the number of ASMs. This conjecture was originally proven a few months earlier by Zeilberger [28]. The eight-vertex solid-on-solid (8VSOS) model [3] is a two parameter generalization of the 6V model. Rosengren [22] specialized the parameters of the 8VSOS model with DWBC in Kuperberg's manner and obtained the three-color model.

Kuperberg [14] later used Tsuchiya's [26] determinant formula for the partition function of the 6V model with DWBC and one reflecting end to give a formula for the number of UASMs, which are ASMs with U-turns on one side. A determinant formula for the partition function of the 8VSOS model with DWBC and one reflecting end was given by Filali [7] in 2011.

Razumov and Stroganov [18] found connections between the supersymmetric XXZ spin chain (i.e., where $\Delta = -1/2$ in the Hamiltonian, see, e.g., [8]) and ASMs. Similar problems for the

supersymmetric XYZ spin chain were studied by Bazhanov and Mangazeev [4, 16] (see also [20]). The ground state eigenvalues of Baxter's Q -operator [2] for the eight-vertex (8V) model as well as the components of the ground state eigenvectors of the supersymmetric XYZ-Hamiltonian can be expressed in terms of certain polynomials which seem to have positive integer coefficients [5, 16]. Further investigation has been done by Brasseur and Hagendorf [6].

In Rosengren's study of the 8VSOS model, certain polynomials with positive coefficients showed up and in [24], Rosengren generalized these polynomials. Zinn-Justin [29] introduced equivalent polynomials and conjectured that Bazhanov's and Mangazeev's polynomials are specializations of these. This led to the suspicion that the combinatorial interpretation of the polynomials of Bazhanov and Mangazeev could have to do with the three-color model.

In this article, as well as in a previous one [9], we assume Zinn-Justin's conjecture, i.e., we use Rosengren's polynomials as the definition of Bazhanov's and Mangazeev's polynomials. In [9], we studied the connection between the three-color model and the polynomials $q_{n-1}(z)$ of Bazhanov and Mangazeev, appearing in the eigenvectors of the Hamiltonian of the supersymmetric XYZ spin chain. By specializing the parameters in the partition function of the 8VSOS model with DWBC and reflecting end in Kuperberg's way, we found an explicit combinatorial expression for $q_{n-1}(z)$, $n \geq 0$, in terms of the partition function of the three-color model on a lattice of size $2n \times n$ with the same boundary conditions.

In [9], we specialized all the spectral parameters in the same way. In the present article, we specialize one of the parameters slightly differently to find a connection between the three-color model on a lattice of size $2n \times n$, and another set of polynomials, which Bazhanov and Mangazeev call $p_{n-1}(z)$, also appearing in the supersymmetric XYZ eigenvectors. Again we consider $n \geq 0$. The expressions that we encounter in this paper get a bit more complicated than in the $q_{n-1}(z)$ case, as the expression includes a sum connected to certain rows of the lattice, which does not appear for $q_{n-1}(z)$.

The paper is organized as follows. In Section 2, we introduce the 8VSOS model as well as the three-color model. In Section 3 we specialize the parameters in Filali's determinant formula and write the partition function in terms of certain polynomials, equivalent to $p_n(z)$. In Section 4 we do the same specialization in the very definition of the partition function and get another expression. In Sections 5 and 6, we combine the two expressions. Using three-colorings, we finally find combinatorial formulas for $p_n(z)$ in terms of three-colorings.

2 Preliminaries

Let τ and η be fixed parameters with $\text{Im}(\tau) > 0$ and $\eta \notin \mathbb{Z} + \tau\mathbb{Z}$. Define $p = e^{2\pi i\tau}$ and $q = e^{2\pi i\eta}$. We then define the theta function

$$\vartheta(x, p) = \prod_{j=0}^{\infty} (1 - p^j x)(1 - p^{j+1}/x).$$

We will often suppress the p and write $\vartheta(x) := \vartheta(x, p)$. We write out the second parameter only when it is not just p . We will also use the short hand notation $\vartheta(bx^{\pm a}) := \vartheta(bx^a)\vartheta(bx^{-a})$. Then we define

$$[x] = q^{-x/2}\vartheta(q^x, p).$$

The most important properties of the theta function are

$$\vartheta(px) = \vartheta(1/x) = -\frac{1}{x}\vartheta(x),$$

and the addition rule

$$\vartheta(x_1 x_3)\vartheta(x_1/x_3)\vartheta(x_2 x_4)\vartheta(x_2/x_4) - \vartheta(x_1 x_4)\vartheta(x_1/x_4)\vartheta(x_2 x_3)\vartheta(x_2/x_3)$$




Figure 1. The 8VSOS model with DWBC and reflecting end in the case $n = 3$. The parameters μ_i and λ_i are the spectral parameters.

$$= \frac{x_2}{x_3} \vartheta(x_1 x_2) \vartheta(x_1/x_2) \vartheta(x_3 x_4) \vartheta(x_3/x_4). \quad (2.1)$$

Also define $\omega = e^{2\pi i/3}$. Later on we will specify to $\eta = -2/3$ (which corresponds to supersymmetry [8]). In this case $q = \omega$.

For any $a \in \mathbb{Z}$ and arbitrary x , we have [22]

$$\begin{aligned} \omega^a &= \omega^{a+3}, \quad 1 + \omega + \omega^2 = 0, \\ \vartheta(p^{1/2}\omega) &= \vartheta(p^{1/2}\omega^2), \end{aligned} \quad (2.2)$$

and

$$\vartheta(x\omega^a) \vartheta(x\omega^{a+1}) \vartheta(x\omega^{a+2}) = \vartheta(x^3, p^3). \quad (2.3)$$

2.1 The 8VSOS model with DWBC and reflecting end

Now we will define the model that we are studying in this paper. Consider a square lattice with $2n \times n$ lines, where the $2n$ horizontal lines are connected pairwise at the left edge as in Figure 1. We equip each line with an orientation. The positive direction is upwards for the vertical lines. For the horizontal double lines, the positive direction goes to the left for the lower part, then it turns and goes to the right on the upper part of the double line. The left side, where the lines turn, is called the reflecting end. To each edge and to each turn we assign a spin ± 1 . A lattice with a spin assigned to each edge and each turn is called a state. Graphically we can describe the orientation of a line with a big arrow at the end of the line, and the spins we describe with arrows on the edges. A positive spin, $+1$, corresponds to an arrow pointing in the positive direction of the line, and a negative spin, -1 , corresponds to an arrow pointing in the opposite direction.

The spins need to follow the so-called ice rule, which demands that at each vertex, two arrows must be pointing inwards and two arrows must be pointing outwards. Because of the ice rule, there can only be six types of different local configurations of the spins at the vertices. These are the vertices in Figure 2. Because of the reflecting end, for every second row in the square lattice, we need to read off the vertices 90 degrees counterclockwise. We assume two possibilities for the turns, depicted in Figure 3. This corresponds to a diagonal reflection matrix [7], and therefore we say that the model has a diagonal reflecting end.

To each vertical line we assign a spectral parameter μ_j , and to each horizontal double line we assign a parameter with the value $-\lambda_i$ on the lower part of the line, and the value λ_i on




Figure 2. The nonzero vertex weights for the 8VSOS model. The spins are indicated with an arrow halfway the edge, where right and up are positive spins, and left and down are negative spins. The vertex weights depend on the spin configurations, the spectral parameters λ and μ , the height z in the upper left face, as well as on the dynamical parameter ρ .



Figure 3. The nonzero boundary weights for the diagonal reflecting end. The weights depend on the spin configurations, the spectral parameter λ , the height z outside the turn, the dynamical parameter ρ , as well as on the boundary parameter ζ .

the upper part of the line, see Figure 1. To each face of the lattice, we assign a height $a \in \mathbb{Z}$. Looking in the direction of each spin arrow, the height to the right of the edge must always be 1 smaller than the height to the left. Fixing the height in one single face of a given state determines the heights of the whole state. We fix the height in the upper left corner to be 0. Finally, we fix a so-called dynamical parameter $\rho \in \mathbb{C}$ and a boundary parameter $\zeta \in \mathbb{C}$.

On the left side of the model we have the reflecting wall. On the remaining three sides, we take DWBC, which means that the spin arrows on the top and the bottom are pointing inwards and the spin arrows on the right boundary are pointing outwards, as in Figure 1. Fixing the height in the upper left corner, all the heights of the faces at the boundaries are determined by the boundary conditions, except for the heights of the faces inside the turns.

2.2 Weights and partition function of the 8VSOS model

To each vertex, we assign a weight which depends on the spins on the adjacent edges, on the height in one of the corners, the dynamical parameter ρ , as well as on the spectral parameters λ_i and μ_j on the lines going through the vertex. Each vertex weight is given by a function w , which is one of the six functions a_{\pm} , b_{\pm} or c_{\pm} , depending on the spin configuration as in Figure 2. To a vertex on the upper part of a double line, with spectral parameters λ_i and μ_j , and height z in the upper left corner, as in Figure 2, we assign the weight $w(\lambda_i - \mu_j, q^{\rho+z})$. On the lower part of the double line, we need to read off the weights 90 degrees counterclockwise and thus consider the

height z in the lower left corner. The weight in this case is $w(\mu_j - (-\lambda_i), q^{\rho+z}) = w(\lambda_i + \mu_j, q^{\rho+z})$. Finally to each turn with spectral parameter λ_i and height z inside the turn, as in Figure 3, we assign a weight $w(\lambda_i, q^{\rho+z}, q^\zeta)$, where w is one of the two functions k_\pm depending on the spins on the turning edge. These local weights are given by

$$\begin{aligned} a_+(\lambda, q^z) &= a_-(\lambda, q^z) = \frac{[\lambda + 1]}{[1]}, \\ b_+(\lambda, q^z) &= \frac{[\lambda][z - 1]}{[z][1]}, & b_-(\lambda, q^z) &= \frac{[\lambda][z + 1]}{[z][1]}, \\ c_+(\lambda, q^z) &= \frac{[z + \lambda]}{[z]}, & c_-(\lambda, q^z) &= \frac{[z - \lambda]}{[z]}, \\ k_+(\lambda, q^z, q^\zeta) &= \frac{[z + \zeta - \lambda]}{[z + \zeta + \lambda]}, & k_-(\lambda, q^z, q^\zeta) &= \frac{[\zeta - \lambda]}{[\zeta + \lambda]}, \end{aligned}$$

and correspond to the different local spin configurations as in Figures 2 and 3. Sometimes when it is the spin configurations around a vertex or at a turn that are of interest, we will refer to an a_\pm , b_\pm , c_\pm vertex or a k_+ (positive) or k_- (negative) turn, without mentioning the different parameters.

The weight of a state is given by the product of all the local weights of the vertices and turns. The partition function is defined as

$$Z_n(q^{\lambda_1}, \dots, q^{\lambda_n}, q^{\mu_1}, \dots, q^{\mu_n}, q^\rho, q^\zeta) = \sum_{\text{states}} \prod_{\text{vertices}} w(\text{vertex}) \prod_{\text{turns}} w(\text{turn}). \quad (2.4)$$

The vertex weights in the partition function are either $w(\lambda_i + \mu_j, q^{\rho+z})$ or $w(\lambda_i - \mu_j, q^{\rho+z})$, where w is one of a_\pm , b_\pm and c_\pm defined above, and the weights of the turns are one of $k_\pm(\lambda_i, q^{\rho+z}, q^\zeta)$. It is easy to see that the weights as well as the partition function (2.4) are well-defined (see [9]). Filali [7] used the Izergin–Korepin method to obtain a determinant formula for the partition function of the 8VSOS model with DWBC and reflecting end, namely,

$$\begin{aligned} &Z_n(q^{\lambda_1}, \dots, q^{\lambda_n}, q^{\mu_1}, \dots, q^{\mu_n}, q^\rho, q^\zeta) \\ &= [1]^{n-2n^2} \prod_{i=1}^n \frac{[2\lambda_i][\zeta - \mu_i][\rho + \zeta + \mu_i][\rho + (2i - n - 2)]}{[\zeta + \lambda_i][\rho + \zeta + \lambda_i][\rho + (n - i)]} \\ &\quad \times \frac{\prod_{i,j=1}^n [\lambda_i + \mu_j + 1][\lambda_i - \mu_j + 1][\lambda_i + \mu_j][\lambda_i - \mu_j]}{\prod_{1 \leq i < j \leq n} [\lambda_i + \lambda_j + 1][\lambda_i - \lambda_j][\mu_j + \mu_i][\mu_j - \mu_i]} \det_{1 \leq i, j \leq n} K, \end{aligned} \quad (2.5)$$

where

$$K_{ij} = \frac{1}{[\lambda_i + \mu_j + 1][\lambda_i - \mu_j + 1][\lambda_i + \mu_j][\lambda_i - \mu_j]}.$$

2.3 The three-color model and its partition function

Another model showing up in this paper is the three-color model. This is a model on a square lattice, with the faces filled with three different colors, which we call color 0, 1, and 2, such that adjacent faces have different colors. A weight t_i is assigned to each face of color i . A state of the three-color model is called a three-coloring.

We study the three-color model on the $2n \times n$ lattice (i.e., a lattice with $(2n+1) \times (n+1)$ faces). If we reduce the heights a of the faces in the 8VSOS model modulo 3, the states of the 8VSOS model can be identified with the states of the three-color model, see Figure 4. The DWBC and the reflecting end in the 8VSOS model correspond to the following rules for the colors in the




Figure 4. A state of the three-color model, for $n = 3$, with colors 0, 1 and 2. The arrows on the edges show the corresponding state in the 8VSOS model. Here $m = 1$.

three-color model. In the upper left corner, we fix color 0. On three of the boundaries, the colors alternate cyclically. Starting from the upper left corner, going to the right, the colors increase in the order $0 < 1 < 2 < 0$, to reach $n \bmod 3$ in the upper right corner. From there, going down, the colors decrease down to $(-n) \bmod 3$ in the lower right corner. Continuing to the left, the colors increase again, up to 0 in the lower left corner. On the left hand side, at the reflecting wall, every second face has color 0. Inside the turns, the colors differ depending on the type of turn in the corresponding state of the 8VSOS model. A negative turn corresponds to color 1, and a positive turn corresponds to color 2. We will henceforth assume these boundary conditions, even if we do not mention them explicitly. The partition function of the three-color model with DWBC and reflecting end, and with color 0 fixed in the upper left corner, is

$$Z_n^{3C}(t_0, t_1, t_2) = \sum_{\text{states}} \prod_{\text{faces}} t_i.$$

Let m be the number of positive turns in a state of the 8VSOS model with DWBC and reflecting end. Specifying m means that we have specified the number of faces with color 2 on the left side. If we specify $m = 0$, the colors on the left side alternate between color 0 and 1. There is a bijection between the three-colorings with $m = 0$ and the vertically symmetric alternating sign matrices of size $(2n + 1) \times (2n + 1)$ [14].

3 Connection to Rosengren's polynomials

The goal in this paper is to express the partition function in terms of certain polynomials p_n , see further Section 6. To do this we specialize the variables in the partition function of the 8VSOS model with DWBC and reflecting end.

First, introduce the variables [24]

$$\psi := \psi(\tau) = \frac{\omega^2 \vartheta(-1) \vartheta(-p^{1/2} \omega)}{\vartheta(-p^{1/2}) \vartheta(-\omega)}, \quad x(z) = \frac{\vartheta(-p^{1/2} \omega)^2 \vartheta(\omega e^{\pm 2\pi i z})}{\vartheta(-\omega)^2 \vartheta(p^{1/2} \omega e^{\pm 2\pi i z})},$$

and define

$$T(x_1, \dots, x_{2n}) = \frac{\prod_{i,j=1}^n G(x_j, x_{n+i})}{\Delta(x_1, \dots, x_n) \Delta(x_{n+1}, \dots, x_{2n})} \det_{1 \leq i, j \leq n} \left(\frac{1}{G(x_j, x_{n+i})} \right), \quad (3.1)$$

where $\Delta(x_1, \dots, x_n) = \prod_{1 \leq i < j \leq n} (x_j - x_i)$ is the Vandermonde polynomial, and

$$G(x, y) = (\psi + 2)xy(x + y) + \psi(2\psi + 1)(x + y) - 2(\psi^2 + 3\psi + 1)xy - \psi(x^2 + y^2).$$

T is a symmetric polynomial [23] as well. For ψ , the following identities hold [22, Lemma 9.1]:

$$2\psi + 1 = \frac{\vartheta(-p^{1/2}\omega)^2 \vartheta(\omega)^2}{\vartheta(-\omega)^2 \vartheta(p^{1/2}\omega)^2}, \quad (3.2)$$

$$\psi + 1 = -\frac{\vartheta(p^{1/2}) \vartheta(-p^{1/2}\omega)}{\vartheta(-p^{1/2}) \vartheta(p^{1/2}\omega)}. \quad (3.3)$$

Another identity we will need, which follows from the addition rules (2.1) and (2.2), is

$$x(z) - x(w) = \frac{\vartheta(-p^{1/2}\omega)^2 \vartheta(p^{1/2}\omega) \vartheta(p^{1/2}) \omega}{\vartheta(-\omega)^2} \frac{e^{-2\pi i w} \vartheta(e^{2\pi i(w \pm z)})}{\vartheta(p^{1/2}\omega e^{\pm 2\pi i z}) \vartheta(p^{1/2}\omega e^{\pm 2\pi i w})}. \quad (3.4)$$

We also need the following lemma, which is Lemma 4.1 of [9], where it is proven.

Lemma 3.1. *We have*

$$\frac{\vartheta(e^{2\pi i(w \pm z)})}{\vartheta(e^{6\pi i(w \pm z)}, p^3)} = \frac{\tilde{C} e^{-4\pi i w}}{\vartheta(p^{1/2}\omega e^{\pm 2\pi i w})^2 \vartheta(p^{1/2}\omega e^{\pm 2\pi i z})^2} \frac{1}{G(x(z), x(w))},$$

with

$$\tilde{C} = \frac{\omega^2 \vartheta(-1) \vartheta(p^{1/2})^3 \vartheta(p^{1/2}\omega)^2 \vartheta(-p^{1/2}\omega)^6}{\vartheta(-\omega)^4 \vartheta(-p^{1/2})}.$$

As in [9], we rewrite Filali's determinant formula (2.5) and specialize to $\eta = -2/3$. Let $z_{n+i} = -2(\lambda_i - 1)/3$ and $z_j = -2\mu_j/3$ for all $0 \leq i, j \leq n$. By using (2.3), (3.4) and Lemma 3.1 we can rewrite Filali's determinant formula in terms of the polynomials T . Details can be found in [9, Section 4]. The partition function becomes

$$\begin{aligned} Z_n(q^{\lambda_1}, \dots, q^{\lambda_n}, q^{\mu_1}, \dots, q^{\mu_n}, \rho, \zeta) \\ = (-1)^{\binom{n}{2}} \omega^{\binom{n+1}{2}} \vartheta(\omega)^{n-2n^2} C^{m^2-n} \tilde{B} \prod_{i=1}^{2n} \vartheta(p^{1/2}\omega e^{\pm 2\pi i z_i})^{n-1} \\ \times \prod_{i=1}^n \frac{\vartheta(\omega e^{4\pi i z_{n+i}}) \vartheta(\zeta e^{-2\pi i z_i}) \vartheta(\rho \zeta e^{2\pi i z_i})}{\vartheta(\zeta \omega e^{2\pi i z_{n+i}}) \vartheta(\rho \zeta \omega e^{2\pi i z_{n+i}})} T(x(z_1), \dots, x(z_{2n})), \end{aligned}$$

where

$$C = \frac{\vartheta(-\omega)^2 \vartheta(-p^{1/2})}{\omega \vartheta(-1) \vartheta(p^{1/2})^2 \vartheta(p^{1/2}\omega) \vartheta(-p^{1/2}\omega)^4}$$

and

$$\tilde{B} = \begin{cases} 1, & \text{for } n \equiv 0, 2 \pmod{3}, \\ \frac{\vartheta(\rho \omega^2)}{\vartheta(\rho)}, & \text{for } n \equiv 1 \pmod{3}. \end{cases}$$

The goal is to express the partition function in terms of the polynomials p_n which correspond to $T(2\psi + 1, \dots, 2\psi + 1, \psi)$, see further Section 6. To do this, specialize $\lambda_i = 1$ for all i , and $\mu_j = 0$ for $1 \leq j \leq n-1$, i.e., $z_i = 0$ for all i except for $i = n$. Recall that $x(0) = 2\psi + 1$ (3.2).

To get $x(z_n) = \psi$, we must have $z_n = -1/6$ and $\mu_n = 1/4$, and hence $q^{\mu_n} = -\omega$. Specializing to $\mu_n = 1/4$ yields

$$\begin{aligned} Z_n(\underbrace{\omega, \dots, \omega}_{n \text{ times}}, \underbrace{1, \dots, 1}_{n-1 \text{ times}}, -\omega, \rho, \zeta) &= (-1)^{\binom{n+1}{2}} \omega^{\binom{n}{2}} \left(\frac{C}{\vartheta(\omega)^2} \right)^{n^2-n} \\ &\times \tilde{B} \vartheta(p^{1/2}\omega)^{2(n-1)(2n-1)} (\vartheta(-p^{1/2}\omega^2) \vartheta(-p^{1/2}))^{n-1} \\ &\times \left(\frac{\vartheta(\zeta) \vartheta(\rho\zeta)}{\vartheta(\zeta\omega) \vartheta(\rho\zeta\omega)} \right)^{n-1} \frac{\vartheta(-\zeta\omega^2) \vartheta(-\rho\zeta\omega)}{\vartheta(\zeta\omega) \vartheta(\rho\zeta\omega)} T(2\psi + 1, \dots, 2\psi + 1, \psi). \end{aligned} \quad (3.5)$$

Now we want to rewrite $\left(\frac{\vartheta(\zeta) \vartheta(\rho\zeta)}{\vartheta(\zeta\omega) \vartheta(\rho\zeta\omega)} \right)^{n-1} \frac{\vartheta(-\zeta\omega^2) \vartheta(-\rho\zeta\omega)}{\vartheta(\zeta\omega) \vartheta(\rho\zeta\omega)}$ in terms of $\frac{\vartheta(\rho\zeta\omega^2)}{\vartheta(\rho\zeta\omega)}$ and $\frac{\vartheta(\zeta\omega^2)}{\vartheta(\zeta\omega)}$. As in [9],

$$\frac{\vartheta(\zeta) \vartheta(\rho\zeta)}{\vartheta(\zeta\omega) \vartheta(\rho\zeta\omega)} = -\frac{\omega \vartheta(\rho\omega)}{\vartheta(\rho)} \frac{\vartheta(\rho\zeta\omega^2)}{\vartheta(\rho\zeta\omega)} - \frac{\omega^2 \vartheta(\rho\omega^2)}{\vartheta(\rho)} \frac{\vartheta(\zeta\omega^2)}{\vartheta(\zeta\omega)}$$

and

$$\frac{\vartheta(-\zeta\omega^2) \vartheta(-\rho\zeta\omega)}{\vartheta(\zeta\omega) \vartheta(\rho\zeta\omega)} = -\frac{\omega \vartheta(-1) \vartheta(-\rho\omega^2)}{\vartheta(\rho) \vartheta(\omega)} \frac{\vartheta(\rho\zeta\omega^2)}{\vartheta(\rho\zeta\omega)} + \frac{\vartheta(-\omega) \vartheta(\rho)}{\vartheta(\rho) \vartheta(\omega)} \frac{\vartheta(\zeta\omega^2)}{\vartheta(\zeta\omega)},$$

because of the addition rule (2.1). Using the binomial theorem we conclude that

$$\begin{aligned} &\left(\frac{\vartheta(\zeta) \vartheta(\rho\zeta)}{\vartheta(\zeta\omega) \vartheta(\rho\zeta\omega)} \right)^{n-1} \frac{\vartheta(-\zeta\omega^2) \vartheta(-\rho\zeta\omega)}{\vartheta(\zeta\omega) \vartheta(\rho\zeta\omega)} \\ &= \frac{(-1)^n}{\vartheta(\rho)^n \vartheta(\omega)} \sum_{m=0}^n \left(\binom{n-1}{m-1} \omega^{2n-m} \vartheta(-1) \vartheta(-\rho\omega^2) \vartheta(\rho\omega)^{m-1} \vartheta(\rho\omega^2)^{n-m} \right. \\ &\quad \left. - \binom{n-1}{m} \omega^{2n-m-2} \vartheta(-\omega) \vartheta(-\rho) \vartheta(\rho\omega)^m \vartheta(\rho\omega^2)^{n-m-1} \right) \\ &\quad \times \left(\frac{\vartheta(\rho\zeta\omega^2)}{\vartheta(\rho\zeta\omega)} \right)^m \left(\frac{\vartheta(\zeta\omega^2)}{\vartheta(\zeta\omega)} \right)^{n-m}. \end{aligned}$$

Inserting this into (3.5) yields

$$\begin{aligned} &Z_n(\omega, \dots, \omega, 1, \dots, 1, -\omega, \rho, \zeta) \\ &= (-\omega)^{\binom{n+1}{2}+n} \left(\frac{C}{\vartheta(\omega)^2} \right)^{n^2-n} T(2\psi + 1, \dots, 2\psi + 1, \psi) \\ &\quad \times \frac{\tilde{B} \vartheta(p^{1/2}\omega)^{2(n-1)(2n-1)} (\vartheta(-p^{1/2}\omega^2) \vartheta(-p^{1/2}))^{n-1}}{\vartheta(\rho)^n \vartheta(\omega)} \\ &\quad \times \sum_{m=0}^n \left(\binom{n-1}{m-1} \omega^{-m} \vartheta(-1) \vartheta(-\rho\omega^2) \vartheta(\rho\omega)^{m-1} \vartheta(\rho\omega^2)^{n-m} - \binom{n-1}{m} \omega^{-m-2} \right. \\ &\quad \left. \times \vartheta(-\omega) \vartheta(-\rho) \vartheta(\rho\omega)^m \vartheta(\rho\omega^2)^{n-m-1} \right) \left(\frac{\vartheta(\rho\zeta\omega^2)}{\vartheta(\rho\zeta\omega)} \right)^m \left(\frac{\vartheta(\zeta\omega^2)}{\vartheta(\zeta\omega)} \right)^{n-m}. \end{aligned} \quad (3.6)$$

In Section 6, we will identify this expression for the partition function with another expression, which we will derive in the next section.

4 The partition function

In the previous section we were able to write Filali's determinant formula of the partition function in terms of the polynomials T . In this section we instead start with the partition function written in the form

$$\sum_{\text{states}} \prod_{\text{vertices}} w(\text{vertex}) \prod_{\text{turns}} w(\text{turn}).$$

We specialize the variables in the same way as in the last section to get an expression in terms of a special case of the three-color model. We follow the same steps as in [9, Section 3], but in the present paper, we specialize one of the variables slightly differently, which forces us to treat one of the columns separately through all computations.

Specialize $\lambda_i = 1$ and $\mu_j = 0$ for all $1 \leq i \leq n$ and $1 \leq j \leq n - 1$ in the partition function. Everywhere, except at the vertices depending on μ_n , the weights are thus $w(1, \rho q^a)$, where a is the height of the face in the upper left or lower left corner as explained in Section 2.2. On the rightmost column of vertices, each vertex weight is either $w(1 + \mu_n, \rho q^a)$ or $w(1 - \mu_n, \rho q^a)$. The weights of the turns are $w(1, \rho, \zeta)$. The partition function can hence be written

$$\begin{aligned} & Z_n(q, \dots, q, 1, \dots, 1, q^{\mu_n}, \rho, \zeta) \\ &= \sum_{\text{states}} \left(\prod_{\text{turns}} w(1, \rho, \zeta) \prod_{\substack{\text{vertices not in the} \\ \text{rightmost column}}} w(1, \rho q^a) \prod_{\substack{\text{vertices in the} \\ \text{rightmost column}}} w(1 \pm \mu_n, \rho q^a) \right). \end{aligned}$$

We first focus on the weights not depending on μ_n . As in [9], we would like to factor things out from the weights, so that we can write all weights with one single formula, depending only on the heights of the adjacent faces of each vertex, as opposed to the original weights which have different formulas depending on the spins around the vertex. The following lemma is similar to [9, Proposition 3.2] or [21, Proposition 7.1].

Lemma 4.1. *Let $\lambda_i = 1$ and $\mu_j = 0$ for all i and $1 \leq j \leq n - 1$. For each vertex, let a, b, c, d denote the heights on the adjacent faces as in Figure 5, and for each turn, let a be the height inside the turn. Let $\nu(w)$ be the number of vertices of type w in a state, and let $\nu_r(w)$ be the number of vertices of type w in the rightmost vertex column. For each state,*

$$\prod_{\substack{\text{vertices not in the} \\ \text{rightmost column}}} w(1, \rho q^a) \prod_{\text{turns}} w(1, \rho, \zeta) = P \prod_{\substack{\text{vertices not in the} \\ \text{rightmost column}}} \frac{\vartheta(\rho q^{\frac{3}{2}a - b + \frac{1}{2}d})}{\vartheta(\rho q^a)} \prod_{\text{turns}} \frac{\vartheta(\rho^{(1-a)/2} \zeta q^{-1})}{\vartheta(\rho^{(1-a)/2} \zeta q)},$$

where

$$P = q^{-n(n-1) + \nu(b_+) - \nu_r(b_+) + \nu(c_-) - \nu_r(c_-) + n} \left(\frac{\vartheta(q^2)}{\vartheta(q)} \right)^{2n(n-1) - \nu(b_+, b_-, c_+, c_-) + \nu_r(b_+, b_-, c_+, c_-)}.$$

Proof. We have

$$\begin{aligned} a_+(1, \rho q^a) &= a_-(1, \rho q^a) = \frac{q^{-1/2} \vartheta(q^2)}{\vartheta(q)}, \\ b_+(1, \rho q^a) &= \frac{q^{1/2} \vartheta(\rho q^{a-1})}{\vartheta(\rho q^a)}, & b_-(1, \rho q^a) &= \frac{q^{-1/2} \vartheta(\rho q^{a+1})}{\vartheta(\rho q^a)}, \\ c_+(1, \rho q^a) &= \frac{q^{-1/2} \vartheta(\rho q^{a+1})}{\vartheta(\rho q^a)}, & c_-(1, \rho q^a) &= \frac{q^{1/2} \vartheta(\rho q^{a-1})}{\vartheta(\rho q^a)}, \\ k_+(1, \rho, \zeta) &= \frac{q \vartheta(\rho \zeta q^{-1})}{\vartheta(\rho \zeta q)}, & k_-(1, \rho, \zeta) &= \frac{q \vartheta(\zeta q^{-1})}{\vartheta(\zeta q)}. \end{aligned}$$



Figure 5. Vertices with heights a , b , c and d on the adjacent faces.

From each vertex weight $w(1, \rho q^a)$, factor out $q^{-1/2}$, and put it in a prefactor P . Then factor out $\frac{\vartheta(q^2)}{\vartheta(q)}$ from the weights of type a_+ and a_- , and factor out q from b_+ and c_- . From each weight of a turn, factor out q . Then we get new vertex weights

$$\begin{aligned} \tilde{a}_+(1, \rho q^a) &= \tilde{a}_-(1, \rho q^a) = 1, \\ \tilde{b}_+(1, \rho q^a) &= \tilde{c}_-(1, \rho q^a) = \frac{\vartheta(\rho q^{a-1})}{\vartheta(\rho q^a)}, & \tilde{b}_-(1, \rho q^a) &= \tilde{c}_+(1, \rho q^a) = \frac{\vartheta(\rho q^{a+1})}{\vartheta(\rho q^a)}, \\ \tilde{k}_+(1, \rho, \zeta) &= \frac{\vartheta(\rho \zeta q^{-1})}{\vartheta(\rho \zeta q)}, & \tilde{k}_-(1, \rho, \zeta) &= \frac{\vartheta(\zeta q^{-1})}{\vartheta(\zeta q)}, \end{aligned}$$

and the prefactor is

$$P = (q^{-1/2})^{2n(n-1)} \left(\frac{\vartheta(q^2)}{\vartheta(q)} \right)^{2n(n-1) - \nu(b_+, b_-, c_+, c_-) + \nu_r(b_+, b_-, c_+, c_-)} q^{\nu(b_+) - \nu_r(b_+) + \nu(c_-) - \nu_r(c_-) + n}.$$

One can now verify that for each vertex of type \tilde{a}_\pm , \tilde{b}_\pm or \tilde{c}_\pm , we have

$$\tilde{w}(1, \rho q^a) = \frac{\vartheta(\rho q^{\frac{3}{2}a - b + \frac{1}{2}d})}{\vartheta(\rho q^a)},$$

where a , b , c and d are the heights on the adjacent faces as in Figure 5. Furthermore,

$$\tilde{k}_\pm(1, \rho, \zeta) = \frac{\vartheta(\rho^{(1-a)/2} \zeta q^{-1})}{\vartheta(\rho^{(1-a)/2} \zeta q)},$$

where a is the height of the face inside the turn. ■

The following lemma is proven in [9].

Lemma 4.2. *For any given state of the 8VSOS model with DWBC and reflecting end, let $\nu(w)$ be the number of vertices or turns of type w . Then we have*

$$\nu(b_+) = \nu(b_-) + \binom{n+1}{2} \quad \text{and} \quad \nu(c_+) + 2\nu(k_-) = \nu(c_-) + n.$$

Because of the boundary conditions, we have $2n$ outgoing arrows on the edges to the right, and ingoing arrows on the edges on the top and the bottom. The ice rule thus forces the arrows on the horizontal edges between the $(n-1)$ th and the n th column (counted from the left) to be right arrows, except for in one place, where the arrow must be a left arrow, see Figure 6. Given the placement of the left arrow in the second to last column, we know the vertex weights on the entire rightmost column of vertices. We have the following lemma.

Lemma 4.3. *For any given state of the 8VSOS model with DWBC and reflecting end, there is exactly one left arrow on the edges between the $(n-1)$ th and the n th column of vertices. Assume*




Figure 6. On the edges between the $(n-1)$ th and n th column of vertices, there is exactly one left arrow in each state. The single left arrow is on row $l = 2k - 1$ in the left lattice, and on row $l = 2k$ in the right lattice, counted from below.

that the left-pointing arrow is in the l th row counted from below. In the case that $l = 2k - 1$ is odd, the contribution to the partition function from the rightmost column of vertices, i.e., where the weights are depending on μ_n , is

$$\begin{aligned} \prod_{\text{vertices in the rightmost column}} w(1 \pm \mu_n, \rho q^a) &= \prod_{i=1}^{k-1} [b_+(1 + \mu_n, \rho q^{-n+2i-1})a_+(1 - \mu_n, \rho q^{-n+2i+1})] \\ &\quad \times c_+(1 + \mu_n, \rho q^{-n+2k-1})b_+(1 - \mu_n, \rho q^{-n+2k-1}) \\ &\quad \times \prod_{i=k+1}^n [a_-(1 + \mu_n, \rho q^{-n+2i-3})b_+(1 - \mu_n, \rho q^{-n+2i-1})]. \end{aligned}$$

In the case that $l = 2k$ is even, the contribution is

$$\begin{aligned} \prod_{\text{vertices in the rightmost column}} w(1 \pm \mu_n, \rho q^a) &= \prod_{i=1}^{k-1} [b_+(1 + \mu_n, \rho q^{-n+2i-1})a_+(1 - \mu_n, \rho q^{-n+2i+1})] \\ &\quad \times b_+(1 + \mu_n, \rho q^{-n+2k-1})c_-(1 - \mu_n, \rho q^{-n+2k-1}) \\ &\quad \times \prod_{i=k+1}^n [a_-(1 + \mu_n, \rho q^{-n+2i-3})b_+(1 - \mu_n, \rho q^{-n+2i-1})]. \end{aligned}$$

Because of Lemma 4.2, we know the difference between the number of b_+ and b_- vertices, and the difference between the number of c_+ and c_- vertices. Because of Lemma 4.3, we know the number of b_{\pm} and c_{\pm} in the rightmost vertex column, and they only depend on whether l is even or odd.

Corollary 4.4. Let $\nu_r(w)$ be the number of vertices of type w in the rightmost vertex column, and let l be the row where the single left arrow of the rightmost column sits. Then

$$\nu_r(b_+) = n, \quad \nu_r(b_-) = 0,$$

and

$$\nu_r(c_+) = \begin{cases} 1, & \text{if } l \text{ is odd,} \\ 0, & \text{if } l \text{ is even,} \end{cases} \quad \nu_r(c_-) = \begin{cases} 0, & \text{if } l \text{ is odd,} \\ 1, & \text{if } l \text{ is even.} \end{cases}$$

With $\eta = -2/3$, the prefactor from Lemma 4.1 thus becomes

$$P = (-1)^{\binom{n+1}{2}-1} \omega^{-\binom{n+1}{2}-2(n-m)+\alpha},$$

where $\alpha = (-1)^l$ and m is the number of k_+ turns. As in [9, 21], specializing $\eta = -2/3$ yields that the weights from Lemma 4.1 now become

$$\tilde{w}(1, \rho q^a) = \frac{\vartheta(\rho q^{\frac{3}{2}a-b+\frac{1}{2}d})}{\vartheta(\rho q^a)} = \frac{\vartheta(\rho q^{-b-d})}{\vartheta(\rho q^a)} = \frac{\vartheta(\rho^3, p^3)}{\vartheta(\rho \omega^a) \vartheta(\rho \omega^b) \vartheta(\rho \omega^d)},$$

since b , d and $-b-d$ are noncongruent modulo 3. Hence for each vertex in the lattice, except for in the rightmost column of vertices, we just need to know the heights of three of the adjacent faces. As in [9, 21], this means that we need to account for each face in the interior three times, but in the present article we need to treat the n th column of faces differently. On the boundary, the number of faces that we need to count differs, but here we know the heights, so we can explicitly compute this contribution. Likewise it differs in the n th column, but given the number l , we can compute this contribution explicitly as well. On the right boundary, we should not count any faces at all, since these faces only have to do with the weights containing μ_n . The faces of each turn should be accounted for only once.

The expression of Lemma 4.1 is now given by

$$\begin{aligned} & \prod_{\substack{\text{vertices not in the} \\ \text{rightmost column}}} w(1, \rho \omega^a) \prod_{\text{turns}} w(1, \rho, \zeta) \\ &= PC \vartheta(\rho^3, p^3)^{2n(n-1)} \prod_{\text{faces}} \frac{1}{\vartheta(\rho \omega^a)^3} \prod_{\text{turns}} \vartheta(\rho \omega^a)^2 \frac{\vartheta(\rho^{(1-a)/2} \zeta \omega^2)}{\vartheta(\rho^{(1-a)/2} \zeta \omega)} \\ &= PC \vartheta(\rho^3, p^3)^{2n(n-1)} \vartheta(\rho \omega)^{2(n-m)} \vartheta(\rho \omega^2)^{2m} \left(\frac{\vartheta(\rho \zeta \omega^2)}{\vartheta(\rho \zeta \omega)} \right)^m \left(\frac{\vartheta(\zeta \omega^2)}{\vartheta(\zeta \omega)} \right)^{n-m} \prod_{\text{faces}} \frac{1}{\vartheta(\rho \omega^a)^3}, \end{aligned}$$

where m is the number of positive turns and C is the correction compensating for the faces that are counted too many times. The correction C is given by

$$\begin{aligned} C &= R \vartheta(\rho)^{n+3} \vartheta(\rho q^{n-1})^2 \vartheta(\rho q^{-n+1})^3 \prod_{i=1}^{n-2} \vartheta(\rho q^i) \prod_{i=-n}^n \vartheta(\rho q^i)^3 \prod_{i=1}^{n-2} \vartheta(\rho q^{-i})^2 \\ &= \vartheta(\rho)^n \vartheta(\rho^3, p^3)^{3n} R \times \begin{cases} \vartheta(\rho)^3 \vartheta(\rho \omega) \vartheta(\rho \omega^2), & n \equiv 0 \pmod{3}, \\ \vartheta(\rho)^5, & n \equiv 1 \pmod{3}, \\ \vartheta(\rho \omega)^2 \vartheta(\rho \omega^2)^3, & n \equiv 2 \pmod{3}, \end{cases} \end{aligned}$$

where R is the contribution to the correction from the second to last column of faces, which depends on l . We have

$$\begin{aligned} R &= \prod_{i=1}^{k-1} (\vartheta(\rho q^{-n+2i}) \vartheta(\rho q^{-n+2i+1})^2) \prod_{i=k}^{n-1} (\vartheta(\rho q^{-n+2i-1})^2 \vartheta(\rho q^{-n+2i})) \\ &\times \begin{cases} \vartheta(\rho q^{-n+2k-2}), & \text{for } l = 2k - 1 \text{ odd,} \\ \vartheta(\rho q^{-n+2k}), & \text{for } l = 2k \text{ even.} \end{cases} \end{aligned}$$

The expression in Lemma 4.3 is

$$\begin{aligned} \prod_{\substack{\text{vertices in the} \\ \text{rightmost column}}} w(1 \pm \mu_n, \rho q^a) &= \frac{\vartheta(\rho q^{-n}) \vartheta(\rho q^{-n+2}) \cdots \vartheta(\rho q^{n-4}) \vartheta(\rho q^{n-2})}{\vartheta(\rho q^{-n+1}) \vartheta(\rho q^{-n+3}) \cdots \vartheta(\rho q^{n-3}) \vartheta(\rho q^{n-1})} \\ &\times \frac{(\vartheta(q^{1+\mu_n}) \vartheta(q^{2-\mu_n}))^{k-1} (\vartheta(q^{2+\mu_n}) \vartheta(q^{1-\mu_n}))^{n-k}}{\vartheta(q)^{2n-1} \vartheta(\rho q^{-n+2k-1})} \\ &\times \begin{cases} \vartheta(\rho q^{-n+2k+\mu_n}) \vartheta(q^{1-\mu_n}), & \text{for } l = 2k - 1 \text{ odd,} \\ q^{1-\mu_n} \vartheta(\rho q^{-n+2k-2+\mu_n}) \vartheta(q^{1+\mu_n}), & \text{for } l = 2k \text{ even.} \end{cases} \end{aligned}$$

Putting all of the above together, we get the following proposition.

Proposition 4.5. *Let $\eta = -2/3$, $\lambda_i = 1$ for all i , and $\mu_j = 0$ for $1 \leq j \leq n - 1$. Then the partition function is*

$$\begin{aligned} Z_n(\omega, \dots, \omega, 1, \dots, 1, q^{\mu_n}, \rho, \zeta) &= (-\omega^2)^{\binom{n+1}{2}+1} \frac{\vartheta(\rho)^{n+1} \vartheta(\rho^3, p^3)^{2n(n+1)-1}}{\vartheta(\omega)^{2n-1}} \tilde{X} \\ &\times \sum_{m=0}^n \frac{\vartheta(\rho\omega)^{2(n-m)} \vartheta(\rho\omega^2)^{2m}}{\omega^{2(n-m)}} \left(\frac{\vartheta(\rho\zeta\omega^2)}{\vartheta(\rho\zeta\omega)} \right)^m \left(\frac{\vartheta(\zeta\omega^2)}{\vartheta(\zeta\omega)} \right)^{n-m} \\ &\times \sum_{l=0}^{2n} (\vartheta(q^{1+\mu_n}) \vartheta(q^{2-\mu_n}))^{k-1} (\vartheta(q^{2+\mu_n}) \vartheta(q^{1-\mu_n}))^{n-k} \\ &\times \vartheta(\rho q^{-n+l}) \vartheta(\rho q^{-n+l-1}) \vartheta(\rho q^{-n+l+1+\mu_n}) \tilde{Y} \sum_{\substack{\text{states with} \\ m \text{ positive turns} \\ \text{and } \leftarrow \text{ on } l\text{th row}}} \prod_{\text{faces}} \frac{1}{\vartheta(\rho\omega^a)^3}, \end{aligned}$$

where k is defined by $l = 2k$ for even l , and by $l = 2k - 1$ for odd l ,

$$\tilde{X} = \begin{cases} \vartheta(\rho)^2, & n \equiv 0 \pmod{3}, \\ \vartheta(\rho) \vartheta(\rho\omega^2), & n \equiv 1 \pmod{3}, \\ \vartheta(\rho\omega) \vartheta(\rho\omega^2), & n \equiv 2 \pmod{3}, \end{cases}$$

and

$$\tilde{Y} = \begin{cases} \vartheta(q^{1-\mu_n}), & \text{for } l = 2k - 1 \text{ odd,} \\ q^{-\mu_n} \vartheta(q^{1+\mu_n}), & \text{for } l = 2k \text{ even.} \end{cases}$$

In the partition function in Proposition 4.5, we now specify $\mu_n = 1/4$. Then

$$\begin{aligned} Z_n(\omega, \dots, \omega, 1, \dots, 1, -\omega, \rho, \zeta) &= (-\omega^2)^{\binom{n+1}{2}+1} \frac{\vartheta(\rho)^{n+3} \vartheta(-1)^{2n-1} \vartheta(\rho^3, p^3)^{2n(n+1)-1}}{\vartheta(\omega)^{2n-1}} X \\ &\times \sum_{m=0}^n \frac{\vartheta(\rho\omega)^{2(n-m)} \vartheta(\rho\omega^2)^{2m}}{\omega^{2(n-m)}} \left(\frac{\vartheta(\rho\zeta\omega^2)}{\vartheta(\rho\zeta\omega)} \right)^m \left(\frac{\vartheta(\zeta\omega^2)}{\vartheta(\zeta\omega)} \right)^{n-m} \\ &\times \sum_{l=0}^{2n} \left((-1)^{l-1} \omega^{l-1} \left(\frac{\vartheta(-\omega)}{\vartheta(-1)} \right)^{l-1} \vartheta(\rho\omega^{-n+l}) \vartheta(\rho\omega^{-n+l-1}) \vartheta(-\rho\omega^{-n+l+2}) \right. \\ &\left. \times \sum_{\substack{\text{states with} \\ m \text{ positive turns} \\ \text{and } \leftarrow \text{ on } l\text{th row}}} \prod_{\text{faces}} \frac{1}{\vartheta(\rho\omega^a)^3} \right), \end{aligned} \tag{4.1}$$

for

$$X = \begin{cases} 1, & n \equiv 0 \pmod{3}, \\ \frac{\vartheta(\rho\omega^2)}{\vartheta(\rho)}, & n \equiv 1 \pmod{3}, \\ \frac{\vartheta(\rho\omega)\vartheta(\rho\omega^2)}{\vartheta(\rho)^2}, & n \equiv 2 \pmod{3}. \end{cases}$$

The 6V model with DWBC and a reflecting end with the unique left arrow of the second to last column fixed to a given edge l has earlier been studied, e.g., in the context of so-called boundary correlation functions (see, e.g., [27]). These functions describe the probability that the second to last column of a state has its unique left arrow on the l th row. It is also related to refined enumerations of UASMs (see, e.g., [19]), where the objective is to count the number of UASMs that have the unique 1 of the rightmost column on the l th row.

5 Identification of terms

In this section, we combine the two expressions (3.6) and (4.1) for the partition function and identify the terms with the same m . This we can do since $\left(\frac{\vartheta(\rho\zeta\omega^2)}{\vartheta(\rho\zeta\omega)}\right)^m \left(\frac{\vartheta(\zeta\omega^2)}{\vartheta(\zeta\omega)}\right)^{n-m}$ are linearly independent as functions of ζ . Then

$$\begin{aligned} & \frac{(-1)^n E^{n^2-n} D^{n-1} B'}{\vartheta(\rho)^{2n^2+4n+3} \vartheta(\rho\omega)^{2n^2+4n-3m} \vartheta(\rho\omega^2)^{2n^2+n+3m}} T(2\psi+1, \dots, 2\psi+1, \psi) \\ & \times \left(\binom{n-1}{m-1} \vartheta(\rho)\vartheta(\rho\omega^2)\vartheta(-\rho\omega^2) - \binom{n-1}{m} \omega \frac{\vartheta(-\omega)}{\vartheta(-1)} \vartheta(\rho)\vartheta(\rho\omega)\vartheta(-\rho) \right) \\ & = \sum_{l=0}^{2n} \left((-1)^l \left(\omega \frac{\vartheta(-\omega)}{\vartheta(-1)} \right)^{l-1} \vartheta(\rho\omega^{-n+l}) \vartheta(\rho\omega^{-n+l+2}) \vartheta(-\rho\omega^{-n+l+2}) \right. \\ & \times \left. \sum_{\substack{\text{states with} \\ m \text{ positive turns} \\ \text{and } \leftarrow \text{ on } l\text{th row}}} \prod_{\text{faces}} \frac{1}{\vartheta(\rho\omega^a)^3} \right), \end{aligned} \quad (5.1)$$

with

$$E = \frac{\vartheta(-\omega)^2 \vartheta(-p^{1/2}) \vartheta(p^{1/2}\omega)^3}{\vartheta(\omega)^2 \vartheta(-1) \vartheta(p^{1/2})^2 \vartheta(-p^{1/2}\omega)^4}, \quad D = \frac{\omega^2 \vartheta(-p^{1/2}\omega) \vartheta(-p^{1/2}) \vartheta(\omega)^2}{\vartheta(p^{1/2}\omega)^2 \vartheta(-1)^2},$$

and

$$B' = \begin{cases} 1, & n \equiv 0, 1 \pmod{3}, \\ \frac{\vartheta(\rho)^2}{\vartheta(\rho\omega)\vartheta(\rho\omega^2)}, & n \equiv 2 \pmod{3}. \end{cases}$$

Focus on the right hand side of (5.1). Let $N_{m,l}(k_0, k_1, k_2)$ be the number of states with m positive turns, the left arrow between the $(n-1)$ th and n th column in the l th row from below, and with k_i faces of color i . Then

$$\sum_{\substack{\text{states with} \\ m \text{ positive turns} \\ \text{and } \leftarrow \text{ on } l\text{th row}}} \prod_{\text{faces}} \frac{1}{\vartheta(\rho\omega^a)^3} = \sum_{(k_0, k_1, k_2) \in \mathbb{Z}^3} N_{m,l}(k_0, k_1, k_2) \frac{1}{\vartheta(\rho)^{3k_0} \vartheta(\rho\omega)^{3k_1} \vartheta(\rho\omega^2)^{3k_2}}.$$

This is the partition function for the three-color model with fixed values of m and l . That a state has its left arrow on the l th row from below means for the three-coloring that when starting from below, the single edge where the color decreases by 1 mod 3 is the l th edge from below.

5.1 Specialization of the dynamical parameter

At this point in the investigation of the polynomials q_n in the previous article [9], we defined the parameters $t_i = 1/\vartheta(\rho\omega^i)^3$ to be able to study the partition function of the three-color model. However, due to the summation over l in the present article, the same substitution in (5.1) leads to an expression with sums, which seems hard to make use of to study the full three-color model partition function. Therefore, we will not use these parameters. Instead, we will consider the special cases $\rho = -\omega^i$, $i = 0, 1, 2$. This allows us to study cases of the three-color model where two colors have equal weight.

Rewrite the sum on the right hand side of (5.1) as three separate sums for $l \equiv n, n+1, n+2 \pmod 3$ respectively. Define

$$\xi = \omega \frac{\vartheta(-\omega)}{\vartheta(-1)}.$$

Then (5.1) becomes

$$\begin{aligned} & \frac{(-1)^n E^{n^2-n} D^{n-1} B'}{\vartheta(\rho)^{2n^2+4n+3} \vartheta(\rho\omega)^{2n^2+4n-3m} \vartheta(\rho\omega^2)^{2n^2+n+3m}} T(2\psi+1, \dots, 2\psi+1, \psi) \\ & \times \left(\binom{n-1}{m-1} \vartheta(\rho) \vartheta(\rho\omega^2) \vartheta(-\rho\omega^2) - \binom{n-1}{m} \xi \vartheta(\rho) \vartheta(\rho\omega) \vartheta(-\rho) \right) \\ & = \sum_{(k_0, k_1, k_2) \in \mathbb{Z}^3} \sum_{l \equiv n \pmod 3} ((-1)^l \xi^{l-1} \vartheta(\rho) \vartheta(\rho\omega^2) \vartheta(-\rho\omega^2) N_{m,l}(k_0, k_1, k_2) \\ & \quad + (-1)^l \xi^l \vartheta(\rho\omega) \vartheta(\rho) \vartheta(-\rho) N_{m,l+1}(k_0, k_1, k_2) \\ & \quad + (-1)^{l+1} \xi^{l+1} \vartheta(\rho\omega^2) \vartheta(\rho\omega) \vartheta(-\rho\omega) N_{m,l+2}(k_0, k_1, k_2)) \\ & \quad \times \frac{1}{\vartheta(\rho)^{3k_0} \vartheta(\rho\omega)^{3k_1} \vartheta(\rho\omega^2)^{3k_2}}. \end{aligned} \tag{5.2}$$

Now specify $\rho = -1$. Terms with $\vartheta(-\rho)$ will then vanish. Divide both sides by $\vartheta(\omega)\vartheta(-1)^2$. Observe that $k_0 + k_1 + k_2 = (2n+1)(n+1)$ and multiply both sides by $\vartheta(-\omega)^{3(2n+1)(n+1)}$. Then we get

$$\begin{aligned} & E^{n^2-n} D^{n-1} \xi^{2n^2+4n+b} \binom{n-1}{m-1} T(2\psi+1, \dots, 2\psi+1, \psi) \\ & = \sum_{(k_0, k_1, k_2) \in \mathbb{Z}^3} \sum_{l \equiv n \pmod 3} (-1)^{n+l} \xi^l (N_{m,l}(k_0, k_1, k_2) + N_{m,l-1}(k_0, k_1, k_2)) \xi^{3k_0}, \end{aligned}$$

where

$$b = \begin{cases} 4, & n \equiv 0, 1 \pmod 3, \\ 2, & n \equiv 2 \pmod 3. \end{cases}$$

We can thus rewrite the sum to depend only on k_0 , and not on k_1 and k_2 . We write

$$\begin{aligned} & E^{n^2-n} D^{n-1} \xi^{2n^2+4n+b} \binom{n-1}{m-1} T(2\psi+1, \dots, 2\psi+1, \psi) \\ & = \sum_{k_0 \in \mathbb{Z}} \sum_{l \equiv n \pmod 3} (-1)^{n+l} \xi^l (N_{m,l}(k_0) + N_{m,l-1}(k_0)) \xi^{3k_0}, \end{aligned} \tag{5.3}$$

where $N_{m,l}(k_i)$ is the number of states with m positive turns, the left arrow between the $(n-1)$ th and n th column in the l th row from below, and with k_i faces of color i .

As in [9],

$$\left(\frac{\vartheta(\omega)^2 \vartheta(-1) \vartheta(p^{1/2})^2 \vartheta(-p^{1/2}\omega)^4}{\vartheta(-\omega)^2 \vartheta(-p^{1/2}) \vartheta(p^{1/2}\omega)^3} \right)^6 = 2^4 \psi^2 (\psi + 1)^8 (2\psi + 1)^6, \quad (5.4)$$

$$\frac{1}{\xi^2} \left(\frac{\vartheta(\omega)^2 \vartheta(-1) \vartheta(p^{1/2})^2 \vartheta(-p^{1/2}\omega)^4}{\vartheta(-\omega)^2 \vartheta(-p^{1/2}) \vartheta(p^{1/2}\omega)^3} \right)^2 = (2\psi(\psi + 1)(2\psi + 1))^2, \quad (5.5)$$

$$\left(\frac{\omega^2 \vartheta(-p^{1/2}\omega) \vartheta(-p^{1/2}) \vartheta(\omega)^2}{\vartheta(p^{1/2}\omega)^2 \vartheta(-1)^2} \right)^3 = \frac{(\psi + 1)(2\psi + 1)^3}{2\psi^5}, \quad (5.6)$$

$$\frac{1}{\xi} \frac{\omega^2 \vartheta(-p^{1/2}\omega) \vartheta(-p^{1/2}) \vartheta(\omega)^2}{\vartheta(p^{1/2}\omega)^2 \vartheta(-1)^2} = \frac{2\psi + 1}{\psi}, \quad (5.7)$$

and

$$\xi^3 = \frac{\psi + 1}{2\psi^2}.$$

Insert (5.4)–(5.7) into (5.3). We need to separate the different cases for $n \pmod 3$, since

$$2n^2 + 4n + b \equiv \begin{cases} 1 \pmod 3, & \text{for } n \equiv 0, 1 \pmod 3, \\ 0 \pmod 3, & \text{for } n \equiv 2 \pmod 3, \end{cases}$$

and

$$n^2 - n \equiv \begin{cases} 0 \pmod 6, & \text{for } n \equiv 0, 1 \pmod 3, \\ 2 \pmod 6, & \text{for } n \equiv 2 \pmod 3. \end{cases}$$

Although the computations are slightly different,

$$E^{n^2-n} D^{n-1} = \xi^{2n^2-n-1} \left(\frac{\psi}{2\psi + 1} \right)^{n^2-2n+1} (\psi + 1)^{2n-2n^2},$$

for all n , where we have used (3.2) and (3.3).

The equation then finally becomes

$$\begin{aligned} & \binom{n-1}{m-1} T(2\psi + 1, \dots, 2\psi + 1, \psi) \\ &= \left(\frac{2\psi + 1}{\psi} \right)^{n^2-2n+1} (\psi + 1)^{2n^2-2n} \sum_{\substack{l \equiv n \pmod 3 \\ k_0 \in \mathbb{Z}}} (-1)^{n+l} (N_{m,l}(k_0) + N_{m,l-1}(k_0)) \xi^{3k_0-4n^2-3n+l-c}, \end{aligned} \quad (5.8)$$

for

$$c = \begin{cases} 3, & n \equiv 0, 1 \pmod 3, \\ 1, & n \equiv 2 \pmod 3. \end{cases}$$

Likewise if we put $\rho = -\omega$ into (5.2) we get

$$\binom{n-1}{m} T(2\psi + 1, \dots, 2\psi + 1, \psi) \quad (5.9)$$

$$= \left(\frac{2\psi+1}{\psi} \right)^{n^2-2n+1} (\psi+1)^{2n^2-2n} \sum_{\substack{l \equiv n \pmod{3} \\ k_2 \in \mathbb{Z}}} (-1)^{n+l} (N_{m,l+1}(k_2) + N_{m,l+2}(k_2)) \xi^{3k_2-4n^2-3m+l-d},$$

and putting $\rho = -\omega^2$ into (5.2) yields

$$\begin{aligned} \binom{n}{m} T(2\psi+1, \dots, 2\psi+1, \psi) & \quad (5.10) \\ &= \left(\frac{2\psi+1}{\psi} \right)^{n^2-2n+1} (\psi+1)^{2n^2-2n} \sum_{\substack{l \equiv n \pmod{3} \\ k_1 \in \mathbb{Z}}} (-1)^{n+l} (N_{m,l}(k_1) + N_{m,l+1}(k_1)) \xi^{3k_1-4n^2-3n+3m+l-d}, \end{aligned}$$

for

$$d = \begin{cases} 0, & n \equiv 0, 1 \pmod{3}, \\ 1, & n \equiv 2 \pmod{3}. \end{cases}$$

6 Formulas for $p_n(z)$

Bazhanov and Mangazeev [4] introduced certain polynomials, describing the ground state eigenvalue of Baxter's Q -operator [2] for the 8V model in the case with $\eta = -2/3$. These polynomials are given by

$$\mathcal{P}_n(x, z) = \sum_{k=0}^n r_k^{(n)}(z) x^k,$$

normalized by $r_n^{(n)}(0) = 1$. Furthermore, they introduced $s_n(z) = r_n^{(n)}(z)$ and $\bar{s}_n(z) = r_0^{(n)}(z)$. In a later paper [16], they connected these polynomials to the ground state eigenvectors of the supersymmetric XYZ-Hamiltonian for spin chains of odd length $2n+1$. Several conjectures about the polynomials were stated, including that $s_n(z)$ factors into certain polynomials which seem to have positive coefficients. Here, polynomials $p_n(z)$ and $q_n(z)$, with $\deg p_n(z) = \deg q_n(z) = n(n+1)$ and $p_n(0) = q_n(0) = 1$, show up as factors of $s_{2n}(z^2)$. In further conjectures, they suggest that certain components of the ground state eigenvectors for the supersymmetric XYZ spin chain can be written in terms of $s_n(z)$, $\bar{s}_n(z)$, $p_n(z)$ and $q_n(z)$. In [9], we investigated the polynomials $q_n(z)$. In this paper, we turn the focus to the polynomials $p_n(z)$.

Zinn-Justin [29] investigated a family of polynomials equivalent to Rosengren's polynomials T . He realized that certain specializations of the parameters in his polynomials seem to yield the polynomials of Bazhanov and Mangazeev. Let [24]

$$\xi_0 = 2\psi + 1, \quad \xi_1 = \frac{\psi}{\psi + 2}, \quad \xi_2 = \frac{\psi(2\psi + 1)}{\psi + 2}, \quad \xi_3 = 1.$$

In the case where all parameters x_i of Rosengren's polynomials T (3.1) are one of the ξ_j 's, the only dependence on a variable is that the ξ_j 's depend on ψ . Therefore we define

$$t^{(k_0, k_1, k_2, k_3)}(\psi) := T(\underbrace{\xi_0, \dots, \xi_0}_{k_0 \text{ times}}, \underbrace{\xi_1, \dots, \xi_1}_{k_1 \text{ times}}, \underbrace{\xi_2, \dots, \xi_2}_{k_2 \text{ times}}, \underbrace{\xi_3, \dots, \xi_3}_{k_3 \text{ times}}),$$

with k_j non-negative integers and $k_0 + k_1 + k_2 + k_3 = 2n$. In [24], this definition is extended to negative values of k_i . In terms of Rosengren's polynomials, the polynomials p_n are, up to a prefactor, equivalent to $t^{(2n+1, 0, 0, -1)}(\psi)$, which in turn corresponds to $T(\underbrace{2\psi+1, \dots, 2\psi+1}_{2n+1 \text{ times}}, \psi)$.

The second correspondence can be seen by using [24, formulas (2.12) and (2.13)]. This together with the expression in [24, Section 5.3] and the symmetries in [24, Proposition 2.2] yield

$$T(\underbrace{2\psi + 1, \dots, 2\psi + 1}_{2n-1 \text{ times}}, \psi) = \left(\frac{\psi}{2\psi + 1} \right)^{n-1} ((\psi + 1)(2\psi + 1)^2)^{n^2-n} p_{n-1} \left(-\frac{1}{2\psi + 1} \right). \quad (6.1)$$

That these polynomials are equivalent to the polynomials of Mangazeev and Bazhanov is still a conjecture, but here we take (6.1) as the definition of p_n .

We insert (6.1) into (5.8) and get

$$\begin{aligned} & \binom{n-1}{m-1} p_{n-1} \left(-\frac{1}{2\psi + 1} \right) \\ &= \left(\frac{\psi + 1}{\psi(2\psi + 1)} \right)^{n^2-n} \sum_{\substack{l \equiv n \pmod{3} \\ k_0 \in \mathbb{Z}}} (-1)^{n+l} (N_{m,l}(k_0) + N_{m,l-1}(k_0)) \xi^{3k_0 - 4n^2 - 3n + l - c}, \end{aligned}$$

where

$$c = \begin{cases} 3, & n \equiv 0, 1 \pmod{3}, \\ 1, & n \equiv 2 \pmod{3}. \end{cases}$$

We can do the same substitution in (5.9) and (5.10). Finally, changing to the variable $z = -1/(2\psi + 1)$ yields the following theorem.

Theorem 6.1. *Let $N_{m,l}(k_i)$ be the number of states with m positive turns, k_i faces of color i , and where the left arrow of the second to last column is on the l th row from below. Formulas for $p_{n-1}(z)$ are then given by*

$$\binom{n-1}{m-1} p_{n-1}(z) = \sum_{\substack{l \equiv n \pmod{3} \\ k_0 \in \mathbb{Z}}} (-1)^{n+l} (N_{m,l}(k_0) + N_{m,l-1}(k_0)) \frac{(z(z-1))^{(3k_0 - n^2 - 6n + l - c)/3}}{(z+1)^{(6k_0 - 5n^2 - 9n + 2l - 2c)/3}},$$

where

$$c = \begin{cases} 3, & n \equiv 0, 1 \pmod{3}, \\ 1, & n \equiv 2 \pmod{3}, \end{cases}$$

$$\binom{n}{m} p_{n-1}(z) = \sum_{\substack{l \equiv n \pmod{3} \\ k_1 \in \mathbb{Z}}} (-1)^{n+l} (N_{m,l}(k_1) + N_{m,l+1}(k_1)) \frac{(z(z-1))^{(3k_1 - n^2 - 6n + 3m + l - d)/3}}{(z+1)^{(6k_1 - 5n^2 - 9n + 6m + 2l - 2d)/3}},$$

and

$$\binom{n-1}{m} p_{n-1}(z) = \sum_{\substack{l \equiv n \pmod{3} \\ k_2 \in \mathbb{Z}}} (-1)^{n+l} (N_{m,l+1}(k_2) + N_{m,l+2}(k_2)) \frac{(z(z-1))^{(3k_2 - n^2 - 3n - 3m + l - d)/3}}{(z+1)^{(6k_2 - 5n^2 - 3n - 6m + 2l - 2d)/3}},$$

where

$$d = \begin{cases} 0, & n \equiv 0, 1 \pmod{3}, \\ 1, & n \equiv 2 \pmod{3}. \end{cases}$$

The polynomials $p_n(z)$ are known to have the symmetry [16]

$$p_n(z) = \left(\frac{1+3z}{2}\right)^{n(n+1)} p_n\left(\frac{1-z}{1+3z}\right),$$

and it is immediate that this holds for the expressions in Theorem 6.1.

Theorem 6.1 gives three expressions for $p_{n-1}(z)$ in terms of three-colorings. The functions $p_{n-1}(z)$ are known to be polynomials, so this must of course hold for the right hand sides of the expressions as well, even though the individual terms in the sums may be rational functions, with possible poles in $z = -1, 0, 1$. Any terms with poles and nonzero coefficients have to cancel. In the corresponding expression for $q_{n-1}(z)$, the individual terms can not contain poles, and this let us, among other things, draw conclusions about the possible values for the numbers k_i [9]. Because of the factors $(-1)^{n+l}$ in the expressions for $p_{n-1}(z)$ in Theorem 6.1, similar conclusions cannot be drawn in this case.

Although we have found combinatorial expressions for $p_{n-1}(z)$, we can not immediately see from the formulas that the coefficients are positive. To draw conclusions about the coefficients, one would need to know something more about the numbers $N_{m,l}(k_i)$.

Acknowledgements

This paper was written during my time as a Ph.D. student at the Department of Mathematical Sciences, University of Gothenburg and Chalmers University of Technology. I am very thankful to my supervisor Hjalmar Rosengren and cosupervisor Jules Lamers for a lot of encouragement and support with my research. I also would like to thank the anonymous referees for nice and useful comments.

References

- [1] Baxter R.J., Three-colorings of the square lattice: a hard squares model, *J. Math. Phys.* **11** (1970), 3116–3124.
- [2] Baxter R.J., Partition function of the eight-vertex lattice model, *Ann. Physics* **70** (1972), 193–228.
- [3] Baxter R.J., Eight-vertex model in lattice statistics and one-dimensional anisotropic Heisenberg chain I. Some fundamental eigenvectors, *Ann. Physics* **76** (1973), 1–24.
- [4] Bazhanov V.V., Mangazeev V.V., Eight-vertex model and non-stationary Lamé equation, *J. Phys. A: Math. Gen.* **38** (2005), L145–L153, [arXiv:hep-th/0411094](https://arxiv.org/abs/hep-th/0411094).
- [5] Bazhanov V.V., Mangazeev V.V., The eight-vertex model and Painlevé VI, *J. Phys. A: Math. Gen.* **39** (2006), 12235–12243, [arXiv:hep-th/0602122](https://arxiv.org/abs/hep-th/0602122).
- [6] Brasseur S., Hagendorf C., Sum rules for the supersymmetric eight-vertex model, *J. Stat. Mech. Theory Exp.* **2021** (2021), 023102, 45 pages, [arXiv:2009.14077](https://arxiv.org/abs/2009.14077).
- [7] Filali G., Elliptic dynamical reflection algebra and partition function of SOS model with reflecting end, *J. Geom. Phys.* **61** (2011), 1789–1796, [arXiv:1012.0516](https://arxiv.org/abs/1012.0516).
- [8] Hagendorf C., Fendley P., The eight-vertex model and lattice supersymmetry, *J. Stat. Phys.* **146** (2012), 1122–1155, [arXiv:1109.4090](https://arxiv.org/abs/1109.4090).
- [9] Hietala L., A combinatorial description of certain polynomials related to the XYZ spin chain, *SIGMA* **16** (2020), 101, 26 pages, [arXiv:2004.09924](https://arxiv.org/abs/2004.09924).
- [10] Izergin A.G., Partition function of the six-vertex model in a finite volume, *Soviet Phys. Dokl.* **32** (1987), 878–879.
- [11] Izergin A.G., Coker D.A., Korepin V.E., Determinant formula for the six-vertex model, *J. Phys. A: Math. Gen.* **25** (1992), 4315–4334.
- [12] Korepin V.E., Calculation of norms of Bethe wave functions, *Comm. Math. Phys.* **86** (1982), 391–418.
- [13] Kuperberg G., Another proof of the alternating-sign matrix conjecture, *Int. Math. Res. Not.* **1996** (1996), 139–150, [arXiv:math.CO/9712207](https://arxiv.org/abs/math.CO/9712207).

-
- [14] Kuperberg G., Symmetry classes of alternating-sign matrices under one roof, *Ann. of Math.* **156** (2002), 835–866, [arXiv:math.CO/0008184](#).
- [15] Lieb E.H., Residual entropy of square ice, *Phys. Rev.* **162** (1967), 162–172.
- [16] Mangazeev V.V., Bazhanov V.V., The eight-vertex model and Painlevé VI equation II: eigenvector results, *J. Phys. A: Math. Theor.* **43** (2010), 085206, 16 pages, [arXiv:0912.2163](#).
- [17] Mills W.H., Robbins D.P., Rumsey Jr. H., Alternating sign matrices and descending plane partitions, *J. Combin. Theory Ser. A* **34** (1983), 340–359.
- [18] Razumov A.V., Stroganov Yu.G., Spin chains and combinatorics, *J. Phys. A: Math. Gen.* **34** (2001), 3185–3190, [arXiv:cond-mat/0012141](#).
- [19] Razumov A.V., Stroganov Yu.G., Refined enumerations of some symmetry classes of alternating-sign matrices, *Theoret. and Math. Phys.* **141** (2004), 1609–1630, [arXiv:math-ph/0312071](#).
- [20] Razumov A.V., Stroganov Yu.G., A possible combinatorial point for the XYZ spin chain, *Theoret. and Math. Phys.* **164** (2010), 977–991, [arXiv:0911.5030](#).
- [21] Rosengren H., An Izergin–Korepin-type identity for the 8VSOS model, with applications to alternating sign matrices, *Adv. in Appl. Math.* **43** (2009), 137–155, [arXiv:0801.1229](#).
- [22] Rosengren H., The three-colour model with domain wall boundary conditions, *Adv. in Appl. Math.* **46** (2011), 481–535, [arXiv:0911.0561](#).
- [23] Rosengren H., Special polynomials related to the supersymmetric eight-vertex model. I. Behaviour at cusps, [arXiv:1305.0666](#).
- [24] Rosengren H., Special polynomials related to the supersymmetric eight-vertex model: a summary, *Comm. Math. Phys.* **340** (2015), 1143–1170, [arXiv:1503.02833](#).
- [25] Sutherland B., Exact solution of a two-dimensional model for hydrogen-bonded crystals, *Phys. Rev. Lett.* **19** (1967), 103–104.
- [26] Tsuchiya O., Determinant formula for the six-vertex model with reflecting end, *J. Math. Phys.* **39** (1998), 5946–5951, [arXiv:solv-int/9804010](#).
- [27] Wang Y.-S., Boundary spontaneous polarization in the six-vertex model with a reflecting boundary, *J. Phys. A: Math. Gen.* **36** (2003), 4007–4013.
- [28] Zeilberger D., Proof of the alternating sign matrix conjecture, *Electron. J. Combin.* **3** (1996), R13, 84 pages, [arXiv:math.CO/9407211](#).
- [29] Zinn-Justin P., Sum rule for the eight-vertex model on its combinatorial line, in Symmetries, Integrable Systems and Representations, *Springer Proc. Math. Stat.*, Vol. 40, Springer, Heidelberg, 2013, 599–637, [arXiv:1202.4420](#).