



Quadrature Domains for the Helmholtz Equation with Applications to Non-scattering Phenomena

Downloaded from: <https://research.chalmers.se>, 2025-05-15 14:02 UTC

Citation for the original published paper (version of record):

Kow, P., Larson, S., Salo, M. et al (2024). Quadrature Domains for the Helmholtz Equation with Applications to Non-scattering Phenomena. *Potential Analysis*, 60(1): 387-424.
<http://dx.doi.org/10.1007/s11118-022-10054-5>

N.B. When citing this work, cite the original published paper.



Quadrature Domains for the Helmholtz Equation with Applications to Non-scattering Phenomena

Pu-Zhao Kow¹ · Simon Larson² · Mikko Salo¹ · Henrik Shahgholian³ 

Received: 5 May 2022 / Accepted: 19 November 2022
© The Author(s) 2022

Abstract

In this paper, we introduce quadrature domains for the Helmholtz equation. We show existence results for such domains and implement the so-called partial balayage procedure. We also give an application to inverse scattering problems, and show that there are non-scattering domains for the Helmholtz equation at any positive frequency that have inward cusps.

Keywords Quadrature domain · Non-scattering phenomena · Mean value theorem · Helmholtz equation · Acoustic equation · Metaharmonic functions · Partial balayage

Mathematics Subject Classification (2010) 35J05 · 35J15 · 35J20 · 35R30 · 35R35

1 Introduction and Main Results

1.1 Background

This work is motivated by a problem in inverse scattering theory, but it raises questions of independent interest in the context of quadrature domains and free boundary problems.

✉ Henrik Shahgholian
henriksh@kth.se

Pu-Zhao Kow
pu-zhao.pz.kow@jyu.fi

Simon Larson
larsons@chalmers.se

Mikko Salo
mikko.j.salo@jyu.fi

¹ Department of Mathematics and Statistics, University of Jyväskylä, P.O. Box 35 (MaD), FI-40014, Jyväskylä, Finland

² Mathematical Sciences, Chalmers University of Technology and the University of Gothenburg, SE-412 96, Göteborg, Sweden

³ Department of Mathematics, KTH Royal Institute of Technology, SE-100 44, Stockholm, Sweden

We recall that a bounded domain $D \subset \mathbb{R}^n$ is called a *quadrature domain* (for harmonic functions), corresponding to a measure μ with $\text{supp}(\mu) \subset D$, if

$$\int_D h(x) dx = \int h(x) d\mu(x) \tag{1.1}$$

for every harmonic function $h \in L^1(D)$. More generally, one can consider distributions $\mu \in \mathcal{E}'(D)$. In the most classical case one is interested in domains D for which μ is supported at finitely many points, so that Eq. 1.1 reduces to a quadrature identity for computing integrals of harmonic functions.

Quadrature domains can be viewed as a generalization of the mean value theorem (MVT) for harmonic functions. Indeed, we can rephrase the MVT for harmonic functions as follows:

$$B_r(a) \text{ is a quadrature domain with } \mu = m(B_r(a))\delta_a,$$

where δ_a is the Dirac measure at a , m denotes the Lebesgue measure in \mathbb{R}^n (i.e. $dm = dx$) and $B_r(a)$ is the ball of radius r centered at a . In general, the boundary of a quadrature domain is a free boundary in an obstacle-type problem (see [39]), and hence near any given point $z \in \partial D$ the domain D is either smooth or D^c has zero density at z . Various examples can be constructed via complex analysis, for example, the cardioid domain in Example 3.3 below. We refer to [19], [42], and [29] for further background.

The inverse scattering problems studied in [45] lead to a related concept, for solutions of the Helmholtz equation $(\Delta + k^2)u = 0$, where $k \geq 0$ is a frequency. This setting gives rise to various interesting questions. We are not aware of earlier work on quadrature domains for $k > 0$, and in this article we only give some first steps. In addition, we show that any quadrature domain is a *non-scattering domain* (cf. Definition 1.8) if it admits an incident wave that is positive on its boundary. In [45] it was observed that in the case $k = 0$ quadrature domains are non-scattering domains, and hence there are non-scattering domains having inward cusps. Corollary 1.9 below provides a similar result valid for all $k > 0$.

1.2 Notation

Here we gather recurring notation and definitions. We also mention here that all functions and measures will be real-valued unless stated otherwise.

- m Lebesgue measure in \mathbb{R}^n
- $B_r(a)$ ball of radius r centered at a
- B_r ball of radius r centered at origin
- \mathbb{D} unit disk
- J_α Bessel function of first kind
- $j_{\alpha,1}$ the first positive zero of the Bessel function J_α
- Y_α Bessel function of second kind
- $R(n, k) = c_n^{\text{ref}} k^{-1}$ maximal length scale
- $\tilde{\Phi}_k = \tilde{\Phi}_{k,R(n,k)}$ a particular fundamental solution of the Helmholtz operator $-(\Delta + k^2)$
- $U_k^\mu = \tilde{\Phi}_k * \mu$ potential of a measure μ
- $\mathcal{F}_k(\mu)$ the class of admissible functions in an obstacle problem
- $c_{n,k,r}^{\text{MVT}}$ constant related to mean value theorem
- $D(\mu)$ saturated set for $Bal_k(\mu)$
- $\omega(\mu)$ non-contact set for an obstacle problem

1.3 Main Results

We begin with a definition generalizing Eq. 1.1.

Definition 1.1 Let $k > 0$. A bounded open set $D \subset \mathbb{R}^n$ (not necessarily connected) is called a *quadrature domain* for $(\Delta + k^2)$, or a *k-quadrature domain*, corresponding to a distribution $\mu \in \mathcal{E}'(D)$, if

$$\int_D w(x) dx = \langle \mu, w \rangle$$

for all $w \in L^1(D)$ satisfying $(\Delta + k^2)w = 0$ in D .

We remark that solutions of $(\Delta + k^2)w = 0$ are sometimes called *metaharmonic functions*, see e.g. [35, Section 4] or [22] for a discussion. It is important that $\text{supp}(\mu)$ has to be a subset of D (see however [32, Lemma 2.8] for a discussion that weakens this assumption for harmonic functions). Indeed, that $\text{supp}(\mu) \subset D$ implies that the distributional pairing $\langle \mu, w \rangle$ is well defined, because solutions of $(\Delta + k^2)w = 0$ are smooth in D . Furthermore, without this requirement the existence of a distribution satisfying the definition would be trivial, indeed one could choose $\mu = \chi_D$.

The first question is whether *k*-quadrature domains even exist for $k > 0$. This is indeed the case. In fact, balls are always *k*-quadrature domains. This is a consequence of a MVT for the Helmholtz equation which goes back to H. Weber [48, 49] (see also [35], [36], or [18, p. 289]). The MVT takes the form

$$\int_{B_r(a)} w(x) dx = c_{n,k,r}^{\text{MVT}} w(a)$$

whenever $w \in L^1(B_r(a))$ and $(\Delta + k^2)w = 0$ in $B_r(a)$. However, unlike for harmonic functions, the constant $c_{n,k,r}^{\text{MVT}}$ has varying sign depending on k, r . In particular, the constant vanishes when $J_{n/2}(kr) = 0$ where J_α denotes the Bessel function of the first kind. More details are given in Appendix A. It follows that unions of disjoint balls are also *k*-quadrature domains corresponding to linear combinations of delta functions. Choosing two balls whose closures intersect at one point furnishes an example of a *k*-quadrature domain whose boundary is not smooth.

In order to make further progress we consider a PDE characterization of *k*-quadrature domains. One can show (see Proposition 2.1) that D is a *k*-quadrature domain corresponding to $\mu \in \mathcal{E}'(D)$ if and only if there is a distribution $u \in \mathcal{D}'(\mathbb{R}^n)$ satisfying

$$\begin{cases} (\Delta + k^2)u = \chi_D - \mu & \text{in } \mathbb{R}^n, \\ u = |\nabla u| = 0 & \text{in } \mathbb{R}^n \setminus D. \end{cases} \tag{1.2}$$

Note that by elliptic regularity the distribution u solving $(\Delta + k^2)u = \chi_D$ near ∂D must be C^1 near ∂D , and thus the condition that u and ∇u vanish in $\mathbb{R}^n \setminus D$ (instead of $\mathbb{R}^n \setminus \overline{D}$) makes sense. The following result is a local version of the above fact, characterizing domains D that are *k*-quadrature domains for some distribution μ . However, there is no reason to expect that μ could be chosen to have support at finitely many points.

Theorem 1.2 *Let $k > 0$, and let D be a bounded open set in \mathbb{R}^n . Then D is a k -quadrature domain for some $\mu \in \mathcal{E}'(D)$ if and only if there is a neighborhood U of ∂D in \mathbb{R}^n and a distribution $u \in \mathcal{D}'(U)$ satisfying*

$$\begin{cases} (\Delta + k^2)u = \chi_D & \text{in } U, \\ u = |\nabla u| = 0 & \text{in } U \setminus D. \end{cases} \tag{1.3}$$

Moreover, if D is a k -quadrature domain for some $\mu \in \mathcal{E}'(D)$, then D is also a k -quadrature domain for some measure $\tilde{\mu}$ having smooth density with respect to Lebesgue measure.

Remark 1.3 If u is as in Theorem 1.2, then clearly

$$\begin{cases} \Delta u = f \chi_D & \text{in } U, \\ u = |\nabla u| = 0 & \text{in } U \setminus D, \end{cases} \tag{1.4}$$

with $f = 1 - k^2u$. Extending u from a neighborhood of ∂D into some distribution in \mathbb{R}^n with $u = |\nabla u| = 0$ in $\mathbb{R}^n \setminus D$ shows that we have an analogue of Eq. 1.2 with $k = 0$ and with χ_D replaced by $f \chi_D$. Thus any k -quadrature domain is a weighted 0-quadrature domain. Since the weight f is positive on ∂D , free boundary regularity results for weighted 0-quadrature domains apply also to k -quadrature domains. In particular, such a domain has locally either smooth boundary or its complement is thin in the sense of minimal diameter (see [39, page 109]). We also remark that when $k = 0$ the Eq. 1.2 is related to harmonic continuation of potentials, see [31] for further information.

Theorem 1.2 has an immediate consequence showing that domains with real-analytic boundary are k -quadrature domains.

Corollary 1.4 *If $k > 0$, then any bounded open set $D \subset \mathbb{R}^n$ with real-analytic boundary is a k -quadrature domain.*

Proof Since ∂D is real-analytic, we can use the Cauchy–Kowalevski theorem to find a real-analytic function u near ∂D satisfying

$$\begin{cases} (\Delta + k^2)u = 1 \text{ near } \partial D, \\ u|_{\partial D} = \partial_\nu u|_{\partial D} = 0, \end{cases}$$

where ∂_ν denotes the derivative in the normal direction to ∂D . We redefine u to be zero outside D . One can directly check that u and ∇u are Lipschitz continuous across ∂D . Hence u will be $C^{1,1}$ near ∂D and will satisfy the condition in Theorem 1.2. This proves that D is a k -quadrature domain. □

The next result gives further examples of k -quadrature domains in two dimensions.

Theorem 1.5 *Let $k > 0$, and let \mathbb{D} be the unit disc in $\mathbb{R}^2 \cong \mathbb{C}$. Suppose that $D = \varphi(\mathbb{D})$ where φ is a complex analytic function in a neighbourhood of $\overline{\mathbb{D}}$ such that $\varphi : \mathbb{D} \rightarrow D$ is bijective. Then D is a k -quadrature domain.*

Domains D as in Theorem 1.5 include cardioid type domains and domains with double points. Examples and further properties of these domains are given in the end of Section 3.

We also study k -quadrature domains from the potential theoretic point of view. More precisely, we construct some k -quadrature domains by using *partial balayage*, that is, given a

non-negative compactly supported Radon measure μ , we construct a measure ν by distributing the mass of μ more uniformly. By investigating the structure of ν , we then construct a k -quadrature domain D with respect to μ . For the case when $k = 0$ this procedure is classical, see e.g. [26–28, 42]. In this paper, we give similar results for $k > 0$ and many of our results and proofs follow those in the case of $k = 0$ as presented in [26, 27]. In this direction, our main goal is to prove the following theorem:

Theorem 1.6 (see also Theorem 7.1) *Let μ be a positive measure supported in a ball of radius $\epsilon > 0$. There exists a constant $c_n > 0$ depending only on the dimension such that if*

$$0 < k < \frac{c_n}{\mu(\mathbb{R}^n)^{1/n}} \quad \text{and} \quad \epsilon < c_n \mu(\mathbb{R}^n)^{1/n}, \tag{1.5}$$

then there exists an open connected set D with real-analytic boundary which is a k -quadrature domain for μ . Moreover, for each $w \in L^1(D) \cap L^1(\mu)$ satisfying $(\Delta + k^2)w \geq 0$ in D we have

$$\int_D w(x) dx \geq \int w(x) d\mu(x). \tag{1.6}$$

Remark 1.7 The assumption $w \in L^1(\mu)$ is in order to ensure that the right-hand side of Eq. 1.6 is well defined.

Finally we consider the relation of k -quadrature domains to the inverse problem of determining the shape of a penetrable obstacle from a single measurement, as discussed in [45]. See [14, 17, 52] for more details about scattering problems. Let $D \subset \mathbb{R}^n$ be a bounded open set, and let $h \in L^\infty(D)$ satisfy $|h| \geq c > 0$ a.e. near ∂D (such a function h is called a *contrast* for D). The pair (D, h) describes a penetrable obstacle D with contrast h .

We now probe the penetrable obstacle (D, h) by some incident field u_0 at frequency $k > 0$. The incident field is a solution of

$$(\Delta + k^2)u_0 = 0 \quad \text{in } \mathbb{R}^n.$$

Let u_{sc} be the corresponding scattered field. That is, the unique function u_{sc} so that the total field $u_{tot} = u_0 + u_{sc}$ satisfies

$$\begin{cases} (\Delta + k^2 + h\chi_D)u_{tot} = 0 & \text{in } \mathbb{R}^n, \\ u_{sc} \text{ satisfies the Sommerfeld radiation condition} & \text{at } |x| \rightarrow \infty. \end{cases} \tag{1.7}$$

Here we recall that a solution u of $(\Delta + k^2)u = 0$ in $\mathbb{R}^n \setminus \overline{B_R}$ (for some $R > 0$) satisfies the Sommerfeld radiation condition if

$$\lim_{|x| \rightarrow \infty} |x|^{\frac{n-1}{2}} (\partial_r u - iku) = 0, \quad \text{uniformly in all directions } \hat{x} = \frac{x}{|x|} \in S^{n-1},$$

where ∂_r denotes the radial derivative. Solutions satisfying the Sommerfeld radiation condition are also called outgoing. The functions u_0, u_{sc} and u_{tot} are allowed to be complex.

The single measurement inverse problem is to determine some properties of the obstacle D from knowledge of the scattered wave $u_{sc}(x)$ when $|x|$ is large. If $D = \emptyset$, then $u_{sc} \equiv 0$, and a related question is to ask whether some nontrivial domain D admits some h and u_0 so that $u_{sc} = 0$ for large x . Such a penetrable obstacle (D, h) would be invisible when probed

by the incident wave u_0 and would look like empty space. Domains D having this property for some h and u_0 will be called non-scattering domains.

Definition 1.8 We say that a bounded open set $D \subset \mathbb{R}^n$ is a *non-scattering domain* if there is some $h \in L^\infty(D)$ with $|h| \geq c > 0$ a.e. near ∂D and some solution u_0 of $(\Delta + k^2)u_0 = 0$ in \mathbb{R}^n such that the corresponding scattered wave u_{sc} satisfies $u_{sc}|_{\mathbb{R}^n \setminus \overline{B}_R} = 0$ for some $R > 0$.

The following result states that k -quadrature domains are also non-scattering domains, at least if there is some incident wave u_0 that is positive on ∂D . By the results in [45] such an incident wave u_0 exists at least when

- D is a C^1 domain (Lipschitz if $n = 2, 3$) so that $\mathbb{R}^n \setminus \overline{D}$ is connected and k^2 is not a Dirichlet eigenvalue of $-\Delta$ in D ; or
- D is contained in a ball of radius $< k^{-1} j_{\frac{n-2}{2}, 1}$ where $j_{\frac{n-2}{2}, 1}$ is the first positive zero of the Bessel function $J_{\frac{n-2}{2}}$.

By combining Theorem 1.2 and [45, Remark 2.4] we deduce the following corollary.

Corollary 1.9 *Let $D \subset \mathbb{R}^n$ be a k -quadrature domain, and assume that there exists u_0 solving $(\Delta + k^2)u_0 = 0$ in \mathbb{R}^n with $u_0|_{\partial D} > 0$. Then D is a non-scattering domain (for the incident wave u_0 and for some contrast h).*

From Theorem 1.5 and Corollary 1.9 we see that there exist non-scattering domains with inward cusps for any $k > 0$, extending the corresponding result for $k = 0$ in [45]. In contrast, domains having suitable corner points cannot be non-scattering domains for any $k > 0$, i.e. “corners always scatter”. This line of research was initiated in [7] and various further results were obtained in [4–6, 15, 16, 21, 38].

1.4 Organization

We prove Theorems 1.2 and 1.5 in §2, respectively §3. In §4, we introduce an obstacle problem, and define the partial balayage in terms of the maximizer of such an obstacle problem. We then study the structure of partial balayage in §5 and §6. Using these properties, we prove Theorem 1.6 in §7. Finally, we provide some details about a real-valued fundamental solution relevant to our construction, some results related to maximum principles, and the mean value theorem (MVT) in Appendix A.

2 PDE Characterization of Quadrature Domains

In this section we will prove Theorem 1.2 from the introduction. We begin with a global PDE characterization of k -quadrature domains.

Proposition 2.1 *Let $k > 0$, and let $D \subset \mathbb{R}^n$ be a bounded open set. Then D is a k -quadrature domain corresponding to $\mu \in \mathcal{E}'(D)$ if and only if there is a distribution $u \in \mathcal{D}'(\mathbb{R}^n)$ satisfying*

$$\begin{cases} (\Delta + k^2)u = \chi_D - \mu & \text{in } \mathbb{R}^n, \\ u = |\nabla u| = 0 & \text{in } \mathbb{R}^n \setminus D. \end{cases} \tag{2.1}$$

Note that even though u is only assumed to be in $\mathcal{D}'(\mathbb{R}^n)$, the equation $(\Delta + k^2)u = \chi_D$ near ∂D and elliptic regularity imply that u is C^1 near ∂D and hence the condition that $u = |\nabla u| = 0$ in $\mathbb{R}^n \setminus D$ is meaningful.

Example 2.2 (When μ is a Dirac mass.) For the case when $D = B_R$ with $R > 0$, and the measure is a constant multiple of the Dirac mass, we can find an explicit solution $u = u_{k,R}$ of Eq. 2.1 in terms of Bessel functions. The general radially symmetric solution of $(\Delta + k^2)u = 1$ in $\mathbb{R}^n \setminus \{0\}$ is

$$u_k(x) = \frac{1}{k^2} + c_1|x|^{1-\frac{n}{2}}J_{\frac{n}{2}-1}(k|x|) + c_2|x|^{1-\frac{n}{2}}Y_{\frac{n}{2}-1}(k|x|).$$

Given a radius $R > 0$ there is a unique choice of the constants c_1, c_2 so that $u_{k,R} := u_k \chi_{B_R} \in C^{1,1}(\mathbb{R}^n \setminus \{0\})$, namely

$$c_1 = \frac{\pi R^{\frac{n}{2}} Y_{\frac{n}{2}}(kR)}{2k} \quad \text{and} \quad c_2 = -\frac{\pi R^{\frac{n}{2}} J_{\frac{n}{2}}(kR)}{2k}.$$

With these choices of coefficients $u_{k,R}$ satisfies

$$(\Delta + k^2)u_{k,R} = \chi_{B_R} - k^{-\frac{n}{2}}(2\pi R)^{\frac{n}{2}}J_{\frac{n}{2}}(kR)\delta \quad \text{in } \mathbb{R}^n, \\ u_{k,R}|_{\mathbb{R}^n \setminus \overline{B_R}} = 0,$$

which gives an example of Proposition 2.1 with $\mu = k^{-\frac{n}{2}}(2\pi R)^{\frac{n}{2}}J_{\frac{n}{2}}(kR)\delta$.

Example 2.3 (When $\mu \equiv 0$.) Let D be a bounded domain in \mathbb{R}^n such that ∂D is homeomorphic to a sphere. The well-known Pompeiu problem [40, 50, 53] asks whether the existence of a nonzero continuous function on \mathbb{R}^n whose integral vanishes on all congruent copies of D implies that D is a ball.

The problem can be reformulated in terms of free boundary problems, or in the context of this paper, in terms of null k -quadrature domains (i.e., $\mu \equiv 0$). Indeed the assumption in Pompeiu problem is equivalent to the existence of a function v solving the free boundary problem

$$\Delta v + \lambda v = \chi_D \text{ in } \mathbb{R}^n, \quad v = 0 \text{ outside } D, \tag{2.2}$$

for some $\lambda > 0$, see [51, Theorem 1] and [50]. If the bounded open set D satisfies the assumptions in the Pompeiu problem and its boundary ∂D is additionally Lipschitz regular, then ∂D is analytic [51]. See also [9, 10, 20] for some related results. The so-far unanswered question is: whether D has to be a ball?

The fact that balls (with appropriate radii depending on $k > 0$) solve this problem is evident from the following simple procedure: take the function $u(x) = |x|^{\frac{2-n}{2}}J_{\frac{n-2}{2}}(k|x|)$ that solves $\Delta u + k^2u = 0$ in \mathbb{R}^n , add a constant to u so that one of the local minima (say $|x| = R$) of u reaches the level zero, and then redefine the function to be zero outside B_R . After multiplying by a suitable constant, this function obviously solves the free boundary formulation of the Pompeiu problem.

An interesting observation is that the solution to the free boundary formulation of the Pompeiu problem thus constructed may change sign. The construction leads to a non-negative solution only if we choose R to be the smallest radius for which u takes a minimum.

The above discussion also gives an indication of the failure of the application of the classical moving-plane technique for this problem.

We will require the following Runge approximation type result, see e.g. [1, Chapter 11] for related results. We will follow the argument in [43, Lemma 5.1].

Proposition 2.4 *Let $k \geq 0$, and let $D \subset \mathbb{R}^n$ be a bounded open set. Let Ψ_k be any fundamental solution of $-(\Delta + k^2)$ and let $\Omega \supset \overline{D}$ be any open set in \mathbb{R}^n . Then the linear span of*

$$F = \{\partial^\alpha \Psi_k(z - \cdot)|_D : z \in \Omega \setminus D, |\alpha| \leq 1\}$$

is dense in

$$H_k L^1(D) = \{w \in L^1(D) : (\Delta + k^2)w = 0 \text{ in } D\}$$

with respect to the $L^1(D)$ topology.

Remark 2.5 If the domain D has sufficiently regular boundary it suffices to take $\alpha = 0$ in F . However, for domains like the slit disk one needs to consider also $\partial^\alpha \Psi_k(z - \cdot)|_D$ for all $|\alpha| = 1$ and $z \in \Omega \setminus D$ below (note that these functions are all in $L^1(D)$). We shall later require a version of this result for sub-solutions (see Proposition 7.5).

Proof of Proposition 2.4 By the Hahn–Banach theorem, it is enough to show that any bounded linear functional ℓ on $L^1(D)$ that satisfies $\ell|_F = 0$ also satisfies $\ell|_{H_k L^1(D)} = 0$. Since the dual of $L^1(D)$ is $L^\infty(D)$, there is a function $f \in L^\infty(D)$ with

$$\ell(w) = \int_D f w \, dx, \quad w \in L^1(D).$$

We extend f by zero to \mathbb{R}^n and consider the function

$$u(z) = -(\Psi_k * f)(z) \quad \text{for all } z \in \Omega.$$

By the assumption $\ell|_F = 0$, the function u satisfies

$$\begin{cases} (\Delta + k^2)u = f & \text{in } \Omega, \\ u = |\nabla u| = 0 & \text{in } \Omega \setminus D. \end{cases}$$

We now consider the zero extension of u , still denoted by u , which satisfies

$$\begin{cases} (\Delta + k^2)u = f & \text{in } \mathbb{R}^n, \\ u = |\nabla u| = 0 & \text{in } \mathbb{R}^n \setminus D. \end{cases}$$

Note that since $f \in L^\infty$, we have $u \in C^{1,\alpha}$ for any $\alpha < 1$. In order to show that $\ell|_{H_k L^1(D)} = 0$, we take some $w \in H_k L^1(D)$ and compute

$$\ell(w) = \int_D f w \, dx = \int_D ((\Delta + k^2)u)w \, dx.$$

We claim that one can integrate by parts and use the condition $(\Delta + k^2)w = 0$ to conclude that

$$\int_D ((\Delta + k^2)u)w \, dx = 0. \tag{2.3}$$

This implies that $\ell|_{H_k L^1(D)} = 0$ and proves the result. However, the proof of Eq. 2.3 is somewhat delicate due to the failure of Calderón–Zygmund estimates when $p = \infty$. In the case $k = 0$, Eq. 2.3 follows from [43, Lemma 5.1]. We will verify that the same argument works for $k > 0$.

First observe that u solves

$$\Delta u = f - k^2 u \text{ in } \mathbb{R}^n.$$

Since f and u are L^∞ , it follows from [25, Theorem 3.9] that ∇u satisfies

$$|\nabla u(x) - \nabla u(y)| \leq C|x - y| \log(1/|x - y|), \quad x, y \in \overline{D}, |x - y| < e^{-2}.$$

Using the condition $u = |\nabla u| = 0$ in $\mathbb{R}^n \setminus D$, this implies that uniformly for $x \in D$ near ∂D one has

$$\begin{aligned} u(x) &= O(\delta(x)^2 \log(1/\delta(x))), \\ \nabla u(x) &= O(\delta(x) \log(1/\delta(x))), \end{aligned}$$

where $\delta(x) = \text{dist}(x, \partial D)$.

As in [43, Lemma 5.1] we introduce the sequence $(\omega_j)_{j=1}^\infty$ of Ahlfors–Bers mollifiers [2, 3] that satisfy $\omega_j \in C^\infty(\mathbb{R}^n)$, $0 \leq \omega_j \leq 1$, $\omega_j = 0$ near ∂D , $\omega_j = 1$ outside a neighborhood of ∂D , $\omega_j(x) \rightarrow 1$ for $x \notin \partial D$, and

$$|\partial^\alpha \omega_j(x)| \leq C_\alpha j^{-1} \delta(x)^{-|\alpha|} (\log 1/\delta(x))^{-1} \text{ for } x \notin \partial D,$$

see [30, Lemma 4]. We now begin the proof of Eq. 2.3. One has

$$\begin{aligned} \int_D ((\Delta + k^2)u)w \, dx &= \lim_{j \rightarrow \infty} \int_D ((\Delta + k^2)u)\omega_j w \, dx \\ &= \lim_{j \rightarrow \infty} \int_D [(\Delta + k^2)(\omega_j u) - 2\nabla \omega_j \cdot \nabla u - (\Delta \omega_j)u] w \, dx. \end{aligned}$$

Using the estimates for u and ω_j , the limits corresponding to the last two terms inside the brackets are zero. Moreover, since w is smooth near $\text{supp}(\omega_j)$, we have

$$\int_D (\Delta + k^2)(\omega_j u)w \, dx = \int_D \omega_j u (\Delta + k^2)w \, dx = 0.$$

This concludes the proof of Eq. 2.3. □

Proof of Proposition 2.1 Let Ψ_k be any fundamental solution of $-(\Delta + k^2)$, i.e. $\Psi_k \in \mathcal{D}'(\mathbb{R}^n)$ solves $-(\Delta + k^2)\Psi_k = \delta_0$ in \mathbb{R}^n . In particular, Ψ_k is smooth away from the origin. If D is a k -quadrature domain corresponding to μ , then

$$\int_D \partial^\alpha \Psi_k(z - x) \, dx = \langle \mu, \partial^\alpha \Psi_k(z - \cdot) \rangle \tag{2.4}$$

whenever $z \in \mathbb{R}^n \setminus D$ and $|\alpha| \leq 1$. Let $u = -\Psi_k * (\chi_D - \mu)$, which is well defined since $\chi_D - \mu$ is a compactly supported distribution. We see that $(\Delta + k^2)u = \chi_D - \mu$ and $u = |\nabla u| = 0$ in $\mathbb{R}^n \setminus D$ as required.

Conversely, suppose that $u \in \mathcal{D}'(\mathbb{R}^n)$ is as in the statement. We easily obtain the quadrature identity for functions w that solve $(\Delta + k^2)w = 0$ near \overline{D} , since by taking a cutoff $\psi \in C_c^\infty(\mathbb{R}^n)$ with $\psi = 1$ near \overline{D} we have

$$\int_D w \, dx - \langle w, \mu \rangle = \langle \chi_D - \mu, \psi w \rangle = \langle -(\Delta + k^2)u, \psi w \rangle = \langle u, -(\Delta + k^2)(\psi w) \rangle = 0,$$

using that the derivatives of ψ vanish near $\text{supp}(u)$.

For general solutions $w \in L^1(D)$ we need another argument. Since u is compactly supported, by the properties of convolution for distributions we have

$$u = \delta_0 * u = -(\Delta + k^2)\Psi_k * u = -\Psi_k * (\Delta + k^2)u = -\Psi_k * (\chi_D - \mu).$$

Using that $u = |\nabla u| = 0$ in $\mathbb{R}^n \setminus D$, we have

$$\int_D \partial^\alpha \Psi_k(z - x) \, dx = \langle \mu, \partial^\alpha \Psi_k(z - \cdot) \rangle \tag{2.5}$$

for all $z \in \mathbb{R}^n \setminus D$ and $|\alpha| \leq 1$. Now let $w \in L^1(D)$ solve $(\Delta + k^2)w = 0$ in D , and use Proposition 2.4 to find a sequence $w_j \in \text{span}\{\partial^\alpha \Psi_k(z - \cdot)|_D : z \in \mathbb{R}^n \setminus D, |\alpha| \leq 1\}$ with $w_j \rightarrow w \in L^1(D)$. In particular, for any $j \geq 1$ we have

$$\int_D w_j \, dx = \langle \mu, w_j \rangle. \tag{2.6}$$

Since $\mu \in \mathcal{E}'(D)$, there is a compact set $K \subset D$ and an integer $m \geq 0$ such that

$$|\langle \mu, \varphi \rangle| \leq C \|\varphi\|_{C^m(K)}, \quad \varphi \in C^\infty(D).$$

By elliptic regularity and Sobolev embedding, any $v \in L^1(D)$ with $(\Delta + k^2)v \in H^{s-2}(D)$ satisfies $v \in C^m(K)$ when $s > m + n/2$. By the closed graph theorem this yields the estimate

$$\|v\|_{C^m(K)} \leq C(\|v\|_{L^1(D)} + \|(\Delta + k^2)v\|_{H^{s-2}(D)}).$$

Applying this estimate to $v = w_j - w$ gives

$$\|w_j - w\|_{C^m(K)} \leq C\|w_j - w\|_{L^1(D)}. \tag{2.7}$$

Thus we may take limits as $j \rightarrow \infty$ in Eq. 2.6 and obtain that

$$\int_D w \, dx = \langle \mu, w \rangle.$$

This shows that D is a k -quadrature domain for μ . □

Proof of Theorem 1.2 If D is a k -quadrature domain corresponding to μ , then taking a neighborhood U of ∂D that is disjoint from $\text{supp}(\mu)$ and restricting the distribution from Proposition 2.1 to U gives the required $u \in \mathcal{D}'(U)$ satisfying Eq. 1.3.

Conversely, assume that $u \in \mathcal{D}'(U)$ satisfies Eq. 1.3. Let $\psi \in C_c^\infty(U)$ satisfy $0 \leq \psi \leq 1$ and $\psi = 1$ near ∂D , and define $\tilde{u} = \psi u \in \mathcal{D}'(\mathbb{R}^n)$. Also define

$$\tilde{\mu} := \chi_D - (\Delta + k^2)\tilde{u}. \tag{2.8}$$

Then \tilde{u} satisfies

$$\begin{cases} (\Delta + k^2)\tilde{u} = \chi_D - \tilde{\mu} & \text{in } \mathbb{R}^n, \\ \tilde{u} = |\nabla \tilde{u}| = 0 & \text{in } \mathbb{R}^n \setminus D. \end{cases}$$

Moreover, $\tilde{\mu} \in \mathcal{D}'(\mathbb{R}^n)$ satisfies $\text{supp}(\tilde{\mu}) \subset D$ by the assumption on u . Then D is a k -quadrature domain by Proposition 2.1. By elliptic regularity u is smooth in $U \cap D$, and thus $\tilde{\mu}$ coincides with a smooth function in D . Since one also has $\text{supp}(\tilde{\mu}) \subset D$, it follows that $\tilde{\mu}$ has a smooth density with respect to Lebesgue measure. □

3 Quadrature Domains with Cusps

This section contains the proof of Theorem 1.5. The proof will employ the following simple fact regarding the vanishing order of solutions. In this section all functions are allowed to be complex valued.

Lemma 3.1 *Let v be a C^∞ function near some $x_0 \in \mathbb{R}^n$ satisfying*

$$\begin{cases} \Delta v = O(|x - x_0|^m) \text{ near } x_0, \\ v|_S = \partial_\nu v|_S = 0, \end{cases}$$

where $m \geq 0$ is an integer, S is a smooth hypersurface through x_0 , and ∂_ν denotes the derivative in the normal direction to S . Then one has $v = O(|x - x_0|^{m+2})$, and more precisely

$$v = \sum_{|\alpha|=m+2} v_\alpha(x)(x - x_0)^\alpha$$

where v_α are smooth near x_0 .

Proof After a rigid motion, we may assume that $x_0 = 0$ and the normal of S satisfies $\nu(0) = e_n$. We use the Taylor formula and write v as

$$v = \sum_{j=0}^{m+1} P_j + R, \quad R = \sum_{|\alpha|=m+2} v_\alpha(x)(x - x_0)^\alpha,$$

where each P_j is a homogeneous polynomial of degree j and each v_α is smooth. Using the assumption $v|_S = \partial_\nu v|_S = 0$ we have $P_0 = P_1 = 0$. Moreover, the assumption for Δv implies that

$$\sum_{j=2}^{m+1} \Delta P_j = O(|x|^m).$$

Since the left hand side is a polynomial of degree $m - 1$, it follows that we must have $\Delta P_j = 0$ for $2 \leq j \leq m + 1$.

Suppose that γ is a smooth curve on S with $\gamma(0) = 0$ and $\dot{\gamma}(0) = \omega$ where $\omega \perp e_n$ and $|\omega| = 1$. Since $v|_S = \partial_\nu v|_S = 0$, we have

$$0 = v(\gamma(t)) = \sum_{j=2}^{m+1} |\gamma(t)|^j P_j(\gamma(t)/|\gamma(t)|) + O(|\gamma(t)|^{m+2}), \tag{3.1}$$

$$0 = \partial_\nu v(\gamma(t)) = \sum_{j=2}^{m+1} |\gamma(t)|^{j-1} \nu(\gamma(t)) \cdot \nabla P_j(\gamma(t)/|\gamma(t)|) + O(|\gamma(t)|^{m+1}). \tag{3.2}$$

Since $\gamma(t) = t\omega + O(t^2)$, we have $\gamma(t)/|\gamma(t)| \rightarrow \omega$ as $t \rightarrow 0$. If one would have $P_2(\omega) \neq 0$, then multiplying Eq. 3.1 by t^{-2} would lead to a contradiction as $t \rightarrow 0$. Similarly $\partial_n P_2(\omega) \neq 0$ would lead to a contradiction with Eq. 3.2. Thus $P_2(\omega) = \partial_n P_2(\omega) = 0$. Varying ω implies that

$$P_2|_{x_n=0} = \partial_n P_2|_{x_n=0} = 0.$$

But since $\Delta P_2 = 0$, unique continuation implies that $P_2 \equiv 0$. Iterating this argument shows that $P_2 \equiv \dots \equiv P_{m+1} = 0$ as required. \square

We are now ready to prove Theorem 1.5. As we shall illustrate in Examples 3.3–3.5 the domain D may have inward cusps and the map φ is not necessarily injective on $\partial\mathbb{D}$, which introduces some technicalities in the argument.

Proof of Theorem 1.5 Let $D = \varphi(\mathbb{D})$ where φ is analytic near $\overline{\mathbb{D}}$ and injective in \mathbb{D} . Note that D is an open set by the open mapping theorem for analytic functions [41, Theorem 10.32]. We claim:

$$\text{if } z_j \in \mathbb{D} \text{ and } d(\varphi(z_j), \partial D) \rightarrow 0, \text{ then } d(z_j, \partial\mathbb{D}) \rightarrow 0. \tag{3.3}$$

¹Lemma 3.1 can also be proved by a simple blow-up argument. Starting with a quadratic blow-up one obtains P_2 in the limit and S becomes a hyperplane $\{x_n = 0\}$, along with the zero Cauchy-data for P_2 . This implies $P_2 \equiv 0$. Repeating this argument, by a cubic scaling we obtain $P_3 \equiv 0$. Iterating this argument we have $P_j \equiv 0$ for all $j \leq m + 1$.

To see Eq. 3.3, we argue by contradiction and assume that $d(\varphi(z_j), \partial D) \rightarrow 0$ but there is $\epsilon > 0$ and a subsequence (z_{j_k}) with $d(z_{j_k}, \partial \mathbb{D}) \geq \epsilon$. After passing to another subsequence also denoted by (z_{j_k}) , we have $z_{j_k} \rightarrow z \in \mathbb{D}$ with $d(z, \partial \mathbb{D}) \geq \epsilon$. However, since $d(\varphi(z_j), \partial D) \rightarrow 0$ we must have $d(\varphi(z), \partial D) = 0$. This contradicts the fact that $\varphi(\mathbb{D}) = D$, proving Eq. 3.3.

Next we show that

$$\varphi(\partial \mathbb{D}) = \partial D. \tag{3.4}$$

We begin the proof of Eq. 3.4 by taking $z \in \partial \mathbb{D}$ and showing that $\varphi(z) \in \partial D$. By continuity $\varphi(z) \in \overline{D}$. If one had $\varphi(z) \in D$, then since φ is bijective $\mathbb{D} \rightarrow D$ there would be some $z' \in \mathbb{D}$ with $\varphi(z') = \varphi(z)$. For any $\epsilon < |z - z'|/2$ we consider the open sets $\varphi(B_\epsilon(z'))$ and $\varphi(B_\epsilon(z) \cap \mathbb{D})$. The point $\varphi(z)$ is contained in the interior of the first set and in the closure of the second, in particular the two sets are not disjoint. This contradicts the assumption that φ is injective, and thus proves that $\varphi(\partial \mathbb{D}) \subset \partial D$. For the converse inclusion, if $x \in \partial D$ and $x_j \rightarrow x$ where $x_j \in D$, then $x_j = \varphi(z_j)$ for some $z_j \in \mathbb{D}$. After passing to a subsequence we may assume that $z_{j_k} \rightarrow z \in \overline{\mathbb{D}}$, and by Eq. 3.3 one must have $z \in \partial \mathbb{D}$. Thus $x = \lim \varphi(z_j) = \varphi(z)$, proving Eq. 3.4.

By Theorem 1.2, the result will follow if we can find a distribution u near ∂D solving

$$\begin{cases} (\Delta + k^2)u = \chi_D & \text{near } \partial D, \\ u = |\nabla u| = 0 & \text{outside } D. \end{cases} \tag{3.5}$$

By the chain rule the equation $(\Delta + k^2)u = 1$ in some set $\varphi(U_1)$, with $U_1 \subset \mathbb{D}$ open, is equivalent to the equation

$$(\Delta + k^2|\varphi'|^2)(u \circ \varphi) = |\varphi'|^2 \text{ in } U_1.$$

Since $|\varphi'|^2 = \partial\bar{\varphi}\bar{\partial}\varphi$ is real-analytic near $\partial \mathbb{D}$, the Cauchy–Kowalevski theorem implies that there exists a neighborhood U of $\partial \mathbb{D}$ and a function \hat{u} which is real-analytic in U such that

$$\begin{cases} (\Delta + k^2|\varphi'|^2)\hat{u} = |\varphi'|^2 & \text{in } U, \\ \hat{u} = \partial_v \hat{u} = 0 & \text{on } \partial \mathbb{D}. \end{cases} \tag{3.6}$$

By Eq. 3.4 and the open mapping theorem we know that $V = \varphi(U)$ is an open neighborhood of ∂D . We define

$$u(x) := \begin{cases} \hat{u}(\varphi^{-1}(x)) & \text{for all } x \in V \cap D, \\ 0 & \text{for all } x \in V \setminus D. \end{cases}$$

The function u is defined piecewise and it satisfies Eq. 3.5 away from ∂D . If we can prove that $u \in C^{1,1}(V)$, then u will satisfy Eq. 3.5 also near ∂D and the proof of the theorem will be concluded. Note that by the inverse function theorem, u is smooth in $V \cap D$. We would like to show that u is continuous up to ∂D . If $x \in \partial D$ and $x_j \in V \cap D$ satisfy $x_j \rightarrow x$, then $x_j = \varphi(z_j)$ for some $z_j \in \mathbb{D}$. Then $d(\varphi(z_j), \partial D) \rightarrow 0$, and Eq. 3.3 ensures that $d(z_j, \partial \mathbb{D}) \rightarrow 0$. It follows that

$$u(x_j) = \hat{u}(z_j) \rightarrow 0$$

since \hat{u} is Lipschitz near $\partial \mathbb{D}$ and $\hat{u}|_{\partial \mathbb{D}} = 0$. This shows that $u \in C^0(V)$.

Next we show that u is C^1 up to ∂D . Let $x \in \partial D$ and $x_j \in D$ with $x_j \rightarrow x$. It is enough to show that for any $\epsilon > 0$ there is j_0 such that $|\nabla u(x_j)| \leq \epsilon$ for $j \geq j_0$. Now $x_j = \varphi(z_j)$ where $z_j \in \mathbb{D}$, and by the chain rule one has

$$\partial u(\varphi(z)) = \frac{\partial \hat{u}(z)}{\varphi'(z)}, \quad \bar{\partial} u(\varphi(z)) = \frac{\bar{\partial} \hat{u}(z)}{\varphi'(z)}.$$

Thus

$$|\nabla u(\varphi(z))| = \frac{|\nabla \hat{u}(z)|}{|\varphi'(z)|}. \tag{3.7}$$

Using Eq. 3.3 we know that $d(z_j, \partial\mathbb{D}) \rightarrow 0$, and thus $\nabla \hat{u}(z_j) \rightarrow 0$ since $\nabla \hat{u}|_{\partial\mathbb{D}} = 0$. However, $\varphi'(z_j)$ may also converge to zero and this requires some care. We start by observing that there are only finitely many points $z_0 \in \partial\mathbb{D}$ with $\varphi'(z_0) = 0$, and near any such z_0 one can write

$$\varphi(z) = \varphi(z_0) + (z - z_0)^2 g(z),$$

for some analytic function g . Since $\varphi: \mathbb{D} \rightarrow D$ is bijective it follows that $\varphi''(z_0) \neq 0$ and hence $g(z_0) \neq 0$ (see Remark 3.2). Thus $|\varphi'(z)|^2 = O(|z - z_0|^2)$. Using Lemma 3.1 we know that

$$\partial^\alpha \hat{u}(z) = O(|z - z_0|^{4-|\alpha|}) \text{ for } |\alpha| \leq 4 \text{ and for all } z \text{ near } z_0. \tag{3.8}$$

By Eq. 3.7, for $z \in \mathbb{D}$ near z_0 we have

$$|\nabla u(\varphi(z))| \leq C \frac{|z - z_0|^3}{|z - z_0|} \leq C|z - z_0|^2.$$

Thus there is $\delta > 0$ such that

$$|\nabla u(\varphi(z))| \leq \epsilon \text{ when } z \in W := \bigcup_{z_0 \in \partial\mathbb{D}, \varphi'(z_0)=0} (B(z_0, \delta) \cap \mathbb{D}).$$

We have $\partial\mathbb{D} \subset W \cup W'$ where W' is some open set with $|\varphi'(z)| \geq c > 0$ for $z \in W'$. We already know that $|\nabla u(\varphi(z_j))| \leq \epsilon$ when $z_j \in W$, and for $z_j \in W'$ the expression Eq. 3.7 gives that

$$|\nabla u(\varphi(z_j))| \leq \frac{1}{c} |\nabla \hat{u}(z_j)|$$

which becomes $\leq \epsilon$ when $j \geq j_0$ for some sufficiently large j_0 by Eq. 3.3. This concludes the proof that $u \in C^1(V)$.

Finally, we use the chain rule again and observe that for $z \in \mathbb{D}$ one has

$$|\nabla^2 u(\varphi(z))| \leq C \left(\frac{|\nabla^2 \hat{u}(z)|}{|\varphi'(z)|^2} + \frac{|\nabla \hat{u}(z)| |\varphi''(z)|}{|\varphi'(z)|^3} \right).$$

As before, the worst case is when z is close to some $z_0 \in \partial\mathbb{D}$ with $\varphi'(z_0) = 0$. By Eq. 3.8, for z near z_0 one has

$$|\nabla^2 u(\varphi(z))| \leq C \left(\frac{|\nabla^2 \hat{u}(z)|}{|z - z_0|^2} + \frac{|\nabla \hat{u}(z)|}{|z - z_0|^3} \right) \leq C.$$

It follows that ∇u is Lipschitz continuous in V . In fact it is Lipschitz in $V \cap D$ and $V \setminus D$, and if $x \in V \cap D$ and $y \in V \setminus D$ we let y_1 be a closest point to x in $\mathbb{R}^n \setminus D$ (so that $y_1 \in \partial D$) and observe that

$$|\nabla u(x) - \nabla u(y)| = |\nabla u(x) - \nabla u(y_1)| \leq C|x - y_1| \leq C|x - y|.$$

This proves that $u \in C^{1,1}(V)$, and therefore concludes the proof of Theorem 1.5. □

Remark 3.2 Let $D = \varphi(\mathbb{D})$ where φ is an analytic function near $\overline{\mathbb{D}}$ which is injective in \mathbb{D} . In this remark we clarify what ∂D looks like. Recall from Eq. 3.4 that $\varphi(\partial\mathbb{D}) = \partial D$. We may divide the boundary points in three categories.

- (i) (Smooth points) If $x_0 \in \partial D$ is of the form $x_0 = \varphi(z_0)$ for a unique $z_0 \in \partial\mathbb{D}$ and $\varphi'(z_0) \neq 0$, then by the inverse function theorem D near x_0 is given by the region above the graph of a real-analytic function.

- (ii) (Inward cusp points) If $x_0 \in \partial D$ is of the form $x_0 = \varphi(z_0)$ for some $z_0 \in \partial \mathbb{D}$ with $\varphi'(z_0) = 0$, then $\varphi''(z_0) \neq 0$ since if φ vanished to higher order the bijectivity in \mathbb{D} would fail in the same way that it does for $z \mapsto z^m, m > 2$, around $z = 0$ (the image of an arbitrary half-plane covers $\mathbb{C} \setminus \{0\}$ more than once). Thus φ behaves near z_0 like $z \mapsto z^2$ which produces an inward cusp.
- (iii) (Double points) If $x_0 \in \partial D$ satisfies $x_0 = \varphi(z_1) = \varphi(z_2)$ for two distinct $z_1, z_2 \in \partial \mathbb{D}$, then by the bijectivity $\varphi'(z_1) \neq 0$ and $\varphi'(z_2) \neq 0$ and there exists an $r > 0$ small enough so that $\partial D \cap B_r(x_0)$ is the union of two analytic arcs whose intersection is $\{x_0\}$ where the arcs touch (by injectivity they do not cross).

Moreover, there are only finitely many points which fail to be in category (i).

This classification of the points on the boundary of D is rather classical. Remark 3.2 is also related to Sakai’s regularity theorem, see [44, Theorem 5.2] as well as [37, Section 3.2].

Example 3.3 (Fig. 1) Let $\varphi(z) = z + \frac{1}{2}z^2$ and $D = \varphi(\mathbb{D})$. Then D is a cardioid whose boundary is smooth except at the point $\varphi(-1) = -1/2$ where it has an inward cusp. It is clear that φ satisfies the conditions of Theorem 1.5. Similarly, if $\varphi(z) = z + \frac{1}{m}z^m$ for integer $m \geq 2$ then D has $m - 1$ inward cusps.

Example 3.4 (Fig. 2) Let $\varphi(z) = z - \frac{2\sqrt{2}}{3}z^2 + \frac{1}{3}z^3$ and $D = \varphi(\mathbb{D})$ (see e.g. [37, equation (1.9)]). Then the corresponding domain D is not a Jordan domain and furthermore its boundary has inward cusps. By Theorem 1.5, the domain D is a k -quadrature domain.

Example 3.5 (Fig. 3) Let $\varphi(z) = (z - 1)^2 - (1 - \frac{i}{2})(z - 1)^3$ and $D = \varphi(\mathbb{D})$. The domain D looks similar to a cardioid, but with an inward cusp which is curved in such a manner that the ∂D cannot locally be represented as the graph of a function. It is also a k -quadrature domain by Theorem 1.5.




Fig. 1 Plot of Example 3.3 (GNU Octave)




Fig. 2 Plot of Example 3.4 (GNU Octave)

4 Partial Balayage Via an Obstacle Problem

In this section we define the partial balayage measure $Bal(\mu)$ with respect to $\Delta + k^2$ for certain sufficiently concentrated measures μ when $k > 0$ is small. For simplicity we will assume that μ is concentrated near the origin, but by translation invariance any other point would do. There are several ways of defining partial balayage, and we will proceed via an obstacle problem (see e.g. [27, Definition 3.2] for $k = 0$).

Let $R(n, k) = \frac{1}{2} j_{\frac{n-2}{2}, 1} k^{-1}$ and let $\tilde{\Phi}_k = \tilde{\Phi}_{k, R(n, k)}$ be the fundamental solution given in Proposition A.1. Here and in what follows we write $U_k^\mu := \tilde{\Phi}_k * \mu$ for any Radon measure μ . In the special case where $\mu = \chi_\Omega m$ for some open set Ω we simply write $U_k^\Omega := \tilde{\Phi}_k * \chi_\Omega$.




Fig. 3 Plot of Example 3.5 (GNU Octave)

We now restrict ourselves to measures μ having a bounded density with respect to Lebesgue measure. Slightly abusing notation we write $\mu \in L^\infty(\Omega)$ to mean that μ has the form

$$\mu = f m,$$

where $f \in L^\infty(\mathbb{R}^n)$ satisfies $f = 0$ outside Ω . Under this assumption, by elliptic regularity (see e.g. [25, Theorem 9.11])

$$U_k^\mu \in \bigcap_{1 < p < \infty} W_{\text{loc}}^{2,p}(\mathbb{R}^n) \subset \bigcap_{0 < \alpha < 1} C^{1,\alpha}(\overline{B_{R(n,k)}}).$$

For $k > 0$ and $\mu \in L^\infty(B_{R(n,k)})$ define

$$\mathcal{F}_k(\mu) = \left\{ v \in H^1(B_{R(n,k)}) \left| \begin{array}{l} (\Delta + k^2)v \geq -1 \text{ in } B_{R(n,k)} \\ v \leq U_k^\mu \text{ in } B_{R(n,k)} \\ v = U_k^\mu \text{ on } \partial B_{R(n,k)} \end{array} \right. \right\}. \tag{4.1}$$

Lemma 4.1 *Let $0 < \gamma < R(n, k)$ and $\mu \in L^\infty(B_\gamma)$. Assume that there exists $r > 0$ such that*

$$r < R(n, k) - \gamma \quad \text{and} \quad c_{n,k,r}^{MVT} \geq \mu_+(\mathbb{R}^n), \tag{4.2}$$

where $c_{n,k,r}^{MVT}$ is the constant appearing in the mean value theorem for the Helmholtz equation (see Eq. A.3). Then $\mathcal{F}_k(\mu)$ contains an element \tilde{u}_k which equals U_k^μ in $B_{R(n,k)} \setminus B_{\gamma+r}$ (note that $\gamma + r < R(n, k)$).

Proof Let

$$\tilde{u}_k := U_k^{\mu_+} * h_r - U_k^{\mu_-} \quad \text{where} \quad h_r := \frac{1}{c_{n,k,r}^{MVT}} \chi_{B_r}. \tag{4.3}$$

Using the mean value theorem in Proposition A.2, we have

$$U_k^{\mu_+ * h_r}(x) \leq U_k^{\mu_+}(x) \text{ for all } x \in \mathbb{R}^n \text{ with equality if } \mu_+(B_r(x)) = 0,$$

which implies

$$\tilde{u}_k \leq U_k^\mu \text{ in } \mathbb{R}^n \quad \text{and} \quad \tilde{u}_k = U_k^\mu \text{ in } \mathbb{R}^n \setminus B_{\gamma+r}.$$

Finally we note that

$$(\Delta + k^2)\tilde{u}_k(x) = (-\mu_+ * h_r + \mu_-)(x) \geq -\mu_+ * h_r(x) = -\frac{\mu_+(B_r(x))}{c_{n,k,r}^{MVT}} \geq -\frac{\mu_+(\mathbb{R}^n)}{c_{n,k,r}^{MVT}} \geq -1,$$

which shows that $\tilde{u}_k \in \mathcal{F}_k(\mu)$. □

For fixed μ we now choose the parameter r in Lemma 4.1 in order to find an explicit range of $k > 0$ for which the lemma applies.

By the definition of $c_{n,k,r}^{MVT}$ the second inequality in Eq. 4.2 is equivalent to

$$k \leq \frac{(2\pi kr)^{\frac{1}{2}} J_{\frac{n}{2}}(kr)^{\frac{1}{n}}}{\mu_+(\mathbb{R})^{\frac{1}{n}}}. \tag{4.4}$$

Since $t \mapsto t^{\frac{1}{2}} J_{\frac{n}{2}}(t)^{\frac{1}{n}}$ is strictly increasing on $[0, j_{\frac{n-2}{2}, 1}]$, we see that in order to maximize the range of k we here want to choose kr as large as possible.

By the definition of $R(n, k)$ we see that the range of r we can consider is given by

$$0 < rk < \frac{1}{2} j_{\frac{n-2}{2}, 1} - \gamma k.$$

Therefore, if we assume that

$$k \leq \frac{j_{\frac{n-2}{2},1}}{4\gamma}$$

we can choose

$$rk = \frac{\frac{1}{2}j_{\frac{n-2}{2},1} - \gamma k}{2} \geq \frac{j_{\frac{n-2}{2},1}}{8}.$$

By the monotonicity of $t \mapsto t^{\frac{1}{2}} J_{\frac{n}{2}}(t)^{\frac{1}{n}}$ we then know that Eq. 4.4 is satisfied for all

$$k \leq \frac{\left(\frac{\pi j_{\frac{n-2}{2},1}}{4}\right)^{\frac{1}{2}} J_{\frac{n}{2}}\left(\frac{j_{\frac{n-2}{2},1}}{8}\right)^{\frac{1}{n}}}{\mu_+(\mathbb{R})^{\frac{1}{n}}}.$$

Consequently, for any

$$c_n \leq \min\left\{\left(\frac{\pi j_{\frac{n-2}{2},1}}{4}\right)^{\frac{1}{2}} J_{\frac{n}{2}}\left(\frac{j_{\frac{n-2}{2},1}}{8}\right)^{\frac{1}{n}}, \frac{j_{\frac{n-2}{2},1}}{4}\right\}$$

we conclude the following lemma:

Lemma 4.2 Fix any $\gamma > 0$ and $\mu \in L^\infty(B_\gamma)$. There exists a positive constant c_n (depending only on the dimension n) such that if

$$0 < k \leq c_n \min\{\gamma^{-1}, \mu_+(\mathbb{R}^n)^{-\frac{1}{n}}\} \tag{4.5}$$

then $\mathcal{F}_k(\mu)$ contains an element \tilde{u}_k with

$$\tilde{u}_k = U_k^\mu \quad \text{near } \mathbb{R}^n \setminus B_{R(n,k)}. \tag{4.6}$$

The following proposition will be used to define partial balayage in terms of the solution of our obstacle problem.

Proposition 4.3 Let μ and $k > 0$ be as in Lemma 4.2. Then there exists a largest element V_k^μ in $\mathcal{F}_k(\mu)$. In addition, the element V_k^μ satisfies

$$\langle 1 + (\Delta + k^2)V_k^\mu, V_k^\mu - U_k^\mu \rangle = 0, \tag{4.7}$$

where $\langle \cdot, \cdot \rangle$ is the $H^{-1}(B_{R(n,k)}) \times H_0^1(B_{R(n,k)})$ duality pairing.

Remark 4.4 Note that Lemma 4.2 implies that there exists $\tilde{u}_k \in \mathcal{F}_k(\mu)$ satisfying $\tilde{u}_k = U_k^\mu$ near $\partial B_{R(n,k)}$. Therefore, if V_k^μ is that largest element in $\mathcal{F}_k(\mu)$ then

$$V_k^\mu = U_k^\mu \quad \text{near } \partial B_{R(n,k)}. \tag{4.8}$$

Therefore, we can extend V_k^μ to the whole \mathbb{R}^n , by defining $V_k^\mu := U_k^\mu$ outside $B_{R(n,k)}$.

The proof of the proposition is based on variational arguments. In particular, we shall need the following elementary lemma several times in the proof.

Lemma 4.5 Fix $k > 0$ and $0 < R < j_{\frac{n-2}{2},1} k^{-1}$. Let $a : H_0^1(B_R) \times H_0^1(B_R) \rightarrow \mathbb{R}$ be the symmetric bilinear form defined by

$$a(u_1, u_2) := \int_{B_R} (\nabla u_1 \cdot \nabla u_2 - k^2 u_1 u_2) \, dm. \tag{4.9}$$

Then a is continuous, positive, and coercive.

Proof That a is a continuous is clear from the definition. To prove that the form is coercive and positive we observe that by assumption k^2 is strictly smaller than the first eigenvalue of Dirichlet Laplacian on B_R (which is exactly $j_{\frac{n-2}{2},1}^2 R^{-2}$). Therefore,

$$a(u, u) = \int_{B_R} (|\nabla u|^2 - k^2|u|^2) \, dm \geq (1 - k^2 R^2 j_{\frac{n-2}{2},1}^{-2}) \int_{B_R} |\nabla u|^2 \, dm.$$

This concludes the proof. □

Proof of Proposition 4.3 Let $\varphi \in H^1(B_{R(n,k)})$ be the unique solution to

$$\begin{cases} (\Delta + k^2)\varphi = -1 & \text{in } B_{R(n,k)}, \\ \varphi = U_k^\mu & \text{on } \partial B_{R(n,k)}, \end{cases} \tag{4.10}$$

and define

$$\tilde{\mathcal{F}}_k(\mu) = \{ w = \varphi - v \mid v \in \mathcal{F}_k(\mu) \} = \left\{ w \in H_0^1(B_{R(n,k)}) \mid \begin{cases} (\Delta + k^2)w \leq 0 & \text{in } B_{R(n,k)} \\ w \geq \varphi - U_k^\mu & \text{in } B_{R(n,k)} \end{cases} \right\}.$$

We claim that there exists a smallest element u_* of $\tilde{\mathcal{F}}_k(\mu)$. If this is the case then

$$V_k^\mu := \varphi - u_* \quad \text{in } B_{R(n,k)} \tag{4.11}$$

is the largest element of $\mathcal{F}_k(\mu)$.

To see that there exists a smallest element in $\tilde{\mathcal{F}}_k(\mu)$ we argue as follows. Let a be the bilinear form defined in Eq. 4.9 with $R = R(n, k)$. By Lemma 4.5 a is symmetric, continuous, and coercive. Define the constraint set

$$\tilde{\mathcal{K}}_k := \{ u \in H_0^1(B_{R(n,k)}) \mid u \geq \varphi - U_k^\mu \}. \tag{4.12}$$

Note that $\varphi - U_k^\mu \in H_0^1(B_{R(n,k)})$ by definition of φ , thus $\tilde{\mathcal{K}}_k$ is nonempty. Since $\tilde{\mathcal{K}}_k$ is a nonempty closed convex subset of $H_0^1(B_{R(n,k)})$, Stampacchia’s theorem [8, Theorem 5.6] implies that there exists a unique $u_* \in \tilde{\mathcal{K}}_k$ that minimizes the functional

$$u \mapsto a(u, u) \tag{4.13}$$

and $u_* \in \tilde{\mathcal{K}}_k$ can also be characterized by

$$a(u_*, u - u_*) \geq 0 \quad \text{for all } u \in \tilde{\mathcal{K}}_k. \tag{4.14}$$

Plugging in $u = u_* + \phi$ with non-negative $\phi \in C_c^\infty(B_{R(n,k)})$ into Eq. 4.14, the definition of a in Eq. 4.9 implies that

$$(\Delta + k^2)u_* \leq 0 \text{ in } B_{R(n,k)}. \tag{4.15}$$

In particular, we conclude that $u_* \in \tilde{\mathcal{F}}_k(\mu)$. Finally, by arguing as in the proof of [34, Theorem II.6.4], one can prove that $u_* \leq v$ in $B_{R(n,k)}$ for all $v \in \tilde{\mathcal{F}}_k(\mu)$. Consequently, we have found the desired smallest element in $\tilde{\mathcal{F}}_k(\mu)$.

Choosing $u = \varphi - U_k^\mu \in H_0^1(B_{R(n,k)})$ in Eq. 4.14, we have

$$\langle (\Delta + k^2)u_*, \varphi - U_k^\mu - u_* \rangle \leq 0. \tag{4.16}$$

Since $u_* \in \tilde{\mathcal{K}}_k$ we know that $u_* \geq \varphi - U_k^\mu$. Along with Eqs. 4.15 and 4.16 this inequality implies

$$\langle (\Delta + k^2)u_*, \varphi - U_k^\mu - u_* \rangle = 0. \tag{4.17}$$

Combining Eq. 4.17 with Eq. 4.11, as well as Eq. 4.10, we obtain

$$0 = \langle (\Delta + k^2)(\varphi - V_k^\mu), V_k^\mu - U_k^\mu \rangle = -\langle 1 + (\Delta + k^2)V_k^\mu, V_k^\mu - U_k^\mu \rangle,$$

which shows that V_k^μ satisfies Eq. 4.7. □

We are now ready to define partial balayage for the Helmholtz operator.

Definition 4.6 Let μ and $k > 0$ be as in Lemma 4.2. The *partial balayage* of μ is defined by

$$\text{Bal}_k(\mu) := -(\Delta + k^2)V_k^\mu \quad \text{in distribution sense,} \tag{4.18}$$

where V_k^μ is given by Proposition 4.3.

We have the following basic properties of the partial balayage measure and the corresponding potential.

Lemma 4.7 Let μ and $k > 0$ be as in Lemma 4.2. Then

$$\text{Bal}_k(\mu) \leq 1 \quad \text{in } \mathbb{R}^n, \tag{4.19a}$$

$$U_k^{\text{Bal}_k(\mu)} \equiv V_k^\mu \quad \text{in } \mathbb{R}^n. \tag{4.19b}$$

We also have

$$U_k^{\text{Bal}_k(\mu)} \leq U_k^\mu \quad \text{in } \mathbb{R}^n \tag{4.19c}$$

and

$$U_k^{\text{Bal}_k(\mu)} = U_k^\mu \quad \text{in a neighborhood of } \mathbb{R}^n \setminus B_{R(n,k)}. \tag{4.19d}$$

Proof If we can show Eq. 4.19b, then Eqs. 4.19a and 4.19c are immediate consequence of Proposition 4.3 and the definition of $\mathcal{F}_k(\mu)$, while Eq. 4.19d is an immediate consequence of Remark 4.4. It remains to prove Eq. 4.19b. Write $u = U_k^\mu - V_k^\mu$, so $-(\Delta + k^2)u = \mu - \text{Bal}_k(\mu)$. Note that u has compact support by Remark 4.4. Thus

$$u = \tilde{\Phi}_k * (-(\Delta + k^2)u) = \tilde{\Phi}_k * (\mu - \text{Bal}_k(\mu)) = U_k^\mu - U_k^{\text{Bal}_k(\mu)}.$$

This proves that $U_k^{\text{Bal}_k(\mu)} = V_k^\mu$. □

We also make the following observation which will be very useful in our construction of k -quadrature domains.

Lemma 4.8 Let μ and $k > 0$ be as in Lemma 4.2. If

$$\text{Bal}_k(\mu) = \chi_D m \quad \text{for some open set } D \tag{4.20}$$

then

$$U_k^D \leq U_k^\mu \quad \text{in } \mathbb{R}^n, \tag{4.21a}$$

$$U_k^D = U_k^\mu \quad \text{in } \mathbb{R}^n \setminus D. \tag{4.21b}$$

Proof We already proved Eq. 4.21a in Lemma 4.7. Set $v = \text{Bal}_k(\mu)$ and rewrite Eq. 4.7 as

$$0 = \int_{B_{R(n,k)}} (U_k^\mu - U_k^D)(1 - \chi_D) dx = \int_{B_{R(n,k)} \setminus D} (U_k^\mu - U_k^D) dx.$$

Combining this equality with Eqs. 4.21a and 4.19d, we conclude Eq. 4.21b. □

We end this section by quickly relating our definition of partial balayage through an obstacle problem to a formulation in terms of energy minimization. Such a formulation is classical in the setting of $k = 0$ (see for instance [27]).

Remark 4.9 (Partial balayage and energy minimization) Let μ and $k > 0$ be as in Lemma 4.2. Using Proposition 5.1, we know that $\nu := \text{Bal}_k(\mu) \in L^\infty(B_{R(n,k)})$. We define the following bilinear form:

$$(\mu_1, \mu_2)_{e,k} := \iint_{B_{R(n,k)} \times B_{R(n,k)}} \tilde{\Phi}_k(x - y) d\mu_1(y) d\mu_2(x) = \int_{B_{R(n,k)}} U_k^{\mu_1}(x) d\mu_2(x)$$

for all $\mu_1, \mu_2 \in L^\infty(B_{R(n,k)})$. Using Lemma 4.7, we can write Eq. 4.7 as $(\nu - \mu, m - \nu)_{e,k} = 0$. Accordingly, for each $\sigma \in L^\infty(B_{R(n,k)})$ with $\sigma \leq m$, we see that

$$(\nu - \mu, \sigma - \nu)_{e,k} = (\nu - \mu, \sigma - m)_{e,k} \geq 0, \tag{4.22}$$

where the inequality follows from Lemma 4.7. By defining the ‘‘energy’’ $E_k(\lambda) := (\lambda, \lambda)_{e,k}$, we see that

$$(\nu - \mu, \sigma - \nu)_{e,k} = -E_k(\nu - \mu) + (\nu - \mu, \sigma - \mu)_{e,k},$$

thus from Eq. 4.22, we have

$$E_k(\nu - \mu) \leq (\nu - \mu, \sigma - \mu)_{e,k} \quad \text{for all } \sigma \in L^\infty(B_{R(n,k)}) \text{ with } \sigma \leq m. \tag{4.23}$$

When $U_k^{\mu_1}, U_k^{\mu_2} \in H_0^1(B_{R(n,k)})$ we can compute

$$\begin{aligned} (\mu_1, \mu_2)_{e,k} &= - \int_{B_{R(n,k)}} U_k^{\mu_1} (\Delta + k^2) U_k^{\mu_2} dx = a(U_k^{\mu_1}, U_k^{\mu_2}), \\ E_k(\mu_1) &= a(U_k^{\mu_1}, U_k^{\mu_1}) \geq 0, \end{aligned}$$

where $a(\cdot, \cdot)$ is the (real) inner product given by Lemma 4.5. Thus the notion of E_k as an energy functional makes sense. Using this observation and the Cauchy-Schwarz inequality, if we restrict σ in Eq. 4.23 to those functions satisfying $U_k^\sigma - U_k^\mu \in H_0^1(B_{R(n,k)})$, then we have

$$\begin{aligned} E_k(\nu - \mu) &\leq a(U_k^{\nu - \mu}, U_k^{\sigma - \mu}) \\ &\leq (a(U_k^\nu - U_k^\mu, U_k^\nu - U_k^\mu))^{\frac{1}{2}} (a(U_k^\sigma - U_k^\mu, U_k^\sigma - U_k^\mu))^{\frac{1}{2}} \\ &\equiv (E_k(\nu - \mu))^{\frac{1}{2}} (E_k(\sigma - \mu))^{\frac{1}{2}}. \end{aligned}$$

Therefore, the partial balayage ν minimizes the energy in the following sense:

$$E_k(\nu - \mu) \leq E_k(\sigma - \mu) \quad \text{for all } \sigma \in L^\infty(B_{R(n,k)}) \text{ with } U_k^\sigma \in \mathcal{F}_k(\mu), \tag{4.24}$$

where $\mathcal{F}_k(\mu)$ is given by Eq. 4.1. Here we also refer to [47, Section 30] for a related discussion.

5 Structure of Partial Balayage

In this section we prove the following proposition which provides information concerning the structure of $\text{Bal}_k(\mu)$. This will in particular be useful when we later on wish to construct k -quadrature domains.

Proposition 5.1 (Structure of partial balayage) *Let μ and $k > 0$ be as in Lemma 4.2 and let $\nu := \text{Bal}_k(\mu)$. Then*

$$\min\{\mu, m\} \leq \nu \leq m \quad \text{in } \mathbb{R}^n. \tag{5.1}$$

Furthermore, if we define the open sets

$$D(\mu) := \mathbb{R}^n \setminus \text{supp}(m - \nu), \text{ and} \tag{5.2}$$

$$\omega(\mu) := \{x \in \mathbb{R}^n \mid U_k^\mu(x) > U_k^\nu(x)\} \tag{5.3}$$

then $\omega(\mu) \subset D(\mu)$ and for each measurable set D with $\omega(\mu) \subset D \subset D(\mu)$ we have

$$v = \chi_D m + \chi_{\mathbb{R}^n \setminus D} \mu. \tag{5.4}$$

Remark 5.2 The corresponding result for $k = 0$ can be found in [26, Theorem 2.3(c)]. We refer also to [24, 27, 46], and in particular [27, Figure 3] for a visualization.

The set $D(\mu)$ is called the saturated set for $v = \text{Bal}_k(\mu)$, which is the largest open set \mathcal{O} in \mathbb{R}^n such that $v|_{\mathcal{O}} = m|_{\mathcal{O}}$. By our assumptions and Lemma 4.7 both $\text{supp}(v)$ and $\text{supp}(\mu)$ are contained in $B_{R(n,k)}$, and hence $D(\mu) \subset B_{R(n,k)}$. We also note that if $\mathcal{O} \subset D(\mu) \setminus \omega(\mu)$ has positive Lebesgue measure then $\mu|_{\mathcal{O}} = v|_{\mathcal{O}} = m|_{\mathcal{O}}$. In particular, if the density of μ is greater than 1 on $\text{supp}(\mu)$ it holds that $m(D(\mu) \setminus \omega(\mu)) = 0$. See also [26, Remark 2.4] for some discussions on the relation between $D(\mu)$ and $\omega(\mu)$ for the case when $k = 0$.

Proof of Proposition 5.1 Step 1: A minimization problem. Let $\xi \in H_0^1(B_{R(n,k)})$ be the unique solution to $-(\Delta + k^2)\xi = (1 - \mu)_+$, and consider the constraint set

$$\hat{\mathcal{K}}_k = \{ w \in H_0^1(B_{R(n,k)}) \mid w \geq \xi - u_* \},$$

where $u_* \in H_0^1(B_{R(n,k)})$ is the function appearing in the proof of Proposition 4.3. We recall that u_* minimizes the functional $a(u, u)$ among all functions in $\tilde{\mathcal{K}}_k$, where a is the bilinear form defined in Eq. 4.9 and $\tilde{\mathcal{K}}_k$ was defined in Eq. 4.12. Note that $\hat{\mathcal{K}}_k$ is nonempty since $\xi - u_* \in \hat{\mathcal{K}}_k$.

By Lemma 4.5 and Stampacchia’s Theorem (see [8, Theorem 5.6]) there exists a unique $w_* \in \hat{\mathcal{K}}_k$ which minimizes the functional $w \mapsto a(w, w)$ among $w \in \hat{\mathcal{K}}_k$. Moreover, the minimizer w_* is characterized by the property

$$a(w_*, w - w_*) = \langle -(\Delta + k^2)w_*, w - w_* \rangle \geq 0 \quad \text{for all } w \in \hat{\mathcal{K}}_k. \tag{5.5}$$

Step 2: Complementarity formulation. Since $w_* \in \hat{\mathcal{K}}_k$, we can in Eq. 5.5 restrict w to those satisfying $w \geq w_*$. The definition of a implies that

$$(\Delta + k^2)w_* \leq 0 \quad \text{in } B_{R(n,k)}. \tag{5.6}$$

Choosing $w = \xi - u_*$ in Eq. 5.5,

$$\langle (\Delta + k^2)w_*, \xi - u_* - w_* \rangle \leq 0,$$

which along with Eq. 5.6 and the fact that $w_* \geq \xi - u_*$ implies

$$\langle (\Delta + k^2)w_*, \xi - u_* - w_* \rangle = 0. \tag{5.7}$$

In fact, if $w_* \in \hat{\mathcal{K}}_k$ satisfies Eqs. 5.6 and 5.7, then

$$\begin{aligned} & \langle (\Delta + k^2)w_*, w - w_* \rangle \\ &= \langle (\Delta + k^2)w_*, w - (\xi - u_*) \rangle + \langle (\Delta + k^2)w_*, \xi - u_* - w_* \rangle \leq 0, \end{aligned}$$

for all $w \in \hat{\mathcal{K}}_k$. Hence the minimizer $w_* \in \hat{\mathcal{K}}_k$ can also be characterized by the complementarity problem Eqs. 5.6 and 5.7.

Step 3: An energy inequality. We can rewrite Eq. 5.7 as

$$\langle (\Delta + k^2)w_*, \xi - w_* \rangle = \langle (\Delta + k^2)w_*, u_* \rangle. \tag{5.8}$$

The inequalities $(\Delta + k^2)\xi = -(1 - \mu)_+ \leq 0$ and $w_* \geq \xi - u_*$ (i.e. $u_* \geq \xi - w_*$) thus imply that

$$\langle (\Delta + k^2)\xi, \xi - w_* \rangle \geq \langle (\Delta + k^2)\xi, u_* \rangle. \tag{5.9}$$

Combining Eqs. 5.8 and 5.9, one finds

$$\begin{aligned} a(\xi - w_*, \xi - w_*) &= \langle -(\Delta + k^2)(\xi - w_*), \xi - w_* \rangle \\ &\leq \langle -(\Delta + k^2)(\xi - w_*), u_* \rangle = a(\xi - w_*, u_*) \\ &\leq a(\xi - w_*, \xi - w_*)^{\frac{1}{2}} a(u_*, u_*)^{\frac{1}{2}}. \end{aligned}$$

By Lemma 4.5 the bilinear form a is positive, and thus we obtain the energy inequality

$$a(\xi - w_*, \xi - w_*) \leq a(u_*, u_*). \tag{5.10}$$

Step 4: Verifying that $w_* = \xi - u_*$. If we can show that $\xi - w_* \in \tilde{\mathcal{K}}_k$, i.e. that it satisfies

$$\xi - w_* \geq \varphi - U_k^\mu \quad \text{in } B_{R(n,k)} \tag{5.11}$$

where φ is the function in Eq. 4.10, then since u_* minimizes $a(u, u)$ among all $u \in \tilde{\mathcal{K}}_k$ the inequality in Eq. 5.10 implies that $\xi - w_* = u_*$ in $B_{R(n,k)}$, in other words $w_* = \xi - u_*$ in $B_{R(n,k)}$.

To prove Eq. 5.11 we argue as follows. Let

$$\phi := \min\{w_*, \xi - (\varphi - U_k^\mu)\} \quad \text{in } B_{R(n,k)}. \tag{5.12}$$

By the definition of ϕ and Proposition A.5,

$$\phi \leq w_*, \quad \phi \in \hat{\mathcal{K}}_k, \quad \text{and} \quad -(\Delta + k^2)\phi \geq 0 \quad \text{in } B_{R(n,k)}. \tag{5.13}$$

Using Eq. 5.13 and the facts that $w_* \in H_0^1(B_{R(n,k)})$ and $-(\Delta + k^2)w_* \geq 0$ in $B_{R(n,k)}$, we have in terms of distributional pairings in $B_{R(n,k)}$ that

$$\begin{aligned} a(\phi, \phi) &= \langle -(\Delta + k^2)\phi, \phi \rangle \\ &\leq \langle -(\Delta + k^2)\phi, w_* \rangle = \langle \phi, -(\Delta + k^2)w_* \rangle \\ &\leq \langle w_*, -(\Delta + k^2)w_* \rangle = a(w_*, w_*). \end{aligned}$$

Since w_* was defined to be the unique minimizer of $a(w, w)$ among $w \in \hat{\mathcal{K}}_k$, we obtain

$$\phi = w_* \quad \text{in } B_{R(n,k)}.$$

By the definition of ϕ this can equivalently be stated as

$$w_* \leq \xi - (\varphi - U_k^\mu) \quad \text{in } B_{R(n,k)}.$$

After rearranging we deduce the desired inequality Eq. 5.11. By the discussion following Eq. 5.11 it holds that

$$w_* = \xi - u_* \quad \text{in } B_{R(n,k)}.$$

Step 5: Proving Eq. 5.1. By 5.1, Eq. 5.6, and the definition of ξ ,

$$(\Delta + k^2)u_* \geq (\Delta + k^2)\xi = -(1 - \mu)_+. \tag{5.14}$$

From Eqs. 4.11 and 4.19b we deduce that

$$u_* = \varphi - U_k^\nu. \tag{5.15}$$

Combining Eqs. 5.14 and 5.15, we obtain that in $B_{R(n,k)}$

$$\begin{aligned} \nu - 1 &\geq -(1 - \mu)_+ = -\max\{1 - \mu, 0\} = \min\{\mu - 1, 0\} \\ &\iff \min\{1, \mu\} \leq \nu. \end{aligned}$$

By the definition of partial balayage $\nu \leq 1$ in $B_{R(n,k)}$, and we have arrived at Eq. 5.1.

Step 6: Proving Eq. 5.4. By the Calderón–Zygmund inequality $U_k^\mu, U_k^\nu \in \bigcap_{p < \infty} W_{loc}^{2,p}(\mathbb{R}^n)$ and hence $\omega(\mu)$ is well defined as an open set. From Eq. 4.7 and Proposition 4.3, it follows that

$$0 \leq \int_{\omega(\mu)} (U_k^\mu - U_k^\nu) d(\mathbf{m} - \nu) \leq \int_{B_{R(n,k)}} (U_k^\mu - U_k^\nu) d(\mathbf{m} - \nu) = 0,$$

and hence

$$\int_{\omega(\mu)} (U_k^\mu - U_k^\nu) d(\mathbf{m} - \nu) = 0. \tag{5.16}$$

Consequently, $\nu|_{\omega(\mu)} = \mathbf{m}|_{\omega(\mu)}$. Since $\omega(\mu)^c = \{x \in \mathbb{R}^n : U_k^\nu(x) = U_k^\mu(x)\}$ and $U_k^\mu, U_k^\nu \in W_{loc}^{2,p}(\mathbb{R}^n)$ it holds that $(\Delta + k^2)(U_k^\nu - U_k^\mu) = 0$ almost everywhere on this set. Therefore, $\nu|_{\omega(\mu)^c} = \mu|_{\omega(\mu)^c}$.

Consequently, for any D as in the proposition we by the definition of $D(\mu)$ have $\nu|_{D \setminus \omega(\mu)} = \mathbf{m}|_{D \setminus \omega(\mu)}$ and thus the claimed decomposition

$$\nu = \chi_D \mathbf{m} + \chi_{\mathbb{R}^n \setminus D} \mu$$

follows. This completes the proof of Proposition 5.1. □

We next deduce the following lemma.

Lemma 5.3 *Let μ and $k > 0$ be as in Lemma 4.2. Suppose there is an open set D such that $\overline{D} \subset B_{R(n,k)}$ and $\text{supp}(\mu) \subset D$ and a distribution u satisfying*

$$\begin{cases} (\Delta + k^2)u = \chi_D - \mu & \text{in } B_{R(n,k)}, \\ u > 0 & \text{in } D, \\ u = 0 & \text{in } B_{R(n,k)} \setminus D. \end{cases} \tag{5.17}$$

Then $\text{Bal}_k(\mu) = \chi_D \mathbf{m}$, $D = \omega(\mu)$ and D is a k -quadrature domain for μ .

Proof of Lemma 5.3 Since u (extended by zero outside $B_{R(n,k)}$) is a compactly supported distribution, we have

$$u = \tilde{\Phi}_k * (-(\Delta + k^2)u) = U_k^\mu - U_k^D.$$

Since u is non-negative, then we know that $U_k^D \in \mathcal{F}_k(\mu)$, where $\mathcal{F}_k(\mu)$ is the collection of functions given in Eq. 4.1. For each $v \in \mathcal{F}_k(\mu)$, since $u = 0$ in $B_{R(n,k)} \setminus D$, we see that

$$w := U_k^D - v = U_k^D - U_k^\mu + U_k^\mu - v \geq 0 \quad \text{in } B_{R(n,k)} \setminus D.$$

On the other hand, we have $(\Delta + k^2)w = -1 - (\Delta + k^2)v \leq 0$ in D . Therefore the maximum principle in Proposition A.4 implies that $w \geq 0$ in D as well. This shows that U_k^D is the largest element in $\mathcal{F}_k(\mu)$, so by the definition of partial balayage Eq. 4.18 we have

$$\text{Bal}_k(\mu) = -(\Delta + k^2)U_k^D = \chi_D \mathbf{m}.$$

By the above we see that $D = \{u > 0\} = \{U_k^\mu > U_k^{\text{Bal}_k(\mu)}\} = \omega(\mu)$.

Since $u \in C^1(\mathbb{R}^n)$ attains its minimum in D^c it holds that $|\nabla u| = 0$ in D^c . Therefore, since by assumption $\text{supp}(\mu) \subset D$, Proposition 2.1 implies that D is a k -quadrature domain for μ . □

6 Performing Balayage in Smaller Steps

Fix $\gamma > 0$ and assume that $\mu_1, \mu_2 \in L^\infty(B_\gamma)$ are non-negative. By Proposition 4.3, there exists a positive constant c_n such that if

$$0 < k < c_n \min\{\gamma^{-1}, \mu_1(\mathbb{R}^n)^{-\frac{1}{n}}\}, \tag{6.1}$$

then $U_k^{\text{Bal}_k(\mu_1)}$ is the largest element in $\mathcal{F}_k(\mu_1)$ (defined as in Eq. 4.1) and $U_k^{\text{Bal}_k(\mu_1)} = U_k^{\mu_1}$ near $\partial B_{R(n,k)}$. Again Proposition 4.3 also implies that if

$$0 < k < c_n \min\{\gamma^{-1}, (\mu_1 + \mu_2)(\mathbb{R}^n)^{-\frac{1}{n}}\}, \tag{6.2}$$

then $U_k^{\text{Bal}_k(\mu_1+\mu_2)}$ is the largest element of $\mathcal{F}_k(\mu_1 + \mu_2)$ and $U_k^{\text{Bal}_k(\mu_1+\mu_2)} = U_k^{\mu_1+\mu_2}$ near $\partial B_{R(n,k)}$.

Finally, if we additionally assume that $\text{supp}(v_1) \subset B_\gamma$ with $v_1 = \text{Bal}_k(\mu_1)$, Proposition 4.3 implies that if

$$0 < k < c_n \min\{\gamma^{-1}, (v_1 + \mu_2)(\mathbb{R}^n)^{-\frac{1}{n}}\}, \tag{6.3}$$

then $U_k^{\text{Bal}_k(v_1+\mu_2)}$ is the largest element of $\mathcal{F}_k(v_1 + \mu_2)$ and $U_k^{\text{Bal}_k(v_1+\mu_2)} = U_k^{v_1+\mu_2}$ near $\partial B_{R(n,k)}$. Using Proposition 5.1 we observe that

$$\text{supp}(\mu_1) \subset \text{supp}(\mu_1 + \mu_2) \subset \text{supp}(v_1 + \mu_2) \subset B_\gamma.$$

We are now ready to prove the following proposition:

Proposition 6.1 *Let $\gamma > 0$ and $\mu_1, \mu_2 \in L^\infty(B_\gamma)$ be non-negative and such that $\text{supp}(\text{Bal}_k(\mu_1)) \subset B_\gamma$. If*

$$0 < k < c_n \min\{\gamma^{-1}, (\mu_1 + \mu_2)(\mathbb{R}^n)^{-\frac{1}{n}}, (\text{Bal}_k(\mu_1) + \mu_2)(\mathbb{R}^n)^{-\frac{1}{n}}\}$$

with c_n (depending only on the dimension n), then

$$\text{Bal}_k(\mu_1 + \mu_2) = \text{Bal}_k(\text{Bal}_k(\mu_1) + \mu_2)$$

and

$$\omega(\mu_1 + \mu_2) = \omega(\mu_1) \cup \omega(\text{Bal}_k(\mu_1) + \mu_2).$$

Proof Note that if k satisfies the inequality in the proposition then k satisfies the inequalities Eqs. 6.1, 6.2, and 6.3.

We begin by showing the equality $\text{Bal}_k(\mu_1 + \mu_2) = \text{Bal}_k(\text{Bal}_k(\mu_1) + \mu_2)$.

Since $U_k^{v_1+\mu_2} = U_k^{v_1} + U_k^{\mu_2}$ and $U_k^{v_1} = U_k^{\mu_1}$ near $\partial B_{R(n,k)}$ it suffices to show that

$$U_k^{\text{Bal}_k(v_1+\mu_2)} = U_k^{\text{Bal}_k(\mu_1+\mu_2)} \quad \text{in } B_{R(n,k)}. \tag{6.4}$$

Step 1: The implication “ \leq ” of Eq. 6.4. Using Lemma 4.7 we observe that

$$\begin{aligned} U_k^{\text{Bal}_k(v_1+\mu_2)} &\leq U_k^{v_1+\mu_2} = U_k^{v_1} + U_k^{\mu_2} \\ &\leq U_k^{\mu_1} + U_k^{\mu_2} = U_k^{\mu_1+\mu_2} \quad \text{in } B_{R(n,k)} \end{aligned}$$

and

$$(\Delta + k^2)U_k^{\text{Bal}_k(v_1+\mu_2)} \geq -1 \quad \text{in } B_{R(n,k)}.$$

Thus $U_k^{\text{Bal}_k(v_1+\mu_2)} \in \mathcal{F}(\mu_1 + \mu_2)$. Since $U_k^{\text{Bal}_k(\mu_1+\mu_2)}$ is the largest element in $\mathcal{F}(\mu_1 + \mu_2)$, we arrive at

$$U_k^{\text{Bal}_k(v_1+\mu_2)} \leq U_k^{\text{Bal}_k(\mu_1+\mu_2)} \quad \text{in } B_{R(n,k)}. \tag{6.5}$$

Step 2: The implication “ \geq ” of Eq. 6.4. Observe that

$$U_k^{\text{Bal}_k(\mu_1+\mu_2)} - U_k^{\mu_2} \leq U_k^{\mu_1+\mu_2} - U_k^{\mu_2} = U_k^{\mu_1} \quad \text{in } B_{R(n,k)}$$

and

$$(\Delta + k^2)(U_k^{\text{Bal}_k(\mu_1+\mu_2)} - U_k^{\mu_2}) \geq -1 + \mu_2 \geq -1 \quad \text{in } B_{R(n,k)}.$$

Thus $U_k^{\text{Bal}_k(\mu_1+\mu_2)} - U_k^{\mu_2} \in \mathcal{F}(\mu_1)$. Since $U_k^{v_1}$ is the largest element in $\mathcal{F}(\mu_1)$, it holds that

$$U_k^{\text{Bal}_k(\mu_1+\mu_2)} - U_k^{\mu_2} \leq U_k^{v_1} \quad \text{in } B_{R(n,k)},$$

and hence

$$U_k^{\text{Bal}_k(\mu_1+\mu_2)} \leq U_k^{v_1} + U_k^{\mu_2} = U_k^{v_1+\mu_2} \quad \text{in } B_{R(n,k)}.$$

Furthermore,

$$(\Delta + k^2)U_k^{\text{Bal}_k(\mu_1+\mu_2)} \geq -1 \quad \text{in } B_{R(n,k)}.$$

Thus $U_k^{\text{Bal}_k(\mu_1+\mu_2)} \in \mathcal{F}(v_1 + \mu_2)$. Since $U_k^{\text{Bal}_k(v_1+\mu_2)}$ is the largest element in $\mathcal{F}(v_1 + \mu_2)$, it follows that

$$U_k^{\text{Bal}_k(v_1+\mu_2)} \geq U_k^{\text{Bal}_k(\mu_1+\mu_2)} \quad \text{in } B_{R(n,k)}. \tag{6.6}$$

Step 3: Conclusion. Combining Eqs. 6.5 and 6.6 implies Eq. 6.4 and completes the proof that

$$\text{Bal}_k(\mu_1 + \mu_2) = \text{Bal}_k(v_1 + \mu_2).$$

In the proof of this equality we established that

$$U_k^{\text{Bal}_k(\mu_1+\mu_2)} = U_k^{\text{Bal}_k(v_1+\mu_2)} \leq U_k^{v_1+\mu_2} = U_k^{v_1} + U_k^{\mu_2} \leq U_k^{\mu_1} + U_k^{\mu_2} = U_k^{\mu_1+\mu_2}.$$

The first inequality is an equality only in $\omega(\text{Bal}_k(\mu_1) + \mu_2)^c$ and the second is an equality only in $\omega(\mu_1)^c$. Therefore, the combined inequality, $U_k^{\text{Bal}_k(\mu_1+\mu_2)}(x) \leq U_k^{\mu_1+\mu_2}(x)$ is an equality only for $x \in \omega(v_1 + \mu_2)^c \cap \omega(\mu_1)^c = (\omega(v_1 + \mu_2) \cup \omega(\mu_1))^c$. By definition, $\omega(\mu_1 + \mu_2)$ is the set where this inequality is strict so the claim follows. This completes the proof of Proposition 6.1. □

7 Construction of k -Quadrature Domains

In this section our aim is to prove the following theorem, which contains the statement of Theorem 1.6.

Theorem 7.1 *Let μ be a positive measure supported in B_ϵ for some $\epsilon > 0$. There exists a constant $c_n > 0$ depending only on the dimension such that if*

$$0 < k < \frac{c_n}{\mu(\mathbb{R}^n)^{1/n}} \quad \text{and} \quad \epsilon < c_n \mu(\mathbb{R}^n)^{1/n}, \tag{7.1}$$

then there exists an open connected set D with real-analytic boundary satisfying $\overline{D} \subset B_{R(n,k)}$ which is a k -quadrature domain for μ . Moreover, for each $w \in L^1(D) \cap L^1(\mu)$ satisfying $(\Delta + k^2)w \geq 0$ in D we have

$$\int_D w(x) dx \geq \int w(x) d\mu(x). \tag{7.2}$$

Remark 7.2 As we shall see in the proof the k -quadrature domain is constructed as the non-contact set ω of the partial balayage of a measure obtained by averaging μ over a small ball. If μ satisfies the assumptions of Lemma 4.2 then $\text{Bal}_k(\mu)$ is well-defined and the k -quadrature domain D we construct is precisely $\omega(\mu)$.

However, our definitions and results concerning $Bal_k(\mu)$ and $\omega(\mu)$ are not valid for every μ as in the statement of the theorem, and as such we need to take some care.

Before we turn to the proof of Theorem 7.1 we prove some preliminary results that we will need in our main argument.

The first is a simple lemma concerning the partial balayage of a multiple of Lebesgue measure restricted to a ball.

Lemma 7.3 *Let $0 < r < r' < \frac{1}{2} j_{\frac{n-2}{2}, 1} k^{-1} = R(n, k)$. Then there exists a positive constant c_n (depending only on the dimension n) such that if*

$$0 < k \leq c_n \min \left\{ \frac{1}{r}, \frac{J_{\frac{n}{2}}(kr)^{\frac{1}{n}}}{J_{\frac{n}{2}}(kr')^{\frac{1}{n}}(rr')^{\frac{1}{2}}} \right\}, \tag{7.3}$$

then

$$Bal_k \left(\frac{c_{n,k,r'}^{MVT}}{c_{n,k,r}^{MVT}} \chi_{B_r} m \right) = \chi_{B_{r'}} m \tag{7.4}$$

and

$$\omega \left(\frac{c_{n,k,r'}^{MVT}}{c_{n,k,r}^{MVT}} \chi_{B_r} m \right) = B_{r'}. \tag{7.5}$$

Remark 7.4 Since $t \mapsto t^{\frac{n}{2}} J_{\frac{n}{2}}(t)$ is strictly increasing on $t \in [0, j_{\frac{n-2}{2}, 1}]$, then we see that

$$\frac{c_{n,k,r'}^{MVT}}{c_{n,k,r}^{MVT}} = \frac{(r')^{n/2} J_{\frac{n}{2}}(kr')}{r^{n/2} J_{\frac{n}{2}}(kr)} > 1. \tag{7.6}$$

Since $t \mapsto t^{-\frac{n}{2}} J_{\frac{n}{2}}(t)$ is a decreasing function on $[0, j_{\frac{n+2}{2}, 1}]$, we find that

$$\begin{aligned} & Bal_k \left(\frac{c_{n,k,r'}^{MVT}}{c_{n,k,r}^{MVT}} \chi_{B_r} m \right) (\mathbb{R}^n) - \frac{c_{n,k,r'}^{MVT}}{c_{n,k,r}^{MVT}} \chi_{B_r} m (\mathbb{R}^n) \\ &= m(B_1) \left((r')^n - \frac{(r')^{n/2} J_{\frac{n}{2}}(kr')}{r^{n/2} J_{\frac{n}{2}}(kr)} r^n \right) \\ &= m(B_1) (r')^{\frac{n}{2}} k^{-\frac{n}{2}} J_{\frac{n}{2}}(kr') \left(\frac{(kr')^{\frac{n}{2}}}{J_{\frac{n}{2}}(kr')} - \frac{(kr)^{\frac{n}{2}}}{J_{\frac{n}{2}}(kr)} \right) \geq 0. \end{aligned} \tag{7.7}$$

Proof of Lemma 7.3 For each $x \in \mathbb{R}^n$, we see that the distribution $y \mapsto \tilde{\Phi}_k(x - y)$ is in $L^1_{loc}(\mathbb{R}^n)$ and satisfies $(\Delta + k^2)\tilde{\Phi}_k(x - \cdot) = -\delta_x \leq 0$ in \mathbb{R}^n . By applying the MVT in Proposition A.2, we have

$$\frac{1}{c_{n,k,r}^{MVT}} U_k^{B_r}(x) = \frac{1}{c_{n,k,r}^{MVT}} \int_{B_r} \tilde{\Phi}_k(x - y) dy \geq \frac{1}{c_{n,k,r'}^{MVT}} \int_{B_{r'}} \tilde{\Phi}_k(x - y) dy = \frac{1}{c_{n,k,r'}^{MVT}} U_k^{B_{r'}}(x) \tag{7.8}$$

for all $x \in \mathbb{R}^n$, and equality holds if and only if $x \in \mathbb{R}^n \setminus B_{r'}$. In other words

$$u = \frac{c_{n,k,r'}^{MVT}}{c_{n,k,r}^{MVT}} U_k^{B_r} - U_k^{B_{r'}} \in C^1(\mathbb{R}^n)$$

satisfies

$$\begin{cases} (\Delta + k^2)u = \chi_{B_{r'}} - \frac{c_{n,k,r'}^{MVT}}{c_{n,k,r}^{MVT}} \chi_{B_r} & \text{in } \mathbb{R}^n \\ u > 0 & \text{in } B_{r'} \\ u = 0 & \text{in } \mathbb{R}^n \setminus B_{r'}. \end{cases} \tag{7.9}$$

The conclusion of the lemma follows by applying Lemma 5.3. □

The second result we require is an analogue of Proposition 2.4 but for sub-solutions of the Helmholtz equation. We again follow the argument in [43, Lemma 5.1] which considered the case $k = 0$.

Proposition 7.5 *Let $k \geq 0$, and let $D \subset \mathbb{R}^n$ be a bounded open set. Let Ψ_k be any fundamental solution of $-(\Delta + k^2)$ and let $\Omega \supset \overline{D}$ be any open set in \mathbb{R}^n . Then the linear span with positive coefficients of*

$$F = \{\pm \partial^\alpha \Psi_k(z - \cdot)|_D : z \in \Omega \setminus D, |\alpha| \leq 1\} \cup \{-\Psi_k(z - \cdot)|_D : z \in D\}$$

is dense in

$$S_k L^1(D) = \{w \in L^1(D) : (\Delta + k^2)w \geq 0 \text{ in } D\}$$

with respect to the $L^1(D)$ topology.

Proof We first show that if any bounded linear functional ℓ on $L^1(D)$ with $\ell|_F \geq 0$ also satisfies

$$\ell|_{S_k L^1(D)} \geq 0, \tag{7.10}$$

then we have $\overline{G} = S_k L^1(D)$, where G is the linear span with positive coefficients of F . Suppose to the contrary that there exists $f_0 \in S_k L^1(D) \setminus \overline{G}$. Using the Hahn-Banach theorem (second geometric form, see e.g. [8, Theorem 1.7]), there exists a closed hyperplane $\{\ell_0 = \alpha\}$ that strictly separates the closed set \overline{G} and the compact set $\{f_0\}$, thus we have

$$\ell_0(f_0) < \alpha < \ell_0(f) \quad \text{for all } f \in G. \tag{7.11}$$

Since $\lambda \ell_0(f) = \ell_0(\lambda f) > \alpha$ for all $\lambda > 0$ and each fixed $f \in G$, we deduce that $\ell_0(f) \geq 0$ for all $f \in G$ and that $\alpha \leq 0$. By Eq. 7.10 and since $f_0 \in S_k L^1(D)$ we know that $\ell_0(f_0) \geq 0$. Combining this with Eq. 7.11 and the fact that $\alpha \leq 0$ gives a contradiction.

Now let ℓ be a bounded linear functional on $L^1(D)$ with $\ell|_F \geq 0$. We need to prove that $\ell|_{S_k L^1(D)} \geq 0$. Since the dual of $L^1(D)$ is $L^\infty(D)$, there is a function $f \in L^\infty(D)$ with

$$\ell(w) = \int_D f w \, dx, \quad w \in L^1(D).$$

We extend f by zero to \mathbb{R}^n and consider the function

$$u(z) = -(\Psi_k * f)(z) \quad \text{for all } z \in \Omega.$$

By the assumption $\ell|_F \geq 0$, the function u satisfies

$$\begin{cases} (\Delta + k^2)u = f & \text{in } \Omega, \\ u = |\nabla u| = 0 & \text{in } \Omega \setminus D, \\ u \geq 0 & \text{in } D. \end{cases}$$

Our aim is to employ the same argument as in the proof of Eq. 2.3 to show that

$$\int_D ((\Delta + k^2)u)w \, dx \geq 0, \quad \text{for all } w \in S_k L^1(D),$$

which implies $\ell|_{S_k L^1(D)} \geq 0$. However, in order to carry out the the integration by parts which concluded that argument we used that solutions of the Helmholtz equation are smooth in the interior of D , this is not necessarily the case for sub-solutions. To circumvent this issue we use a classical mollification argument. Fix a non-negative $\psi \in C_c^\infty(\mathbb{R}^n)$ with support in B_1 and $\|\psi\|_{L^1(\mathbb{R}^n)} = 1$. For $\epsilon > 0$ set $\psi_\epsilon(x) = \epsilon^{-n}\psi(x/\epsilon)$. For $w \in S_k L^1(D)$ set $w_\epsilon = w * \psi_\epsilon$, which is well defined and C^∞ near any compact subset K of D if $\epsilon < \text{dist}(K, D^c)$. Then $w_\epsilon \rightarrow w$ in $L^1(K)$ as $\epsilon \rightarrow 0^+$. We claim that $(\Delta + k^2)w_\epsilon(x) \geq 0$ in K for $\epsilon < \text{dist}(K, D^c)$. Indeed, since $(\Delta + k^2)w(x) \geq 0$ in D this is in particular the case in $B_\epsilon(y)$ for any $y \in K$. The claim follows by differentiating under the integral sign and using the non-negativity of ψ . With this approximation in hand the argument can be completed as in the proof of Eq. 2.3 by appealing to the L^1 convergence of w_ϵ to w in the support of the cutoff function ω_j , and applying the integration by parts argument with w replaced by w_ϵ . \square

Finally we need the following result which is in the spirit of Proposition 2.1. This result can be interpreted as saying that D is a quadrature domain for sub-solutions w satisfying $(\Delta + k^2)w \geq 0$ in D (i.e. D is a quadrature domain for metasubharmonic functions).

Corollary 7.6 *Let $k > 0$, and let $D, \Omega \subset \mathbb{R}^n$ be bounded open sets such that $\bar{D} \subset \Omega$, and let $\mu \in L^\infty(D)$ be a non-negative measure with $\text{supp}(\mu) \subset D$. If*

$$U_k^D = U_k^\mu \quad \text{in } \Omega \setminus D, \tag{7.12a}$$

$$U_k^D \leq U_k^\mu \quad \text{in } \Omega, \tag{7.12b}$$

then for each $w \in L^1(D)$ satisfying $(\Delta + k^2)w \geq 0$ in D we know that

$$\int_D w(x) dx \geq \int w(x) d\mu(x). \tag{7.13}$$

Proof By Eqs. 7.12a and 7.12b $U_k^\mu - U_k^D \geq 0$ with equality in $\Omega \setminus D$. By Calderón–Zygmund estimates $U_k^\mu, U_k^D \in C^1(\Omega)$. Since $U_k^\mu - U_k^D$ attains its minimum in $\Omega \setminus D$ it holds that $\nabla U_k^\mu = \nabla U_k^D$ in $\Omega \setminus D$. When combined with Eqs. 7.12a and 7.12b we conclude that

$$\int_D \partial^\alpha \tilde{\Phi}_k(z-x) dx = \int \partial^\alpha \tilde{\Phi}_k(z-x) d\mu(x) \quad \text{for all } z \in B_{R(n,k)} \setminus D, |\alpha| \leq 1, \tag{7.14a}$$

$$\int_D \tilde{\Phi}_k(z-x) dx \leq \int \tilde{\Phi}_k(z-x) d\mu(x) \quad \text{for all } z \in D. \tag{7.14b}$$

Let w be the function as in the statement of the lemma, and use Proposition 7.5 to find a sequence

$$w_j \in \text{span}_+ \left(\begin{aligned} & \{ \pm \partial^\alpha \tilde{\Phi}_k(z-\cdot)|_D : z \in B_{R(n,k)} \setminus D, |\alpha| \leq 1 \} \\ & \cup \{ -\tilde{\Phi}_k(z-\cdot)|_D : z \in D \} \end{aligned} \right)$$

with $w_j \rightarrow w \in L^1(D)$. From Eqs. 7.14a and 7.14b, we know that

$$\int_D w_j(x) dx \geq \int w_j(x) d\mu(x) \quad \text{for all } j. \tag{7.15}$$

Since $\mu \in L^\infty(D)$ Hölder’s inequality implies that

$$\left| \int (w_j(x) - w(x)) d\mu(x) \right| \leq \|\mu\|_{L^\infty(D)} \|w - w_j\|_{L^1(D)}.$$

Taking the limit $j \rightarrow \infty$ in Eq. 7.15 we therefore arrive at

$$\int_D w(x) dx \geq \int w(x) d\mu(x).$$

This is the desired inequality Eq. 7.13, and thus completes the proof. □

We are now ready to prove Theorem 7.1.

Proof of Theorem 7.1 Step 1: Constructing the k -quadrature domain. For $\epsilon < \delta < R(n, k)$ to be chosen set

$$h_\delta = \frac{1}{c_{n,k,\delta}^{MVT}} \chi_{B_\delta}. \tag{7.16}$$

Then

$$\mu * h_\delta(x) = \frac{\mu(B_\delta(x))}{c_{n,k,\delta}^{MVT}} \tag{7.17}$$

is non-negative and supported in $B_{\epsilon+\delta}$. Furthermore for all $x \in B_{\delta-\epsilon}$,

$$\mu * h_\delta(x) = \frac{\mu(B_\delta(x))}{c_{n,k,\delta}^{MVT}} = \frac{\mu(\mathbb{R}^n)}{c_{n,k,\delta}^{MVT}}. \tag{7.18}$$

Let us for $\kappa \geq 1$ and $r > 0$ define

$$\mu_{\kappa,r} := \kappa \chi_{B_r} m.$$

Set

$$\mu_1 := \mu_{\kappa,r} \quad \text{and} \quad \mu_2 := \mu * h_\delta - \mu_{\kappa,r}$$

with

$$1 < \kappa \leq \frac{\mu(\mathbb{R}^n)}{c_{n,k,\delta}^{MVT}} \quad \text{and} \quad 0 < r \leq \delta - \epsilon \tag{7.19}$$

to be chosen. Note that these choices of r, κ imply that the measures μ_1, μ_2 are non-negative. Furthermore, both measures have bounded densities with respect to Lebesgue measure and

$$\mu_1(\mathbb{R}^n) = \kappa m(B_r) \quad \text{and} \quad \mu_2(\mathbb{R}^n) = \frac{\mu(\mathbb{R}^n)m(B_\delta)}{c_{n,k,\delta}^{MVT}} - \kappa m(B_r).$$

Our aim is to appeal to Proposition 6.1 and perform an initial balayage of μ_1 by utilizing Lemma 7.3. To this end we choose

$$\kappa = \frac{c_{n,k,r'}^{MVT}}{c_{n,k,r}^{MVT}} = \frac{(r')^{\frac{n}{2}} J_{\frac{n}{2}}(kr')}{r^{\frac{n}{2}} J_{\frac{n}{2}}(kr)}.$$

for some $r < r' < R(n, k)$. By Lemma 7.3, if

$$0 < k < c_n \min \left\{ \frac{1}{r}, \frac{J_{\frac{n}{2}}(kr)^{\frac{1}{n}}}{J_{\frac{n}{2}}(kr')^{\frac{1}{n}}(rr')^{\frac{1}{2}}} \right\} = c_n \min \left\{ \frac{1}{r}, \frac{1}{\kappa^{\frac{1}{n}} r} \right\} = \frac{c_n}{r}, \tag{7.20}$$

then $\text{Bal}_k(\mu_1) = \chi_{B_{r'}} m$ and by Eq. 7.7 it holds that $\text{Bal}_k(\mu_1)(\mathbb{R}^n) \geq \mu_1(\mathbb{R}^n) = \kappa |B_r|$.

Consequently, $(\mu_1 + \mu_2)(\mathbb{R}^n) \leq (\text{Bal}_k(\mu_1) + \mu_2)(\mathbb{R}^n)$ and thus if

$$0 < k < c_n \min \left\{ \frac{1}{r'}, \frac{1}{\delta + \epsilon}, \frac{1}{((c_{n,k,\delta}^{MVT})^{-1} \mu(\mathbb{R}^n) \delta^n + (r')^n - \kappa r^n)^{1/n}} \right\}, \tag{7.21}$$

then Eq. 7.20 is valid, since $r' > r$, and furthermore Proposition 6.1 implies that

$$\text{Bal}_k(\mu * h_\delta) = \text{Bal}_k(\mu_1 + \mu_2) = \text{Bal}_k(\text{Bal}_k(\mu_1) + \mu_2) = \text{Bal}_k(\mu_{1,r'} + \mu * h_\delta - \mu_{\kappa,r}) \tag{7.22}$$

and

$$\omega(\mu * h_\delta) = \omega(\mu_1) \cup \omega(\text{Bal}_k(\mu_1) + \mu_2) \supset B_{r'}, \tag{7.23}$$

where we also used that $\omega(\mu_1) = B_{r'}$ by Lemma 7.3.

By construction

$$\mu_{1,r'} + \mu * h_\delta - \mu_{\kappa,r} = 0 \quad \text{outside } B_{r'} \cup B_{\epsilon+\delta} \tag{7.24}$$

and

$$\mu * h_\delta - \mu_{\kappa,r} \geq 0. \tag{7.25}$$

Therefore,

$$\mu_{1,r'} + \mu * h_\delta - \mu_{\kappa,r} \geq 1 \quad \text{in } B_{r'}. \tag{7.26}$$

Consequently, if we choose our parameters to satisfy

$$r' > \epsilon + \delta, \tag{7.27}$$

then the fact that $\mu * h_\delta \in L^\infty(\mathbb{R}^n)$ combined with Proposition 5.1 implies

$$\text{Bal}_k(\mu_{1,r'} + \mu * h_\delta - \mu_{\kappa,r}) = \chi_{D(\mu_{1,r'} + \mu * h_\delta - \mu_{\kappa,r})} \mathfrak{m}, \tag{7.28}$$

and $B_{r'} \subset D(\mu_{1,r'} + \mu * h_\delta - \mu_{\kappa,r})$.

Combining Eqs. 7.22 and 7.28, we have

$$\text{Bal}_k(\mu * h_\delta) = \text{Bal}_k(\mu_{1,r'} + \mu * h_\delta - \mu_{\kappa,r}) = \chi_{D(\mu_{1,r'} + \mu * h_\delta - \mu_{\kappa,r})} \mathfrak{m}.$$

Using Eq. 7.23 and that we shall choose our parameters so that $r' > \epsilon + \delta$ we find

$$\text{supp}(\mu * h_\delta) \subset B_{r'} \subset \omega(\mu * h_\delta),$$

therefore Lemma 5.3 implies that²

$$\text{Bal}_k(\mu * h_\delta) = \chi_{\omega(\mu * h_\delta)} \mathfrak{m}. \tag{7.29}$$

Using that only one connected component of $\omega(\mu * h_\delta)$ intersects $\text{supp}(\mu * h_\delta)$, we can argue as in [26, Corollary 2.3] to prove that $\omega(\mu * h_\delta)$ is connected.

By Lemma 4.8 and the definition of $\omega(\mu * h_\delta)$,

$$U_k^{\omega(\mu * h_\delta)} = U_k^{\mu * h_\delta} = \tilde{\Phi}_k * \mu * h_\delta \quad \text{in } B_{R(n,k)} \setminus \omega(\mu * h_\delta), \text{ and} \tag{7.30a}$$

$$U_k^{\omega(\mu * h_\delta)} < U_k^{\mu * h_\delta} = \tilde{\Phi}_k * \mu * h_\delta \quad \text{in } \omega(\mu * h_\delta). \tag{7.30b}$$

Under our assumptions, Corollary 7.6 implies that $\omega(\mu * h_\delta)$ is a k -quadrature domain for $\mu * h_\delta$ and furthermore we have the quadrature inequality for sub-solutions.

The MVT (Proposition A.2) implies that $\tilde{\Phi}_k * h_\delta(y) \leq \tilde{\Phi}_k(y)$ for all $y \in \mathbb{R}^n$ and equality holds if $|y| \geq \delta$. Therefore, by the non-negativity of μ ,

$$U_k^{\mu * h_\delta}(x) = \tilde{\Phi}_k * \mu * h_\delta(x) = \int (\tilde{\Phi}_k * h_\delta)(x - y) d\mu(y) \leq \tilde{\Phi}_k * \mu(x) = U_k^\mu(x)$$

with equality if $\text{dist}(x, \text{supp}(\mu)) \geq \delta$. In particular, since under our assumptions

$$\{x \in \mathbb{R}^n : \text{dist}(x, \text{supp}(\mu)) < \delta\} \subset B_{\epsilon+\delta} \subset B_{r'} \subset \omega(\mu * h_\delta)$$

²Note that $\mu * h_\delta$ does not necessarily have a density which is greater than 1 on its support and so the structure of its partial balayage in Eq. 7.29 does not follow directly from Proposition 5.1.

we have equality for $x \in B_{R(n,k)} \setminus \omega(\mu * h_\delta)$. We have thus arrived at³

$$U_k^{D(\mu * h_\delta)} = U_k^\mu \quad \text{in } B_{R(n,k)} \setminus \omega(\mu * h_\delta), \text{ and} \tag{7.31a}$$

$$U_k^{D(\mu * h_\delta)} < U_k^\mu \quad \text{in } \omega(\mu * h_\delta). \tag{7.31b}$$

We now show that we can choose the parameters r, δ, κ appropriately only depending on the measure μ , specifically we shall choose them depending on $\epsilon, \mu(\mathbb{R}^n)$. We choose $\delta = 2\epsilon$ and $r = \epsilon$, and let $\gamma = k\epsilon$.

Since $t \mapsto t^{-\frac{n}{2}} J_{\frac{n}{2}}(t)$ is a decreasing function on $[0, j_{\frac{n-2}{2},1}]$ satisfying that $\lim_{y \rightarrow 0+} y^{-\frac{n}{2}} J_{\frac{n}{2}}(y) = \frac{2^{-\frac{n}{2}}}{\Gamma(1+\frac{n}{2})}$, we by using the explicit form of $c_{n,k,r}^{MVT}$ find that

$$(c_{n,k,2\epsilon}^{MVT})^{-1} (2\epsilon)^n = \frac{\gamma^{\frac{n}{2}}}{\pi^{\frac{n}{2}} J_{\frac{n}{2}}(2\gamma)} \geq \frac{\Gamma(1 + \frac{n}{2})}{\pi^{\frac{n}{2}}}. \tag{7.32}$$

The required bound on k Eq. 7.21 is then valid if

$$0 < k < c_n \min \left\{ \frac{1}{r'}, \frac{1}{\epsilon}, \frac{1}{(\pi^{-\frac{n}{2}} \Gamma(1 + \frac{n}{2}) \mu(\mathbb{R}^n) + (r')^n)^{1/n}} \right\}. \tag{7.33}$$

Assume that $c_n \leq j_{\frac{n-2}{2},1}/4$ so that $0 \leq \gamma \leq j_{\frac{n-2}{2},1}/4$. Then, since $t \mapsto t^{\frac{n}{2}} J_{\frac{n}{2}}(t)$ is strictly increasing on $[0, j_{\frac{n-2}{2},1}]$ we can choose $r' = 4r$ which ensures that Eq. 7.27 is satisfied since

$$r' = 4r = 4\epsilon > 3\epsilon = \epsilon + \delta,$$

and we have

$$\kappa = \frac{c_{n,k,4r}^{MVT}}{c_{n,k,r}^{MVT}} = \frac{(4r)^{\frac{n}{2}} J_{\frac{n}{2}}(4\gamma)}{r^{\frac{n}{2}} J_{\frac{n}{2}}(\gamma)} = \frac{2^n J_{\frac{n}{2}}(4\gamma)}{J_{\frac{n}{2}}(\gamma)} > 1.$$

We now want to find a sufficient condition so that Eq. 7.19 holds. By the choice of κ and the definition of γ what we need to verify is the inequality

$$\kappa = \frac{2^n J_{\frac{n}{2}}(4\gamma)}{J_{\frac{n}{2}}(\gamma)} \leq \frac{\mu(\mathbb{R}^n)}{c_{n,k,2\epsilon}^{MVT}} = \frac{\mu(\mathbb{R}^n) \gamma^{\frac{n}{2}}}{(2\epsilon)^n \pi^{\frac{n}{2}} J_{\frac{n}{2}}(2\gamma)}, \tag{7.34}$$

or equivalently,

$$4^n \pi^{\frac{n}{2}} \frac{J_{\frac{n}{2}}(4\gamma) J_{\frac{n}{2}}(2\gamma)}{J_{\frac{n}{2}}(\gamma) \gamma^{\frac{n}{2}}} \epsilon^n \leq \mu(\mathbb{R}^n).$$

Since the function

$$\gamma \mapsto \frac{J_{\frac{n}{2}}(4\gamma) J_{\frac{n}{2}}(2\gamma)}{J_{\frac{n}{2}}(\gamma) \gamma^{\frac{n}{2}}}$$

is continuous it is bounded from above for all $\gamma \in [0, j_{\frac{n-2}{2},1}/4]$ by a constant depending only on n . Therefore, the required bound Eq. 7.19 holds if we assume that

$$\epsilon \leq c_n \mu(\mathbb{R}^n)^{1/n} \tag{7.35}$$

³If $\mu \in L^\infty(B_{R(n,k)})$ then Eqs. 7.31a and 7.31b combined with an application of Lemma 5.3 implies that

$$\omega(\mu) = \omega(\mu * h_\delta) \quad \text{and} \quad \text{Bal}_k(\mu) = \text{Bal}_k(\mu * h_\delta) = \chi_{\omega(\mu)} \mathbf{m}.$$

provided c_n is chosen sufficiently small, specifically so that

$$c_n \leq \min_{\gamma \in [0, J_{\frac{n-2}{2}, 1}^{n-2, 1}/4]} \left(\frac{J_{\frac{n}{2}}(\gamma) \gamma^{\frac{n}{2}}}{4^n \pi^{\frac{n}{2}} J_{\frac{n}{2}}(4\gamma) J_{\frac{n}{2}}(2\gamma)} \right)^{1/n}.$$

Since we chose our parameters so that $r' = 4\epsilon$, the require bound on k Eq. 7.33 (and thus also Eqs. 7.21 and 7.20) is valid if

$$0 < k < \frac{c_n}{\mu(\mathbb{R}^n)^{1/n}}. \tag{7.36}$$

Consequently, all our requirements are met provided

$$0 < k < \frac{c_n}{\mu(\mathbb{R}^n)^{1/n}} \quad \text{and} \quad \epsilon \leq c_n \mu(\mathbb{R}^n)^{1/n} \tag{7.37}$$

for some constant c_n depending only on n .

From here and on we let h denote the function h_δ with the particular choice $\delta = 2\epsilon$ in the discussion above. By the construction we have that

$$\text{Bal}_k(\mu * h) = \chi_{\omega(\mu * h)} m \tag{7.38}$$

and

$$\text{supp}(\mu) \subset \overline{B_\epsilon} \subset \text{supp}(\mu * h) \subset B_{4\epsilon} \subset \omega(\mu * h). \tag{7.39}$$

Step 2: $\omega(\mu * h)$ is a k -quadrature domain with respect to μ . Equations 7.31a and 7.31b imply that $U_k^\mu - U_k^{\omega(\mu * h)} \geq 0$ with equality in $B_{R(n,k)} \setminus \omega(\mu * h)$. By elliptic regularity $U_k^{\omega(\mu * h)}$ is $C^1(B_{R(n,k)})$ and U_k^μ is smooth away from $\text{supp}(\mu)$. Thus $U_k^\mu - U_k^{\omega(\mu * h)}$ attains its minimum in $B_{R(n,k)} \setminus \omega(\mu * h)$, and consequently

$$\nabla U_k^{\omega(\mu * h)} = \nabla U_k^\mu \quad \text{in } B_{R(n,k)} \setminus \omega(\mu * h). \tag{7.40}$$

Since $\text{supp}(\mu) \subset \omega(\mu * h) \subset B_{R(n,k)}$ the extension by zero of $U_k^\mu - U_k^{\omega(\mu * h)}$ to all of \mathbb{R}^n satisfies the assumptions of Proposition 2.1 and hence $\omega(\mu * h)$ is a k -quadrature domain with respect to μ .

As noted above, Eqs. 7.30a, 7.30b and Corollary 7.6 imply that

$$\int_{\omega(\mu * h)} w(x) dx \geq \int w(x) d(\mu * h)(x)$$

for all $w \in L^1(\omega(\mu * h))$ satisfying $(\Delta + k^2)w(x) \geq 0$ in $\omega(\mu * h)$. Assume further that $w \in L^1(\mu)$. By Fubini and since by construction $B_\delta(y) \subset \omega(\mu * h)$ for all $y \in \text{supp}(\mu)$ the integral above can be rewritten as

$$\begin{aligned} \int_{\omega(\mu * h)} w(x) d(\mu * h)(x) &= \int \left(\int_{\omega(\mu * h)} w(x) h_\delta(x - y) dx \right) d\mu(y) \\ &= \int \left(\frac{1}{c_{n,k,\delta}^{\text{MVT}}} \int_{B_\delta(y)} w(x) dx \right) d\mu(y). \end{aligned}$$

Since $B_\delta(y) \subset \omega(\mu * h)$ for all $y \in \text{supp}(\mu)$ the function $w(x)$ is a sub-solution of the Helmholtz equation on $B_\delta(y)$ for all $y \in \text{supp}(\mu)$. Therefore, the mean value inequality for sub-solutions of the Helmholtz equation implies that the expression within the parenthesis is greater than $w(y)$ for each $y \in \text{supp}(\mu)$. Since $w \in L^1(\mu)$ and μ is non-negative this proves Eq. 7.2.

Step 3: Regularity of $\partial\omega(\mu * h)$. We now show that $\omega(\mu * h)$ has real-analytic boundary $\partial\omega(\mu * h)$ by using the moving plane technique as in [27, Theorem 5.4]. Set

$u := U_k^{\mu * h} - U_k^{\omega(\mu * h)} \in \bigcap_{0 < \alpha < 1} C^{1, \alpha}(\overline{B_{R(n, k)}})$. By Proposition 4.3, we know that u is the smallest among all $w \in H_0^1(B_{R(n, k)})$ satisfying

$$w \geq 0 \quad \text{and} \quad (\Delta + k^2)w \leq -\mu * h + 1 \quad \text{in } B_{R(n, k)}. \tag{7.41}$$

Moreover, Lemma 4.8 and the fact that $\text{Bal}_k(\mu * h) = \chi_{\omega(\mu * h)} \mathbf{m}$ implies

$$u = 0 \quad \text{in } B_{R(n, k)} \setminus \omega(\mu * h). \tag{7.42}$$

Given $x_0 \in \partial\omega(\mu * h)$, there by Eq. 7.39 exists a hyperplane that separates $\text{supp}(\mu * h)$ and x_0 . Since Laplacian is translation and rotation invariant, without loss of generality, we may assume that the hyperplane is

$$\{x \in \mathbb{R}^n \mid x_n = 0\}, \tag{7.43}$$

and that $\text{supp}(\mu * h) \subset \{x \in \mathbb{R}^n \mid x_n < 0\}$. We define

$$(\omega(\mu * h))_{\text{loc}} = \omega(\mu * h) \cap \{x \in \mathbb{R}^n \mid x_n > 0\}$$

and

$$(\partial\omega(\mu * h))_{\text{loc}} = \partial\omega(\mu * h) \cap \{x \in \mathbb{R}^n \mid x_n > 0\}.$$

Let u^* be the reflection of u with respect to the hyperplane Eq. 7.43, that is, $u^*(x', x_n) = u(x', -x_n)$. We now define

$$v := u - \inf\{u, u^*\} = (u - u^*)_+.$$

Since $(\Delta + k^2)u \leq -\mu * h + 1 \leq 1$ in $B_{R(n, k)}$, we have $(\Delta + k^2)u^* \leq 1$ in $B_{R(n, k)}$. Since there exists a unique $\phi \in H_0^1(B_{R(n, k)})$ such that $(\Delta + k^2)\phi = 1$ in $B_{R(n, k)}$, using Proposition A.5, we know that

$$(\Delta + k^2) \inf\{u, u^*\} \leq 1 \quad \text{in } B_{R(n, k)}.$$

From $(\Delta + k^2)u = 1$ in $(\omega(\mu * h))_{\text{loc}}$, we have

$$(\Delta + k^2)v \geq 0 \quad \text{in } (\omega(\mu * h))_{\text{loc}}.$$

The boundary condition $v = 0$ on $\partial((\omega(\mu * h))_{\text{loc}})$ and using maximum principle in Proposition A.4 yield $v \leq 0$ in $(\omega(\mu * h))_{\text{loc}}$, and hence

$$v = 0 \quad \text{in } (\omega(\mu * h))_{\text{loc}},$$

because $v \geq 0$ by its definition. Thus we have

$$\frac{\partial u}{\partial x_n} \leq 0 \quad \text{on } \{x \in \mathbb{R}^n \mid x_n = 0\}. \tag{7.44}$$

From Eq. 7.42, we know that $\frac{\partial u}{\partial x_n} = 0$ on $(\partial\omega(\mu * h))_{\text{loc}}$. On the other hand, we know that

$$(\Delta + k^2)u = 1 \quad \text{in } (\omega(\mu * h))_{\text{loc}}.$$

Hence $\frac{\partial u}{\partial x_n} \in C^\infty((\omega(\mu * h))_{\text{loc}}) \cap \bigcap_{0 < \alpha < 1} C^{0, \alpha}(\overline{(\omega(\mu * h))_{\text{loc}}})$ and it satisfies

$$(\Delta + k^2) \frac{\partial u}{\partial x_n} = \frac{\partial}{\partial x_n} (\Delta + k^2)u = 0 \quad \text{in } (\omega(\mu * h))_{\text{loc}}.$$

Applying the strong maximum principle in Proposition A.4 on $\frac{\partial u}{\partial x_n}$, we obtain $\frac{\partial u}{\partial x_n} < 0$ in $(\omega(\mu * h))_{\text{loc}}$ (because $\frac{\partial u}{\partial x_n} \not\equiv 0$ in $\omega(\mu * h)$).

Since x_0 can be separated from $\text{supp}(\mu * h)$ by hyperplanes whose normals form an open convex cone, this argument implies that in a neighbourhood of x_0 the function u is decreasing in a cone of directions. We deduce that in a neighbourhood of x_0 the free boundary $\partial\omega(\mu * h)$ is the graph of a Lipschitz function. Since the choice of $x_0 \in \partial\omega(\mu * h)$ was

arbitrary, we conclude that the free boundary $\partial\omega(\mu * h)$ is locally a Lipschitz graph. Using [11–13], we know that $\partial\omega(\mu * h)$ is C^1 , and then from [33] we conclude that $\partial\omega(\mu * h)$ is real-analytic. We also refer to the monograph [23] for the general regularity theory for free boundaries. \square

Appendix A: Auxiliary Propositions

The results in this appendix are well-known, and the proofs can found at arXiv:2204.13934.

A.1 A Real-Valued Fundamental Solution

In this section we give an exact expression for a real-valued radial fundamental solution to the Helmholtz equation. This solution is positive in a ball with suitable radius, which is crucial for our construction of k -quadrature domains.

Proposition A.1 Fix $k > 0$ and $n \geq 2$. For any $R > 0$, let $\tilde{\Phi}_{k,R}$ be given by

$$\tilde{\Phi}_{k,R}(x) = \frac{k^{\frac{n-2}{2}}}{4(2\pi)^{\frac{n-2}{2}} J_{\frac{n-2}{2}}(kR)} |x|^{-\frac{n-2}{2}} \left(Y_{\frac{n-2}{2}}(kR) J_{\frac{n-2}{2}}(k|x|) - J_{\frac{n-2}{2}}(kR) Y_{\frac{n-2}{2}}(k|x|) \right). \tag{A.1}$$

Then the distribution $\tilde{\Phi}_{k,R} \in L^1_{loc}(\mathbb{R}^N)$ is radial, smooth outside the origin and satisfies

$$\begin{cases} (\Delta + k^2)\tilde{\Phi}_{k,R} = -\delta_0 & \text{in } \mathcal{D}'(\mathbb{R}^n), \\ \tilde{\Phi}_{k,R}(x) = 0 & \text{for } x \in \partial B_R(0). \end{cases} \tag{A.2}$$

Furthermore, in the case when $0 < R < j_{\frac{n-2}{2}} k^{-1}$, the distribution $\tilde{\Phi}_{k,R}$ is positive in $B_R(0)$.

A.2 The Mean Value Theorem

Proposition A.2 Let $n \geq 2$ be an integer, and let $R > 0$ be any constant. If $u \in L^1(B_R(x_0))$ is a solution to

$$(\Delta + k^2)u = 0 \quad \text{in } B_R(x_0),$$

then

$$\int_{B_R(x_0)} u(x) dx = c_{n,k,R}^{MVT} u(x_0) \quad \text{with} \quad c_{n,k,R}^{MVT} = (2\pi)^{n/2} \frac{R^{\frac{n}{2}} J_{\frac{n}{2}}(kR)}{k^{\frac{n}{2}}}. \tag{A.3}$$

In addition, if we assume that $0 < R < j_{\frac{n-2}{2},1} k^{-1}$ and $u \in L^1(B_R(x_0))$ is a sub-solution of the Helmholtz equation,

$$(\Delta + k^2)u \geq 0 \quad \text{in } B_R(x_0),$$

then, provided x_0 is a Lebesgue point for u ,

$$\int_{B_R(x_0)} u(x) dx \geq c_{n,k,R}^{MVT} u(x_0) \tag{A.4}$$

with equality if and only if $(\Delta + k^2)u = 0$ in $B_R(x_0)$. In addition, the mapping

$$r \mapsto \frac{1}{c_{n,k,r}^{MVT}} \int_{B_r(x_0)} u(x) \, dx$$

is monotone increasing on $(0, R)$ unless there exists an $0 < R' \leq R$ such that $(\Delta + k^2)u = 0$ in $B_{R'}(x_0)$ in which case the mapping is constant on $(0, R')$ and increasing on (R', R) .

Remark A.3 In particular,

$$\begin{aligned} \text{when } n = 2, \quad c_{2,k,R}^{MVT} &= \frac{2\pi R J_1(kR)}{k}, \\ \text{when } n = 3, \quad c_{3,k,R}^{MVT} &= \frac{4\pi(\sin(kR) - kR \cos(kR))}{k^3}. \end{aligned}$$

Unlike the mean value theorem for harmonic functions, there are radii for which $c_{n,k,R}^{MVT}$ is zero or even negative.

A.3 Maximum Principle

We will need the following (generalized) maximum principle and properties of sub/super-solutions in small domains.

Proposition A.4 Fix $n \geq 2$, let $U \subset \mathbb{R}^n$ be a bounded open set, and let $\lambda_1(U)$ denote the first eigenvalue of the Dirichlet Laplacian on U , that is

$$\lambda_1(U) := \inf_{u \in H_0^1(U)} \frac{\|\nabla u\|_{L^2(U)}^2}{\|u\|_{L^2(U)}^2}. \tag{A.5}$$

Given any $0 < k^2 < \lambda_1(U)$. If $w \in H^1(U)$ satisfies $w|_{\partial U} \leq 0$ (i.e. $w_+ := \max\{w, 0\} \in H_0^1(U)$) and $(\Delta + k^2)w \geq 0$ in the sense of $H^{-1}(U)$, then $w \leq 0$ in U . If we additionally assume that $w \in C(U)$, then in each connected component of U we have either $w < 0$ or $w \equiv 0$.

Proposition A.5 Fix $n \geq 2$, $k > 0$, and let $U \subset \mathbb{R}^n$ be a bounded open set. If $w_1, w_2 \in H^1(U)$ satisfy $(\Delta + k^2)w_j \leq 0$ in the sense of $H^{-1}(U)$ for $j = 1$ and 2 , then the same is true for $w = \min\{w_1, w_2\}$.

Remark A.6 Note that $\lambda_1(B_R) = j_{\frac{n-2}{2},1}^2 R^{-2}$. Therefore, the condition $k^2 < \lambda_1(U)$ is satisfied if $U \subset B_R$ with $0 < R < j_{\frac{n-2}{2},1} k^{-1}$.

Acknowledgments This project was finalized while the authors stayed at Institute Mittag Leffler (Sweden), during the program Geometric aspects of nonlinear PDE. The authors would like to express their gratitude to Lavi Karp and Aron Wennman for helpful comments. We would also like to give special thanks to Björn Gustafsson and the anonymous referee for a careful reading of the manuscript and several detailed suggestions that have improved the presentation. In particular, the comments from the anonymous referee have led to improvements in our results. Kow and Salo were partly supported by the Academy of Finland (Centre of Excellence in Inverse Modelling and Imaging, 312121) and by the European Research Council under Horizon 2020 (ERC CoG 770924). Larson was supported by Knut and Alice Wallenberg Foundation grant KAW 2021.0193. Shahgholian was supported by Swedish Research Council.

Funding Open access funding provided by Royal Institute of Technology.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Adams, D.R., Hedberg, L.I.: Function spaces and potential theory volume 314 of Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer-Verlag, Berlin (1996). MR1411441, <https://doi.org/10.1007/978-3-662-03282-4>
- Ahlfors, L.V.: Finitely generated kleinian groups. *Amer. J. Math.* **86**, 413–429 (1964). MR0167618, <https://doi.org/10.2307/2373173>
- Bers, L.: An approximation theorem. *J. Analyse Math.* **14**, 1–4 (1965). MR0178287, <https://doi.org/10.1007/BF02806376>
- Blåsten, E.: Nonradiating sources and transmission eigenfunctions vanish at corners and edges. *SIAM J. Math. Anal.* **50**(6), 6255–6270 (2018). MR3885754, <https://doi.org/10.1137/18M1182048>, arXiv: 1803.10917
- Blåsten, E., Liu, H.: On vanishing near corners of transmission eigenfunctions. *J. Funct. Anal.* **273**(11), 3616–3632 (2017). MR3706612, <https://doi.org/10.1016/j.jfa.2017.08.023>, arXiv: 1701.07957, Addendum: arXiv: 1710.08089
- Blåsten, E., Liu, H.: On corners scattering stably and stable shape determination by a single far-field pattern. *Indiana Univ. Math. J.* **70**(3), 907–947 (2021). MR4284101, <https://doi.org/10.1512/IUMJ.2021.70.8411>, arXiv: 1611.03647
- Blåsten, E., Päivärinta, L., Sylvester, J.: Corners always scatter. *Comm. Math. Phys.* **331**(2), 725–753 (2014). MR3238529, <https://doi.org/10.1007/s00220-014-2030-0>
- Brezis, H.: Functional analysis, Sobolev spaces and partial differential equations. Universitext. Springer, New York (2011). MR2759829, <https://doi.org/10.1007/978-0-387-70914-7>
- Brown, L., Kahane, J.-P.: A note on the Pompeiu problem for convex domains. *Math. Ann.* **259**(1), 107–110 (1982). MR0656655, <https://doi.org/10.1007/BF01456832>
- Brown, L., Schreiber, B.M., Taylor, B.A.: Spectral synthesis and the Pompeiu problem. *Ann. Inst. Fourier (Grenoble)* **23**(3), 125–154 (1973). MR0352492, <https://doi.org/10.5802/aif.474>
- Caffarelli, L.A.: The regularity of free boundaries in higher dimensions. *Acta Math.* **139**(3–4), 155–184 (1977). MR0454350, <https://doi.org/10.1007/BF02392236>
- Caffarelli, L.A.: Compactness methods in free boundary problems. *Comm. Partial Differential Equations* **5**(4), 427–448 (1980). MR0567780, <https://doi.org/10.1080/0360530800882144>
- Caffarelli, L.A.: The obstacle problem revisited. *J. Fourier Anal. Appl.* **4**(4–5), 383–402 (1998). MR1658612, <https://doi.org/10.1007/BF02498216>
- Cakoni, F., Colton, D., Haddar, H.: Inverse scattering theory and transmission eigenvalues volume 88 of regional conference series in applied mathematics society for industrial and applied mathematics (SIAM), Philadelphia, PA. MR3601119, <https://doi.org/10.1137/1.9781611974461> (2016)
- Cakoni, F., Vogelius, M.S.: Singularities almost always scatter: Regularity results for non-scattering inhomogeneities arXiv: 2104.05058 (2021)
- Cakoni, F., Xiao, J.: On corner scattering for operators of divergence form and applications to inverse scattering. *Comm. Partial Differential Equations* **46**(3), 413–441 (2021). MR4232500, <https://doi.org/10.1080/03605302.2020.1843489>, arXiv: 1905.02558
- Colton, D., Kress, R.: Inverse acoustic and electromagnetic scattering theory, volume 93 of Applied Mathematical Sciences. Springer, Cham, fourth edition. MR3971246, <https://doi.org/10.1007/978-3-030-30351-8> (2019)
- Courant, R., Hilbert, D.: Methods of mathematical physics (vol II). Wiley Classics Library. A Wiley-Interscience Publication John Wiley & Sons, Inc. New York Reprint of the 1989 original. MR1013360, <https://doi.org/10.1002/9783527617234> (1962)
- Davis, P.J.: The Schwarz function and its applications, volume 17 of The Carus Mathematical Monographs. The Mathematical Association of America. Buffalo, N.Y. (1974). MR0407252, <https://doi.org/10.5948/9781614440178>

20. Deng, J.: Some results on the Schiffer conjecture in \mathbb{R}^2 . *J. Differential Equations* **253**(8), 2515–2526 (2012). MR2950461, <https://doi.org/10.1016/j.jde.2012.06.002>, arXiv: 1112.0207
21. Elschner, J., Hu, G.: Corners and edges always scatter. *Inverse Prob.* **31**(1), 015003 (2015). 17 pp, MR3302364, <https://doi.org/10.1088/0266-5611/31/1/015003>
22. Friedman, A.: On n -metaharmonic functions and harmonic functions of infinite order. *Proc. Amer. Math. Soc.* **8**, 223–229 (1957). MR0085430, <https://doi.org/10.2307/2033716>
23. Friedman, A.: Variational principles and free-boundary problems Robert E. Krieger Publishing Co., Inc., Malabar, FL, second edition, MR1009785 (1988)
24. Gardiner, S., Sjödin, T.: Partial balayage and the exterior inverse problem of potential theory. In: Potential theory and stochastics in Albac, vol 11, pp 111–123, Theta, Bucharest, Theta Ser. Adv. Math. MR2681841 (2009)
25. Gilbarg, D., Trudinger, N.S.: Elliptic partial differential equations of second order (reprint of the 1998 edition), volume 224 of Classics in Mathematics. Springer-Verlag, Berlin Heidelberg (2001). MR1814364, <https://doi.org/10.1007/978-3-642-61798-0>
26. Gustafsson, B.: On quadrature domains and an inverse problem in potential theory. *J. Analyse Math.* **55**, 172–216 (1990). MR1094715, <https://doi.org/10.1007/BF02789201>
27. Gustafsson, B.: Lectures on balayage. In: Clifford algebras and potential theory, vol 7 of Univ. Joensuu Dept. Math. Rep. Ser., pp 17–63. Univ. Joensuu, Joensuu, MR2103705, diva2:492834 (2004)
28. Gustafsson, B., Roos, J.: Partial balayage on Riemannian manifolds. *J. Math. Pures Appl.* **118**(9), 82–127 (2018). MR3852470, <https://doi.org/10.1016/j.matpur.2017.07.013>, arXiv:1605.03102
29. Gustafsson, B., Shapiro, H.S.: What is a quadrature domain? In: Quadrature domains and their applications, vol 156 of Oper. Theory Adv. Appl., pp 1–25. Birkhäuser, Basel (2005). MR2129734, https://doi.org/10.1007/3-7643-7316-4_1
30. Hedberg, L.L.: Approximation in the mean by solutions of elliptic equations. *Duke Math. J.* **40**, 9–16 (1973). MR0312071, <https://doi.org/10.1215/S0012-7094-73-04002-7>
31. Isakov, V.: Inverse source problems, volume 34 of Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI. MR1071181, <https://doi.org/10.1090/surv/034> (1990)
32. Karp, L., Margulis, A.: Newtonian potential theory for unbounded sources and applications to free boundary problems. *J. Anal. Math.* **70**, 1–63 (1996). MR1444257, <https://doi.org/10.1007/BF02820440>
33. Kinderlehrer, D., Nirenberg, L.: Regularity in free boundary problems. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)* **4**(2), 373–391 (1977). MR0440187, NUMDAM
34. Kinderlehrer, D., Stampacchia, G.: An introduction to variational inequalities and their applications, volume 31 of Classics in Applied Mathematics. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2000. Reprint of the original. MR1786735, <https://doi.org/10.1137/1.9780898719451> (1980)
35. Kuznetsov, N.: Mean value properties of harmonic functions and related topics (a survey). *J. Math. Sci. (N.Y.)* **242**, 177–199 (2019). MR4002408, <https://doi.org/10.1007/s10958-019-04473-w>, arXiv: 1904.08312
36. Kuznetsov, N.: Metaharmonic functions: mean flux theorem, its converse and related properties. *Algebra i Analiz* **33**(2), 82–98 (2021). MR4240827, Mi aa1749, arXiv: 2004.03433
37. Lee, S.Y., Makarov, N.G.: Topology of quadrature domains. *J. Amer. Math. Soc.* **29**(2), 333–369 (2016). MR3454377, <https://doi.org/10.1090/jams828>, arXiv: 1307.0487
38. Päivärinta, L., Salo, M., Vesalainen, E.V.: Strictly convex corners scatter. *Rev. Mat Iberoam.* **33**(4), 1369–1396 (2017). MR3729603, <https://doi.org/10.4171/RMI/975>, arXiv: 1404.2513
39. Petrosyan, A., Shahgholian, H., Uraltseva, N.: Regularity of free boundaries in obstacle-type problems, volume 136 of Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, MR2962060, <https://doi.org/10.1090/gsm/136> (2012)
40. Pompeiu, P.: Sur certains systèmes d'équations linéaires et sur une propriété intégrale des fonctions de plusieurs variables. *Comptes Rendus de l'académie des Sciences. série I, Mathématique* **188**, 1138–1139 (1929)
41. Rudin, W. Real and complex analysis, 3rd edition. McGraw-Hill Book Co, New York (1987). MR0924157
42. Sakai, M.: Applications of variational inequalities to the existence theorem on quadrature domains. *Trans. Amer. Math. Soc.* **276**(1), 267–279 (1983). MR0684507, <https://doi.org/10.1090/S0002-9947-1983-0684507-2>
43. Sakai, M.: Solutions to the obstacle problem as green potentials. *J. Analyse Math.* **44**, 97–116 (1984). MR0801289, <https://doi.org/10.1007/BF02790192>
44. Sakai, M.: Regularity of a boundary having a schwarz function. *Acta Math.* **166**(3–4), 263–297 (1991). MR1097025, <https://doi.org/10.1007/BF02398888>

45. Salo, M., Shahgholian, H.: Free boundary methods and non-scattering phenomena. Res. Math. Sci. vol 8(4) Paper No.58, MR4323345, <https://doi.org/10.1007/s40687-021-00294-z>, arXiv: 2106.15154 (2021)
46. Sjödin, T.: On the structure of partial balayage. Nonlinear Anal. **67**(1), 94–102 (2007). MR2313881, <https://doi.org/10.1016/j.na.2006.05.001>
47. Trèves, F.: Basic linear partial differential equations, volume 62 of Pure and Applied Mathematics. Academic Press Harcourt Brace Jovanovich, Publishers, New York-London (1975). MR0447753
48. Weber, H.: Ueber einige bestimmte Integrale. J. Reine Angew. Math. **69**, 222–237 (1868). MR1579416, <https://doi.org/10.1515/crll.1868.69.222>
49. Weber, H.: Ueber die integration der partiellen differentialgleichung: $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + k^2 u = 0$. Math. Ann. **1**(1), 1–36 (1869). MR1509609, <https://doi.org/10.1007/BF01447384>
50. Williams, S.A.: A partial solution of the Pompeiu problem. Math. Ann. **223**(2), 183–190 (1976). MR0414904, <https://doi.org/10.1007/BF01360881>
51. Williams, S.A.: Analyticity of the boundary for Lipschitz domains without the Pompeiu property. Indiana Univ. Math. J. **30**(3), 357–369 (1981). MR0611225
52. Yafaev, D.R.: Mathematical scattering theory. Analytic theory, volume 158 of Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, MR2598115, <https://doi.org/10.1090/surv/158> (2010)
53. Zalcman, L.: A bibliographic survey of the Pompeiu problem. In: Approximation by solutions of partial differential equations (Hanstholm, 1991), vol 365 of NATO Adv. Sci. Inst. Ser. C: Math. Phys. Sci., pp 185–194. Kluwer Acad. Publ., Dordrecht, MR1168719 (1992)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.