

# Outbreak of OXA-48-producing Enterobacteriaceae in a neonatal intensive care unit in Western Sweden

Downloaded from: https://research.chalmers.se, 2024-05-03 12:01 UTC

Citation for the original published paper (version of record):

Hallbäck, E., Johnning, A., Myhrman, S. et al (2023). Outbreak of OXA-48-producing Enterobacteriaceae in a neonatal intensive care unit in Western Sweden. European Journal of Clinical Microbiology and Infectious Diseases, 42(5): 597-605. http://dx.doi.org/10.1007/s10096-023-04584-y

N.B. When citing this work, cite the original published paper.

research.chalmers.se offers the possibility of retrieving research publications produced at Chalmers University of Technology. It covers all kind of research output: articles, dissertations, conference papers, reports etc. since 2004. research.chalmers.se is administrated and maintained by Chalmers Library

#### **ORIGINAL ARTICLE**



# Outbreak of OXA-48-producing *Enterobacteriaceae* in a neonatal intensive care unit in Western Sweden

Erika Tång Hallbäck<sup>1,2</sup> · Anna Johnning<sup>3,4,5</sup> · Sofia Myhrman<sup>6</sup> · Marie Studahl<sup>1,7</sup> · Elisabet Hentz<sup>8,9</sup> · Anders Elfvin<sup>8,9</sup> · Ingegerd Adlerberth<sup>1,2,5</sup>

Received: 21 January 2023 / Accepted: 28 February 2023 © The Author(s) 2023

#### Abstract

In 2015, an outbreak caused by OXA-48-producing *Enterobacteriaceae* affected a neonatal intensive care unit at a Swedish University Hospital. The aim was to explore the transmission of OXA-48-producing strains between infants and the transfer of resistance plasmids between strains during the outbreak. Twenty-four outbreak isolates from ten suspected cases were whole-genome sequenced. A complete assembly was created for the index isolate (*Enterobacter cloacae*) and used as a mapping reference to detect its plasmids in the remaining isolates (17 *Klebsiella pneumoniae*, 4 *Klebsiella aerogenes*, and 2 *Escherichia coli*). Strain typing was performed using core genome MLST and SNP analysis. As judged from sequencing and clinical epidemiological data, the outbreak involved nine cases (two developed sepsis) and four OXA-48-producing strains: *E. cloacae* ST1584 (index case), *K. pneumoniae* ST25 (eight cases), *K. aerogenes* ST93 (two cases), and *E. coli* ST453 (2 cases). Two plasmids from the index strain, p*Ecl*A2 and p*Ecl*A4, carrying *bla*<sub>OXA48</sub> and *bla*<sub>CMY4</sub>, respectively, were traced to all *K. pneumoniae* ST25 isolates. *Klebsiella aerogenes* ST93 and *E. coli* ST453 harboured either only p*Ecl*A2, or both p*Ecl*A2 and p*Ecl*A4. One suspected case harbouring OXA-162-producing *K. pneumoniae* ST37 could be excluded from the outbreak. Once initiated by an *E. cloacae* strain, the outbreak was caused by the dissemination of a *K. pneumoniae* ST25 strain and involved inter-species horizontal transfer of two resistance plasmids, one of which carried *bla*<sub>OXA-48</sub>. To our knowledge, this is the first description of an outbreak of OXA-48-producing *Enterobacteriaceae* in a neonatal setting in northern Europe.

Keywords  $Enterobacteriaceae \cdot OXA-48 \cdot Antibiotic resistance \cdot Whole-genome sequencing \cdot New-born infant \cdot Neonatal intensive care \cdot Outbreak \cdot Plasmid transmission$ 

Erika Tång Hallbäck erika.tang.hallback@gu.se

- <sup>1</sup> Department of Infectious Diseases, Institute of Biomedicine, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden
- <sup>2</sup> Department of Clinical Microbiology, Sahlgrenska University Hospital, Region Västra Götaland, Gothenburg, Sweden
- <sup>3</sup> Department of Mathematical Sciences, Chalmers University of Technology and University of Gothenburg, Gothenburg, Sweden
- <sup>4</sup> Department of Systems and Data Analysis, Fraunhofer-Chalmers Centre, Gothenburg, Sweden

- <sup>5</sup> Centre for Antibiotic Resistance Research in Gothenburg (CARe), Gothenburg, Sweden
- <sup>6</sup> Department of Clinical Microbiology, Infection Control Unit, Sahlgrenska University Hospital, Region Västra Götaland, Gothenburg, Sweden
- <sup>7</sup> Department of Infectious Diseases, Sahlgrenska University Hospital, Region Västra Götaland, Gothenburg, Sweden
- <sup>8</sup> Department of Pediatrics, Institute of Clinical Sciences, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden
- <sup>9</sup> Department of Pediatrics, The Queen Silvia Children's Hospital, Sahlgrenska University Hospital, Region Västra Götaland, Gothenburg, Sweden

During the last decades, there has been a worrying rise in Gram-negative intestinal bacteria producing carbapenemases, i.e. b-lactamases that hydrolyse carbapenems and usually also other b-lactam antibiotics. These bacteria are particularly troubling since they are often resistant to several additional classes of antibiotics which further limits available treatment options. The incidence of carbapenem-producing Enterobacteriaceae is low but increasing in Sweden: from 0.2 to 1.3 cases/100,000 inhabitants between 2012 and 2021 [1]. Some of the most frequently encountered carbapenemases are those of the OXA-48 group [2]. These enzymes hydrolyse carbapenems only at a low level and show weak activity against cephalosporins, but high-level resistance to carbapenems can be achieved when combined with permeability defects and/or co-expression of extended-spectrum b-lactamases (ESBLs) or AmpC b-lactamases [3, 4]. Carbapenemases of the OXA-48 group, hence, constitute a threat to our ability to treat bacterial infections and it is, therefore, important to understand how *bla*<sub>OXA-48-like</sub> genes are disseminated in bacterial populations.

Neonatal care units (NICUs) frequently experience spread of Enterobacteriaceae [5] and outbreaks involving OXA-48-producing Enterobacteriaceae strains in this setting have been reported [6-8]. Due to sparse colonization by anaerobic bacteria during the first weeks of life, members of the Enterobacteriaceae family establish and reach high abundance levels in the neonatal gut. As the gut microbiota matures, the proliferation of Enterobacteriaceae and other facultative anaerobic bacteria is suppressed [9]. The acquisition of anaerobes is severely delayed in preterm infants, partly related to comprehensive antibiotic treatments, rendering this population prone to heavy colonization by Enterobac*teriaceae* [10]. A high concentration of these bacteria in the gut lumen may favour bacterial conjugation and plasmid transfer, which requires close contact between bacterial cells[11–13]. The intestines of preterm infants could therefore represent a hotspot for horizontal transfer of bla<sub>OXA-48</sub> and other plasmid-borne resistance genes, as also indicated by a recent experimental study [14].

In this study, we describe an outbreak in a NICU at a Swedish University Hospital. The outbreak involved ten suspected cases and four different OXA-48-producing *Enterobacteriaceae* species: *Enterobacter cloacae*, *Klebsiella pneumoniae*, *Klebsiella aerogenes*, and *Escherichia coli*. We used whole-genome sequencing to identify both clonal spread between cases and the dissemination of plasmids, originating in the index strain (*E. cloacae*), to other strains. The results strongly indicate horizontal transfer in vivo between both species and genera—of two different plasmids encoding the carbapenemase OXA-48 and the AmpC-type b-lactamase CMY-4, respectively.

# **Materials and methods**

## The setting, data collection, and definitions

The outbreak occurred in 2015 in an NICU of a tertiary care University Hospital in Western Sweden. Comprised of 14 beds allocated to six rooms, the ward provides specialist care for approximately 1000 term and preterm infants (born from gestational week 22 + 0) annually. In 2015, rectal screening cultures were obtained from all patients at admission to the ward and weekly thereafter. If carriage of ESBL- or carbapenemase-producing bacteria was suspected or confirmed by the Clinical Microbiology laboratory, the patient was cared for in a single-bed room by a dedicated nurse.

Details on contact tracing and sampling were retrospectively collected from the patient charts and reports from the infection prevention and control team. A suspected case was defined as a patient within the NICU from whom an *Enterobacteriaceae* strain positive for carbapenemases of the OXA-48 group was isolated. The outbreak period was defined from the identification of the first suspected case i.e. the sampling date of the case's first positive isolate—to the identification of the last suspected case. No additional infants colonized or infected by *Enterobacteriaceae* positive for carbapenemases of the OXA-48-group were identified at the NICU past this period within the following year. (Fig. 1, Supplementary Fig. S1).

The study was approved by the Regional Ethical Review Board in Gothenburg, Sweden (no. 997–18 and no. 228–15).

# Culturing, bacterial identification and characterization, and selection of isolates

During the outbreak period, ten infants were colonized and/or infected by Enterobacteriaceae positive for OXA-48-like carbapenemases. The methods used for culturing, bacterial identification, antimicrobial susceptibility testing, and identification of resistance determinants were those routinely applied in the clinical microbiology laboratory at the University Hospital (Supplementary Material). A total of 155 rectal screening samples were obtained from the ten suspected outbreak cases during and after the outbreak period, of which 78 were assessed as positive for Enterobacteriaceae bacteria producing carbapenemases of the OXA-48 group (Supplementary Figure S1). In addition, four clinical samples-from two of the suspected cases (two blood cultures, two nasopharyngeal cultures) ---were positive for such bacteria (Supplementary Figure S1). In total, 116 isolates assessed as positive for carbapenemases of the OXA-48 group were obtained from the 82 positive samples and 74/116 isolates were saved and stored frozen ( $-80 \,^{\circ}C$ ) at the Clinical Microbiology laboratory.



Fig. 1 Timeline of the outbreak with OXA-48-producing *Enterobacteriaceae* in a neonatal unit. Isolation day, species identity and carriage of the outbreak plasmids are shown for the isolates selected from the ten suspected cases, with the index isolate (Case 1) set as day 0. Twin pair one (\*) and twin pair two (\*\*) are marked with asterisks. Horizontal lines represent the duration of hospital care, and their colour indicates the ward caring for each case until case identifi-

For this study, we selected 24 of the stored isolates with confirmed carriage of genes encoding a carbapenemase of the OXA-48 group. For each suspected case, the first identified isolate of each OXA-48-positive species colonizing that infant was selected (n = 14). For Case 2, the first identified isolate was a blood culture isolate, and both the blood isolate and the rectal screening isolate from the following day were selected for analysis. Furthermore, from three persistently colonized cases, nine isolates were selected that showed deviations in phenotypic resistance tests and/or meropenem MIC-values from previous isolates of that species in the colonized infant. In total, 1 *E. cloacae*, 17 K. *pneumoniae*, 4 K. *aerogenes*, and 2 *E. coli* isolates were selected for wholegenome sequencing. (Table 1, Supplementary Figure S1).

#### DNA extraction and whole-genome sequencing

Genomic DNA was extracted using the QIAamp DNA Mini Kit (Qiagen) for short-read sequencing and the Marmur protocol for long-read sequencing [15]. DNA quality and quantity were measured with a Nanodrop<sup>TM</sup> ND-1000 Spectrophotometer (Thermo Fisher) and a Qubit 2.0 Fluorometer (Invitrogen), respectively. The species identity was confirmed by 16S rRNA Sanger sequencing (GATC Biotech). The isolates were whole genome sequenced using either IonTorrent (isolates *Ecl1A*, *Kpn2A2*, *Kpn5B*, *Kpn6A*, *Kpn7C*, *Kpn8A1*, *Kpn9A2*, *Kpn10A1*) or Illumina (isolates *Kpn2A1*, *Kpn3A1*, *Kae3B2*, *Kpn3B3*, *Kpn3B4*, *Kae3D5*, *Kae3D6*, *Kpn4A*, *Eco8A2*, *Kpn8A3*, *Kpn8E4*, *Kae9A1*, *Kpn10A2*, *Kpn10A3*, *Eco10A4*, *Kpn10E5*) technology. The

cation: red = Ward A, blue = Ward B, black = Ward C, orange = Ward D. Symbols represent species: star = E. cloacae, square = K. *pneumoniae*, triangle = K. *aerogenes*, and circle = E. *coli*. Carriage of outbreak plasmids is indicated by symbol colour: green = carriage of *pEclA2*, yellow = carriage of *pEclA4*, white = no carriage of outbreak plasmids

index isolate (*Ecl1A*) was also sequenced by Nanopore (full details in Supplementary material). All sequenced isolates were deposited at the Culture Collection University of Gothenburg (CCUG; Table 1).

#### Sequence data analysis

The Illumina sequence data were trimmed from adapters and low-quality sequences using Trim galore! (www.bioin formatics.babraham.ac.uk/projects/trim\_galore/; v. 0.4.3, parameters -stringency 3 -paired -retain unpaired). The shortread data for all but the index isolate was assembled using SPAdes (v.3.14.1, Illumina parameters: -isolate -cov-cutoff 5; IonTorrent parameter: -iontorrent -k 21,33,55,77,99,127 -isolate -cov-cutoff 5) [16]. The short- and long-read data for the index isolate were assembled using Unicycler (v. 0.4.4, default parameters) [17] which produced complete circular assemblies of the chromosome and all plasmids. This Whole Genome Shotgun project has been deposited at DDBJ/ ENA/GenBank under the accession JAGPNS00000000-JAGPOP000000000. The versions described in this paper are versions JAGPNS010000000.1- JAGPOP010000000.1 (Table 1).

The assemblies were annotated for antimicrobial resistance genes using ResFinder followed by manual inspection (v. 4.1, 90% identity, 20% coverage) [18] and the incompatibility groups of the index isolate plasmids were identified using PlasmidFinder (v. 2.1, default parameters). To investigate the transfer of the index isolate's plasmids, the reads of the remaining isolates were mapped to the plasmid

Table 1 Iso	lates pos	itive for carbapene	emases of ti	he OXA-48 grou	p selected	for whole-ge	nome sequencing											
Isolate ID <sup>a</sup>	Case	Species	Location	Isolation day <sup>b</sup>	Ward $^{\rm c}$	CCUG no	NCBI accession no	MIC <sup>d</sup>	Resi	stance p	henoty	pes <sup>e</sup>					β-lac	tamases <sup>f</sup>
								MP	CT	CAZ	MP	TZP	TM	CI ,	LS ]	P C	0 ESB	AmpC
$Ecl1A^*$	1	E. cloacae	Faeces	0	A	67,822	JAGPNS0000000000	4	R	R	I	R	1	R	- 	S	0	1
Kpn2A1	5	K. pneumoniae	Blood	17	A	67,824	JAGPNT00000000000	0.125	R	R	S	R	S	S	~	1	1	1
Kpn2A2	2	K. pneumoniae	Faeces	18	A	67,823	JAGPNU000000000000000000000000000000000000	1	R	R	S	R	S	S	~	s	1	1
Kpn3A1	б	K. pneumoniae	Faeces	18	A	75,156	JAGPNV000000000000000000000000000000000000	2	R	R	S	R	S	S	~	s	1	I
Kae3A2	б	K. aerogenes	Faeces	28	A	67,825	JAGPNW000000000	32	R	I	R	R	S	S		ו צ	I	I
Kpn3B3	ŝ	K. pneumoniae	Faeces	73	В	75,157	JAGPNX0000000000	32	R	R	R	R	S	S	~	1	I	1
Kpn3B4	б	K. pneumoniae	Faeces	73	В	75,158	JAGPNY0000000000	2	R	R	S	R	S	S	~	1	1	1
Kae3D5	б	K. aerogenes	Faeces	130	D	75,159	JAGPNZ0000000000	8	I	S	I	R	S	S	~	S	1	1
Kae3D6	б	K. aerogenes	Faeces	130	D	75,160	JAGPOA00000000000	4	К	R	I	R	S	S	~	S	1	1
Kpn4A	4	K. pneumoniae	Faeces	18	А	75,168	JAGPOB00000000000	1	R	R	S	R	S	S	č	1	1	1
Kpn5B	5	K. pneumoniae	Faeces	18	В	67,830	JAGPOC0000000000	0.25	R	R	S	R	S	S	č	1	1	1
Kpn6A	9	K. pneumoniae	Faeces	22	в	67,831	JAGPOD000000000000000000000000000000000000	1	Я	R	S	R	S	S	í n	S	1	1
Kpn7C	7	K. pneumoniae	Faeces	29	C	74,849	JAGPOE000000000000000000000000000000000000	2	R	R	I	R	S	S	0		1	I
Kpn8A1	8	K. pneumoniae	Faeces	35	A	67,826	JAGPOF0000000000	1	Я	R	S	R	S	S	č	1	1	1
Eco8A2	8	$E. \ coli$	Faeces	74	А	75,163	JAGPOG00000000000000000000000000000000000	0.5	Я	R	S	R	S	S	M	1	Ι	1
Kpn8A3	8	K. pneumoniae	Faeces	145	А	75,161	JAGPOH0000000000	32	Я	R	R	R	S	s	M	1	I	1
Kpn8E4	8	K. pneumoniae	Faeces	433	Е	75,162	JAGPOI000000000	4	Я	R	I	R	S	s	0	' _	Ι	1
Kae9A1	6	K. aerogenes	Faeces	39	А	67,827	JAGPOJ0000000000	32	К	I	R	R	S	S	2	RS	I	I
Kpn9A2	6	K. pneumoniae	Faeces	39	А	67,828	JAGPOK000000000000000000000000000000000000	0.25	R	R	S	R	S	S	~	S	1	1
Kpn10A1	10	K. pneumoniae	Faeces	49	Α	67,829	JAGPOL0000000000	0.25	R	R	S	R	S	S	~	1	1	1
Kpn10A2	10	K. pneumoniae	Faeces	67	A	75,164	JAGPOM000000000000000000000000000000000000	4	R	R	I	R	S	S	0	S	1	1
Kpn10A3	10	K. pneumoniae	Faeces	67	A	75,165	JAGPON0000000000	16	R	R	R	R	S	s	~	s S	I	1
Ecol0A4	10	E. coli	Faeces	88	Α	75,166	JAGPOO00000000000000000000000000000000000	0.125	S	S	S	R	S	S	0	S	Ι	I
Kpn10E5	10	K. pneumoniae	Faeces	433	ш	75,167	JAGPOP0000000000	7	R	R	S	R	S	S	~		1	1
<sup>a</sup> The index cared for wh cefotaxime (	isolate nen the s (CT), ce	is marked with an ample yielding the fitazidime (CAZ),	asterisk (* e isolate wa meropenen	). <sup>b</sup> The day who is taken: Ward A, 1 (MP), piperacil	en the sam , Ward B, <sup>1</sup> llin-tazoba	ple yielding Ward C, War ctam (TZP),	the isolate was taken. ' I D, and Ward E. <sup>d</sup> Min tobramycin (TM), cipro	The index timum inh offoxacin	i samj nibitor (CI),	ple date y conce trimetho	is set ntratio prim/s	o day ( n of me ulfame	). <sup>c</sup> The ropene hoxaz	e hosj m. <sup>e</sup> J ole (T	pital v Pheno S), ir	vard w typic r nipene	here the esistance m (IP), a	infant was results for nd colistin
(CU). Pher	notypic t	est results for ESE	SL and Amp	oc production: 1	= positive,	U=negative.	, and $=$ not tested											

assemblies in the following way. First, the IonTorrent and the trimmed Illumina data were error-corrected using SPAdes (v 0.4.3, parameters: Illumina, -only-error-correction; IonTorrent, -only-error-correction -iontorrent) [16]. The trimmed and error-corrected paired reads were mapped to each of the five plasmids of the index isolate using BowTie2 (v. 2.3.0, default parameters) [19] and the coverage was calculated for each position using samtools (v 1.6) [20]. The complete plasmid profiles of isolates other than the index isolate were not investigated.

Epidemiological typing of all sequenced isolates was performed using MLST schemes from PubMLST.org [21]. All *K. pneumoniae* sequences were also submitted to the Institute Pasteur MLST-2.0 server and analysed with the BIGSdb database for curation and publication of core genome MLST [22]. Epidemiological strain typing of *E. coli* and *K. aerogenes* isolates was done by SNP analysis using the references *E. coli* IAI1 (accession number ENA\_CU928160 and *K. aerogenes* KCTC 2190 (NC\_015663.1) in CLC Genomics Workbench 12 (Qiagen; full details in Supplementary materials).

# Results

#### **Outbreak description**

The outbreak timeline and the follow-up period are shown in Fig. 1 and Supplementary Figure S1. During the outbreak, *Enterobacteriaceae* strains producing carbapenemases of the OXA-48 group were isolated from two twin pairs and six singlets, with a mean gestational age of 27 weeks (clinical characteristics of the 10 preterm infants are shown in Supplementary Table S1). All suspected cases except the index case (Case 1) yielded a *K. pneumoniae* isolate positive for OXA-48-like carbapenemases at the time of case identification. Case 2 was identified from a positive blood culture and the others were identified by rectal screening.

The index case was identified at Ward A, where an *E. cloacae* isolate positive for the OXA-48 group (*Ecl*1A) was detected in a rectal screening culture collected on day 0 (Table 1, Fig. 1, Supplementary Fig. S1). Due to overcrowding, the patient was initially cared for in a multi-bedroom by a dedicated nurse before being transferred to a single-bed room on day 5. Since isolation routines had not been followed, rectal screening was enhanced to twice weekly from day 5, including all neonates at the ward. The index case remained on Ward A for 10 days, but no follow-up cultures were obtained. The second case was identified on day 17 on Ward A, by a blood culture yielding a *K. pneumoniae* isolate positive for the OXA-48 group (*Kpn*2A1). Rectal screening the following day detected gut colonization by similar

OXA-48-producing *K. pneumoniae* in Cases 2–5. At this time point, Case 5 had been transferred to a different unit (Ward B), where Case 6 was discovered 4 days later (day 22). On day 29, Case 7 was identified at a third unit (Ward C). Three additional cases (8–10) were identified on Ward A between days 35 and 49: Cases 8 and 10 were a twin pair and Case 9 was the twin sibling of Case 3. Thus, all suspected cases except for Case 7 had a spatial link to either Ward A or Ward B and/or another case.

Screening cultures from the two twin pairs also yielded gut colonization with other enterobacterial species positive for OXA-48-like carbapenemases. *Klebsiella aerogenes* was isolated from Case 3 (*Kae*3B2) on day 28 and 11 days later from the sibling, Case 9, who, thus, harboured both *K. aerogenes* (*Kae*9A1) and *K. pneumoniae* (*Kpn*9A2) strains positive for the OXA-48 group at the time of case identification. A positive *E. coli* strain was first isolated from Case 8 (*Eco*8A2) on day 74, i.e. after the outbreak period, and 14 days later from its sibling, Case 10 (*Eco*10A4).

The infection prevention and control team identified overcrowding, deficient routine practices for milk preparation, inadequate handling of shared medical equipment, inexperienced healthcare workers, and cluttering of non-medical utensils in the patient areas as risk factors for continued transmission. Hence, control measures were focused on adherence to existing routines for infection prevention and control and included the education of both healthcare workers and parents. The extended rectal screening program was continued until day 60.

Two infants, Case 2 and Case 8, developed sepsis at 28 and 39 days of age, respectively, with blood cultures yielding *K. pneumoniae* positive for OXA-48-like carbapenemases. Of the six cases that were screened more than one year after the outbreak, three were still colonized by OXA-48-producing *Enterobacteriaceae*: two with *K. pneumoniae* (Cases 8 and 10) and one with *K. aerogenes* (Case 3). Later follow-up samples from these infants were negative (Supplementary Figure S1).

### The index isolate and its plasmids

The index isolate *E. cloacae* CCUG 67,822 (*Ecl*1A) belonged to a novel sequence type (ST), ST1584 [23]. All its plasmids, except for the smallest one (p*Ecl*A5), harboured antimicrobial resistance genes (Table 2). The  $bla_{OXA-48}$  gene was located as the only resistance gene on a 64 kb long IncL plasmid (p*Ecl*A2), which was near identical (two substitutions, both non-synonymous, in *traN* and *traX*) to the wide-spread OXA-48-encoding plasmid characterized by Hamprecht et al. [24]. This plasmid locates  $bla_{OXA-48}$  between two identical insertion sequences, IS1999, members of the transposon (Tn)1999 family [2].

Plasmid ID	Size (bp)	Multiplicity <sup>a</sup>	Inc group	Antimicrobial resistance genes	Closest GenBank match [%nt identity/%coverage]
pEclA1	202 089	1.88x	IncF	bla <sub>TEM-1B</sub> , aac(3)-IIa, qnrS1, sul2	CP058188.1 [98.84/55]
pEclA2	63 589	2.11x	IncL	$bla_{\rm OXA-48}$	LR025098.1 [100/100]
pEclA3	61 783	2.00x	IncR	sul1, aadA1, dfrA15, tet(D), catA2, bla <sub>TEM-1D</sub>	LR697132.1 [99.71/58]
p <i>Ecl</i> A4	6 926	9.24x	IncQ	$bla_{\rm CMY-4}$	KP205272.1 [99.94/100]
pEclA5	3 019	8.59x	-	_	CP050768.1 [100/98.54]

Table 2 Plasmids in the index isolate Enterobacter cloacae CCUG 67822 (isolate ID Ecl1A)

<sup>a</sup> Plasmid multiplicity relative to the chromosome

The plasmid pEclA4 also carried a single resistance gene, bla<sub>CMY-4</sub>, and was almost identical (one substitution and one insertion, both in an intergenic region) to the CMY-4-encoding IncQ plasmid described by Kotsakis et al. [25]. The largest plasmid, pEclA1, carried genes encoding resistance towards  $\beta$ -lactams (*bla*<sub>TEM-1</sub>), sulphonamides (sul2), aminoglycosides (aac(3)-IIa), and fluoroquinolones (qnrS1). Finally, the plasmid pEclA3 harboured resistance genes towards  $\beta$ -lactams ( $bla_{\text{TEM-1}}$ ), sulphonamides (sul1) aminoglycosides (aadA1), trimethoprim (dfrA15), tetracyclines (*tet*(D)), and chloramphenicol (*catA2*; Table 2).

# Detection of plasmids pEcIA2 and pEcIA4 in other outbreak isolates

By mapping the short reads of the 23 sequenced non-index isolates onto the complete plasmid sequences of the index isolate, we could conclude that all isolates harboured a plasmid resembling pEclA2, i.e. the  $bla_{OXA-48}$  plasmid (Fig. 1,

Table 3 Molecular typing of the sequenced isolates	Isolate ID <sup>a</sup>	MLST <sup>b</sup>	OXA- 48/ CMY-4 <sup>c</sup>	Additional antimicrobial resistance genes
	Ecl1A*	ST1584	+/+	bla <sub>CMH-3-like</sub> , fosA, aadA1, ere(A), bla <sub>TEM-1B</sub> , aac(3)-IIa, qnrS1, sul2, sul1, aadA1, dfrA15, tet(D), catA2, bla <sub>TEM-1D</sub>
	Kpn2A1	ST25	+/+	fosA, bla <sub>SHV-11</sub> , dfrA12, aadA2, sul1, mph(A)
	Kpn2A2	ST25	+/+	fosA, bla <sub>SHV-11</sub> , dfrA12, aadA2, sul1, mph(A)
	Kpn3A1	ST25	+/+	fosA, bla <sub>SHV-11</sub> , dfrA12, aadA2, sul1, mph(A)
	Kae3B2	ST93	+/-	fosA
	Kpn3B3	ST25	+/+	fosA, bla <sub>SHV-11</sub> , dfrA12, aadA2, sul1
	Kpn3B4	ST25	+/+	fosA, bla <sub>SHV-11</sub> , dfrA12, aadA2, sul1
	Kae3D5	ST93	+/-	fosA
	Kae3D6	ST93	+/+	fosA
	Kpn4A	ST25	+/+	fosA, bla <sub>SHV-11</sub> , dfrA12, aadA2, sul1, mph(A), sul2, catA1, tet(D)
	Kpn5B	ST25	+/+	fosA, bla <sub>SHV-11</sub> , dfrA12, aadA2, sul1, mph(A), sul2, catA1, tet(D)
	Kpn6A	ST25	+/+	fosA, bla <sub>SHV-11</sub>
	Kpn7C	ST37	- / -	fosA, bla <sub>SHV-11</sub> , bla <sub>CTX-M-15</sub> , bla <sub>OXA-162</sub>
	Kpn8A1	ST25	+/+	fosA, bla <sub>SHV-11</sub> , dfrA12, aadA2, sul1, mph(A), sul2, catA1, tet(D)
	Eco8A2	ST453	+/+	<i>bla</i> <sub>TEM-1B</sub> , <i>tet</i> (A)
	Kpn8A3	ST25	+/+	fosA, bla <sub>SHV-11</sub>
	Kpn8E4	ST25	+/+	fosA, bla <sub>SHV-11</sub>
	Kae9A1	ST93	+/-	fosA
	Kpn9A2	ST25	+/+	fosA, bla <sub>SHV-11</sub> , dfrA12, aadA2, sul1, mph(A)
	Kpn10A1	ST25	+/+	fosA, bla <sub>SHV-11</sub> , dfrA12, aadA2, sul1, mph(A), sul2, catA1
	Kpn10A2	ST25	+/+	fosA, bla <sub>SHV-11</sub>
	Kpn10A3	ST25	+/+	fosA, bla <sub>SHV-11</sub> , dfrA12, aadA2, sul1, mph(A), sul2, catA1, tet(D)
	Eco10A4	ST453	+/-	<i>bla</i> <sub>TEM-1B</sub> , <i>tet</i> (A)
	Kpn10E5	ST25	+/+	fosA, bla <sub>SHV-11</sub>

<sup>a</sup> The index isolate is marked with an asterisk (\*). <sup>b</sup> Multi-locus sequence type. <sup>c</sup> Carriage of the plasmids encoding OXA-48 and CMY-4, pEcl1A2 and pEcl1A4, respectively

Tables 2 and 3). All isolates had a uniform mapping coverage over p*Ecl*A2, except for the *K. pneumoniae* isolate from Case 7 (*Kpn*7C), in which the plasmid lacked a 12 kb region (position 31 267–43 887) including, e.g. genes involved in conjugation (*mobAB* and *traHIJ*), as well as a 1.5-kb region (position 46 682–46 792) encoding a primase. Further, the plasmid of isolate *Kpn*7C did not harbour *bla*<sub>OXA-48</sub> but *bla*<sub>OXA-162</sub>, another carbapenemase of the OXA-48 group. This isolate also lacked p*Ecl*A4, the IncQ plasmid of the index isolate, which was detected in all other *K. pneumoniae* isolates, one *K. aerogenes*, and one *E. coli* isolate (Fig. 1, Table 3, Fig. 2). None of the other plasmids of the index isolate was found in the other outbreak isolates.

#### **Epidemiological strain typing**

The MLST analysis revealed that all *K. pneumoniae* isolates belonged to ST25 except for isolate *Kpn*7C (Case 7) which belonged to ST37 (Table 3). Further analyses using core genome MLST (573 loci) showed no more than two allele differences between the ST25 isolates, strongly indicating strain identity. Isolate *Kpn*7C clearly belonged to a separate strain as it differed in at least 475 core alleles when compared to the ST25 isolates. Based on these data, and the fact that the *bla*<sub>OXA-162</sub> plasmid of isolate *Kpn*7C did not fully



**Fig.2** A schematic figure of the proposed bacterial transmission and plasmid transfer events during the outbreak. Grey squares represent the *K. pneumoniae* ST25 outbreak strain in sources where we have no proof of its presence

resemble p*Ecl*A2, we could conclude that Case 7 was not part of the outbreak.

The four *K. aerogenes* isolates (*Kae*3B2, *Kae*3D5, *Kae*3D6, and *Kae*9A1) from twin pair 1, Case 3 and 9, belonged to ST93 (Table 3), and the SNP analysis of these isolates identified 1–7 SNPs difference using *K. aerogenes* KCTC 2190 as the reference and 0–4 SNPs using *Kae*3D5 as the reference, demonstrating that the four isolates belonged to the same strain. Finally, the two *E. coli* isolates (*Eco*8A2 and *Eco*10A4) from twin pair 2, Cases 8 and 10, both belonged to ST453 (Table 3), and the SNP analysis did not reveal any true SNPs when comparing the two isolates.

# Discussion

In this study, we describe in detail an outbreak caused by OXA-48-producing *Enterobacteriaceae*, involving nine preterm infants at a Swedish NICU. To the best of our knowledge, this is the first description of an outbreak of OXA-48-producing *Enterobacteriaceae* in a neonatal setting in northern Europe.

Interestingly, all cases identified in the present outbreak were preterm infants (gestational age at birth 24–34 weeks). Both prematurity per se, but also increased exposure to antibiotics in this group may hamper colonization by protective anaerobic gut bacteria [10]. This predisposes to overgrowth by *Enterobacteriaceae*, and high population counts of these bacteria might promote bacterial conjugation and horizontal gene transfer [14, 26].

The extended rectal screening program, initiated after the identification of the index case, ensured early detection of each new case and in-depth molecular data was obtained using whole-genome sequencing. The approach of combining genomic and clinical epidemiological data allowed us to create a likely scenario for the plasmid transfer and strain dissemination (Fig. 2).

Thus, we propose that the *E. cloacae* index strain transferred two plasmids, p*Ecl*A2 and p*Ecl*A4, to the outbreak strain *K. pneumoniae* ST25. This likely occurred in the gut of Case 1, who was identified as both the index and the primary case in this outbreak. However, since Case 1 was excluded from further screening after the isolation of the *E. cloacae* strain, carriage of *K. pneumoniae* ST25 could not be established for this patient. Therefore, an unidentified carrier in whom the plasmid transfer occurred, either from *E. cloacae* ST1584 to *K. pneumoniae* ST25 or the reverse, cannot be excluded, although the meticulous rectal screening program reduces the likelihood of an undetected neonatal case within Ward A. It is also unlikely that the plasmid transfer occurred outside a human host since high bacterial densities are required for conjugation [26].

This outbreak lasted for 50 days; hence, multiple incidents would have contributed to the continued spread of the K. pneumoniae ST25 strain. However, the simultaneous identification of Cases 2-5, after several negative screening cultures, indicates that they obtained the K. pneumoniae ST25 strain at approximately the same time, although the different resistance profiles of their isolates could imply several sources of transmission (Table 2). As Case 1 had been absent from the ward for 7 days at the time of their identification, we hypothesize that K. pneumoniae ST25 spread from Case 1 to Cases 2–5 through one or several contaminated fomites within Ward A. This could not be further investigated as no environmental cultures were collected during the outbreak. However, the isolation of outbreak clones of K. pneumoniae from the NICU environment has been reported in several studies [7, 27–31].

Within Ward B, *K. pneumoniae* ST25 was likely transmitted from Cases 5 to Case 6 since there was a clear spatial and temporal link between these cases. The source and route of spread to Case 8 were obscure, while the discovery of Cases 9 and 10 was expected due to the spread from their respective twin sibling.

The genetic analysis of isolate *Kpn*7C separated Case 7 from the outbreak. This illustrates the potential clinical benefits of access to rapid epidemiological genetic typing in future outbreak investigations, as unnecessary resource-consuming investigations may be avoided at an early stage of an outbreak.

Although we could only speculate on when and where the plasmids pEclA2 and pEclA4 have transferred between the index E. cloacae strain and the K. pneumoniae outbreak strain, other plasmid transfer events were more clearly indicated. Based on the genetic analysis and timeline of followup isolates from the two twin pairs, in vivo horizontal transfer of the plasmids pEclA2 and pEclA4 from the outbreak K. pneumoniae strain to other enterobacterial species was strongly suggested (Fig. 1). We hypothesize that an initial transfer of pEclA2, followed by pEclA4 from K. pneumoniae ST25 to the K. aerogenes ST93 strain occurred in the gut of Case 3, and that both the K. pneumoniae ST25 strain (with both plasmids) and the K. aerogenes ST93 strain (with plasmid pEclA2) were transmitted to its sibling, Case 9. Similarly, we propose that pEclA2 and pEclA4 were transferred from the K. pneumoniae ST25 strain to the E. coli ST453 strain in the gut of Case 8. Somewhat later the E. coli ST453 strain carrying only the pEclA2 plasmid was isolated from its twin sibling, Case 10. This could either be a result of the transfer of pEclA2 from K. pneumoniae ST25 to E. coli ST453 in the gut of Case 10, or the spread of an E. coli ST453 variant carrying only pEclA2 from Cases 8 to Case 10.

Our data, therefore, give strong indications, but no proof of plasmid transfer. However, transfer of resistance

plasmids in vivo between *E. coli* strains in the gut microbiota of infants has previously been demonstrated [13, 32], and in vitro conjugation experiments have shown horizontal transfer of carbapenemase-encoding outbreak plasmids between gut bacteria [24, 33, 34]. Further, a recent experimental study suggests a high potential for horizontal gene transfer in the gut microbiota of preterm neonates [14].

In summary, strain typing and plasmid tracing using whole-genome sequencing in combination with clinical epidemiology data proved to be a useful tool for the analysis of the transmission of bacterial strains and in vivo transfer of plasmids during this outbreak. Our methods allowed the exclusion of a strain producing an OXA-48-like carbapenemase unrelated to the outbreak, while several enterobacterial strains of different species could be identified as part of the outbreak, due to their carriage of identical OXA-48-encoding plasmids. Although rapid less discriminating diagnostic methods are crucial for an early outbreak response, the ability to discriminate non-related cases at an early stage is important for effective resource allocation. Even more so in high-prevalence settings where incidental findings of carbapenemase-producing *Enterobacteriaceae* are common.

Supplementary Information The online version contains supplementary material available at https://doi.org/10.1007/s10096-023-04584-y.

Acknowledgements The authors acknowledge the assistance of the Culture Collection University of Gothenburg (CCUG) Lab. The CCUG is supported by the Department of Clinical Microbiology at the Sahlgrenska University Hospital. We thank the Institute Pasteur teams for the curation and maintenance of BIGSdb-Pasteur databases at bigsdb. pasteur.fr.

Funding Open access funding provided by University of Gothenburg.

**Data Availability** The genome assemblies of the included isolates have been submitted til NCBI Assembly. The accession numbers for the assemblies are listed in Table 1 together with additional information on the included isolates.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

# References

 Institute PHAoSaNV, Swedres Svarm- sales of antibiotics and occurrence of antibiotic resistance in Sweden, in Swedres Svarm, R.O.a.W.T. Olov Aspevall, Public health agency of Sweden. Oskar Nilsson and Märit Pringle, National Veterinary Institute., Editor. 2021, Public Health Agency of Sweden and National Veterinary Institute: Sweden

- Pitout JDD, Peirano G, Kock MM, Strydom KA, Matsumura Y (2019) The global ascendency of OXA-48-Type Carbapenemases. Clin Microbiol Rev 33(1):e00102-19. https://doi.org/10.1128/ CMR.00102-19
- Poirel L, Potron A, Nordmann P (2012) OXA-48-like carbapenemases: the phantom menace. J Antimicrob Chemother 67(7):1597–1606
- Mansour W et al (2017) Outbreak of colistin-resistant carbapenemase-producing Klebsiella pneumoniae in Tunisia. J Glob Antimicrob Resist 10:88–94
- 5. Flannery DD et al (2022) Neonatal multidrug-resistant gramnegative infection: epidemiology, mechanisms of resistance, and management. Pediatr Res 91(2):380–391
- Taoufik L et al (2019) Emergence of OXA-48 carbapenemase producing Klebsiella pneumoniae in a neonatal intensive care unit in Marrakech. Morocco Clin Med Insights Pediatr 13:1179556519834524
- Adler A et al (2013) Epidemiological and microbiological characteristics of an outbreak caused by OXA-48-producing Enterobacteriaceae in a neonatal intensive care unit in Jerusalem. Israel J Clin Microbiol 51(9):2926–2930
- Gona F et al (2019) Emergence of two novel sequence types (3366 and 3367) NDM-1- and OXA-48-co-producing K pneumoniae in Italy. Eur J Clin Microbiol Infect Dis 38(9):1687–1691
- Beller L, Deboutte W, Falony G, Vieira-Silva S, Tito RY, Valles-Colomer M, Rymenans L, Jansen D, Van Espen L, Papadaki MI, Shi C, Yinda CK, Zeller M, Faust K, Van Ranst M, Raes J, Matthijnssens J (2021) Successional Stages in Infant Gut Microbiota Maturation. mBio 12(6):e0185721. https://doi.org/10.1128/mBio. 01857-21
- Aguilar-Lopez M et al (2021) A systematic review of the factors influencing microbial colonization of the preterm infant gut. Gut Microbes 13(1):1–33
- Lerminiaux NA, Cameron ADS (2019) Horizontal transfer of antibiotic resistance genes in clinical environments. Can J Microbiol 65(1):34–44
- Neil K, Allard N, Rodrigue S (2021) Molecular mechanisms influencing bacterial conjugation in the intestinal microbiota. Front Microbiol 12:673260
- 13. Gumpert H et al (2017) Transfer and persistence of a multi-drug resistance plasmid in situ of the infant gut microbiota in the absence of antibiotic treatment. Front Microbiol 8:1852
- Hagbo M et al (2020) Experimental support for multidrug resistance transfer potential in the preterm infant gut microbiota. Pediatr Res 88(1):57–65
- Francisco Salvà-Serra, Margarita Gomila, Liselott Svensson-Stadler et al (2018) A protocol for extraction and purification of high-quality and quantity bacterial DNA applicable for genome sequencing: a modified version of the Marmur procedure. PRO-TOCOL (Version 1) available at Protocol Exchange. https://doi. org/10.1038/protex.2018.084
- Bankevich A et al (2012) SPAdes: A new genome assembly algorithm and its applications to single-cell sequencing. J Comput Biol 19(5):455–477
- Wick RR et al (2017) Unicycler: resolving bacterial genome assemblies from short and long sequencing reads. PLoS Comput Biol 13(6):e1005595

- 18. Zankari E et al (2012) Identification of acquired antimicrobial resistance genes. J Antimicrob Chemother 67(11):2640–2644
- Langmead B, Salzberg SL (2012) Fast gapped-read alignment with Bowtie 2. Nat Methods 9(4):357
- Li H et al (2009) The sequence alignment/map format and SAMtools. Bioinformatics 25(16):2078–2079
- Larsen MV et al (2012) Multilocus sequence typing of totalgenome-sequenced bacteria. J Clin Microbiol 50(4):1355–1361
- Jolley KA, Bray JE, Maiden MCJ (2018) Open-access bacterial population genomics: BIGSdb software, the PubMLST org website and their applications. Wellcome Open Res 3:124
- 23. Miyoshi-Akiyama T et al (2013) Multilocus sequence typing (MLST) for characterization of Enterobacter cloacae. PLoS ONE 8(6):e66358
- 24. Hamprecht A et al (2019) Pathogenicity of Clinical OXA-48 Isolates and impact of the OXA-48 IncL plasmid on virulence and bacterial fitness. Front Microbiol 10:2509
- Kotsakis SD et al (2015) Characterization of a mobilizable IncQ plasmid encoding cephalosporinase CMY-4 in Escherichia coli. Antimicrob Agents Chemother 59(5):2964–2966
- Stecher B et al (2012) Gut inflammation can boost horizontal gene transfer between pathogenic and commensal Enterobacteriaceae. Proc Natl Acad Sci USA 109(4):1269–1274
- Haller S et al (2015) What caused the outbreak of ESBL-producing Klebsiella pneumoniae in a neonatal intensive care unit, Germany 2009 to 2012? Reconstructing transmission with epidemiological analysis and whole-genome sequencing. BMJ Open 5(5):e007397
- Rettedal S et al (2012) First outbreak of extended-spectrum betalactamase-producing Klebsiella pneumoniae in a Norwegian neonatal intensive care unit; associated with contaminated breast milk and resolved by strict cohorting. APMIS 120(8):612–621
- 29. Giuffre M et al (2013) Successful control of an outbreak of colonization by Klebsiella pneumoniae carbapenemase-producing K. pneumoniae sequence type 258 in a neonatal intensive care unit. Italy. J Hosp Infect. 85(3):233–6
- 30. Saporito L et al (2021) Efficacy of a coordinated strategy for containment of multidrug-resistant Gram-negative bacteria carriage in a Neonatal Intensive Care Unit in the context of an active surveillance program. Antimicrob Resist Infect Control 10(1):30
- 31. Cantey JB, Sreeramoju P, Jaleel M, Treviño S, Gander R, Hynan LS, Hill J, Brown C, Chung W, Siegel JD, Sánchez PJ (2013) Prompt control of an outbreak caused by extended-spectrum β-lactamase-producing Klebsiella pneumoniae in a neonatal intensive care unit. J Pediatr 163(3):672–9.e1–3. https://doi.org/ 10.1016/j.jpeds.2013.03.001
- 32. Porse A et al (2017) Genome Dynamics of Escherichia coli during antibiotic treatment: transfer, loss, and persistence of genetic elements in situ of the infant gut. Front Cell Infect Microbiol 7:126
- Tofteland S et al (2013) A long-term low-frequency hospital outbreak of KPC-producing Klebsiella pneumoniae involving Intergenus plasmid diffusion and a persisting environmental reservoir. PLoS ONE 8(3):e59015
- Pitart C et al (2011) First outbreak of a plasmid-mediated carbapenem-hydrolyzing OXA-48 beta-lactamase in Klebsiella pneumoniae in Spain. Antimicrob Agents Chemother 55(9):4398–4401

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.