



## **Split-marker-mediated genome editing improves homologous recombination frequency in the CTG clade yeast *Candida intermedia***

Downloaded from: <https://research.chalmers.se>, 2025-12-04 23:24 UTC

Citation for the original published paper (version of record):

Peri, K., Da Silva Faria Oliveira, F., Larsson, A. et al (2023). Split-marker-mediated genome editing improves homologous recombination frequency in the CTG clade yeast *Candida intermedia*. FEMS Yeast Research, 23. <http://dx.doi.org/10.1093/femsyr/foad016>

N.B. When citing this work, cite the original published paper.

# Split-marker-mediated genome editing improves homologous recombination frequency in the CTG clade yeast *Candida intermedia*

Kameshwara V. R. Peri<sup>1</sup>, Fábio Faria-Oliveira<sup>1</sup>, Adam Larsson<sup>1</sup>, Alexander Plovie<sup>1</sup>, Nicolas Papon<sup>2</sup>, Cecilia Geijer<sup>1,\*</sup>

<sup>1</sup>Chalmers University of Technology, Department of Life Sciences, Division of Industrial Biotechnology, Kemigården 1, 41296 Gothenburg, Sweden

<sup>2</sup>University of Angers, Univ Brest, IRF, SFR ICAT, F-49000 Angers, France

\*Corresponding author. Chalmers University of Technology, Department of Life Sciences, Division of Industrial Biotechnology, 41296 Gothenburg, Sweden.

E-mail: [cecilia.geijer@chalmers.se](mailto:cecilia.geijer@chalmers.se)

Editor: [Carol Munro]

## Abstract

Genome-editing toolboxes are essential for the exploration and exploitation of nonconventional yeast species as cell factories, as they facilitate both genome studies and metabolic engineering. The nonconventional yeast *Candida intermedia* is a biotechnologically interesting species due to its capacity to convert a wide range of carbon sources, including xylose and lactose found in forestry and dairy industry waste and side-streams, into added-value products. However, possibilities of genetic manipulation have so far been limited due to lack of molecular tools for this species. We describe here the development of a genome editing method for *C. intermedia*, based on electroporation and gene deletion cassettes containing the *Candida albicans* NAT1 dominant selection marker flanked by 1000 base pair sequences homologous to the target loci. Linear deletion cassettes targeting the ADE2 gene originally resulted in <1% targeting efficiencies, suggesting that *C. intermedia* mainly uses nonhomologous end joining for integration of foreign DNA fragments. By developing a split-marker based deletion technique for *C. intermedia*, we successfully improved the homologous recombination rates, achieving targeting efficiencies up to 70%. For marker-less deletions, we also employed the split-marker cassette in combination with a recombinase system, which enabled the construction of double deletion mutants via marker recycling. Overall, the split-marker technique proved to be a quick and reliable method for generating gene deletions in *C. intermedia*, which opens the possibility to uncover and enhance its cell factory potential.

**Keywords:** nonhomologous end-joining, nonconventional yeast, cell factory development

## Introduction

A biobased society relies on efficient utilization of renewable raw materials for production of fuels, chemicals, and materials. In this regard, microbial cell factories that can convert various carbon sources found in industrial waste and side-streams into valuable products are highly sought after as key enablers of sustainable and circular production processes. Whereas the industrial workhorse *Saccharomyces cerevisiae* can only ferment a subset of monosaccharides, several so called nonconventional yeast species display broad substrate ranges, often including various monosaccharides as well as disaccharides and sometimes even more complex oligo- and polysaccharides (Ramírez-Orozco et al. 2001, Lobs et al. 2017, Baptista and Domingues 2022, Šuchová et al. 2022). To develop a nonconventional yeast into a cell factory, detailed metabolic characterization on a molecular level remains an essential prerequisite. This typically requires genome editing tools that enable deletion of specific genes of interest and determination of genotype–phenotype relationships (Catlett et al. 2003). Moreover, targeted mutagenesis and expression of heterologous genes can further improve traits of a production host (Fonseca et al. 2011, Westfall et al. 2012).

*Candida intermedia* is a nonconventional yeast with an unusually broad substrate range (Shen et al. 2018). Its potential as a

cell factory has been highlighted by the production of ethanol (Gardonyi et al. 2003, Fonseca et al. 2011, Geijer et al. 2020), xylitol, and single-cell protein (Wu et al. 2018) from xylose derived from lignocellulosic biomass. *Candida intermedia* has also been used for bioremediation of lactose found in the abundant industrial side stream cheese whey (Yonten and Aktas 2014), as well as production of antimicrobial peptides with application in wine preservation (Acuña-Fontecilla et al. 2017, Pena and Ganga 2019). We have previously *de novo* genome-sequenced the *C. intermedia* strain CBS 141442, which allows us to study the yeast at a genomic level (Geijer et al. 2020). Our analysis showed that *C. intermedia* is a haploid yeast belonging to the CTG clade of ascomycetous yeast, where the universal CUG codon is predominantly translated into a serine rather than a leucine (Geijer et al. 2020, Muhlhausen et al. 2021). The CTG clade comprises many relatively well-studied yeasts including human pathogens *Candida albicans*, *Candida tropicalis*, *Candida parapsilosis*, and industrially attractive yeasts such as *Debaryomyces hansenii*, *Metschnikowia pulcherrima*, and *Scheffersomyces stipitis* (Enkler et al. 2016, Cao et al. 2018, Muhlhausen et al. 2021, Strucko et al. 2021, Zhang et al. 2022). Genome editing tools have been developed for many of these species, which means that several CTG clade-compatible plasmids containing selectable marker genes and reporter sys-

Received: November 23, 2022. Revised: February 2, 2023. Accepted: March 8, 2023

© The Author(s) 2023. Published by Oxford University Press on behalf of FEMS. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

tems are available (Papon et al. 2012, Lobs et al. 2017, Uthayakumar et al. 2021). These plasmids can be employed in the development of genetic engineering toolboxes for other yeast such as *C. intermedia*.

Genome editing often involves integration of a foreign DNA fragment in the host genome through the action of a double-strand break (DSB) repair mechanism, typically by nonhomologous end-joining (NHEJ) or homologous recombination (HR). The classical NHEJ mechanism is aided by the ku70/ku80 heterodimer and DNA ligase IV and involves error prone DSB repair where short (1–4 nucleotides) homologous fragments are used to repair the DSB. Ku-independent mechanisms are also known to exist, including microhomology-mediated end-joining (MMEJ) and single-strand annealing (SSA) that involve more extended homologies flanking the DSB (Wilson 2007). In contrast, HR involves DSB repair templated by the homologous sister chromosome, making it an error-free mechanism (Krejci et al. 2012, Sfeir and Symington 2015). Whereas NHEJ results in random integration of linear DNA fragments into the genome, HR targets DNA to a specific genomic locus (Orr-Weaver et al. 1981). NHEJ and HR coexist in most organisms but the balance in activity between them varies among species and cell types. *Saccharomyces cerevisiae* uses almost exclusively HR (Detloff et al. 1991) and as little as 20 bp genomic homology on each side of the selective marker gene is sufficient for targeted deletions (Corrigan et al. 2013). On the other hand, many members of the CTG clade seem to display a predominance of ectopic, NHEJ-mediated integrations (Lobs et al. 2017) necessitating long stretches of homology (0.5–2 kbp) to help direct the DNA fragment to the intended genome locus in many species (Nelson et al. 2003, Moreno-Beltran et al. 2021). Additionally, targeted gene deletions in species with low HR/NHEJ ratios typically require screening of many transformants to identify the desired mutant(s) (Fraczek et al. 2018).

Different approaches have been used to increase the HR frequency to make identification of mutants more efficient. Impairment of the NHEJ machinery by disrupting one of the genes encoding the ku70/ku80 heterodimer or other central NHEJ components has been shown to significantly increase HR in various yeast and fungal species (Kooistra et al. 2004, da Silva Ferreira et al. 2006, Haarmann et al. 2008, Kretzschmar et al. 2013). However, creating strains that are defective in the NHEJ pathway may be time consuming and labor intensive, and mutations in this pathway are known to compromise genome integrity maintenance (Barnes and Rio 1997, Dudášová et al. 2004, Kretzschmar et al. 2013, Raschmanová et al. 2018). Another approach to increase HR frequency, where NHEJ is left intact, is to benefit from the well-conserved cell cycle-dependent oscillation in HR/NHEJ preference of cells (Barnes and Rio 1997). Hydroxyurea treatment arrests cells in the S-phase of the cell cycle, where cells are more prone to use HR as sister chromatids are available as templates for repair. This way, targeting rates have been improved in yeast such as *Yarrowia lipolytica*, *S. cerevisiae*, *Kluyveromyces lactis*, and *Pichia pastoris* (Tsakraklides et al. 2015). A third way to promote HR is to use a split-marker transformation protocol, which was originally developed for rapid, gap repair-mediated, one step cloning in *S. cerevisiae*. Here, the homologous sequences of the target gene are fused to a selectable marker gene that is truncated in two different fragments with partial overlap in sequence, and only transformants in which HR between the overlapping parts has resulted in the constitution of a complete selection marker gene will be able to grow in selective medium. It has been shown that HR assembly of the marker fragments also promotes HR integration of the DNA cassette in the genome, resulting

in a high frequency of targeted gene disruptions (Fairhead et al. 1996).

In many nonconventional yeasts, the limited availability of auxotrophic and antibiotic resistance markers is a real bottleneck for generation of multiple genome modifications in the same strain background. One solution is marker recycling, for example by making use of a flippase (Flp) adapted for CTG clade yeast (Morschhauser et al. 1999, Mancera et al. 2019). In this approach, the selectable marker is combined with the *FLP* gene and the whole cassette is flanked on either side by Flp recognition target (*FRT*) sequences. Flp catalyzes recombination of the *FRT* sequences and excision of the cassette, leaving only a small scar at the target locus. Another, more recently developed solution for markerless, multiple genome editing is the clustered regularly interspaced short palindromic repeat (CRISPR) and CRISPR-associated protein 9 (Cas9) (Jinek et al. 2012). Although the list of species that can be modified by this method expands every year, development of CRISPR/Cas9-based tools for new yeast species is not trivial and often requires species-specific optimization of gene expression, transformation, and selection protocols (Vyas et al. 2015, Ng and Dean 2017, Ennis et al. 2021, Strucko et al. 2021).

The aim of this study was to develop a first set of genome-editing tools for *C. intermedia*. The work included identification of a functional transformation protocol and deletion cassettes that enabled both single and double gene deletions in this yeast. Moreover, cell synchronization and in particular a split-marker approach improved the frequency of HR-mediated targeted genome integration of deletion cassettes, which may be useful also for future CRISPR/Cas9-based engineering of *C. intermedia* and other yeasts.

## Materials and methods

### Plasmids

#### CaNAT1-containing plasmids

Integrative plasmids pCaNAT1 (pJK795) and pFLP\_CaNAT1 (pJK863) were provided by Dr. Julia Köhler, Boston Children's Hospital, USA. Both plasmids are derivatives of plasmid pJK799 (Shen et al. 2005) and contain the CaNAT1 selectable marker encoding nourseothricin acetyltransferase that has been shown to confer nourseothricin (cloNAT) resistance to *C. albicans* and other *Candida* species. The CaNAT1 marker is expressed from the constitutive *TEF1* promoter and terminator in *Ashbya gossypii*. In addition to the CaNAT1 marker, plasmid pFLP\_CaNAT1 contains a *FLP* gene encoding a FLP-recombinase driven by *C. albicans* *SAP2* promoter, which is induced by growing the yeast in medium with bovine serum albumin (BSA) as the sole nitrogen source (Morschhauser et al. 1999).

#### Construction of ADE2 disruption plasmid

To generate the ADE2 disruption cassette with CaNAT1 marker, the ADE2 gene from *C. intermedia* was PCR amplified along with 1000 bp upstream and downstream sequences and subsequently cloned into the multiple cloning site of a pBluescript II KS + vector using restriction enzymes KpnI and SacII and T4 DNA ligase (Thermo Fisher, Waltham, MA, USA) according to manufacturer's instructions. The resulting plasmid was then digested using ClaI restriction enzyme (Thermo Fisher) that cuts in the middle of the ADE2 gene. The CaNAT1 marker, PCR-amplified from the pCaNAT1 plasmid, was thereafter inserted at the cut site, creating plasmid pADE2disr\_CaNAT1.

## Construction of additional gene deletion cassettes

Construction of deletion cassettes (including the ADE2 deletion cassette with pFLP\_CaNAT1) was done using molecular cloning: plasmids pCaNAT1 or pFLP\_CaNAT1 were linearized and used as backbone for cloning the upstream and downstream homologous flanking region of the target gene next to the CaNAT1 marker. Plasmid pCaNAT1 was linearized using KpnI (for inserting an upstream homology sequence) or EcoRV (for inserting a downstream homology sequence) whereas pFLP\_CaNAT1 was cut with KpnI (for upstream homology) or SacI (for downstream homology). Primer design and cloning conditions were performed using NEBuilder® HiFi DNA assembly master mix and NEBuilder® Assembly tool (New England Biolabs, Ipswich, MA, USA), as per manufacturer's protocol. Primers were designed to PCR amplify the homology arms (~1000 bp) on either side of the target gene from genomic DNA isolated from *C. intermedia* strain CBS 141442. Amplified homology arms were cloned into the respective linearized plasmid resulting in two plasmids, each containing the same backbone but different homology arms.

## Transformation of *Escherichia coli*

All cloned plasmids were transformed into *E. coli* DH5 $\alpha$  chemical competent cells (Sambrook and Russell 2006). Positive clones were identified using colony PCR, and propagated overnight in LB medium (1% tryptone, 1% NaCl and 0.5% yeast extract) supplemented with ampicillin (100  $\mu$ g/ml) at 37°C and 1  $\times$  g followed by plasmid isolation using the GENEJet Plasmid miniprep kit (Thermo Fisher).

## Preparation of linear and split-marker DNA for yeast transformation

Prior to yeast transformation, plasmid pCaNAT1 was linearized by restriction digestion with KpnI enzyme resulting in linearized plasmid, lin\_CaNAT1.

Linearized cassette targeting the ADE2 gene was PCR amplified using pADE2disr\_CaNAT1 as template with primers annealing to the homologous arms. The resulting linear cassette contained CaNAT1 marker flanked on either side by 1000 bp homologous arms to the ADE2 gene in *C. intermedia*.

The split-marker fragments were generated using PCR such that the primers annealed to the homologous arms and in the markers, resulting in two fragments each containing up/downstream homology to the target gene fused to a part of the CaNAT1 marker. Split-marker fragments had an overlap of 391 bp in the CaNAT1 gene. Split-marker fragments derived from pCaNAT1 and pFLP\_CaNAT1 are referred to as splitCaNAT1 and splitFLP\_CaNAT1, respectively.

## Yeast transformation

The transformation procedure was adapted from the protocol developed by De Backer et al. (1999) for the transformation of *C. albicans* by electroporation. In short, an overnight culture of *C. intermedia* CBS 141442 grown at 30°C and 1  $\times$  g shaking was used to start a culture at OD<sub>600</sub> = 0.1 in 100 ml YPD (1% yeast extract, 2% bacto peptone, and 2% glucose). Once cells reached OD<sub>600</sub> = 1, the culture was centrifuged (2600  $\times$  g for 3 min) and washed twice with ice-cold sterile deionized water and once with ice-cold 1 M sorbitol. Cells were centrifuged once again followed by resuspension in 160  $\mu$ l of 1 M sorbitol. The resuspended cells were divided into 40  $\mu$ l aliquots in microcentrifuge tubes. To each aliquot, 8  $\mu$ l of DNA (with an optimal amount of 2  $\mu$ g) was added. The transformation mix was transferred to a chilled electroporation cuvette

(0.2 cm gap BioRad MicroPulser electroporation cuvettes, BioRad Laboratories, Hercules, CA, USA) and electroporated with a BioRad MicroPulser (BioRad Laboratories) at 1.5 kV. Cells were thereafter gently transferred to 1 ml of 1 M sorbitol in YPD and incubated at 30°C for 3 h with 1  $\times$  g of shaking. Cultures were spread on YPD agar (2%) plates with cloNAT (200  $\mu$ g/ml) for selection and left to grow at 30°C for 48 h or until colonies emerged.

## Validation of yeast deletion mutant strains

For visual screening of *ade2* deletion strains, cloNAT resistant transformants were incubated at 4°C for 7–10 days on YPD agar plates and observed for pink/red colonies. For faster screening, transformants were replica plated onto agar plates with YNB medium (0.17% yeast nitrogen base, 0.5% of (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 2% glucose, and 0.079% complete synthetic medium) without adenine, on which *C. intermedia ade2* mutants could not grow but turned dark red within 24 h.

For screening of mutants without visible phenotypes, colony PCR was performed. Individual colonies of transformants were lysed in 15  $\mu$ l deionized water and heated to 98°C for 10 min. PCR analysis was performed using PHIRE II polymerase (Thermo Fisher) with a primer pair designed to anneal upstream and downstream of the homologous flanking regions of the deletion cassette as well as a third control primer that anneal to the target gene. In this way, a successful PCR reaction resulted in either a band of ~1200 bp identifying true deletion mutants, or a band of ~1500 bp in false positive transformants where the target gene was still intact.

## Cell synchronization

Liquid cultures of *C. intermedia* CBS 141442 were prepared by inoculating 20 ml of YPD with fresh cells from YPD plates. Cultures were incubated overnight at 30°C with 1  $\times$  g of shaking. These precultures were then used to inoculate fresh cultures at OD<sub>600</sub> = 0.2 and left to grow for 3 h at 30°C with 1  $\times$  g shaking. Cells were thereafter harvested and resuspended in 16 ml of YPD. To each culture, 4 ml of sterile solution containing deionized water and hydroxyurea was added to obtain the desired concentrations of hydroxyurea (0–200 mM). Cultures were incubated at 30°C with 1  $\times$  g of shaking and OD<sub>600</sub> was measured over 3.5 h. Microscopy pictures were also taken at regular intervals using a Leica AF 6000 inverted microscope (Leica Microsystems, Wetzlar, Germany) and processed with the Leica Application Suite (LAS-X) software (version 3.4.2).

## Induction of FLP recombinase gene expression

Yeast carbon base (YCB) media (2.34%) with BSA (0.4%) was used to induce expression of the FLP recombinase gene from the integrated FLP\_CaNAT1 deletion cassette. Transformed colonies were cultured in YCB with BSA and grown over 96 h before being plated on YPD agar plates. After 48 h of growth, colonies were replica plated onto solid YPD containing cloNAT to check for regained antibiotic sensitivity due to FLP recombinase induction and activity.

## Results

### The CaNAT1 dominant marker gene can be used for selection of *C. intermedia* transformants

First, we set out to identify a working transformation protocol and a functional selectable marker gene for *C. intermedia*. Like many other CTG clade yeast (Vyas et al. 2015, Norton et al. 2017, En-



nis et al. 2021, Strucko et al. 2021), *C. intermedia* displayed sensitivity to cloNAT in the concentration of 10–1000 µg/ml (data not shown), of which 200 µg/ml was chosen in further experiments. We transformed *C. intermedia* using electroporation and a linearized CaNAT1 marker cassette that has proven effective in numerous yeast including *C. albicans*, *C. parapsilosis*, and *C. intermedia*'s close relative *Clavispora lusitanae* (outlined in Fig. 1A) (De Backer et al. 1999, Shen et al. 2005). Transformations using 1, 2, and 10 µg of linearized plasmid DNA resulted in different number of transformants, where 2 µg DNA provided the highest number (average 78) of transformants per µg DNA (Fig. 1B) and was, therefore, used in subsequent transformations. A total of 16 randomly selected cloNAT resistant colonies all contained the CaNAT1 cassette, as judged by the PCR products of expected size using primers that anneal to the CaNAT1 gene (data not shown). As the plasmid DNA contained no significant homology to the *C. intermedia* genome, these results suggest that the DNA integrated randomly in the genome or possibly remained as linear or recircularized DNA in the nucleus. In either case, the experiment provides evidence that the transformation protocol resulted in the introduction and expression of the CaNAT1 marker gene in *C. intermedia* cells, enabling selection of transformants.

To develop a protocol for targeted genome editing, we next created a cassette to disrupt the *ADE2* gene, where the CaNAT1 cassette was flanked on both sides by 1000 bp of homology to the *ADE2* locus. In *ade2* mutants, phosphoribosyl aminoimidazole accumulates causing a red pigmentation of yeast cells (Dujon et al. 2004), which enables visual screening of the desired *ade2* mutants. Three rounds of transformations yielded a total of 274 cloNAT-resistant transformants, of which only two (<1%) turned red. PCR was used to confirm that the antibiotic-resistant, pink/red colonies had the CaNAT1 marker integrated in the *ADE2* locus (data not shown). Although we have not investigated the mechanism(s) behind the DNA integrations observed, these results point to that *C. intermedia* preferentially uses NHEJ rather than HR to integrate foreign DNA in the genome, despite long homologous flanking regions of the deletion cassette.

### Cell synchronization increases the HR/NHEJ ratio in *C. intermedia*

Having established a transformation protocol and a functional selection marker, we tried to increase the HR efficiency of *C. intermedia* by synchronizing cells in the S-phase using hydroxyurea (Fig. 2A). Both 100- and 200-mM concentrations of hydroxyurea were sufficient to block the cell cycle, as determined by microscopic examination (cells arrested in the large-budded stage) and optical density measurements (cease of biomass increase over time) (Fig. 2B and C). Synchronized cells (using 200 mM hydroxyurea) transformed with the linearized *ade2\_CaNAT1* disruption cassette yielded 164 colonies, of which 3.7% proved to be pink/red *ade2* mutants. Thus, cell cycle synchronization resulted in a ~5-fold improvement in HR efficiency compared to nonsynchronized cells (Fig. 2D).

### The split-marker approach has a large positive effect on gene targeting efficiency in *C. intermedia*

To further improve gene deletion efficiency in *C. intermedia*, we thereafter employed the split-marker approach where the CaNAT1 marker was divided into two partly overlapping fragments, each flanked by 1000 bp homologous target sequences (Fig. 3A). Both S-phase synchronized, and nonsynchronized cells


were transformed with the split-marker fragments targeting the *ADE2* gene, to test the hypothesis that cells in the synchronized population would utilize the HR pathway to a larger extent than the nonsynchronized counterparts. Surprisingly, the HR efficiency increased to as much as 63.9% for synchronized and 55.7% for nonsynchronized populations, with the total number of CaNAT1 resistant transformants being only marginally higher for the synchronized cells (119) than for nonsynchronized cells (115) (Fig. 3B and C). Thus, we can conclude that with our experimental setup, the split-marker approach had a greater positive impact on the HR/NHEJ ratio than cell synchronization. As cell synchronization extends the overall transformation time and hydroxyurea is classified as a toxic compound that must be handled with great care, we decided to skip this step in subsequent transformations.

As proof of concept, we used the split-marker approach to delete three additional loci in the *C. intermedia* genome. Targeting efficiencies were similar to *ADE2*: 71% for gene *XYL1\_2* (gene size 948 bp), 67% for *LAC9* (2097 bp) encoding proteins with accession numbers SGZ56686 and SGZ49734, and 89% for a gene cluster of four genes (6577 bp) that is unique to *C. intermedia*, respectively (Fig. 3C). We speculate that the differences observed in targeting efficiency could be attributed to the genomic loci targeted, gene size and day to day experimental variation. We also assessed if short homology arms of 50 bp (split\_ade2Δ\_50 bp) were sufficient to direct the deletion cassette to the *ADE2* locus. However, only 3/78 (4%) transformants turned red, indicating the need for longer homology arms for efficient targeting using the split-marker approach (Fig. 3C). The split-marker approach repeatedly resulted in fewer transformants compared to linear deletion cassettes, and transformation of only the first of the two split-marker fragments (split\_ade2Δ\_UP) resulted in the absence of selectable transformants (Fig. 3C). Thus, transformants where HR between the marker gene fragments did not occur seem efficiently counter-selected for on the cloNAT plates. Overall, these experiments demonstrate the potential of the split-marker approach in improving HR efficiencies and enabling targeted gene deletions in *C. intermedia*.


### Marker recycling enables construction of double deletions in *C. intermedia*

A drawback of the developed gene deletion protocol in *C. intermedia* is that the marker is integrated in the genome and cannot be recycled for subsequent deletions. To achieve a markerless engineered strain that can be used for additional genome editing or for industrial purposes where the microbial cell factories cannot contain antibiotic resistance markers, we combined the split-marker approach with a FLP recombinase. The splitFLP\_CaNAT1 deletion cassettes were constructed using a pFLP\_CaNAT1 plasmid containing FLP recombinase gene in combination with the CaNAT1 marker located between flippase recognition target (FRT) sites and flanked by a homologous region (Fig. 4A) (Shen et al. 2005). For easy comparison of the targeting efficiency, we deleted again *ADE2*, *XYL1\_2* as well as genes *LAC9\_2* and *LADA* (with protein accession numbers SGZ49734 and SGZ50177). Surprisingly, the targeting efficiency dropped to ~10% for all tested loci (Fig. 4B). While further studies are needed to determine the cause of this marked decrease, one may hypothesize that the considerable larger size of the FLP\_CaNAT1 deletion cassette compared to the CaNAT1 marker alone (3202 vs. 6121 bp) has a negative effect on the HR/NHEJ ratio (Štafa et al. 2017).

When applying the pFLP\_CaNAT1 plasmid in *C. albicans*, the SAP2 promoter that drives the expression of the FLP gene in the




**Figure 1.** Transformation of *C. intermedia*. **(A)** Overview of the transformation protocol of *C. intermedia* using electroporation. Competent yeast cells were transformed with linearized plasmid (*lin\_CaNAT1*), containing *CaNAT1* selection marker driven by *A. gossypii* *TEF1* promoter and terminator, resulting in *cloNAT* resistant colonies on selection plate. **(B)** Graph depicting the number of transformants obtained per µg of linearized plasmid DNA plotted against different amount of transformed DNA (µg) tested. Error bars represent standard deviation based on biological triplicates.



**Figure 2.** Cell synchronization using hydroxyurea. **(A)** Schematic representation of cell synchronization. Yeast cells arrested in S-phase of the cell cycle using hydroxyurea were transformed with *lin\_ADE2disr\_CaNAT1* cassette. **(B)** Photomicrograph showing cells arrested in their S-phase (also called the budding stage) after treatment with hydroxyurea. **(C)** Graphical representation of the change in biomass (OD<sub>600</sub>) over time (hours) of yeast cultures treated with different concentrations of hydroxyurea. **(D)** Targeting efficiencies achieved upon transformation of linear cassette, *lin\_ADE2disr\_CaNAT1*, targeting the *ADE2* gene in both nonsynchronized (nonsync) and synchronized (sync) cultures of *C. intermedia*. Legend shows targeted *ade2Δ* and *ADE2* transformants (*WT ADE2*). A break in the y-axis has been introduced to better visualize lower values for targeting efficiencies.

deletion cassette is activated upon growth in medium with protein as the sole nitrogen source (Morschhauser et al. 1999). To test if the *CaSAP2* promoter functions in the same way in *C. intermedia*, and thereby can be used to flip out the *FLP\_CaNAT1* cassette, the *C. intermedia* transformants were grown in medium containing BSA as protein source, streaked for single colonies on YPD plates without selection and thereafter transferred to *cloNAT*-containing plates. Like wild-type cells, transformants in which the deletion cassette had flipped could no longer grow on *cloNAT*-containing medium (Fig. 4C). In this way, we successfully constructed several transformants with marker-free deletions. However, it should be noted that we had to screen multiple colonies to identify cells where the *FLP\_CaNAT1* cassette had flipped, and for some loci we even failed

to flip out the cassette despite multiple attempts. This suggests that the *C. albicans* *SAP2* promoter or the *FLP* recombinase are not fully functional in *C. intermedia*. Nevertheless, we have managed to create a double deletion mutant by first deleting a gene cluster (the *GAL* gene cluster, 7027 bp) using the *splitFLP\_CaNAT1* cassette (targeting efficiency of 4%), followed by deletion of the *XYL1\_2* gene using the *splitCaNAT1* deletion cassette with a targeting efficiency of 60%. Although further optimization such as replacing the *CaSAP2* promoter with an endogenous inducible promoter may likely improve the overall efficiency of the process, our results clearly show that *FLP* recombinase technique in combination with split-marker cassette enable construction of double deletion mutants in *C. intermedia*.




**Figure 3.** Split-marker method for genome editing in *C. intermedia*. **(A)** Schematic drawing of the integration of split-marker cassette at a target locus by HR. Split-marker cassettes contain either upstream/downstream (represented as UP/DOWN respectively) homology to the target gene and 391 bp homology at the *CaNAT1* marker. Homology between different fragments is represented by dotted lines. **(B)** A representative transformation plate displaying pink/red *ade2* mutants and white *ADE2* transformants growing on cloNAT selective medium. **(C)** Bar graph depicting targeting efficiencies using split-marker approach. Labels on x-axis: split-marker deletion with synchronized (split\_sync\_ade2Δ), nonsynchronized cells (split\_ade2Δ), 50 bp homology split fragment targeting *ADE2* gene (split\_ade2Δ\_50 bp), only one of the split fragments (split\_ade2Δ\_UP), *XYL1\_2* gene deletion (split\_xyl1\_2Δ), *LAC9* deletion (split\_lac9Δ), and gene cluster deletion (split\_geneclusterΔ) targeted in *C. intermedia*. Error bars represent standard deviation based on biological triplicates and statistical significance is determined using student's t-test and is represented as "\*" for  $P \leq .05$ , "\*\*\*" for  $P \leq .01$ , and "\*\*\*\*" for  $P \leq .001$ .

## Discussion

The ability to manipulate chromosomally encoded genes within cells is a fundamental tool for dissecting the biology of an organism and to engineer microbial cell factories for bioproduction. In this study, we have developed a molecular toolbox for *C. intermedia*, including identification and optimization of a functional transformation protocol and deletion cassettes. Importantly, we improved the HR frequency from an original <1% to 55%–70%, enabling straight-forward targeted gene deletions and screening of transformants.

*De novo* development of genome-editing tools in new yeast species requires identification of functional transformation protocols and screening of cells that have taken up, integrated, and expressed the foreign DNA. Even though we benefitted greatly from previous work and tools developed for other CTG-clade yeast (De Backer et al. 1999, Morschhauser et al. 1999, Shen et al. 2005), each step of the process had to be tested and optimized for *C. intermedia*. Transformation of yeast is most often done using LiAc and

heat shock or electroporation protocols (Gietz and Schiestl 2007, Kawai et al. 2010). In our hands, electroporation resulted in hundreds of *C. intermedia* transformants whereas the assessed LiAc-coupled heat shock protocol gave rise to 10-fold less colonies on the selection plates (unpublished results). The deletion cassettes employed in this study, in which the constitutive promoter *AgTEF1* drives the expression of selection marker *CaNAT1* that confers resistance to cloNAT seems to work in many CTG clade yeast (Shen et al. 2005, Norton et al. 2017, Strucko et al. 2021), and so also in *C. intermedia*. To identify additional markers, we tested also a CTG-compatible kanMX marker shown to function in several yeast (Hara et al. 2000, Millerioux et al. 2011). Unfortunately, transformation of the kanMX cassette did not result in any geneticin-resistant *C. intermedia* colonies. We also tried to generate uracil biosynthesis deficient strain by UV-mutagenesis followed by selection on 5-fluoroorotic acid-containing plates (Kohler and Fink 1996) but this experiment failed, as all screened colonies turned out to be auxotrophs for uracil (data not shown). This left us with



**Figure 4.** Gene deletion and marker recycling using FLP recombinase technique. **(A)** Schematic representation of HR-mediated integration of the splitFLP\_CaMAT1 at target locus in *C. intermedia* genome followed by the induction of FLP-recombinase expression resulting in FLP-CaMAT1 deletion cassette "flip out" and recombination of the FRT sites. Thus, the resulting transformant is marker free. **(B)** Targeting efficiencies using splitFLP-CaMAT1 compared to the splitCaMAT1 deletion cassette. A break in the y-axis has been introduced to better visualize lower values for targeting efficiencies. Error bars represent standard deviation based on biological triplicates and statistical significance is determined using student's t-test and is represented as "\*" for  $P \leq .05$ , "\*\*\*" for  $P \leq .01$ , and "\*\*\*\*" for  $P \leq .001$ . **(C)** Plates depicting the selection for marker free transformants following removal of the FLP-CaMAT1 cassette enabling marker recycling. Wild-type and a nonflipped colony are taken as controls for selection on cloNAT containing medium.

only one selectable marker to work with and shows how difficult it can be to transfer knowledge and tools from one yeast to another.

Many yeast, including *C. intermedia*, display a high preference for ectopic integrations of foreign DNA molecules. In this study, we stepwise improved the HR frequency, where our first attempt to synchronize cells in the S-phase only helped to some degree (an increase from <1% to 3.7%). This method has proved efficient in other yeast and could possibly have been optimized further also for *C. intermedia*. However, the split-marker approach had a much larger positive effect on the HR/NHEJ ratio, as it increased efficiency by 50-fold as compared to linear DNA fragments. Such high HR efficiency using the split-marker approach circumvents the need to construct *ku70/80* mutants, which is both labor-intensive and limits molecular characterization in other strain backgrounds. It also avoids potential loss of fitness associated with NHEJ deficient mutants in some organisms, which

may interfere with phenotyping of mutants in additional genes (Fraczek et al. 2018).

Unfortunately, the positive effect of the split-marker approach on HR frequency was largely lost when we applied it together with the FLP-CaMAT1 cassette that enables marker-recycling and creation of marker-free transformants. Štafa et al. (2017) have shown that the size-ratio of transforming DNA to that of the homology arms can be a determining factor for the difference in targeting efficiency in *S. cerevisiae*. This might explain why we observed a significant difference in targeting efficiency when we switched from 1000 bp to 50 bp homology arms using the splitCaMAT1 cassette as well as from the splitCaMAT1 cassette (3200 bp in size) to the larger splitFLP-CaMAT1 cassette (6100 bp), both flanked by homology arms of 1000 bp on each side of the cassette. Optimization of this approach could, therefore, involve downsizing the cassette and possibly flanking it with even longer homology arms. Nevertheless, we successfully constructed double deletion mutants



using the FLP-CaMAT1 cassette and further use of this cassette could be to construct auxotrophic strains that can be used as a background in targeting other genes.

CRISPR/Cas9 technology holds great potential to revolutionize marker-free genome editing in nonconventional yeast, although development of these tools for new species can still be a highly challenging task. Besides having tools and protocols for transformation and selection, there is also a need to express functional and properly localized Cas9 proteins and sgRNAs, for which positive controls are often missing. Efficient CRISPR/Cas9-based genome editing is also limited by a low HR/NHEJ ratio in many species, forcing researchers to construct and screen for NHEJ-deficient strains. Using the split-marker approach for initial deletion of *ku70* or *ku80* can, therefore, pave the way for more efficient CRISPR/Cas9-based protocols, not limited to use in *C. intermedia*.

To summarize, in this study we have presented a simple gene deletion method based on the split-marker approach, which can be applicable for genome editing in virtually any yeast with preference for NHEJ over HR. Collectively, the results expand our knowledge on genome editing in yeast in general and *C. intermedia* in particular. The genetic tools developed herein will facilitate detailed molecular characterization of this yeast, both to decipher the genetics underlying *C. intermedia*'s capacity to metabolize different sugars and to develop it into an efficient cell factory for industrial use.

## Authors' contributions

Conceptualization: K.V.R.P., F.F.O., and C.G.; methodology: K.V.R.P., F.F.O., N.P., and C.G.; investigation: all; original manuscript draft preparation: K.V.R.P. and C.G.; and manuscript review and editing: all.

## Acknowledgments

The authors would like to thank Julia Köhler for generously providing the CaMAT1-containing integrative plasmids that made this work possible.

**Conflict of interest statement.** The authors declare no conflict of interest.

## Funding

This research was funded by Formas, grant number 2017-01417.

## Data availability statement

Not applicable.

## References

- Acuña-Fontecilla A, Silva-Moreno E, Ganga MA et al. Evaluation of antimicrobial activity from native wine yeast against food industry pathogenic microorganisms. *CyTA J Food* 2017;**15**: 457–65.
- Baptista M, Domingues L. *Kluyveromyces marxianus* as a microbial cell factory for lignocellulosic biomass valorisation. *Biotechnol Adv* 2022;**60**:108027.
- Barnes G, Rio D. DNA double-strand-break sensitivity, DNA replication, and cell cycle arrest phenotypes of Ku-deficient *Saccharomyces cerevisiae*. *Proc Natl Acad Sci USA* 1997;**94**:867–72.
- Cao M, Gao M, Ploessl D et al. CRISPR-mediated genome editing and gene repression in *Scheffersomyces stipitis*. *Biotechnol J* 2018;**13**:e1700598.
- Catlett NL, Lee B-N, Yoder O et al. Split-marker recombination for efficient targeted deletion of fungal genes. *Fung Genet Rep* 2003;**50**: 9–11.
- Corrigan MW, Kerwin-Iosue CL, Kuczmarski AS et al. The fate of linear DNA in *Saccharomyces cerevisiae* and *Candida glabrata*: the role of homologous and non-homologous end joining. *PLoS ONE* 2013;**8**:e69628.
- da Silva Ferreira ME, Kress MR, Savoldi M et al. The *akuB*<sup>KU80</sup> mutant deficient for nonhomologous end joining is a powerful tool for analyzing pathogenicity in *Aspergillus fumigatus*. *Eukaryot Cell* 2006;**5**:207–11.
- De Backer MD, Maes D, Vandoninck S et al. Transformation of *Candida albicans* by electroporation. *Yeast* 1999;**15**:1609–18.
- Detloff P, Sieber J, Petes TD. Repair of specific base pair mismatches formed during meiotic recombination in the yeast *Saccharomyces cerevisiae*. *Mol Cell Biol* 1991;**11**:737–45.
- Dudášová Z, Dudáš A, Chovanec M. Non-homologous end-joining factors of *Saccharomyces cerevisiae*. *FEMS Microbiol Rev* 2004;**28**:581–601.
- Dujon B, Sherman D, Fischer G et al. Genome evolution in yeasts. *Nature* 2004;**430**:35–44.
- Enkler L, Richer D, Marchand AL et al. Genome engineering in the yeast pathogen *Candida glabrata* using the CRISPR-Cas9 system. *Sci Rep* 2016;**6**:35766.
- Ennis CL, Hernday AD, Nobile CJ. A markerless CRISPR-mediated system for genome editing in *Candida auris* reveals a conserved role for Cas5 in the caspofungin response. *Microbiol Spectr* 2021;**9**:e0182021.
- Fairhead C, Llorente B, Denis F et al. New vectors for combinatorial deletions in yeast chromosomes and for gap-repair cloning using 'split-marker' recombination. *Yeast* 1996;**12**:1439–57.
- Fonseca C, Olofsson K, Ferreira C et al. The glucose/xylose facilitator Gxf1 from *Candida intermedia* expressed in a xylose-fermenting industrial strain of *Saccharomyces cerevisiae* increases xylose uptake in SSCF of wheat straw. *Enzyme Microb Technol* 2011;**48**:518–25.
- Fraczek MG, Naseeb S, Delneri D. History of genome editing in yeast. *Yeast* 2018;**35**:361–8.
- Gardonyi M, Osterberg M, Rodrigues C et al. High capacity xylose transport in *Candida intermedia* PYCC 4715. *FEMS Yeast Res* 2003;**3**:45–52.
- Geijer C, Faria-Oliveira F, Moreno AD et al. Genomic and transcriptomic analysis of *Candida intermedia* reveals the genetic determinants for its xylose-converting capacity. *Biotechnol Biofuels* 2020;**13**:48.
- Gietz RD, Schiestl RH. High-efficiency yeast transformation using the LiAc/SS carrier DNA/PEG method. *Nat Protoc* 2007;**2**:31–34.
- Haarmann T, Lorenz N, Tudzynski P. Use of a nonhomologous end joining deficient strain ( $\Delta ku70$ ) of the ergot fungus *Claviceps purpurea* for identification of a nonribosomal peptide synthetase gene involved in ergotamine biosynthesis. *Fung Genet Biol* 2008;**45**:35–44.
- Hara A, Ueda M, Misawa S et al. A mutated hygromycin resistance gene is functional in the n-alkane-assimilating yeast *Candida tropicalis*. *Arch Microbiol* 2000;**173**:187–92.
- Jinek M, Chylinski K, Fonfara I et al. A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. *Science* 2012;**337**:816–21.
- Kawai S, Hashimoto W, Murata K. Transformation of *Saccharomyces cerevisiae* and other fungi: methods and possible underlying mechanism. *Bioeng Bugs* 2010;**1**:395–403.

- Kohler JR, Fink GR. *Candida albicans* strains heterozygous and homozygous for mutations in mitogen-activated protein kinase signaling components have defects in hyphal development. *Proc Natl Acad Sci USA* 1996;**93**:13223–8.
- Kooistra R, Hooykaas PJ, Steensma HY. Efficient gene targeting in *Kluyveromyces lactis*. *Yeast* 2004;**21**:781–92.
- Krejci L, Altmannova V, Spirek M et al. Homologous recombination and its regulation. *Nucleic Acids Res* 2012;**40**:5795–818.
- Kretzschmar A, Otto C, Holz M et al. Increased homologous integration frequency in *Yarrowia lipolytica* strains defective in non-homologous end-joining. *Curr Genet* 2013;**59**:63–72.
- Lobs AK, Schwartz C, Wheeldon I. Genome and metabolic engineering in non-conventional yeasts: current advances and applications. *Synth Syst Biotechnol* 2017;**2**:198–207.
- Mancera E, Frazer C, Porman AM et al. Genetic modification of closely related *Candida* species. *Front Microbiol* 2019;**10**:357.
- Millerioux Y, Clastre M, Simkin AJ et al. Drug-resistant cassettes for the efficient transformation of *Candida guilliermondii* wild-type strains. *FEMS Yeast Res* 2011;**11**:457–63.
- Moreno-Beltran M, Gore-Lloyd D, Chuck C et al. Variation among *Metschnikowia pulcherrima* isolates for genetic modification and homologous recombination. *Microorganisms* 2021;**9**:290.
- Morschhauser J, Michel S, Staib P. Sequential gene disruption in *Candida albicans* by FLP-mediated site-specific recombination. *Mol Microbiol* 1999;**32**:547–56.
- Muhlhausen S, Schmitt HD, Plessmann U et al. Proteogenomics analysis of CUG codon translation in the human pathogen *Candida albicans*. *BMC Biol* 2021;**19**:258.
- Nelson RT, Pryor BA, Lodge JK. Sequence length required for homologous recombination in *Cryptococcus neoformans*. *Fung Genet Biol* 2003;**38**:1–9.
- Ng H, Dean N. Dramatic improvement of CRISPR/Cas9 editing in *Candida albicans* by increased single guide RNA expression. *mSphere* 2017;**2**:e00385–16.
- Norton EL, Sherwood RK, Bennett RJ. Development of a CRISPR-Cas9 system for efficient genome editing of *Candida lusitanae*. *mSphere* 2017;**2**:e00217–17.
- Orr-Weaver TL, Szostak JW, Rothstein RJ. Yeast transformation: a model system for the study of recombination. *Proc Natl Acad Sci USA* 1981;**78**:6354–8.
- Papon N, Courdavault V, Clastre M et al. *Deus ex Candida* genetics: overcoming the hurdles for the development of a molecular toolbox in the CTG clade. *Microbiology* 2012;**158**:585–600.
- Pena R, Ganga MA. Novel antimicrobial peptides produced by *Candida intermedia* LAMAP1790 active against the wine-spoilage yeast *Brettanomyces bruxellensis*. *Antonie Van Leeuwenhoek* 2019;**112**:297–304.
- Ramírez-Orozco M, Hernández-Saavedra NY, Ochoa J-L. *Debaryomyces hansenii* growth in nonsterile seawater CIO2 - peptone-containing medium. *Can J Microbiol* 2001;**47**:676–9.
- Raschmanová H, Weninger A, Glieder A et al. Implementing CRISPR-Cas technologies in conventional and non-conventional yeasts: current state and future prospects. *Biotechnol Adv* 2018;**36**:641–65.
- Sambrook J, Russell DW. Preparation and transformation of competent *E. coli* using calcium chloride. *CSH Protoc* 2006;**2006**:pdb.prot3932.
- Sfeir A, Symington LS. Microhomology-mediated end joining: a backup survival mechanism or dedicated pathway? *Trends Biochem Sci* 2015;**40**:701–14.
- Shen J, Guo W, Kohler JR. CaNAT1, a heterologous dominant selectable marker for transformation of *Candida albicans* and other pathogenic *Candida* species. *Infect Immun* 2005;**73**:1239–42.
- Shen XX, Opulente DA, Kominek J et al. Tempo and mode of genome evolution in the budding yeast subphylum. *Cell* 2018;**175**:1533–45e1520.
- Štafa A, Miklenić MS, Zandona A et al. In *Saccharomyces cerevisiae* gene targeting fidelity depends on a transformation method and proportion of the overall length of the transforming and targeted DNA. *FEMS Yeast Res* 2017;**17**:fox041.
- Strucko T, Andersen NL, Mahler MR et al. A CRISPR/Cas9 method facilitates efficient oligo-mediated gene editing in *Debaryomyces hansenii*. *Synth Biol* 2021;**6**:ysab031.
- Šuchová K, Fehér C, Ravn JL et al. Cellulose- and xylan-degrading yeasts: enzymes, applications and biotechnological potential. *Biotechnol Adv* 2022;**59**:107981.
- Tsakraklides V, Brevnova E, Stephanopoulos G et al. Improved gene targeting through cell cycle synchronization. *PLoS ONE* 2015;**10**:e0133434.
- Uthayakumar D, Sharma J, Wensing L et al. CRISPR-based genetic manipulation of *Candida* species: historical perspectives and current approaches. *Front Genome Ed* 2021;**2**:606281.
- Vyas VK, Barrasa MI, Fink GR. A *Candida albicans* CRISPR system permits genetic engineering of essential genes and gene families. *Sci Adv* 2015;**1**:e1500248.
- Westfall PJ, Pitera DJ, Lenihan JR et al. Production of amorpha-14:15-diene in yeast, and its conversion to dihydroartemisinic acid, precursor to the antimalarial agent artemisinin. *Proc Natl Acad Sci USA* 2012;**109**:E111–8.
- Wilson TE. Nonhomologous end-joining: mechanisms, conservation and relationship to illegitimate recombination. In: Aguilera A, Rothstein R (eds), *Molecular Genetics of Recombination*. Berlin, Heidelberg: Springer, 2007, 487–513.
- Wu J, Hu J, Zhao S et al. Single-cell protein and xylitol production by a novel yeast strain *Candida intermedia* FL023 from lignocellulosic hydrolysates and xylose. *Appl Biochem Biotechnol* 2018;**185**:163–78.
- Yonten V, Aktas N. Exploring the optimum conditions for maximizing the microbial growth of *Candida intermedia* by response surface methodology. *Prep Biochem Biotechnol* 2014;**44**:26–39.
- Zhang G, Zbed HM, An Y et al. Biocatalytic conversion of a lactose-rich dairy waste into D-tagatose, D-arabitol and galactitol using sequential whole cell and fermentation technologies. *Bioresour Technol* 2022;**358**:127422.