



On the parabolic Cauchy problem for quantum graphs with vertex noise

Downloaded from: <https://research.chalmers.se>, 2025-12-04 09:45 UTC

Citation for the original published paper (version of record):

Kovacs, M., Sikolya, E. (2023). On the parabolic Cauchy problem for quantum graphs with vertex noise. *Electronic Journal of Probability*, 28. <http://dx.doi.org/10.1214/23-EJP962>

N.B. When citing this work, cite the original published paper.

On the parabolic Cauchy problem for quantum graphs with vertex noise*

Mihály Kovács[#] Eszter Sikolya[§]

Abstract

We investigate the parabolic Cauchy problem associated with quantum graphs including Lipschitz or polynomial type nonlinearities and additive Gaussian noise perturbed vertex conditions. The vertex conditions are the standard continuity and Kirchhoff assumptions in each vertex. In the case when only Kirchhoff conditions are perturbed, we can prove existence and uniqueness of a mild solution with continuous paths in the standard state space \mathcal{H} of square integrable functions on the edges. We also show that the solution is Markov and Feller. Furthermore, assuming that the vertex values of the normalized eigenfunctions of the self-adjoint operator governing the problem are uniformly bounded, we show that the mild solution has continuous paths in the fractional domain space associated with the Hamiltonian operator, \mathcal{H}_α for $\alpha < \frac{1}{4}$. This is the case when the Hamiltonian operator is the standard Laplacian perturbed by a potential. We also show that if noise is present in both type of vertex conditions, then the problem admits a mild solution with continuous paths in the fractional domain space \mathcal{H}_α with $\alpha < -\frac{1}{4}$ only. These regularity results are the quantum graph analogues obtained by da Prato and Zabczyk [9] in case of a single interval and classical boundary Dirichlet or Neumann noise.

Keywords: quantum graph; Hamiltonian operator; white-noise vertex conditions.

MSC2020 subject classifications: Primary 81Q35; 60H15; 35R60, Secondary 35R02; 47D06.

Submitted to EJP on July 7, 2022, final version accepted on May 13, 2023.

Supersedes arXiv:2207.01944.

*M. Kovács acknowledges the support of the Marsden Fund of the Royal Society of New Zealand through grant no. 18-UOO-143, the Swedish Research Council (VR) through grant no. 2017-04274 and the National Research, Development, and Innovation Fund of Hungary under Grant no. TKP2021-NVA-02 and Grant no. K-131545. E. Sikolya was supported by the OTKA grant no. 135241.

[†]Department of Mathematical Sciences, Chalmers University of Technology and University of Gothenburg, Gothenburg, Sweden.

[‡]Department of Differential Equations, Budapest University of Technology and Economics, Budapest, Hungary and Faculty of Information Technology and Bionics, Pázmány Péter Catholic University, Budapest, Hungary.

E-mail: mihaly@chalmers.se

[§]Institute of Mathematics, Eötvös Loránd University, Budapest, Hungary.

E-mail: eszter.sikolya@ttk.elte.hu

1 Introduction

We consider a so-called quantum graph; that is, is a metric graph G , equipped with a diffusion operator on each edge and certain vertex conditions. Our terminology follows [3, Chap. 1] (see also [21]), we list here only the most important concepts. The graph G consists of a finite set of vertices $V = \{v\}$ and a finite set $E = \{e\}$ of edges connecting the vertices. We denote by $m = |E|$ the number of edges and by $n = |V|$ the number of vertices. In general, a metric graph is assumed to have directed edges; that is edges having an origin and a terminal vertex. In our case, dealing with self-adjoint operators, we can just consider undirected edges. Each edge is assigned a positive length $\ell_e \in (0, +\infty)$, and we denote by $x \in [0, \ell_e]$ a coordinate of G . We assume that G is simple; that is, there are no multiple edges connecting two vertices, and there are no loops at any of the vertices in G .

The metric graph structure enables one to speak about functions u on G , defined along the edges such that for any coordinate x , the function takes its value $u(x)$. If we emphasize that x is taken from the edge e , we write $u_e(x)$. Thus, a function u on G can be regarded as a vector of functions that are defined on the edges, therefore we will also write

$$u = (u_e)_{e \in E},$$

and consider it as an element of a product function space.

To write down the vertex conditions in the form of equations, for a given function u on G and for each $v \in V$, we introduce the following notation. For any $v \in V$, we denote by E_v the set of edges incident to the vertex v , and by $d_v = |E_v|$ the degree of v . Let $u_e(v)$ denote the value of u in v along the edge e in the case $e \in E_v$. Let $E_v = \{e_1, \dots, e_{d_v}\}$, and define

$$U(v) = (u_e(v))_{e \in E_v} = \begin{pmatrix} u_{e_1}(v) \\ \vdots \\ u_{e_{d_v}}(v) \end{pmatrix} \in \mathbb{R}^{d_v}, \quad (1.1)$$

the vector of the function values in the vertex v .

Let I_v be the bi-diagonal matrix

$$I_v = \begin{pmatrix} 1 & -1 & & \\ & \ddots & \ddots & \\ & & 1 & -1 \end{pmatrix} \in \mathbb{R}^{(d_v-1) \times d_v}.$$

It is easy to see that if we set

$$I_v U(v) = 0_{\mathbb{R}^{d_v-1}}, \quad (1.2)$$

this means that all the function values coincide in v . If it is satisfied for each vertex $v \in V$ for a function u on G which is continuous on each edge, including the one-sided continuity at the endpoints, then we call u is continuous on G .

Similarly, for a function u on G which is differentiable on each edge, that is, u'_e exists for each $e \in E$ including the one-sided derivatives at the endpoints, we set

$$U'(v) = (u'_e(v))_{e \in E_v} = \begin{pmatrix} u'_{e_1}(v) \\ \vdots \\ u'_{e_{d_v}}(v) \end{pmatrix} \in \mathbb{R}^{d_v}, \quad (1.3)$$

the vector of the function derivatives in the vertex v . We will assume throughout the paper that derivatives are taken in the directions away from the vertex v (i.e. into the edge), see [3, Sec. 1.4.].

First we aim to analyse the existence, uniqueness and regularity of solutions of the problem written formally as

$$\begin{cases} \dot{u}_e(t, x) = (c_e u'_e)'(t, x) - p_e(x) u_e(t, x) + F_e(u_e(t, x)), & x \in (0, \ell_e), t \in (0, T], e \in E, \quad (a) \\ 0 = I_v U(t, v), & t \in (0, T], v \in V, \quad (b) \\ \dot{\beta}_v(t) = C(v)^\top U'(t, v), & t \in (0, T], v \in V, \quad (c) \\ u_e(0, x) = u_{0,e}(x), & x \in [0, \ell_e], e \in E, \quad (d) \end{cases} \quad (1.4)$$

where $F_e: \mathbb{R} \rightarrow \mathbb{R}$ is (globally) Lipschitz and $T > 0$ is arbitrary but fixed. However, at the end of the paper we comment on the non-Lipschitz case as well.

Here \dot{u}_e and u'_e denote the time and space derivative, respectively, of u_e . For each $v \in V$, $U(t, v)$ and $U'(t, v)$ denote the vector of the function values in v introduced in (1.1) and (1.3), respectively, for the function on G defined as

$$u(t, \cdot) = (u_e(t, \cdot))_{e \in E}.$$

Analogously to (1.1), for each $v \in V$ and $E_v = \{e_1, \dots, e_{d_v}\}$ the 1-row matrix in (1.6c) is defined by

$$C(v)^\top = (c_{e_1}(v), \dots, c_{e_{d_v}}(v)) \in \mathbb{R}^{1 \times d_v}.$$

Equations (1.6b) assume continuity of the function $u(t, \cdot)$ on the metric graph G , cf. (1.2).

It is clear by definition, that (1.6b) consists of

$$\sum_{v \in V} (d_v - 1) = 2m - n \quad (1.5)$$

equations. At the same time, (1.6c) consists of n equations. Hence, we have altogether $2m$ (boundary or vertex) conditions in the vertices.

Here $(\Omega, \mathcal{F}, \mathbb{P})$ is a complete probability space endowed with a right-continuous filtration $\mathbb{F} = (\mathcal{F}_t)_{t \in [0, T]}$, whilst the process

$$(\beta(t))_{t \in [0, T]} = \left((\beta_v(t))_{t \in [0, T]} \right)_{v \in V},$$

is an \mathbb{R}^n -valued Brownian motion (Wiener process) with covariance matrix

$$\tilde{Q} \in \mathbb{R}^{n \times n}$$

with respect to the filtration \mathbb{F} ; that is, $(\beta(t))_{t \in [0, T]}$ is $(\mathcal{F}_t)_{t \in [0, T]}$ -adapted and for all $t > s$, $\beta(t) - \beta(s)$ is independent of \mathcal{F}_s .

The functions c_e are (variable) diffusion coefficients or conductances, and we assume that

$$c_e \in C[0, \ell_e], \quad 0 < c_0 \leq c_e(x), \quad \text{for all } x \in [0, \ell_e], \text{ for all } e \in E.$$

The functions p_e are non-negative, bounded functions, hence

$$0 \leq p_e \in L^\infty(0, \ell_e), \quad e \in E.$$

In equation (1.6d) we pose the initial conditions on the edges.

In Section 4 we omit the nonlinear drift term in (1.4a) and perturb (1.4b) using a white-noise-term $\dot{\Phi}_v$ for all $v \in V$ as, in this case, it turns out that the solution process even to the linear equation is too irregular in space to include a Nemytskii-type nonlinearity. Hence, we investigate the problem

$$\begin{cases} \dot{u}_e(t, x) = (c_e u'_e)'(t, x) - p_e(x) u_e(t, x), & x \in (0, \ell_e), t \in (0, T], e \in E, \quad (a) \\ \dot{\Phi}_v(t) = I_v U(t, v), & t \in (0, T], v \in V, \quad (b) \\ \dot{\beta}_v(t) = C(v)^\top U'(t, v), & t \in (0, T], v \in V, \quad (c) \\ u_e(0, x) = u_{0,e}(x), & x \in [0, \ell_e], e \in E. \quad (d) \end{cases} \quad (1.6)$$

Here the process

$$\Theta := \begin{pmatrix} (\Phi_v)_{v \in V} \\ (\beta_v)_{v \in V} \end{pmatrix}$$

is an \mathbb{R}^{2m} -valued Brownian motion (Wiener process) with covariance matrix

$$Q \in \mathbb{R}^{2m \times 2m} \quad (1.7)$$

with respect to the filtration \mathbb{F} . The main motivation to consider (1.4) and (1.6) is to generalize the classical concept of boundary noise, see, for example, [6, 9] for Gaussian noise and [23, Sec. 15.1] for Lévy noise, to quantum graphs, see also [5].

The paper is organized as follows. In Section 2 we first investigate the linear deterministic version of (1.4) (that is, of (1.6)). We rewrite the system in the form of an abstract Cauchy problem governed by the operator A and prove well-posedness by showing that A generates a strongly continuous, analytic contraction semigroup on the Hilbert space \mathcal{H} of L^2 -functions on the edges, see Proposition 2.3. The statement of Proposition 2.2 is the existence and boundedness of the so-called Dirichlet-operator. Using this and results from [9], we can treat two versions of the stochastic problem.

In Section 3 we consider the problem (1.4). In Theorem 3.5 we prove the existence and uniqueness of the mild solution for this problem with continuous paths in the Hilbert space \mathcal{H} . We also verify that this solution is Markov and Feller. Assuming that the vertex values of the eigenfunctions of A are uniformly bounded, we can show in Theorem 3.8 that the mild solution has continuous paths in the fractional domain space of order $\alpha < \frac{1}{4}$. This is the case, for example, if all diffusion coefficients and all edge lengths are constant and equal 1, see Example 3.10. In Remark 3.11 we treat the case of odd-degree polynomial type nonlinearities.

In Section 4 we briefly investigate the problem (1.6) where all the boundary conditions are perturbed by some noise, but due to the low regularity shown, only in the linear case. In Theorem 4.1 we prove that the stochastic convolution process has values in the fractional domain space of A of order $\alpha < -\frac{1}{4}$ only. Therefore, Nemytskii type nonlinearities, as in the previous section, cannot be considered as point evaluation is not well-defined anymore.

The regularity results obtained in this paper are the analogues of the ones obtained by Da Prato and Zabczyk in [9] in case of a single interval and classical boundary Dirichlet or Neumann noise.

2 Heat equation on a network

2.1 The abstract Cauchy problem

We start with the deterministic problem

$$\begin{cases} \dot{u}_e(t, x) = (c_e u'_e)'(t, x) - p_e(x) u_e(t, x), & x \in (0, \ell_e), t > 0, e \in E, \quad (a) \\ 0 = I_v U(t, v), & t > 0, v \in V, \quad (b) \\ 0 = C(v)^\top U'(t, v), & t > 0, v \in V, \quad (c) \\ u_e(0, x) = u_{0,e}(x), & x \in [0, \ell_e], e \in E. \quad (d) \end{cases} \quad (2.1)$$

where 0 denotes the constant 0 vector of dimension $d_v - 1$ on the left-hand-side of (2.1b).

We would like to rewrite our system in the form of an abstract Cauchy problem. First we consider the Hilbert space

$$\mathcal{H} := \prod_{e \in E} L^2(0, \ell_e)$$

as the *state space* of the edges, endowed with the natural inner product

$$\langle f, g \rangle_{\mathcal{H}} := \sum_{e \in E} \int_0^{\ell_e} f_e(x) g_e(x) dx, \quad f = (f_e)_{e \in E}, \quad g = (g_e)_{e \in E} \in \mathcal{H}.$$

On \mathcal{H} we define the operator

$$A_{\max} := \text{diag} \left(\frac{d}{dx} \left(c_e \frac{d}{dx} \right) - p_e \right)_{e \in E} \quad (2.2)$$

with maximal domain

$$D(A_{\max}) = \mathcal{H}^2 := \prod_{e \in E} H^2(0, \ell_e). \quad (2.3)$$

We also introduce the *boundary space*

$$\mathcal{Y} := \ell^2(\mathbb{R}^{2m}) \cong \mathbb{R}^{2m}. \quad (2.4)$$

Notice that for fixed $u \in D(A_{\max})$, the boundary (or vertex) conditions can be written as

$$0_{\mathbb{R}^{d_v-1}} = I_v U(v), \quad 0 = C(v)^{\top} U'(v), \quad v \in V, \quad (2.5)$$

cf. (2.1b) and (2.1c).

Remark 2.1. Define A_v as the square matrix that arises from I_v by inserting an additional row containing only 0's; that is,

$$A_v = \begin{pmatrix} 1 & -1 & & \\ & \ddots & \ddots & \\ & & 1 & -1 \\ 0 & \dots & 0 & 0 \end{pmatrix} \in \mathbb{R}^{d_v \times d_v}.$$

Furthermore, let B_v be the square matrix defined by

$$B_v = \begin{pmatrix} 0 & \dots & 0 \\ \vdots & & \vdots \\ 0 & \dots & 0 \\ c_{e_1}(v) & \dots & c_{e_{d_v}}(v) \end{pmatrix} \in \mathbb{R}^{d_v \times d_v}.$$

It is straightforward that for a fixed $v \in V$, equations (2.5) have the form

$$A_v U(v) + B_v U'(v) = 0_{\mathbb{R}^{d_v}},$$

where the $d_v \times (2d_v)$ matrix (A_v, B_v) has maximal rank. Thus, our vertex conditions have the form as in [3, Sec. 1.4.1].

We now define the feedback operator $B: D(A_{\max}) \rightarrow \mathcal{Y}$ by

$$\begin{aligned} D(B) &= D(A_{\max}); \\ Bu &= \begin{pmatrix} (I_v U(v))_{v \in V} \\ (C(v)^{\top} U'(v))_{v \in V} \end{pmatrix}, \end{aligned} \quad (2.6)$$

where the first “block” of Bu has $\sum_{v \in V} (d_v - 1) = 2m - n$ coordinates (see also (1.5)), while the second “block” has n coordinates. Hence, B maps indeed into \mathbb{R}^{2m} .

With these notations, we can finally rewrite (2.1) in form of an abstract Cauchy problem. Define

$$A := A_{\max} \quad (2.7)$$

$$D(A) := \{u \in D(A_{\max}) : Bu = 0_{\mathcal{Y}}\}. \quad (2.8)$$

Using this, (2.1) becomes

$$\begin{cases} \dot{u}(t) = Au(t), & t > 0, \\ u(0) = u_0, \end{cases}$$

with $u_0 = (u_{0,1}, \dots, u_{0,m})^\top$.

2.2 Well-posedness of the abstract Cauchy problem

Proposition 2.2. *The operators A_{\max} and B satisfy the conditions in [23, Section 15.1] and [9, Section 1], that is*

1. *the operator A defined in (2.7) generates a C_0 -semigroup on \mathcal{H} ;*
2. *for $\lambda \in \rho(A)$ there exists a bounded operator $D_{B,\lambda}: \mathcal{Y} \rightarrow \mathcal{H}$ such that*

$$D_{B,\lambda} = (B|_{\text{Ker}(\lambda - A_{\max})})^{-1}. \quad (2.9)$$

Proof. 1. Follows directly by [11, Sec. 3.2 and Rem. 2.9] or [22, Rem. 3.6].

2. We will show that the assumptions (a)–(d) of [13, (1.13)] are satisfied for A_{\max} and B . Then, by [13, Lemma 1.2] the existence of the bounded operator of $D_{B,\lambda}$ in (2.9) follows for every $\lambda \in \rho(A)$.

The operator $(A_{\max}, D(A_{\max}))$ is densely defined and closed on \mathcal{H} , hence assumption (a) is satisfied. The boundary operator as a mapping

$$B: (D(A_{\max}), \|\cdot\|_{\mathcal{H}^2}) \rightarrow \mathcal{Y}$$

is bounded by Sobolev embedding, which is assumption (b). The statement of (d) is exactly 1. above.

It remains only to prove (c); that is, $\text{Im } B = \mathbb{R}^{2m}$, which is the assertion of Proposition A.1. \square

Proposition 2.3. *The operator $(-A, D(A))$, where A is defined in (2.7), is the operator associated with the form*

$$\begin{aligned} \mathfrak{a}(u, v) &= \sum_{e \in E} \int_0^{\ell_e} c_e(x) u'_e(x) v'_e(x) dx + \sum_{e \in E} \int_0^{\ell_e} p_e(x) u_e(x) v_e(x) dx, \\ D(\mathfrak{a}) &= \{u \in \mathcal{H}^1 : I_v U(v) = 0, v \in V\}, \end{aligned} \quad (2.10)$$

where

$$\mathcal{H}^1 := \prod_{e \in E} H^1(0, \ell_e),$$

in the following sense:

$$\begin{aligned} D(A) &= \{u \in D(\mathfrak{a}) : \exists h \in \mathcal{H} \text{ s.t. } \mathfrak{a}(u, v) = \langle h, v \rangle_{\mathcal{H}} \text{ for all } v \in D(\mathfrak{a})\}, \\ -Au &= h. \end{aligned}$$

The form $(\mathfrak{a}, D(\mathfrak{a}))$ is symmetric, densely defined, continuous, closed and accretive. The operator $(A, D(A))$ is densely defined, dissipative, sectorial and self-adjoint with $(0, +\infty) \subset \rho(A)$. The strongly continuous semigroup $(S(t))_{t \geq 0}$ generated by $(A, D(A))$ is analytic, positive and contractive.

Proof. All the statements follow by [22, Sec. 3] (see also [17, Prop. 2.3–2.6]). \square

3 Stochastic perturbation of the Kirchhoff-Neumann vertex conditions

In this section we consider the problem

$$\begin{cases} \dot{u}_e(t, x) = (c_e u'_e)'(t, x) - p_e(x) u_e(t, x) + F_e(u_e(t, x)), & x \in (0, \ell_e), t \in (0, T], e \in E, \\ 0 = I_v U(t, v), & t \in (0, T], v \in V, \\ \dot{\beta}_v(t) = C(v)^\top U'(t, v), & t \in (0, T], v \in V, \\ u_e(0, x) = u_{0,e}(x), & x \in [0, \ell_e], e \in E, \end{cases} \quad (3.1)$$

where $F_e: \mathbb{R} \rightarrow \mathbb{R}$ is (globally) Lipschitz, see (1.4).

To treat this problem, on \mathcal{H} we define the maximal operator $(A_{K,\max}, D(A_{K,\max}))$ in a slightly different way than it has been done in (2.2) and (2.3). Namely, we put the continuity conditions in its domain (see also [17]), that is, we set

$$A_{K,\max} := \text{diag} \left(\frac{d}{dx} \left(c_e \frac{d}{dx} \right) - p_e \right)_{e \in E}, \\ D(A_{K,\max}) := \{u \in \mathcal{H}^2: I_v U(v) = 0, v \in V\}.$$

Accordingly, we have to modify the boundary space as

$$\mathcal{Y}_K := \ell^2(\mathbb{R}^n) \cong \mathbb{R}^n,$$

where $n = |V|$ the number of vertices in G , cf. (2.4). The feedback operator becomes $B_K: D(A_{K,\max}) \rightarrow \mathcal{Y}_K$,

$$D(B_K) = D(A_{K,\max}); \\ B_K u := (C(v)^\top U'(v))_{v \in V},$$

see (2.6).

Defining

$$A := A_{K,\max} \quad (3.2)$$

$$D(A) := \{u \in D(A_{K,\max}): B_K u = 0_{\mathcal{Y}_K}\} \quad (3.3)$$

we obtain the same operator as in (2.7).

Mimicking the proof of Proposition 2.2 one obtains the following result.

Proposition 3.1. *The operators $A_{K,\max}$ and B_K satisfy the conditions in [23, Section 15.1] and [9, Section 1], that is*

1. *the operator A defined in (3.2) – that is, in (2.7) – generates a C_0 -semigroup on \mathcal{H} ;*
2. *for $\lambda > 0$ there exists a bounded operator $D_{B_K, \lambda}: \mathcal{Y}_K \rightarrow \mathcal{H}$ such that*

$$D_{B_K, \lambda} = (B_K|_{\text{Ker}(\lambda - A_{K,\max})})^{-1}. \quad (3.4)$$

In order to define the so-called mild solution to (3.1) we first study the stochastic convolution process defined by

$$Z_K(t) := \int_0^t (\lambda - A) S(t-s) D_K d\beta(s), \quad t \in [0, T], \quad (3.5)$$

where $\lambda > 0$ is fixed, $D_K := D_{B_K, \lambda}$ and

$$\beta = (\beta_v)_{v \in V},$$

is an \mathbb{R}^n -valued Brownian motion (Wiener process) with covariance matrix

$$\tilde{Q} \in \mathbb{R}^{n \times n}.$$

To this aim we first introduce the fractional domain spaces of the generator. Since A generates a contractive analytic semigroup, we can define its fractional powers for $\lambda > 0$ and $\alpha \in (0, 1)$. In particular, the fractional domain spaces

$$\mathcal{H}_\alpha := D((\lambda - A)^\alpha), \quad \|u\|_\alpha := \|(\lambda - A)^\alpha u\|, \quad u \in D((\lambda - A)^\alpha) \quad (3.6)$$

are Banach spaces. We fix $\mathcal{H}_0 := \mathcal{H}$.

For $\alpha \in (-1, 0)$ we define the extrapolation spaces \mathcal{H}_α as the completion of \mathcal{H} under the norms $\|u\|_\alpha := \|(\lambda - A)^\alpha u\|$, $u \in \mathcal{H}$.

It is well-known (see e.g. [12, §II.4–5.]) that up to equivalent norms, these spaces are independent of the choice of $\lambda > 0$.

Remark 3.2. Since the semigroup $(S(t))_{t \geq 0}$ is contractive, hence bounded, then by [14, Prop. 3.1.7] we can choose $\lambda = 0$ in (3.6). That is,

$$\mathcal{H}_\alpha \cong D((-A)^\alpha), \quad \alpha \in [0, 1),$$

when $D((-A)^\alpha)$ is equipped with the graph norm.

Theorem 3.3. *The stochastic convolution $Z_K(\cdot)$ given by (3.5) is well-defined,*

$$Z_K \in C([0, T], L^2(\Omega, \mathcal{H}))$$

and has an \mathcal{H} -valued continuous version.

Proof. First we prove that the stochastic process $Z_K(\cdot)$ in (3.5) is well-defined in \mathcal{H} , that is,

$$\int_0^T \left\| (\lambda - A)S(t)D_K \tilde{Q}^{\frac{1}{2}} \right\|_{\text{HS}(\mathcal{Y}_K, \mathcal{H})}^2 dt < +\infty, \quad (3.7)$$

where HS denotes the Hilbert–Schmidt-norm between the appropriate spaces. As we have

$$\left\| (\lambda - A)S(t)D_K \tilde{Q}^{\frac{1}{2}} \right\|_{\text{HS}(\mathcal{Y}_K, \mathcal{H})}^2 \leq \|(\lambda - A)S(t)D_K\|_{\text{HS}(\mathcal{Y}_K, \mathcal{H})}^2 \cdot \text{Tr}(\tilde{Q}),$$

it is enough to verify that

$$\int_0^T \|(\lambda - A)S(t)D_K\|_{\text{HS}(\mathcal{Y}_K, \mathcal{H})}^2 dt < +\infty. \quad (3.8)$$

Using a standard energy argument (or, alternatively, Parseval’s formula), it follows that

$$\int_0^T \|(\lambda - A)S(t)D_K\|_{\text{HS}(\mathcal{Y}_K, \mathcal{H})}^2 dt \leq c_T \cdot \left\| (\lambda - A)^{\frac{1}{2}} D_K \right\|_{\text{HS}(\mathcal{Y}_K, \mathcal{H})}^2. \quad (3.9)$$

Since

$$D_K: \mathcal{Y}_K \rightarrow D(A_{K, \max})$$

and

$$D(A_{K, \max}) = \{u \in \mathcal{H}^2: I_\nu U(\nu) = 0, \nu \in \mathcal{V}\} \hookrightarrow D(\mathfrak{a}),$$

see (2.10), D_K can be regarded as a bounded operator into $D(\mathfrak{a})$.

On the other hand, we obtain that the form

$$\mathfrak{a}_\lambda(u, v) := \mathfrak{a}(u, v) + \lambda \cdot \langle u, v \rangle_{\mathcal{H}}, \quad u, v \in D(\mathfrak{a}) \quad (3.10)$$

is coercive, symmetric and continuous, see Proposition 2.3 and [14, Rem. 7.3.3]. It is straightforward that the operator associated with \mathfrak{a}_λ is $\lambda - A$. For the form-domain $D(\mathfrak{a}_\lambda) = D(\mathfrak{a})$, see (2.10), equipped with the usual \mathcal{H}^1 -norm, we have that

$$D((\lambda - A)^{\frac{1}{2}}) \cong D(\mathfrak{a}) \quad (3.11)$$

holds with equivalence of norms, see e.g. [1, Prop. 5.5.1].

Hence D_K is a bounded linear operator into $D((\lambda - A)^{\frac{1}{2}})$. Since the range of D_K is finite dimensional,

$$\left\| (\lambda - A)^{\frac{1}{2}} D_K \right\|_{\text{HS}(\mathcal{Y}_K, \mathcal{H})} < +\infty$$

holds. Thus, using (3.9), the assertion in (3.8), hence (3.7) holds, the stochastic convolution $Z_K(\cdot)$ is well-defined in \mathcal{H} .

Finally, since the semigroup $(S(t))_{t \geq 0}$ is contractive, [15, Rem. 1] implies the continuity of the trajectories. \square

Next, we define the mild solution of (3.1) as in [23, Sec. 15.1] (and also [9, Sec. 1]). Let

$$u(t)(\cdot) = u(t, \cdot) = (u_e(t, \cdot))_{e \in E}, \quad F = (F_e)_{e \in E}, \quad F(u(s)) = (F_e(u_e(s)))_{e \in E}$$

Definition 3.4. If $X_K(\cdot) \in C([0, T], L^2(\Omega, \mathcal{H}))$, $\{X_K(t)\}_{t \in [0, T]}$ is $(\mathcal{F}_t)_{t \in [0, T]}$ -adapted and for all $t \in [0, T]$,

$$X_K(t) = X_K(t, u_0) = S(t)u_0 + \int_0^t S(t-s)F(X_K(s))ds + Z_K(t), \quad (3.12)$$

\mathbb{P} -almost surely, where Z_K is defined by (3.5), then X_K is called a mild solution of the problem (3.1).

Theorem 3.5. For all $u_0 \in \mathcal{H}$, equation (3.1) has a unique mild solution. The mild solution has an \mathcal{H} -valued continuous version and it is Markov and Feller.

Proof. Existence and uniqueness of a mild solution of (3.1) follow by a simple standard fixed point argument in the closed subspace of $(\mathcal{F}_t)_{t \in [0, T]}$ -adapted processes of the Banach space $C([0, T], L^2(\Omega, \mathcal{H}))$ using Theorem 3.3 which asserts that $Z_K \in C([0, T], L^2(\Omega, \mathcal{H}))$ (Z_K is clearly $(\mathcal{F}_t)_{t \in [0, T]}$ -adapted). Since, also by Theorem 3.3, Z_K has an \mathcal{H} -valued continuous version, it can be easily seen in view of by (3.12), that so does X_K .

To prove that X_K is Markov, we may use the same reasoning as in the proof of [10, Thm. 9.21] with $E = \mathcal{H}$ noting that Z_K has a continuous version by Theorem 3.3.

Finally, let $u_0, v_0 \in \mathcal{H}$ arbitrary given initial values. Then, by (3.12), the contractivity of $(S(t))_{t \geq 0}$ and the global Lipschitz continuity of F , a simple Gronwall argument shows that

$$\sup_{t \in [0, T]} \|X_K(t, u_0) - X_K(t, v_0)\|_{\mathcal{H}}^2 \leq C(T) \cdot \|u_0 - v_0\|_{\mathcal{H}}^2, \text{ almost surely,}$$

and thus X_K is Feller (c.f. [23, Rem. 9.33]). \square

Remark 3.6. Observe that $D(A) \subset \mathcal{H}^1$, and by the Rellich-Kondrachov theorem, $\mathcal{H}^1 \hookrightarrow \mathcal{H}$ is a continuous, compact embedding. If $\lambda \in \rho(A)$ is arbitrary, for the resolvent operator $\text{ran}(R(\lambda, A)) \subset D(A)$ holds, and we obtain that $R(\lambda, A): \mathcal{H} \rightarrow \mathcal{H}$ is bounded and compact. That is, A has compact resolvent and thus A has only point spectrum. Since A is self-adjoint and dissipative, its eigenvalues $(\lambda_k)_{k \in \mathbb{N}}$ form a sequence of negative real numbers and

$$\lambda_k \rightarrow -\infty, \quad k \rightarrow \infty.$$

We may then choose a set $(f_k)_{k \in \mathbb{N}} \subset D(A)$ of eigenfunctions such that

$$Af_k = \lambda_k f_k, \quad k \in \mathbb{N}, \quad (3.13)$$

and the functions $(f_k)_{k \in \mathbb{N}}$ form a complete orthonormal system in \mathcal{H} .

Remark 3.7. In the view of [9, (9)–(11)] we can also consider adding a space-time white noise term to the first equations of (3.1). To obtain the statements of Theorem 3.5 for this new equation, by [9, (17)] it is enough to show that

$$\int_0^T \|S(t)\|_{\text{HS}}^2 dt < \infty.$$

Let $(f_k)_{k \in \mathbb{N}}$ be the complete orthonormal system consisting of eigenfunctions of A , see Remark 3.6. Then we have

$$\begin{aligned} \int_0^T \|S(t)\|_{\text{HS}}^2 dt &= \int_0^T \sum_{k=1}^{\infty} \|S(t)f_k\|^2 dt = \int_0^T \sum_{k=1}^{\infty} e^{-2\lambda_k t} dt \\ &= \int_0^T 1 dt + \int_0^T \sum_{k=2}^{\infty} e^{-2\lambda_k t} dt = T + \sum_{k=2}^{\infty} \frac{1}{2\lambda_k} (1 - e^{-2\lambda_k T}) \\ &\leq T + \sum_{k=2}^{\infty} \frac{1}{2\lambda_k} < \infty \end{aligned}$$

by Proposition B.1.

It turns out that if we assume uniform boundedness for the vertex values of the eigenfunctions (f_k) , the mild solution has a continuous version in \mathcal{H}_α for $\alpha < \frac{1}{4}$.

We introduce now the boundary operator $L: D(L) \rightarrow \mathcal{Y}_K$ defined by

$$\begin{aligned} D(L) &= \left\{ u \in \prod_{e \in E} C[0, \ell_e] : I_v U(v) = 0, v \in V \right\}, \\ Lu &:= (u(v))_{v \in V} \in \mathcal{Y}_K, \end{aligned} \quad (3.14)$$

where $u(v)$ denotes the common vertex value of the function $u \in D(L)$ on the edges incident to v . That is, L assigns to each function u that is continuous on G the vector of the vertex values of u . Observe that $D(A) \subset D(L)$ holds.

Theorem 3.8. Suppose that there exists $c > 0$ such that

$$\|Lf_k\|_{\mathcal{Y}_K}^2 \leq c, \quad k \in \mathbb{N}, \quad (3.15)$$

where $(f_k)_{k \in \mathbb{N}}$ is the complete orthonormal system consisting of eigenfunctions of A (see Remark 3.6) and L is the operator (3.14).

Then, for $\alpha < \frac{1}{4}$ the stochastic convolution process $Z_K(\cdot)$ defined in (3.5) has a continuous version in \mathcal{H}_α .

Proof. By a straightforward modification of [9, Thm. 2.3] to include the covariance matrix \tilde{Q} (see also, [9, Thm. 5.9]), we have to show that if $\alpha < \frac{1}{4}$, then for a fixed $\lambda > 0$ and $D_K = D_{B_K, \lambda}$, there exists $\gamma > 0$ such that

$$\int_0^T t^{-\gamma} \left\| (\lambda - A)S(t)D_K \tilde{Q}^{\frac{1}{2}} \right\|_{\text{HS}(\mathcal{Y}_K, \mathcal{H}_\alpha)}^2 dt < +\infty,$$

As noted before,

$$\left\| (\lambda - A)S(t)D_K \tilde{Q}^{\frac{1}{2}} \right\|_{\text{HS}(\mathcal{Y}_K, \mathcal{H}_\alpha)}^2 \leq \|(\lambda - A)S(t)D_K\|_{\text{HS}(\mathcal{Y}_K, \mathcal{H}_\alpha)}^2 \cdot \text{Tr}(\tilde{Q})$$

and thus it is enough to prove that

$$\int_0^T t^{-\gamma} \|(\lambda - A)S(t)D_K\|_{\text{HS}(\mathcal{Y}_K, \mathcal{H}_\alpha)}^2 dt < +\infty. \quad (3.16)$$

Using Remark 3.2 and proceeding as in [9, Sec. 3], we obtain that (3.16) holds if

$$\sum_{k=1}^{\infty} (\lambda - \lambda_k)^{2\alpha+\gamma+1} \|D_K^* f_k\|_{\mathcal{Y}_K}^2 < \infty. \quad (3.17)$$

We will show that under the assumptions,

$$\|D_K^* f_k\|_{\mathcal{Y}_K}^2 \leq \frac{c}{(\lambda - \lambda_k)^2}, \quad k \in \mathbb{N}, \quad (3.18)$$

where $c \in \mathbb{R}$ is the constant from (3.15). By Proposition B.1, there exist constants $l_1, l_2 > 0$ such that

$$l_1 \cdot k^2 \leq \lambda - \lambda_k \leq l_2 \cdot k^2, \quad k \in \mathbb{N}. \quad (3.19)$$

Hence, if (3.18) holds, by (3.19) the terms of the series (3.17) can be estimated as

$$(\lambda - \lambda_k)^{2\alpha+\gamma+1} \|D_K^* f_k\|_{\mathcal{Y}_K}^2 \leq C \cdot \frac{k^{4\alpha+2\gamma+2}}{k^4}$$

for an appropriate constant $C > 0$. Thus, the series (3.17) converges if and only if

$$4\alpha + 2\gamma + 2 - 4 < -1 \iff 4\alpha + 2\gamma < 1.$$

This means that there exists appropriate $\gamma > 0$ if and only if

$$\alpha < \frac{1}{4}.$$

Now we turn to the proof of (3.18). First notice that

$$\|D_K^* f_k\|_{\mathcal{Y}_K}^2 = \sum_{i=1}^n \langle e_i, D_K^* f_k \rangle_{\mathcal{Y}_K}^2, \quad (3.20)$$

where

$$e_i = \begin{pmatrix} 0 \\ \vdots \\ 1 \\ \vdots \\ 0 \end{pmatrix} \leftarrow i^{\text{th}} \text{ row}$$

the usual i th basis vector in \mathbb{R}^n and $\langle \cdot, \cdot \rangle_{\mathcal{Y}_K}$ denotes the scalar product in \mathcal{Y}_K .

By definition, for each $i = 1, \dots, n$,

$$\langle e_i, D_K^* f_k \rangle_{\mathcal{Y}_K} = \langle D_K e_i, f_k \rangle_{\mathcal{H}} = \langle u^i, f_k \rangle_{\mathcal{H}}, \quad (3.21)$$

where $u^i \in D(A_{K,\max})$

$$A_{K,\max} u^i = \lambda u^i, \quad B_K u^i = e_i, \quad (3.22)$$

see (3.4). Integration by parts then yields

$$\begin{aligned}
 \langle u^i, f_k \rangle_{\mathcal{H}} &= \frac{1}{\lambda} \langle A_{K, \max} u^i, f_k \rangle_{\mathcal{H}} \\
 &= \frac{1}{\lambda} \sum_{e \in E} \int_0^{\ell_e} (c_e(u_e^i)')'(x) \cdot f_{k,e}(x) dx - \frac{1}{\lambda} \sum_{e \in E} \int_0^{\ell_e} p_e(x) \cdot u_e^i(x) \cdot f_{k,e}(x) dx \\
 &= \frac{1}{\lambda} \sum_{e \in E} [c_e(x) \cdot (u_e^i)'(x) \cdot f_{k,e}(x)]_0^{\ell_e} \\
 &\quad - \frac{1}{\lambda} \sum_{e \in E} \left(\int_0^{\ell_e} c_e(x) \cdot (u_e^i)'(x) \cdot f_{k,e}'(x) dx + \int_0^{\ell_e} p_e(x) \cdot u_e^i(x) \cdot f_{k,e}(x) dx \right) \\
 &= \frac{1}{\lambda} \left(\sum_{e \in E} [c_e(x) \cdot (u_e^i)'(x) \cdot f_{k,e}(x)]_0^{\ell_e} - \alpha(u^i, f_k) \right) \\
 &= \frac{1}{\lambda} \left(\sum_{e \in E} [c_e(x) \cdot (u_e^i)'(x) \cdot f_{k,e}(x)]_0^{\ell_e} + \langle u^i, A f_k \rangle_{\mathcal{H}} \right) \\
 &= \frac{1}{\lambda} \left(\sum_{e \in E} [c_e(x) \cdot (u_e^i)'(x) \cdot f_{k,e}(x)]_0^{\ell_e} + \lambda_k \cdot \langle u^i, f_k \rangle_{\mathcal{H}} \right),
 \end{aligned}$$

where we have used Proposition 2.3 and (3.13). This implies that

$$\langle u^i, f_k \rangle_{\mathcal{H}} = \frac{1}{\lambda - \lambda_k} \sum_{e \in E} [c_e(x) \cdot (u_e^i)'(x) \cdot f_{k,e}(x)]_0^{\ell_e}.$$

Since $f_k \in D(L)$, we have by similar calculations as in the proof of [18, Prop. A.1]

$$\langle u^i, f_k \rangle_{\mathcal{H}} = \frac{1}{\lambda - \lambda_k} \langle B_K u^i, L f_k \rangle_{\mathcal{Y}_K} = \frac{1}{\lambda - \lambda_k} \langle e_i, L f_k \rangle_{\mathcal{Y}_K}, \quad (3.23)$$

where we used (3.22).

Hence by (3.15), (3.20), (3.21) and (3.23) we obtain

$$\|D_K^* f_k\|_{\mathcal{Y}_K}^2 = \frac{1}{(\lambda - \lambda_k)^2} \sum_{i=1}^n \langle e_i, L f_k \rangle_{\mathcal{Y}_K}^2 = \frac{\|L f_k\|_{\mathcal{Y}_K}^2}{(\lambda - \lambda_k)^2} \leq \frac{c}{(\lambda - \lambda_k)^2},$$

which is exactly (3.18) and the proof is complete. \square

Corollary 3.9. *Under the hypothesis of Theorem 3.8, if $u_0 \in \mathcal{H}_\alpha$ for some $\alpha < \frac{1}{4}$, then the unique mild solution of equation (3.1) has a version with continuous paths in \mathcal{H}_α .*

Proof. By Theorem 3.8 Z_K has a version with continuous paths in \mathcal{H}_α for $\alpha < \frac{1}{4}$, and in particular, this version is \mathcal{H} -continuous. For all $\omega \in \Omega$ we then obtain a unique solution $\tilde{X}_K(\cdot, \omega)$ of (3.12) using a standard fixed point argument in the Banach space $C([0, T], \mathcal{H})$. By uniqueness, $\tilde{X}_K(\cdot, \omega)$ is a version of the mild solution of (3.1). On the right-hand side of (3.12) the first and the last terms are elements of $C([0, T], \mathcal{H}_\alpha)$ for α for all $\omega \in \Omega$. Finally, the convolution term on the right-hand-side of (3.12), with X_K replaced by \tilde{X}_K , is an element of the space $C([0, T], \mathcal{H}_\eta)$ by [25, Lem. 3.6] for all $\eta < 1$, as $\tilde{X}_K(\cdot, \omega) \in C([0, T], \mathcal{H})$, F has linear growth and the semigroup $(S(t))_{t \geq 0}$ is analytic. Therefore, \tilde{X}_K has continuous paths in \mathcal{H}_α . \square

Example 3.10. Let $c_e \equiv 1$ and $\ell_e \equiv 1$, $e \in E$ in (3.1). Then (3.1) has a mild solution of the form (3.12) with continuous paths in \mathcal{H}_α for $\alpha < \frac{1}{4}$.

Proof. By [7, Lem. 2.1] and [16, Sec. 2], we have that

$$\|Lf_k\|_{\mathcal{Y}_K}^2 \leq c, \quad k \in \mathbb{N},$$

hence (3.15) holds with an appropriate positive c . Thus, by Theorem 3.8, problem (3.1) has a mild solution of the form (3.12) with continuous paths in \mathcal{H}_α for $\alpha < \frac{1}{4}$. \square

Remark 3.11. In the first equations of (3.1) we may also consider odd-degree polynomial nonlinearities of the form

$$F_e(x) = -x^{2k_e+1} + \sum_{j=0}^{2k_e} a_{e,j}x^j, \quad e \in E$$

as in [9, Example after Thm. 4.2] provided that the hypothesis of Theorem 3.8 is satisfied. Then the mapping $F = (F_e)_{e \in E}$ satisfies the conditions of [9, Thm. 4.2] with

$$K := \prod_{e \in E} L^{2(2k_e+1)}(0, \ell_e), \quad J := \prod_{e \in E} L^{2(2k_e+1)^2}(0, \ell_e).$$

Using Theorem 3.8 and Sobolev's embedding, we obtain that the stochastic convolution process $Z_K(\cdot)$ has a J -continuous version. Therefore, by [9, Thm. 4.2], equation (3.1), has a \mathcal{H} -continuous mild solution. Uniqueness in this case follows the same way as in the proof of [8, Thm. 4.8(i)] using the one sided Lipschitz property

$$\langle F(u) - F(v), u - v \rangle_{\mathcal{H}} \leq C \|u - v\|_{\mathcal{H}}^2, \quad u, v \in \mathcal{H}, \quad C > 0,$$

and the dissipativity of A .

4 Stochastic perturbation of the vertex conditions

Here we briefly investigate the problem where all the boundary conditions are perturbed by some noise, but due to the low regularity shown below, only in the linear case. That is, we consider the problem

$$\begin{cases} \dot{u}_e(t, x) = (c_e u'_e)'(t, x) - p_e(x) u_e(t, x), & x \in (0, \ell_e), \quad t \in (0, T], \quad e \in E, \\ \dot{\Phi}_v(t) = I_v U(t, v), & t \in (0, T], \quad v \in V, \\ \dot{\beta}_v(t) = C(v)^\top U'(t, v), & t \in (0, T], \quad v \in V, \\ u_e(0, x) = u_{0,e}(x), & x \in [0, \ell_e], \quad e \in E, \end{cases}$$

see (1.6). The process

$$\Theta := \begin{pmatrix} (\Phi_v)_{v \in V} \\ (\beta_v)_{v \in V} \end{pmatrix}.$$

is an \mathbb{R}^{2m} -valued Brownian motion (Wiener process) with covariance matrix

$$Q \in \mathbb{R}^{2m \times 2m}. \quad (4.1)$$

For the sake of simplicity we fix again $\lambda > 0$, denote $D := D_{B,\lambda}$, and analogously to (3.5) we define the stochastic convolution process by

$$Z(t) := \int_0^t (\lambda - A) S(t-s) D d\Theta(s).$$

We next prove that $Z(\cdot)$ has values in the fractional domain space of A of order $\alpha < -\frac{1}{4}$ only. Therefore, Nemytskii type nonlinearities, as in the previous section, cannot be considered as point evaluation is not well-defined any more.

Theorem 4.1. *The stochastic convolution process $Z(\cdot)$ has continuous paths in \mathcal{H}_α for $\alpha < -\frac{1}{4}$.*

Proof. Let $\delta > \frac{1}{4}$ arbitrary fixed and we will prove the result for $\alpha := -\delta$. Without loss of generality we may assume that δ is close to $\frac{1}{4}$.

By a straightforward modification of [9, Thm. 2.3] to include the covariance matrix Q , see (4.1), we have to show that for some $\gamma > 0$,

$$\int_0^T t^{-\gamma} \left\| (\lambda - A)S(t)DQ^{\frac{1}{2}} \right\|_{\text{HS}(\mathcal{Y}, \mathcal{H}_{-\delta})}^2 dt < +\infty, \quad (4.2)$$

where HS denotes the Hilbert–Schmidt-norm between the appropriate spaces (see also, [9, Theorem 5.9]). As

$$\left\| (\lambda - A)S(t)DQ^{\frac{1}{2}} \right\|_{\text{HS}(\mathcal{Y}, \mathcal{H}_{-\delta})}^2 \leq \|(\lambda - A)S(t)D\|_{\text{HS}(\mathcal{Y}, \mathcal{H}_{-\delta})}^2 \cdot \text{Tr}(Q),$$

it suffices to prove that

$$\int_0^T t^{-\gamma} \|(\lambda - A)S(t)D\|_{\text{HS}(\mathcal{Y}, \mathcal{H}_{-\delta})}^2 dt < +\infty. \quad (4.3)$$

First we will prove that the Dirichlet operator D maps \mathcal{Y} into \mathcal{H}_ρ for $0 < \rho < \frac{1}{4}$. By (3.11) and Remark 3.2, we have

$$D(\mathfrak{a}) \cong D((\lambda - A)^{\frac{1}{2}}) \cong D((-A)^{\frac{1}{2}})$$

with equivalent norms. Using this, similarly as in [17, Lem. 3.6] one can show that

$$D((-A)^{\frac{1}{2}}) \cong \prod_{e \in E} H_0^1(0, \ell_e) \times \mathbb{R}^n.$$

By [24, Sec. 4.3.3] we have that if $\theta < \frac{1}{2}$, for the complex interpolation space

$$\left[D((-A)^{\frac{1}{2}}), \mathcal{H} \right]_\theta \cong \prod_{e \in E} H^\theta(0, \ell_e) \times \{0_{\mathbb{R}^n}\} \quad (4.4)$$

holds – see also [4] –, where we have used $\mathcal{H} \cong \mathcal{H} \times \{0_{\mathbb{R}^n}\}$ and [2, Thm. 4.2.2]. Furthermore, [14, Thm. 6.6.9] implies that for $0 < \theta < 1$,

$$\left[D((-A)^{\frac{1}{2}}), \mathcal{H} \right]_\theta \cong [D(A), \mathcal{H}]_{\frac{\theta}{2}}. \quad (4.5)$$

Using [1, Thm. in § 4.7.3] and [1, Prop. in § 4.4.10], we obtain that for any $0 < \rho < 1$,

$$[D(A), \mathcal{H}]_\rho \cong D((-A)^\rho) = \mathcal{H}_\rho. \quad (4.6)$$

Combining (4.4), (4.5) and (4.6) yields that for $0 < \rho < \frac{1}{4}$

$$\prod_{e \in E} H^{2\rho}(0, \ell_e) \times \{0_{\mathbb{R}^n}\} \cong \mathcal{H}_\rho \quad (4.7)$$

holds. Hence, by (2.3) and (4.7) we have that for $0 < \rho < \frac{1}{4}$

$$D: \mathcal{Y} \rightarrow D(A_{\max}) \hookrightarrow \prod_{e \in E} H^{2\rho}(0, \ell_e) \cong \prod_{e \in E} H^{2\rho}(0, \ell_e) \times \{0_{\mathbb{R}^n}\} \cong \mathcal{H}_\rho. \quad (4.8)$$

Now we are in the position to prove (4.3). We take a small $\varepsilon > 0$, to be specified later, and estimate the integral in the following way, where we use the analyticity of the semigroup $(S(t))_{t \geq 0}$ from Proposition 2.3,

$$\begin{aligned} & \int_0^T t^{-\gamma} \|(\lambda - A)S(t)D\|_{\text{HS}(\mathcal{Y}, \mathcal{H}_{-\delta})}^2 dt \leq c \cdot \int_0^T t^{-\gamma} \|(\lambda - A)^{1-\delta}S(t)D\|_{\text{HS}(\mathcal{Y}, \mathcal{H})}^2 dt \\ & = c \cdot \int_0^T t^{-\gamma} \|(\lambda - A)^\varepsilon S(t)(\lambda - A)^{1-\delta-\varepsilon}D\|_{\text{HS}(\mathcal{Y}, \mathcal{H})}^2 dt \\ & \leq C_T \cdot \int_0^T t^{-\gamma-2\varepsilon} dt \cdot \|(\lambda - A)^{1-\delta-\varepsilon}D\|_{\text{HS}(\mathcal{Y}, \mathcal{H})}^2. \end{aligned} \quad (4.9)$$

In the last expression,

$$\int_0^T t^{-\gamma-2\varepsilon} dt < \infty \iff \gamma + 2\varepsilon < 1 \iff \varepsilon < \frac{1}{2} - \frac{\gamma}{2}. \quad (4.10)$$

Since the range of D is finite dimensional, if we take a suitable $\varepsilon > 0$ such that $(\lambda - A)^{1-\delta-\varepsilon}D$ is a bounded operator, then the Hilbert-Schmidt norm in (4.9) is finite. To satisfy this, by (4.8),

$$0 < 1 - \delta - \varepsilon < \frac{1}{4} \quad (4.11)$$

should hold. On the other hand, by (4.10),

$$1 - \delta - \varepsilon > \frac{1}{2} + \frac{\gamma}{2} - \delta.$$

Hence, we have to find $\gamma > 0$ such that

$$\frac{1}{2} + \frac{\gamma}{2} - \delta < \frac{1}{4} \iff \frac{1}{4} + \frac{\gamma}{2} < \delta$$

holds true. Since, by assumption, $\delta > \frac{1}{4}$, an appropriate $0 < \gamma < 1$ can be chosen. Taking any $\varepsilon > 0$ satisfying (4.10) and (4.11), that is

$$\frac{3}{4} - \delta < \varepsilon < \frac{1}{2} - \frac{\gamma}{2},$$

the estimate (4.9) yields a finite bound for the left-hand-side of (4.3). This means that for γ chosen this way, (4.3), hence (4.2) is satisfied, and the proof is complete. \square

A Surjectivity of the boundary operator B

In this section we complete the proof of Theorem 2.2.

Proposition A.1. *For the operator $(B, D(B))$ defined in (2.6), $\text{Im } B = \mathbb{R}^{2m}$ holds.*

Proof. We show that for $z \in \mathbb{R}^{2m}$ given arbitrarily, there exists $u \in D(A_{\max}) = \mathcal{H}^2$ (see (2.3)) such that

$$Bu = z. \quad (A.1)$$

We will seek $u \in \mathcal{H}^2$ in the form

$$u(x) = \begin{pmatrix} \alpha_1 e^{-\gamma x} + \beta_1 e^{-\gamma(\ell_1-x)} \\ \vdots \\ \alpha_m e^{-\gamma x} + \beta_m e^{-\gamma(\ell_m-x)} \end{pmatrix} \quad (A.2)$$

for suitable vectors $\alpha = (\alpha_1, \dots, \alpha_m)^\top$, $\beta = (\beta_1, \dots, \beta_m)^\top$ and constant $\gamma > 0$, where ℓ_1, \dots, ℓ_m denote the edges lengths in the graph G .

We introduce the notation

$$u(\ell) := \begin{pmatrix} u(\ell_1) \\ \vdots \\ u(\ell_m) \end{pmatrix}, \quad u'(\ell) := \begin{pmatrix} u'(\ell_1) \\ \vdots \\ u'(\ell_m) \end{pmatrix}, \quad u \in \mathcal{H}^2,$$

and

$$z_C := \begin{pmatrix} z_1 \\ \vdots \\ z_{2m-n} \end{pmatrix}, \quad z_K := \begin{pmatrix} z_{2m-n+1} \\ \vdots \\ z_{2m} \end{pmatrix}$$

for the two “blocks” of the vector $z \in \mathbb{R}^{2m}$. Using the idea of the proof of [19, Prop. 3.2], there exist $(2m-n) \times m$ matrices V_0 and V_1 and $n \times m$ matrices W_0 and W_1 such that equation (A.1) can be rewritten as

$$V_0 u(0) + V_1 u(\ell) = z_C, \quad W_0 u'(0) - W_1 u'(\ell) = z_K. \quad (\text{A.3})$$

A straightforward computation shows that for the function in (A.2), equations (A.3) turn into

$$\begin{aligned} (V_0 + V_1 E_\gamma) \cdot \alpha + (V_0 E_\gamma + V_1) \cdot \beta &= z_C, \\ (-\gamma W_0 + \gamma W_1 E_\gamma) \cdot \alpha + (\gamma W_0 E_\gamma - \gamma W_1) \cdot \beta &= z_K \end{aligned} \quad (\text{A.4})$$

where E_γ is the diagonal matrix

$$E_\gamma := \begin{pmatrix} e^{-\gamma \ell_1} & & \\ & \ddots & \\ & & e^{-\gamma \ell_m} \end{pmatrix}$$

We rewrite now (A.4) with $(2m) \times (2m)$ block-matrices as

$$\begin{pmatrix} V_0 & V_1 \\ -\gamma W_0 & -\gamma W_1 \end{pmatrix} \cdot \begin{pmatrix} \alpha \\ \beta \end{pmatrix} + \begin{pmatrix} V_1 & V_0 \\ \gamma W_1 & \gamma W_0 \end{pmatrix} \cdot \begin{pmatrix} E_\gamma & 0 \\ 0 & E_\gamma \end{pmatrix} \cdot \begin{pmatrix} \alpha \\ \beta \end{pmatrix} = z. \quad (\text{A.5})$$

Denoting by

$$N_\gamma := \begin{pmatrix} V_0 & V_1 \\ -\gamma W_0 & -\gamma W_1 \end{pmatrix}, \quad \tilde{N}_\gamma := \begin{pmatrix} V_1 & V_0 \\ \gamma W_1 & \gamma W_0 \end{pmatrix}, \quad F_\gamma := \begin{pmatrix} E_\gamma & 0 \\ 0 & E_\gamma \end{pmatrix},$$

we have to show that

$$\text{Im} \left(N_\gamma + \tilde{N}_\gamma \cdot F_\gamma \right) = \mathbb{R}^{2m}. \quad (\text{A.6})$$

First we show that N_γ is invertible. Using again ideas from the proof of [19, Prop. 3.2], we can permute rows and columns of N_γ such that we obtain a block diagonal matrix $M_\gamma \in \mathbb{R}^{(2m) \times (2m)}$, consisting of n blocks of size $d_v \times d_v$ for $v \in V$. Denoting by M_v the block corresponding to vertex v in M_γ we have that if $d_v = 1$ then $M_v = -\gamma \cdot c_e(v)$ for $E_v = \{e\}$. Otherwise,

$$M_v = \begin{pmatrix} 1 & -1 & & \\ & \ddots & \ddots & \\ & & 1 & -1 \\ -\gamma \cdot c_{e_1}(v) & \dots & \dots & -\gamma \cdot c_{e_{d_v}}(v) \end{pmatrix}$$

for $E_v = \{e_1, \dots, e_{d_v}\}$. A straightforward computation yields that

$$\det M_v = -\gamma \cdot (c_{e_1}(v) + \dots + c_{e_{d_v}}(v)) \neq 0 \quad (\text{A.7})$$

because of the assumption on the c_e 's. Hence, we obtain that

$$\det M_\gamma = \prod_{v \in V} \det M_v = (-\gamma)^n \cdot K_c \neq 0$$

with

$$K_c = \prod_{\substack{v \in V \\ E_v = \{e_1, \dots, e_{d_v}\}}} (c_{e_1}(v) + \dots + c_{e_{d_v}}(v)).$$

Hence, M_γ is invertible. Since permutations do not change the determinant of a matrix, we also have that N_γ is invertible. That is, in (A.6) we have

$$N_\gamma + \tilde{N}_\gamma \cdot F_\gamma = N_\gamma \cdot (\text{Id} + N_\gamma^{-1} \cdot \tilde{N}_\gamma \cdot F_\gamma) \quad (\text{A.8})$$

with $\text{Id} = \text{Id}_{(2m) \times (2m)}$. If for an appropriate $\gamma > 0$,

$$\|N_\gamma^{-1} \cdot \tilde{N}_\gamma \cdot F_\gamma\|_{\max} < 1 \quad (\text{A.9})$$

is satisfied, then the matrix (A.8) is invertible, hence (A.6) holds.

First we estimate the max-norm of N_γ^{-1} , or, which is the same, the max-norm of M_γ^{-1} . Clearly, M_γ^{-1} is the block-diagonal matrix of blocks M_v^{-1} . Each cofactor $M_{k,l}$ of M_v is the determinant of a matrix of the same type as M_v itself (but having dimension $(d_v - 1) \times (d_v - 1)$), except those in the last row of M_v^{-1} which are all equal to 1. Thus we obtain by (A.7) that for γ big enough,

$$\|M_v^{-1}\|_{\max} = \frac{1}{|\det M_v|} \cdot \max_{k,l=1,\dots,d_v} |M_{k,l}| = \frac{\gamma \cdot L_{c,v}}{\gamma \cdot (c_{e_1}(v) + \dots + c_{e_{d_v}}(v))} =: K_{c,v}$$

with constants $L_{c,v}, K_{c,v} > 0$. Hence, for γ big enough,

$$\|N_\gamma^{-1}\|_{\max} = \|M_\gamma^{-1}\|_{\max} = \max_{v \in V} \|M_v^{-1}\|_{\max} = \max_{v \in V} K_{c,v} =: K_c. \quad (\text{A.10})$$

Similarly as above, we can permute rows and columns of \tilde{N}_γ such that we obtain a block diagonal matrix \tilde{M}_γ consisting of 1×1 blocks $\{\gamma \cdot c_e(v)\}$ and blocks of size $d_v \times d_v$ (for $d_v > 1$)

$$\tilde{M}_v = \begin{pmatrix} 1 & -1 & & \\ & \ddots & \ddots & \\ & & 1 & -1 \\ \gamma \cdot c_{e_1}(v) & \dots & \dots & \gamma \cdot c_{e_{d_v}}(v) \end{pmatrix}.$$

Hence, if γ is big enough,

$$\|\tilde{N}_\gamma\|_{\max} = \|\tilde{M}_\gamma\|_{\max} = \gamma \cdot \max_{\substack{v \in V \\ e \in E_v}} \{c_e(v)\} =: \gamma \cdot \tilde{K}_c. \quad (\text{A.11})$$

Equations (A.10) and (A.11) imply that we have

$$\|N_\gamma^{-1} \cdot \tilde{N}_\gamma \cdot F_\gamma\|_{\max} \leq e^{-\gamma \cdot l} \cdot \gamma \cdot K_c \cdot \tilde{K}_c < 1, \text{ if } \gamma \text{ is big enough,}$$

where l is defined as

$$l := \min_{e \in E} \ell_e > 0$$

Thus for γ big enough (A.9) holds. This implies that (A.6) is true; that is, for arbitrary $z \in \mathbb{R}^{2m}$ there exist $\gamma > 0, \alpha, \beta \in \mathbb{R}^m$ such that (A.5) is satisfied. Hence, by defining u as in (A.2) with the constants $\gamma, \alpha_j, \beta_j, j = 1, \dots, m$, we obtain

$$u \in D(A_{\max}) \text{ and } Bu = z,$$

and the proof is complete. \square

B Asymptotics of the spectrum of A

Proposition B.1. *Let $(\lambda_k)_{k \in \mathbb{N}}$ be the sequence of eigenvalues of the generator $(A, D(A))$ from Remark 3.6. Then for any $\lambda > 0$ there exist constants $l_1, l_2 > 0$ such that*

$$l_1 \cdot k^2 \leq \lambda - \lambda_k \leq l_2 \cdot k^2, \quad k \in \mathbb{N}. \quad (\text{B.1})$$

Proof. Let us fix $\lambda > 0$ and recall that $\lambda - A$ is the operator associated with the coercive, symmetric, continuous form \mathfrak{a}_λ defined in (3.10) by

$$\begin{aligned} \mathfrak{a}_\lambda(f, g) &= \mathfrak{a}(f, g) + \lambda \cdot \langle f, g \rangle_{\mathcal{H}}, \\ D(\mathfrak{a}_\lambda) &= \{u \in \mathcal{H}^1 : I_\nu U(\nu) = 0, \nu \in V\}. \end{aligned}$$

Let $\nu_1 \in V$ arbitrary, and define the form $(\mathfrak{a}^1, D(\mathfrak{a}^1))$ acting in the same way as \mathfrak{a}_λ but having Dirichlet condition in the vertex ν_1 in its domain. That is,

$$\begin{aligned} \mathfrak{a}^1(f, g) &= \mathfrak{a}_\lambda(f, g), \\ D(\mathfrak{a}^1) &= \{u \in \mathcal{H}^1 : I_\nu U(\nu) = 0, \nu \neq \nu_1, U(\nu_1) = 0_{\mathbb{R}^{d_{\nu_1}}}\}, \end{aligned}$$

cf. (1.1). Denote by $(A_1, D(A_1))$ the operator associated with $(\mathfrak{a}^1, D(\mathfrak{a}^1))$. We can now carry out the proof of [3, Thm. 3.1.8] applied to the forms \mathfrak{a}_λ and \mathfrak{a}^1 . We only have to use the facts that $D(\mathfrak{a}^1)$ is a subspace of co-dimension 1 of $D(\mathfrak{a}_\lambda)$, \mathfrak{a}_λ and \mathfrak{a}^1 agree on $D(\mathfrak{a}^1)$, and the “min-max principle” for coercive, symmetric forms holds, see e.g. [20, Thm. 6.5]. Thus we obtain that denoting by $(\lambda_k^1)_{k \in \mathbb{N}}$ the eigenvalues of $(A_1, D(A_1))$ labelled in non-increasing order,

$$\lambda - \lambda_{k+1} \leq \lambda_k^1 \leq \lambda - \lambda_k, \quad k \in \mathbb{N}$$

holds. Continuing this process by defining the finite sequence of symmetric and accretive forms \mathfrak{a}^i , $i = 1, \dots, n$ for an ordering of the vertices $V = \{\nu_1, \dots, \nu_n\}$ such that

$$\begin{aligned} \mathfrak{a}^{i+1}(f, g) &= \mathfrak{a}^i(f, g), \\ D(\mathfrak{a}^{i+1}) &= \{u \in D(\mathfrak{a}^i) : U(\nu_{i+1}) = 0\}, \end{aligned}$$

we can apply the proof of [3, Thm. 3.1.8] for each pair of forms \mathfrak{a}^i and \mathfrak{a}^{i+1} . Hence, if we denote by $(\lambda_k^i)_{k \in \mathbb{N}}$ the eigenvalues of $(A_i, D(A_i))$, the operator associated to $(\mathfrak{a}^i, D(\mathfrak{a}^i))$, labelled in non-increasing order, we obtain that

$$\lambda_{k+1}^i \leq \lambda_k^{i+1} \leq \lambda_k^i, \quad k \in \mathbb{N}, \quad i = 1, \dots, n-1. \quad (\text{B.2})$$

Clearly,

$$\begin{aligned} \mathfrak{a}^n(f, g) &= \mathfrak{a}_\lambda(f, g), \\ D(\mathfrak{a}^n) &= \prod_{e \in E} H_0^1(0, \ell_e). \end{aligned}$$

It is straightforward that the operator $(A_n, D(A_n))$ associated with \mathfrak{a}^n is the operator acting as $\lambda - A$ with Dirichlet conditions in all vertices. Using that $c_e > 0$, $p_e \geq 0$, $e \in E$, and the min-max principle holds, we have that for the set $(\lambda_k^n)_{k \in \mathbb{N}}$ of eigenvalues of A_n there exist constants $l_1, l_2 > 0$ such that

$$l_1 \cdot k^2 \leq \lambda_k^n \leq l_2 \cdot k^2, \quad k \in \mathbb{N},$$

see also [20, Probl. 6.1]. Thus, by (B.2) also (B.1) holds which finishes the proof. \square

References

- [1] Wolfgang Arendt, *Semigroups and evolution equations: functional calculus, regularity and kernel estimates*, Evolutionary equations. Vol. I, Handb. Differ. Equ., North-Holland, Amsterdam, 2004, pp. 1–85. MR2103696
- [2] Jöran Bergh and Jörgen Löfström, *Interpolation spaces. An introduction*, Grundlehren der Mathematischen Wissenschaften, No. 223, Springer-Verlag, Berlin-New York, 1976. MR0482275
- [3] Gregory Berkolaiko and Peter Kuchment, *Introduction to quantum graphs*, Mathematical Surveys and Monographs, vol. 186, American Mathematical Society, Providence, RI, 2013. MR3013208
- [4] Stefano Bonaccorsi, Carlo Marinelli, and Giacomo Ziglio, *Stochastic FitzHugh-Nagumo equations on networks with impulsive noise*, Electron. J. Probab. **13** (2008), no. 49, 1362–1379. MR2438810
- [5] Stefano Bonaccorsi and Delio Mugnolo, *Existence of strong solutions for neuronal network dynamics driven by fractional Brownian motions*, Stoch. Dyn. **10** (2010), no. 3, 441–464. MR2671386
- [6] Stefano Bonaccorsi and Giacomo Ziglio, *A semigroup approach to stochastic dynamical boundary value problems*, Systems, control, modeling and optimization, IFIP Int. Fed. Inf. Process., vol. 202, Springer, New York, 2006, pp. 55–65. MR2241696
- [7] Robert Carlson and Vyacheslav Pivovarchik, *Ambarzumian’s theorem for trees*, Electron. J. Differential Equations (2007), No. 142, 9. MR2349970
- [8] Giuseppe Da Prato, *Kolmogorov equations for stochastic PDEs*, Advanced Courses in Mathematics. CRM Barcelona, Birkhäuser Verlag, Basel, 2004. MR2111320
- [9] Giuseppe Da Prato and Jerzy Zabczyk, *Evolution equations with white-noise boundary conditions*, Stochastics Stochastics Rep. **42** (1993), no. 3-4, 167–182. MR1291187
- [10] Giuseppe Da Prato and Jerzy Zabczyk, *Stochastic equations in infinite dimensions*, second ed., Encyclopedia of Mathematics and its Applications, vol. 152, Cambridge University Press, Cambridge, 2014. MR3236753
- [11] Klaus-Jochen Engel and Marjeta Kramar Fijavž, *Waves and diffusion on metric graphs with general vertex conditions*, Evol. Equ. Control Theory **8** (2019), no. 3, 633–661. MR3985968
- [12] Klaus-Jochen Engel and Rainer Nagel, *One-parameter semigroups for linear evolution equations*, Graduate Texts in Mathematics, vol. 194, Springer-Verlag, New York, 2000, With contributions by S. Brendle, M. Campiti, T. Hahn, G. Metafune, G. Nickel, D. Pallara, C. Perazzoli, A. Rhandi, S. Romanelli and R. Schnaubelt. MR1721989
- [13] Günther Greiner, *Perturbing the boundary conditions of a generator*, Houston J. Math. **13** (1987), no. 2, 213–229. MR904952
- [14] Markus Haase, *The functional calculus for sectorial operators*, Operator Theory: Advances and Applications, vol. 169, Birkhäuser Verlag, Basel, 2006. MR2244037
- [15] Erika Hausenblas and Jan Seidler, *A note on maximal inequality for stochastic convolutions*, Czechoslovak Math. J. **51(126)** (2001), no. 4, 785–790. MR1864042
- [16] Márton Kiss, *Spectral determinants and an Ambarzumian type theorem on graphs*, Integral Equations Operator Theory **92** (2020), no. 3, Paper No. 24, 11. MR4109188
- [17] Mihály Kovács and Eszter Sikolya, *On the stochastic Allen-Cahn equation on networks with multiplicative noise*, Electron. J. Qual. Theory Differ. Equ. (2021), Paper No. 7, 1–24. MR4204948
- [18] Mihály Kovács and Eszter Sikolya, *Stochastic reaction-diffusion equations on networks*, J. Evol. Equ. **21** (2021), no. 4, 4213–4260. MR4350573
- [19] Marjeta Kramar Fijavž and Aleksandra Puchalska, *Semigroups for dynamical processes on metric graphs*, Philos. Trans. Roy. Soc. A **378** (2020), no. 2185, 20190619, 16. MR4176394
- [20] Stig Larsson and Vidar Thomée, *Partial differential equations with numerical methods*, Texts in Applied Mathematics, vol. 45, Springer-Verlag, Berlin, 2003. MR1995838

- [21] Delio Mugnolo, *Semigroup methods for evolution equations on networks*, Understanding Complex Systems, Springer, Cham, 2014. MR3243602
- [22] Delio Mugnolo and Silvia Romanelli, *Dynamic and generalized Wentzell node conditions for network equations*, Math. Methods Appl. Sci. **30** (2007), no. 6, 681–706. MR2301840
- [23] Szymon Peszat and Jerzy Zabczyk, *Stochastic partial differential equations with Lévy noise*, Encyclopedia of Mathematics and its Applications, vol. 113, Cambridge University Press, Cambridge, 2007. MR2356959
- [24] Hans Triebel, *Interpolation theory, function spaces, differential operators*, North-Holland Mathematical Library, vol. 18, North-Holland Publishing Co., Amsterdam-New York, 1978. MR503903
- [25] J. M. A. M. van Neerven, Mark Christiaan Veraar and Lutz Weis, *Stochastic evolution equations in UMD Banach spaces*, J. Funct. Anal. **255** (2008), no. 4, 940–993. MR2433958

Acknowledgments. The authors would like to thank the anonymous referee for the careful reading of the manuscript and for the useful comments that helped them to improve the presentation and the results of the paper significantly.