

# Weyl asymptotics for Poincaré–Steklov eigenvalues in a domain with Lipschitz boundary

Downloaded from: https://research.chalmers.se, 2024-04-27 13:08 UTC

Citation for the original published paper (version of record):

Rozenblioum, G. (2023). Weyl asymptotics for Poincaré–Steklov eigenvalues in a domain with Lipschitz boundary. Journal of Spectral Theory, 13(3): 755-803. http://dx.doi.org/10.4171/JST/475

N.B. When citing this work, cite the original published paper.

research.chalmers.se offers the possibility of retrieving research publications produced at Chalmers University of Technology. It covers all kind of research output: articles, dissertations, conference papers, reports etc. since 2004. research.chalmers.se is administrated and maintained by Chalmers Library

# The spectrum of self-adjoint extensions associated with exceptional Laguerre differential expressions

Dale Frymark and Jessica Stewart Kelly

**Abstract.** Exceptional Laguerre-type differential expressions form an infinite class of Schrödinger operators having rational potentials and one limit-circle endpoint. In this manuscript, the spectrum of all self-adjoint extensions for a general exceptional Laguerre-type differential expression is given in terms of the Darboux transformations which relate the expression to the classical Laguerre differential expression. The spectrum is extracted from an explicit Weyl *m*-function, up to a sign.

The construction relies primarily on two tools: boundary triples, which parameterize the self-adjoint extensions and produce the Weyl *m*-functions, and manipulations of Maya diagrams and partitions, which classify the seed functions defining the relevant Darboux transforms. Several examples are presented.

### Contents

1.	Introduction	991
2.	Preliminaries	994
3.	Motivating example	1003
4.	Exceptional Laguerre operators	1008
5.	Manipulation of Maya diagrams	1012
6.	Initial condition normalizations	1023
7.	Weyl <i>m</i> -functions	1028
8.	Example	1035
9.	Conclusions	1039
Re	ferences	1040

## 1. Introduction

Introduced in [33, 34] in 2009, exceptional orthogonal polynomials (XOPs) describe a class of Sturm–Liouville polynomial families where some (exceptional) degrees

<sup>2020</sup> Mathematics Subject Classification. Primary 34L05; Secondary 33D45, 47E05, 47A10. *Keywords*. Exceptional orthogonal polynomials, self-adjoint Extensions, Weyl *m*-function, spectral theory, boundary triples.

of polynomials are missing. These XOPs form eigenfunctions of rational Sturm– Liouville equations, but fall outside of the restrictions of the classical Böchner theorem [4] that characterizes the classical orthogonal polynomial (COP) systems of Hermite, Laguerre and Jacobi. The properties of XOPs have been widely studied over the past 13 years [18, 20, 22, 29] as this area has brought forth interesting connections between XOP families and their COP relatives as well as surprising observations such as the fact that although a finite number of degrees are omitted from the sequence of XOP solutions, each family is complete in the associated natural Hilbert space setting.

The motivation of [18] to study these families was twofold—first, in response to a generalized Böchner problem posed in [35] and second, an interest in quantum mechanics. As such, there are numerous applications to mathematical physics. Perhaps most significantly, XOPs are, up to a gauge factor, eigenfunctions of exactly solvable potentials obtained by taking Darboux transforms of potentials, called rational extensions, from quantum mechanics [23, 30, 36]. In particular, the harmonic oscillator, isotonic oscillator and trigonometric Darboux–Pösch–Teller potential have rational extensions defined by Hermite, Laguerre and Jacobi polynomials, respectively. Many of these rational extensions are translationally shape invariant [16], and their corresponding Darboux transforms are usually called supersymmetric quantum mechanical (SUSY QM) partnerships.

Specific types of XOPs were initially introduced individually; well-known XOPs include Hermite XOPs [17], Type I-III Laguerre XOPs [26] and Type I and II Jacobi XOPs [25]. The spectral properties of the operators in each case have also been determined [2, 17, 25–27]. However, it was shown in [14] that there are an infinite number of distinct XOP classes. Although all XOP families are derived from a COP family, most of them do not fit into the finite pre-existing "type" classifications. In this manuscript, the focus will be on Laguerre-type XOP families and in particular, the associated spectral analysis.

Generally speaking, every XOP family is related to a COP family by a sequence of Darboux transformations [19, 21, 24, 37]. When a suitable Darboux transformation is applied to a COP, the result is an XOP that is an eigenfunction for an exceptional eigenvalue problem. Darboux transformations are often called "state-deleting" because when applied to an operator with a set of eigenfunctions (i.e. an operator with COP solutions) they remove one or more of the eigenfunctions from this set. With regard to the spectrum, which consists only of eigenvalues, the term "state-deleting" is misleading as the set may remain the same or even include new points after a Darboux transform is performed. Also, it is not immediately clear how the Darboux transformation impacts different self-adjoint extensions of the same symmetric differential expression; this is the effect of the Darboux transformation on the spectrum that we aim to study. For instance, we can ask whether another extension compensates for these removed eigenfunctions in some way? Perturbation theory gives two immediate restrictions. First, because the spectrum of one self-adjoint extension is discrete, the spectrum of all self-adjoint extensions must be discrete, see e.g. [38, 39]. Second, the eigenvalues of extensions must change continuously with the parameterization of the self-adjoint extension.

To study the effect of the Darboux transformation on self-adjoint expressions, we exploit the intertwining nature of the COP and XOP expressions. If the classical Laguerre expression is given in Schrödinger form, Darboux transforms can be classified via their changes to the potential thanks to the well-known Darboux–Crum formulas [8]. These changes are given by a Wronskian whose entries are eigenfunctions of the COP expression. The process of determining the effect of the Darboux transformation on the spectrum requires manipulating general Wronskians into a standard form from which information can be extracted. Fortunately, this study may utilize shape invariance where many multi-step Darboux transformations can be shown to be equivalent up to a constant. The manuscripts [6] and [16] both give equivalence relations for Wronskians, and the methods used therein can be adapted for our purposes.

To exploit spectral information from the manipulated Wronskians, we use the framework of boundary triples. This perspective is unique as compared to other spectral analyses of XOP families. The theory of boundary triples provides a natural parameterization of all self-adjoint extensions in order to produce a Weyl *m*-function. The Weyl *m*-function associated with an operator provides all of the relevant spectral data. Here we are primarily focused on only the location of eigenvalues, but is possible to extract the spectral measure and even eigenfunctions [7]. The challenge of applying the theory in this context is that solutions to the general XOP must be normalized. This normalization simplifies to evaluating the transformed solution at x = 0, which is achievable thanks to a standard form resulting from manipulation of the general Wronskians. Crucially, boundary triples for general XOP expressions are very easy to obtain, as they will carry the same structure as in the COP Laguerre expression with adjusted parameters.

#### Structure of the paper

The contents of this paper are as follows. Section 2 reviews some essential facts about the formulation of XOP families via Darboux transforms as well as the basics of boundary triples and Weyl *m*-functions as they pertain to XOP expressions. Maya diagrams are introduced as they are important to organizing information about the Wronskians associated with a Darboux transformation. Section 3 contains a relatively simple example, the Type I Laguerre XOP differential equation. Boundary triples are used to complete the spectral analysis of the Type I Laguerre XOP family of operators. This serves as a motivating example for the construction of the general methods in

later sections. Note that some of the results in this example overlap with those of [2, 26].

A number of assumptions are made throughout the text as their need arises, but for the convenience of the reader we collect them here: Assumptions 4.2, 4.5, 5.1, 5.2 and 7.2. Section 4 introduces the general notation and concepts necessary to discuss the seed functions and Wronskians that will define solutions of the XOP family. The Wronskians are defined by two Maya diagrams of seed functions:  $M_1$  and  $M_2$ . This section also sets the parameters for the underlying Hilbert space of the XOP family in terms of these Maya diagrams.

Section 5 begins with the Frobenius analysis showing that the XOP expression has the limit-circle endpoint x = 0 for similar parameter choices as the COP expression. Then we prove two central results: Theorems 5.3 and 5.5. These theorems detail the shifting of the Maya diagrams from a general position into canonical and conjugate canonical positions, tracking constants and parameters along the way. The result of this shifting process is two solutions in standard form. Section 6 takes the Wronskians, which represent solutions of the XOP family, in these standard forms, and evaluates them at x = 0. Effectively, dividing the solutions by these constants normalizes them with respect to the sesquilinear form. The resulting constants depend on many parameters, including  $M_1$ ,  $M_2$ ,  $\alpha$  and the total number of seed functions in each position. Section 7 combines the results of the previous two sections to properly augment the solutions of the XOP family in order to build a boundary triple. Construction of a general deficiency element then yields Weyl m-functions for each self-adjoint extension in the family, as shown in Corollary 7.7. A number of other observations and remarks are made in order to properly interpret the resulting expressions. Section 8 illustrates the results of the manuscript by explicitly applying them to a previously unknown XOP family. The spectral properties of this family are surprising and, although any inverse spectral theory remains elusive, encourages future investigation. Lastly, a general summary of the work and closing remarks are made in Section 9.

#### 2. Preliminaries

Generally, a XOP sequence  $\{p_n\}_{\mathbb{N}_0\setminus A}$ , where  $p_n$  is a polynomial of degree n and  $A \subseteq \mathbb{N}_0$  is a non-empty finite set, satisfies the following properties:

- (i) each  $y = p_n$  is a solution to the associated second-order eigenvalue problem;
- (ii) the associated eigenvalue problem admits no polynomial eigenfunctions having degree  $n \in A$ ;

- (iii) the sequence  $\{p_n\}_{\mathbb{N}_0\setminus A}$  is orthogonal on an open interval with respect to a positive measure w; and
- (iv) all moments  $\{\mu_n\}_{n=0}^{\infty}$  of w exist and are finite.

Each XOP expression may be written as the composition of first-order operators or Darboux transformations; these first-order operators are derived from COP expressions. We outline the process below.

Suppose  $\ell$  is a second-order expression of the form

$$\ell[y](x) = p(x)y''(x) + q(x)y'(x) + r(x)$$
(2.1)

whose associated eigenvalue equation is

$$\ell[y] = \lambda y, \tag{2.2}$$

where  $\lambda \in \mathbb{C}$ . Define rational functions

$$P(x) = \exp\left(\int \frac{x}{p(z)} \frac{q(z)}{p(z)} dz\right), \quad W(x) = \frac{P(x)}{p(x)}, \quad R(x) = r(x)W(x).$$

Multiplying (2.2) by W(x) yields the associated symmetric Sturm–Liouville form of (2.1)

$$(Py')' + Ry = \lambda Wy.$$

It follows that W(x) is the weight function associated with the expression  $\ell$ .

Two second-order expressions  $\ell$  and  $\hat{\ell}$  are *gauge-equivalent* if there exists a rational function  $\sigma$  such that

 $\hat{\ell} = \sigma \ell \sigma^{-1}.$ 

In order to derive the XOP operator from its COP counterpart, we first need to rewrite  $\ell$  as a composition of two first-order operators. A quasi-rational function  $\phi$  is a *quasi-rational eigenfunction* for the expression  $\ell$  if  $\ell[\phi] = \lambda \phi$  for  $\lambda \in \mathbb{C}$ . For example, in the case of the Laguerre COP, the quasi-rational eigenfunctions that, under certain admissibility conditions, produce XOP systems are

$$\phi_1(x) = L_m^{\alpha}(x), \tag{2.3}$$

$$\phi_2(x) = e^x L_m^{\alpha}(-x), \tag{2.4}$$

$$\phi_3(x) = x^{-\alpha} L_m^{-\alpha}(x), \tag{2.5}$$

$$\phi_4(x) = e^x x^{-\alpha} L_m^{\alpha}(-x).$$
(2.6)

The quasi-rational eigenfunction (2.5) produces the Type I Laguerre XOP family. The reader is directed to Section 8 for more details into the decomposition of the COP Laguerre expression and formulation of the Type I expression.

**Proposition 2.1** ([14, Proposition 3.5]). For a second-order differential operator  $\ell[y]$  having rational coefficients, let  $\phi$  be a quasi-rational eigenfunction of  $\ell$  with eigenvalue  $\lambda$ , and let b(x) be an arbitrary, non-zero rational function. Define rational functions

$$w = \frac{\phi'}{\phi}, \quad \hat{b} = \frac{p}{b}, \quad \hat{w} = -w - \frac{q}{p} + \frac{\hat{b}}{b}$$

and first-order operators A and B by

$$A[y] = b(y' - wy) \quad and \quad B[y] = \hat{b}(y' - \hat{w}y).$$
(2.7)

Then the operators A and B of (2.7) are used to form a rational factorization

$$\ell = BA + \lambda$$

of  $\ell$ .

Given a rational factorization, the *partner operator*  $\hat{\ell}$  is defined via

$$\hat{\ell} = AB + \lambda$$

The operator  $\hat{\ell}$  will be a second–order operator of the form

$$\hat{\ell}[y](x) = p(x)y''(x) + Q(x)y' + \mathcal{R}(x)y(x)$$
(2.8)

where Q and  $\mathcal{R}$  are rational functions defined by

$$\begin{split} &\mathcal{Q} = q + p' - \frac{2pb'}{b}, \\ &\mathcal{R} = -p(\hat{w}' + \hat{w}^2) - \mathcal{Q}\hat{w} \\ &= r + q' + wp' - \frac{b'}{b}(q + p') + \left(2\left(\frac{b'}{b}\right)^2 - \frac{b''}{b} + 2w'\right)p, \\ &\mathcal{W} = \frac{pW}{b^2} = \frac{P}{b^2}. \end{split}$$

When *b* and  $\phi$  are suitably chosen and  $\ell$  is a COP expression, the partner operator  $\hat{\ell}$  produces an XOP operator having weight function *W*. The reader is directed to [14, Proposition 3.6] for the suitable conditions on *b*. The polynomial part of *b* is important for future calculations and is a generalized Laguerre polynomial in the context of Laguerre XOP families, see (4.9).

The transformation  $\mathcal{A} := \ell \to \hat{\ell}$  is a (rational) *Darboux transformation*. In fact, with appropriate assumptions, this process may be iterated  $n \in \mathbb{N}$  times to produce a factorization chain or a Darboux transformation of size *n*. These complex Darboux

transformations are used to generate XOP systems. We emphasize that factorizations of a COP are not unique; and herein provides the opportunity for many XOP families.

One crucial property that arises from this construction is the following.

**Lemma 2.2** (Intertwining property [14, Proposition 3.2]). If  $\ell$  and  $\hat{\ell}$  are related via a rational Darboux transformation, then the following intertwining relations hold:

$$A\ell = \hat{\ell}A, \quad \ell B = B\hat{\ell}.$$

In particular, if  $y_{\lambda}$  is an eigenfunction of  $\ell[y] = \lambda y$ , then  $\tilde{y}_{\lambda} := A[y_{\lambda}]$  satisfies  $\hat{\ell}[\tilde{y}_{\lambda}] = \lambda \tilde{y}_{\lambda}$ .

To simplify our discussion, we will refer to all self-adjoint extensions of a symmetric operator as a "family." XOP and COP families thus refer to families that contain one extension which possesses a set of XOPs or COPs. The notation  $\ell$  and  $\hat{\ell}$  will often be used to denote the classical and exceptional symmetric differential expressions, respectively, that generate families. The general Laguerre COP expression is dependent upon a parameter  $\alpha$  and it follows that both  $\ell$  and  $\hat{\ell}$  also have dependency upon this parameter. When emphasis on the parameter is needed, we will use  $\ell^{\alpha}$  and  $\hat{\ell}^{\alpha}$  for  $\ell$  and  $\hat{\ell}$ , but for ease of notation, the parameter will often be suppressed. Additionally, when relevant, a superscript of I, II or III may be included to indicate the XOP Laugerre type.

Of course, when the differential expressions  $\ell$  or  $\hat{\ell}$  are associated with a specific domain they are operators—and whether they are self-adjoint or not depends on this domain. Two important symmetric (and not self-adjoint) operators associated with a general differential expression are the maximal and minimal operators. The following setup is centered around the XOP expression  $\hat{\ell}$ , as this is the most common setting we will use, but analogous definitions for  $\ell$  hold.

**Definition 2.3** ([31, Section 17.2]). The *maximal domain* of  $\hat{\ell}[\cdot]$  is given by

$$\mathcal{D}_{\max} = \mathcal{D}_{\max}(\hat{\ell}) := \{ f : (0, \infty) \to \mathbb{C} \mid f, Pf' \in AC_{loc}(0, \infty); \\ f, \hat{\ell}[f] \in L^2[(0, \infty), W] \}$$

The associated maximal operator

$$\mathcal{L}_{\max}: \mathcal{D}_{\max} \to L^2[(0,\infty); \mathcal{W}]$$

is defined to be

$$\mathcal{L}_{\max}[f] = \hat{\ell}[f], \quad f \in \mathcal{D}_{\max}$$

The designation of "maximal" is appropriate in this case because  $\mathcal{D}_{\text{max}}$  is the largest possible subspace that  $\hat{\ell}$  maps back into  $L^2[(0,\infty), \mathcal{W}]$ .

The sesquilinear form associated to the expression  $\hat{\ell}$  can be explicitly calculated by integration by parts for  $f, g \in \mathcal{D}_{max}$  as

$$[f,g]_{\hat{\ell}}(x) = P(x)[f(x)g'(x) - f'(x)g(x)].$$
(2.9)

**Theorem 2.4** ([31, Section 17.2]). The two limits  $[f, g]_{\hat{\ell}}(\infty) := \lim_{x \to \infty^-} [f, g]_{\hat{\ell}}(x)$ and  $[f, g]_{\hat{\ell}}(0) := \lim_{x \to 0^+} [f, g]_{\hat{\ell}}(x)$  exist and are finite for  $f, g \in \mathcal{D}_{max}$ .

**Definition 2.5** ([31, Section 17.2]). The *minimal domain* of  $\hat{\ell}[\cdot]$  is given by

$$\mathcal{D}_{\min} = \mathcal{D}_{\min}(\hat{\ell}) := \{ f \in \mathcal{D}_{\max} : [f, g]_{\hat{\ell}} |_{0}^{\infty} = 0 \ \forall g \in \mathcal{D}_{\max} \}.$$

The associated minimal operator

$$\mathcal{L}_{\min}: \mathcal{D}_{\min} \to L^2[(0,\infty); \mathcal{W}]$$

is defined to be

$$\mathcal{L}_{\min}[f] = \hat{\ell}[f], \quad f \in \mathcal{D}_{\min}.$$

The maximal and minimal operators  $\mathcal{L}_{max}$  and  $\mathcal{L}_{min}$  are defined as the symmetric operators acting via  $\hat{\ell}$  on the domains  $\mathcal{D}_{max}(\hat{\ell})$  and  $\mathcal{D}_{min}(\hat{\ell})$ , respectively. By [31, Section 17.2], these operators are adjoints of one another, i.e.  $(\mathcal{L}_{min})^* = \mathcal{L}_{max}$  and  $(\mathcal{L}_{max})^* = \mathcal{L}_{min}$ . Self-adjoint operators associated with  $\hat{\ell}$  are defined via the theory of boundary triples in the next subsection.

#### 2.1. Boundary triples

The principal tools for extracting spectral information from self-adjoint extensions of symmetric differential operators in this manuscript are the Weyl *m*-functions that are generated by boundary triples and the parameterization of all self-adjoint extensions. For completeness, some basics from the theory of boundary triples are included here. The content of this subsection is also found in [3, 7], which the interested reader may consult for more details. For the purposes of this paper, we restrict ourselves to the simplified case of Sturm–Liouville differential operators rather than more general linear relations. Additionally, we further narrow the scope of discussion as the XOP Laguerre differential expressions will have deficiency indices (1, 1), see Section 4.

Let  $\mathscr{L}_{\text{max}}$  and  $\mathscr{L}_{\text{min}}$  denote the symmetric (and not self-adjoint) XOP Laguerre operators defined by the expression  $\hat{\ell}$  acting on the domains given in Definitions 2.3 and 2.5, as above, with  $\mathscr{H} = L^2[(0,\infty), \mathscr{W}(x)dx]$ , with  $\mathscr{W}(x) > 0$  a.e. in  $(0,\infty)$  and  $\mathscr{W}(x) \in L^1_{\text{loc}}[(0,\infty), dx]$ .

**Definition 2.6** ([3, Definition 2.1.1]). A *boundary triple* for  $\mathscr{L}_{max}$ , denoted { $\mathbb{C}$ ,  $\Gamma_0$ ,  $\Gamma_1$ }, is composed of the boundary space  $\mathbb{C}$  and two surjective linear maps

$$\Gamma_0, \Gamma_1: \mathcal{D}_{max} \to \mathbb{C}$$

for which the Green's identity

$$\langle \mathcal{L}_{\max}f,g \rangle_{\mathcal{H}} - \langle f,\mathcal{L}_{\max}g \rangle_{\mathcal{H}} = \langle \Gamma_1f,\Gamma_0g \rangle_{\mathbb{C}} - \langle \Gamma_0f,\Gamma_1g \rangle_{\mathbb{C}},$$
(2.10)

holds for all  $f, g \in \mathcal{D}_{max}$ .

Notice that the left-hand side of (2.10) is simply  $[\cdot, \cdot]_{\hat{\ell}}$ .

**Definition 2.7** ([3, Definition 1.4.9]). Let  $\zeta \in \mathbb{C}$ . The space

$$\mathfrak{N}_{\boldsymbol{\zeta}}(\boldsymbol{\mathcal{L}}_{\max}) := \ker(\boldsymbol{\mathcal{L}}_{\max} - \boldsymbol{\zeta}),$$

is called the *defect subspace* of  $\mathcal{L}_{\min}$  at the point  $\zeta \in \mathbb{C}$ .

Elements of these defect subspaces are referred to as deficiency elements. The usual positive and negative defect spaces are then simply  $\mathfrak{N}_i(\mathcal{L}_{\max})$  and  $\mathfrak{N}_{-i}(\mathcal{L}_{\max})$ , respectively. It should be clear that boundary triples are not typically unique. Indeed, given a self-adjoint extension H of  $\mathcal{L}_{\min}$ ,  $\mathcal{L}_{\max}$  can be decomposed into a direct sum of dom(H) and a defect space. This allows maps  $\Gamma_0$  and  $\Gamma_1$  to be defined so that dom(H) = ker( $\Gamma_0$ ) and { $\mathbb{C}$ ,  $\Gamma_0$ ,  $\Gamma_1$ } form a boundary triple, see [3, Theorem 2.4.1].

In our context, the boundary triples for  $\mathcal{L}_{max}$  will be formed via quasi-derivatives.

**Definition 2.8.** Let u and v be linearly independent real solutions of the equation  $(\mathcal{L}_{\max} - \zeta_0)y = 0$  for some  $\zeta_0 \in \mathbb{R}$  and assume that the solutions are normalized by  $[u, v]_{\hat{\ell}} = 1$ . For  $f \in \mathcal{D}_{\max}$ , the quasi-derivatives of f are induced by the normalized solutions u, v and defined as complex functions on  $(0, \infty)$  given by

$$f^{[0]} := [f, v]_{\hat{\ell}}$$
 and  $f^{[1]} := -[f, u]_{\hat{\ell}}$ .

Note, that in practice, the minus sign outside the sesquilinear form in the second quasi-derivative is absorbed into the definition of u. These quasi-derivatives naturally define two maps and self-adjoint extensions for exceptional Laguerre operators:

$$\Gamma_0 f := f^{[0]}(0), \quad \mathcal{D}_{\infty} := \{ f \in \mathcal{D}_{\max} : f \in \ker(\Gamma_0) \},$$
(2.11a)

$$\Gamma_1 f := f^{[1]}(0), \quad \mathcal{D}_0 := \{ f \in \mathcal{D}_{\max} : f \in \ker(\Gamma_1) \}.$$
 (2.11b)

Recall that for all  $f \in \mathcal{D}_{\text{max}}$  the quasi-derivatives  $f^{[0]}(0)$  and  $f^{[1]}(0)$  are well defined due to Theorem 2.4. All self-adjoint extensions of  $\mathcal{L}_{\min}$  are in one-to-one correspondence with

$$\mathcal{D}_{\tau} := \{ f \in \mathcal{D}_{\max} : \tau \Gamma_0 f = \Gamma_1 f \},$$
(2.12)

where  $\tau \in \mathbb{R} \cup \{\infty\}$ . The case  $\tau = \infty$  is interpreted as representing the operator  $\mathcal{L}_{\infty}$ ; this case technically represents the linear relation  $\{0, \mathbb{R}\}$ . The only other onedimensional linear relation that is not an operator,  $\{\mathbb{R}, 0\}$ , clearly corresponds to  $\mathcal{L}_0$ . The definitions can be made rigorous by appealing to the corresponding semi-bounded forms of the operators; interested readers should consult [7, Remark 3.7]. The above definitions are adequate for our purposes.

Also note that the extension  $\mathcal{L}_0$  is identified as the Friedrichs extension when u is the principal solution by [28]: the extension whose semi-bounded form has the greatest lower bound. On the other hand, if  $\mathcal{L}_{max}$  is positive and v is a non-principal solution,  $\mathcal{L}_{\infty}$  is the Krein–von Neumann extension, see [3, Definition 5.4.2]. This structure immediately allows for the definition of a Weyl *m*-function.

**Definition 2.9** ([3, Definition 2.3.1, 2.3.4]). Let { $\mathbb{C}$ ,  $\Gamma_0$ ,  $\Gamma_1$ } be a boundary triple for  $\mathcal{L}_{max}$  and  $\lambda \in \mathbb{C}$ . Then

$$\rho(\mathscr{L}_{\infty}) \ni \lambda \mapsto M_{\infty}(\lambda) = \Gamma_1(\Gamma_0 \upharpoonright \mathfrak{N}_{\lambda}(\mathscr{L}_{\max}))^{-1},$$

where  $\uparrow$  denotes the restriction, is called the *Weyl m-function* associated with the boundary triple { $\mathbb{C}, \Gamma_0, \Gamma_1$ }.

In this context, the spectrum of  $\mathcal{L}_{\infty}$  is discrete and the difference of the resolvents of  $\mathcal{L}_{\infty}$  and  $\mathcal{L}_{\tau}$  is an operator of rank one. Thus, the spectrum of the self-adjoint operator  $\mathcal{L}_{\tau}$  is also discrete. The following proposition can be found in various forms in [9, Proposition 1], [13, Proposition 3.5], or [3, Theorem 2.6.5], but in our context has a slightly altered notation.

**Proposition 2.10.** Let  $\{\mathbb{C}, \Gamma_0, \Gamma_1\}$  be the boundary triple for  $\mathcal{L}_{max}$ ,  $\tau \in \mathbb{R} \cup \{\infty\}$  and  $\lambda \in \rho(\mathcal{L}_{\infty})$ . Then

$$\lambda \in \sigma_p(\mathcal{L}_{\tau}) \iff 0 \in \sigma_p(\tau - M_{\infty}(\lambda)).$$

*Moreover, we have* dim ker $(\mathcal{L}_{\tau} - \lambda) = \dim \ker(\tau - M_{\infty}(\lambda)).$ 

For  $\lambda \in \rho(\mathcal{L}_0) \cap \rho(\mathcal{L}_\infty)$ , the spectral properties of  $\mathcal{L}_\tau$  can thus be described with the help of the function

$$M_{\tau}(\lambda) = (\tau - M_{\infty}(\lambda))^{-1}, \qquad (2.13)$$

see [3, (3.8.7)].

The relationship  $M_0 = -M_{\infty}^{-1}$  is immediately apparent. Note that  $M_0(\lambda)$  can be obtained without exploiting (2.13) by simply switching the definitions of the maps  $\Gamma_0$  and  $\Gamma_1$ . A similar construction that the reader may find useful for more general Sturm–Liouville operators is available in [7, 12].

Now, fix a fundamental system  $(u_1(\cdot, \lambda); u_2(\cdot, \lambda))$  for the equation

$$(\mathcal{L}_{\max} - \lambda)f = 0$$

by the initial conditions

$$\begin{pmatrix} u_1^{[0]}(0,\lambda) & u_2^{[0]}(0,\lambda) \\ u_1^{[1]}(0,\lambda) & u_2^{[1]}(0,\lambda) \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$
 (2.14)

Linear combinations of these solutions then define general elements of  $\mathfrak{N}_{\lambda}(\mathcal{L}_{max})$  in a standard way.

**Proposition 2.11** ([3, Proposition 6.4.9]). Let  $\Gamma_0$  and  $\Gamma_1$  be defined as in (2.11). Then  $\{\mathbb{C}, \Gamma_0, \Gamma_1\}$  is a boundary triple for  $\mathcal{L}_{max}$ . Moreover, if  $\lambda \in \mathbb{C} \setminus \mathbb{R}$  and  $\chi(x, \lambda)$  is a nontrivial element in  $\mathfrak{N}_{\lambda}(\mathcal{L}_{max})$ , then  $\chi^{[0]}(0, \lambda) \neq 0$  and the Weyl *m*-function is given by

$$M_{\infty}(\lambda) = \frac{\chi^{[1]}(0,\lambda)}{\chi^{[0]}(0,\lambda)}.$$

It is then possible to transform the boundary triple so that  $M_{\tau}(\lambda)$  naturally emerges from Proposition 2.11 by using [3, (6.4.8)]:

$$M_{\tau}(\lambda) = \frac{1 + \tau M_{\infty}(\lambda)}{\tau - M_{\infty}(\lambda)}, \quad \lambda \in \mathbb{C} \setminus \mathbb{R}.$$
 (2.15)

This means that a corresponding Weyl m-function can be accessed for each selfadjoint extension in the XOP family, and a spectral analysis can be completed by finding the poles of these m-functions.

#### 2.2. Maya diagrams

Maya diagrams will be used to indicate the indices of seed functions within the Wronskians that define Darboux transformations. These Maya diagrams are vitally important to the construction of XOPs and will be manipulated throughout the manuscript. We include a brief overview here that mostly follows [6]. More information on Maya diagrams, their history, Young diagrams, and partitions relating to XOP families may be found in [6, 16, 32].

A Maya diagram M is an infinite subset of integers that contains finitely many non-negative integers and excludes finitely many negative integers. Visually, imagine an infinite row of boxes with a fixed origin. These boxes are either filled (the integer belongs to M) or empty (the integer does not belong to M). Thus, M is characterized by two finite sequences of integers: the positive integers included in M and the negative integers excluded by M. Each of the non-negative integers  $n_i$  in M are labeled to create a decreasing sequence by

$$n_1 > n_2 > \cdots > n_{r_1} \ge 0$$
,

where  $r_1$  is the number of filled boxes to the right of the origin. If no boxes are filled to the right of the origin, the sequence is empty and  $r_1 = 0$ . Likewise, the number of empty boxes to the left of the origin can be denoted as  $r_4$ , the negative integers k corresponding to these boxes assigned values via n' = -k - 1 and

$$n'_1 > n'_2 > \dots > n'_{r_4} \ge 0.$$

The Maya diagram can then be succinctly described by these two sequences as

$$M = (n'_1, n'_2, \dots, n'_{r_4} \mid n_1, n_2, \dots, n_{r_1}).$$

This notation comes from the Frobenius symbol of an integer partition and agrees with the classical notation when  $r_1 = r_4$ .

As an example, let *M* be the Maya diagram M = (5, 2, 1 | 4, 3, 1), whose graphical representation may be seen in Figure 1.



Figure 1. Maya diagram.

All boxes to the left of those shown in Figure 1 are filled, while those to the right are empty. The origin of a Maya diagram may be shifted to the left or right by adding or subtracting, respectively. The Maya diagram  $\tilde{M} = M + 3$  is represented by Figure 2.



Figure 2. Shifted Maya diagram.

In particular, the length of the two subsequences,  $r_1 + r_4$ , is not necessarily stable under such translations: M has 6 indices and  $\tilde{M}$  has 5.

A Maya diagram is in *canonical form* if there are no empty boxes to the left of the origin, but the first box to the right is empty. This form is useful for comparing diagrams and can always be obtained in the current context by applying a shift. If M + t puts a Maya diagram M into canonical form for some  $t \in \mathbb{Z}$ , we denote this shift as t = t(M). In the context of the above example,  $t(M) = 6 = n'_1 + 1$ .

Alternatively, the *conjugate canonical form* of a Maya diagram is the position when there are no filled boxes to the right of the origin, but the first box to the left is filled. The required shift to put M into conjugate canonical form is denoted by t' = t'(M). In the above example,  $t'(M) = -(n_1 + 1) = -5$  and we have the associated representation in Figure 3.



Figure 3. Conjugate canonical form Maya diagram.

One of the goals of Section 5 will be to manipulate the Maya diagrams  $M_1$  and  $M_2$  of (4.5) and (4.6) so that they are either in canonical or conjugate canonical position and can therefore each be described by a single finite decreasing sequence of non-negative integers. Naturally, such sequences describe a partition, see (4.12) and (4.13). Hence, any given Maya diagram gives rise to a partition, when shifted into canonical form, and a conjugate partition, when shifted into conjugate canonical form.

#### 3. Motivating example

We begin with an example: the Type I Laguerre exceptional family. The spectral properties of the extension  $\mathcal{L}_0$  are already known from [26], where explicit knowledge of the XOP system and the respective polynomials were used. However, the authors find it instructive to present the construction of the boundary triple and *m*-functions in this specific context before moving on to the general setting.

The classical Laguerre differential expression  $\ell^{\alpha}$  acts via

$$\ell^{\alpha}[y](x) := -xy''(x) + (x - \alpha - 1)y'(x).$$
(3.1)

For fixed  $m \in \mathbb{N}$ ,  $\ell^{\alpha}[y](x)$  has the rational factorization

$$-\ell^{\alpha} = B_m^{I,\alpha} \circ A_m^{I,\alpha} + \alpha + m + 1, \qquad (3.2)$$

where

$$A_m^{I,\alpha}[y] := L_m^{\alpha}(-x)y'(x) - L_m^{\alpha+1}(-x)y(x),$$
(3.3)

$$B_m^{I,\alpha}[y] := \frac{xy'(x) + (1+\alpha)y(x)}{L_m^{\alpha}(-x)},$$
(3.4)

where  $L_m^{\alpha}$  is the classical Laguerre polynomial of degree *m*. The rational factorization in (3.2) is used to define the differential expression  $\hat{\ell}_m^{I,\alpha}$  for the Type I Laguerre XOP family

$$\begin{split} \hat{\ell}_{m}^{I,\alpha}[y](x) &= -(A_{m}^{I,\alpha-1} \circ B_{m}^{I,\alpha-1}[y](x) + \alpha + m) \\ &= -xy''(x) + \left(x - \alpha - 1 + 2x \frac{(L_{m}^{\alpha-1}(-x))'}{L_{m}^{\alpha-1}(-x)}\right)y'(x) \\ &+ \left(\frac{2\alpha(L_{m}^{\alpha-1}(-x))'}{L_{m}^{\alpha-1}(-x)} - m\right)y(x). \end{split}$$

Observe that, as opposed to the classical Laguerre differential expression  $\ell^{\alpha}$ , the exceptional expression  $\hat{\ell}_m^{I,\alpha}$  has rational coefficients. As a result, in the exceptional case,  $\hat{\ell}_m^{I,\alpha}[y] = \lambda y$  contains no polynomial solutions of degrees  $0, 1, \ldots, m-1$ . The weight function associated with the XOP family is

$$W_m^{I,\alpha} := \frac{x^{\alpha} e^{-x}}{[L_m^{\alpha-1}(-x)]^2}.$$

In the Type I Laguerre XOP case,  $\alpha - 1 > -1$  to ensure that  $L_m^{\alpha-1}(-x)$  has no roots in  $[0, \infty)$  and all moments of  $\mathcal{W}_m^{I,\alpha}$  exist and are finite. For the remainder of this motivating example,  $\alpha > 0$ . The associated maximal operator  $\mathcal{L}_{\text{max}}$  acts via  $\hat{\ell}_m^{I,\alpha}$  on the maximal domain

$$\mathcal{D}_{\max} := \{ f : (0, \infty) \to \mathbb{C} \mid f, x^{\alpha+1} e^{-x} f' \in AC_{\text{loc}}(0, \infty); \\ f, \hat{\ell}_m^{I,\alpha}[f] \in L^2[(0, \infty), W_m^{I,\alpha}] \}.$$

For  $f, g \in \mathcal{D}_{\text{max}}$  and  $x \in (0, \infty)$ , the sesquilinear form is given by

$$[f,g]_{\hat{\ell}}(x) := \frac{x^{\alpha+1}e^{-x}}{[L_m^{\alpha-1}(-x)]^2} [f(x)g'(x) - f'(x)g(x)].$$

Frobenius analysis shows the endpoint x = 0 is limit-circle for  $\hat{\ell}_m^{I,\alpha}$  when  $\alpha \in (0, 1)$ ; and the endpoint  $x = \infty$  is limit-point for  $\alpha > 0$  [26, Theorem 3.3]. Observe that the functions

$$\tilde{y}_1 := -\frac{[L_m^{\alpha-1}(0)]^2}{\alpha}$$
 and  $\tilde{y}_2 := x^{-\alpha}$ ,

are both in  $L^2[(0,\infty), W_m^{I,\alpha}]$  and that  $\tilde{y}_2$  is a particular solution to  $\hat{\ell}_m^{I,\alpha}[y] = 0$ .

**Remark 3.1.** The other particular solution to  $\hat{\ell}_m^{I,\alpha}[y] = 0$  is actually  $L_m^{\alpha-1}(x)$ , not  $\tilde{y}_1$ . However,  $\tilde{y}_1$  does capture the correct asymptotic behavior of the solution near 0. Indeed, when defining maps using quasi-derivatives with solutions that are written as power series, it is easy to see that only the lowest-order term plays any role, see, i.e., [12, Section 3.1]. The use of  $\tilde{y}_1$ , as opposed to  $L_m^{\alpha-1}(x)$ , therefore makes all calculations much simpler.

Together, these two functions define maps using the quasi-derivatives:

$$\Gamma_0 f := f^{[0]}(0) = [f, \tilde{y}_2]_{\hat{\ell}}(0) = \lim_{x \to 0^+} -\frac{\alpha f(x) + x f'(x)}{[L_m^{\alpha-1}(0)]^2}$$

and

$$\Gamma_1 f := f^{[1]}(0) = [f, \tilde{y}_1]_{\hat{\ell}}(0) = \lim_{x \to 0^+} \frac{x^{\alpha+1}}{\alpha} f'(x).$$

Using results of Subsection 2.1, it is easy to show that  $\{\mathbb{C}, \Gamma_0, \Gamma_1\}$  is a boundary triple for  $\mathcal{L}_{max}$ . Note that the choice of  $\tilde{y}_1$  immediately yields  $[\tilde{y}_1, \tilde{y}_2]_{\hat{\ell}}(0) = 1$ . The boundary triple naturally defines two self-adjoint extensions  $\mathcal{L}_0$  and  $\mathcal{L}_\infty$  (suppressing the dependence on *m* and  $\alpha$ ) that act via  $\hat{\ell}_m^{I,\alpha}$  on the domains

$$dom(\mathcal{L}_0) = \{ f \in \mathcal{D}_{max} : \Gamma_1(f) = 0 \},\ dom(\mathcal{L}_\infty) = \{ f \in \mathcal{D}_{max} : \Gamma_0(f) = 0 \},\$$

respectively.

The two linearly independent solutions of the Laguerre differential equation  $\ell^{\alpha-1}$  are given by confluent hypergeometric functions:

$$M(-\lambda, \alpha, x) = {}_{1}F_{1}(-\lambda, \alpha, x) \quad \text{(for } \alpha \notin -\mathbb{N}_{0}\text{)}, \tag{3.5}$$

$$x^{-\alpha+1}M(1-\lambda-\alpha,2-\alpha,x) \quad \text{(for } \alpha \notin \mathbb{N} > 1\text{)}. \tag{3.6}$$

As a consequence of Lemma 2.2, the Type I Laguerre XOP expression is therefore intertwined with the classical Laguerre expression  $\ell^{\alpha-1}$ . Therefore, a fundamental system for the equation  $(\mathcal{L}_{max} - \lambda)f = 0$  can be found by plugging the solutions

of (3.5) and (3.6) into the Darboux transformation  $-A^{I,\alpha-1}$  of (3.3). Thus

$$\Omega_{1,m}^{\alpha}(x,\lambda) := -A^{I,\alpha-1}[M(-\lambda,\alpha,x)]$$
  
=  $\frac{\Gamma(m+\alpha)}{\Gamma(m+1)\Gamma(\alpha)} \Big[ \frac{\alpha+m}{\alpha} M(-m,\alpha+1,-x)M(-\lambda,\alpha,x) + \frac{\lambda}{\alpha} M(-m,\alpha,-x)M(-\lambda+1,\alpha+1,x) \Big],$ 

and

$$\Omega_{2,m}^{\alpha}(x,\lambda) := -A^{I,\alpha-1}[x^{-\alpha+1}M(1-\lambda-\alpha,2-\alpha,x)]$$
  
=  $\frac{\Gamma(m+\alpha+1)}{\Gamma(m+1)\Gamma(\alpha+1)}x^{-\alpha}$   
 $\times \left[\frac{\alpha(1-\alpha)}{m+\alpha}M(-m,\alpha,-x)M(1-\lambda-\alpha,1-\alpha,x)\right]$   
 $+ xM(-m,\alpha+1,-x)M(1-\lambda-\alpha,2-\alpha,x)\right]$ 

are two linearly independent solutions of  $(\mathcal{L}_{\max} - \lambda) f = 0$ . Furthermore, a straightforward calculation shows

$$\begin{pmatrix} (\Omega_{1,m}^{\alpha})^{[0]}(0,\lambda) & (\Omega_{2,m}^{\alpha})^{[0]}(0,\lambda) \\ (\Omega_{1,m}^{\alpha})^{[1]}(0,\lambda) & (\Omega_{2,m}^{\alpha})^{[1]}(0,\lambda) \end{pmatrix} = \begin{pmatrix} 1/C_m^{\alpha} & 0 \\ 0 & 1/D_m^{\alpha} \end{pmatrix},$$
(3.7)

where

$$C_m^{\alpha}(\lambda) := -\frac{\Gamma(m+\alpha)}{(\lambda+m+\alpha)\Gamma(\alpha)\Gamma(m+1)} \quad \text{and} \quad D_m^{\alpha}(\lambda) := -\frac{\Gamma(m+1)\Gamma(\alpha)}{(1-\alpha)\Gamma(m+\alpha)}.$$

The fundamental system composed of  $u_1(x, \lambda)$  and  $u_2(x, \lambda)$  must satisfy both  $(\mathcal{L}_{\max} - \lambda) f = 0$  and the initial conditions

$$\begin{pmatrix} u_1^{[0]}(0,\lambda) & u_2^{[0]}(0,\lambda) \\ u_1^{[1]}(0,\lambda) & u_2^{[1]}(0,\lambda) \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

If we suppress dependence on  $\alpha$  and m, this can be accomplished by setting

$$u_1(x,\lambda) := C_m^{\alpha}(\lambda)\Omega_{1,m}^{\alpha}(x,\lambda),$$
  
$$u_2(x,\lambda) := D_m^{\alpha}(\lambda)\Omega_{2,m}^{\alpha}(x,\lambda).$$

It remains only to compute the explicit Weyl *m*-function and extract spectral information for each self-adjoint extension. A general deficiency element associated with the subspace  $\mathfrak{N}_{\lambda}(\ell^{\alpha-1})$ , see Definition 2.7, satisfying  $(\ell^{\alpha-1} - \lambda)f = 0$  and denoted by  $\chi(x, \lambda)$  may be written via the Tricomi confluent hypergeometric function [5, (13.2.42)] as

$$U(-\lambda, \alpha, x) = \frac{\Gamma(1-\alpha)}{\Gamma(1-\alpha-\lambda)} M(-\lambda, \alpha, x) + \frac{\Gamma(\alpha-1)}{\Gamma(-\lambda)} x^{-\alpha+1} M(1-\lambda-\alpha, 2-\alpha, x).$$

The intertwining property in Lemma 2.2 implies that a general deficiency element  $\chi_{\hat{\ell}}$  of  $(\mathcal{L}_{\text{max}} - \lambda)f = 0$  can be written as

$$\begin{split} \chi_{\hat{\ell}}(x,\lambda) &= -A^{I,\alpha-1}[U(-\lambda,\alpha,x)] \\ &= \frac{\Gamma(1-\alpha)}{\Gamma(1-\alpha-\lambda)}(-A^{I,\alpha-1}[M(-\lambda,\alpha,x)]) \\ &+ \frac{\Gamma(\alpha-1)}{\Gamma(-\lambda)}(-A^{I,\alpha-1}[x^{-\alpha+1}M(1-\lambda-\alpha,2-\alpha,x)]) \\ &= \frac{\Gamma(1-\alpha)}{\Gamma(1-\alpha-\lambda)}\Omega_{1,m}^{\alpha}(x,\lambda) + \frac{\Gamma(\alpha-1)}{\Gamma(-\lambda)}\Omega_{2,m}^{\alpha}(x,\lambda) \\ &= \frac{\Gamma(1-\alpha)}{C_{m}^{\alpha}(\lambda)\Gamma(1-\alpha-\lambda)}u_{1}(x,\lambda) + \frac{\Gamma(\alpha-1)}{D_{m}^{\alpha}\Gamma(-\lambda)}u_{2}(x,\lambda). \end{split}$$

Hence, the initial conditions mean

r - 1

$$\chi_{\hat{\ell}}^{[0]}(0,\lambda) = \frac{\Gamma(1-\alpha)}{C_m^{\alpha}(\lambda)\Gamma(1-\alpha-\lambda)} \quad \text{and} \quad \chi_{\hat{\ell}}^{[1]}(0,\lambda) = \frac{\Gamma(\alpha-1)}{D_m^{\alpha}\Gamma(-\lambda)}.$$

Proposition 2.11 implies that if  $\lambda \in \rho(\mathcal{L}_{\infty})$  the Weyl *m*-function for the extension  $\mathcal{L}_{\infty}$  is given by

$$M_{\infty}(\lambda) = \frac{\chi_{\hat{\ell}}^{[1]}(0,\lambda)}{\chi_{\hat{\ell}}^{[0]}(0,\lambda)} = \left(\frac{\Gamma(m+\alpha)}{\Gamma(m+1)\Gamma(\alpha)}\right)^2 \frac{(1-\alpha)\Gamma(\alpha-1)\Gamma(1-\alpha-\lambda)}{(\lambda+m+\alpha)\Gamma(-\lambda)\Gamma(1-\alpha)}$$
$$= -[L_m^{\alpha-1}(0)]^2 \frac{\Gamma(\alpha)\Gamma(1-\alpha-\lambda)}{(\lambda+m+\alpha)\Gamma(-\lambda)\Gamma(1-\alpha)}.$$

The spectrum of the self-adjoint operator  $\mathscr{L}_{\infty}$  are those points which are poles of  $M_{\infty}(\lambda)$ . Recall that the Gamma function has no zeros, but does have simple poles at zero and the negative integers. Therefore,  $M_{\infty}$  has poles at  $\sigma(\mathscr{L}_{\infty}) = (-m - \alpha) \cup \{n + 1 - \alpha\}_{n \in \mathbb{N}_0}$  and  $\sigma(\mathscr{L}_{\infty})$  is the spectrum of the self-adjoint operator  $\mathscr{L}_{\infty}$ . The corresponding self-adjoint extension of the COP expression has the same spectrum with the point  $-m - \alpha$  replaced by  $-\alpha$ .

Likewise, the *m*-function of  $\mathcal{L}_0$ , denoted  $M_0$ , can be found by computing the value of  $-1/M_{\infty}(\lambda)$  for  $\lambda \in \rho(\mathcal{L}_{\infty}) \cup \rho(\mathcal{L}_0)$ . This extension contains the Type I Laguerre XOP polynomials and is easily seen to have eigenvalues  $\lambda = \{n\}_{n \in \mathbb{N}_0}$ .

The *m*-function for any self-adjoint extension can also be written down in a standard way. Let  $\tau \in \mathbb{R} \cup \{\infty\}$  and  $\mathcal{L}_{\tau}$  refer to the self-adjoint operator acting via  $\hat{\ell}_{m}^{I,\alpha}$  on

$$\operatorname{dom}(\mathcal{L}_{\tau}) := \{ f \in \mathcal{D}_{\max} : f \in \ker(\Gamma_1 - \tau \Gamma_0) \} = \{ f \in \mathcal{D}_{\max} | f^{[1]}(0) = \tau f^{[0]}(0) \}.$$

The corresponding Weyl *m*-function for  $\lambda \in \rho(\mathcal{L}_{\infty}) \cup \rho(\mathcal{L}_{\tau})$ , as in (2.15), is

$$M_{\tau}(\lambda) = \frac{1 + \tau M_{\infty}(\lambda)}{\tau - M_{\infty}(\lambda)}$$
  
=  $\frac{(\lambda + m + \alpha)\Gamma(-\lambda)\Gamma(1 - \alpha) - \tau [L_m^{\alpha - 1}(0)]^2\Gamma(\alpha)\Gamma(1 - \alpha - \lambda)}{\tau(\lambda + m + \alpha)\Gamma(-\lambda)\Gamma(1 - \alpha) + [L_m^{\alpha - 1}(0)]^2\Gamma(\alpha)\Gamma(1 - \alpha - \lambda)}.$ 

Hence, if  $\tau \notin \{0\} \cup \{\infty\}$ ,  $\mathcal{L}_{\tau}$  will have eigenvalues precisely when

$$\tau(\lambda+m+\alpha)\Gamma(-\lambda)\Gamma(1-\alpha) = -[L_m^{\alpha-1}(0)]^2\Gamma(\alpha)\Gamma(1-\alpha-\lambda).$$

Solving for  $\tau$  shows that eigenvalues of  $\mathcal{L}_{\tau}$  are just the level curves where  $M_{\infty}(\lambda) = \tau$ . Our example is now complete.

There are two main obstacles to generalizing the example that the reader should keep in mind. The first is the manipulation of the general solutions to the XOP expression so that they are in a practical format. This step is not necessary in the above example as all calculations are explicit. The second is the determination of the normalizations for initial conditions; this is found by a simple calculation in the example and stated in (3.7). In the general case, these challenges are the main subject of Sections 5 and 6, respectively.

#### 4. Exceptional Laguerre operators

In order to determine the spectrum of an XOP operator associated with  $\hat{\ell}$ , it is first necessary to determine the general solutions to the formal eigenvalue problem  $(\hat{\ell} - \lambda) f = 0$ . These solutions can be found through the intertwining property of Lemma 2.2. By [16, Section 3] the solutions may be written as the Wronskian of a sequence of seed functions along with a solution to the eigenvalue problem of the COP operator.

Seed functions are the quasi-rational eigenfunctions of the Laguerre COP differential expression in (2.3)–(2.6) and will be indexed via

$$f_j(x) = L_{n_j}^{\alpha}(x), \qquad j = 1, \dots, r_1, \qquad (4.1)$$

$$f_{r_1+j}(x) = e^x L^{\alpha}_{m_j}(-x), \qquad j = 1, \dots, r_2, \qquad (4.2)$$

$$f_{r_1+r_2+j}(x) = x^{-\alpha} L_{m'_j}^{-\alpha}(x), \qquad j = 1, \dots, r_3, \qquad (4.3)$$

$$f_{r_1+r_2+r_3+j}(x) = e^x x^{-\alpha} L_{n'_j}^{-\alpha}(-x), \qquad j = 1, \dots, r_4, \qquad (4.4)$$

with  $r_1 + r_2 + r_3 + r_4 = r$ ,  $n_1 > \cdots > n_{r_1} \ge 0$ ,  $m_1 > \cdots > m_{r_2} \ge 0$ ,  $n'_1 > \cdots > n'_{r_1} \ge 0$ and  $m'_1 > \cdots > m'_{r_4} \ge 0$ . The degrees of seed functions appearing in the Wronskian can be better described via two Maya diagrams, see Subsection 2.2. Denote these Maya diagrams by

$$M_1 = (n'_1, \dots, n'_{r_4} \mid n_1, \dots, n_{r_1}),$$
(4.5)

and

$$M_2 = (m'_1, \dots, m'_{r_3} \mid m_1, \dots, m_{r_2}).$$
(4.6)

The two general solutions of the classical Laguerre differential expression  $\ell^{\alpha}$  given by (3.1) are denoted

$$h^{\alpha}(x,\lambda) := M(-\lambda,\alpha+1,x) = {}_{1}F_{1}(-\lambda,\alpha+1,x) \quad \text{for } \alpha+1 \notin -\mathbb{N}_{0}, \quad (4.7)$$

and

 $\tilde{h}^{\alpha}(x,\lambda) := x^{-\alpha} M(-\lambda - \alpha, 1 - \alpha, x) \quad \text{for } \alpha + 1 \notin \mathbb{N} > 1.$ (4.8)

A generalized Laguerre polynomial,  $\Omega^{\alpha}_{M_1,M_2}$ , is a Wronskian of seed functions with the same parameter  $\alpha$  and distinct degrees. A prefactor is required to make it a polynomial. Let  $\Omega^{\alpha}_{M_1,M_2}$  be defined by

$$\Omega^{\alpha}_{M_1,M_2}(x) = e^{-(r_2 + r_4)x} x^{(\alpha + r_1 + r_2)(r_3 + r_4)} \operatorname{Wr}[f_1, \dots, f_r],$$
(4.9)

where  $f_1, \ldots, f_r$  are as in (4.1)-(4.4). If both Maya diagrams are trivial or the associated partitions are empty, the generalized Laguerre polynomial is a constant function.

Not every set of seed functions will produce a valid XOP expression. The following notion will determine admissibility; ensuring that the generalized Laguerre polynomial corresponds to a valid factorization chain as in Subsection 2. The Maya diagrams  $M_1$  and  $M_2$  of (4.5) and (4.6) can be presented as partitions, denoted respectively by  $\mu$  and  $\nu$  in canonical position or  $\mu'$  and  $\nu'$  in conjugate canonical position. The lengths of these partitions, denoted  $r(\mu)$  and  $r(\nu)$  or  $r(\mu')$  and  $r(\nu')$ , are determined by using the values  $t_1$ ,  $t_2$  and  $t'_1$ ,  $t'_2$  that needed to shift  $M_1$  and  $M_2$  into their canonical and conjugate canonical positions. In particular,

$$r(\mu) = \begin{cases} r_1 + n'_1 + 1 - r_4, & t_1 > 0, \\ r_1, & t_1 = 0, \\ r_1 - \min\{k \in \mathbb{N}_0 : k \notin \{n_i\}_{i=1}^{r_1}\}, & t_1 < 0, \end{cases}$$
(4.10)

and

$$r(\nu) = \begin{cases} r_2 + m'_1 + 1 - r_3, & t_2 > 0, \\ r_2, & t_2 = 0, \\ r_2 - \min\{k \in \mathbb{N}_0 : k \notin \{m_i\}_{i=1}^{r_2}\}, & t_2 < 0. \end{cases}$$
(4.11)

The lengths of the conjugate partitions,  $r(\mu')$  and  $r(\nu')$ , can be determined similarly but will not play an important role in our calculations.

In canonical position, the Maya diagram can thus be relabeled and written as a partition defined by

$$\mu = (\mu_1, \dots, \mu_{r(\mu)}), \quad \mu_j = n_j - r(\mu) + j, \quad j = 1, \dots, r(\mu), \tag{4.12}$$

and

$$v = (v_1, \dots, v_{r(v)}), \quad v_j = m_j - r(v) + j, \quad j = 1, \dots, r(v),$$
 (4.13)

where  $r(\mu) + r(\nu) = r$ . Conjugate partitions are relabeled so that  $r(\mu') + r(\nu') = r'$ and defined analogously. These values of  $n_j$  and  $m_j$  should not be confused with the entries of general Maya diagrams; they correspond only to those diagrams that are in canonical position. Partitions also define Young diagrams, where the number of boxes in each row corresponds to the entry of the partition. Conjugate partitions are then obtained by reflection over the main diagonal of the corresponding Young diagram.

**Definition 4.1.** A partition  $v = (v_1, ..., v_r)$  with  $v_r \ge 1$  is *even* if r is even and  $v_{2j-1} = v_{2j}$  for every j = 1, ..., r/2.

In order for a Maya diagram to represent a valid Laugerre XOP expression, we make the following assumption.

Assumption 4.2. The partition  $\mu$  is even.

It follows from [10, 11] that the polynomial  $\Omega^{\alpha}_{\mu,\nu}$  has no zeros on  $[0, \infty)$  if and only if  $\mu$  is an even partition. It is for this reason that we require an even partition for admissibility. Convention states that the empty partition (when r = 0) is even. For the remainder of the manuscript, it is assumed that all Maya diagrams will satisfy Assumption 4.2.

In order to apply the Darboux transform to the COP solutions of (4.7) and (4.8), it is necessary to add another entry to the corresponding Wronskian and adjust the prefactor. In particular, denote

$$\Omega_{M_1,M_2}[h^{\alpha}(x,\lambda)] = e^{-(r_2+r_4)x} x^{(\alpha+r_1+r_2+1)(r_3+r_4)} \operatorname{Wr}[f_1,\ldots,f_r,h^{\alpha}(x,\lambda)],$$
(4.14)

and

$$\Omega_{M_1,M_2}[\tilde{h}^{\alpha}(x,\lambda)] = e^{-(r_2+r_4)x} x^{(\alpha+r_1+r_2)(r_3+r_4+1)} \operatorname{Wr}[f_1,\ldots,f_r,\tilde{h}^{\alpha}(x,\lambda)].$$
(4.15)

For the sake of convenience, we will often refer to (4.14) as a solution of the first kind and (4.15) as a solution of the second kind. The parameter  $\alpha$  is written only for

the solution, but it should be understood that this is also the parameter for all seed functions in the Wronskian.

**Remark 4.3.** It is clear that adding a column with a solution  $h^{\alpha}(x, \lambda)$  or  $\tilde{h}^{\alpha}(x, \lambda)$  to the Wronskian changes the expression from a polynomial to an infinite series. However, consider the case  $\lambda = n$  in (4.14) where  $n \in \mathbb{N}_{\mu,\nu}$ , the natural numbers without a subset that is determined by  $\mu$  and  $\nu$ , see [6, Definition 2.9] for details. Then the series terminates, and if multiplied by a normalizing constant will yield an exceptional Laguerre polynomial  $L^{\alpha}_{\mu,\nu,n}$ . These polynomials can emerge from either solution, and will be discussed further in Assumption 7.2 and Remark 7.3.

We now direct our attention to the behavior of solutions near the singular endpoints x = 0 and  $x = \infty$  for the XOP Laguerre expression found in (2.8). Note that  $r(\mu) + r(\nu) = r$  here corresponds to the number of elements in the two partitions  $\mu$ and  $\nu$ , not in the original Maya diagrams.

**Theorem 4.4.** For  $\alpha + r > -1$ , let  $\hat{\ell}^{\alpha+r}$  be the XOP differential expression on the interval  $(0, \infty)$ .

- (a.)  $\hat{\ell}^{\alpha+r}$  is in the limit-circle case at x = 0 for  $-1 < \alpha + r < 1$  and is in the limit-point case at x = 0 when  $\alpha + r \ge 1$ .
- (b.)  $\hat{\ell}^{\alpha+r}$  is in the limit-point case at  $x = \infty$  for any  $\alpha + r > -1$ .

*Proof.* The point x = 0 is, in the sense of Frobenius, a regular singular endpoint of the XOP expression  $(\hat{\ell}^{\alpha+r} - \lambda) f = 0$  for any value  $\lambda \in \mathbb{C}$ . Using the classical Laguerre expression (3.1) to derive  $\hat{\ell}^{\alpha+r}$  associated with (2.8) yields an indicial equation of

$$k(k + \alpha + r) = 0.$$

This indicial equation holds for any admissible choice of  $\phi$  and b. Meanwhile, the point  $x = \infty$  is an irregular singular point and a reduction of order method must be used to find two linearly independent solutions. The result may be shown using the techniques within the proof of [26, Theorem 3.3].

Since  $\hat{\ell}^{\alpha+r}$  is in the limit-circle case at x = 0 for  $-1 < \alpha + r < 1$ , Glazman–Krein–Naimark theory requires that one appropriate boundary condition be imposed in order to generate a self-adjoint extension of the minimal operator. Thus, if  $-1 < \alpha + r < 1$ , the deficiency index of  $\mathcal{L}_{\min}$  is (1, 1). Meanwhile, if  $\alpha + r \ge 1$ , the deficiency index is (0, 0).

Assumption 4.5. The XOP Laguerre expression  $\hat{\ell}^{\alpha+r}$  is assumed to have parameter  $\alpha + r > -1$ . Furthermore,  $\alpha \notin \mathbb{Z}$ .

The assumption on  $\alpha$  follows from the definitions of the confluent hypergeometric solutions in equations (4.7) and (4.8).

Under our Assumptions 4.2 and 4.5, these Laguerre XOPs form a complete set of orthogonal polynomials on the positive real line [10, 11].

**Lemma 4.6.** Suppose  $\alpha + r > -1$  and  $\mu$  is an even partition. Then the polynomials  $L^{\alpha}_{\mu,\nu,n}$  for  $n \in \mathbb{N}_{\mu,\nu}$  are orthogonal on  $[0,\infty)$  with respect to the positive weight function associated with  $\hat{\ell}^{\alpha+r}$ :

$$W^{\alpha}_{\mu,\nu}(x) = \frac{x^{\alpha+r}e^{-x}}{[\Omega^{\alpha}_{\mu,\nu}(x)]^2}, \quad x > 0.$$

That is, if  $n, m \in \mathbb{N}_{\mu,\nu}$  with  $n \neq m$ , then

$$\int_{0}^{\infty} L^{\alpha}_{\mu,\nu,n}(x) L^{\alpha}_{\mu,\nu,m}(x) W^{\alpha}_{\mu,\nu}(x) dx = 0.$$

Moreover, they form a complete orthogonal set in  $L^2([0,\infty), W^{\alpha}_{\mu,\nu}(x)dx)$ .

Note that the assumptions of Lemma 4.6 are not the best possible conditions, but they are convenient and straightforward to apply. More specific conditions for admissibility can be found in [10, 11].

Also, the solution of the second kind defined by (4.15) does not currently possess the correct asymptotic behavior near 0; it should have asymptotic behavior of order  $x^{-\alpha-r}$  to fit the Frobenius analysis of the corresponding exceptional Laguerre expression in Theorem 4.4. The current formulation of this solution of the second kind as a polynomial is simply more convenient to work with and avoids further complicating notation. This discrepancy will be rectified in Section 6, where other behavior near x = 0 is needed to ensure the solution is in the proper Hilbert space.

Finally, the construction of a general exceptional Laguerre expression requires beginning with the classical Laguerre expression, which has a parameter  $\alpha$ , and applying Darboux transforms. In this process,  $\alpha$  may be shifted. Therefore, the parameter  $\alpha$  in Lemma 4.6 should not be thought of as simply the same  $\alpha$  from the original COP expression. One of the main goals of Section 5 is to track the necessary changes to the classical  $\alpha$  as operations are performed.

#### 5. Manipulation of Maya diagrams

Information must be extracted from the two solutions of the XOP Laguerre expression given in (4.14) and (4.15) in order to build the Weyl *m*-function. The results follow the general methods of [6, Theorem 4.2, Lemma 4.4], which may be consulted for additional insight. The presentation here involves significant additions. Most importantly,

it is necessary to track changes of the spectral parameter through the shifts as well as any constants produced by the spectral parameter. The format of these solutions can vary significantly based on how the Maya diagrams are shifted. In this section, the Maya diagrams are translated into standard positions where the necessary information can be extracted. Solutions of the first kind have their Maya diagrams shifted into canonical position, while solutions of the second kind are shifted into conjugate canonical position. A few additional assumptions are necessary.

Assumption 5.1. Without loss of generality, we assume that any shifts necessary to bring  $M_1$  to canonical form are taken prior to any shifts that may be necessary for  $M_2$ . Indeed, the steps of the shifting process outlined in Theorems 5.3 and 5.5 may be applied to  $M_1$  and  $M_2$  in a variety of orders, but this assumption minimizes notational complexity.

Assumption 5.2. Factors of  $(-1)^d$  for some  $d \in \mathbb{N}$  that arise in calculations are omitted throughout the remainder of the manuscript. Tracking this factor requires substantial notation and does not meaningfully contribute to results. Such a factor appears in the equations of Theorems 5.3, 5.5, 6.1, Corollaries 5.7, 6.2, 6.3, and the *m*-functions of Section 7. Therefore, all equalities in these results are correct up to a change of sign. The cumulative effect of these factors throughout the construction of the Weyl *m*-function can still be determined by enforcing the property that this function is a so-called Nevanlinna–Herglotz function: an analytic self-map of the upper half-plane. See i.e. [15, Section 6] for examples where this property is utilized in a similar context.

We begin by shifting the Maya diagrams defining a solution of the first kind into canonical position. The notation

$$x^{(n)} = x(x+1)\cdots(x+n-1)$$

and

$$x_{(n)} = x(x-1)\cdots(x-n+1)$$

is used to denote the rising and falling factorials, respectively.

**Theorem 5.3.** Let  $f_1, \ldots, f_r$  be as in (4.1)-(4.4),  $M_1$  and  $M_2$  be as in (4.5) and (4.6), and  $\mu$  and  $\nu$  be their associated partitions after shifts  $t_1$  and  $t_2$  to canonical position, respectively. Then for  $\lambda \neq n_i$  with  $i = 1, \ldots, r_1$ ,

$$\Omega_{M_1,M_2}[h^{\alpha}(x,\lambda)] = C_1(\alpha,\lambda,M_1)C_2(\alpha,\lambda,M_2)C_3(\alpha,M_1,M_2)\Omega_{\mu,\nu}[h^{\alpha-t_1-t_2}(x,\lambda+t_1)],$$

where

$$C_{1}(\alpha, \lambda, M_{1}) = \begin{cases} \frac{(-\lambda)^{(lt_{1}|)}}{(\alpha+1)^{(lt_{1}|)}} & \text{for } t_{1} < 0, \\ 1 & \text{for } t_{1} = 0, \\ \frac{(\alpha)_{(t_{1})}}{\prod_{k_{1}}(-\lambda-k_{1}-1)} & \text{for } t_{1} > 0, \end{cases}$$
(5.1)

for  $k_1 \in \{0, \dots, t_1 - 1\}$  such that  $k_1 \notin \{n'_j\}_{j=1}^{r_4}$ ;

$$C_{2}(\alpha, \lambda, M_{2}) = \begin{cases} \frac{(\alpha+1+\lambda)^{(l_{2})}}{(\alpha-t_{1}+1)^{(l_{2})}} & \text{for } t_{2} < 0, \\ 1 & \text{for } t_{2} = 0, \\ \frac{(\alpha-t_{1})_{(t_{2})}}{\prod_{k_{2}}(\lambda+\alpha-k_{2})} & \text{for } t_{2} > 0, \end{cases}$$
(5.2)

for  $k_2 \in \{0, ..., t_2 - 1\}$  such that  $k_2 \notin \{m'_j\}_{j=1}^{r_3}$ ; and

$$C_{3}(\alpha, M_{1}, M_{2}) = \prod_{j=1}^{r_{1}} \prod_{k=1}^{r_{3}} (m'_{k} - \alpha - n_{j}) \prod_{j=1}^{r_{2}} \prod_{k=1}^{r_{4}} (n'_{k} - \alpha - m_{j}) \\ \times \prod_{j=1}^{r_{1}} \prod_{k=1}^{r_{4}} (n_{j} + n'_{k} + 1) \prod_{j=1}^{r_{2}} \prod_{k=1}^{r_{3}} (m_{j} + m'_{k} + 1).$$
(5.3)

The constants from Theorem 5.3 will be collectively denoted as

$$C := C_1(\alpha, \lambda, M_1) C_2(\alpha, \lambda, M_2) C_3(\alpha, M_1, M_2).$$
(5.4)

The proof of Theorem 5.3 requires several intermediate steps. Recall the following derivative identities:

$$\frac{d}{dx}(L_n^{\alpha}(x)) = -L_{n-1}^{\alpha+1}(x),$$
(5.5a)

$$\frac{d}{dx}(e^{x}L_{n}^{\alpha}(-x)) = e^{x}L_{n}^{\alpha+1}(-x),$$
(5.5b)

$$\frac{d}{dx}(x^{-\alpha}L_n^{-\alpha}(x)) = (n-\alpha)x^{-\alpha-1}L_n^{-\alpha-1}(x), \qquad (5.5c)$$

$$\frac{d}{dx}(x^{-\alpha}e^{x}L_{n}^{-\alpha}(-x)) = (n+1)x^{-\alpha-1}e^{x}L_{n+1}^{-\alpha-1}(-x).$$
(5.5d)

The derivatives of (5.5) involve normalizing constants, which result in significant cancellations, but otherwise are special cases of the following derivative identities for

confluent hypergeometric functions

$$\frac{d}{dx}(M(a,b,x)) = \frac{a}{b}M(a+1,b+1,x),$$
(5.6a)

$$\frac{d}{dx}(e^{x}M(a,b,-x)) = \frac{b-a}{b}e^{x}M(a,b+1,-x),$$
(5.6b)

$$\frac{d}{dx}(x^{b-1}M(a,b,x)) = (b-1)x^{b-2}M(a,b-1,x),$$
(5.6c)

$$\frac{d}{dx}(x^{b-1}e^x M(a,b,-x)) = -(b-1)x^{b-2}e^x M(a-1,b-1,-x).$$
(5.6d)

The results of (5.6) will be used to take derivatives of h and  $\tilde{h}$ .

Additionally, the following elementary Wronskian identities that will be useful for calculations. Assume that the functions  $f_1, \ldots, f_r, g$ , and h are all sufficiently differentiable. Then

$$Wr[gf_1, \dots, gf_r] = (g(x))^r Wr[f_1, \dots, f_r],$$
(5.7)  
$$Wr[f_1 \circ h, \dots, f_r \circ h] = (h'(x))^{\frac{r(r-1)}{2}} Wr[f_1, \dots, f_r] \circ h(x).$$

To prove Theorem 5.3, we begin with a lemma that involves simplifications when 0 appears in the encoding of one of the Maya diagrams  $M_1$  or  $M_2$ . This is an alteration of [6, Lemma 4.4].

**Lemma 5.4.** Let  $M_1$  and  $M_2$  be given by (4.5) and (4.6).

(a) If 
$$n_{r_1} = 0$$
, then  

$$\Omega_{M_1,M_2}[h^{\alpha}(x,\lambda)]$$

$$= \left(\frac{-\lambda}{\alpha+1}\prod_{i=1}^{r_3}(m'_i - \alpha)\prod_{i=1}^{r_4}(n'_i + 1)\right)\Omega_{M_1 - 1,M_2}[h^{\alpha+1}(x,\lambda-1)].$$
 (5.8)

(b) If  $n'_{r_4} = 0$ , then

$$\Omega_{M_1,M_2}[h^{\alpha}(x,\lambda)] = \left(\alpha \prod_{i=1}^{r_1} (n_i+1) \prod_{i=1}^{r_2} (m_i+\alpha)\right) \Omega_{M_1+1,M_2}[h^{\alpha-1}(x,\lambda+1)].$$
(5.9)

(c) If  $m_{r_2} = 0$ , then

$$\Omega_{M_1,M_2}[h^{\alpha}(x,\lambda)] = \left(\frac{\alpha+1+\lambda}{\alpha+1}\prod_{i=1}^{r_3}(m'_i+1)\prod_{i=1}^{r_4}(n'_i-\alpha)\right)\Omega_{M_1,M_2-1}[h^{\alpha+1}(x,\lambda)].$$
(5.10)

(d) If  $m'_{r_3} = 0$ , then

$$\Omega_{M_1,M_2}[h^{\alpha}(x,\lambda)] = \left(\alpha \prod_{i=1}^{r_1} (n_i + \alpha) \prod_{i=1}^{r_2} (m_i + 1)\right) \Omega_{M_1,M_2+1}[h^{\alpha-1}(x,\lambda)].$$
(5.11)

*Proof.* We give only the proof of (b) as the other claims follow from using the same procedure. Let  $n'_{r_4} = 0$ . Then  $f_r = e^x x^{-\alpha}$ . Removing a factor of  $e^x x^{-\alpha}$  from each column via (5.7) and expanding about the second to last column yields

$$\begin{aligned} \Omega_{M_1,M_2}[h^{\alpha}(x,\lambda)] \\ &= (e^x x^{-\alpha})^{(r+1)} e^{-(r_2+r_4)x} x^{(\alpha+r_1+r_2+1)(r_3+r_4)} \\ &\times \operatorname{Wr}[e^{-x} x^{\alpha} f_1, \dots, e^{-x} x^{\alpha} f_{r-1}, 1, e^{-x} x^{\alpha} h^{\alpha}(x,\lambda)] \\ &= (-1)^{r-1} (e^x x^{-\alpha})^{(r+1)} e^{-(r_2+r_4)x} x^{(\alpha+r_1+r_2+1)(r_3+r_4)} \\ &\times \operatorname{Wr}\left[\frac{d}{dx} (e^{-x} x^{\alpha} f_1), \dots, \frac{d}{dx} (e^{-x} x^{\alpha} f_{r-1}), \frac{d}{dx} (e^{-x} x^{\alpha} h^{\alpha}(x,\lambda))\right]. \end{aligned}$$

Let  $\tilde{M}_1 = M_1 + 1$  and let  $\tilde{f}_1, \ldots, \tilde{f}_{\tilde{r}}$  be the functions associated with the Maya diagram  $\tilde{M}_1$  and parameter  $\tilde{\alpha} = \alpha - 1$ . Also let  $\tilde{r}_1 = r_1, \tilde{r}_2 = r_2, \tilde{r}_3 = r_3, \tilde{r}_4 = r_4 - 1$  and  $\tilde{r} = r - 1$ . Equations (5.5) and (5.6) imply that

$$\frac{d}{dx}(e^{-x}x^{\alpha}f_j) = c_j e^{-x}x^{\alpha-1}\tilde{f_j} \quad \text{for } j = 1, \dots, r-1,$$

and

$$\frac{d}{dx}(e^{-x}x^{\alpha}M(-\lambda,\alpha+1,x)) = \alpha e^{-x}x^{\alpha-1}M(-\lambda-1,\alpha,x)$$
$$= \alpha e^{-x}x^{\alpha-1}h^{\alpha-1}(x,\lambda+1),$$

where  $c_j = n_j + 1$  for  $j = 1, ..., r_1$ ;  $c_{r_1+j} = m_j + \alpha$  for  $j = 1, ..., r_2$ ;  $c_{r_1+r_2+j} = -1$  for  $j = 1, ..., r_3$ ; and  $c_{r_1+r_2+r_3+j} = 1$  for  $j = 1, ..., r_4 - 1$ . Each column of the Wronskian has a corresponding constant removed, and then the common factor  $e^{-x}x^{\alpha-1}$  is removed by (5.7), yielding

$$\begin{aligned} \Omega_{M_1,M_2}[h^{\alpha}(x,\lambda)] \\ &= \tilde{c}_1(e^x x^{-\alpha})^{(r+1)}(e^{-x} x^{\alpha-1})^{(r)}e^{-(r_2+r_4)x} x^{(\alpha+r_1+r_2+1)(r_3+r_4)} \\ &\quad \times \operatorname{Wr}[\tilde{f_1},\ldots,\tilde{f_{r-1}},h^{\alpha-1}(x,\lambda+1)] \\ &= \tilde{c}_1 e^{-(r_2+r_4-1)x} x^{(\alpha+r_1+r_2+1)(r_3+r_4)-\alpha-r} \\ &\quad \times \operatorname{Wr}[\tilde{f_1},\ldots,\tilde{f_{r-1}},h^{\alpha-1}(x,\lambda+1)], \end{aligned}$$

where

$$\tilde{c}_1 = \alpha (-1)^{r-1} \prod_{j=1}^{r_4-1} c_j = \alpha (-1)^{r_1+r_2+r_4-1} \prod_{j=1}^{r_1} (n_j+1) \prod_{j=1}^{r_2} (m_j+\alpha).$$

Note the analogous step in the proof of [6, Lemma 4.4] omits a  $-\alpha$  term from the exponent of x but the end result is unchanged. Claim (b) follows because

$$(\alpha + r_1 + r_2 + 1)(r_3 + r_4) - \alpha - r = (\alpha + r_1 + r_2)(r_3 + r_4 - 1)$$
$$= (\tilde{\alpha} + \tilde{r}_1 + \tilde{r}_2 + 1)(\tilde{r}_3 + \tilde{r}_4).$$

Collectively, (a)–(d) of Lemma 5.4 describe how the spectral parameter  $\lambda$  and  $\alpha$  change along with the constants produced with any shift of  $M_1$  or  $M_2$  one unit left or right. We will continue to track of these changes while iterating the shifts until the Maya diagrams are in canonical or conjugate canonical form. Note that we will appeal to [6, Section 4.4] to show the exact form of the products in the statement of Theorem 5.3, as there are no alterations to that argument, we simply keep track of additional parameters.

*Proof of Theorem* 5.3. To begin, we verify (5.1) and (5.2). These results only consider contributions of non-product coefficient factors found in (5.8)-(5.11). These product terms are collected and addressed in the verification of (5.3).

We begin by shifting  $M_1$  into canonical form. If  $t_1 = 0$ , the Maya diagram  $M_1$  is already in canonical form and no shift is required. If  $t_1 < 0$ , then  $|t_1|$  boxes to the immediate right of the origin are filled; that is,  $n_{r_1-j} = j$  for  $j = 0, ..., |t_1| - 1$ . Hence, (5.8) will be applied  $|t_1|$  times to  $M_1$ . Therefore,  $|t_1|$  applications of (5.8) yield coefficient factor

$$\frac{(-\lambda)^{(|t_1|)}}{(\alpha+1)^{(|t_1|)}}.$$

Note that for  $t_1 < 0$ , the parameters of h are now  $\alpha - t_1$  and  $\lambda + t_1$ .

If  $t_1 > 0$ , denote

$$\tilde{M}_1 = M_1 + 1: (\tilde{n}'_1, \dots, \tilde{n}'_{\tilde{r}_4} \mid \tilde{n}_1, \dots, \tilde{n}_{\tilde{r}_1})$$

Determining the appropriate equation of Lemma 5.4 to apply depends upon whether the box immediately to the left of the origin of  $M_1$  is or is not filled. There are two cases.

(i) If the box to the left of the origin is empty, then  $n'_{r_4} = 0$ ,  $\tilde{r}_4 = r_4 - 1$ ,  $\tilde{r}_1 = r_1$ , and

$$\widetilde{M}_1: (n'_1 - 1, \dots, n'_{r_4 - 1} - 1 \mid n_1 + 1, \dots, n_{r_1} + 1)$$

(ii) If the box to the left of the origin is filled, then  $\tilde{n}_{r_1+1} = 0$ ,  $\tilde{r}_4 = r_4$ ,  $\tilde{r}_1 = r_1 + 1$ , and

$$\widetilde{M}_1: (n'_1 - 1, \dots, n'_{r_4-1} - 1 \mid n_1 + 1, \dots, n_{r_1} + 1, 0).$$

In total,  $M_1$  requires  $t_1$  shifts in order to be in canonical form. There will be  $r_4$  boxes to the left of the origin that are empty; hence we are in case (i) of above and (5.9) will be applied  $r_4$  times. As a result, each application of (5.9) lowers  $\alpha$  by one and increases  $\lambda$  by one. Additionally, for each  $i \in \{1, \ldots, r_4\}$ , a factor of  $(\alpha - n'_i)$  will be contributed to the coefficient. Case (ii) occurs  $t_1 - r_4$  times and (5.8) will be applied. As  $M_1$  needs to be shifted to the right, (5.8) must be restated as

$$\Omega_{M_1,M_2}[h^{\alpha}(x,\lambda)] = \left[ \left( \frac{-\lambda - 1}{\alpha} \right) \prod_{i=1}^{r_3} (m'_i - \alpha + 1) \prod_{i=1}^{r_4} n'_i \right]^{-1} \Omega_{M_1 + 1,M_2}[h^{\alpha - 1}(x,\lambda + 1)].$$

Each application of (5.8) also lowers  $\alpha$  by one and increases  $\lambda$  by one. A factor of  $(\alpha - k_1)/(-\lambda - k_1 - 1)$  for each  $k_1 \in \{0, \ldots, t_1 - 1\}$  such that  $k_1 \notin \{n'_j\}_{j=1}^{r_4}$  will be collected by the coefficient.

Together contributions from all case (i) and case (ii) shifts produce a coefficient of

$$\frac{(\alpha)_{(t_1)}}{\prod_{k_1}(-\lambda-k_1-1)},$$

and the parameters of h are now  $\alpha - t_1$  and  $\lambda + t_1$ . Equation (5.1) has been verified.

Now, we consider the adjustments required to shift  $M_2$  into canonical form. As  $M_1$  is first put into canonical form, the parameters for h are now  $\alpha - t_1$  and  $\lambda + t_1$  for  $t_1 \in \mathbb{Z}$ . If  $t_2 = 0$ , the Maya diagram  $M_2$  is already in canonical form and no shift is required. If  $t_2 < 0$ , then  $|t_2|$  boxes to the immediate right of the origin are filled; that is,  $m_{r_2-j} = j$  for  $j = 0, \ldots, |t_2| - 1$ . Hence, (5.10) will be applied  $|t_2|$  times to  $M_2$ . Therefore,  $|t_2|$  applications of (5.10) yields additional coefficient factors

$$\frac{(\alpha + 1 + \lambda)^{(|t_2|)}}{(\alpha - t_1 + 1)^{|t_2|}}.$$

For  $t_2 > 0$ , verification follows in a similar fashion as the  $M_1$  case except it relies on the reductions, and therefore the constants, from (5.10) and its restatement:

$$\Omega_{M_1,M_2}[h^{\alpha}(x,\lambda)] = \left[ \left(\frac{\alpha+\lambda}{\alpha}\right) \prod_{i=1}^{r_3} m'_i \prod_{i=1}^{r_4} (n'_i - \alpha + 1) \right]^{-1} \Omega_{M_1,M_2+1}[h^{\alpha-1}(x,\lambda)].$$

The contribution of the products found in (5.8)–(5.11) is contained in  $C_3$  and the proof is identical to [6, Section 4.4].

The method of proof of Theorem 5.3 may be repeated with a solution of the second kind, but there are a few changes. First, it is necessary to state the result with conjugate partitions; shifting the Maya diagrams to the conjugate canonical position so that the partitions are determined by the eigenfunctions in (4.3) and (4.4). This is due to the fact that the solution  $\tilde{h}$  has the same format and similar derivative formula as the seed function (4.3)—a shift to canonical position would not change this fact. The conjugate canonical position, on the other hand, still allows the use of partitions. Second, there is no formula giving the product of constants from the shifting functions (stated as  $C_3$  in Theorem 5.3) in the literature. It is therefore necessary to prove such a formula.

**Theorem 5.5.** Let  $f_1, \ldots, f_r$  be as in (4.1)–(4.4),  $M_1$  and  $M_2$  be as in (4.5) and (4.6), and  $\mu', \nu'$  be their associated conjugate partitions after shifts  $t'_1$  and  $t'_2$  to conjugate canonical position, respectively. Then for  $\lambda \neq m'_i$  with  $i = 1, \ldots, r_3$ ,

$$\begin{aligned} \Omega_{M_1,M_2}[\hat{h}^{\alpha}(x,\lambda)] \\ &= D_1(\alpha,\lambda,M_1)D_2(\alpha,\lambda,M_2)D_3(\alpha,M_1,M_2)\Omega_{\mu',\nu'}[\tilde{h}^{\alpha-t_1'-t_2'}(x,\lambda+t_1')], \end{aligned}$$

where

$$D_{1}(\alpha, \lambda, M_{1}) = \begin{cases} \frac{(-\alpha)_{(|t_{1}'|)}}{\prod_{k_{1}'} (\lambda - k_{1}')} & \text{for } t_{1}' < 0, \\ 1 & \text{for } t_{1}' = 0, \\ \frac{(\lambda + 1)^{(t_{1}')}}{(1 - \alpha)^{(t_{1}')}} & \text{for } t_{1}' > 0, \end{cases}$$
(5.12)

for  $k'_1 \in \{0, \dots, |t'_1| - 1\}$  such that  $k'_1 \notin \{n_j\}_{j=1}^{r_1}$ ;

$$D_{2}(\alpha, \lambda, M_{2}) = \begin{cases} \frac{(-\alpha + t_{1}')_{(|t_{2}'|)}}{\prod_{k_{2}'}(-\lambda - \alpha - 1 - k_{2}')} & \text{for } t_{2}' < 0, \\ 1 & \text{for } t_{2}' = 0, \\ \frac{(-\lambda - \alpha)^{(t_{2}')}}{(1 - \alpha + t_{1}')^{(t_{2}')}} & \text{for } t_{2}' > 0, \end{cases}$$
(5.13)

for  $k'_2 \in \{0, \dots, |t'_2| - 1\}$  such that  $k'_2 \notin \{m_j\}_{j=1}^{r_2}$ ;

$$D_{3}(\alpha, M_{1}, M_{2}) = \prod_{j=1}^{r_{1}} \prod_{k=1}^{r_{3}} (n_{j} + \alpha - m_{k}') \prod_{j=1}^{r_{2}} \prod_{k=1}^{r_{4}} (m_{j} + \alpha - n_{k}')$$
$$\times \prod_{j=1}^{r_{1}} \prod_{k=1}^{r_{4}} (n_{j} + n_{k}' + 1) \prod_{j=1}^{r_{2}} \prod_{k=1}^{r_{3}} (m_{j} + m_{k}' + 1).$$
(5.14)

The constants from Theorem 5.5 are collectively denoted as

$$D := D_1(\alpha, \lambda, M_1) D_2(\alpha, \lambda, M_2) D_3(\alpha, M_1, M_2),$$
(5.15)

and will be used in Section 7.

The change in parameters and constants is now tracked for a shift of one unit in either direction for the Maya diagrams. These are similar to those in Lemma 5.4 but with extra contributions emerging from  $\tilde{h}$ .

**Lemma 5.6.** Let  $M_1$  and  $M_2$  be given by (4.5) and (4.6).

(a) If 
$$n_{r_1} = 0$$
, then  

$$\Omega_{M_1,M_2}[\tilde{h}^{\alpha}(x,\lambda)] = \left((-\alpha)\prod_{i=1}^{r_3} (m'_i - \alpha)\prod_{i=1}^{r_4} (n'_i + 1)\right)\Omega_{M_1 - 1,M_2}[\tilde{h}^{\alpha + 1}(x,\lambda - 1)].$$

(b) If  $n'_{r_4} = 0$ , then

$$\Omega_{M_1,M_2}[\tilde{h}^{\alpha}(x,\lambda)] = \left(\frac{\lambda+1}{1-\alpha}\prod_{i=1}^{r_1}(n_i+1)\prod_{i=1}^{r_2}(m_i+\alpha)\right)\Omega_{M_1+1,M_2}[\tilde{h}^{\alpha-1}(x,\lambda+1)].$$
(5.16)

(c) If 
$$m_{r_2} = 0$$
, then

$$\Omega_{M_1,M_2}[\tilde{h}^{\alpha}(x,\lambda)] = \left((-\alpha)\prod_{i=1}^{r_3} (m'_i+1)\prod_{i=1}^{r_4} (n'_i-\alpha)\right)\Omega_{M_1,M_2-1}[\tilde{h}^{\alpha+1}(x,\lambda)]$$

(d) If 
$$m'_{r_3} = 0$$
, then

$$\Omega_{M_1,M_2}[\tilde{h}^{\alpha}(x,\lambda)] = \left(\frac{-\lambda-\alpha}{1-\alpha}\prod_{i=1}^{r_1}(n_i+\alpha)\prod_{i=1}^{r_2}(m_i+1)\right)\Omega_{M_1,M_2+1}[\tilde{h}^{\alpha-1}(x,\lambda)].$$
(5.17)

*Proof.* It is sufficient to use the derivative formulas in (5.6) to calculate the changed parameters and constants in each of the situations. Otherwise, the proof is completely analogous to that of Lemma 5.4.

Verifying (5.12) and (5.13), the formulas for  $D_1$  and  $D_2$ , follow the same procedure as the proof of Theorem 5.3, but relies on the equations found within Lemma 5.6 and so proofs of these formulas are omitted here. The method also differs slightly from that of Theorem 5.3 as the Maya diagrams are to be shifted into the conjugate canonical form. However, as no existing result addresses the formulation of (5.14), the proof is included below. The method closely follows the result of [6, Section 4.4].

*Proof of Theorem* 5.5. We note that the shifting procedure for  $t_1 < 0$  depends on the Maya diagram for  $M_1$  and whether the box of immediately to the right of the origin is or is not filled. Then, for  $\tilde{M}_1 = M_1 - 1$ , there are two cases to consider.

(i) If the box to the right of the origin is filled then  $n_{r_1} = 0$ ,  $\tilde{r}_1 = r_1 - 1$ ,  $\tilde{r}_4 = r_4$ , and

$$\tilde{M}_1: (n'_1 + 1, \dots, n'_{r_4} + 1 \mid n_{r_1 - 1} - 1, \dots, n_{r_1} - 1).$$
(5.18)

(ii) If the box to the right of the origin is empty then  $\tilde{n}'_{r_4+1} = 0$ ,  $\tilde{r}_4 = r_4 + 1$ ,  $\tilde{r}_1 = r_1$ , and

$$\widetilde{M}_1: (n'_1 + 1, \dots, n'_{r_4} + 1, 0 \mid n_1 - 1, \dots, n_{r_1} - 1).$$
(5.19)

In case (ii), (5.16) is rewritten as

$$\Omega_{M_1,M_2}[\tilde{h}^{\alpha}(x,\lambda)] = \left[ \left(\frac{\lambda}{-\alpha}\right) \prod_{i=1}^{r_1} n_i \prod_{i=1}^{r_2} (m_i + \alpha + 1) \right]^{-1} \Omega_{M_1 - 1,M_2}[\tilde{h}^{\alpha + 1}(x,\lambda - 1)], \quad (5.20)$$

before application.

Equation (5.14) will be shown using induction on the value of  $T' = |t'_1| + |t'_2|$ . For T' = 0, no shift of either  $M_1$  or  $M_2$  is required. As a result of  $r_1 = r_2 = 0$ , all products are empty and equal to 1. Therefore,  $D_3 = 1$  and the base case is shown. Let T' > 0 and assume that (5.14) holds whenever  $|t'_1| + |t'_2| = T' - 1$ . Considering now  $|t'_1| + |t'_2| = T'$ , there are a number of cases that may occur. When both  $t'_1, t'_2 > 0$ , as in the case when T' = 0,  $r_1 = r_2 = 0$  and all products are equal to 1. Therefore, we assume that either  $t'_1 < 0$  or  $t'_2 < 0$ . Without loss of generality, consider  $t'_1 < 0$ . The following cases can happen:

- (i)  $t'_1 < 0$  and  $\tilde{M}_1$  is given by (5.18) and
- (ii)  $t'_1 < 0$  and  $\tilde{M}_1$  is given by (5.19).

The treatment of each case is similar, so we will prove only case (ii). In this situation, (5.20) holds and

$$D_3(\alpha, M_1, M_2) = D_3(\alpha + 1, M_1 - 1, M_2) \left(\prod_{i=1}^{r_1} n_i \prod_{i=1}^{r_2} (m_i + \alpha + 1)\right)^{-1}.$$
 (5.21)

The induction hypothesis states

$$D_{3}(\alpha + 1, M_{1} - 1, M_{2})$$

$$= \prod_{j=1}^{\tilde{r}_{1}} \prod_{k=1}^{r_{3}} (\tilde{n}_{j} + (\alpha + 1) - m'_{k}) \prod_{j=1}^{r_{2}} \prod_{k=1}^{\tilde{r}_{4}} (m_{j} + (\alpha + 1) - \tilde{n}'_{k})$$

$$\times \prod_{j=1}^{\tilde{r}_{1}} \prod_{k=1}^{\tilde{r}_{4}} (\tilde{n}_{j} + \tilde{n}'_{k} + 1) \prod_{j=1}^{r_{2}} \prod_{k=1}^{r_{3}} (m_{j} + m'_{k} + 1).$$

The first three double products can be rewritten as

$$\prod_{j=1}^{\tilde{r}_1} \prod_{k=1}^{r_3} (\tilde{n}_j + (\alpha + 1) - m'_k) = \prod_{j=1}^{r_1} \prod_{k=1}^{r_3} (n_j + \alpha - m'_k),$$
  
$$\prod_{j=1}^{r_2} \prod_{k=1}^{\tilde{r}_4} (m_j + (\alpha + 1) - \tilde{n}'_k) = \prod_{j=1}^{r_2} \prod_{k=1}^{r_4} (m_j + \alpha - n'_k) \prod_{j=1}^{r_2} (m_j + \alpha + 1),$$
  
$$\prod_{j=1}^{\tilde{r}_1} \prod_{k=1}^{\tilde{r}_4} (\tilde{n}_j + \tilde{n}'_k + 1) = \prod_{j=1}^{r_1} \prod_{k=1}^{r_4} (n_j + n'_k + 1) \prod_{j=1}^{r_1} n_j.$$

The extra products now cancel in (5.21) and the formula for  $D_3$  in (5.14) holds.

Theorems 5.3 and 5.5 can both be simplified in the case where only the generalized Laguerre polynomial given by (4.9) is considered, i.e. there is no general solution to  $\ell^{\alpha}$  in the last column of the Wronskian. This is precisely (4.9). The constants generated by shifting this generalized Laguerre polynomial appear in Section 7 but will not affect any spectral properties of the exceptional operators; they are identified here for convenience. Note that the first half of the result was shown in [6, Theorem 4.2].

**Corollary 5.7.** Let  $f_1, \ldots, f_r$  be as in (4.1)–(4.4),  $M_1$  and  $M_2$  be as in (4.5) and (4.6),  $\mu$  and  $\nu$  be their associated partitions after shifts  $t_1$  and  $t_2$  to canonical position, and  $\mu'$ ,  $\nu'$  be their associated partitions after shifts  $t'_1$  and  $t'_2$  to conjugate canonical position. Recall that  $C_3$  and  $D_3$  are defined by (5.3) and (5.14), respectively. Then,

$$\Omega^{\alpha}_{M_1,M_2}(x) = C_3(\alpha, M_1, M_2) \Omega^{\alpha - t_1 - t_2}_{\mu,\nu}(x) = D_3(\alpha, M_1, M_2) \Omega^{\alpha - t_1' - t_2'}_{\mu',\nu'}(x)$$

*Proof.* The terms  $C_1$  and  $C_2$  from Theorem 5.3 are clearly produced solely by the solution  $h^{\alpha}(x, \lambda)$  in the last column of the Wronskian when the shifts to canonical position are performed. When this last column is removed, the cumulative effect is only determined by  $C_3$ . Analogously,  $D_3$  is the cumulative effect of shifting the Maya diagrams to conjugate canonical position.

#### 6. Initial condition normalizations

Theorems 5.3 and 5.5 allow for solutions of the first and second kind to be written in terms of Wronskians indexed by partitions. This section will extract information from these simplified Wronskians to determine how to form a fundamental system of solutions  $(u_1(x, \lambda) \text{ and } u_2(x, \lambda))$  for  $(\mathcal{L}_{\max} - \lambda) f = 0$  that is properly normalized according to (2.14). We leave the exact definitions of this system of solutions and the relevant quasi-derivatives to Section 7; now it is enough to calculate the value of these solutions at 0.

It is necessary to recall some notation and identities. Classical Laguerre polynomials satisfy

$$L_n^{\alpha}(0) = \frac{(\alpha+1)^{(n)}}{n!} = \frac{(\alpha+1)(\alpha+2)\cdots(\alpha+n)}{n!}$$

for all  $n \in \mathbb{Z}^+$ . Additionally,  $L_0^{\alpha}(0) = 1$  and we set  $L_n^{\alpha}(0) = 0$  when  $n \in \mathbb{Z}^-$ . The Vandermonde determinant will be denoted as

$$\Delta(a_1, a_2, \dots, a_r) = \prod_{i < j} (a_j - a_i).$$
(6.1)

Partitions will be defined via (4.12) and (4.13) and corresponding indices, e.g.  $n'_i$ , used here do not correspond to the starting general Maya diagram for solutions of the first kind, but to the indices of the associated canonical position Maya diagram (or conjugate canonical position when solutions of the second kind are discussed). Note that in canonical form,  $r_3 = r_4 = 0$  so there are no n' or m'-terms in the Maya diagram; and in the conjugate canonical form,  $r_1 = r_2 = 0$  so there are no n or m-terms in the Maya diagram.

Solutions of the first kind, after canceling some terms with the prefactor and using the derivative rules in (5.5) and (5.6), can be written as

$$\Omega_{\mu,\nu}[h^{\alpha}(x,\lambda)] = \begin{vmatrix} L_{n_{1}}^{\alpha}(x) & \dots & L_{m_{r_{2}}}^{\alpha}(-x) & M(-\lambda,\alpha+1,x) \\ (-1)L_{n_{1}-1}^{\alpha+1}(x) & \dots & L_{m_{r_{2}}}^{\alpha+1}(-x) & \frac{-\lambda}{\alpha+1}M(-\lambda+1,\alpha+2,x) \\ \vdots & \ddots & \vdots & \vdots \\ (-1)^{r}L_{n_{1}-r}^{\alpha+r}(x) & \dots & L_{m_{r_{2}}}^{\alpha+r}(-x) & \frac{(-\lambda)^{(r)}}{(\alpha+1)^{(r)}}M(-\lambda+r-1,\alpha+r,x) \end{vmatrix}$$

Recall that  $r = r(\mu) + r(\nu)$ . It may occur that  $n_i < r$  for some *i*, which would mean that some of the entries in the column are 0. For this reason, let s > 0 denote the smallest natural number such that  $n_i < r$  for all  $i \ge s$  and  $n_i \ge r$  for all i < s. Recall

that the sequence  $\{n_i\}_{i=1}^{r(\mu)}$  is associated with a Maya diagram and thus is assumed to be strictly decreasing so the Wronskian has distinct entries. Likewise, we let s' > 0 be the smallest natural number such that  $m_i < r$  for all  $j \ge s'$  and  $m_i \ge r$  for all j < s'.

The following result can be viewed as a generalization of [10, Lemma 5.1]. Recall that Assumptions 4.2 and 5.2 hold.

**Theorem 6.1.** Given partitions  $\mu$  and  $\nu$  defined via (4.12) and (4.13) and s, s' defined above, then,  $\Omega_{\mu,\nu}[h^{\alpha}(0,\lambda)]$  is equal to

$$\frac{\prod_{k=1}^{r+1} (\alpha+k)^{(r+1-k)} \prod_{i=1}^{s-1} (\alpha+r+1)^{(n_i-r)} \prod_{j=1}^{s'-1} (\alpha+r+1)^{(m_j-r)} \Delta(n_{\mu}, m_{\nu}, -\lambda)}{(\alpha+1)^{(r)} \prod_{i=s}^{r(\mu)} (\alpha+1+n_i)^{(r-n_i)} \prod_{j=s'}^{r(\nu)} (\alpha+1+m_j)^{(r-m_j)} \prod_{i=1}^{r(\mu)} n_i! \prod_{j=1}^{r(\nu)} m_j!},$$

where

$$\Delta(n_{\mu}, m_{\nu}, \lambda) = \Delta(-n_{r(\mu)}, \dots, -n_{2}, -n_{1}, \alpha + 1 + m_{1}, \alpha + 1 + m_{2}, \dots, \alpha + 1 + m_{r(\nu)}, -\lambda).$$

*Proof.* Separate the Wronskian  $\Omega_{\mu,\nu}[h^{\alpha}(0,\lambda)]$  into two submatrices, one containing columns given by seed functions indexed by  $\mu$  and the solution  $h^{\alpha}(0,\lambda)$ ; the other with columns given by seed functions indexed by  $\nu$ . Each submatrix has r + 1 rows. Splitting the Wronskian matrix  $\Omega_{\mu,\emptyset}[h^{\alpha}(0,\lambda)]$  into these submatrices allows us to conveniently perform necessary column operations to each submatrix and then recombine the two submatrices before performing row operations. The resulting Wronskian will be a Vandermonde determinant and allow for  $\Omega_{\mu,\nu}[h^{\alpha}(0,\lambda)]$  to be evaluated.

Begin by fixing s > 0 as above and considering the submatrix of the Wronskian with columns given by seed functions indexed by  $\mu$  and the solution  $h^{\alpha}(0, \lambda)$ . This matrix has  $r(\mu)$  columns and r + 1 rows. By pulling out factors of (-1), rearrange the columns so that the indices of the  $n_i$  are in ascending order. We find that this submatrix is equal to

$$\begin{pmatrix} \frac{(\alpha+1)^{(n_r(\mu))}}{n_r(\mu)!} & \dots & \frac{(\alpha+1)^{(n_s)}}{n_s!} & \dots & \frac{(\alpha+1)^{(n_1)}}{n_1!} & 1 \\ \vdots & & & & \\ (-1)^{n_r(\mu)+1} & \vdots & & & \\ 0 & \ddots & & \vdots & & \vdots & \\ & & (-1)^{n_s+1} & & & \\ \vdots & & & & \\ 0 & \dots & 0 & \dots & (-1)^r \frac{(\alpha+r+1)^{(n_1-r)}}{(n_1-r)!} & (-1)^r \frac{(-\lambda)^{(r)}}{(\alpha+1)^{(r)}} \end{pmatrix}$$

Using column operations, we can remove a factor of  $1/(\alpha + 1)^{(r)}$  from the last column, a factor  $(\alpha + r + 1)^{(n_i-r)}/(n_i)!$  from columns where i < s and a factor  $1/((\alpha + 1 + n_i)^{(r-n_i)}(n_i)!)$  from columns where  $i \ge s$ . This leaves the submatrix as

$$\begin{pmatrix} (\alpha+1)^{(r)} & \dots & (\alpha+1)^{(r)} & \dots & (\alpha+1)^{(r)} & (\alpha+1)^{(r)} \\ \vdots & & & & \\ (\alpha+1+n_{r(\mu)})^{(r-n_{r(\mu)})}(-n_{r(\mu)})! & \vdots & & \\ 0 & \ddots & & \vdots & \vdots \\ & & (\alpha+1+n_{s})^{(r-n_{s})}(-n_{s})! & & \\ \vdots & & 0 & \ddots & \\ & & & \vdots & \\ 0 & \dots & 0 & \dots & (-n_{1})^{(r)} & (-\lambda)^{(r)} \end{pmatrix}.$$

Of course, the factorial  $(-n_j)! = (-1)^{n_j} ((n_j)!)$  and Assumption 5.2 should be kept in mind here.

We now perform row operations that we will also apply to the other submatrix of the Wronskian. We remove a factor of  $(\alpha + k)^{(r+1-k)}$  from each row k where k = 1, ..., r + 1. Collecting the terms removed from this submatrix yields

$$\frac{\prod_{k=1}^{r+1} (\alpha + k)^{(r+1-k)} \prod_{i=1}^{s-1} (\alpha + r + 1)^{(n_i - r)}}{(\alpha + 1)^{(r)} \prod_{i=s}^{r(\mu)} (\alpha + 1 + n_i)^{(r-n_i)} \prod_{i=1}^{r(\mu)} n_i!} \\
\times \begin{pmatrix} 1 & \dots & 1 & 1 \\ \vdots & & \\ (n_{r(\mu)})! & & \\ 0 & \ddots & \vdots & \vdots \\ \vdots & & \\ 0 & \dots & (-n_1)^{(r)} & (-\lambda)^{(r)} \end{pmatrix}$$

Now, consider the second submatrix of the Wronskian with columns given by seed functions indexed by v and r + 1 rows. Set s' > 0 as above. Fortunately, there will be no zeros in the matrix, but the value s' will still play a role in removing terms. Calculations show that the evaluation of this submatrix at 0 is equal to

$$\begin{pmatrix} \frac{(\alpha+1)^{(m_1)}}{m_1!} & \frac{(\alpha+1)^{(m_2)}}{m_2!} & \cdots & \frac{(\alpha+1)^{(m_r(\nu))}}{m_{r(\nu)}!} \\ \frac{(\alpha+2)^{(m_1)}}{m_1!} & \frac{(\alpha+2)^{(m_2)}}{m_2!} & \cdots & \frac{(\alpha+2)^{(m_r(\nu))}}{m_{r(\nu)}!} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{(\alpha+r+1)^{(m_1)}}{m_1!} & \frac{(\alpha+r+1)^{(m_2)}}{m_2!} & \cdots & \frac{(\alpha+r+1)^{(m_r(\nu))}}{(m_{r(\nu)})!} \end{pmatrix}$$

Using column operations, we can remove a factor of  $(\alpha + r + 1)^{(m_j - r)}/m_j!$  from all columns where j < s' and a factor of  $1/((\alpha + 1 + m_j)^{(r-m_j)}(m_j)!)$  from remaining columns. The submatrix is then

$$\begin{pmatrix} (\alpha+1)^{(r)} \cdot 1 & \dots & (\alpha+1)^{(r)} \cdot 1 \\ (\alpha+2)^{(r-1)}(\alpha+1+m_1) & \dots & (\alpha+2)^{(r-1)}(\alpha+1+m_{r(\nu)}) \\ \vdots & \ddots & \vdots \\ 1 \cdot (\alpha+1+m_1)^{(r)} & \dots & 1 \cdot (\alpha+1+m_{r(\nu)})^{(r)} \end{pmatrix}.$$

We now apply the same row operations as we did for the other submatrix and remove a factor of  $(\alpha + k)^{(r+1-k)}$  from each row k = 1, ..., r + 1. Altogether, the second submatrix simplifies to

$$\frac{\prod_{k=1}^{r+1} (\alpha + k)^{(r+1-k)} \prod_{j=1}^{s'-1} (\alpha + r + 1)^{(m_j - r)}}{\prod_{j=s'}^{r(\nu)} (\alpha + 1 + m_j)^{(r-m_j)} \prod_{j=1}^{r(\nu)} m_j!} \times \begin{pmatrix} 1 & \dots & 1 \\ (\alpha + 1 + m_1) & \dots & (\alpha + 1 + m_{r(\nu)}) \\ \vdots & \ddots & \vdots \\ (\alpha + 1 + m_1)^{(r(\nu)-1)} & \dots & (\alpha + 1 + m_{r(\nu)})^{(r(\nu)-1)} \end{pmatrix}.$$

The solution of the first kind  $\Omega_{\mu,\nu}[h^{\alpha}(0, \lambda)]$  now merges the two submatrices analyzed above, with some columns rearranged. The result is a power of (-1) as an additional factor. Applying elementary row operations, the remaining determinant equals a partially simplified Vandermonde determinant with zeros in the bottom left corner.

A standard proof of the Vandermonde determinant formula in (6.1) begins with the last row and subtracts the previous row multiplied by  $-n_{r(\mu)}$ . Expanding about the first column and removing common factors yields a smaller Vandermonde matrix. The zeros in the bottom left corner of our Wronskian do not change this process and thus will not affect (6.1). Collecting all of the above terms thus yields the result.

The solution of the second kind can similarly be evaluated at 0 using the methods of Theorem 6.1. We define **s** to be the analog of *s* for the sequence  $\{n_i'\}_{i=1}^{r(\mu')}$  and **s'** the analog for the sequence  $\{m_i'\}_{i=1}^{r(\nu')}$ .

**Corollary 6.2.** Given partitions  $\mu'$  and  $\nu'$  defined via (4.12) and (4.13) and s, s' defined above, then,  $\Omega_{\mu',\nu'}[\tilde{h}^{\alpha}(0,\lambda)]$  is equal to

where

$$\mathfrak{M}_{1} := \prod_{i=1}^{r+1} (-\alpha + k)^{(r+1-k)} \prod_{j=1}^{\mathbf{s}'-1} (-\alpha + r + 1)^{(m'_{j}-r)} \prod_{i=1}^{\mathbf{s}-1} (-\alpha + r + 1)^{(n'_{i}-r)} \times \Delta(m'_{\nu'}, n'_{\mu'}, -\lambda - \alpha)$$

and

$$\mathfrak{D}_{2} := (1-\alpha)^{(r)} \prod_{j=s'}^{r(v')} (-\alpha + 1 + m'_{j})^{(r-m'_{j})} \prod_{i=s}^{r(\mu')} (-\alpha + 1 + n'_{i})^{(r-n'_{i})} \times \prod_{j=1}^{r(v')} m'_{j}! \prod_{i=1}^{r(\mu')} n'_{i}!,$$

and where

$$\Delta(m'_{\nu'}, n'_{\mu'}, -\lambda - \alpha) = \Delta(-m'_{r(\nu)'}, \dots, -m'_2, -m'_1, -\alpha + 1 + n'_1, -\alpha + 1 + n'_2, \dots, -\alpha + 1 + n'_{r(\mu')}, -\lambda - \alpha).$$

*Proof.* The solution of the second kind  $\Omega_{\mu',\nu'}[\tilde{h}^{\alpha}(x,\lambda)]$  by definition has a prefactor of  $x^{\alpha(r'+1)}$ . Removing a factor of  $x^{-\alpha}$  from each column cancels with this prefactor and yields a Wronskian with only functions of the types given in (4.1), indexed by  $\{m'_j\}_{j=1}^{r(\nu')}$ , and (4.2), indexed by  $\{n'_i\}_{i=1}^{r(\mu')}$ , with the function  $M(-\lambda - \alpha, 1 - \alpha, x)$  as the input for the last column. The partition indices have thus been changed to  $(\nu', \mu')$ . It is then possible to apply Theorem 6.1 to evaluate this Wronskian. Note the parameter for the seed functions is now  $-\alpha$ , as opposed to  $\alpha$  from Theorem 6.1. The result follows.

The proof of Theorem 6.1 can be modified to fit the simplified case where the generalized Laguerre polynomial is considered. This is essentially [10, Lemma 5.1] in the notation of this manuscript. Here, *s* and its analog must be slightly altered as there are r - 1 instead of *r* derivatives taken. Let s > 0 be the smallest natural number such that  $n_i < r - 1$  for all  $i \ge s$  and  $n_i \ge r - 1$  for all i < s. If such an *s* does not exist, set  $s = r(\mu) + 1$ . Define *s'*, **s** and **s'** analogously.

**Corollary 6.3.** Given partitions  $\mu$  and  $\nu$  defined via (4.12) and (4.13) and s, s' defined above, then

$$\Omega_{\mu,\nu}^{\alpha}(0) = \frac{\prod_{k=1}^{r} (\alpha+k)^{(r-k)} \prod_{i=1}^{s-1} (\alpha+r)^{(n_i-r+1)} \prod_{j=1}^{s'-1} (\alpha+r)^{(m_j-r+1)} \Delta(n_{\mu}, m_{\nu})}{\prod_{i=s}^{r(\mu)} (\alpha+1+n_i)^{(r-1-n_i)} \prod_{j=s'}^{r(\nu)} (\alpha+1+m_j)^{(r-1-m_j)} \prod_{i=1}^{r(\mu)} n_i! \prod_{j=1}^{r(\nu)} m_j!}.$$

*Proof.* It is necessary to slightly alter the column operations of Theorem 6.1 in order to account for the fact that the Wronskian only has r rows and columns. Hence, there are r - 1 derivatives taken and this is reflected in the altered rising factorials and terms in the numerators of the expressions. The results follow after these alterations.

#### 7. Weyl *m*-functions

Sections 5 and 6 have provided the tools necessary to obtain spectral information about self-adjoint extensions of exceptional Laguerre expressions given by two general Maya diagrams.

Let  $\hat{\ell}$  be a general Laguerre-type differential expression acting on the weighted space  $L^2[(0, \infty), W(x)dx]$  having maximal domain  $\mathcal{D}_{max}$ . Given fixed  $M_1$  and  $M_2$ , the partitions  $\mu$ ,  $\nu$  are determined by (4.12) and (4.13) after shifts to canonical and conjugate canonical forms, respectively. The values  $r(\mu)$ ,  $r(\nu)$  are then determined by (4.10) and (4.11). The weight function possesses a factor of  $e^{-x}x^{\alpha}$ , where  $\alpha = \alpha' + r(\mu) + r(\nu) = \alpha' + r$  is fixed due to Lemma 4.6. It is assumed that  $\alpha > -1$ so  $\hat{\ell}$  is in the limit-circle case at x = 0. The value  $\alpha' = \alpha - t_1 - t_2$  is determined by the shifts  $t_1$  and  $t_2$  of the partitions from Theorem 5.3. The weight function W(x)depends on a generalized Laguerre polynomial  $\Omega^{\alpha}_{M_1,M_2}(x)$ , which we identify with a valid factorization chain from Section 2.

**Remark 7.1.** The attentive reader may question why  $\alpha'$  is used in the definition of  $\alpha$  for the order of the underlying Hilbert space instead of  $\alpha - t'_1 - t'_2$ , which appears in Theorem 5.5. Is the order  $\alpha - t'_1 - t'_2$  even useful? Indeed, Theorem 5.3 shows that multiplication of the Wronskian by a certain factor can change the order of the parameter  $\alpha$  within the Wronskian, causing some ambiguity. The solution of the second kind in Theorem 5.5 could be shifted to canonical position, also yielding a parameter  $\alpha - t_1 - t_2$ . However, it is not possible to perform the evaluation in Corollary 6.2 with this setup. Hence, the order  $\alpha - t'_1 - t'_2$  is still necessary but not used here.

The Frobenius analysis of Section 4 for general Laguerre XOP expressions indicates that there are two linearly independent solutions at the regular singular point x = 0:  $y_1(x) = 1$  and  $y_2(x) = x^{-\alpha}$ . These two functions are in the maximal domain  $\mathcal{D}_{\text{max}}$  as they both belong to  $L^2[(0, \infty), \mathcal{W}(x)]$  and are solutions to the equation  $(\mathcal{L}_{\text{max}} - \lambda) f = 0$ . For the sake of normalization, we will use the functions

$$\tilde{y}_1(x) := -\frac{\Omega_{M_1,M_2}^{\alpha}(0)}{\alpha}$$
 and  $\tilde{y}_2(x) := \Omega_{M_1,M_2}^{\alpha}(0)x^{-\alpha}$ 

instead of the particular solutions  $y_1(x) = 1$  and  $y_2(x) = x^{-\alpha}$ , see also Remark 3.1. Note that  $\Omega^{\alpha}_{M_1,M_2}(x)$  is a polynomial with real coefficients so  $\Omega^{\alpha}_{M_1,M_2}(0)$  exists, and it was already assumed that  $\Omega^{\alpha}_{M_1,M_2}(0) \neq 0$ . Hence,  $\tilde{y}_1$  and  $\tilde{y}_2$  are well defined and non-zero but slightly different from the formulations found in Section 3.

The sesquilinear form associated to the expression  $\hat{\ell}$  is defined via (2.9). Let  $f, g \in \mathcal{D}_{max}$  and define

$$\Gamma_0 f = f^{[0]}(0) := [f, \tilde{y}_2]_{\hat{\ell}}(0) = \lim_{x \to 0^+} -\frac{\alpha f(x) + x f'(x)}{\Omega^{\alpha}_{M_1, M_2}(0)},$$
  
$$\Gamma_1 f = f^{[1]}(0) := [f, \tilde{y}_1]_{\hat{\ell}}(0) = \lim_{x \to 0^+} \frac{x^{\alpha+1}}{\alpha \Omega^{\alpha}_{M_1, M_2}(0)} f'(x).$$

The choice of the functions  $\tilde{y}_1$  and  $\tilde{y}_2$  immediately imply that  $[\tilde{y}_1, \tilde{y}_2]_{\hat{\ell}}(0) = 1$ . The functions  $\tilde{y}_1$  and  $\tilde{y}_2$  adhere to Definition 2.8. Therefore, Proposition 2.11 implies that  $\{\mathbb{C}, \Gamma_0, \Gamma_1\}$  is a boundary triple for  $\mathcal{D}_{max}$ . Explicitly, it is easy to calculate that

$$\langle \Gamma_1 f, \Gamma_0 g \rangle - \langle \Gamma_0 f, \Gamma_1 g \rangle = [f, g]_{\hat{\ell}}(0).$$

As in (2.11), this boundary triple naturally defines two self-adjoint extensions of the exceptional Laguerre operator  $\mathcal{L}_{\min}$ :  $\mathcal{L}_{\infty}$  and  $\mathcal{L}_{0}$ .

The solutions of the first and second kind can be modified to form a fundamental system for the equation  $(\mathcal{L}_{max} - \lambda)f = 0$ . Define

$$u_{1}(x,\lambda) := -\frac{\Omega_{M_{1},M_{2}}^{\alpha}(0)\Omega_{M_{1},M_{2}}[h^{\alpha}(x,\lambda)]}{\alpha\Omega_{M_{1},M_{2}}[h^{\alpha}(0,\lambda)]} = -\frac{\Omega_{M_{1},M_{2}}^{\alpha}(0)\Omega_{\mu,\nu}[h^{\alpha'}(x,\lambda+t_{1})]}{\alpha\Omega_{\mu,\nu}[h^{\alpha'}(0,\lambda+t_{1})]},$$
(7.1)

and

$$u_{2}(x,\lambda) := -\frac{\Omega_{M_{1},M_{2}}^{\alpha}(0)x^{-\alpha}\Omega_{M_{1},M_{2}}[\bar{h}^{\alpha}(x,\lambda)]}{\Omega_{M_{1},M_{2}}[\bar{h}^{\alpha}(0,\lambda)]} = -\frac{\Omega_{M_{1},M_{2}}^{\alpha}(0)x^{-\alpha}\Omega_{\mu',\nu'}[\bar{h}^{\alpha-t_{1}'-t_{2}'}(x,\lambda+t_{1}')]}{\Omega_{\mu',\nu'}[\bar{h}^{\alpha-t_{1}'-t_{2}'}(0,\lambda+t_{1}')]},$$
(7.2)

where each equality follows from shifting the Maya diagrams in both the numerator and denominator, with the constants from Theorems 5.3 and 5.5 canceling out. Notice that if the prefactor of  $\Omega_{M_1,M_2}[\tilde{h}^{\alpha}(x,\lambda)]$  is ignored, then the result is guaranteed to be in  $L^2[(0,\infty), W(x)dx]$  by the intertwining property of the Darboux transform. The multiplication by  $x^{-\alpha}$  above ensures this remains the same as the case where the standard prefactor is included.

Assumption 7.2. We assume that the  $x^{-\alpha}$  term is multiplied to the solution of the second kind and therefore defines the behavior of  $u_2(x, \lambda)$  near x = 0.

**Remark 7.3.** Multiplication by the  $x^{-\alpha}$  term discussed in Assumption 7.2 may instead be applied to the solution of the first kind, but this would change our notation significantly. In particular, this alternative convention can be translated into what we use here by changing the Maya diagrams and treating the function of the second kind as the principal solution, which can be written as a power series by the notation of Section 4. The alternate convention has been used to state examples like the Type III Laguerre XOPs described in [26], where the eigenvalues generated by the polynomials can all be of the form  $n - \alpha$  for  $n \in \mathbb{N}_0$ .

**Corollary 7.4.** The fundamental system  $(u_1(x, \lambda), u_2(x, \lambda))$  for the equation

$$(\mathcal{L}_{\max} - \lambda)f = 0,$$

given in (7.1) and (7.2), satisfies the initial conditions

$$\begin{pmatrix} u_1^{[0]}(0,\lambda) & u_2^{[0]}(0,\lambda) \\ u_1^{[1]}(0,\lambda) & u_2^{[1]}(0,\lambda) \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

*Proof.* Recall that both  $\Omega_{M_1,M_2}[h^{\alpha}(x,\lambda)]$  and  $\Omega_{M_1,M_2}[\tilde{h}^{\alpha}(x,\lambda)]$  are given by power series and only the lowest order terms make an impact on the quasi-derivatives. The identities are straightforward to verify by plugging  $u_1(x,\lambda)$  and  $u_2(x,\lambda)$  into the quasi-derivatives.

Let  $\chi(x, \lambda)$  be a general deficiency element formally satisfying  $(\ell - \lambda)\chi(x, \lambda) = 0$ . Then,  $\chi$  can be written as a linear combination of the solutions which compose the fundamental system. In the case of the classical Laguerre operator,  $\chi$  is the Tricomi confluent hypergeometric function which can be expressed via [5, (13.2.42)] as

$$U(-\lambda, \alpha + 1, x) = \frac{\Gamma(-\alpha)}{\Gamma(-\lambda - \alpha)} M(-\lambda, \alpha + 1, x) + \frac{\Gamma(\alpha)}{\Gamma(-\lambda)} x^{-\alpha} M(-\lambda - \alpha, 1 - \alpha, x).$$
(7.3)

The intertwining property of the Darboux transform then allows us to define a general deficiency element of the exceptional differential expression.

**Proposition 7.5.** Let  $\chi_{\hat{\ell}}(x,\lambda)$  be a general deficiency element of  $\mathcal{L}_{\max}$  such that

$$(\mathcal{L}_{\max} - \lambda)\chi_{\hat{\ell}}(x,\lambda) = 0.$$

*Then,*  $\chi_{\hat{\ell}}(x,\lambda)$  *can be written as* 

$$\chi_{\hat{\ell}}(x,\lambda) = \mathfrak{C}u_1(x,\lambda) + \mathfrak{D}u_2(x,\lambda),$$

where

$$\begin{split} \mathfrak{C} &= -\frac{\alpha C \, \Gamma(-\alpha) \Omega_{\mu,\nu} [h^{\alpha'}(0,\lambda+t_1)]}{\Gamma(-\lambda-\alpha) \Omega^{\alpha}_{M_1,M_2}(0)},\\ \mathfrak{D} &= -\frac{D \, \Gamma(\alpha) \Omega_{\mu',\nu'} [\tilde{h}^{\alpha-t_1'-t_2'}(0,\lambda+t_1')]}{\Gamma(-\lambda) \Omega^{\alpha}_{M_1,M_2}(0)} \end{split}$$

In particular, we have that  $\chi_{\hat{\ell}}^{[0]}(x,\lambda) = \mathfrak{C}$  and  $\chi_{\hat{\ell}}^{[1]}(x,\lambda) = \mathfrak{D}$ .

*Proof.* Let  $A_{M_1,M_2}$  denote the multi-step Darboux transform having seed functions indexed by  $M_1$  and  $M_2$ . The Wronskian is linear within its entries because differentiation and multiplication are. Applying  $A_{M_1,M_2}$  to (7.3) then yields

$$\begin{split} \chi_{\hat{\ell}}(x,\lambda) &= A_{M_1,M_2}[\chi(x,\lambda)] \\ &= A_{M_1,M_2} \Big[ \frac{\Gamma(-\alpha)}{\Gamma(-\lambda-\alpha)} h^{\alpha}(x,\lambda) + \frac{\Gamma(\alpha)}{\Gamma(-\lambda)} \tilde{h}^{\alpha}(x,\lambda) \Big] \\ &= \frac{\Gamma(-\alpha)}{\Gamma(-\lambda-\alpha)} \Omega_{M_1,M_2}[h^{\alpha}(x,\lambda)] + \frac{\Gamma(\alpha)}{\Gamma(-\lambda)} x^{-\alpha} \Omega_{M_1,M_2}[\tilde{h}^{\alpha}(x,\lambda)]. \end{split}$$

Recall the constants stemming from Theorems 5.3 and 5.5, and stated in (5.4) and (5.15), are denoted by *C* and *D*. Then,

$$\chi_{\hat{\ell}}(x,\lambda) = C \frac{\Gamma(-\alpha)}{\Gamma(-\lambda-\alpha)} \Omega_{\mu,\nu} [h^{\alpha'}(x,\lambda+t_1)] + D \frac{\Gamma(\alpha)}{\Gamma(-\lambda)} x^{-\alpha} \Omega_{\mu',\nu'} [\tilde{h}^{\alpha-t_1'-t_2'}(x,\lambda+t_1')].$$

The formulation of  $\chi_{\hat{\ell}}(x,\lambda)$  from the statement of the proposition follows. The initial conditions of Corollary 7.4 yield the values of the quasi-derivatives of this deficiency element and the result follows.

The quasi-derivatives of all functions  $f \in \mathcal{D}_{\text{max}}$  are guaranteed to exist and be finite by Theorem 2.4. However, when these quasi-derivatives are applied to  $\chi_{\hat{\ell}}$  in Proposition 7.5 there are many terms in the denominator which may seem to cause singularities for certain values of  $\lambda$ . These singularities may cause eigenvalues of self-adjoint extensions, so it is worth investigating the quasi-derivatives from Proposition 7.5 further.

**Proposition 7.6.** The quasi-derivatives  $\chi_{\hat{\ell}}^{[0]}(x,\lambda) = \mathfrak{C}$  and  $\chi_{\hat{\ell}}^{[1]}(x,\lambda) = \mathfrak{D}$  from *Proposition 7.5 are well defined and finite for all values of x and*  $\lambda$ *.* 

*Proof.* We prove the proposition for  $\chi_{\hat{\ell}}^{[0]}(x,\lambda) = \mathfrak{C}$ ; the proof for  $\chi_{\hat{\ell}}^{[1]}(x,\lambda) = \mathfrak{D}$  follows analogously. Note that some care should be taken to distinguish a Maya diagram

entry to the left of the origin while in conjugate canonical form, k, with its associated partition entry, -k - 1.

Begin by noticing the terms in the denominator of  $\mathfrak{C}$ ;  $\Gamma(-\lambda - \alpha)$  is analytic and  $\Omega^{\alpha}_{M_1,M_2}(0)$  has no zeros in the interval  $[0,\infty)$  by definition. Additionally, the terms from  $\Omega_{\mu,\nu}[h^{\alpha'}(0,\lambda+t_1)]$  that involve  $\lambda$  are all in the numerator. Hence, the only singularities possible in  $\mathfrak{C}$  are terms that may come from the denominator of C.

Theorem 5.3 says that terms involving  $\lambda$  may appear in the denominator of C only if  $t_1 > 0$  and  $t_2 > 0$ . Terms arising from Theorem 5.3 when  $t_1 > 0$  are  $\prod_{k_1} (-\lambda - k_1 - 1)$  for  $k_1 \in \{0, \ldots, t_1 - 1\}$  such that  $k_1 \notin \{n'_j\}_{j=1}^{r_4}$ . Let k be a general index that meets the criteria for  $k_1$  so a term  $-\lambda - k - 1$  is included in  $C_1$ .

The partition entry associated with the index k in  $\mu$  after  $M_1$  is shifted to canonical position is  $-k - 1 + t_1$ . The Vandermonde determinant in  $\Omega_{\mu,\nu}[h^{\alpha'}(0, \lambda + t_1)]$  will produce a term associated with this partition index which is

$$-\lambda - t_1 - (-(-k - 1 + t_1)) = -\lambda - k - 1.$$

Hence, this factor in the numerator cancels the contribution from the denominator of  $C_1$ .

Now, consider terms that may come from the denominator of  $C_2$  when  $t_2 > 0$ :  $\prod_{k_2} (\lambda + \alpha - k_2)$  for  $k_2 \in \{0, \dots, t_2 - 1\}$  such that  $k_2 \notin \{m'_j\}_{j=1}^{r_3}$ . Let  $\tilde{k}$  be a general index that meets the criteria for  $k_2$  so that a term  $\lambda + \alpha - \tilde{k}$  is included in  $C_2$ .

The partition entry associated with the index  $\tilde{k}$  in  $\nu$  after  $M_2$  is shifted to canonical position is  $-\tilde{k} - 1 + t_2$ . The Vandermonde determinant, however, again produces a term associated with this partition index:

$$-\lambda - t_1 - (\alpha - t_1 - t_2 + 1 - \tilde{k} - 1 + t_2) = -\lambda - \alpha - \tilde{k}.$$

This numerator factor also cancels the contribution from the denominator of  $C_2$ . We conclude that there can be no terms in the denominator of C which include  $\lambda$  and are not canceled out by identical terms from  $\Omega_{\mu,\nu}[h^{\alpha'}(0, \lambda + t_1)]$ . The quasi-derivative  $\chi_{\hat{i}}^{[0]}(x, \lambda)$  is thus well defined and finite.

Let  $\tau \in \mathbb{R} \cup \{\infty\}$  and  $\mathcal{L}_{\tau}$  refer to the self-adjoint operator acting via (2.12). The values of the quasi-derivatives of a general deficiency element allows for the definition of a Weyl *m*-function for every self-adjoint extension  $\mathcal{L}_{\tau}$ .

**Corollary 7.7.** For  $\lambda \in \rho(\mathcal{L}_{\infty})$ ,

$$M_{\infty}(\lambda) = \frac{\chi_{\hat{\ell}}^{[1]}(0,\lambda)}{\chi_{\hat{\ell}}^{[0]}(0,\lambda)} = \frac{\mathfrak{D}}{\mathfrak{C}}$$

$$= \frac{D\Gamma(-\lambda-\alpha)\Gamma(\alpha)\Omega_{\mu',\nu'}[\tilde{h}^{\alpha-t_1'-t_2'}(0,\lambda+t_1')]}{\alpha C\Gamma(-\lambda)\Gamma(-\alpha)\Omega_{\mu,\nu}[h^{\alpha'}(0,\lambda+t_1)]}.$$
(7.4)

For  $\lambda \in \rho(\mathcal{L}_{\infty}) \cup \rho(\mathcal{L}_{0})$ ,

$$M_{0}(\lambda) = -\frac{\chi_{\hat{\ell}}^{[0]}(0,\lambda)}{\chi_{\hat{\ell}}^{[1]}(0,\lambda)} = -\frac{\mathfrak{C}}{\mathfrak{D}}$$

$$= -\frac{\alpha C \Gamma(-\lambda) \Gamma(-\alpha) \Omega_{\mu,\nu} [h^{\alpha'}(0,\lambda+t_{1})]}{D \Gamma(-\lambda-\alpha) \Gamma(\alpha) \Omega_{\mu',\nu'} [\tilde{h}^{\alpha-t_{1}'-t_{2}'}(0,\lambda+t_{1}')]}.$$
(7.5)

For  $\lambda \in \rho(\mathcal{L}_{\infty}) \cup \rho(\mathcal{L}_{\tau})$ ,

$$M_{\tau}(\lambda) = \frac{1 + \tau M_{\infty}(\lambda)}{\tau - M_{\infty}(\lambda)} = \frac{\mathfrak{N}_2}{\mathfrak{D}_2},\tag{7.6}$$

where

$$\mathfrak{M}_{2} := \boldsymbol{\alpha} C \Gamma(-\lambda) \Gamma(-\alpha) \Omega_{\mu,\nu} [h^{\alpha'}(0, \alpha + t_{1})] + \tau D \Gamma(-\lambda - \alpha) \Gamma(\alpha) \Omega_{\mu',\nu'} [\tilde{h}^{\alpha - t_{1}' - t_{2}'}(0, \alpha + t_{1}')]$$

and

$$\mathfrak{D}_{2} := \tau \alpha C \Gamma(-\lambda) \Gamma(-\alpha) \Omega_{\mu,\nu} [h^{\alpha'}(0,\alpha+t_{1})] - D \Gamma(-\lambda-\alpha) \Gamma(\alpha) \Omega_{\mu',\nu'} [\tilde{h}^{\alpha-t_{1}'-t_{2}'}(0,\alpha+t_{1}')]$$

*Proof.* An application of Proposition 2.11 to the boundary triple { $\mathbb{C}$ ,  $\Gamma_0$ ,  $\Gamma_1$ } and the deficiency element  $\chi_{\hat{\ell}}(x, \lambda)$  yields (7.4). Likewise, the identity

$$M_0(\lambda) = -1 \backslash M_\infty(\lambda)$$

yields (7.5). The *m*-function of  $\mathcal{L}_{\tau}$  is obtained from (2.15).

The spectrum of the self-adjoint operator  $\mathcal{L}_{\infty}$  consists of the points that are poles of  $M_{\infty}$ . The function  $\Gamma(-\lambda - \alpha)$  is responsible for the eigenvalues  $\lambda = n - \alpha$ , for  $n \in \mathbb{N}_0$ . These are exactly the eigenvalues of the extension  $\mathcal{L}_{\infty}$  for the classical Laguerre operator. In that case, both partitions are empty so *C*, *D* and the two evaluations at 0 are all equal to 1.

The self-adjoint extension  $\mathcal{L}_0$  will possess the exceptional solution of the first kind and therefore have the family of exceptional polynomial eigenfunctions in its domain by Lemma 4.6. Eigenvalues of  $\mathcal{L}_0$  are simply the poles of  $M_0$ . The corresponding extension of the classical Laguerre expression contains the Laguerre polynomials and has eigenvalues  $\lambda = n$  for  $n \in \mathbb{N}_0$ . The function  $\Gamma(-\lambda)$  is responsible for these eigenvalues in (7.5).

Comparing these expressions to the classical Laguerre operator, we conclude that eigenvalues  $\lambda$  not stemming from  $\Gamma(-\lambda)$  or  $\Gamma(-\lambda - \alpha)$  are manufactured by the Darboux transform. The evaluations

$$\Omega_{\mu,\nu}[h^{\alpha'}(0,\lambda+t_1)]$$
 and  $\Omega_{\mu',\nu'}[\tilde{h}^{\alpha-t_1'-t_2'}(0,\lambda+t_1')]$ 

. .

do not depend on  $\lambda$  except for the Vandermonde determinants  $\Delta(n_{\mu}, m_{\nu}, -\lambda)$  and  $\Delta(m'_{\nu'}, n'_{\mu'}, -\lambda - \alpha)$ , respectively in the numerators. The values *C* and *D* can also have factors of  $\lambda$  in their numerators or denominators depending on whether the shifts are positive or negative.

The only true knowledge about how these terms combine to create eigenvalues in the Weyl *m*-functions from Corollary 7.7 thus comes from Proposition 7.6: terms in the denominators of *C* and *D* actually cancel with contributions from the numerator of Vandermonde determinants. The similar question of whether terms in the numerators of *C* and *D* can cancel with terms coming from the opposite Vandermonde determinant cannot be answered so easily. Technically this is possible, but the terms involved are not directly related in any way. For example, if  $t_1 < 0$ , then  $C_1$  will include terms from  $(-\lambda)^{(|t_1|)}$  which, in some cases, may cancel with terms  $-\lambda - t'_1 - 1 - n'_j$  from  $\Omega_{\mu',\nu'}[\tilde{h}^{\alpha-t'_1-t'_2}(0, \alpha + t'_1)]$ , where  $n'_j$  is from a general Maya diagram and not associated with the partition after the shift to conjugate canonical form.

Fortunately, because the Weyl *m*-functions of Corollary 7.7 and all the included formulas are explicit, it is always possible to directly calculate the eigenvalues and such an analysis is not necessary.

**Remark 7.8.** The Friedrichs extension of  $\mathcal{L}_{\min}$  is identified as the self-adjoint extension whose boundary condition is given by  $[f, u]_{\hat{\ell}}(0) = 0$  for a principal solution u by [28]. The two linearly independent 'particular solutions' (near 0) of  $\hat{\ell}$  used in our boundary triple were  $\tilde{y}_1$  and  $\tilde{y}_2$ . In practice, which solution is principal thus depends on the parameter  $\alpha$ . Hence, one of  $\mathcal{L}_0$  and  $\mathcal{L}_\infty$  will be the Friedrichs extension and, if  $\hat{\ell} > 0$ , the other will be the Krein–von Neumann extension, see e.g. [3, (5.4.26)], but identifying which is not always straightforward.

The easiest way to distinguish these important extensions is to observe that the Friedrichs extension will have the greatest first eigenvalue because it has the greatest lower bound among all extensions, see, e.g., [3, Proposition 5.3.6]. Any dependence of the lowest eigenvalues on  $\alpha$  may then result in the Friedrichs extension switching from  $\mathcal{L}_0$  to  $\mathcal{L}_{\infty}$ , or vice versa.

**Remark 7.9.** Section 3 analyzes the Type I Laguerre exceptional expression  $\hat{\ell}_m^{I,\alpha}$  and finds that  $\sigma(\mathcal{L}_0) = n$  and  $\sigma(\mathcal{L}_\infty) = (-m - \alpha) \cup (n + 1 - \alpha)$  for  $n \in \mathbb{N}_0$  and fixed  $m \in \mathbb{N}$ . These eigenvalues can also be recovered from (7.4) and (7.5). Note it is possible to match the *m*-functions from these two methods, but we focus only on the spectra here for the sake of simplicity.

The Darboux transform in this case is identified by [6, Section 5.2] to consist of seed functions indexed by  $\mu = \emptyset$ ,  $\nu = (m)$  and starting parameter  $\alpha - 1$ . Hence,  $t_1 = t_2 = t'_1 = 0$ ,  $t'_2 = -m - 1$  and  $\alpha = (\alpha - 1) + 1 = \alpha$ . Theorem 5.5 sets  $C_1 = C_2 = 0$  and Theorem 6.1 finds that  $\Omega_{\mu,\nu}[h^{\alpha-1}(0,\lambda)]$  has a factor of  $-\lambda - (\alpha - 1) - 1 - m =$ 



Figure 4. Original Maya diagrams.

 $-\lambda - \alpha - m$  in the numerator stemming from the Vandermonde determinant. The only dependence on  $\lambda$  in  $\mathfrak{C}$  is therefore a factor of  $(-\lambda - \alpha - m)\Gamma(-\lambda)$ .

After shifting to conjugate canonical position,  $M_2 = (m - 1, ..., 0)$ . Theorem 5.5 sets  $D_1 = 0$  and  $D_2$  has a factor of  $\prod_{k'} (-\lambda - (\alpha - 1) - 1 - k')$  with k' = 0, ..., m - 1 in the denominator. Corollary 6.2 also generates a factor, for k = 1, ..., m,

$$\prod_{k}(-\lambda - [(\alpha - 1) - (-m - 1)] + k) = \prod_{k}(-\lambda - \alpha - m + k),$$

in the numerator of  $\Omega_{\mu',\nu'}[\tilde{h}^{\alpha-1}(0,\lambda)]$ . These contributions cancel and the only dependence on  $\lambda$  in  $\mathfrak{D}$  is therefore a factor of  $\Gamma(-\lambda - \alpha + 1)$ .

Generating the expressions for  $M_0$  and  $M_\infty$  via (7.4) and (7.5) allows for easy identification of the eigenvalues, which agree with the conclusions of Section 3.

**Remark 7.10.** Perturbation theory says that the spectrum of  $\mathcal{L}_0$  will smoothly shift to the spectrum of  $\mathcal{L}_\infty$  as the parameter  $\tau$  increases and eigenvalues of all self-adjoint extensions will be simple, see, e.g., [7, 38, 39].

#### 8. Example

To illustrate the results of Theorem 5.3, we present the following example. Consider Maya diagrams  $M_1 = (\emptyset \mid 3, 2)$  and  $M_2 = (1, 0 \mid \emptyset)$ , illustrated in Figure 4.

From  $M_1$  and  $M_2$ , it follows that n = (3, 2),  $m = \emptyset$ ,  $n' = \emptyset$ , and m' = (1, 0)as well as  $r_1 = r_3 = 2$  and  $r_2 = r_4 = 0$ . Observe that  $M_1$  is already in canonical form; thus  $t_1 = 0$ . To put  $M_2$  into canonical form, a shift of  $t_2 = 2$  is required. Thus,  $t_1(M_1) = (\emptyset \mid 3, 2)$  and  $t_2(M_2) = (\emptyset \mid \emptyset)$ . The visual representation for the canonical form of  $M_1$  and  $M_2$  may be seen in Figure 5, where all boxes to the left of the pictured subdiagram are filled and those to the right are empty.



Figure 5. Canonical Maya diagrams.

Using (4.1) and (4.2) in (4.14) yields

$$\Omega_{M_1,M_2}[h^{\alpha}(x,\lambda)] = x^{2(\alpha+3)} \operatorname{Wr}[L_3^{\alpha}(x), L_2^{\alpha}(x), x^{-\alpha}L_1^{-\alpha}(x), x^{-\alpha}L_0^{-\alpha}(x), M(-\lambda,\alpha+1,x)].$$
(8.1)

Following the procedure outlined in the proof of Theorem 5.3, equation (5.11) of Lemma 5.4 will be applied twice to shift  $M_1$  and  $M_2$  into  $t_1(M_1) = \mu$  and  $t_2(M_2) = \nu$ . To demonstrate how Theorem 5.3 works, we illustrate the first application of the shift on  $M_2$ .

Beginning with (8.1), we first factor out  $x^{-\alpha}$  from each column

$$\Omega_{M_1,M_2}[h^{\alpha}(x,\lambda)] = x^{2(\alpha+3)}x^{-5\alpha}\operatorname{Wr}[x^{\alpha}L_3^{\alpha}(x), x^{\alpha}L_2^{\alpha}(x), L_1^{-\alpha}(x), 1, x^{\alpha}M(-\lambda, \alpha+1, x)].$$

To compute the Wronskian, we expand along the first row, fourth column entry, using appropriate derivative rules

$$\Omega_{M_1,M_2}[h^{\alpha}(x,\lambda)] = x^{2(\alpha+3)} x^{-5\alpha} \operatorname{Wr}[(3+\alpha)x^{\alpha-1}L_3^{\alpha-1}(x), (2+\alpha)x^{\alpha-1}L_2^{\alpha-1}(x), -L_0^{-\alpha+1}(x), \alpha x^{\alpha-1}M(-\lambda,\alpha,x)].$$

Factoring out constants and  $x^{\alpha-1}$  from each factor in the Wronskian yields

$$\Omega_{M_1,M_2}[h^{\alpha}(x,\lambda)] = -\alpha(\alpha+2)(\alpha+3)x^{\alpha+2} \times \operatorname{Wr}[L_3^{\alpha-1}(x), L_2^{\alpha-1}(x), x^{-\alpha+1}L_0^{-\alpha+1}(x), M(-\lambda,\alpha,x)] = -\alpha(\alpha+2)(\alpha+3)\Omega_{\tilde{M}_1,\tilde{M}_2}[h^{\alpha-1}(x,\lambda)],$$

where  $\tilde{M}_1 = M_1 = (\emptyset \mid 3, 2)$  and  $\tilde{M}_2 = M_2 + 1 = (0 \mid \emptyset)$ . Observe that just as (5.11) states, the parameter  $\alpha$  has been shifted down by one while  $\lambda$  does not change. One

additional application of (5.11) gives

$$\begin{aligned} \Omega_{M_1,M_2}[h^{\alpha}(x,\lambda)] &= -\alpha(\alpha+2)(\alpha+3)x^{-3\alpha+6} \\ &\times \operatorname{Wr}[x^{\alpha-1}L_3^{\alpha-1}(x), x^{\alpha-1}L_2^{\alpha-1}(x), 1, x^{\alpha-1}M(-\lambda,\alpha,x)] \\ &= -(\alpha-1)\alpha(\alpha+1)(\alpha+2)^2(\alpha+3) \\ &\times \operatorname{Wr}[L_3^{\alpha-2}(x), L_2^{\alpha-2}(x), 1, x^{\alpha-1}M(-\lambda,\alpha-1,x)] \\ &= C_1C_2C_3\Omega_{\mu,\nu}^{\alpha-2}(x,\lambda), \end{aligned}$$

. .

where  $C_1 = 1$ ,  $C_2 = \alpha(\alpha - 1)$ , and  $C_3 = (\alpha + 1)(\alpha + 2)^2(\alpha + 3)$ . Keep in mind that the signs of coefficients  $C_1$ ,  $C_2$ , and  $C_3$  are suppressed. This result is consistent with that guaranteed by Theorem 5.3.

Now, we aim to adjust  $M_1$  and  $M_2$  into conjugate canonical form, see Figure 6 below. Observe that  $M_1$  must be shifted left by four units; thus  $t'_1 = -4$ . To put  $M_2$ into conjugate canonical form, a shift of  $t'_2 = 2$  is required. Note that in this example,  $M_2$  has the same canonical and conjugate canonical form. Thus,  $t'_1(M_1) = (3, 2 \mid \emptyset)$  and  $t'_2(M_2) = (\emptyset \mid \emptyset)$ . Following the procedure outlined in Theorem 5.5, equation (5.17) of Lemma 5.6 is applied twice to shift  $M_2$  into conjugate canonical form, and the inverse of (5.16) is applied twice to  $M_1$ , followed by two applications of (5.16) to shift  $M_1$  into conjugate canonical form. Note that there is some freedom as to how the shifts are applied. For example,  $M_1$  may be put into canonical form before  $M_2$ ; however, the order of the shifts applied to  $M_1$  must be done in the prescribed order.



Figure 6. Conjugate canonical Maya diagrams.

Consequently, (4.15) may be written, up to a change of sign, as

$$\Omega_{M_1,M_2}[\tilde{h}^{\alpha}(x,\lambda)] = D_1 D_2 D_3 \Omega_{\mu',\nu'}[\tilde{h}^{\alpha+2}(x,\lambda-4)],$$

where

$$D_1 = \frac{\alpha(\alpha+1)(\alpha+2)(\alpha+3)}{\lambda(\lambda-1)}, \quad D_2 = \frac{(\lambda+\alpha)(\lambda+\alpha+1)}{(\alpha+2)(\alpha+3)},$$

and  $D_3 = (\alpha + 1)(\alpha + 2)^2(\alpha + 3)$ .

Now, we turn our attention to calculating the value of  $\Omega_{M_1,M_2}[h^{\alpha}(0,\lambda)]$  and  $\Omega_{M_1,M_2}[\tilde{h}^{\alpha}(0,\lambda)]$ , which requires the use of Theorem 6.1 and Corollary 6.2 respectively. For the shift to canonical position,  $\mu = (2,2)$  and  $\nu = \emptyset$ ;  $r(\mu) = 2$  and  $r(\nu) = 0$ ; and s = 3 and s' = 1. Note that  $\mu$  is even, satisfying Assumption 4.2. Applying Theorem 6.1 gives

$$\Omega_{\mu,\nu}[h^{\alpha-2}(0,\lambda)] = \frac{\alpha(\alpha+1)(3-\lambda)(2-\lambda)}{12}$$

For the shift to conjugate canonical position,  $\mu' = (2, 2)$  and  $\nu' = \emptyset$ ;  $r(\mu') = 2$  and  $r(\nu') = 0$ ; and s = 3 and s' = 1. Applying Corollary 6.2 gives

$$\Omega_{\mu',\nu'}[\tilde{h}^{\alpha+2}(0,\lambda-4)] = \frac{(\alpha-1)\alpha(\lambda-1)\lambda}{12}$$

The value  $\alpha = \alpha - 0 - 2 + 2 + 0 = \alpha$  satisfies the requirement that  $\alpha > -1$  because  $\alpha > -1$ . Hence, the exceptional Laguerre differential expression  $\hat{\ell}$  is limit-circle at x = 0 and limit-point at  $x = \infty$ . Now, it is possible to explicitly write out  $\mathfrak{C}$  and  $\mathfrak{D}$  as

$$\mathfrak{C} = \frac{(\alpha-1)\alpha^3(\alpha+1)^2(\alpha+2)^2(\alpha+3)(3-\lambda)(2-\lambda)\Gamma(-\alpha)}{12\Gamma(-\lambda-\alpha)[\Omega^{\alpha}_{M_1,M_2}(0)]^2}$$

and

$$\mathfrak{D} = \frac{(\alpha - 1)\alpha^2(\alpha + 1)^2(\alpha + 2)^2(\alpha + 3)(\lambda + \alpha)(\lambda + \alpha + 1)\Gamma(\alpha)}{12\Gamma(-\lambda)[\Omega^{\alpha}_{M_1,M_2}(0)]^2}.$$

The quotient of D and C yields

$$M_{\infty}(\lambda) = \frac{(\lambda + \alpha)(\lambda + \alpha + 1)\Gamma(\alpha)\Gamma(-\lambda - \alpha)}{\alpha(3 - \lambda)(2 - \lambda)\Gamma(-\alpha)\Gamma(-\lambda)}.$$
(8.2)

To find the spectrum of the self-adjoint operator  $\mathcal{L}_{\infty}$ , consider where (8.2) has poles. The poles occur for eigenvalues  $\lambda = n - \alpha$  for  $n \in \mathbb{N}_0$  as well as  $\lambda = 2$  and  $\lambda = 3$ . However, the pole at  $\lambda = -\alpha$  is canceled by the factor of  $\lambda + \alpha$  in the numerator. Thus,  $\sigma(\mathcal{L}_{\infty}) = \{n - \alpha\}_{n \in \mathbb{N}} \cup \{2, 3\}.$ 

Considering the reciprocal of  $M_{\infty}$ , we get

$$M_0(\lambda) = \frac{\alpha(3-\lambda)(2-\lambda)\Gamma(-\alpha)\Gamma(-\lambda)}{(\lambda+\alpha)(\lambda+\alpha+1)\Gamma(\alpha)\Gamma(-\lambda-\alpha)}$$

which has poles at  $\lambda = n$  for all  $n \in \mathbb{N}_0$  as well as  $\lambda = -\alpha$  and  $\lambda = -\alpha - 1$ . The poles at  $\lambda = 2, 3$  are canceled by factors in the numerator. Thus,  $\sigma(\mathcal{L}_0) = \{n\}_{n \in \mathbb{N}_0 / \{2,3\}} \cup \{-\alpha, -\alpha - 1\}.$ 

**Remark 8.1.** The removal of the eigenvalues  $\lambda = 2, 3$  from  $\sigma(\mathcal{L}_0)$  and  $\lambda = -\alpha$  from  $\sigma(\mathcal{L}_\infty)$  is crucial, otherwise  $\sigma(\mathcal{L}_0) \cap \sigma(\mathcal{L}_\infty)$  is nontrivial. Assume some  $\lambda_0 \in \mathbb{R}$  belongs to  $\sigma(\mathcal{L}_0) \cap \sigma(\mathcal{L}_\infty)$ . Then the spectral measures of the two extensions are not mutually singular, violating the Aronszajn–Donoghue theorem for rank-one perturbations, see e.g. [1,38]

Furthermore, Remark 7.8 identifies that  $\mathcal{L}_{\infty}$  is the Friedrichs extension for all values of  $\alpha > -1$ , as comparing the first eigenvalues shows  $1 - \alpha > -1 - \alpha$ .

#### 9. Conclusions

Exceptional Laguerre-type symmetric expressions  $\hat{\ell}$ , which have XOPs as eigenfunctions for a self-adjoint extension, may be derived from the classical Laguerre symmetric operator with parameter  $\alpha$  by applying Darboux transforms. These are written as Wronskians of seed functions that meet Assumption 4.2 and the seed functions are described via the Maya diagrams  $M_1$  and  $M_2$  or, in special situations, as partitions  $\mu$ ,  $\nu$ ,  $\mu'$  or  $\nu'$ .

The expression  $\hat{\ell}$  acts on functions in the Hilbert space  $L^2[(0, \infty), W(x)]$  that satisfy standard differentiability criteria. The weight function

$$\mathcal{W}(x) = \frac{x^{\alpha} e^{-x}}{[\Omega_{\mu,\nu}^{\alpha'}(x)]^2} \quad \text{for } x > 0,$$

where  $\alpha' = \alpha - t_1 - t_2$ ,  $\alpha = \alpha' + r$ , and  $\Omega^{\alpha}_{\mu,\nu}(x)$  is defined by (4.9). The expression  $\hat{\ell}$  is limit-circle at x = 0 and limit-point at  $x = \infty$  when  $\alpha$  is chosen so that  $-1 < \alpha < 1$ .

Laguerre-type XOPs of order  $n \in \mathbb{N}_{\mu,\nu}$  are written as  $L^{\alpha}_{\mu,\nu,n}$  and the sequence forms a complete orthogonal set in  $L^2[(0,\infty), W]$  by Lemma 4.6, see Remark 4.3 for more details. These polynomials all belong to a family of eigenfunctions given by  $\Omega_{M_1,M_2}[h^{\alpha}(x,\lambda)]$  and are in a self-adjoint extension of  $\mathcal{L}_{\min}$  denoted  $\mathcal{L}_0$ .

Eigenvalues of self-adjoint extensions of  $\mathcal{L}_{\min}$ , denoted  $\mathcal{L}_{\tau}$  in general, can be extracted from an explicit Weyl *m*-function given by (7.6) generated by a boundary triple. The resulting *m*-function can be compared to the *m*-function of self-adjoint extensions of the classical Laguerre symmetric operator. Differences stem from two important processes: shifting Maya diagrams into canonical/conjugate canonical position, and evaluating Wronskians with these shifted diagrams at x = 0. Other spectral information, such as the strength of the point masses in the spectral measure, can also be obtained from the *m*-function, see e.g., [7].

The examples presented in Sections 3 and 8 provide insight into how eigenvalues are changed by different types of Maya diagrams. Inverse spectral theory, where a set of eigenvalues is given and a corresponding set of seed functions within the Wronskian is determined (thereby fixing the differential equation and self-adjoint extension), however, remains unknown in general and warrants further investigation.

Acknowledgments. The authors would like to thank Niels Bonneux for correspondence that provided a detailed proof of [10, Lemma 5.1]. This correspondence was instrumental in the proofs of Theorem 6.1 and Corollary 6.2. The authors would also like to thank the referees for suggesting many detailed and thoughtful improvements to the presentation of the paper.

#### References

- N. Aronszajn, On a problem of Weyl in the theory of singular Sturm-Liouville equations. Amer. J. Math. 79 (1957), 597–610 Zbl 0079.10802 MR 88623
- [2] M. J. Atia, L. L. Littlejohn, and J. Stewart, Spectral theory of X<sub>1</sub>-Laguerre polynomials. Adv. Dyn. Syst. Appl. 8 (2013), no. 2, 181–192 MR 3162140
- J. Behrndt, S. Hassi, and H. de Snoo, *Boundary value problems, Weyl functions, and differential operators*. Monogr. Math. 108, Birkhäuser/Springer, Cham, 2020
   Zbl 1457.47001 MR 3971207
- [4] S. Bochner, Über Sturm-Liouvillesche Polynomsysteme. Math. Z. 29 (1929), no. 1, 730– 736 Zbl 55.0260.01 MR 1545034
- [5] R. F. Boisvert, C. W. Clark, H. S. Cohl, D. W. Lozier, M. A. McClain, B. R. Miller, A. B. Olde Daalhuis, F, W. J. Olver, B. V. Saunders, B. I. Schneider (eds.), NIST digital library of mathematical functions. Release 1.0.22 of 2019-03-15 https://dlmf.nist.gov/ visited on 20 November 2023
- [6] N. Bonneux and A. B. J. Kuijlaars, Exceptional Laguerre polynomials. *Stud. Appl. Math.* 141 (2018), no. 4, 547–595 Zbl 1408.33019 MR 3879969
- M. Bush, D. Frymark, and C. Liaw, Singular boundary conditions for Sturm-Liouville operators via perturbation theory. *Canad. J. Math.* 75 (2023), no. 4, 1110–1146
   Zbl 07720708 MR 4620317
- [8] M. M. Crum, Associated Sturm-Liouville systems. Quart. J. Math. Oxford Ser. (2) 6 (1955), 121–127 Zbl 0065.31901 MR 72332
- [9] V. A. Derkach and M. M. Malamud, Generalized resolvents and the boundary value problems for Hermitian operators with gaps. *J. Funct. Anal.* 95 (1991), no. 1, 1–95 Zbl 0748.47004 MR 1087947
- [10] A. J. Durán, Exceptional Meixner and Laguerre orthogonal polynomials. J. Approx. Theory 184 (2014), 176–208 Zbl 1295.33013 MR 3218798
- [11] A. J. Durán and M. Pérez, Admissibility condition for exceptional Laguerre polynomials. J. Math. Anal. Appl. 424 (2015), no. 2, 1042–1053 Zbl 1305.33019 MR 3292716

- [12] D. Frymark, Boundary triples and Weyl *m*-functions for powers of the Jacobi differential operator. J. Differential Equations 269 (2020), no. 10, 7931–7974 Zbl 1493.34080 MR 4113193
- [13] D. Frymark and C. Liaw, Spectral properties of singular Sturm–Liouville operators via boundary triples and perturbation theory. J. Differential Equations 363 (2023), 391–421 Zbl 07684541 MR 4568769
- [14] M. Á. García-Ferrero, D. Gómez-Ullate, and R. Milson, A Bochner type characterization theorem for exceptional orthogonal polynomials. J. Math. Anal. Appl. 472 (2019), no. 1, 584–626 Zbl 1421.42014 MR 3906391
- [15] F. Gesztesy, L. L. Littlejohn, and R. Nichols, On self-adjoint boundary conditions for singular Sturm-Liouville operators bounded from below. J. Differential Equations 269 (2020), no. 9, 6448–6491 Zbl 1458.34143 MR 4107059
- [16] D. Gómez-Ullate, Y. Grandati, and R. Milson, Shape invariance and equivalence relations for pseudo-Wronskians of Laguerre and Jacobi polynomials. J. Phys. A 51 (2018), no. 34, article no. 345201 Zbl 1400.33017 MR 3835328
- [17] D. Gómez-Ullate, Y. Grandati, and R. Milson, Spectral theory of exceptional Hermite polynomials. In From operator theory to orthogonal polynomials, combinatorics, and number theory—a volume in honor of Lance Littlejohn's 70th birthday, pp. 173–196, Oper. Theory Adv. Appl. 285, Birkhäuser/Springer, Cham, 2021 Zbl 1493.33007 MR 4367467
- [18] D. Gómez-Ullate, N. Kamran, and R. Milson, An extended class of orthogonal polynomials defined by a Sturm-Liouville problem. J. Math. Anal. Appl. 359 (2009), no. 1, 352–367 Zbl 1183.34033 MR 2542180
- [19] D. Gómez-Ullate, N. Kamran, and R. Milson, Exceptional orthogonal polynomials and the Darboux transformation. J. Phys. A 43 (2010), no. 43, article no. 434016 Zbl 1202.81036 MR 2727790
- [20] D. Gómez-Ullate, N. Kamran, and R. Milson, An extension of Bochner's problem: exceptional invariant subspaces. J. Approx. Theory 162 (2010), no. 5, 987–1006 Zbl 1214.34079 MR 2610341
- [21] D. Gómez-Ullate, N. Kamran, and R. Milson, Two-step Darboux transformations and exceptional Laguerre polynomials. J. Math. Anal. Appl. 387 (2012), no. 1, 410–418 Zbl 1232.33013 MR 2845760
- [22] D. Gómez-Ullate, N. Kamran, and R. Milson, A conjecture on exceptional orthogonal polynomials. *Found. Comput. Math.* 13 (2013), no. 4, 615–666 Zbl 1282.33019 MR 3085680
- [23] Y. Grandati, Multistep DBT and regular rational extensions of the isotonic oscillator. Ann. Physics 327 (2012), no. 10, 2411–2431 Zbl 1255.81169 MR 2970643
- [24] J. S. Kelly, C. Liaw, and J. Osborn, Moment representations of exceptional X<sub>1</sub> orthogonal polynomials. J. Math. Anal. Appl. 455 (2017), no. 2, 1848–1869 Zbl 1370.33010 MR 3671258
- [25] C. Liaw, L. Littlejohn, and J. S. Kelly, Spectral analysis for the exceptional X<sub>m</sub>-Jacobi equation. *Electron. J. Differential Equations* (2015), article no. 194 Zbl 1320.33017 MR 3386555

- [26] C. Liaw, L. L. Littlejohn, R. Milson, and J. Stewart, The spectral analysis of three families of exceptional Laguerre polynomials. J. Approx. Theory 202 (2016), 5–41 Zbl 1333.33018 MR 3440071
- [27] C. Liaw, L. L. Littlejohn, J. Stewart, and Q. Wicks, A spectral study of the second-order exceptional X<sub>1</sub>-Jacobi differential expression and a related non-classical Jacobi differential expression. J. Math. Anal. Appl. 422 (2015), no. 1, 212–239 Zbl 1303.33012 MR 3263455
- [28] M. Marletta and A. Zettl, The Friedrichs extension of singular differential operators. J. Differential Equations 160 (2000), no. 2, 404–421 Zbl 0954.47012 MR 1736997
- [29] I. Marquette and C. Quesne, New families of superintegrable systems from Hermite and Laguerre exceptional orthogonal polynomials. J. Math. Phys. 54 (2013), no. 4, article no. 042102 Zbl 1282.70036 MR 3088222
- [30] B. Midya and B. Roy, Exceptional orthogonal polynomials and exactly solvable potentials in position dependent mass Schrödinger Hamiltonians. *Phys. Lett. A* 373 (2009), no. 45, 4117–4122 Zbl 1234.81072 MR 2552404
- [31] M. A. Naimark, *Linear differential operators*. Part II: Linear differential operators in Hilbert space. Frederick Ungar, New York, 1968 Zbl 0227.34020 MR 0262880
- [32] M. Noumi, *Painlevé equations through symmetry*. Transl. Math. Monogr. 223, American Mathematical Society, Providence, RI, 2004 Zbl 1077.34003 MR 2044201
- [33] S. Odake and R. Sasaki, Infinitely many shape invariant potentials and new orthogonal polynomials. *Phys. Lett. B* 679 (2009), no. 4, 414–417 MR 2569488
- [34] S. Odake and R. Sasaki, Another set of infinitely many exceptional  $(X_{\ell})$  Laguerre polynomials. *Phys. Lett. B* **684** (2010), no. 2-3, 173–176 MR 2588057
- [35] G. Post and A. Turbiner, Classification of linear differential operators with an invariant subspace of monomials. *Russian J. Math. Phys.* 3 (1995), no. 1, 113–122
   Zbl 0913.34069 MR 1337491
- [36] C. Quesne, Exceptional orthogonal polynomials, exactly solvable potentials and supersymmetry. J. Phys. A 41 (2008), no. 39, article no. 392001 Zbl 1192.81174 MR 2439200
- [37] R. Sasaki, S. Tsujimoto, and A. Zhedanov, Exceptional Laguerre and Jacobi polynomials and the corresponding potentials through Darboux-Crum transformations. J. Phys. A 43 (2010), no. 31, article no. 315204 Zbl 1200.33013 MR 2665675
- B. Simon, Spectral analysis of rank one perturbations and applications. In *Mathematical quantum theory*. II. Schrödinger operators (Vancouver, BC, 1993), pp. 109–149, CRM Proc. Lecture Notes 8, American Mathematical Society, Providence, RI, 1995
   Zbl 0824.47019 MR 1332038
- [39] B. Simon, *Trace ideals and their applications*. Second edn., Math. Surveys Monogr. 120, American Mathematical Society, Providence, RI, 2005 Zbl 1074.47001 MR 2154153

Received 10 October 2022; revised 17 July 2023.

#### Dale Frymark

Institut für Angewandte Mathematik, Technische Universität Graz, Steyrergasse 30, 8010 Graz, Austria; frymark@math.tugraz.at

#### Jessica Stewart Kelly

Department of Mathematics, Christopher Newport University, 1 Avenue of the Arts, Newport News, VA 23606, USA; jessica.kelly@cnu.edu