



## **Broadening the selection criteria for Astronauts undertaking long-term space travel**

Downloaded from: <https://research.chalmers.se>, 2024-04-25 21:28 UTC

Citation for the original published paper (version of record):

Yasuda, H., Sihver, L. (2022). Broadening the selection criteria for Astronauts undertaking long-term space travel. *Frontiers in Nuclear Medicine*, 2. <http://dx.doi.org/10.3389/fnume.2022.997718>

N.B. When citing this work, cite the original published paper.



## OPEN ACCESS

## EDITED BY

Gabriel G. De La Torre,  
University of Cádiz, Spain

## REVIEWED BY

Cyril Schandorf,  
University of Ghana, Ghana  
Livio Narici  
University of Rome Tor Vergata, Italy

## \*CORRESPONDENCE

Hiroshi Yasuda  
hyasuda@hiroshima-u.ac.jp

## SPECIALTY SECTION

This article was submitted to Dosimetry and Radiation Safety, a section of the journal Frontiers in Nuclear Medicine

RECEIVED 19 July 2022

ACCEPTED 12 October 2022

PUBLISHED 04 November 2022

## CITATION

Yasuda H and Sihver L (2022) Broadening the selection criteria for Astronauts undertaking long-term space travel. *Front. Nucl. Med.* 2:997718. doi: 10.3389/fnume.2022.997718

## COPYRIGHT

© 2022 Yasuda and Sihver. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

# Broadening the selection criteria for Astronauts undertaking long-term space travel

Hiroshi Yasuda<sup>1\*</sup> and Lembit Sihver<sup>2,3,4</sup>

<sup>1</sup>Research Institute for Radiation Biology and Medicine, Hiroshima University, Hiroshima, Japan,

<sup>2</sup>Technische Universität Wien - Institute of Atomic and Subatomic Physics, Wien, Austria,

<sup>3</sup>Department of Physics, Chalmers University of Technology, Gothenburg, Sweden, <sup>4</sup>Department of Radiation Dosimetry, Nuclear Physics Institute of the CAS, Prague, Czech Republic

## KEYWORDS

astronaut, space travel, radiation exposure, health risk, deep space mission

One of the major concerns in future deep space missions to the moon and Mars is the increased radiological risk of astronauts. They will be exposed to enhanced levels of ionizing radiation from natural sources, such as galactic cosmic radiation, radiation from the Sun including high-energy charged particles at solar particle events (SPEs), and radiation belts surrounding the Earth (1, 2). The accumulated radiation dose over a long-term mission to Mars is estimated to reach 1 Sv or more, depending on duration, shielding and time in the solar cycle (3). Although it is a rare event, SPE particles could further increase their doses to a serious level up to as high as 10 Gy (4), which is far beyond the dose limit for radiation workers (5), and could induce severe acute deterministic effects such as deterioration of blood-forming function (6), reproductive potential (7), cataract (8) and even death by acute radiation syndrome.

With these concerns, space agencies such as the National Aeronautics and Space Administration (NASA) of the USA and Japan Aerospace Exploration Agency (JAXA) have developed dose-limitation criteria for controlling the space radiation exposure of astronauts below an acceptable level. **Table 1** shows the previous career dose limits for NASA (1) and the current limits for JAXA (9) astronauts involved in low-Earth-orbit missions. The limits of NASA were determined to constrain the increasing cancer risk incurred by an astronaut to 3%; more precisely, the limits for NASA astronauts were not to exceed 3% risk of exposure-induced death from fatal cancers at a 95% confidence level based on a statistical assessment of the uncertainties in the risk projections (10). As the unit-dose cancer risk generally increases with age (5, 11, 12), the dose limit became higher for older astronauts than that for young ones. In addition, at the same age range, a limit value for female astronauts was higher than that for male ones, reflecting the fact that the breast has a notably higher radiosensitivity (5, 11, 12).

As inferred from the values in **Table 1**, these dose-limitation criteria allowed older male astronauts to have more opportunities of space travel than young or female astronauts, which could be regarded as a problem of inequality. Then, the National Academy of Sciences (NAS) in United States recently made a recommendation of applying a 600 mSv age and gender independent career limit of effective dose based on a median estimate to reach 3% cancer fatality for 35-year-old females (13), withdrawing the age and gender specific limits. This recommendation is expected to allow equivalent flight opportunities for all male/female astronauts of different ages

(13). Following this recommendation of NAS, NASA updated the standards for crew health in 2022 (14). In the new standards, the total career effective dose of an astronaut due to space flight radiation exposure shall be less than 600 mSv and this limit is universal for all ages and sexes. It is also required that radiation exposure from all sources below the limit shall be further minimized following the principle of “as low as reasonably achievable (ALARA)”.

This concept has been criticized by some researchers for the reason that it could have negative impacts on crew health and safety and violate the principles of radiological protection (15). It is worried also that such a lower annual dose limit would make a Mars mission unfeasible since the cumulative dose in one mission is expected to far exceed 600 mSv. While, recognizing the fact that the risk of radiation-induced cancer can considerably change among individuals and also under different radiation dose rates, the National Aeronautics and Space Administration (NASA) had requested National Council on Radiation Protection and Measurements (NCRP) in United States to evaluate the risk of radiation-induced lung cancer in populations exposed to chronic or fractionated radiation to learn whether differences exist when exposures occur gradually over years contrasted with the acute exposure received by the Japanese atomic-bomb survivors. In response to the request from NASA, NCRP launched a scientific committee and have been working to prepare a commentary (16) on this issue with accompanying recommendations for NASA.

Considering such a fluid situation on the radiological protection criteria for astronauts, the authors like to present here a different viewpoint which might mitigate the ongoing discussion on radiological protection of astronauts. Apart from the possible ethical issue of discriminating people by age or gender, it is known that elderly people have generally more health problems related to aging. Any person inevitably becomes vulnerable with age through various types of deteriorative changes due to many causes (17, 18), although the pace of aging varies among individuals (19, 20). Some of the typical aging symptoms that could be commonly experienced before age 65 years (general retirement age in many countries) are as follows:

- Loss of muscle mass and strength (21)
- Weakening and embrittlement of bones (osteoporosis) (22)
- Loss of arterial elasticity (atherosclerosis) and other cardiovascular changes (23)
- Difficulty in focusing eyes on close objects (presbyopia)
- Lowering hearing ability of ears (presbycusis)
- Cardiovascular diseases
- Menopause associated with hot flash, disruption in sleep, subsequent osteoporosis, etc.

In addition, risks for cognitive impairment (dementia, Alzheimer’s disease, Parkinson’s disease, etc.) (24), cataracts (8), and carcinogenesis (12, 23) increase with age. With advancing

**TABLE 1** Previous career dose limits for the astronauts of NASA (USA) (1) and current limits for JAXA (Japan) (9) involved in low-earth-orbit space missions.


Space agency (country)	Male (Sv)	Female (Sv)
NASA (USA)	1.5 Sv for 25 y	1.0 Sv for 25 y
	2.5 Sv for 35 y	1.75 Sv for 35 y
	3.25 Sv for 45 y	2.5 Sv for 45 y
	4.0 Sv for 55 y	3.0 Sv for 55 y
JAXA (Japan)	0.6 Sv for 27 y	0.5 Sv for 27 y
	0.7 Sv for 31 y	0.6 Sv for 31 y
	0.8 Sv for 36 y	0.65 Sv for 36 y
	0.95 Sv for 41 y	0.75 Sv for 41 y
	1.0 Sv for 46 y	0.8 Sv for 46 y

The limit value is given as effective dose for the age of the first flight exposure.

age, individuals also tend to have difficulty coping with various stresses such as strenuous exercise and environmental changes; and those stresses tend to cause functional deterioration of some of their organs such as the heart, urinary organs, and brain. The senescence-associated health deterioration can be different between male and female people, as women have some biological advantage related to their ability to bear a child and the physiological systems that permit pregnancy (25).

For preventing the occurrence of a serious problem caused by the aging of astronauts in long-term deep-space mission, it is desirable to carefully consider all possible age- and gender-related deterioration of health when selecting the astronauts who will take up difficult tasks during a long space mission for up to few years. While young astronauts are generally tough and swiftly acting, experienced older astronauts are more knowledgeable and prudent, which would make them more reliable when facing unexpected troubles. The current criteria of NASA on crew health management (14) does not clearly indicate how to evaluate and balance the unique competences of individual astronauts who will work together for a long period in the same mission. According to these facts, the authors propose to deal with the age- and gender-dependent radiation sensitivity as one of the major qualifications required for astronauts involved in a deep space mission.

The conceptual basis of this idea can be illustrated with a radar chart as shown in **Figure 1**. This chart has six axes of major requirements on qualifications of astronauts with two example patterns of typical scores of young and older astronauts. The requirements assumed here are (1) physical strength related to muscles, bones, and cartilage; (2) physiological soundness related to cardiovascular, renal, digestive, respiratory, and immune systems; (3) sensory capability related to perceptions with eyes, ears, nose, and nerves; (4) cognition and memory related to the neuroimaging functions mainly controlled by brain; (5) knowledge and judgement supported by acquired intellectual base and experience; and (6) radiological health which could be quantified as an inverse quantity of radiation sensitivity regarding carcinogenesis. Older astronauts could have higher



scores on the knowledge, judgement and radiological health, while their scores on physical strength, physiological soundness, sensory capability, and cognitive functions would be lower.

For the success of future deep-space missions, it will be crucial to carefully select healthy astronauts who can show strength against not only psychosomatic stress including radiation-induced cancers, but also the inevitable aging effects during a long traveling period of up to few years; possible appearance of different-quality age-associated symptoms should be projected in the process of crew selection. In this sense, a routine health surveillance programme based on the general principles of occupational health will take a vital role for assessing the initial and continuing fitness of the astronauts for achieving their intended tasks as a team in a specific space mission. With these efforts, it is expected that a well-balanced team of male and female astronauts covering a broad range of age will be formed, so that the scores regarding all requirements as shown in [Figure 1](#) could be maximized as a whole.

In conclusion, the diversity in formation of a team of astronauts is preferably to be pursued for successful deep space missions in the future. For this, further studies for

overcoming various aging-related health issues are needed to smash the current highest score.

## Author contributions

HY: conceptualization, investigation and original draft preparation. LS: review, editing and supervision. All authors contributed to the article and approved the submitted version.

## Acknowledgments

The authors acknowledge continuous support of colleagues from Hiroshima University.

## Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their

affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

## References

1. National Council on Radiation Protection and Measurements (NCRP). *Recommendations of dose limits for low earth orbit*. Bethesda, MD: NCRP Report No. 132. NCRP (2000).
2. International Commission on Radiological Protection (ICRP). *Assessment of radiation Exposure of astronauts in space*. London: ICRP Publication 123. Ann. ICRP 42(4). SAGE (2013).
3. Cucinotta FA, Schimmerling W, Wilson JW, Peterson LE, Badhwar GD, Saganti PB, et al. Space radiation cancer risks and uncertainties for Mars missions. *Radiat Res* (2001) 156:682–8. doi: 10.1667/0033-7587(2001)156[0682:SRCRAU]2.0.CO;2
4. Townsend LW, Stephens Jr. DL, Hoff JL, Zapp EN, Moussa HM, Miller TM, et al. The Carrington event: possible doses to crews in space from a comparable event. *Adv Space Res* (2006) 38:226–31. doi: 10.1016/j.asr.2005.01.111
5. International Commission on Radiological Protection (ICRP). 2007 Recommendations of the International Commission on Radiological Protection. ICRP Publication 103. ICRP Publication 103, Ann. ICRP 37 (2–4). Elsevier (2007).
6. National Council on Radiation Protection and Measurements (NCRP). *The relative biological effectiveness of radiations of different quality*. NCRP Report No. 104. NCRP. (1990).
7. Wang B, Yasuda H. Relative biological effectiveness of high LET particles on the reproductive system and fetal development. *Life*. (2020) 10:298. doi: 10.3390/life10110298
8. International Commission on Radiological Protection (ICRP). ICRP statement on tissue reactions/early and late effects of radiation in normal tissues and organs—threshold doses for tissue reactions in a radiation protection context. ICRP Publication 118, Ann. ICRP 41(1–2). Pergamon Press (2012).
9. Japan Aerospace Exploration Agency (JAXA). *Rules on radiation exposure management for the astronauts involved in the international space station*. Tokyo: JAXA (2013). [http://iss.jaxa.jp/med/research/radiation/pdf/kitei\\_130626\\_a.pdf](http://iss.jaxa.jp/med/research/radiation/pdf/kitei_130626_a.pdf) [in Japanese].
10. Cucinotta FA, Kim MY, Chappell L. *Space radiation cancer risk projections and uncertainties-2012*. Houston, TX: NASA TP 2013-217375. NASA (2013).
11. National Research Council Committee on Health Effects of Exposure to Low Levels of Ionizing Radiations (BEIR VII). *Health effects of exposure to low levels of ionizing radiations*. Washington, DC: National Academies Press (1998).
12. United Nations Scientific Committee on the Effects of Atomic Radiation (UNSCEAR). *Effects of ionizing radiation. UNSCEAR 2006 report to the general assembly with scientific annexes volume I. Annex A: Epidemiological studies of radiation and cancer*. Vienna: UNSCEAR (2008).
13. National Academy of Sciences (NAS). *Space radiation and astronaut health: managing and communicating risk*. Washington, DC: National Academy Press (2021).
14. National Aeronautics and Space Administration (NASA). *NASA Space Flight Human-System Standard; Volume 1: Crew Health*. NASA-STD-3001, V. 1 Rev. B. NASA (2022).
15. Cucinotta FA, Schimmerling W, Blakely EA, Hei TK. A proposed change to astronaut exposures limits is a giant leap backwards for radiation protection. *Life Sci Space Res*. (2021) 31:59–70. doi: 10.1016/j.lssr.2021.07.005
16. National Council on Radiation Protection and Measurements (NCRP). *SC 1-27: Evaluation of sex-specific differences in lung cancer radiation risks and recommendations for use in transfer and projection models*. Bethesda, MD: NCRP (2022). Available from: <https://ncrponline.org/program-areas/sc-1-27-evaluation-of-sex-specific-differences-in-lung-cancer-radiation-risks-and-recommendations-for-use-in-transfer-and-projection-models/> (Accessed August 23, 2022)
17. Crimmins EM. Lifespan and healthspan: past, present, and promise. *Gerontologist*. (2015) 55:901–11. doi: 10.1093/geron/gnv130
18. Liochev SI. Which is the most significant cause of aging? *Antioxidants*. (2015) 4:793–810. doi: 10.3390/antiox4040793
19. Partridge L, Deelen J, Slagboom PE. Facing up to the global challenges of ageing. *Nature*. (2018) 561:45–56. doi: 10.1038/s41586-018-0457-8
20. Elliott ML, Caspi A, Houts RM, Ambler A, Broadbent JM, Hancox RJ, et al. Disparities in the pace of biological aging among midlife adults of the same chronological age have implications for future frailty risk and policy. *Nat Aging*. (2021) 1:295–308. doi: 10.1038/s43587-021-00044-4
21. Ryall JG, Schertzer JD, Lynch GS. Cellular and molecular mechanisms underlying age-related skeletal muscle wasting and weakness. *Biogerontology*. (2008) 9:213–28. doi: 10.1007/s10522-008-9131-0
22. Ensrud KE. Epidemiology of fracture risk with advancing age. *J Gerontol A Biol Sci Med Sci*. (2013) 68:1236–42. doi: 10.1093/gerona/glt092
23. Driver JA, Djoussé L, Logroscino G, Gaziano JM, Kurth T. Incidence of cardiovascular disease and cancer in advanced age: prospective cohort study. *Br Med J*. (2008) 337:a2467. doi: 10.1136/bmj.a2467
24. Grady CL, Craik FIM. Changes in memory processing with age. *Curr Opin Neurobiol*. (2000) 10:224–31. doi: 10.1016/S0959-4388(00)00073-8
25. Vlassoff C. Gender differences in determinants and consequences of health and illness. *J Health Popul Nutr*. (2007) 25:47–61. doi: 10.17226/26155