



Environmental assessment of diets: overview and guidance on indicator choice

Downloaded from: <https://research.chalmers.se>, 2025-07-02 03:40 UTC

Citation for the original published paper (version of record):

Ran, Y., Cederberg, C., Jonell, M. et al (2024). Environmental assessment of diets: overview and guidance on indicator choice. *The Lancet Planetary Health*, 8(3): e172-e187.
[http://dx.doi.org/10.1016/S2542-5196\(24\)00006-8](http://dx.doi.org/10.1016/S2542-5196(24)00006-8)

N.B. When citing this work, cite the original published paper.

Environmental assessment of diets: overview and guidance on indicator choice



Ylva Ran, Christel Cederberg, Malin Jonell, Kristina Bergman, Imke J M De Boer, Rasmus Einarsson, Johan Karlsson, Hanna Karlsson Potter, Michael Martin, Geneviève S Metson, Thomas Nemecek, Kimberly A Nicholas, Åsa Strand, Pernilla Tidåker, Hayo Van der Werf, Davy Vanham, Hannah H E Van Zanten, Francesca Veronesi, Elin Rööf

Comprehensive but interpretable assessment of the environmental performance of diets involves choosing a set of appropriate indicators. Current knowledge and data gaps on the origin of dietary foodstuffs restrict use of indicators relying on site-specific information. This Personal View summarises commonly used indicators for assessing the environmental performance of diets, briefly outlines their benefits and drawbacks, and provides recommendations on indicator choices for actors across multiple fields involved in activities that include the environmental assessment of diets. We then provide recommendations on indicator choices for actors across multiple fields involved in activities that use environmental assessments, such as health and nutrition experts, policy makers, decision makers, and private-sector and public-sector sustainability officers. We recommend that environmental assessment of diets should include indicators for at least the five following areas: climate change, biosphere integrity, blue water consumption, novel entities, and impacts on natural resources (especially wild fish stocks), to capture important environmental trade-offs. If more indicators can be handled in the assessment, indicators to capture impacts related to land use quantity and quality and green water consumption should be used. For ambitious assessments, indicators related to biogeochemical flows, stratospheric ozone depletion, and energy use can be added.

Introduction

Environmental assessments of food products and diets are increasingly used by a wide variety of actors, including the research community across disciplines;¹ public policy makers;^{2,3} non-governmental organisations;^{4,5} and private-sector food industries, retailers, and consumers.^{6–8} Multiple indicators for assessing the environmental performance of individual food items and agricultural systems have been developed,^{9,10} but few are in regular use to assess diets,¹¹ which commonly comprise of hundreds of foods sourced from many locations, with little information on exact origin or production system.

When selecting indicators for environmental assessment of diets, researchers and other actors usually (and understandably) choose those that are well established, which include those related to climate change, water use, and land use;^{11,12} however, to capture the overall environmental performance of diets, including potential trade-offs between different environmental aspects, a broader set of indicators is required.¹³ If assessment methods for such indicators require data that are commonly unavailable, and give results that cannot easily be communicated, they risk being impractical and ineffective for decision making for a broad audience, so simplification is needed.¹⁴

Food items in diets are commonly sourced from different locations globally, with effects that vary depending on site-specific conditions and management practices; however, identifying the exact origin or production system for the multitude of foods available and the ingredients that make up these foods is often difficult. This difficulty prevents the use of indicators suggested in current standards for product assessment for assessing diets (eg, the Product Environmental Footprint developed in the EU).¹⁵ Beyond individual food items, providing

specific guidance on indicator choice for assessment of environmental sustainability of diets is needed to find the right balance between type and number of indicators to use, and what they can reveal.

In this Personal View we provide an overview of commonly used dietary environmental performance indicators, identify aspects that are missing, and guide readers to a relevant indicator choice that captures the key aspects of the environmental impact. Although we focus on indicators that assess the environmental component of sustainability, we acknowledge that the concept of sustainability goes beyond environmental sustainability and includes a range of socioeconomic aspects that also need consideration when striving to feed future generations sustainably.¹⁶

Commonly used environmental indicators

We searched the Web of Science Core Collection and Scopus for reviews on indicators for sustainable diets using the Boolean search phrase “indicator* AND sustainab* AND diet*”. Two authors (YR and ER) screened reviews by title and abstract. Full-text and data extraction was done by the same two authors. The search method, screening process, and results are described in more detail in the appendix (pp 3–43). The search generated 109 records after deduplication, of which seven reviews^{11,13,14,17–20} met our criteria and assessed the environmental sustainability of diets and synthesised multiple indicators. After identifying relevant reviews, we extracted the indicators identified in each study (appendix pp 31–43). We included indicators “used to assess, compare, and control the impacts on the environment”,²⁰ and excluded indices that aggregate several environmental and other sustainability impacts. We categorised indicators according to the planetary

Lancet Planet Health 2024;
8: e172–87

Department of Energy and Technology, Swedish University of Agricultural Sciences, Uppsala, Sweden (Y Ran PhD, R Einarsson PhD, J Karlsson PhD, H Karlsson Potter PhD, P Tidåker PhD, E Rööf PhD); Division of Physical Resource Theory, Department of Space, Earth and Environment, Chalmers University of Technology, Göteborg, Sweden (Prof C Cederberg PhD); Global Economic Dynamics and the Biosphere, Royal Swedish Academy of Science, Stockholm, Sweden (M Jonell PhD); Stockholm Resilience Centre, Stockholm University, Stockholm, Sweden (M Jonell); KTH Royal Institute of Technology, Department of Sustainable Development, Environmental Science and Engineering, Stockholm, Sweden (K Bergman MSc); Animal Production Systems Group, Wageningen University & Research, Wageningen, Netherlands (Prof J M de Boer PhD); IVL Swedish Environmental Research Institute, Stockholm, Sweden (M Martin PhD, Å Strand PhD); Department of Geography and Environment, Social Sciences Centre, University of Western Ontario, London, ON, Canada (G S Meston PhD); Ecological and Environmental Modeling Division, Department of Physics, Chemistry and Biology, Linköping University, Linköping, Sweden (G S Meston); Agroscope, Life Cycle Assessment Research Group, Zurich, Switzerland (T Nemecek PhD); Lund University Centre for Sustainability Studies, Lund, Sweden (K A Nicholas PhD); French National Research Institute for Agriculture, Food and Environment, l'Institut Agro Rennes-Angers, Rennes, France (H Van der Werf PhD MG); Ispra, Italy (D Vanham PhD); Farming Systems Ecology

Group, Wageningen University and Research, Wageningen, Netherlands (Prof H H E Van Zanten PhD); Department of Global Development, College of

boundaries framework²¹ to relate them to biophysical earth system processes, and excluded duplicate environmental indicators with similar definitions and units but different names (eg, land use and land occupation). A summary of the most commonly applied

indicators are listed in table 1. Some indicators relate to several boundaries (eg, grey water use relates to both novel entities and biogeochemical flows,²³ and acidification potential relates to atmospheric aerosol loading and biogeochemical flows). Some overlaps are

	Main influence from the food system	Pressure indicators	Impact indicators
Climate change			
Warming of the atmosphere has effects such as increasing the frequency and amplitude of extreme weather events and threats to humans, other species, and ecosystems in oceans, coastal regions, and land	Use of land, animals, energy, and fuels that generate GHG emissions (mainly CO ₂ , CH ₄ , and N ₂ O), and land use change (eg, deforestation) that results in GHG emissions and changes in albedo (reflection of solar radiation)	Not commonly used†	Carbon footprint (measured in kg CO ₂ eq)—ie, aggregation of different GHGs into CO ₂ equivalents considering their different warming dynamics
Land system change			
Changes in land cover destroy ecosystems, lead to decline and extinction of species, and affect local and global climates and precipitation patterns (eg, the Amazon rainforest risks becoming a semi-arid savannah); land suitable for agricultural production is a finite resource	Expansion of agricultural land through deforestation and conversion of natural grasslands into intensively managed grasslands and cropland, and intensification of land use	Use of cropland or total agricultural land (measured in m ² × year)—ie, the amount of cropland (and pasture in terms of total agricultural land) during a certain time (often 1 year) needed to produce the food or diet	Scarcity-related land footprint (measured in m ² land eq × year)—ie, the amount of land used for production adjusted for local stress or scarcity over land resources
Freshwater use			
Freshwater use and withdrawal causes local and regional water stress and competition for water resources	Cultivation of food and feed crops that uses rainwater infiltrated as soil moisture (green water) and liquid freshwater resources (blue water) for irrigation, animal drinking, and servicing; water is also used in food and feed processing and preparing	Blue water, green water, or total water footprint (measured in L)—ie, the volume of water used in production	Stress-related water footprint (measured in L water eq)—ie, the volume of water used adjusted for local water stress or water scarcity, as in LCA; and water stress assessment (eg, using SDG indicator 6.4.2 ²²) as conducted in environmental footprint assessment
Biogeochemical flows			
Eutrophication of terrestrial, freshwater, and marine ecosystems leads to excessive growth of algae that causes oxygen depletion and death of fish and other freshwater or marine species; and terrestrial eutrophication causes loss of species, changes to ecosystem structure and functioning, and homogenisation of vegetation	Addition of reactive N and P by conversion of atmospheric N ₂ into reactive N (mainly through synthetic fertilisers but also cultivation of leguminous crops) and P mining; considerable amounts of N and P end up in the environment, causing eutrophication	New N and P input (measured in kg N and P)—ie, the addition of new N and P to agricultural land; N footprint or N losses (measured in kg N)—ie, reactive N emissions to the environment; P footprint (measured in kg P)—ie, P emissions to the environment; and greywater footprint (measured in L)—ie, the volume of water needed to assimilate a pollutant load that reaches a water body	Eutrophication potential (measured in kg PO ₄ ³⁻ eq)—ie, aggregation of potential impact from N and P emissions based on amounts of these nutrients needed to build phytoplankton biomass; marine eutrophication potential assesses N increase in water by converting emissions of eutrophication substances into kg N eq; and freshwater eutrophication potential assesses P increase in freshwater by converting emissions of eutrophication substances into emissions of kg P eq
Atmospheric aerosol loading			
Atmospheric aerosol loading affects the climate system (through cooling and warming) and hydrological cycles (eg, a shift in Asian monsoon circulation), and harms human health (eg, by contributing to cardiopulmonary disease, acute respiratory infections, etc)§	Use of fuels, chemicals, and other compounds that cause air pollution with primary aerosols (fine particulate matter [PM ₁₀] in air) by emissions of black carbon (soot) from activities such as cooking, land clearing, burning biomass, and heating with biofuels	Not commonly used	Particulate matter formation potential (measured in PM _{2.5} eq) measures impacts on human health caused by emissions of fine particulate matter and its precursors (eg, NH ₃ , NO _x , or SO ₂)§
Atmospheric aerosol loading causes damage to vegetation and loss of freshwater fish from acidic precipitation, and harms human health (eg, by contributing to cardiopulmonary disease, acute respiratory infections, etc)§	Emissions of NH ₃ from manure management and fertiliser application (and some smaller contributions from SO ₂ and NO _x in fossil fuel combustion); secondary aerosols are formed (eg, SO ₂ and NO _x from fossil fuel combustion and from NH ₃ from manure management and fertiliser application)	Emissions of NH ₃ , SO ₂ , or NO _x , or a combination of these	Acidification potential (measured in g SO ₂ eq)—ie, aggregation of acidifying substances converted to equivalents of the acidification potential of SO ₂ ; particulate matter formation potential (measured in PM _{2.5} eq) measures impacts on human health caused by emissions of fine particulate matter and its precursors (eg, NH ₃ , NO _x , or SO ₂)§
Atmospheric aerosol loading also causes damage to vegetation (eg, by exposure to ozone) and harms human health (eg, by causing reduced lung function)§	Tropospheric, or ground-level, ozone is formed from NO _x and VOCs (eg, those emitted from engines or power plants) in the presence of sunlight	Emissions of NO _x	Photochemical ozone creation potential (measured in kg ethylene eq/m ² ppm) evaluates the contribution of individual substances to ozone formation potential by converting their emissions in parts per million per area to ethylene ozone formation potential equivalents

(Table 1 continues on next page)

therefore inevitable,^{11,14} as indicator categories are not mutually exclusive, which must be considered in indicator choice and interpretation.

We applied the Driver–Pressure–State–Impact–Response (DPSIR) framework²⁴ to categorise identified indicators, as this framework enables analysis of indicator types along a cause–effect chain in a structured way (appendix pp 46–48). The framework is well suited to our overall objective of providing recommendations to decision makers because it is already used in the context of public policy instruments.^{14,25} For environmental diet assessments, primarily the three middle categories of indicators (Pressure, State, and Impact) are relevant to this Personal View, as Drivers include large-scale socioeconomic factors such as farm subsidies, and Responses include policies formulated to address impacts, both of which are beyond the scope of identifying indicators for assessing the environmental sustainability of diets. Pressure indicators measure environmental load or stress caused by human activities through emissions of environmentally damaging substances such as greenhouse gases (GHGs) and

nitrogen (N), and use of resources such as land, water, and minerals. State indicators refer to physical, biological, and chemical conditions in an area, for example, concentrations of pollutants in lake water (panel 1). Impact indicators aim to capture what consequences such pressures have on the environment (eg, climate change or eutrophication of waterways³⁶), but vary in how far along the cause–effect chain they measure the impact (figure 1, panel 1, appendix p 49). Some indicators, such as how different GHGs contribute to climate change (called midpoint impact indicators in lifecycle assessments [LCAs]), merely aggregate different pressures using an equivalence relationship, whereas others (called endpoint impact indicators in LCAs) aim to quantify the actual impact caused (eg, in terms of damages to ecosystems or human health). Some authors, including van Dooren and colleagues,¹⁴ categorise LCA midpoint impact indicators as pressure indicators, as they do not measure the actual impact of any of the endpoint areas of protection (ie, human health, natural environment, and natural resources). We refer to them as midpoint impact indicators

Agriculture and Life Sciences, and Cornell Atkinson Center for Sustainability, Cornell University, Ithaca, NY, USA (Prof H H E Van Zanten); Industrial Ecology Programme, Department of Energy and Process Engineering, Norwegian University of Science and Technology, Trondheim, Norway (Prof F Verones PhD)

Correspondence to: Dr Ylva Ran, Department of Energy and Technology, Swedish University of Agricultural Sciences, Uppsala 75007, Sweden ylva.ran@slu.se

See Online for appendix

	Main influence from the food system	Pressure indicators	Impact indicators
(Continued from previous page)			
Biosphere integrity			
Destruction of ecosystems, extinction of species, and decline in genetic diversity reduce the resilience of the biosphere and cause declines in functions provided by biodiversity (eg, pollination)	Fishing, hunting, and use of land for agriculture diminishes wild stocks and harms ecosystems, and contributes to pollution from use of fertilisers, pesticides, and other chemicals	Use of cropland or total agricultural land (m ² × year)—ie, the amount of cropland (and pasture in the case of total agricultural land) during a certain time (often 1 year) that is needed to produce the food or diet	Biodiversity damage potential, expressed as potentially disappeared fraction of species—ie, the fraction of species potentially lost locally, regionally, or globally; potential species loss (number of species)—ie, the number of potential species that will eventually be lost locally, regionally, or globally
Novel entities**			
Release of new substances, new forms of existing substances, and modified life forms negatively affects ecosystems and human health	Use of pesticides and other inputs that include toxic or polluting substances such as microplastics and polycyclic aromatic hydrocarbons	Amount of pesticides used (measured in kg active substance)—ie, the amount of pesticides used to produce the food or diet; greywater footprint (measured in L)—ie, the volume of water needed to assimilate a pollutant load that reaches a water body	Freshwater, marine, and terrestrial ecotoxicity (measured in comparative toxic unit for ecosystems, kg 1,4-dichlorobenzene eq, or kg triethylene glycol eq) estimates the effect of use of toxic substance on organisms
Release of new substances, new forms of existing substances, and modified life forms negatively affects ecosystems and human health	Use of nanomaterials and plastics	Not commonly used	Not commonly used
Stratospheric ozone depletion			
Thinning of the protective ozone layer of the stratosphere causes negative effects on marine organisms and ecosystems and human health	Emissions of CFCs used as a cooling medium in food storage and transport, and emissions of N ₂ O from soils and manure management; CFCs and N ₂ O trigger ozone-destroying reactions	Not commonly used	Stratospheric ozone depletion (measured in kg CFC-11 eq)—ie, the aggregation of ozone-depleting substances based on their depletion potential relative to CFC-11
<p>Examples of indicators that are currently used for assessment of environmentally sustainable diets were identified in our synthesis of six key reviews.^{11–14,16,17} For a complete list of diet-related environmental indicators, see van Dooren and colleagues.¹⁴ All indicators can be based on different units of assessment (eg, per kg of food); here the denominator was per diet (potentially adjusted to calorie intake or similar). CFC-11= trichlorofluoromethane. CFCs=chlorofluorocarbons. eq=equivalent. GHG=greenhouse gas. LCA=lifecycle assessment. N=nitrogen. NO_x=nitrogen oxides. P=phosphorous. ppm=parts per million. SDG=Sustainable Development Goal. VOCs=volatile organic compounds. *Following Steffen and colleagues,¹⁹ ocean acidification is not included as it is a consequence of CO₂ emissions, covered by climate change. †In environmental footprint assessment, carbon footprint is identified as pollution (pressure) footprint.²³ ‡New N is nitrogen added using synthetic fertiliser or leguminous crops, and new P is added mined phosphorus. New N/P excludes N/P in manure and other organic amendments in which N/P only circulates within the agricultural system. Emissions of N and P occur throughout the food system, also from recirculating nutrients in manure, waste, and crop residues, but high emissions in the long term are only possible through continued inputs of new N and P. Mitigation potential is therefore distributed throughout the food system; however, new N and P inputs are useful indicators of total pressure. §Concerns and indicators related to effects on human health were outside of scope of this study. ¶[Small particles of solid or liquid suspended in air. Can be calculated as endpoints based on the aggregated impact from other environmental issues (climate change, land use change, disturbances to biogeochemical flows all cause biodiversity loss eventually) or as separate endpoints from different pressures, most commonly land use (discussed further in the section on biosphere integrity and in panel 1). **Novel entities are new substances, new forms of existing substances, and modified life forms with potentially unwanted geophysical and biological effects.¹⁹</p>			
Table 1: Environmental issues for each planetary boundary, * influences, and indicators			

to distinguish between indicators that merely measure use of a natural resource (eg, water or land) or an emission (eg, NH_3) and indicators that consider some type of impact

along the cause–effect chain. The DPSIR framework represents a cyclic process where a driver (in this case, food consumption) is directly linked to the endpoint

Panel 1: Tools and concepts for environmental performance assessment

Lifecycle assessment and environmental footprint assessment

The environmental performance of diets and foods is usually quantified using lifecycle assessment (LCA) or environmental footprint assessment (EFA).^{11,13} LCA and EFA of food usually focus on the environmental issues associated with a certain food product (eg, 1 kg of tomatoes) or macronutrient (eg, 1 kg of protein). By multiplying the amounts of foods in a certain diet by per-kilogram LCA or footprint of individual food products, the environmental impacts or pressures of a certain diet can be assessed.

LCA is a well established method by which emissions (eg, CO_2 and NH_3) and resource use (eg, land, water, and minerals) along the product's whole supply chain, or lifecycle, are quantified and categorised into a set of midpoint impact categories (eg, global warming, eutrophication, ecotoxicity etc) using mostly deterministic and linear cause–effect chains (figure 1).

Optionally, but more rarely in the case of diets, the analysis is extended along the cause–effect chain to include the actual final impact on humans and ecosystems (called endpoint impact categories). Such analysis enables aggregation of impacts caused by different damage categories such as global warming, eutrophication, or eco-toxicity into so-called areas of protection:²⁶ human Health, ecosystem quality,²⁷ and natural resources.²⁸

Although LCA has been standardised generally^{29,30} and more specifically for different food items, how LCAs are performed, the environmental impacts included, and the indicators used to describe them still vary considerably.

In EFA, environmental footprint is an umbrella term for the different footprint concepts developed in recent decades, including carbon, water, and land footprint.³¹ Footprints are commonly indicators of the pressure of human activities on the environment and, as in LCA, cover emissions and resource use along the whole supply chain from producer to consumer, and sometimes to waste management. Footprints and LCA impact categories sometimes overlap; for example, carbon footprint is the same as the global warming LCA midpoint impact category. On the other hand, water footprint is used to denote water use without considering the impact of water consumption in the landscape,³² but is also used by the LCA community to denote water scarcity-adjusted use of water.³³

Driver–Pressure–State–Impact–Response framework

The Driver–Pressure–State–Impact–Response framework (figure 2) developed by the European Environment Agency can be used to illustrate where indicators are placed along the cause–effect chain.²⁴ A driver indicator could be, for example, the amount of animal protein consumed, as livestock production is a major driver of many environmental impacts.

Pressure indicators relate to emissions and natural resource use, which put pressure on ecosystems that leads to different types of damage in different locations and contexts. For example, use of land is a pressure indicator because land is a finite resource and appropriation of land for agriculture puts major pressures on ecosystems in many regions, but to varying extents depending on the location. State indicators measure the properties of ecosystems, such as lake water pH. Impact indicators aim to capture the environmental impacts caused by pressures, such as the temperature change due to greenhouse gas emissions. Response indicators measure the policy responses to impacts caused, such as the ratio of emissions included in tax schemes or cap-and-trade systems.

Impact indicators are usually considered preferable for environmental assessments, as they measure the actual effect caused by drivers and pressures; however, they usually require information on where production takes place and site-specific data for that location that are usually not available for all foods in diets, such as soil and water status. Pressure indicators on the dietary level can still be highly valuable as decision support because they indicate the direction of change needed, but they can be misleading if not appropriately selected for the case at hand (panel 3). Even driver indicators can be useful, and sometimes preferable, due to their ease of assessment and interpretation. For example, the amount or proportion of animal-sourced foods or protein in a diet captures many of the environmental impacts and is easy to calculate and communicate. The framework represents a cyclic cause–effect chain linking drivers, such as animal-sourced food consumption, all the way to the societal response. Different responses, or the absence of a response, to environmental impacts can then be fed back and affect drivers, pressures, states, and impacts, ultimately affecting food availability and diets.

Planetary boundaries

The planetary boundaries framework defines environmental limits within which humanity can operate with low risk of causing destabilisation of the Earth system.^{21,34} The framework is based around nine biophysical processes that are important for regulating the stability of the planet, enabling human civilisations as we know them today. The concept has been influential in shaping academic and policy debate globally by reviving the discussion on planetary limits.³⁵ Although all human activities, including those related to food, housing, energy, industry, and transport, need to be encompassed within the planetary boundaries, Willett and colleagues²² presented boundaries of the global food system specifically for six Earth system processes, that is, the share of the overall operating space that food systems specifically should respect.

impact and response, which, represented here as damage to ecosystems, can ultimately limit the availability of food and thus, affect the driver of the system (figure 2, appendix p 50).

Our analysis showed that pressure indicators are mostly used for processes that result in local environmental concerns such as water and land use. Using pressure indicators for such local issues can be problematic, as effects are highly dependent on prevailing conditions (panel 2); however, their use is understandable because impact assessment typically requires complicated assessment models and detailed data on locations, production methods, and the local environment, which are seldom available for foods sourced from complex global supply chains.

Indicators related to climate change, land system change, freshwater use, and biogeochemical flows were the most

commonly assessed planetary boundaries in the dataset. Midpoint impact indicators were commonly used for climate change, atmospheric aerosol loading, stratospheric ozone, and novel entities, and pressure indicators were used for land system change, freshwater use, and biogeochemical flows (figure 3). The least frequently assessed boundaries were biosphere integrity, stratospheric ozone, and novel entities, confirming previous findings.³⁶ For biosphere integrity, widely different indicators were used, including indicators of global extinction rates,^{22,50} remaining fish stocks,^{28,31} and forest cover loss.⁵¹

State of the art and recommendations for indicator use

In addition to consulting the literature, topic experts were identified to discuss the state of the art and formulate recommendations for indicator use. We conducted




Figure 1: Environmental assessment of foods and examples of indicators along the lifecycle

The production lifecycle of the food product is first investigated (top). System boundaries differ, but in LCA studies, emissions and resource use from production to inputs, farming, manufacturing, and transport up to the retail gate are commonly included in the inventory (middle). From the inventory data, pressure indicators can be formulated. In the impact assessment phase (bottom), the midpoint or endpoint impacts, or both, that are caused by the emissions and resource use are modelled and expressed as impact indicators. Illustration by Gunilla Hagström for the Swedish University of Agricultural Sciences. End-I=endpoint indicator. LCA=lifecycle assessment. Mid-I=midpoint indicator. PI=pressure indicator.




Figure 2: Examples of Driver-Pressure-State-Impact-Response indicators and policy responses to diets

Illustration by Gunilla Hagström for the Swedish University of Agricultural Sciences, adapted from Kristensen.³⁷ DI=driver indicators. End-I=endpoint indicators. Mid-I=midpoint indicators. PI=pressure indicators. RI=response indicators. SI=state indicators.

a workshop with invited experts that covered the key earth system processes and different types of indicators (LCA-based indicators, environmental footprints, etc). This process is described more in the appendix (pp 45–46), and the results from the consultation are outlined in the following sections.

Climate change

State of the art

Many studies assessing environmental performance of diets use the carbon footprint indicator (also called GHG emissions, climate impact, or global warming impact) with global warming potential (GWP, measured as CO₂ equivalents), which is generally used to aggregate impacts from different GHGs. GWP is defined as integrated change in radiative forcing (ie, change in energy flux in the atmosphere, which determines warming) over a specific period after the emission of a certain GHG in the present-day atmosphere, relative to the same quantity of CO₂.⁵² The GWP metric requires a choice of time period over which to compare gases. This choice strongly influences the results for products with high methane emissions (eg, rice, dairy products, and ruminant meat), as methane is shorter-lived but has higher radiative efficiency than CO₂. A 100-year period (GWP₁₀₀) is used in most LCAs, international climate reporting, and many standards,^{53–55} but the use of GWP₁₀₀ for foods and diets is debatable (panel 3).

Recommendation

No metric can adequately capture all climate effects of foods, so choice of metric is heavily dependent on the study aim. For foods and diets, we recommend the use of GWP₁₀₀, but sensitivity of results to metric choice (including different GWP time periods) should be investigated (panel 3). If the choice of metric changes the overall conclusions, this must be clearly stated and discussed.

Land system change

State of the art

Although under the planetary boundaries framework, land system change focuses primarily on land affecting biogeophysical processes that regulate climate,²¹ diet assessments commonly focus on the use of agricultural land used for producing the foods in the diet (called land use, land occupation, land footprint, cropland use, nature occupation, etc). This indicator measures the amount of agricultural land, or a specific land category (eg, cropland or pasture), required during a certain period for production of foods in a diet, which is expressed as hectares×years or m²×years (table 1). This indicator shows that diets require large amounts of agricultural land, a finite natural resource.²⁸ Some studies calculate total cropland use relative to global cropland availability, to determine whether a diet is within the sustainable production capacity of cropland.^{22,50,66} Land use is relatively

Panel 2: Examples of the importance of choosing appropriate indicators

Example 1: Blue or green water consumption, or both?

Beef production uses a large amount of water. Measuring the environmental impact of this water consumption requires careful indicator selection. When the water footprint of beef is reported as total water consumption, without differentiation between blue, green, and grey water and without considering water scarcity (ie, using a total water footprint),³² estimates commonly reach 15 000 L of water per kilogram of beef produced, and for pasture-based systems, estimates reach as high as 25 000 L/kg beef produced, compared with 6000 L/kg for pig meat produced and 4000 L/kg for chicken.³⁸ A dietary recommendation based on this outlook would conclude that it is preferable to eat pork and chicken meat to beef; however, if the cattle are kept on grasslands that are not irrigated and are unsuitable for crop production for human consumption or bioenergy production, the total water footprint of beef then mostly consists of green water that has few other uses for human activities. Use of rainwater under such circumstances would, therefore, not contribute to water shortages for humans. On closer inspection, of the 15 000 L of water required for the production of 1 kg of beef, only about 4% is blue water, and the amounts of blue water required for production of pig and chicken meat are 9% and 7%, respectively. Thus, from the perspective of blue water consumption only, the production of beef requires similar amounts of water as the production of pig meat. However, high consumption of green water indicates that the beef production system captures water that could have been used by natural ecosystems.³⁹ Hence, indicators for blue and green water consumption provide different, complementary information.

Example 2: Organic comes up short when the perspective is too narrow

Organic agriculture strives to be a production system that sustains the health of soils, ecosystems, and people, and relies on ecological processes, biodiversity, and resource use that is adapted to local conditions.⁴⁰ Under this definition, organic agriculture applies a broad sustainability perspective, with restrictions on chemical inputs and a focus on management practices that enhance soil quality and biodiversity. By only assessing the most common indicators for climate change, land use, and freshwater use, studies tend to favour intensive, high-yielding agricultural systems that produce more food

per unit of land and water, as opposed to less intensive systems⁴¹ with lower greenhouse gas (GHG) emissions.⁴² Positive ecological feedback, such as regained ecosystem services, that would contribute to sustainable crop yields in the long term are generally not accounted for.⁴³ Therefore, a diet that contains a high amount of organic products, with potentially lower harmful effects on biodiversity, a smaller contribution to chemical pollution, and better soil fertility, would perform worse in an environmental assessment than a diet with more conventional foods. Conversely, if indicators for novel entities and soil health were included, a diet rich in foods from organic production systems could show more favourable results.

Example 3: Baltic Sea herring—climate-friendly and overfished

The environmental impacts from foods from aquatic animals, algae, and plants (farmed and fished, freshwater and ocean) is generally lower than for terrestrial animal products;^{44,45} however, the diversity of aquatic foods is enormous, with more than 2500 species farmed or fished globally.⁴⁶ Environmental impacts vary substantially among species and production systems. Some types of aquatic foods generally show particularly good performance with low GHG emissions and small effects with respect to other dimensions traditionally measured with lifecycle assessment. Farmed filter feeding organisms (including mussels), seaweed (algae), and wild-caught small pelagic fish (such as mackerel or anchovy) tend to have the least environmental impact in assessments. Small pelagic fish are caught with midwater trawls, requiring little fuel from fishing vessels, and therefore have a small carbon footprint per kilogram of edible weight. Clupeoids (including sardines, anchovies, and herring) are characterised by high but fluctuating recruitment rates and are among the most important commercial fish species globally.⁴⁷ Although the carbon footprint is low per kilogram of edible weight, fishing can have other severe effects. The central and western herring stocks in the Baltic Sea have been declining for a long time and are now at record low amounts.⁴⁸ Despite this, Baltic Sea herring destined for both human consumption and fish meal and oil is in high demand.⁴⁹ This example illustrates that choosing climate-friendly fish does not guarantee that the fish stock is well managed and that biodiversity is considered, indicating the need for a larger set of indicators in diet assessment.

straightforward to calculate, using data on aggregated yields of food and feed crops, which are widely available in databases such as the Food and Agriculture Organization of the UN FAOSTAT database and the Eurostat database. More sophisticated land use indicators consider factors such as land productivity in different regions⁶⁷ or geographical scarcity of land.⁶⁸ For grazing land, a variety of accounting methods are used.⁶⁹

Human appropriation of land drives biodiversity loss,⁷⁰ so land use is commonly used as a pressure

indicator for biodiversity loss and ecosystem damage.⁷¹ Endpoint impact methods capturing global biodiversity impacts from land use (eg, number of species at risk of extinction) require the amount of land used as input.^{72,73}

Recommendation

Amount of land required to produce a diet (eg, measured in m²×year) is an indicator that can be computed with reasonable accuracy on the basis of existing data, and is easy to interpret. We recommend

For more on the **FAOSTAT database** see fao.org/faostat

For more on the **Eurostat database** see ec.europa.eu/eurostat/web/main/data/database




Figure 3: Assessed planetary boundaries by indicator type
Number of studies in the review by Harrison and colleagues in 2022¹¹ assessing each planetary boundary,²¹ categorised according to Driver–Pressure–State–Impact–Response indicator type.²⁴ Some studies assessed several boundaries. Ocean acidification is not included, as it is a consequence of CO₂ emissions covered by climate change.

using this indicator to capture the resource-use aspect of land use. Different types of land (importantly, cropland and pasture), however, should be kept separate. To capture biodiversity impacts specifically related to land use, we recommend the use of endpoint indicators of actual impacts on biodiversity, such as the number of species affected.⁷² Capturing biodiversity impacts is highly complex (as discussed in the section on biosphere integrity) and substantial work is needed to improve biodiversity indicators.

Freshwater use

State of the art

Water use for diets is often assessed using pressure indicators that measure the total volume of water (in litres) required to produce foods in a diet.^{74–76} In this Personal View, we refer to consumptive water use, or water consumption (ie, the part of water withdrawn that is evaporated, transpired, incorporated into products or crops, or consumed by humans or livestock).⁷⁷ By contrast, non-consumptive water refers to the amount of water that, despite being withdrawn from the body of water, is later returned to the water system. Water consumption can be assessed as either blue water (eg, surface water and groundwater) consumption, or blue and green water (rainwater available as soil moisture) consumption for crop growth.^{78,79} Crops and grass are both major water consumers, as water is essential for plant growth.⁷⁹ Rainfed crops receive only green water, whereas irrigated crops receive both blue and green water. Grazing animals require large amounts of green water embedded in grazed biomass and other feedstuffs. Researchers have suggested global sustainable limits for both blue²² and green⁸⁰ water consumption against which diets can be benchmarked; however, preciseness of these limits is highly uncertain, particularly at a local level. In supply

chain assessments, including those for diets, blue water consumption (eg, as blue water footprint) is generally used as an indicator to avoid any double-counting of non-consumptive water uses.⁸¹

To account for environmental impacts of water consumption, indicators that compute water stress can be used;^{81,82} however, data availability on where different foods are produced can restrict computation for whole diets, as water stress is highly localised.⁸³ Average country values on water stress are available,⁸² but can obscure the spatial dimension of water stress, which can render country-specific comparisons meaningless as foods can be produced in areas of both low and high water stress within the same country.⁸⁴

Freshwater use estimations can also include a proxy for water pollution as a grey water footprint, which is calculated by how much water is needed to dilute polluted water to a threshold concentration.³² As grey water is an indicator of pollution rather than water consumption, however, it is better categorised under the planetary boundaries of biogeochemical flows and novel entities.

Recommendation

Blue and green water consumption are meaningful, complementary indicators (panel 2), but the grey water footprint is better positioned under the planetary boundary of novel entities, as it is a theoretical proxy for water pollution rather than actual water consumption. We recommend using both blue and green water consumption indicators but presenting them separately. To relate blue and green water consumption to environmental effects, indicators capturing water stress or scarcity, such as Sustainable Development Goal indicator 642 on water stress⁸¹ or available water remaining factors, can be used;⁸² however, using these indicators requires detailed data on food production area, rendering water stress or scarcity indicators less useful. Because data on blue water use are much more widely available than data on green water use,³⁹ there is a slight preference to focus on blue water consumption.

Biogeochemical flows

State of the art

Effects on biogeochemical flows are most commonly measured by pressure indicators,¹¹ such as kilograms of N and P (phosphorous) required to produce foods in a diet.⁵⁰ Such indicators (sometimes also called N and P application) commonly include only new N and P added (ie, N captured from air in synthetic fertilisers or by legumes and P mined from rock). The rationale for focusing on new N and P is that, although emissions also occur from recirculating so-called old nutrients in manure, waste, and residues, in the long term, emissions are only possible due to the continued input of new N and P to replace the nutrients lost. Therefore, although emissions of N and P do not necessarily occur at the time

Panel 3: Assessing impacts on climate change

Limitations of the GWP₁₀₀ metric had already been acknowledged when it was introduced.⁵⁶ One of the most discussed drawbacks is the metric's weighting of short-lived greenhouse gases (GHGs) such as CH₄ and long-lived GHGs such as CO₂. Although a large fraction of CO₂ stays in the atmosphere for millennia, CH₄ is naturally removed with a half-life of approximately 10 years. Due to the integral nature of global warming potential (GWP), it has a long memory of short-lived GHG emissions after their effect has decayed.⁵⁷ Alternative metrics have been developed to overcome such limitations, including global temperature change potential (GTP).⁵⁸ GTP represents the effect of an emission pulse of a certain GHG on global average surface temperature at a specified point in time after the emission relative to the same quantity of CO₂, thus limiting the so-called memory component of the metric. GTP has its own limitations, however; for example, it only considers warming at a specific time, not before or after that timepoint. For food products from ruminant animals, metric choice is important due to the associated CH₄ emissions. In 2019, Lynch and colleagues⁵⁹ showed that beef production using grassfed systems tended to create larger emissions than beef production from non-grassfed systems when using GWP₁₀₀ but the opposite when using GTP₁₀₀. However, beef remains vastly higher in climate impact than other meats or plant-based products, independent of metric choice. When using GWP or GTP, the choice of time horizon, a normative decision based on the value placed on near-term versus long-term impacts, strongly affects the results.⁶⁰

The GWP* metric has been proposed to better represent the fundamental difference in short-lived and long-lived GHGs.⁶¹ GWP* compares a one-off emission of CO₂ with a change of rate

in CH₄ emissions, which has been shown to better reflect the temperature change contribution of different gases than when these are aggregated using GWP;⁶² however, calculating the change in rate of short-lived GHG emissions for GWP* introduces new challenges in relating current emissions to emissions in the past. On a global scale, this challenge is unproblematic, but for smaller entities such as a country, farm, or product, it introduces a need to consider equity or fairness principles.⁶³ For example, if the rate of change in short-lived emissions is considered on a country level, countries with historically large CH₄ emissions would be favoured. This favouring is equivalent to grandfathering, a principle in climate ethics that is strongly criticised for being unfair.⁶⁴ In 2019, Rogelj and Schleussner⁶⁵ suggested that GWP* could be used for countries under different equity concepts (eg, by an equal per capita division of emissions or warming) but this usage also reflects moral value judgements, as with choice of time horizon for GWP and GTP. Whether and how GWP* could be meaningfully applied at product level to be useful for decision making related to foods and diets is unclear.

Another aspect to consider when comparing results from different studies is that GWP factors are regularly updated as the composition of the atmosphere changes and methods to establish such factors are improved. For example, the GWP factors in the latest Intergovernmental Panel on Climate Change Working Group 1 report⁵² are 27 for biogenic CH₄ and 273 for N₂O, compared with 28 and 265, respectively, in the previous Intergovernmental Panel on Climate Change report that was released 10 years earlier.⁶⁵

and place of new N and P inputs, the new N and P inputs are useful proxies of probable emissions.

Another pressure indicator is the N footprint, defined as total reactive N emissions per unit of consumption.^{85,86} An analogous P footprint approach exists.^{87,88} These footprint indicators have the advantage over new input of N and P in that they explicitly consider emissions of N and P regardless of whether they are old or new, and therefore more directly account for environmental pressures. Moreover, N footprints are sometimes usefully disaggregated by chemical form or location.⁸⁵ Grey water footprint has also been used as a proxy for pollution from biogeochemical flows to water ecosystems in diets.^{32,89,90}

These pressure indicators do not measure actual impacts on terrestrial and aquatic ecosystems, but are reasonable proxies for potential impacts, considering the large differences between product categories (beef, pork, chicken, root vegetables, cereals, etc).⁸⁵ Considering mined P also captures some geopolitical resource concerns about these non-substitutable nutrients.^{91,92}

Actual impacts of N and P emissions can differ widely depending on local and regional conditions, for example,

initial water eutrophication status and factors other than application of nutrients to agricultural land.⁹³ Assessing impacts from N and P use requires spatially detailed data on emissions and other environmental variables,⁹⁴ which are usually unavailable for most foods in a diet. Thus, use of impact indicators for biogeochemical flows from diets (eg, using indicators such as marine, freshwater, or terrestrial eutrophication⁹⁵ or Eutrophication potential⁹⁶) is not meaningful without data on food origin and local conditions (table 1).

Recommendation

Impact indicators that measure eutrophication, acidification, and other impacts of N and P flows are preferable, as impacts are highly influenced by local conditions; however, for these to accurately reflect impacts compared with pressure indicators, detailed data with high spatial resolution on factors such as the origin of foods and site-specific conditions are required but commonly unavailable. Simpler pressure indicators (eg, inputs of new N and P, N and P footprint) are useful to capture differences across food groups and diets, but should not be used for detailed comparisons within food groups.

Atmospheric aerosol loading

State of the art

The effect of atmospheric aerosol loading from diets is usually assessed as negative effects on human health, rather than environmental effects,^{11,97} using the LCA midpoint impact indicator of fine particulate matter (PM_{2.5}) pollution or respiratory organics (measured in kg PM_{2.5} equivalents [eq]; table 1). Acidification is also a concern classified under the planetary boundary of atmospheric aerosol loading, and is included in some studies as acidification potential (measured in kg SO₂ eq; table 1).^{98,99} To capture the creation of ground-level ozone, the indicators of photochemical ozone creation (measured in kg ethylene eq) is used, among others.⁹⁸

Recommendation

How choice of diet drives emissions of fine particulate matter is unknown, which is a reason to include it in future studies on diets. Acidification impacts vary substantially on the basis of the amount of animal products in the diet being assessed, but this aspect will be captured in carbon footprint and land use. NH₃ emissions are commonly covered by biogeochemical flow indicators. Acidification could, therefore, be excluded from environmental assessments of diets to reduce the number of indicators; however, because NH₃ emissions drive processes such as eutrophication, acidification, and fine particulate matter formation, an NH₃ emissions indicator could be warranted, provided overlaps with indicators for biogeochemical flows are avoided.

Biosphere integrity

State of the art

Biosphere integrity (ie, biodiversity) differs from other planetary boundaries in that it is an endpoint affected by all other pressures. Climate change, habitat change, and disturbances to biogeochemical flows, among other factors, all ultimately affect biosphere integrity. Lifecycle impact assessment can be used to model the aggregated effect of different pressures on the ecosystem quality endpoint impact indicator (species×years lost or potentially disappeared fraction×year), combining endpoint impact indicators for climate change, eutrophication, acidification, land use, and water consumption, for example (using methods such as the ReCiPe method,⁹⁵ LC-IMPACT,¹⁰⁰ and ImpactWorld+).¹⁰¹

Many models for individual impact categories also exist. For land use, one of the most relevant pressures, methods exist for estimating how different species are affected by land management (eg, cropping or pasture) in relation to wild habitats.^{72,73} These methods account for impacts of land transformation (eg, land clearance), land occupation (ie, continuous use of land for cropping or grazing that prevents regrowth of natural vegetation), and, in some cases, land fragmentation. Exploitation through fishing, the second largest driver of

biodiversity loss,¹⁰² can be modelled with the endpoint impact model recently developed by Helias and colleagues.³

Indicators or methods for biosphere integrity give coarse results associated with major uncertainties, due to complex cause–effect chains and spatial variation in biodiversity impacts. For example, many different taxa are affected by many pressures. The results are also affected by the normative choices made in these methods.¹⁰³ With some exceptions,^{50,104,105} biodiversity assessments are uncommon in studies of diets. Biodiversity aspects related to seafood are commonly not included—a major omission.

Despite uncertainties and limitations, methods with global coverage⁷² can be useful in highlighting biodiversity impact hotspots in diets that would be missed if focusing only on one aspect, such as climate change. In a 2020 study on Swedish diets, Moberg and colleagues⁵⁰ showed that coffee, tea, and other tropical products made relatively small contributions to the diet's overall climate impact compared with the other food products in the diet, but had large effects on biodiversity.

Coarse indicators cannot capture local biodiversity impacts. For example, the Chaudhary–Brooks⁷² method shows higher biodiversity impacts for organic coffee production due to greater land use compared with conventional production,¹⁰⁶ although studies on the actual measured biodiversity on site often show the opposite.¹⁰⁷

Biodiversity assessment methods operationalised for LCAs and footprinting of complete diets with broad geographical coverage usually only cover species richness, although overall biodiversity comprises three levels: genetic, species, and ecosystem. For some biodiversity drivers (invasive species, noise, light pollution, and overfishing), emerging methods have been applied only in a few case studies.¹⁰⁸ Meaningfully capturing such aspects of biosphere integrity at the food product or diet level might be impossible, as these aspects of biosphere integrity might be too indirect and complex to link to the consumption of specific foods.

Recommendation

Considering the range of aspects affecting biosphere integrity, methods that include many causes of biodiversity loss (eg, lifecycle impact assessment methods) are preferable, but also complex and data-intensive. Methods based on land use with global coverage are straightforward to use and can capture effects of land use in biodiversity hotspots; however, they are not able to capture the nuances of biodiversity impacts associated with different production systems. We recommend their use with the highest possible resolution. Interpretation should acknowledge uncertainties and focus on magnitude and direction of impact, rather than comparing numbers or similar products. Methods based on land use should also be complemented by methods considering biodiversity in freshwater and marine ecosystems.

Novel entities

State of the art

Few environmental assessments of diets include indicators for novel entities, probably due to a shortage of data and the complexity in using models to assess the effects caused, although food production is a major driver of issues such as chemical pollution. Indicators for chemical pollution from agriculture comprise pressure-type indicators measuring use of pesticides by kilogram of active substance applied, without considering the compound used,¹⁰⁹ and impact assessment models in LCAs evaluating the effect of toxic substances on terrestrial or aquatic ecosystems.^{110,111} Grey water footprint has also been used as a proxy for pollution to water ecosystems in diets.^{32,89,90}

Thousands of novel entities, with specific characteristics and unique effects on organisms and ecosystems, are emitted from global food production systems, and calculating ecotoxicity factors for these is data-intensive and complex.¹¹² Using pressure indicators for pesticide application only requires data on the total amount of pesticides used, not individual substances and their different factors;²⁷ however, even the use of pesticides per hectare for different crops is difficult to establish, as these data are not regularly monitored. Moreover, some pesticides are used in very small amounts but are highly toxic (eg, pyrethroids), so the amount used can be a misleading indicator.¹¹³ This misleading indicator is partly avoided by using indicators such as treatment frequency index, based on mean number of treatments weighted by the ratio of pesticide dose used to recommended dose.¹¹⁴ If indicators for chemical pollution are excluded from the environmental assessment of diets, the benefits of low pesticide use in alternative (eg, organic) farming systems are not captured, which could lead to unfair comparisons⁴¹ (panel 2).

Veterinary antibiotics are a novel entity released from livestock production and aquaculture.¹¹⁵ Data on toxicological properties are needed for estimating robust impact indicators for antibiotics.¹¹⁶ Novel entities are also emitted from other processes related to food production, including electricity use.¹¹⁷ Aggregating such effects with those from pesticide use requires impact indicators, which can be difficult to establish.

Plastic particles varying in size from large particles to nanoparticles are now ubiquitous in nature. Effects on species vary greatly, from physical effects from entanglement or ingestion to toxic impacts.¹¹⁸ Indicators for occurrence and effects of plastic contamination are emerging,^{119,120} but not currently used in assessments of diets.

Recommendation

Indicators that capture toxicity are preferable, as the environmental effect of novel entities varies on the basis of local conditions and substance. For pressure indicators such as pesticide use in kilogram of active substance,

results can be misleading because some low-dose pesticides can have high toxicity; however, we still recommend use of pesticide treatment frequency indices or application rates as indicators for novel entities, rather than omitting these completely, especially for assessments including production systems with differing pesticide use (eg, organic and conventional).

Stratospheric ozone

State of the art

Destruction of the ozone layer is seldom included in environmental assessments of diets. When included,^{71,98,121} it is often assessed using the midpoint impact indicator of ozone layer depletion potential (measured as CFC-11 eq).¹²²

Recommendation

Agriculture is currently a top contributor to ozone depletion due to N₂O emissions.¹²³ However, N₂O emissions are captured by carbon footprint (N₂O is a major GHG and contributor to global warming), so we recommend omitting ozone layer depletion from assessments of sustainable diets to limit indicator numbers.

Missing aspects: soils, blue foods, and depletion of natural resources

State of the art

Preventing further degradation of agricultural land is an urgent challenge for long-term food security,¹²⁴ but is seldom included in assessments of diets. Kraamwinkel and colleagues¹²⁵ suggest soil degradation as the tenth planetary boundary, and soil organic carbon change as an indicator for soil quality. Soil organic carbon changes have been suggested within the EU Soil Mission as one of eight indicators for soil health.¹²⁶ How different agricultural practices influence soil organic carbon is highly dependent on factors such as crops, management practices, soil, and weather conditions,¹²⁷ for which site-specific information is commonly unavailable; however, gross default factors for different crop types (eg, annual and perennial) can be used. Moberg and colleagues⁵⁰ included carbon emissions and sequestration in soil in the carbon footprint but did not report soil organic carbon changes separately as an indicator for soil quality.

The planetary boundary framework includes use of three finite natural resources (water, land, and P); however, other resources should be considered. For example, many sectors compete for renewable energy, and energy use in agriculture and aquatic production is highly variable across diets.¹²⁸

Sustainability issues specifically related to blue foods are often omitted in evaluations of diets.¹²⁹ These issues include extraction of wild species, seafloor damage, and competition for space for aquaculture.¹³⁰ Extraction of wild species through fishing should be included in assessments of environmental impacts of diets, as it

Justification		Section
Basic assessment		
Carbon footprint	Well established indicator with relatively good data availability	Climate change, panel 3
Blue water consumption	Captures trade-offs with carbon footprint, as some foods with a low carbon footprint have high blue water consumption (eg, some nuts, fruits, and vegetables); if precise production location for foods in the diet is known, indicators that consider water scarcity or stress provide complementary information	Freshwater use
Biodiversity impact from land use	Captures trade-offs with carbon footprint, as some foods with a low carbon footprint have high biodiversity impacts from land use (eg, products grown in tropical regions)	Biosphere integrity
Ecotoxicity or pesticide use	Capture differences between factors such as organic and conventional production or crops with varying pesticide intensity (eg, forage crops or legumes); impact indicators that cater for toxicity in substances should be selected if data on the substances used are available; otherwise, indicators on amounts or frequency of pesticide use can be used	Novel entities, panel 2
Exploitation of wild fish stock	Captures impact on wild stocks from fishing, which is an important cause of biodiversity loss, and can highlight trade-offs between overfishing and carbon footprint of diets	Missing aspects, panel 2
For more comprehensive assessment		
Land use	Captures use of agricultural land from a resource perspective; if no indicator for biodiversity loss from land use is used, then land use can also function as a pressure indicator of biodiversity impact	Land system change
Soil quality	Captures how the diet affects land quality in terms of soil fertility	Missing aspects
Green water consumption	Complements blue water consumption by accounting also for the rainwater needed for producing the diet; however, green water is correlated with carbon and land footprints on diet level, so green water consumption can be omitted when the number of indicators needs to be reduced	Freshwater use, panel 2
For high-ambition assessment		
New N and P input, N and P footprint, acidification, or NH ₃ emissions, or a combination of these	New N and P input or footprint act as a gross proxy of impacts on biogeochemical flows, and NH ₃ emissions can be used as a proxy somewhat more specific to acidification, air pollution, and terrestrial eutrophication	Biogeochemical flows
Energy use	Competition for renewable energy justifies keeping track of the total energy used to produce a certain diet	Missing aspects
Impacts on seafloor	Complements the land use indicator and captures an important source of damage to oceans from diets that is not captured by other indicators	Missing aspects

N=nitrogen. P=phosphorous.

Table 2: Recommendations on choice of indicators for assessing the environmental performance of diets

poses a major threat to biosphere integrity globally and exploits a natural resource (panel 2).⁷⁰ Fisheries that affect non-target species and ecosystems and the target stock can, if managed poorly, be overfished and eventually collapse. LCA indicators that capture effects of wild-caught seafood on overfishing risks and global species loss do exist,^{3,131–133} and include methods that consider the depleted stock fraction to quantify effects on ecosystem quality and that assess the maximum sustainable yield to quantify overfishing and lost

potential yield. These methods have not, however, been applied in diet assessments. Seafloor impacts related to fisheries can also be quantified using existing metrics,^{132,133} and development of additional LCA methods and indicators is ongoing. Diet assessments generally focus on terrestrial food production, but should also cover aquatic food production.

Recommendation

We encourage use of indicators for total energy use (eg, cumulative energy demand)¹³⁴ to capture the energy efficiency of diets. For wild-caught seafood, overfishing and impacts on biosphere integrity and seafloors should be considered using available methods.^{3,131–133} We also recommend considering how soils and land quality are affected by diets, but acknowledge the challenge involved considering limitations in methods, traceability, and data.

Conclusions

Our recommendations on indicator choice for diet assessment reflect three levels of ambition (table 2). Unlike Doreen and colleagues¹⁴ and Cimini and Moresi,⁹⁷ who considered indicators of GHG emissions and land use sufficient for capturing most environmental performance of diets, we believe that environmental assessments of diets should also include indicators relating to biosphere integrity, water consumption, novel entities, and exploitation of wild fish stocks, to capture important trade-offs (table 2). If more indicators can be handled in the assessment, we recommend use of indicators that capture impacts related to the quantity and quality of land use and green water consumption to complement blue water consumption. For an ambitious assessment, indicators related to biogeochemical flows, energy use, and seafloor damage can be added. This ranking does not reflect the importance of different environmental issues; for example, disturbances to biogeochemical flows are a very serious problem.²¹ However, the set of indicators we recommend can capture differences in environmental impacts across diets and highlight trade-offs between dietary choices.

Uncertainty in research assessments stemming from a shortage of representative data and variation in impacts from different production systems should be acknowledged and communicated when presenting results (eg, when comparing different diets), by better reporting ranges in primary data^{135,136} or performing sensitivity analysis on the most uncertain and impactful assumptions. Statistical inferential methods, such as hypothesis testing or confidence intervals, cannot, however, be used on simulated data.¹³⁷ Furthermore, care is needed when communicating indicator results to avoid giving an impression of high certainty (eg, reporting in many digits or ranking diets when the difference in impact is small, and to correctly communicate what the indicator shows (eg, pressure indicators should not be interpreted as reflecting the actual impact)).

The promotion of healthy and sustainable diets is urgently needed, and assessing the sustainability of diets should incorporate environmental sustainability. In this Personal View, we identify, analyse, and discuss indicator choice and provide recommendations and guidance as to which indicators to choose, and in conjunction with assessment ambition level, with the aim to unpack some of the complexity of the assessment of the environmental sustainability of diets for those carrying out assessments.

Although challenging, complex, and labour-intensive to implement correctly, inclusion of environmental dimensions in assessments of sustainable and healthy diets for a broad audience is important. We should not let perfect be the enemy of good. Instead, we need to work with what is available under given circumstances in terms of indicators and data. Furthermore, food system actors should join forces to collect more of the data needed for environmental assessments, including data on pesticide use and traceability of food items. Awareness of trade-offs in terms of accuracy and relevance is needed when choosing so-called simpler methods, and better tools for communication, interpretation, and implementation of research results need to be developed.

Contributors

YR, ER, MJ, and CC designed the study and performed the analysis. All authors interpreted results and discussed findings. YR and ER coordinated the development of the first manuscript draft. All authors commented on and edited the draft version and approved the submission of the final version.

Declaration of interests

We declare no competing interests.

Acknowledgments

This research was funded by the Swedish Environmental Protection Agency (grant number 2020-00076).

References

- Guo AN, Bryngelsson S, Strid A, Bianchi M, Winkvist A, Hallstrom E. Choice of health metrics for combined health and environmental assessment of foods and diets: a systematic review of methods. *J Clean Prod* 2022; **365**: 132622.
- Dawkins E, Kløcker Larsen R, André K, Axelsson K. Do footprint indicators support learning about sustainable consumption among Swedish public officials? *Ecol Indic* 2021; **120**: 106846.
- Hélias A, Stanford-Clark C, Bach V. A new impact pathway towards ecosystem quality in life cycle assessment: characterisation factors for fisheries. *Int J Life Cycle Assess* 2023; **28**: 367–79.
- Food and Agriculture Organization of the United Nations, WHO. Sustainable healthy diets – guiding principles. 2019. <https://www.fao.org/3/ca6640en/ca6640en.pdf> (accessed March 8, 2023).
- WWF. Footprint calculator. 2023. <https://footprint.wwf.org.uk/#/> (accessed March 8, 2023).
- Coop. Regelverk för Coops hållbarhetsdeklaration. March, 2022. https://www.coop.se/contentassets/a5d47855b7ac4f3d8f265f1ad2c06efb/coop_regelverk_for_hallbarhetsdeklaration_version_4.2_mars_2022.pdf (accessed March 8, 2023).
- Sainsbury's. Sainsbury's cuts five years from target to become Net Zero by 2035 in its own operations. Oct 26, 2021. <https://www.about.sainsburys.co.uk/news/latest-news/2021/26-10-2021-sainsburys-cuts-five-years-from-target-to-become-net-zero-by-2035-in-its-own-operations#:~:text=Sainsbury's%20has%20today%20strengthened%20its,global%20warming%20to%201.5%20degrees.> (accessed March 8, 2023).
- TESCO. Climate change. 2023. <https://www.tescopl.com/sustainability/planet/climate-change/> (accessed Feb 10, 2024).
- Pintér L, Hardi P, Martinuzzi A, Hall J. Bellagio STAMP: principles for sustainability assessment and measurement. *Ecol Indic* 2012; **17**: 20–28.
- Soulé E, Michonneau P, Michel N, Bockstaller C. Environmental sustainability assessment in agricultural systems: a conceptual and methodological review. *J Clean Prod* 2021; **325**: 129291.
- Harrison MR, Palma G, Buendia T, Bueno-Tarodo M, Quell D, Hachem F. A scoping review of indicators for sustainable healthy diets. *Front Sustain Food Syst* 2022; **5**: 822263.
- Jones AD, Hoey L, Blesh J, Miller L, Green A, Shapiro LF. A systematic review of the measurement of sustainable diets. *Adv Nutr* 2016; **7**: 641–64.
- Aldaya MM, Ibañez FC, Domínguez-Lacueva P, et al. Indicators and recommendations for assessing sustainable healthy diets. *Foods* 2021; **10**: 999.
- van Dooren C, Aiking H, Vellinga P. In search of indicators to assess the environmental impact of diets. *Int J Life Cycle Assess* 2018; **23**: 1297–314.
- Manfredi S, Allacker K, Chomkhamrui K, Pelletier N, de Souza DM. Product environmental footprint (PEF) guide. Ispra: European Commission Joint Research Centre, 2012.
- Biesbroek S, Kok FJ, Tufford AR, et al. Toward healthy and sustainable diets for the 21st century: Importance of sociocultural and economic considerations. *Proc Natl Acad Sci USA* 2023; **120**: e2219272120.
- Bôto JM, Rocha A, Miguéis V, Meireles M, Neto B. Sustainability dimensions of the Mediterranean diet: a systematic review of the indicators used and its results. *Adv Nutr* 2022; **13**: 2015–38.
- Eme PE, Douwes J, Kim N, Foliaki S, Burlingame B. Review of methodologies for assessing sustainable diets and potential for development of harmonised indicators. *Int J Environ Res Public Health* 2019; **16**: 1184.
- Forbes S, Bicknell E, Guilovica L, Wingrove K, Charlton K. A rapid review of the environmental impacts associated with food consumption in Australia and New Zealand. *Curr Nutr Rep* 2021; **10**: 334–51.
- Hatjithanassiadou M, Rolim PM, Seabra LMAJ. Nutrition and its footprints: using environmental indicators to assess the nexus between sustainability and food. *Front Sustain Food Syst* 2023; **6**: 1078997.
- Steffen W, Richardson K, Rockström J, et al. Sustainability. Planetary boundaries: guiding human development on a changing planet. *Science* 2015; **347**: 1259855.
- Willett W, Rockström J, Loken B, et al. Food in the Anthropocene: the EAT–Lancet Commission on healthy diets from sustainable food systems. *Lancet* 2019; **393**: 447–92.
- Vanham D, Gawlik BM, Bidoglio G. Cities as hotspots of indirect water consumption: the case study of Hong Kong. *J Hydrol* 2019; **573**: 1075–86.
- Smeets E, Weterings R. Environmental indicators: typology and overview. Sept 7, 1999. <https://www.eea.europa.eu/publications/TEC25> (accessed Feb 10, 2024).
- Carnohan SA, Trier X, Liu S, et al. Next generation application of DPSIR for sustainable policy implementation. *Curr Res Environ Sustain* 2023; **5**: 100201.
- Hauschild MZ, Huijbregts MAJ. Life cycle impact assessment. Dordrecht: Springer, 2015.
- Moberg E, Sall S, Hansson PA, Roos E. Taxing food consumption to reduce environmental impacts-identification of synergies and goal conflicts. *Food Policy* 2021; **101**: 102090.
- Sonderegger T, Dewulf J, Fantke P, et al. Towards harmonizing natural resources as an area of protection in life cycle impact assessment. *Int J Life Cycle Assess* 2017; **22**: 1912–27.
- International Organization for Standardization. ISO 14044:2006: environmental management: life cycle assessment: requirements and guidelines. Geneva: International Organization for Standardization, 2006.
- International Organization for Standardization. ISO 14040:2006: environmental management: life cycle assessment: principles and framework. Geneva: International Organization for Standardization, 2006.
- Turner GM, Larsen KA, Candy S, et al. Squandering Australia's food security—the environmental and economic costs of our unhealthy diet and the policy Path We're On. *J Clean Prod* 2018; **195**: 1581–99.

- 32 Hoekstra AY, Chapagain AK, Aldaya MM, Mekonnen MM. The water footprint assessment manual: setting the global standard. London: Earthscan, 2011.
- 33 Ridoutt BG, Pfister S. A revised approach to water footprinting to make transparent the impacts of consumption and production on global freshwater scarcity. *Glob Environ Change* 2010; **20**: 113–20.
- 34 Rockström J, Steffen W, Noone K, et al. A safe operating space for humanity. *Nature* 2009; **461**: 472–75.
- 35 Biermann F, Kim RE. The boundaries of the planetary boundary framework: a critical appraisal of approaches to define a “safe operating space” for humanity. *Annu Rev Environ Resour* 2020; **45**: 497–521.
- 36 Vanham D, Leip A, Galli A, et al. Environmental footprint family to address local to planetary sustainability and deliver on the SDGs. *Sci Total Environ* 2019; **693**: 133642.
- 37 Kristensen P. The DPSIR Framework. 2004. [http://fis.freshwatertools.eu/files/MARS_resources/Info_lib/Kristensen\(2004\)DPSIR%20Framework.pdf](http://fis.freshwatertools.eu/files/MARS_resources/Info_lib/Kristensen(2004)DPSIR%20Framework.pdf) (accessed Feb 10, 2024).
- 38 Mekonnen MM, Hoekstra AY. A global assessment of the water footprint of farm animal products. *Ecosystems* 2012; **15**: 401–15.
- 39 Schyns JF, Hoekstra AY, Booij MJ, Hogeboom RJ, Mekonnen MM. Limits to the world’s green water resources for food, feed, fiber, timber, and bioenergy. *Proc Natl Acad Sci USA* 2019; **116**: 4893–98.
- 40 IFOAM Organics International. Definition of organic agriculture. <https://www.ifoam.bio/why-organic/organic-landmarks/definition-organic> (accessed April 25, 2023).
- 41 van der Werf HMG, Knudsen MT, Cederberg C. Towards better representation of organic agriculture in life cycle assessment. *Nat Sustain* 2020; **3**: 419–25.
- 42 Searchinger TD, Wiersma S, Beringer T, Dumas P. Assessing the efficiency of changes in land use for mitigating climate change. *Nature* 2018; **564**: 249–53.
- 43 Schneider K, Barreiro-Hurle J, Rodriguez-Cerezo E. Pesticide reduction amidst food and feed security concerns in Europe. *Nat Food* 2023; **4**: 746–50.
- 44 Bianchi M, Hallström E, Parker RWR, Mifflin K, Tyedmers P, Ziegler F. Assessing seafood nutritional diversity together with climate impacts informs more comprehensive dietary advice. *Commun Earth Environ* 2022; **3**: 188.
- 45 Gephart JA, Henriksson PJG, Parker RWR, et al. Environmental performance of blue foods. *Nature* 2021; **597**: 360–65.
- 46 Food and Agriculture Organization of the UN. Global production by production source quantity (1950–2021). https://www.fao.org/fishery/statistics-query/en/global_production/global_production_quantity (accessed Feb 10, 2024).
- 47 Food and Agriculture Organization of the UN. The state of world fisheries and aquaculture 2022: towards blue transformation. Rome: Food and Agriculture Organization of the UN, 2022.
- 48 International Council for the Exploration of the Sea. Baltic fisheries assessment working group. 2021. <https://www.ices.dk/community/groups/pages/wgbfas.aspx> (accessed Feb 10, 2024).
- 49 Pihlajamäki M, Sarkki S, Karjalainen TP. Food or feed? The contribution of Baltic herring fisheries to food security and safety. In: Olsson AIS, Araújo SM, Vieira FM, eds. Food futures: ethics, science and culture. Wageningen: Wageningen Academic Publishers, 2016: 239–43.
- 50 Moberg E, Karlsson Potter H, Wood A, Hansson P-A, Rös E. Benchmarking the Swedish diet relative to global and national environmental targets—identification of indicator limitations and data gaps. *Sustainability* 2020; **12**: 1407.
- 51 Galway LP, Acharya Y, Jones AD. Deforestation and child diet diversity: a geospatial analysis of 15 sub-Saharan African countries. *Health Place* 2018; **51**: 78–88.
- 52 Intergovernmental Panel on Climate Change. Climate change 2021: the physical science basis. 2021. <https://www.ipcc.ch/report/ar6/wg1/> (accessed March 4, 2023).
- 53 International Organization for Standardization. ISO/TS 14067:2013. Geneva: International Organization for Standardization, 2013.
- 54 UN Environment Programme, Life Cycle Initiative. Global guidance for life cycle impact assessment indicators. Jan 14, 2017. <https://www.lifecycleinitiative.org/training-resources/global-guidance-lcia-indicators-v-1/> (accessed Feb 10, 2024).
- 55 The European Commission. Commission recommendation (EU) 2021/2279 of 15 December 2021 on the use of the Environmental Footprint methods to measure and communicate the life cycle environmental performance of products and organisations. Brussels: European Commission, 2021.
- 56 Houghton JT, Jenkins GJ, Ephraums JJ. Climate change, the IPCC scientific assessment. Cambridge: Cambridge University Press, 1990.
- 57 Shine KP. The global warming potential—the need for an interdisciplinary retrieval. *Clim Change* 2009; **96**: 467–72.
- 58 Shine KP, Fuglestad JS, Hailemariam K, Stuber N. Alternatives to the Global Warming Potential for comparing climate impacts of emissions of greenhouse gases. *Clim Change* 2005; **68**: 281–302.
- 59 Lynch J. Availability of disaggregated greenhouse gas emissions from beef cattle production: a systematic review. *Environ Impact Assess Rev* 2019; **76**: 69–78.
- 60 Persson UM, Johansson DJA, Cederberg C, Hedenus F, Bryngelsson D. Climate metrics and the carbon footprint of livestock products: where’s the beef? *Environ Res Lett* 2015; **10**: 034005.
- 61 Allen MR, Fuglestad JS, Shine KP, Reisinger A, Pierrehumbert RT, Forster PM. New use of global warming potentials to compare cumulative and short-lived climate pollutants. *Nat Clim Change* 2016; **6**: 773–76.
- 62 Cain M, Lynch J, Allen MR, Fuglestad JS, Frame DJ, Macey AH. Improved calculation of warming-equivalent emissions for short-lived climate pollutants. *NPJ Clim Atmos Sci* 2019; **2**: 29.
- 63 Rogelj J, Schleussner C-F. Unintentional unfairness when applying new greenhouse gas emissions metrics at country level. *Environ Res Lett* 2019; **14**: 114039.
- 64 Caney S. Justice and the distribution of greenhouse gas emissions. *J Glob Ethics* 2009; **5**: 125–46.
- 65 Intergovernmental Panel on Climate Change. Climate Change 2013: the physical science basis. Cambridge: Cambridge University Press, 2013.
- 66 Springmann M, Clark M, Mason-D’Croz D, et al. Options for keeping the food system within environmental limits. *Nature* 2018; **562**: 519–25.
- 67 Scherer L, Behrens P, Tukker A. Opportunity for a dietary win-win in nutrition, environment, and animal welfare. *One Earth* 2019; **1**: 349–60.
- 68 Font Vivanco D, Sprecher B, Hertwich E. Scarcity-weighted global land and metal footprints. *Ecol Indic* 2017; **83**: 323–27.
- 69 Vanham D, Bruckner M, Schwarzmüller F, Schyns J, Kastner T. Multi-model assessment identifies livestock grazing as a major contributor to variation in European Union land and water footprints. *Nat Food* 2023; **4**: 575–84.
- 70 Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. Global assessment report on biodiversity and ecosystem services. Bonn: IPBES, 2019.
- 71 Saxe H. The New Nordic Diet is an effective tool in environmental protection: it reduces the associated socioeconomic cost of diets. *Am J Clin Nutr* 2014; **99**: 1117–25.
- 72 Chaudhary A, Brooks TM. Land use intensity-specific global characterization factors to assess product biodiversity footprints. *Environ Sci Technol* 2018; **52**: 5094–104.
- 73 Kuipers KJJ, May R, Veronesi F. Considering habitat conversion and fragmentation in characterisation factors for land-use impacts on vertebrate species richness. *Sci Total Environ* 2021; **801**: 149737.
- 74 Aleksandrowicz L, Green R, Joy EJM, et al. Environmental impacts of dietary shifts in India: a modelling study using nationally-representative data. *Environ Int* 2019; **126**: 207–15.
- 75 Hoekstra AY. The water footprint: the relation between human consumption and water use. In: Antonelli M, Greco F, eds. The water we eat: combining virtual water and water footprints. Cham: Springer, 2015: 35–48.
- 76 Mekonnen MM, Gerbens-Leenes W. The water footprint of global food production. *Water* 2020; **12**: 2696.
- 77 Falkenmark M, Lannerstad M. Consumptive water use to feed humanity - curing a blind spot. *Hydrol Earth Syst Sci* 2005; **9**: 15–28.
- 78 Food and Agriculture Organization of the UN. Land-water linkages: a synopsis. 1995. <https://www.fao.org/3/v5400e/v5400e06.htm#land%20water%20linkages:%20a%20synopsis> (accessed Feb 10, 2024).

- 79 Falkenmark M, Rockstrom J. The new blue and green water paradigm: breaking new ground for water resources planning and management. *J Water Resour Plan Manage* 2006; **132**: 129–32.
- 80 Wang-Erlandsson L, Tobian A, van der Ent RJ, et al. A planetary boundary for green water. *Nat Rev Earth Environ* 2022; **3**: 380–92.
- 81 Vanham D, Hoekstra AY, Wada Y, et al. Physical water scarcity metrics for monitoring progress towards SDG target 6.4: an evaluation of indicator 6.4.2 “Level of water stress”. *Sci Total Environ* 2018; **613–614**: 218–32.
- 82 Boulay A-M, Bare J, Benini L, et al. The WULCA consensus characterization model for water scarcity footprints: assessing impacts of water consumption based on available water remaining (AWARE). *Int J Life Cycle Assess* 2018; **23**: 368–78.
- 83 Vanham D, Alfieri L, Flörke M, et al. The number of people exposed to water stress in relation to how much water is reserved for the environment: a global modelling study. *Lancet Planet Health* 2021; **5**: e766–74.
- 84 Vanham D. Water resources for sustainable healthy diets: state of the art and outlook. *Water* 2020; **12**: 3224.
- 85 Einarsson R, Cederberg C. Is the nitrogen footprint fit for purpose? An assessment of models and proposed uses. *J Environ Manage* 2019; **240**: 198–208.
- 86 Leach K, Gerrard CL, Kudahl AB, et al. Assessing the sustainability of EU dairy farms with different management systems and husbandry practices. September 14, 2012. <https://orgrprints.org/id/eprint/21761/7/21761.pdf> (accessed Feb 10, 2024).
- 87 Metson GS, MacDonald GK, Leach AM, Compton JE, Harrison JA, Galloway JN. The U.S. consumer phosphorus footprint: where do nitrogen and phosphorus diverge? *Environ Res Lett* 2020; **15**: 1–15.
- 88 Oita A, Wirasenjaya F, Liu J, Webeck E, Matsubae K. Trends in the food nitrogen and phosphorus footprints for Asia’s giants: China, India, and Japan. *Resour Conserv Recycling* 2020; **157**: 104752.
- 89 Blas A, Garrido A, Unver O, Willaarts B. A comparison of the Mediterranean diet and current food consumption patterns in Spain from a nutritional and water perspective. *Sci Total Environ* 2019; **664**: 1020–29.
- 90 Blas A, Garrido A, Willaarts BA. Evaluating the water footprint of the Mediterranean and American diets. *Water* 2016; **8**: 448.
- 91 Brownlie WJ, Spears BM, Heal KV, et al. Towards our phosphorus future. In: Brownlie WJ, Sutton MA, Heal KV, Reay DS, Spears BM, eds. *Our phosphorous future*. Edinburgh: UK Centre for Ecology & Hydrology, 2022: 339–69.
- 92 Sutton MA, Reis S, Riddick SN, et al. Towards a climate-dependent paradigm of ammonia emission and deposition. *Philos Trans R Soc Lond B Biol Sci* 2013; **368**: 20130166.
- 93 Henryson K, Kätterer T, Tidåker P, Sundberg C. Soil N₂O emissions, N leaching and marine eutrophication in life cycle assessment - a comparison of modelling approaches. *Sci Total Environ* 2020; **725**: 138332.
- 94 Henryson K, Hansson P-A, Sundberg C. Spatially differentiated midpoint indicator for marine eutrophication of waterborne emissions in Sweden. *Int J Life Cycle Assess* 2018; **23**: 70–81.
- 95 Huijbregts MAJ, Steinmann ZJN, Elshout PMF, et al. ReCiPe2016: a harmonised life cycle impact assessment method at midpoint and endpoint level. *Int J Life Cycle Assess* 2017; **22**: 138–47.
- 96 Bruijn H, Duin R, Huijbregts MAJ, et al. Handbook on life cycle assessment: operational guide to the ISO standards. Dordrecht: Kluwer Academic Publishers, 2002.
- 97 Cimini A, Moresi M. Are the present standard methods effectively useful to mitigate the environmental impact of the 99% EU food and drink enterprises? *Trends Food Sci Technol* 2018; **77**: 42–53.
- 98 Davis J, Sonesson UG, Baumgartner DU, Nemecek T. Environmental impact of four meals with different protein sources: case studies in Spain and Sweden. *Food Res Int* 2010; **43**: 1874–84.
- 99 Adhikari B, Prapasongsa T. Environmental sustainability of food consumption in Asia. *Sustainability* 2019; **11**: 3749.
- 100 Verones F, Hellweg S, Antón A, et al. LC-IMPACT: a regionalized life cycle damage assessment method. *J Ind Ecol* 2020; **24**: 1201–19.
- 101 Bulle C, Margni M, Patouillard L, et al. IMPACT World+: a globally regionalized life cycle impact assessment method. *Int J Life Cycle Assess* 2019; **24**: 1653–74.
- 102 Jaureguiberry MS, Venturino A. Nutritional and environmental contributions to autism spectrum disorders: focus on nutrigenomics as complementary therapy. *Int J Vitam Nutr Res* 2022; **92**: 248–66.
- 103 Crenna E, Marques A, La Notte A, Sala S. Biodiversity assessment of value chains: state of the art and emerging challenges. *Environ Sci Technol* 2020; **54**: 9715–28.
- 104 Portilho F, Castañeda M, de Castro IRR. [Food in the contemporary context: consumption, political action and sustainability]. *Cien Saude Colet* 2011; **16**: 99–106 (in Portuguese).
- 105 Rös E, Karlsson HK, Withöft CM, Sundberg C. Evaluating the sustainability of diets—combining environmental and nutritional aspects. *Environ Sci Policy* 2015; **47**: 157–66.
- 106 Eneroth H, Karlsson Potter K, Rös E. Environmental effects of coffee, tea and cocoa - data collection for a consumer guide for plant-based foods. Uppsala: Department of Energy and Technology, Swedish University of Agricultural Sciences, 2022.
- 107 Gomiero T, Pimentel D, Paoletti MG. Environmental impact of different agricultural management practices: conventional vs. organic agriculture. *Crit Rev Plant Sci* 2011; **30**: 95–124.
- 108 Crenna E, Sinkko T, Sala S. Biodiversity impacts due to food consumption in Europe. *J Clean Prod* 2019; **227**: 378–91.
- 109 Marlow HJ, Harwatt H, Soret S, Sabaté J. Comparing the water, energy, pesticide and fertilizer usage for the production of foods consumed by different dietary types in California. *Public Health Nutr* 2015; **18**: 2425–32.
- 110 Fantke PE, Bijster M, Guignard C, et al. USEtox 2.0 documentation (version 1.1). January, 2017. <https://usetox.org/model/documentation> (accessed March 3, 2023).
- 111 Sala S, Biganzoli F, Mengual ES, Saouter E. Toxicity impacts in the environmental footprint method: calculation principles. *Int J Life Cycle Assess* 2022; **27**: 587–602.
- 112 Gentil C, Fantke P, Mottes C, Basset-Mens C. Challenges and ways forward in pesticide emission and toxicity characterization modeling for tropical conditions. *Int J Life Cycle Assess* 2020; **25**: 1290–306.
- 113 Nordborg M, Cederberg C, Berndes G. Modeling potential freshwater ecotoxicity impacts due to pesticide use in biofuel feedstock production: the cases of maize, rapeseed, salix, soybean, sugar cane, and wheat. *Environ Sci Technol* 2014; **48**: 11379–88.
- 114 Pelosi C, Toutout L, Chiron F, et al. Reduction of pesticide use can increase earthworm populations in wheat crops in a European temperate region. *Agric Ecosyst Environ* 2013; **181**: 223–30.
- 115 Persson L, Carney Almroth BM, Collins CD, et al. Outside the safe operating space of the planetary boundary for novel entities. *Environ Sci Technol* 2022; **56**: 1510–21.
- 116 Nyberg O, Rico A, Guinée JB, Henriksson PJG. Characterizing antibiotics in LCA—a review of current practices and proposed novel approaches for including resistance. *Int J Life Cycle Assess* 2021; **26**: 1816–31.
- 117 Zira S, Rös E, Ivarsson E, Hoffmann R, Rydhmer L. Social life cycle assessment of Swedish organic and conventional pork production. *Int J Life Cycle Assess* 2020; **25**: 1957–75.
- 118 Woods JS, Verones F, Joliet O, Vázquez-Rowe I, Boulay A-M. A framework for the assessment of marine litter impacts in life cycle impact assessment. *Ecol Indic* 2021; **129**: 107918.
- 119 Høiøberg MA, Woods JS, Verones F. Global distribution of potential impact hotspots for marine plastic debris entanglement. *Ecol Indic* 2022; **135**: 108509.
- 120 Corella-Puertas E, Guieu P, Aufoujal A, Bulle C, Boulay A-M. Development of simplified characterization factors for the assessment of expanded polystyrene and tire wear microplastic emissions applied in a food container life cycle assessment. *J Ind Ecol* 2022; **26**: 1882–94.
- 121 Jensen JD, Saxe H, Denver S. Cost-effectiveness of a new nordic diet as a strategy for health promotion. *Int J Environ Res Public Health* 2015; **12**: 7370–91.
- 122 Farinha C, de Brito J, Veiga MD. Life cycle assessment. In: Farinha C, Jankovic J, Veiga MD, eds. *Eco-efficient rendering mortars: use of recycled materials*. Cambridge: Woodhead Publishing, 2021: 205–34.
- 123 Kanter DR, Wagner-Riddle C, Groffman PM, et al. Improving the social cost of nitrous oxide. *Nat Clim Chang* 2021; **11**: 1008–10.

- 124 Intergovernmental Panel on Climate Change. Climate Change and land. 2019. <https://www.ipcc.ch/srccl/> (accessed Feb 10, 2024).
- 125 Kraamwinkel CT, Beaulieu A, Dias T, Howison RA. Planetary limits to soil degradation. *Commun Earth Environ* 2021; 2: 249.
- 126 European Commission. Caring for soils is caring for life. Sept 22, 2020. https://research-and-innovation.ec.europa.eu/knowledge-publications-tools-and-data/publications/all-publications/caring-soil-caring-life_en (accessed Feb 10, 2024).
- 127 Basile-Doelsch I, Balesdent J, Pellerin S. Reviews and syntheses: the mechanisms underlying carbon storage in soil. *Biogeosciences* 2020; 17: 5223–42.
- 128 Meier T, Christen O. Environmental impacts of dietary recommendations and dietary styles: Germany as an example. *Environ Sci Technol* 2013; 47: 877–88.
- 129 Bogard JR, Farmery AK, Little DC, Fulton EA, Cook M. Will fish be part of future healthy and sustainable diets? *Lancet Planet Health* 2019; 3: e159–60.
- 130 Sanchez-Jerez P, Karakassis I, Massa F, et al. Aquaculture's struggle for space: the need for coastal spatial planning and the potential benefits of Allocated Zones for Aquaculture (AZAs) to avoid conflict and promote sustainability. *Aquacult Environ Interact* 2016; 8: 41–54.
- 131 Emanuelsson A, Ziegler F, Pihl L, Sköld M, Sonesson U. Accounting for overfishing in life cycle assessment: new impact categories for biotic resource use. *Int J Life Cycle Assess* 2014; 19: 1156–68.
- 132 Woods JS, Verones F. Ecosystem damage from anthropogenic seabed disturbance: a life cycle impact assessment characterisation model. *Sci Total Environ* 2019; 649: 1481–90.
- 133 Ziegler F, Nilsson P, Mattsson B, Walther Y. Life cycle assessment of frozen cod fillets including fishery-specific environmental impacts. *Int J Life Cycle Assess* 2003; 8: 39–47.
- 134 Frischknecht R, Wyss F, Büsser Knöpfel S, Lützkendorf T, Balouktsi M. Cumulative energy demand in LCA: the energy harvested approach. *Int J Life Cycle Assess* 2015; 20: 957–69.
- 135 Henriksson PJG, Guinée JB, Heijungs R, de Koning A, Green DM. A protocol for horizontal averaging of unit process data—including estimates for uncertainty. *Int J Life Cycle Assess* 2014; 19: 429–36.
- 136 Ruett J, Hennes L, Teubler J, Braun B. How compatible are western European dietary patterns to climate targets? Accounting for uncertainty of life cycle assessments by applying a probabilistic approach. *Sustainability* 2022; 14: 14449.
- 137 von Brömssen C, Röös E. Why statistical testing and confidence intervals should not be used in comparative life cycle assessments based on Monte Carlo simulations. *Int J Life Cycle Assess* 2020; 25: 2101–05.

Copyright © 2024 The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY 4.0 license.