



CHALMERS
UNIVERSITY OF TECHNOLOGY

Fraïssé theory for Cuntz semigroups

Downloaded from: <https://research.chalmers.se>, 2026-04-21 07:59 UTC

Citation for the original published paper (version of record):

Cantier, L., Vilalta Vila, E. (2024). Fraïssé theory for Cuntz semigroups. *Journal of Algebra*, 658: 319-364. <http://dx.doi.org/10.1016/j.jalgebra.2024.05.052>

N.B. When citing this work, cite the original published paper.



Contents lists available at ScienceDirect

Journal of Algebra

journal homepage: www.elsevier.com/locate/jalgebra



Research Paper

Fraïssé theory for Cuntz semigroups [☆]

Laurent Cantier ^{a,b,*}, Eduard Vilalta ^c



^a *Departament de Matemàtiques, Universitat Autònoma de Barcelona, 08193 Bellaterra, Spain*

^b *Institute of Mathematics, Czech Academy of Sciences, Žitná 25, 115 67 Praha 1, Czechia*

^c *Department of Mathematical Sciences, Chalmers University of Technology and University of Gothenburg, Chalmers Tvärgata 3, SE-412 96 Gothenburg, Sweden*

ARTICLE INFO

Article history:

Received 18 March 2024

Available online 19 June 2024

Communicated by Volodymyr Mazorchuk

Keywords:

Fraïssé theory

Cuntz semigroup

Cauchy sequences

Cu-distance

ABSTRACT

We develop a theory of Cauchy sequences and intertwining morphisms for Cuntz semigroups, which generalizes all past approaches to study metric-like properties of the invariant. Further, the techniques presented here can be applied to all known refinements of the Cuntz semigroup, including those that may be used in new classification results.

As a particular application, we introduce a Fraïssé theory for abstract Cuntz semigroups akin to the theory of Fraïssé categories developed by Kubiś. We also show that any (Cuntz) Fraïssé category has a unique Fraïssé limit which is both universal and homogeneous. Several examples of such categories and their Fraïssé limits are given throughout the paper.

© 2025 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

[☆] Both authors were partially supported by MINECO (grant No. PID2020-113047GB-I00), and by the Departament de Recerca i Universitats de la Generalitat de Catalunya (grant No. 2021-SGR-01015). The first author was also supported by the Spanish Ministry of Universities and the European Union-NextGenerationEU through a Margarita Salas grant. The second author was also supported by MINECO grant No. PRE2018-083419 and by the Fields Institute for Research in Mathematical Sciences.

* Corresponding author.

E-mail addresses: laurent.cantier@uab.cat (L. Cantier), vilalta@chalmers.se (E. Vilalta).

URLs: <https://www.laurencantier.fr> (L. Cantier), <https://www.eduardvilalta.com> (E. Vilalta).

1. Introduction

Fraïssé Theory was introduced in [29] by Fraïssé in the context of model theory with the intent of giving a generic method to construct countable homogeneous structures from their finitely-generated substructures. Since then, several adaptations of this method have appeared. These include, but are not limited to, projective Fraïssé theory ([35]), Fraïssé theory for metric structures ([60]) and, of late, Fraïssé categories ([32,40]). The overall idea is to build a ‘large object’, called the *Fraïssé limit*, which is unique, universal, and homogeneous with respect to a class of ‘small sub-objects’, known as the *Fraïssé class*. All these variations of Fraïssé theory have produced interesting results in their corresponding areas. For example, well-known topological spaces such as the Cantor set, the pseudo-arc, the Lelek fan or the Menger curve have been expressed as (projective) Fraïssé limits. (See [9,10,35,40].) Similarly, as shown in [40], the Gurarii space is the Fraïssé limit of the class of finite dimensional Banach spaces. In graph theory, the countable random graph also arises as the Fraïssé limit of the class of finite graphs and, in model theory, Ehrenfeucht-Fraïssé-like games can be built using Fraïssé classes.

Recently, Fraïssé theory has been applied to the field of C^* -algebras (for example, in [25,36,42,56]) and, in that setting, many well known C^* -algebras have been constructed as Fraïssé limits. As proof, the Jiang-Su algebra \mathcal{Z} —introduced in [37], and which plays a central role in the classification of simple C^* -algebras— can be seen as a Fraïssé limit ([42]). Further, as noted in [31], this construction can be used to (re)prove in simpler ways some of the properties of the algebra, such as its strong self-absorption. Among other examples, several stably projectionless C^* -algebras were also built as Fraïssé limits in [36], and the existence of a universal AF-algebra was proved in [32].

In light of the recent discoveries connecting Fraïssé theory and C^* -algebras, it is high time to explore Fraïssé categories of (abstract) Cuntz semigroups. First introduced by Cuntz in [24], the *Cuntz semigroup* is a powerful invariant for C^* -algebras that codifies how positive elements are compared. In [22], Coward, Elliott and Ivanescu introduced the category Cu of *abstract Cuntz semigroups*, or *Cu-semigroups* for short. This rich subcategory of positively ordered monoids has been studied extensively (see, among many others, [2,6–8,13,14,21]) and has yielded new results for C^* -algebras, such as the ones obtained in [3,17,18,26,43,51,53,55,59]. Further, the category Cu allows one to view the Cuntz semigroup as a continuous functor for C^* -algebras.

The aim of this paper is twofold: To develop a general theory of metric-like properties for (abstract) Cuntz semigroups, and to use such results to introduce a Fraïssé theory in the category Cu , while also giving examples and studying its relations to its C^* -algebraic counterpart.

The first obstruction that one finds when trying to mimic the past approaches is the general lack of a non-trivial, enriched distance in the Hom-sets of Cu . Although such a distance does exist for specific Cu -semigroups (and has been exploited successfully in a number of situations; see [17–19,43,46]), this approach is still too restrictive for our purposes. Instead, we will compare morphisms in Cu by using *finite-set comparison*, an

idea that had appeared implicitly in the past (e.g. [19,46]) but that was first given a name in [15].

Using this notion of comparison, we introduce *Cauchy sequences* of morphisms (Definition 3.4) and show that all such sequences converge.

Theorem (3.8). *Let S, T be Cu-semigroups. Then any Cauchy sequence $(\alpha_i)_i$ in $\text{Hom}_{\text{Cu}}(S, T)$ converges towards a unique Cu-morphism.*

Further, we also develop a comprehensive theory of *intertwinings* in Cu; see Definitions 3.10/3.13.

Theorem (3.17). *Let $(S_i, \sigma_{i,j})_{i \in \mathbb{N}}$ and $(T_i, \tau_{i,j})_{i \in \mathbb{N}}$ be two inductive sequences in Cu with respective limits $(S, \sigma_{i,\infty})$ and $(T, \tau_{i,\infty})$. Assume that there exists a two-sided approximate intertwining $(\alpha_i: S_i \rightarrow T_{\varphi(i)}, \beta_i: T_i \rightarrow S_{\psi(i)})_i$.*

Then there exists a Cu-isomorphism $\alpha: S \cong T$ induced by $(\alpha_i)_i$ whose inverse is induced by $(\beta_i)_i$.

Both of the aforementioned theorems generalize all previous results on the subject and provide a unified picture of the metric-like structure that Cuntz semigroups enjoy. Moreover, as noted in Paragraph 3.18, the ideas and techniques developed here can be applied to a vast family of generalizations of the Cuntz semigroup (including, in particular, all of its refinements considered in the past). We predict that these metric flavoured statements will play a key role in future classification and structure results for possibly non-simple C^* -algebras.

With all these tools at our disposal, we are able to develop a *Fraïssé theory* for Cu-semigroups in Part B of Section 3 and obtain the following theorem.

Theorem (3.27). *Let $\mathfrak{c} \subseteq \text{Cu}$ be a Fraïssé category of Cu-semigroups. Then any Fraïssé sequence $(S_i, \sigma_{i,j})_i$ has a $\bar{\mathfrak{c}}$ -limit $(S, \sigma_{i,\infty})_i$ such that*

- (i) *S is unique up to isomorphism, that is, S does not depend on the Fraïssé sequence chosen.*
- (ii) *The set $\text{Hom}_{\bar{\mathfrak{c}}}(D, S)$ is nonempty whenever D is countably-based.*

Further, assume that \mathfrak{c} is contained in a category \mathfrak{d} where every \mathfrak{c} -inductive sequence has a limit, and where every \mathfrak{d} -object is the limit of a \mathfrak{c} -sequence. If $\mathfrak{c} \subseteq \mathfrak{d}$ satisfies the almost factorization property, then

- (iii) *For any $C \in \mathfrak{c}$, any $\alpha, \beta \in \text{Hom}_{\mathfrak{d}}(C, S)$ and any finite set $F \subseteq C$, there exists a \mathfrak{d} -isomorphism $\eta_F: S \xrightarrow{\cong} S$ such that $\eta_F \circ \alpha \simeq_F \beta$.*

In Part C of Section 3 we study the relations between this theorem and the Fraïssé theory of C^* -algebras, while in Section 4 we provide a number of examples, listed below:

- (1) Any dimension Cu-semigroup S_p of infinite type is the Fraïssé limit of the Fraïssé category \mathfrak{s}_p .
- (2) There exists a universal dimension Cu-semigroup \mathcal{S} which is the Fraïssé limit of the Fraïssé category \mathfrak{s}_{\dim} .
- (3) There exist countably many Fraïssé categories \mathfrak{e}_n whose Fraïssé limits are simple, non-stably finite, not weakly purely infinite Cu-semigroups.
- (4) The Cu-semigroup $\text{Lsc}(2^{\mathbb{N}}, \overline{\mathbb{N}})$, where $2^{\mathbb{N}}$ denotes the Cantor set, is the Fraïssé limit of the Fraïssé category $\mathcal{K}_{2^{\mathbb{N}}}$.
- (5) The Cu-semigroup $\text{Lsc}(\mathbb{P}, \overline{\mathbb{N}})$, where \mathbb{P} denotes the pseudo-arc, is the Fraïssé limit of the Fraïssé category $\mathcal{K}_{\mathbb{P}}$.
- (6) The Cuntz semigroup of the Jiang-Su algebra is the Fraïssé limit of the Fraïssé category \mathcal{K}_Z .

Some of the notions in this paper and in [50] seem to hint at the right direction to develop a model theory of Cu-semigroups, similar to the model theory of C^* -algebras from [27,28]. We do not pursue this here, since this is an elaborate task that will be done elsewhere.

Organization of the paper. Section 2 recalls the categorical Fraïssé theory developed in [40], where the reader can find the definition of Fraïssé categories, Fraïssé sequences, and the fact that any Fraïssé category admits a unique Fraïssé sequence, up to two-sided approximate intertwining.

We introduce (*Cuntz*) *Fraïssé categories* and their *limit* in Section 3. We start by recalling some preliminaries about the category Cu and the notion of *finite-set comparison* for Cu-morphisms (Definition 3.2). We show that Cauchy sequences with respect to finite-set comparison have a unique *limit* (Theorem 3.8), and we study approximate intertwinings in the category Cu (Theorem 3.17). Then, we define the ‘Cuntz analogue’ of a Fraïssé category (Definition 3.19), a Fraïssé sequence (Definition 3.20), and the fact that any Fraïssé category admits a unique Fraïssé sequence, up to two-sided approximate intertwining (Theorem 3.21). Finally, a characterization of Fraïssé limits is given (Theorem 3.27). We finish the section by exploring the relations between Fraïssé categories of C^* -algebras and Fraïssé categories of Cuntz semigroups. (Part C.)

Section 4 is divided in Parts A–E, which deal with the aforementioned examples. In Section 5 we define the *Thomsen semigroup of a Cu-semigroup* (Paragraph 5.2) by using the generator \mathbb{G} of the category Cu. This allows us to define metrics on any Hom_{Cu} -set (Definition 5.8). We then explore the properties and several examples of such metrics (Examples 5.10–5.14 and Proposition 5.17).

Acknowledgments. This research started when the second author visited the first author at the Czech Academy of Sciences. They are both grateful to the IMCAS for its hospitality and for providing a great working environment. The first author would also like to thank W. Kubiś for introducing him to Fraïssé Theory.

Both authors thank the anonymous referee for their helpful comments.

2. Preliminaries on Fraïssé categories

The aim of this section is to gather the main definitions and results about *Fraïssé categories*. These differ slightly, but include, the model-theoretical tools originally developed in [29]. The categorical approach described below has been developed by Kubiś, and we follow his notation and definitions from [40]. (See also [9] and [32].)

As mentioned in the introduction, the core idea of Fraïssé theory is to produce ‘large’ objects that are universal and homogeneous in a generic sense —these objects will be called *Fraïssé limits*— for a given category of ‘small’ objects —these categories will be called *Fraïssé categories*—.

Definition 2.1. Let \mathfrak{c} be a small category. We say that \mathfrak{c} is *metric-enriched* if

- (i) For any $A, B \in \mathfrak{c}$, the set $\text{Hom}_{\mathfrak{c}}(A, B)$ is equipped with a metric $d_{(A,B)}$.
- (ii) For any $\alpha_1, \alpha_2 \in \text{Hom}_{\mathfrak{c}}(A, B)$ and $\beta \in \text{Hom}_{\mathfrak{c}}(B, C)$, we have

$$d_{(A,C)}(\beta \circ \alpha_1, \beta \circ \alpha_2) \leq d_{(A,B)}(\alpha_1, \alpha_2).$$

- (iii) For any $\alpha \in \text{Hom}_{\mathfrak{c}}(A, B)$ and $\beta_1, \beta_2 \in \text{Hom}_{\mathfrak{c}}(B, C)$, we have

$$d_{(A,C)}(\beta_1 \circ \alpha, \beta_2 \circ \alpha) \leq d_{(B,C)}(\beta_1, \beta_2).$$

Whenever the context is clear, we write $d_{\mathfrak{c}}$ instead of $d_{(A,B)}$.

Definition 2.2. Let \mathfrak{c} be a metric-enriched category. We say that \mathfrak{c}

- (JEP) satisfies the *joint embedding property* if, for any $A_1, A_2 \in \mathfrak{c}$, there exists $B \in \mathfrak{c}$ such that both $\text{Hom}_{\mathfrak{c}}(A_1, B)$ and $\text{Hom}_{\mathfrak{c}}(A_2, B)$ are nonempty.
- (NAP) satisfies the *near amalgamation property* if, for any $\varepsilon > 0$, and any \mathfrak{c} -morphisms $\alpha_1 \in \text{Hom}_{\mathfrak{c}}(A, B_1)$ and $\alpha_2 \in \text{Hom}_{\mathfrak{c}}(A, B_2)$, there exist $C \in \mathfrak{c}$ and \mathfrak{c} -morphisms $\beta_1 \in \text{Hom}_{\mathfrak{c}}(B_1, C)$ and $\beta_2 \in \text{Hom}_{\mathfrak{c}}(B_2, C)$ such that $d_{\mathfrak{c}}(\beta_1 \circ \alpha_1, \beta_2 \circ \alpha_2) < \varepsilon$.
- (SEP) is *separable* if there exists a countable dominating subcategory $\mathfrak{s} \subseteq \mathfrak{c}$, that is,
 - the set of \mathfrak{s} -morphisms is countable. (A fortiori, so is the set of \mathfrak{s} -objects.)
 - for any $A \in \mathfrak{c}$ there exists $S \in \mathfrak{s}$ such that $\text{Hom}_{\mathfrak{c}}(A, S)$ is nonempty.
 - for any $\varepsilon > 0$ and any \mathfrak{c} -morphism $\sigma: S \rightarrow A$ with $S \in \mathfrak{s}$, there exist $T \in \mathfrak{s}$ and $\alpha \in \text{Hom}_{\mathfrak{c}}(A, T)$ and $\nu \in \text{Hom}_{\mathfrak{s}}(S, T)$ such that $d_{\mathfrak{c}}(\alpha \circ \sigma, \nu) < \varepsilon$.

We say that \mathfrak{c} is a *Fraïssé category* if \mathfrak{c} satisfies (JEP), (NAP) and (SEP).

Remark 2.3. Note that, despite the term ‘Joint Embedding Property’, (JEP) does not ask the homomorphisms involved to be embeddings.

Definition 2.4. Let \mathfrak{c} be a metric-enriched category. An inductive sequence $(F_i, \sigma_{i,j})_{i \in \mathbb{N}}$ is called a *Fraïssé sequence* if, for any $\varepsilon > 0$ and any \mathfrak{c} -morphism $\gamma_i: F_i \rightarrow C$, there exists a \mathfrak{c} -morphism $\gamma_j: C \rightarrow F_j$ for some $j \geq i$ such that $d_{\mathfrak{c}}(\sigma_{i,j}, \gamma_j \circ \gamma_i) < \varepsilon$.

Theorem 2.5 (cf. [40]). *Let \mathfrak{c} be a Fraïssé category. Then \mathfrak{c} admits a Fraïssé sequence which is unique up to two-sided approximate intertwining.*

Let us now recall the notion of *Fraïssé limits*. As stated before, these objects are often ‘large’, in the sense that they do not belong to the Fraïssé category \mathfrak{c} at hand. Instead, they are built as inductive limits of \mathfrak{c} -objects. (Particularly, as inductive limits of Fraïssé sequences.)

More concretely, this amounts to the fact that a Fraïssé category \mathfrak{c} need not have inductive limits. Because of this, one often considers an inclusion of categories of the form $\mathfrak{c} \subseteq \mathfrak{d}$, where \mathfrak{d} does have inductive limits. However, the category \mathfrak{d} cannot be ‘too’ large, since this inclusion is asked to satisfy the following *almost factorization property*.

Definition 2.6. Let $\mathfrak{c}, \mathfrak{d}$ be metric-enriched categories such that \mathfrak{d} has inductive limits and $\mathfrak{c} \subseteq \mathfrak{d}$. We say that the inclusion $\mathfrak{c} \subseteq \mathfrak{d}$ has the *almost factorization property* if, for any inductive system $(A_i, \sigma_{i,j})_{i \in \mathbb{N}}$ in \mathfrak{c} with \mathfrak{d} -limit $(A, \sigma_{i,\infty})_i$, any $B \in \mathfrak{c}$, any \mathfrak{d} -morphism $\beta: B \rightarrow A$, and any $\varepsilon > 0$, there exist $i \in \mathbb{N}$ and a \mathfrak{c} -morphism $\beta_\varepsilon: B \rightarrow A_i$ such that $d_{\mathfrak{d}}(\sigma_{i,\infty} \circ \beta_\varepsilon, \beta) < \varepsilon$.

Theorem 2.7 (cf. [40]). *Let \mathfrak{c} be a Fraïssé category included in a category \mathfrak{d} which admits inductive limits, and such that any object in \mathfrak{d} is a limit of a \mathfrak{c} -sequence. Then, any Fraïssé sequence in \mathfrak{c} has a \mathfrak{d} -limit F satisfying the following properties:*

- (i) F is unique up to isomorphism.
- (ii) For any $D \in \mathfrak{d}$, the set $\text{Hom}_{\mathfrak{d}}(D, F)$ is nonempty.

If, additionally, $\mathfrak{c} \subseteq \mathfrak{d}$ satisfies the almost factorization property, then

- (iii) For any $\varepsilon > 0$, any $C \in \mathfrak{c}$, and any $\alpha_1, \alpha_2 \in \text{Hom}_{\mathfrak{d}}(C, F)$, there exists a \mathfrak{d} -isomorphism $\eta: F \xrightarrow{\cong} F$ such that $d_{\mathfrak{d}}(\eta \circ \alpha_1, \alpha_2) < \varepsilon$.

Remark 2.8. The category of C^* -algebras C^* is metric-enriched by the usual norm-distance between $*$ -homomorphisms. Therefore, the definitions given here can be applied to C^* directly (as done in [32]). However, it is more common to compare $*$ -homomorphisms metrically on finite sets. (See e.g. [56].) For example, (NAP) in Definition 2.2 gets changed to: for any $\varepsilon, \alpha_1, \alpha_2$, and finite set $F \subseteq A$, there exist $*$ -homomorphism β_1, β_2 such that $\|\beta_1 \circ \alpha_1(x) - \beta_2 \circ \alpha_2(x)\| < \varepsilon$ for every $x \in F$. A similar change is done in the third condition of (SEP) and the definition of Fraïssé sequence.

As we will discuss in Remark 3.32, for our purposes it would also be interesting to study Fraïssé categories of C^* -algebras with respect to approximate unitary equivalence. In that version, (NAP) would be changed to: for any ε , α_1, α_2 , and finite set $F \subseteq A$, there exist $*$ -homomorphism β_1, β_2 and a unitary $u \in \tilde{C}$ such that $\|u^* \beta_1 \circ \alpha_1(x) u - \beta_2 \circ \alpha_2(x)\| < \varepsilon$ for every $x \in F$. Analogous changes would be made to the other definitions.

3. Fraïssé categories of Cuntz semigroups

As explained in the previous section, the approach to Fraïssé categories from [40] requires each Hom-set to be equipped with a right- and left-enriched metric. This rarely happens in the category Cu : Every Hom_{Cu} -set admits a natural metric (defined and studied in the last section of this paper), but such a metric is seldom left-enriched. This was already the case for the specific instances of this metric considered in the past. (See e.g. [15] and [19].)

To overcome these constraints, we introduce a theory of Fraïssé categories for Cu -semigroups where, instead of using a metric on the Hom-sets, we compare morphisms on finite sets. This allows us to bypass any sort of enrichment property. In the course of our investigations, we also define an analog of Cauchy sequences for Cu -morphisms, which are shown to have a unique limit. Further, building on the results from [15], we develop a general theory of one- and two-sided approximate intertwinings in Cu . These tools allow us to define and obtain analogous notions and results to those of Kubiś. We finish the section by describing the relation between these theories in the context of C^* -algebras and concrete Cuntz semigroups.

First, let us recall some preliminaries about the category Cu .

3.1 (Cu-semigroups). Let x, y be elements in a partially ordered set P . We write $x \ll y$ if, for every increasing sequence $(z_n)_n$ which has a supremum such that $y \leq \sup_n z_n$, there exists $n \in \mathbb{N}$ such that $x \leq z_n$.

As defined in [22], a positively ordered monoid S is said to be a *Cu-semigroup* if S satisfies the following properties:

- (O1) Every increasing sequence in S has a supremum.
- (O2) Every element in S can be written as the supremum of a \ll -increasing sequence.
- (O3) The addition and the \ll -relation are compatible.
- (O4) Suprema of increasing sequences and the addition are compatible.

A monoid morphism between Cu -semigroups is a *Cu-morphism* if it preserves the order, the \ll -relation, and suprema of increasing sequences. We denote the category of Cu -semigroups and Cu -morphisms by Cu . (See e.g. [5] or [30] for a more detailed exposition.)

The *Cuntz semigroup* of a C^* -algebra A , denoted by $\text{Cu}(A)$, is the quotient $(A \otimes \mathcal{K})_+ / \sim$ equipped with the addition induced by diagonal addition and the order induced by \preceq , where the relations \preceq and \sim are defined as follows:

$$a \preceq b : \iff a = \lim_n r_n b r_n^* \text{ for some sequence } (r_n)_n \subseteq A \otimes \mathcal{K}.$$

$$a \sim b : \iff a \preceq b \text{ and } b \preceq a.$$

The Cuntz semigroup of A , first considered in [24], is always a Cu-semigroup. (See [22].) Further, every $*$ -homomorphism from A to B induces a Cu-morphism from $\text{Cu}(A)$ to $\text{Cu}(B)$.

The relation between concrete Cuntz semigroups and the abstract category Cu has been studied extensively. (See [2,3,6,13,14,21,51,53] among many others.) For instance, it is known that the category Cu has direct limits and that the functor Cu is continuous. (See [5], [22].)

A Cu-semigroup S is *countably-based* if S contains a countable, sup-dense subset. Examples include the Cuntz semigroup of any separable C^* -algebra.

A. Comparison of Cu-morphisms

As mentioned at the beginning of this section, it is not clear when a set of Cu-morphisms can be equipped with a (meaningful) enriching metric. In order to overcome this issue and work in the general setting, we will compare Cu-morphisms on finite sets. This notion was introduced explicitly in [15], although the idea had also appeared implicitly in the past when working with specific families of Cu-morphisms. (See e.g. [17–19,41,43].)

Definition 3.2 ([15, Definition 3.9]). Given a pair of Cu-morphisms $\alpha, \beta : S \rightarrow T$ and a finite subset $F \subseteq S$, we say that α and β *compare on F* , and we write $\alpha \simeq_F \beta$, if for any pair $x', x \in F$ with $x' \ll x$, we have

$$\alpha(x') \leq \beta(x) \quad \text{and} \quad \beta(x') \leq \alpha(x).$$

Remark 3.3. As observed in [15], the following are equivalent:

- (i) $\alpha = \beta$.
- (ii) $\alpha \simeq_F \beta$ for any finite subset $F \subseteq S$.
- (iii) $\alpha \simeq_{\{s',s\}} \beta$ for any $s', s \in S$ with $s' \ll s$.

Note that finite-set comparison can also be used for weaker forms of morphisms between Cu-semigroups. (See [17, Definition 2.2].)

With this notion of comparison at hand, we can define Cauchy sequences, limits, and approximate intertwinings in the category Cu.

Definition 3.4. Let $(\alpha_i)_i$ be a sequence of Cu-morphisms in $\text{Hom}_{\text{Cu}}(S, T)$. We say that $(\alpha_i)_i$ is a *Cauchy sequence* if, for any finite subset $F \subseteq S$, there exists an index i_F such that $\alpha_j \simeq_F \alpha_k$ whenever $j, k \geq i_F$.

The following definition of convergence was introduced in [41, Definition 5.1] for concrete Cuntz morphisms. We give here the definition for any sequence of Cu-morphisms.

Definition 3.5. Let $(\alpha_i)_i$ be a sequence of Cu-morphisms in $\text{Hom}_{\text{Cu}}(S, T)$. We say that $(\alpha_i)_i$ *converges towards a Cu-morphism* $\alpha : S \rightarrow T$ if, for any finite subset $F \subseteq S$, there exists an index i_F such that $\alpha_j \simeq_F \alpha$ whenever $j \geq i_F$.

Remark 3.6. In view of our previous remark, note that $(\alpha_i)_i$ converges to α if, for any pair of elements $x', x \in S$ with $x' \ll x$, there exists an index i_0 such that $\alpha_j(x') \leq \alpha(x)$ and $\alpha(x') \leq \alpha_j(x)$ whenever $j \geq i_0$.

Example 3.7. If a sequence of $*$ -homomorphisms $(\varphi_i : A \rightarrow B)_i$ converges in point-norm topology to a morphism $\varphi : A \rightarrow B$, then the sequence $(\text{Cu}(\varphi_i))_i$ converges to $\text{Cu}(\varphi)$.

Indeed, given $[a], [b] \in \text{Cu}(A)$ with $[a] \ll [b]$, we can find $\varepsilon > 0$ small enough such that $[a] \leq [(b - \varepsilon)_+]$. Moreover, we can find $i \in \mathbb{N}$ big enough such that $\|\varphi_j(a) - \varphi(a)\| < \varepsilon$ for every $j \geq i$. Therefore, we have $\varphi_j(a) \lesssim \varphi_j((b - \varepsilon)_+) = (\varphi_j(b) - \varepsilon)_+ \lesssim \varphi(b)$ and $\varphi(a) \lesssim \varphi((b - \varepsilon)_+) = (\varphi(b) - \varepsilon)_+ \lesssim \varphi_j(b)$.

Theorem 3.8. *Let S, T be Cu-semigroups. Then any Cauchy sequence $(\alpha_i)_i$ in $\text{Hom}_{\text{Cu}}(S, T)$ converges towards a unique Cu-morphism.*

Proof. Let us first assume that S is countably-based, so that there exists a \subseteq -increasing sequence $(B_n)_n$ of finite sets of S such that $B := \bigcup_{n \in \mathbb{N}} B_n$ is sup-dense in S . Let $(\alpha_i)_i$ be a Cauchy sequence in $\text{Hom}_{\text{Cu}}(S, T)$. We can find a strictly increasing map $\varphi : \mathbb{N} \rightarrow \mathbb{N}$ such that $\alpha_j \simeq_{B_n} \alpha_k$ for any $j, k \geq \varphi(n)$.

Let $\text{Seq}_{\ll}(B)$ denote the set of \ll -increasing sequence in B , which we may think of as maps $f : \mathbb{N} \rightarrow B, i \mapsto f_i$, such that $f_i \ll f_{i+1}$ in S for each i . With this notation, there exists a map $\psi : \mathbb{N} \times \text{Seq}_{\ll}(B) \rightarrow \mathbb{N}$ such that

- (i) $\psi(\cdot, f) : \mathbb{N} \rightarrow \mathbb{N}$ is strictly increasing for every fixed f .
- (ii) $\{f_0, \dots, f_{l+1}\} \subseteq B_{\psi(l, f)}$ for each $l \in \mathbb{N}$.

Note that the map $\varphi \circ \psi(\cdot, f)$ is strictly increasing for any $f \in \text{Seq}_{\ll}(B)$.

Now fix $l \in \mathbb{N}$. By the definition of φ we have that $\alpha_j \simeq_{B_{\psi(l, f)}} \alpha_k$ for any $j, k \geq \varphi(\psi(l, f))$. Further, it follows from the construction of ϕ that

$$\alpha_j(f_i) \leq \alpha_k(f_{i+1}) \quad \text{and} \quad \alpha_k(f_i) \leq \alpha_j(f_{i+1})$$

for any $0 \leq i \leq l$ and any $j, k \geq \varphi(\psi(l, f))$.

In particular, for $j = \varphi(\psi(l, f))$, $k = \varphi(\psi(l + 1, f))$ and $i = l$, we obtain

$$\alpha_{\varphi(\psi(l, f))}(f_l) \leq \alpha_{\varphi(\psi(l+1, f))}(f_{l+1}).$$

In other words, the sequence $(\alpha_{\varphi(\psi(l, f))}(f_l))_l$ is increasing. Consequently, we can define the following map

$$\begin{aligned} \alpha_{\text{Seq}} : \text{Seq}_{\ll}(B) &\longrightarrow T \\ f &\longmapsto \sup_l \alpha_{\varphi(\psi(l, f))}(f_l) \end{aligned}$$

We aim to construct a Cu-morphism $\alpha : S \longrightarrow T$ induced by α_{Seq} . For this, we will need the following claim.

Claim. Let $f, g \in \text{Seq}_{\ll}(B)$ be such that $\sup f \leq \sup g$. Then $\alpha_{\text{Seq}}(f) \leq \alpha_{\text{Seq}}(g)$.

Proof of the Claim. Let f, g be as in the statement. For any $l \in \mathbb{N}$, there exists $m \in \mathbb{N}$ such that $f_l \ll f_{l+1} \ll g_m$. Since $\varphi \circ \psi(\cdot, g)$ is strictly increasing, we can assume that m is large enough so that $\varphi(\psi(l, f)) \leq \varphi(\psi(m, g))$. By construction, we have $\alpha_{\varphi(\psi(l, f))}(f_l) \leq \alpha_k(f_{l+1})$ for any $k \geq \varphi(\psi(l, f))$. Thus, we compute

$$\alpha_{\varphi(\psi(l, f))}(f_l) \leq \alpha_{\varphi(\psi(m, g))}(f_{l+1}) \leq \alpha_{\varphi(\psi(m, g))}(g_m) \leq \alpha_{\text{Seq}}(g)$$

which implies that $\alpha_{\text{Seq}}(f) \leq \alpha_{\text{Seq}}(g)$ and proves the claim.

Since B is dense in S , we are now able to construct the following order-preserving map

$$\begin{aligned} \alpha : S &\longrightarrow T \\ s &\longmapsto \alpha_{\text{Seq}}((s_i)_i) \end{aligned}$$

where $(s_i)_i$ is any \ll -increasing sequence in the basis B obtained from (O2), whose supremum is s . (The claim shows that α is well-defined, i.e. α does not depend on the sequence $(s_i)_i$, and also that α preserves the order.)

Further, using (O4), it is readily checked that α preserves the addition. Using a diagonal-type argument (see e.g. the proof of [15, Lemma 3.12]), it can also be shown that α preserves suprema of increasing sequences.

We are left to show that α preserves the compact-containment relation. Let $f, g \in \text{Seq}_{\ll}(B)$ be such that $\sup f \ll \sup g$. Then, there exists $m \in \mathbb{N}$ such that $f_l \ll g_{m-2} \ll g_{m-1} \ll g_m$ for any $l \in \mathbb{N}$. Find l_0 big enough such that $\varphi(\psi(m, g)) \leq \varphi(\psi(l, f))$ for any $l \geq l_0$. By construction, we have

$$\alpha_{\varphi(\psi(l, f))}(f_l) \ll \alpha_{\varphi(\psi(l, f))}(g_{m-2}) \leq \alpha_{\varphi(\psi(m, g))}(g_{m-1}) \ll \alpha_{\varphi(\psi(m, g))}(g_m) \leq \alpha_{\text{Seq}}(g)$$

whenever $l \geq l_0$. In particular, we get $\alpha_{\text{Seq}}(f) \ll \alpha_{\text{Seq}}(g)$ and, hence, α preserves the \ll -relation. This shows that α is a well-defined Cu-morphism.

Finally, let us prove that $(\alpha_i)_i$ converges to α . Let $x', x \in S$ be such that $x' \ll x$. By density of B in S , there exists $f \in \text{Seq}_{\ll}(B)$ such that $x' \ll \sup f \ll x$. On the one

hand, note that there exists $l \in \mathbb{N}$ big enough such that $x' \ll f_l \ll f_{l+1} \ll x$. We deduce that, for any $i \geq \varphi(\psi(l + 1, f))$, we have

$$\alpha_i(x') \leq \alpha_i(f_l) \leq \alpha_{\varphi(\psi(l+1,f))}(f_{l+1}) \leq \alpha_{\text{Seq}}(f) = \alpha(\sup f) \leq \alpha(x).$$

On the other hand, note that $\alpha(x') \ll \alpha_{\text{Seq}}(f)$ and, hence, there exists $m \in \mathbb{N}$ big enough such that $\alpha(x') \ll \alpha_{\varphi(\psi(m,f))}(f_m)$. We deduce that, for any $i \geq \varphi(\psi(m, f))$, we have

$$\alpha(x') \leq \alpha_{\varphi(\psi(m,f))}(f_m) \leq \alpha_i(f_{m+1}) \leq \alpha_i(x).$$

This shows that $(\alpha_i)_i$ converges towards α . Since such an α is unique, this also proves that α does not depend on the basis B chosen.

Now let us show that the result holds for any (possibly not countably-based) Cu-semigroup S . Let $(\alpha_i)_i$ be a Cauchy sequence in $\text{Hom}_{\text{Cu}}(S, T)$. Any countably-based sub-Cu-semigroup H of S (i.e. H is Cu-semigroup that order-embeds into S) naturally induces a Cauchy sequence $(\alpha_{i|_H})_i$ by restriction. We can thus construct its limit as above, which we denote by α_H .

Let $x \in S$. It follows from [50, Lemma 5.1] that there exists a (possibly not unique) countably-based sub-Cu-semigroup H_x such that $x \in H_x$. Let H_1, H_2 be countably-based sub-Cu-semigroups that contain x . By construction, there exist \ll -increasing sequences $(x_n)_n$ and $(x'_n)_n$ in H_1 and H_2 respectively with supremum x such that $\alpha_{H_1}(x) = \sup_n \alpha_n(x_n)$ and $\alpha_{H_2}(x) = \sup_n \alpha_n(x'_n)$. Further, we can choose these sequences so that, for any $m \geq n$, we have

$$\alpha_n(x_n) \leq \alpha_m(x_{n+1}) \quad \text{and} \quad \alpha_n(x'_n) \leq \alpha_m(x'_{n+1}).$$

Let $n \in \mathbb{N}$, and find $m \geq n$ such that $x_{n+1} \ll x'_m$. Then, we get

$$\alpha_n(x_n) \leq \alpha_m(x_{n+1}) \leq \alpha_m(x'_m) \leq \alpha_{H_2}(x).$$

Taking supremum over n we obtain $\alpha_{H_1}(x) \leq \alpha_{H_2}(x)$ and, by a symmetric argument, we also get $\alpha_{H_2}(x) \leq \alpha_{H_1}(x)$. We conclude that $\alpha_{H_1}(x) = \alpha_{H_2}(x)$ for any $x \in S$ and, consequently, that the following map is well-defined

$$\begin{aligned} \alpha: S &\longrightarrow T \\ x &\longmapsto \alpha_{H_x}(x) \end{aligned}$$

Using the techniques from [50] one can check that α is a Cu-morphism and that the sequence $(\alpha_i)_i$ converges to α (by construction). This ends the proof. \square

Notation 3.9. We have just shown that any Cauchy sequence $(\alpha_i)_i$ in $\text{Hom}_{\text{Cu}}(S, T)$ converges towards a unique Cu-morphism $\alpha: S \longrightarrow T$. We will say that α is the *limit* of the sequence, and write $\lim_i \alpha_i = \alpha$.

We are now able to define and study approximate intertwinings in the category Cu . This generalizes the concepts introduced in [15] for the specific case of uniformly-based Cu -semigroups.

Definition 3.10. Let $(S_i, \sigma_{i,j})_{i \in \mathbb{N}}$ and $(T_i, \tau_{i,j})_{i \in \mathbb{N}}$ be two inductive sequences in Cu . Assume that there exists a strictly increasing map $\varphi: \mathbb{N} \rightarrow \mathbb{N}$ together with a sequence of Cu -morphisms $(\alpha_i: S_i \rightarrow T_{\varphi(i)})_i$ satisfying the following property:

For any finite subset $F \subseteq S_i$, there exists an index $i_F \geq i$ such that, for any $j \geq i_F$ and any $k \geq j$, the diagram

$$\begin{CD} S_i @>\sigma_{i,j}>> S_j @>>\sigma_{j,k}>> S_k \\ @. @V\alpha_jVV @VV\alpha_kV \\ T_{\varphi(j)} @>>\tau_{\varphi(j),\varphi(k)}>> T_{\varphi(k)} \end{CD}$$

approximately commutes within F , that is, $\alpha_k \circ \sigma_{i,k} \simeq_F \tau_{\varphi(j),\varphi(k)} \circ \alpha_j \circ \sigma_{i,j}$.

We say that $(\alpha_i)_i$ is a *one-sided approximate intertwining*.

Proposition 3.11. Let $(S_i, \sigma_{i,j})_{i \in \mathbb{N}}$ and $(T_i, \tau_{i,j})_{i \in \mathbb{N}}$ be two inductive sequences in Cu with respective limits $(S, \sigma_{i,\infty})$ and $(T, \tau_{i,\infty})$. Let $(\alpha_i: S_i \rightarrow T_{\varphi(i)})_i$ be a one-sided approximate intertwining. Then there exists a Cu -morphism $\alpha: S \rightarrow T$ such that, for any finite subset $F \subseteq S_i$, there exists $i_F \geq i$ such that, for any $j \geq i_F$, the diagram

$$\begin{CD} S_i @>\sigma_{i,j}>> S_j @>>\sigma_{j,\infty}>> S \\ @. @V\alpha_jVV @VV\exists\alpha V \\ T_{\varphi(j)} @>>\tau_{\varphi(j),\infty}>> T \end{CD}$$

approximately commutes within F , that is, $\alpha \circ \sigma_{i,\infty} \simeq_F \tau_{\varphi(j),\infty} \circ \alpha_j \circ \sigma_{i,j}$.

Proof. For any $i \in \mathbb{N}$ the sequence $(\tau_{\varphi(j),\infty} \circ \alpha_j \circ \sigma_{i,j})_j$ is Cauchy. It follows from Theorem 3.8 that the sequence has a limit, which we denote by $\eta_i: S_i \rightarrow T$. Using that this limit is unique, it is readily checked that $\eta_i = \eta_j \circ \sigma_{i,j}$ for any $i \leq j$. By the universal property of direct limits, this induces a Cu -morphism $\alpha: S \rightarrow T$ such that $\eta_i = \alpha \circ \sigma_{i,\infty}$ for any $i \in \mathbb{N}$. Finally, note that for any finite subset $F \subseteq S_i$, we can find an index i_F such that $\eta_i \simeq_F \tau_{\varphi(j),\infty} \circ \alpha_j \circ \sigma_{i,j}$ for any $j \geq i_F$. Thus, we have

$$\alpha \circ \sigma_{i,\infty} = \eta_i \simeq_F \tau_{\varphi(j),\infty} \circ \alpha_j \circ \sigma_{i,j}$$

which ends the proof. \square

Example 3.12. Let $(\psi_i: A_i \rightarrow B_{\varphi(i)})_i$ be a one-sided approximate intertwining of C^* -algebras from $(A_i, \varphi_{i,j})_{i \in \mathbb{N}}$ to $(B_i, \phi_{i,j})_{i \in \mathbb{N}}$. It follows from Example 3.7 that the induced

maps $(\text{Cu}(\psi_i))_i$ define a one-sided approximate intertwining from $(\text{Cu}(A_i), \text{Cu}(\varphi_{i,j}))_i$ to $(\text{Cu}(B_i), \text{Cu}(\phi_{i,j}))_i$.

Definition 3.13. Let $(S_i, \sigma_{i,j})_{i \in \mathbb{N}}$ and $(T_i, \tau_{i,j})_{i \in \mathbb{N}}$ be two inductive sequences in Cu . Assume that there exist two strictly increasing maps $\varphi, \psi: \mathbb{N} \rightarrow \mathbb{N}$ together with two sequences of Cu -morphisms $(\alpha_i: S_i \rightarrow T_{\varphi(i)})_i$ and $(\beta_i: T_i \rightarrow S_{\psi(i)})_i$ satisfying the following property:

For any finite sets $F \subseteq S_i$ and $G \subseteq T_i$, there exist indices $i_F, i_G \geq i$ such that, for any $j \geq i_F, j' \geq i_G$ and any $k \geq \varphi(j), k' \geq \psi(j')$, the diagrams

$$\begin{array}{ccc}
 S_i & \xrightarrow{\sigma_{i,j}} & S_j & \xrightarrow{\sigma_{j,\psi(k)}} & S_{\psi(k)} & & S_{\psi(j')} & \xrightarrow{\sigma_{\psi(j'),k'}} & S_{k'} \\
 & & \alpha_j \downarrow & & \uparrow \beta_k & & \beta_{j'} \uparrow & & \downarrow \alpha_{k'} \\
 & & T_{\varphi(j)} & \xrightarrow{\tau_{\varphi(j),k}} & T_k & & T_i & \xrightarrow{\tau_{i,j'}} & T_{j'} & \xrightarrow{\tau_{j',\varphi(k')}} & T_{\varphi(k')}
 \end{array}$$

approximately commute within F and G respectively, that is,

$$\sigma_{i,\psi(k)} \simeq_F \beta_k \circ \tau_{\varphi(j),k} \circ \alpha_j \circ \sigma_{i,j} \quad \text{and} \quad \tau_{i,\varphi(k')} \simeq_G \alpha_{k'} \circ \sigma_{\psi(j'),k'} \circ \beta_{j'} \circ \tau_{i,j'}.$$

We say that $(\alpha_i, \beta_i)_i$ is a *two-sided approximate intertwining*.

Remark 3.14. Each of the sequences $(\alpha_i)_i$ and $(\beta_i)_i$ that define a two-sided approximate intertwining induce a one-sided approximate intertwining.

Throughout the paper, when considering a finite set F for comparison of Cu -morphisms, we will often need to construct a larger finite set \tilde{F} which is *finer* than F in the following sense.

Definition 3.15. Let F, \tilde{F} be (finite) subsets of a Cu -semigroup and let $n \in \mathbb{N}$. We will say that \tilde{F} is an *n-refinement* of F , or that \tilde{F} *refines* F *n-times*, if

- (i) $F \subseteq \tilde{F}$.
- (ii) For any $f', f \in F$ such that $f' \ll f$, there exist n elements $g_1, \dots, g_n \in \tilde{F}$ such that $f' \ll g_1 \ll \dots \ll g_n \ll f$.

Note that, for any $n \geq 1$ and any finite set F of a Cu -semigroup, we can always find a *finite n-refinement* of F .

Remark 3.16. One of the reasons why the previous notion is needed is that \simeq_F is not a transitive relation, that is, $\alpha \simeq_F \beta \simeq_F \gamma$ does not imply $\alpha \simeq_F \gamma$. Instead, what we do have is that, if \tilde{F} is an *n-refinement* of F , then $\alpha \simeq_{\tilde{F}} \alpha_1 \simeq_{\tilde{F}} \dots \simeq_{\tilde{F}} \alpha_n \simeq_{\tilde{F}} \beta$ implies $\alpha \simeq_F \beta$.

Theorem 3.17. *Let $(S_i, \sigma_{i,j})_{i \in \mathbb{N}}$ and $(T_i, \tau_{i,j})_{i \in \mathbb{N}}$ be two inductive sequences in Cu with respective limits $(S, \sigma_{i,\infty})$ and $(T, \tau_{i,\infty})$. Assume that there exists a two-sided approximate intertwining $(\alpha_i: S_i \rightarrow T_{\varphi(i)}, \beta_i: T_i \rightarrow S_{\psi(i)})_i$.*

Then there exists a Cu -isomorphism $\alpha: S \cong T$ induced by $(\alpha_i)_i$ whose inverse is induced by $(\beta_i)_i$.

Proof. Our approach is similar to that of [15, Theorem 3.16], but we proceed with additional care since our setting is more general.

Arguing as in the proof of Proposition 3.11, we know that for any $i \in \mathbb{N}$ the sequences $(\tau_{\varphi(j),\infty} \circ \alpha_j \circ \sigma_{i,j})_j$ and $(\sigma_{\psi(j'),\infty} \circ \beta_{j'} \circ \tau_{i,j'})_{j'}$ are Cauchy, and we denote their respective limits by $\eta_i: S_i \rightarrow T$ and $\nu_i: T_i \rightarrow S$. Furthermore, these limits induce Cu -morphisms $\alpha: S \rightarrow T$ and $\beta: T \rightarrow S$. In order to show that α and β are inverses of one another, it suffices to show that $\beta \circ \eta_i = \sigma_{i,\infty}$ and $\alpha \circ \nu_i = \tau_{i,\infty}$ for any $i \in \mathbb{N}$.

Let F be a finite subset of S_i and let \tilde{F} be a 2-refinement of F . Since η_i is the limit of $(\tau_{\varphi(j),\infty} \circ \alpha_j \circ \sigma_{i,j})_j$, we know that there exists $j \geq i$ big enough such that $\eta_i \simeq_{\tilde{F}} \tau_{\varphi(j),\infty} \circ \alpha_j \circ \sigma_{i,j}$. Post-composing with β , we obtain

$$\beta \circ \eta_i \simeq_{\tilde{F}} \nu_{\varphi(j)} \circ \alpha_j \circ \sigma_{i,j}. \tag{1}$$

Consider $\tilde{G} := \alpha_j \circ \sigma_{i,j}(\tilde{F}) \subseteq T_{\varphi(j)}$. Since $\nu_{\varphi(j)}$ is the limit of $(\sigma_{\psi(k),\infty} \circ \beta_k \circ \tau_{\varphi(j),k})_k$, we know that there exists $k \geq \varphi(j)$ big enough such that $\nu_{\varphi(j)} \simeq_{\tilde{G}} \sigma_{\psi(k),\infty} \circ \beta_k \circ \tau_{\varphi(j),k}$. Precomposing by $\alpha_j \circ \sigma_{i,j}$, this implies

$$\nu_{\varphi(j)} \circ \alpha_j \circ \sigma_{i,j} \simeq_{\tilde{F}} \sigma_{\psi(k),\infty} \circ \beta_k \circ \tau_{\varphi(j),k} \circ \alpha_j \circ \sigma_{i,j}. \tag{2}$$

Finally, since $(\alpha_i, \beta_i)_i$ is a two-sided approximate intertwining, we also have

$$\sigma_{\psi(k),\infty} \circ \beta_k \circ \tau_{\varphi(j),k} \circ \alpha_j \circ \sigma_{i,j} \simeq_{\tilde{F}} \sigma_{i,\infty} \tag{3}$$

whenever j and k are big enough.

It follows from the construction of \tilde{F} and a combination of (1)-(3) that $\beta \circ \eta_i \simeq_F \sigma_{i,\infty}$. Since this holds for any finite subset, we must have $\beta \circ \eta_i = \sigma_{i,\infty}$. Therefore, we get $\beta \circ \alpha = \text{id}_S$. The fact that $\alpha \circ \beta = \text{id}_T$ follows from a symmetric argument. \square

3.18 (Comparison and approximate intertwining in Cu^*). Note that none of the proofs above uses the fact that the ordered monoids under consideration are positively ordered. Thus, all the results in this section are still valid for the larger category Cu^* introduced in [16] (loosely, this is the category of not necessarily positively ordered Cu -semigroups).

Many refinements of the Cuntz semigroup have Cu^* as their target category, and thus are amenable to the techniques developed here. We predict that this will play an important role when one such variant of the Cuntz semigroup is used to classify morphisms between certain C^* -algebras.

B. Fraïssé categories of Cuntz semigroups

We are now ready to introduce a version of Fraïssé Theory for abstract Cuntz semigroups, analogous to that of [40]. As stated earlier, we use finite-set comparison of Cu-morphisms to bypass the need of enriched metrics.

Definition 3.19. Let \mathfrak{c} be a subcategory of Cu. We say that \mathfrak{c}

- (JEP_{Cu}) satisfies the (*Cuntz*) *joint embedding property* if, for any $A_1, A_2 \in \mathfrak{c}$, there exists $B \in \mathfrak{c}$ such that both $\text{Hom}_{\mathfrak{c}}(A_1, B)$ and $\text{Hom}_{\mathfrak{c}}(A_2, B)$ are nonempty.
- (NAP_{Cu}) satisfies the (*Cuntz*) *near amalgamation property* if, for any pair of \mathfrak{c} -morphisms $\alpha_1 \in \text{Hom}_{\mathfrak{c}}(A, B_1)$ and $\alpha_2 \in \text{Hom}_{\mathfrak{c}}(A, B_2)$, and any finite subset $F \subseteq A$, there exist $C \in \mathfrak{c}$ and \mathfrak{c} -morphisms $\beta_1 \in \text{Hom}_{\mathfrak{c}}(B_1, C)$ and $\beta_2 \in \text{Hom}_{\mathfrak{c}}(B_2, C)$ such that $\beta_1 \circ \alpha_1 \simeq_F \beta_2 \circ \alpha_2$.
- (SEP_{Cu}) is (*Cuntz*) *separable* if there exists a countable dominating subcategory $\mathfrak{s} \subseteq \mathfrak{c}$, that is,
 - any object $S \in \mathfrak{s}$ is a countably-based Cu-semigroup.
 - the set of \mathfrak{s} -morphisms is countable.
 - for any $A \in \mathfrak{c}$ there exist $S \in \mathfrak{s}$ such that $\text{Hom}_{\mathfrak{c}}(A, S)$ is nonempty.
 - for any \mathfrak{c} -morphism $\sigma: S \rightarrow A$ with $S \in \mathfrak{s}$ and any finite subset $F \subseteq S$, there exists $T \in \mathfrak{s}$ and $\alpha \in \text{Hom}_{\mathfrak{c}}(A, T)$ and $\tau \in \text{Hom}_{\mathfrak{s}}(S, T)$ such that $\alpha \circ \sigma \simeq_F \tau$.

We say that \mathfrak{c} is a (*Cuntz*) *Fraïssé category* if \mathfrak{c} satisfies (JEP_{Cu}), (NAP_{Cu}) and (SEP_{Cu}).

As clarified in Remark 2.3, the Cu-morphisms involved need not be embeddings but we chose to be faithful to the historical name anyways. Next, we define a notion of (*Cuntz*) *Fraïssé sequences* and show that any (Cuntz) Fraïssé category admits such a sequence, which is unique up to two-sided approximate intertwining.

Definition 3.20. Let \mathfrak{c} be a subcategory of Cu. An inductive sequence $(S_i, \sigma_{i,j})_{i \in \mathbb{N}}$ is called a (*Cuntz*) *Fraïssé sequence* if

- (i) Every S_i is a countably-based Cu-semigroup.
- (ii) For any finite subset $F \subseteq S_i$ and any \mathfrak{c} -morphism $\alpha: S_i \rightarrow C$, there exists a \mathfrak{c} -morphism $\beta_F: C \rightarrow S_j$ for some $j \geq i$ such that $\beta_F \circ \alpha \simeq_F \sigma_{i,j}$.

Theorem 3.21 (*Existence and Uniqueness*). *Let $\mathfrak{c} \subseteq \text{Cu}$ be a Fraïssé category. Then \mathfrak{c} admits a Fraïssé sequence which is unique up to two-sided approximate intertwining in \mathfrak{c} .*

Proof. The proof that such a sequence exists is analogous to that of [40, Theorem 3.3]. We reproduce the proof here with our language of finite-set comparison for the sake of completeness. In contrast, the uniqueness part of the proof differs from [40].

[Existence] Without loss of generality, we may assume that the countable dominating subcategory $\mathfrak{s} \subseteq \mathfrak{c}$ satisfies the joint embedding property and the near amalgamation property.

Now consider the partially ordered set $\mathcal{S} := \{\text{finite inductive sequences in } \mathfrak{s}\}$ with the end-extension order, i.e. $(S_i, \sigma_{i,j})_{i,j \leq n} \leq (T_i, \tau_{i,j})_{i,j \leq m}$ in \mathcal{S} whenever $n \leq m$ and $(T_i, \tau_{i,j})_{i,j \leq n} = (S_i, \sigma_{i,j})_{i,j \leq n}$. For any $S \in \mathfrak{s}$, fix a basis $B := \bigcup_k B_k$ such that $(B_k)_k$ is a \subseteq -increasing sequence of finite sets of S . For any \mathfrak{s} -morphism $\alpha: S \rightarrow T$ and $n, k \in \mathbb{N}$, we let $D_{n,\alpha,k}$ be the subset of \mathcal{S} of all elements $(S_i, \sigma_{i,j})_{i,j \leq m}$ satisfying the following

- $m > n$.
- $\text{Hom}_{\mathfrak{s}}(S, S_i) \neq \emptyset$ for some i .
- if $S = S_n$, then there exist $j > n$ and $\beta \in \text{Hom}_{\mathfrak{s}}(T, S_j)$ such that $\beta \circ \alpha \simeq_{B_k} \sigma_{n,j}$.


Using the joint embedding property and the near amalgamation property, it is readily checked that all sets of the form $D_{n,\alpha,k}$ are cofinal in \mathcal{S} with respect to the end-extension order, that is, for any triple n, α, k and any $s \in \mathcal{S}$ there exists $d \in D_{n,\alpha,k}$ such that $s \leq d$.

Next, we argue similarly as in the proof of [40, Theorem 3.3] and make use of the following, commonly known as the Rasiowa-Sikorski lemma: Given an ordered set P and a countable family of cofinal subsets $\{D_n\}_{n \in \mathbb{N}}$, then there exists an increasing sequence $(d_n)_n$ in P with $d_n \in D_n$.


Let us fix an ordering $\varphi: \mathbb{N} \rightarrow \{(n, \alpha, k) \mid n, k \in \mathbb{N}, \alpha \in \mathfrak{s}\}$. Then, we can use the Rasiowa-Sikorski lemma to find a \leq -increasing sequence $(c_l)_l$ where $c_l \in D_{\varphi(l)}$. Note that the supremum of $(c_l)_l$ is in fact a well-defined inductive sequence in \mathfrak{s} , which we write as $(S_i, \sigma_{i,j})_i$. By construction, $(S_i, \sigma_{i,j})_i$ is Fraïssé for \mathfrak{s} . Finally, arguing again as in the proof of [40, Theorem 3.3], we deduce that $(S_i, \sigma_{i,j})_i$ is in fact a Fraïssé sequence for \mathfrak{c} .

[Uniqueness] Let $(S_i, \sigma_{i,j})_i$ and $(T_i, \tau_{i,j})_i$ be two Fraïssé sequences in \mathfrak{c} . We are going to recursively construct a two-sided approximate intertwining between them.

First, recall that all the Cu-semigroups involved are countably-based (by definition). Therefore, for each i , we can fix two \subseteq -increasing sequences $(B_n^i)_n, (C_n^i)_n$ of finite subsets of S_i, T_i respectively, such that their unions over n are sup-dense in S_i and T_i . Now, using the joint embedding property first, and then the Fraïssé sequence property twice, we construct the maps β_0 and $\alpha_{\psi(0)}$ which make the following diagram approximately commute.



Our aim is to construct strictly increasing map $\psi, \varphi: \mathbb{N} \rightarrow \mathbb{N}$ together with large enough subsets $\tilde{B}_{\psi(i)}, \tilde{C}_{\varphi(i)}$ of $S_{\psi(i)}, T_{\varphi(i)}$ and \mathfrak{c} -morphisms $\beta_{\varphi(i)}: T_{\varphi(i)} \rightarrow S_{\psi(i+1)}$ and $\alpha_{\psi(i+1)}: S_{\psi(i+1)} \rightarrow T_{\varphi(i+1)}$ producing the following approximately commutative diagram.



To do this, we proceed by induction from the initial data $B_0^0, C_0^0, \beta_0, \alpha_{\psi(0)}$. Assume that

- the numbers $\psi(0), \varphi(0), \dots, \psi(i), \varphi(i)$
- the finite sets $B_0^0, C_0^0, \dots, \tilde{B}_{\psi(i-1)}, \tilde{C}_{\varphi(i-1)}$
- the \mathfrak{c} -morphisms $\beta_0, \alpha_{\psi(0)}, \dots, \beta_{\varphi(i-1)}, \alpha_{\psi(i)}$

have been constructed for some $i \geq 0$. (By convention, we have fixed $\psi(-1) = \varphi(-1) = 0$ and $\tilde{B}_0 := B_0^0, \tilde{C}_0 := C_0^0$.)

In what follows, a *path* is any \mathfrak{Cu} -morphism in the above diagram that can be expressed as the composition of finitely many maps among $\sigma_{j,k}, \tau_{j,k}, \alpha_{\psi(j)}$ and $\beta_{\varphi(j)}$. Let us start by choosing $\tilde{B}_{\psi(i)} \subseteq S_{\psi(i)}$ and $\tilde{C}_{\varphi(i)} \subseteq T_{\varphi(i)}$ such that

- (i) $B_{\psi(i)}^{\psi(i)} \subseteq \tilde{B}_{\psi(i)}$ and $C_{\varphi(i)}^{\varphi(i)} \subseteq \tilde{C}_{\varphi(i)}$.
- (ii) $\tilde{B}_{\psi(i)}$ refines $\left\{ \pi(b) \in S_{\psi(i)} \mid b \in \bigcup_{0 < l \leq i} (\tilde{B}_{\psi(i-l)} \cup \tilde{C}_{\varphi(i-l)}) , \pi \text{ a path} \right\}$.
- (iii) $\tilde{C}_{\varphi(i)}$ refines $\left\{ \pi(c) \in T_{\varphi(i)} \mid c \in \bigcup_{0 < l \leq i} (\tilde{B}_{\psi(i-l)} \cup \tilde{C}_{\varphi(i-l)}) , \pi \text{ a path} \right\}$.

We apply successively the Fraïssé sequence property twice. First, we obtain an index $\psi(i+1) > \psi(i)$ together with a \mathfrak{c} -morphism $\beta_{\varphi(i)}: T_{\varphi(i)} \rightarrow S_{\psi(i+1)}$ such that

$$\beta_{\varphi(i)} \circ \alpha_{\psi(i)} \simeq_{\tilde{B}_{\psi(i)}} \sigma_{\psi(i), \psi(i+1)}$$

and then we get an index $\varphi(i+1) > \varphi(i)$ together with a \mathfrak{c} -morphism $\alpha_{\psi(i+1)}: S_{\psi(i+1)} \rightarrow T_{\varphi(i+1)}$ such that

$$\alpha_{\psi(i+1)} \circ \beta_{\varphi(i)} \simeq_{\tilde{C}_{\varphi(i)}} \tau_{\varphi(i),\varphi(i+1)}$$

which finishes the inductive argument.

We will now check that the sequences of \mathfrak{c} -morphisms that we have just constructed induce a two-sided approximate intertwining. Let us first prove the following.

Claim. Let $i, l \in \mathbb{N}$. For any elements $b^-, b, b^+ \in \tilde{B}_{\psi(i)}$ such that $b^- \ll b \ll b^+$, we have

$$\begin{cases} \sigma_{\psi(i),\psi(i+l+1)}(b^-) \leq \beta_{\varphi(i+l)} \circ \tau_{\varphi(i),\varphi(i+l)} \circ \alpha_{\psi(i)}(b^+) \\ \beta_{\varphi(i+l)} \circ \tau_{\varphi(i),\varphi(i+l)} \circ \alpha_{\psi(i)}(b^-) \leq \sigma_{\psi(i),\psi(i+l+1)}(b^+) \end{cases}$$

Proof of the Claim. Note that $\sigma_{\psi(i),\psi(i+1)}(b^-) \leq \beta_{\varphi(i)} \circ \alpha_{\psi(i)}(b)$. Moreover, it follows from the construction of $\tilde{B}_{\psi(i+1)}$ (see (ii) above) that there exists $b_2 \in \tilde{B}_{\psi(i+1)}$ such that $\sigma_{\psi(i),\psi(i+1)}(b^-) \ll b_2 \ll \beta_{\varphi(i)} \circ \alpha_{\psi(i)}(b)$. Thus, we have that

$$\sigma_{\psi(i),\psi(i+2)}(b^-) \leq \beta_{\varphi(i+1)} \circ \alpha_{\psi(i+1)}(b_2) \leq \beta_{\varphi(i+1)} \circ \alpha_{\psi(i+1)}(\beta_{\varphi(i)} \circ \alpha_{\psi(i)}(b)).$$

Proceeding successively in this fashion, we obtain

$$\sigma_{\psi(i),\psi(i+l+1)}(b^-) \leq (\beta_{\varphi(i+l)} \circ \alpha_{\psi(i+l)}) \circ \dots \circ (\beta_{\varphi(i)} \circ \alpha_{\psi(i)})(b). \tag{4}$$

A similar argument involving the pair $\alpha_{\psi(i)}(b)$ and $\alpha_{\psi(i)}(b^+)$ shows that

$$(\alpha_{\psi(i+l)} \circ \beta_{\varphi(i+l-1)}) \circ \dots \circ (\alpha_{\psi(i+1)} \circ \beta_{\varphi(i)})(\alpha_{\psi(i)}(b)) \leq \tau_{\varphi(i),\varphi(i+l)}(\alpha_{\psi(i)}(b^+)). \tag{5}$$

Post-composing (5) by $\beta_{\varphi(i+l)}$ and combining it with (4) gives us the first inequality of the claim. The other inequality follows from a symmetric argument, which proves the claim.

Finally, let F be a finite subset of S_i . From the construction of the $\tilde{B}_{\psi(i)}$'s (see (i) above), there exists an index $i_F \geq i$ such that, for any $f', f \in F$ with $f' \ll f$ and any $j \geq i_F$, we can find $b^-, b, b^+ \in \tilde{B}_{\psi(j)}$ such that $\sigma_{i,\psi(j)}(f') \ll b^- \ll b \ll b^+ \ll \sigma_{i,\psi(j)}(f)$. Applying the claim, we get

$$\begin{cases} \sigma_{i,\psi(k+1)}(f') \leq \beta_{\varphi(k)} \circ \tau_{\varphi(j),\varphi(k)} \circ \alpha_{\psi(j)} \circ \sigma_{i,\psi(j)}(f) \\ \beta_{\varphi(k)} \circ \tau_{\varphi(j),\varphi(k)} \circ \alpha_{\psi(j)} \circ \sigma_{i,\psi(j)}(f') \leq \sigma_{i,\psi(k+1)}(f) \end{cases}$$

for any $j \geq i_F$ and any $k \geq j + 1$. In other words, $\sigma_{i,\psi(k+1)} \simeq_F \beta_{\varphi(k)} \circ \tau_{\varphi(j),\varphi(k)} \circ \alpha_{\psi(j)} \circ \sigma_{i,\psi(j)}$. Setting $\alpha'_j := \alpha_{\psi(j)} \circ \sigma_{j,\psi(j)}$ and $\beta'_k := \beta_{\varphi(k)} \circ \tau_{k,\varphi(k)}$, we obtain

$$\sigma_{i,\psi(k+1)} \simeq_F \beta'_k \circ \tau_{\varphi(j),k} \circ \alpha'_j \circ \sigma_{i,j}.$$


The second property of Definition 3.13 is proved by using a symmetric argument and we are done. \square

On the road towards our version of Theorem 2.7, we follow with some technical lemmas.

Lemma 3.22 (Universality). *Let $\mathfrak{c} \subseteq \text{Cu}$ be a Fraïssé category. Let $(S_i, \sigma_{i,j})_{i \in \mathbb{N}}$ be a Fraïssé sequence and let $(T_i, \tau_{i,j})_{i \in \mathbb{N}}$ be a \mathfrak{c} -inductive sequence of countably-based \mathfrak{c} -objects.*

Then, there exists a one-sided approximate intertwining $(\alpha_i: T_i \rightarrow S_{\varphi(i)})_i$.

Proof. We are going to construct the desired one-sided approximate intertwining recursively following a similar argument to that of Theorem 3.21. Recall that all the Cu -semigroups involved are countably-based by definition. Therefore, for each i , we can fix a \subseteq -increasing sequence $(B_n^i)_n$ of finite subsets of T_i such that its unions over n are sup-dense in T_i . Now, using the joint embedding property together with the fact that $(S_i, \sigma_{i,j})_i$ is a Fraïssé sequence, we construct a \mathfrak{c} -morphism $\alpha_0: T_0 \rightarrow S_{\varphi(0)}$.



Our aim is to construct a strictly increasing map $\varphi: \mathbb{N} \rightarrow \mathbb{N}$ together with large enough subsets \tilde{B}_i of T_i and \mathfrak{c} -morphisms $\alpha_{i+1}: T_{i+1} \rightarrow S_{\varphi(i+1)}$ such that $\sigma_{\varphi(i), \varphi(i+1)} \circ \alpha_i \simeq_{\tilde{B}_i} \alpha_{i+1} \circ \tau_{i, i+1}$. To do this, we proceed by induction from the initial data B_0^0, α_0 . Assume that


- the numbers $\varphi(0), \dots, \varphi(i)$
- the finite sets $B_0^0, \tilde{B}_1, \dots, \tilde{B}_{i-1}$
- the \mathfrak{c} -morphisms $\alpha_0, \dots, \alpha_i$

have been constructed for some $i \geq 0$. (By convention, we have fixed $\tilde{B}_{-1} := B_0^0$.)

Let us start by choosing $\tilde{B}_i \subseteq T_i$ such that

- (i) $B_i^i \subseteq \tilde{B}_i$.
- (ii) \tilde{B}_i is a 2-refinement of $\left\{ \tau_{i-l, i}(b) \in T_i \mid b \in \bigcup_{0 < l \leq i} \tilde{B}_{i-l} \right\}$.

We then use the near amalgamation property to construct \mathfrak{c} -morphisms $\xi_{i+1}^u, \theta_{i+1}$ such that the quadrilateral in the diagram below approximately commutes within \tilde{B}_i . Lastly, we use the Fraïssé sequence property to get a \mathfrak{c} -morphism ξ_{i+1}^d such that the triangle underneath approximately commutes within $\alpha_i(\tilde{B}_i)$. Define $\alpha_{i+1} := \xi_{i+1}^d \circ \theta_{i+1}$.



Hence, we have obtained a sequence of \mathfrak{c} -morphisms $(\alpha_i : T_i \rightarrow S_{\varphi(i)})_i$, and we are left to show that this is a one-sided approximate intertwining.

Claim. Let $i, j \in \mathbb{N}$. For any elements $b', b_1^-, b_1^+, b \in \tilde{B}_i$ such that $b' \ll b_1^- \ll b_1^+ \ll b$, we have

$$\begin{cases} \sigma_{\varphi(i), \varphi(i+j)} \circ \alpha_i(b') \leq \alpha_{i+j} \circ \tau_{i, i+j}(b) \\ \alpha_{i+j} \circ \tau_{i, i+j}(b') \leq \sigma_{\varphi(i), \varphi(i+j)} \circ \alpha_i(b) \end{cases}$$

Proof of the claim. Note that

$$\sigma_{\varphi(i), \varphi(i+1)} \circ \alpha_i(b') \leq \xi_{i+1}^d \circ \xi_{i+1}^u \circ \alpha_i(b_1^-) \leq \xi_{i+1}^d \circ \theta_{i+1} \circ \tau_{i, i+1}(b_1^+) = \alpha_{i+1} \circ \tau_{i, i+1}(b_1^+).$$

From the construction of the \tilde{B}_i 's (see (ii) above), we know that we can find $b_2^-, b_2^+ \in \tilde{B}_{i+1}$ such that $\tau_{i, i+1}(b_1^+) \ll b_2^- \ll b_2^+ \ll \tau_{i, i+1}(b)$. Therefore, we have

$$\sigma_{\varphi(i+1), \varphi(i+2)} \circ \alpha_{i+1} \circ \tau_{i, i+1}(b_1^+) \leq \xi_{i+2}^d \circ \xi_{i+2}^u \circ \alpha_{i+1}(b_2^-) \leq \alpha_{i+2} \circ \tau_{i+1, i+2}(b_2^+).$$

Proceeding successively in this fashion, we obtain elements $b_1^+, b_2^+, \dots, b_j^+$ such that $b_l^+ \in \tilde{B}_{i+l-1}$, and $b_l^+ \ll \tau_{i, i+l-1}(b)$ for any $l \leq j$. We compute

$$\begin{aligned} \sigma_{\varphi(i), \varphi(i+1)} \circ \alpha_i(b') &\leq \alpha_{i+1} \circ \tau_{i, i+1}(b_1^+) \\ \sigma_{\varphi(i+1), \varphi(i+2)} \circ \alpha_{i+1}(\tau_{i, i+1}(b_1^+)) &\leq \alpha_{i+2} \circ \tau_{i+1, i+2}(b_2^+) \\ \sigma_{\varphi(i+2), \varphi(i+3)} \circ \alpha_{i+2}(\tau_{i+1, i+2}(b_2^+)) &\leq \alpha_{i+3} \circ \tau_{i+2, i+3}(b_3^+) \\ &\vdots \\ \sigma_{\varphi(i+j-1), \varphi(i+j)} \circ \alpha_{i+j-1}(\tau_{i+j-2, i+j-1}(b_{j-1}^+)) &\leq \alpha_{i+j} \circ \tau_{i+j-1, i+j}(b_j^+). \end{aligned}$$

This proves the first inequality of the claim. The other inequality is shown using a symmetric argument.

Finally let F be a finite subset of T_i . From the construction of the \tilde{B}_i 's (see (i) above), there exists an index $i_F \geq i$ such that, for any $j \geq i_F$ and any pair $f', f \in F$ with $f' \ll f$, we can find $b', b_1^-, b_1^+, b \in \tilde{B}_j$ such that $\tau_{i, j}(f') \ll b' \ll b_1^- \ll b_1^+ \ll b \ll \tau_{i, j}(f)$. Applying the claim, we readily obtain

$$\begin{cases} \sigma_{\varphi(j),\varphi(k)} \circ \alpha_j \circ \tau_{i,j}(f') \leq \alpha_k \circ \tau_{i,k}(f) \\ \alpha_k \circ \tau_{i,k}(f') \leq \sigma_{\varphi(j),\varphi(k)} \circ \alpha_j \circ \tau_{i,j}(f) \end{cases}$$

for any $j \geq i_F$ and any $k > j$. In other words, $\alpha_k \circ \tau_{i,k} \simeq_F \sigma_{\varphi(j),\varphi(k)} \circ \alpha_j \circ \tau_{i,j}$, as required. \square


Lemma 3.23 (Homogeneity). *Let $\mathfrak{c} \subseteq \text{Cu}$ be a Fraïssé category and let $(S_i, \sigma_{i,j})_{i \in \mathbb{N}}$ be a Fraïssé sequence. Then, for any \mathfrak{c} -morphisms $\alpha: C \rightarrow S_l, \beta: C \rightarrow S_l$ and any finite subset $F \subseteq C$, there exists a two-sided approximate intertwining $(\eta_i: S_i \rightarrow S_{\varphi(i)}, \nu_i: S_i \rightarrow S_{\psi(i)})_{i \geq l}$ such that*

$$\sigma_{l,\psi(i)} \circ \alpha \simeq_F \nu_i \circ \sigma_{l,i} \circ \beta \quad \text{and} \quad \sigma_{l,\varphi(i)} \circ \beta \simeq_F \eta_i \circ \sigma_{l,i} \circ \alpha$$

for any $i \geq l$.

Proof. We will build the approximate intertwining following the structure of the previous proofs.

First, let us consider a 4-refinement \tilde{F} of F . (That is, \tilde{F} is a finite subset of C that contains F and is such that, for any $f', f \in F$ with $f' \ll f$, there exist $g', g^-, g^+, g \in \tilde{F}$ satisfying $f' \ll g' \ll g^- \ll g^+ \ll g \ll f$.) Then, using the near amalgamation property together with the Fraïssé sequence property, we construct \mathfrak{c} -morphisms $\nu_l, \eta_{\psi(l)}$ such that the following diagram approximately commutes.



Using the ideas and techniques from Theorem 3.21, it is readily verified that

$$\begin{cases} \sigma_{l,\varphi(l)} \circ \beta(f') \leq \sigma_{l,\varphi(l)} \circ \beta(g') \leq \eta_{\psi(l)} \circ \sigma_{l,\psi(l)} \circ \alpha(g) \leq \eta_{\psi(l)} \circ \sigma_{l,\psi(l)} \circ \alpha(f) \\ \sigma_{l,\psi(l)} \circ \alpha(f') \leq \sigma_{l,\psi(l)} \circ \alpha(g') \leq \nu_l \circ \beta(g) \leq \nu_l \circ \beta(f) \end{cases}$$

Finally, from the initial data $\tilde{B}_l := \alpha(\tilde{F}), \tilde{C}_l := \beta(\tilde{F}), \nu_l, \eta_{\psi(l)}$, one can construct a two-sided approximate intertwining following the proof of Theorem 3.21 (starting at l instead of 0). Such an intertwining will enjoy the desired properties. \square

We now have all the tools that we need to obtain Fraïssé limits in the category Cu . Let us first introduce the almost factorization property adapted to our setting.

Definition 3.24. Let $\mathfrak{c} \subseteq \mathfrak{d}$ be an inclusion of categories in Cu such that \mathfrak{d} has inductive limits. We say that the inclusion $\mathfrak{c} \subseteq \mathfrak{d}$ has the (Cuntz) almost factorization property

if, for any $C \in \mathfrak{c}$, any \mathfrak{c} -inductive system $(S_i, \sigma_{i,j})_{i,j \in I}$ with \mathfrak{d} -limit $(S, \sigma_{i,\infty})_i$, any \mathfrak{d} -morphism $\alpha: C \rightarrow S$, and any finite subset $F \subseteq C$, there exist an index i_F and a \mathfrak{c} -morphism $\alpha_F: C \rightarrow S_{i_F}$ satisfying $\sigma_{i_F, \infty} \circ \alpha_F \simeq_F \alpha$.

Adapting the definition of the Ind-completion (see, for example, [38, Chapter VI]), we define:

Definition 3.25. Let \mathfrak{c} be a subcategory of Cu . The *completion* of \mathfrak{c} , denoted by $\bar{\mathfrak{c}}$, is the subcategory of Cu whose

- (i) objects are Cu -limits of inductive sequences in \mathfrak{c} .
- (ii) morphisms are induced by *some* one-sided approximate intertwining.

More specifically, a Cu -morphism $\alpha: S \rightarrow T$ between $\bar{\mathfrak{c}}$ -objects S, T is a $\bar{\mathfrak{c}}$ -morphism if for any \mathfrak{c} -inductive sequence $(S_i, \sigma_{i,j})_i$ whose Cu -limit object is S , there exists a \mathfrak{c} -inductive sequence $(T_i, \tau_{i,j})_i$ whose Cu -limit object is T together with a one-sided approximate intertwining $(\alpha_i: S_i \rightarrow T_{\varphi(i)})_i$ in \mathfrak{c} which induces α in the sense of Proposition 3.11.

Remark 3.26. The following two properties of $\bar{\mathfrak{c}}$ are readily verified. (For example, they can be adapted from the results and references from [38, Chapter VI].)

- (i) The category $\bar{\mathfrak{c}}$ is a well-defined subcategory of Cu containing \mathfrak{c} as a subcategory.
- (ii) Any inductive sequence in \mathfrak{c} has an inductive limit in $\bar{\mathfrak{c}}$ which coincides with its inductive limit in Cu . In particular, any object $S \in \bar{\mathfrak{c}}$ can be written as the $\bar{\mathfrak{c}}$ -limit object of an inductive sequence in \mathfrak{c} .

Further, note that $\mathfrak{c} \subseteq \bar{\mathfrak{c}}$ may not satisfy the almost factorization property. However, this will be the case for all our examples.

Theorem 3.27. *Let \mathfrak{c} be a Fraïssé category. Then any Fraïssé sequence $(S_i, \sigma_{i,j})_i$ has a $\bar{\mathfrak{c}}$ -limit $(S, \sigma_{i,\infty})_i$ such that*

- (i) *S is unique up to isomorphism, that is, S does not depend on the Fraïssé sequence chosen.*
- (ii) *The set $\text{Hom}_{\bar{\mathfrak{c}}}(D, S)$ is nonempty whenever D is countably-based.*

Assume that \mathfrak{c} is contained in a category \mathfrak{d} where every \mathfrak{c} -inductive sequence has a limit, and where every \mathfrak{d} -object is the limit of a \mathfrak{c} -sequence. If $\mathfrak{c} \subseteq \mathfrak{d}$ satisfies the almost factorization property, then

- (iii) *For any $C \in \mathfrak{c}$, any $\alpha, \beta \in \text{Hom}_{\mathfrak{d}}(C, S)$ and any finite set $F \subseteq C$, there exists a \mathfrak{d} -isomorphism $\eta_F: S \xrightarrow{\cong} S$ such that $\eta_F \circ \alpha \simeq_F \beta$.*

Proof. (i) follows immediately from Theorem 3.21 together with Theorem 3.17, while (ii) follows as a combination of Lemma 3.22 and Proposition 3.11.

To see (iii), let $\alpha, \beta \in \text{Hom}_{\mathfrak{D}}(C, S)$ and let $F \subseteq C$ be a finite subset. Construct a 3-refinement \tilde{F} of F . Let $(S_i, \sigma_{i,j})_i$ be a Fraïssé sequence. Then, by the almost factorization property, there exist morphisms $\alpha_{\tilde{F}}, \beta_{\tilde{F}}: C \rightarrow S_l$ such that $\alpha \simeq_{\tilde{F}} \sigma_{l,\infty} \circ \alpha_{\tilde{F}}$ and $\beta \simeq_{\tilde{F}} \sigma_{l,\infty} \circ \beta_{\tilde{F}}$. Using Lemma 3.23 and Theorem 3.17, we see that there exists an isomorphism $\eta_F: S \rightarrow S$ satisfying the desired condition. \square

3.28 (*Fraïssé Categories of Cu^* -semigroups*). Following the discussion from Paragraph 3.18, we note that all the results above also do not use the fact that the underlying ordered monoids have a positive order. Thus, we have in fact developed a Fraïssé theory for Cu^* -semigroups.

C. C^* -algebras and Fraïssé categories of Cuntz semigroups

In this last subsection we study under which assumptions the functor Cu induces a Fraïssé category of Cuntz semigroups when applied to a Fraïssé category of separable C^* -algebras. A natural (but rather strong) assumption to consider is that Cu classifies $*$ -homomorphisms of the Fraïssé category $\mathfrak{c} \subseteq C^*$ at hand. We will see that, under an additional mild assumption, this is sufficient to deduce that $\text{Cu}(\mathfrak{c})$ is a Fraïssé category. We will conclude with some remarks on the link between these Fraïssé categories, where we discuss a weak converse of our result and ways to considerably strengthen the statement.

Let us start by recalling the definition of classifying morphisms. (See, for example, [43] or [17] for more details.)

Definition 3.29. Let \mathfrak{c} and \mathfrak{d} be subcategories of separable C^* -algebras. We say that the functor Cu classifies $*$ -homomorphisms from \mathfrak{c} to \mathfrak{d} if, for any A in \mathfrak{c} , any B in \mathfrak{d} and any scaled Cu -morphism $\alpha: \text{Cu}(A) \rightarrow \text{Cu}(B)$, there exists a $*$ -homomorphism $\chi: A \rightarrow B$, unique up to approximate unitary equivalence, such that $\text{Cu}(\chi) = \alpha$.

We will say that Cu classifies $*$ -homomorphisms of \mathfrak{c} whenever $\mathfrak{c} = \mathfrak{d}$.

In the setting of Definition 3.29 above, the term *scaled Cu -morphism* simply means that there exist strictly positive elements $s_A \in A$ and $s_B \in B$ such that $\alpha([s_A]) \leq [s_B]$. This notion fits in the general framework of *scales* in Cu -semigroups and their associated morphisms, as developed in [6, Section 4]. We refer the reader to that paper for a more detailed exposition.

Lemma 3.30. *Let \mathfrak{c} be a category of separable C^* -algebras which is either full or replete.¹ Assume that Cu classifies $*$ -homomorphisms of \mathfrak{c} . Then $\text{Cu}(\mathfrak{c})$ is a subcategory of Cu .*

¹ Most of the Fraïssé categories that we will consider are not full (for example, one usually considers injective maps). However, they will all be replete, i.e. for any $A \in \mathfrak{c}$ and any $*$ -isomorphism $\alpha: A \rightarrow B$, then both B and α are in \mathfrak{c} .

Proof. In both cases, the fact that the identity of every $\text{Cu}(c)$ -object is a $\text{Cu}(c)$ -morphism, and that both domain and codomain of a $\text{Cu}(c)$ -morphism are in $\text{Cu}(c)$ is immediate. The non-trivial part is to check that $\text{Cu}(\mathfrak{c})$ is closed under composition of morphisms.

Let $\varphi: A \rightarrow B$ and $\psi: B' \rightarrow C$ be $*$ -homomorphisms such that $\text{Cu}(B) = \text{Cu}(B')$. Write $\alpha := \text{Cu}(\psi) \circ \text{Cu}(\varphi): \text{Cu}(A) \rightarrow \text{Cu}(C)$. Note that α is still a scaled morphism, that is, it maps the class of a strictly positive element in A below the class of a strictly positive element in C .

Assume that \mathfrak{c} is full. Since Cu classifies $*$ -homomorphisms of \mathfrak{c} , we know that there exists a $*$ -homomorphism $\chi_\alpha: A \rightarrow C$, which is a \mathfrak{c} -morphism by fullness of \mathfrak{c} , such that $\text{Cu}(\chi_\alpha) = \alpha$. In other words, $\alpha \in \text{Cu}(\mathfrak{c})$.

Now assume that \mathfrak{c} is replete. Since Cu classifies $*$ -homomorphisms of \mathfrak{c} and $\text{Cu}(B) = \text{Cu}(B')$, we know that there exists a $*$ -isomorphism $\chi: B \cong B'$ lifting $\text{id}_{\text{Cu}(B)}$. Since \mathfrak{c} is replete, we also know that χ is in fact a \mathfrak{c} -morphism. Now define $\chi_\alpha := \psi \circ \chi \circ \varphi: A \rightarrow C$. From construction, χ_α is a \mathfrak{c} -morphism and $\text{Cu}(\chi_\alpha) = \alpha$. In other words, $\alpha \in \text{Cu}(\mathfrak{c})$. \square

Theorem 3.31. *Let \mathfrak{c} be a category of separable C^* -algebras which is either full or replete. Assume that Cu classifies $*$ -homomorphisms of \mathfrak{c} .*

If \mathfrak{c} is a Fraïssé category whose Fraïssé limit is A , then $\text{Cu}(\mathfrak{c})$ is a Fraïssé category whose Fraïssé limit is $\text{Cu}(A)$.

Proof. The joint embedding property immediately follows applying the functor Cu .

Let us prove the near amalgamation property in $\text{Cu}(\mathfrak{c})$. First, we note the following fact.

Fact. Let A be a C^* -algebra and let $F \subseteq \text{Cu}(A)$ be a finite subset. Then there exists a small enough $\varepsilon > 0$ such that, for any $[a], [b] \in F$ with $[a] \ll [b]$, then $[a] \ll [(b - \varepsilon)_+]$.

Let $\text{Cu}(\phi_1): \text{Cu}(A) \rightarrow \text{Cu}(B_1)$ and $\text{Cu}(\phi_2): \text{Cu}(A') \rightarrow \text{Cu}(B_2)$ be \mathfrak{c} -morphisms with $\text{Cu}(A) = \text{Cu}(A')$. Using the same arguments as above, we may assume $A = A'$. Let $F \subseteq \text{Cu}(A)$ be finite and let $\varepsilon > 0$ be the constant given by the previous fact. Now, using the near amalgamation property in \mathfrak{c} , we know that there exist \mathfrak{c} -morphisms $\psi_1: B_1 \rightarrow C$ and $\psi_2: B_2 \rightarrow C$ such that $d_{C^*}(\psi_1 \circ \phi_1, \psi_2 \circ \phi_2) < \varepsilon$. We are left to show that $\text{Cu}(\psi_1 \circ \phi_1) \simeq_F \text{Cu}(\psi_2 \circ \phi_2)$. Let $[a], [b] \in F$ be such that $[a] \ll [b]$. We know that $\|\psi_1 \circ \phi_1(b) - \psi_2 \circ \phi_2(b)\| < \varepsilon$ and that $[a] \ll [(b - \varepsilon)_+]$ which implies that

$$\begin{cases} \text{Cu}(\psi_1 \circ \phi_1)([a]) \ll \text{Cu}(\psi_1 \circ \phi_1)([(b - \varepsilon)_+]) \leq \text{Cu}(\psi_2 \circ \phi_2)([b]) \\ \text{Cu}(\psi_2 \circ \phi_2)([a]) \ll \text{Cu}(\psi_2 \circ \phi_2)([(b - \varepsilon)_+]) \leq \text{Cu}(\psi_1 \circ \phi_1)([b]) \end{cases}$$

as desired.

That $\text{Cu}(\mathfrak{c})$ is separable is shown similarly, by also using the fact that any C^* -algebra in \mathfrak{c} is separable, and thus gives rise to a countably-based Cu -semigroup.

We deduce that $\text{Cu}(\mathfrak{c})$ is a Fraïssé category. Further, given any Fraïssé sequence in \mathfrak{c} , we can use that Cu classifies $*$ -homomorphisms of \mathfrak{c} to show that the induced sequence

in Cu is Fraïssé in $\text{Cu}(\mathfrak{c})$. Thus, by continuity of the functor Cu , it is readily checked that $\text{Cu}(A)$ is the Fraïssé limit of $\text{Cu}(\mathfrak{c})$. \square

Some remarks are in order:

Remark 3.32. (i) Assume that \mathfrak{c} is such that Cu classifies $*$ -homomorphisms from \mathfrak{c} to stable rank one C^* -algebras or, more generally, to a category \mathfrak{d} closed under ultraproducts. Then, it follows from [43, Theorem 3.3.1] that the previous theorem has a weak converse: $\text{Cu}(\mathfrak{c})$ is Fraïssé if and only if \mathfrak{c} is Fraïssé with respect to approximate unitary equivalences (in the sense of Remark 2.8).

(ii) The assumption of classification of $*$ -homomorphisms is rather strong in general. For instance, it is proved in [17] that the functor Cu does not classify $*$ -homomorphisms of circle algebras. Nevertheless, we do not use the full force of the assumption, neither in Lemma 3.30 nor in Theorem 3.31. Explicitly, one only needs to assume the following much weaker condition:

For every $A, A', B, B' \in \mathfrak{c}$ such that $\text{Cu}(A) \cong^\alpha \text{Cu}(A')$ and $\text{Cu}(B) \cong^\beta \text{Cu}(B')$, and any \mathfrak{c} -morphism $\varphi: A \rightarrow B$, there exists a \mathfrak{c} -morphism $\phi: A' \rightarrow B'$ such that $\text{Cu}(\phi) \circ \alpha = \beta \circ \text{Cu}(\varphi)$.

Note that, in particular, this holds whenever our category has a single object. (See Remark 4.37.)

To obtain a weak converse (as in (i)), the additional property that one needs is:

For every C^ -algebra A in \mathfrak{c} , any finite set $F \subseteq A$ and any $\varepsilon > 0$ there exists a finite subset $G \subseteq \text{Cu}(A)$ such that, whenever two \mathfrak{c} -morphisms $\varphi_1, \varphi_2: A \rightarrow B$ satisfy $\text{Cu}(\varphi_1) \simeq_G \text{Cu}(\varphi_2)$, then there exists $u \in \tilde{B}$ such that $\|u^* \varphi_1(x)u - \varphi_2(x)\| < \varepsilon$ for every $x \in F$.*

(iii) A number of examples of C^* -algebraic Fraïssé subcategories \mathfrak{c} have injective $*$ -homomorphisms as morphisms. Note that the theorem above does not imply that \mathfrak{c} induces a category $\text{Cu}(\mathfrak{c})$ whose morphisms are order-embeddings, since injective $*$ -homomorphisms do not generally induce injective Cu -morphisms. For example, the diagonal map $\mathbb{C} \oplus \mathbb{C} \rightarrow M_2(\mathbb{C})$ is injective, but the induced Cu -morphism maps both $[(1, 0)]$ and $[(0, 1)]$ to $[1 \oplus 0]$. Conversely, a $*$ -homomorphism that induces an order-embedding may not be injective.

4. Examples

In this section we exhibit natural examples of (Cuntz) Fraïssé categories together with their Fraïssé limit. This allows us to deduce several generic properties about the Cu -semigroups at play. More explicitly, we show that the Cuntz semigroup of any UHF-

algebra and that the Cuntz semigroup of the universal AF-algebra are the Fraïssé limits of some well-chosen Fraïssé categories. We also show that there are countably many Fraïssé categories of elementary Cu-semigroups whose Fraïssé limits are not purely infinite, non-stably finite, simple Cu-semigroups. Finally, we prove that both Cu-semigroups $\text{Lsc}(X, \overline{\mathbb{N}})$, where X is either the Cantor set $2^{\mathbb{N}}$ or the pseudo-arc \mathbb{P} , can also be written as Fraïssé limits.

In all our examples the inclusion $\mathfrak{c} \subseteq \overline{\mathfrak{c}}$ satisfies the almost factorization property.

A. Dimension Cu-semigroups of infinite type as Fraïssé limits

In what follows, we show that the Cuntz semigroup of any UHF-algebra arises as a Fraïssé limit. Following [5], recall that a Cu-semigroup S is said to be *simplicial* whenever $S \cong \overline{\mathbb{N}}^r$ for some $r \in \mathbb{N}$, and that an inductive limit of simplicial Cu-semigroups is called a *dimension Cu-semigroup*.

Let p be a prime number and consider the semigroup $S_p := \mathbb{N}[\frac{1}{p}] \sqcup (0, \infty]$, where the mixed sum and mixed order are defined as follows:

- (+) $x_c + y_s = x_s + y_s \in (0, \infty]$ for any pair $x_c = k/p^l \in \mathbb{N}[\frac{1}{p}]$ and $y_s \in (0, \infty]$.
- (≤) retaining the same notation, $x_s \leq x_c \ll x_c \leq x_s + \varepsilon$ for any $\varepsilon > 0$.


Recall that S_p is the Cuntz semigroup of the UHF-algebra M_{p^∞} . (See, e.g. [5, Proposition 7.4.3].)

4.1. The category \mathfrak{s}_p is the category whose (single) object is $\overline{\mathbb{N}}$ and whose morphisms are powers of p . (That is, a \mathfrak{s}_p -morphism is given by $\overline{\mathbb{N}} \xrightarrow{\times p^k} \overline{\mathbb{N}}$ for some $k \in \mathbb{N}$.)

Theorem 4.2. *The category \mathfrak{s}_p is Fraïssé and its limit is S_p .*

Proof. The joint embedding property and separability are clear and the amalgamation property follows from the commutativity in $\text{Hom}_{\mathfrak{s}_p}(\overline{\mathbb{N}}, \overline{\mathbb{N}})$. Therefore \mathfrak{s}_p is a Fraïssé category.

Let us now show that the sequence given by $(\overline{\mathbb{N}}, \xrightarrow{\times p})_n$ is Fraïssé. For any \mathfrak{s}_p -morphism $\alpha: \overline{\mathbb{N}} \xrightarrow{\times p^k} \overline{\mathbb{N}}$ there exists a large enough index (consisting of $k + 1$ steps further from the domain of α) such that the following diagram commutes



It follows that the sequence is Fraïssé and, hence, its limit S_p is the Fraïssé limit of \mathfrak{s}_p . □

Remark 4.3. An analogous statement to that of Theorem 4.2 works for the Cuntz semigroup of any UHF-algebra. However, such semigroups may not be as easily computed as those of infinite type. Also, let us remark that we could have deduced the previous result from Theorem 3.31.

B. The universal dimension Cu-semigroup as a Fraïssé limit

Based on the work done in [32], we now exhibit a *universal dimension Cu-semigroup* \mathcal{S} as a Fraïssé limit of a well-chosen Fraïssé category \mathfrak{s}_{\dim} containing all simplicial Cu-semigroups. As a matter of fact, we will see that \mathcal{S} is the concrete Cuntz semigroup of the universal AF-algebra \mathcal{A} constructed in [32]. Therefore, \mathcal{S} enjoys analogous properties to those of \mathcal{A} , but in the category Cu. Before we begin, we need to recall and prove some results about *retractions* in the category Cu.

4.4 (Retractions). We aim to define a Fraïssé category \mathfrak{s}_{\dim} that will be built out of retractable Cu-morphisms. Recall that an ordered monoid morphism $\alpha: S \rightarrow T$ between two Cu-semigroups is said to be a *generalized Cu-morphism* if α preserves suprema of increasing sequences.

Definition 4.5 ([51, Definition 3.14]). Let S, T be Cu-semigroups. We say that S is a *retract* of T if there exists a Cu-morphism $\iota: S \rightarrow T$ together with a generalized Cu-morphism $\rho: T \rightarrow S$ such that $\rho \circ \iota = \text{id}_S$.

We say that ι is *retractable* and that ρ *retracts* ι .

Following [21], a submonoid I of a Cu-semigroup S is said to be an *ideal* if I is downward-hereditary and closed under suprema of increasing sequences. Given any ideal I of S , one can construct the quotient Cu-semigroup S/I ; see [5, Lemma 5.1.2]. Ideals and quotients of a C^* -algebra A are in bijective correspondence with the ideals and quotients of its Cuntz semigroup $\text{Cu}(A)$; see [5, Section 5].

Proposition 4.6. *Let S be a Cu-semigroup.*

- (i) *If S is countably-based and satisfies (O5)-(O7), then any ideal of S is a retract of S .*
- (ii) *If $(S, \sigma_{i,\infty})_i$ is the limit of an inductive sequence $(S_i, \sigma_{i,j})_i$ such that $\sigma_{i,i+1}$ is retractable for each $i \in \mathbb{N}$, then $\sigma_{i,\infty}$ is retractable for any $i \in \mathbb{N}$.*

Proof. (i) Let I be an ideal of a countably-based Cu-semigroup S and let $\iota: I \rightarrow S$ be the canonical order-embedding. Observe that I is a countably-based Cu-semigroup. Thus, I has a greatest element that we denote by ∞_I . Using [2, Theorem 2.4], we know that the infimum $x \wedge \infty_I$ exists for any $x \in S$. Further, such an infimum is always in I , since $x \wedge \infty_I \leq \infty_I$.

Now consider the map $\rho: S \rightarrow I$ that sends $x \mapsto x \wedge \infty_I$. It follows directly from [2, Theorem 2.5] that ρ is a generalized Cu-morphism that satisfies $\rho \circ \iota = \text{id}_I$, as desired.

(ii) Let $(S_i, \sigma_{i,j})_i$ be an inductive sequence with retractable maps, and let $(S, \sigma_{i,\infty})_i$ be its inductive limit. Denote by $\rho_{i+1,i}$ the retract of $\sigma_{i,i+1}$. For each pair $i \leq j$, let $\rho_{j,i}: S_j \rightarrow S_i$ be the composition $\rho_{i+1,i} \circ \dots \circ \rho_{j,j-1}$, which trivially retracts $\sigma_{i,j}$.

Now fix $i \in \mathbb{N}$ and let $s \in S$. Using the characterization of sequential inductive limits in the category Cu (see e.g. [43, Section 2.1]), we know that there exists a sequence $(s_n)_{n \geq i}$ with $s_n \in S_n$ such that $\sigma_{n,n+1}(s_n) \ll s_{n+1}$ and $\sup_{n \geq i} \sigma_{n,\infty}(s_n) = s$. Applying the retract at each step gives us that $s_n \leq \rho_{n+1,n}(s_{n+1})$ and, hence, $\rho_{n,i}(s_n) \leq \rho_{n+1,i}(s_{n+1})$ for any $n \geq i$.

Set $\rho_{\infty,i}(s) := \sup_{n \geq i} \rho_{n,i}(s_n)$. A standard argument shows that $\rho_{\infty,i}: S \rightarrow S_i$ does not depend on the sequence $(s_n)_{n \geq i}$ chosen. This also proves that $\rho_{\infty,i}$ is additive, preserves the order, and suprema of increasing sequences. (See e.g. the argument in [15, Lemma 3.12] or in [1, Lemma 7.3].)

Finally, let $x \in S_i$ and let $(x_n)_n$ be a \ll -increasing sequence in S_i with supremum x . For each $n \geq i$, set $s_n := \sigma_{i,n}(x_n)$. Since $\rho_{\infty,i}$ does not depend on the sequence $(s_n)_n$ chosen, we have $\rho_{\infty,i}(\sigma_{i,\infty}(x)) = \sup_n \rho_{n,i}(s_n) = \sup_n \rho_{n,i}(\sigma_{i,n}(x_n)) = \sup_n x_n = x$ which ends the proof. \square

The following is based on [32, Proposition 3.8].

Proposition 4.7. *Let \mathfrak{c} be a subcategory of Cu closed under direct sums. Let $(S_i, \sigma_{i,j})_i$ be an inductive sequence in \mathfrak{c} with $\bar{\tau}$ -limit $(S, \sigma_{i,\infty})_i$.*

Then, there exists an inductive sequence $(T_i, \tau_{i,j})_i$ in \mathfrak{c} with $\bar{\tau}$ -limit $(T, \tau_{i,\infty})_i$ such that $\tau_{i,j}$ is retractable for every pair $i \leq j$, and such that $S \cong T/J$ for some ideal J of T .

Proof. Let $i \in \mathbb{N}$ and set $T_i := S_0 \oplus \dots \oplus S_{i-1} \oplus S_i$. Let $\tau_{i,i+1}: T_i \rightarrow T_{i+1}$ be the Cu -morphism defined by

$$\tau_{i,i+1}(s_0, \dots, s_{i-1}, s_i) := (s_0, \dots, s_{i-1}, s_i, \sigma_{i,i+1}(s_i)).$$

Note that $\tau_{i,i+1}: T_i \rightarrow T_{i+1}$ is retractable by the projection $\pi_{i+1,i}: T_{i+1} \rightarrow T_i$ onto the first i components of T_i . More particularly, we have $\pi_{i+1,i}(s_0, \dots, s_i, s_{i+1}) = (s_0, \dots, s_i)$. It follows that every composition $\tau_{i,j} := \tau_{j-1,j} \circ \dots \circ \tau_{i,i+1}$ is also retractable for any $j \geq i + 1$.

Let $(T, \tau_{i,\infty})_i$ be the limit of the inductive system of $(T_i, \tau_{i,j})_i$ and set

$$J_0 := \{x \in T \mid x = \tau_{i,\infty}(s_0, \dots, s_{i-1}, 0) \text{ for some } i \in \mathbb{N} \text{ and } s_0 \in S_0, \dots, s_{i-1} \in S_{i-1}\}.$$

Note that J_0 is a submonoid of T . Thus, we can construct its sup-closure $J := \overline{J_0}^{\text{sup}}$. (See e.g. [50, Definition 4.6].) Further, given $y \in J$ and $x \in T$ such that $x \leq y$, take $x' \in T$ such that $x' \ll x$. Since J is the closure of J_0 and $x' \ll x$, there exist $i \in \mathbb{N}$, $t \in T_i$, and $(s_0, \dots, s_{i-1}, 0) \in T_i$ such that

$$x' \ll \tau_{i,\infty}(t) \ll x, \quad \text{and} \quad t \ll (s_0, \dots, s_{i-1}, 0).$$

In particular, we get $t \in S_0 \oplus \dots \oplus S_{i-1} \oplus 0$ and, hence, $\tau_{i,\infty}(t) \in J_0$. This shows that x can be written as the supremum of elements in J_0 or, equivalently, that x is in J . Thus, J is an ideal of T . It is now clear that $S \cong T/J$. \square

4.8. *The category \mathfrak{s}_{\dim} is the category whose objects are the simplicial Cu-semigroups and whose morphisms are retractable Cu-morphisms.*

Theorem 4.9. *The category \mathfrak{s}_{\dim} is Fraïssé and its limit \mathcal{S} is the Cuntz semigroup of the universal AF-algebra.*

Proof. First, note that any retractable Cu-morphism between simplicial Cu-semigroups can always be retracted by a Cu-morphism. Indeed, let $\iota: S \rightarrow T$ be a Cu-morphism between simplicial Cu-semigroups, and let $\rho: T \rightarrow S$ be a generalized Cu-morphism such that $\rho \circ \iota = \text{id}_S$.

Define $\rho': T \rightarrow S$ by $\rho'(x) := \rho(x \wedge \infty \iota(1_S))$. This map is the composition of generalized Cu-morphisms, and so it itself is a generalized Cu-morphism. To see that it preserves the \ll -relation, simply take $x \in T$ such that $x \ll x$. Then, one gets $x \wedge \infty \iota(1_S) \ll x \wedge \infty \iota(1_S)$ and, consequently, that $x \wedge \infty \iota(1_S) \leq n \iota(1_S)$ for some $n \geq 1$. This implies $\rho'(x) \leq n 1_S$, which is equivalent to $\rho'(x) \ll \rho'(x)$, as desired.

Further, since Cu-morphisms map compact elements to compact elements, all maps involved can be seen as scaled Cu-morphisms between finite-dimensional C^* -algebras of appropriate size.

The result now follows from [32] and Theorem 3.31, combined with the well-known fact that the scaled Cuntz semigroup classifies $*$ -homomorphisms of finite dimensional C^* -algebras. \square

Corollary 4.10. *Let \mathcal{S} be the universal dimension Cu-semigroup and let S be a (countably-based) dimension Cu-semigroup. Then, there exists a surjective Cu-morphism $\mathcal{S} \rightarrow S$.*

Proof. By Proposition 4.7 we know that S is isomorphic to a quotient of the form T/J , where T is the limit of some inductive system in \mathfrak{s}_{\dim} . Further, by universality of the Fraïssé limit, there exists a retractable Cu-morphism $\iota: T \rightarrow \mathcal{S}$ whose retract $\rho: \mathcal{S} \rightarrow T$ is a Cu-morphism. The composition of ρ with the quotient map $T \rightarrow T/J$ gives the desired Cu-morphism. \square

C. Elementary Fraïssé categories

As defined in [52, 8.1], a Cu-semigroup S is said to be *elementary* if S is simple and contains a minimal, nonzero element that is finite.² Assume further that S has finitely many elements. If S satisfies (O5) and (O6), it follows from [5, Proposition 5.1.19] that

² This differs slightly from the definition given in [5], and was adjusted to not include the Cuntz semigroup of simple, purely infinite C^* -algebras.

$S \cong E_n := \{0, 1, \dots, n, \infty\}$ for some $n \in \mathbb{N}$. It is well known that no such Cu-semigroup is the Cuntz semigroup of a C^* -algebra.³

The following lemma shows that the Cu-morphisms between elementary Cu-semigroups are well understood. By an *order-embedding* between Cu-semigroups we will always mean a Cu-morphism that is also an order-embedding.

Lemma 4.11. *Let $n, m \in \mathbb{N}$. A map $\alpha: \{0, 1, \dots, n, \infty\} \rightarrow \{0, 1, \dots, m, \infty\}$ is*

- (i) *a Cu-morphism if and only if $(n + 1)\alpha(1) = \infty$ and $\alpha(k1) = k\alpha(1)$ for every $k \leq n$.*
- (ii) *an order-embedding if and only if $\alpha(1) \neq 0$ and $n\alpha(1) \neq \infty$.*

In particular, there exist choices of natural numbers $n, m \in \mathbb{N}$ such that there is no or only one order-embedding from E_n to E_m . For example, we know that $\alpha(1) = k1$ must satisfy $m/(n + 1) < k \leq m/n$. Thus, if $m = n(n + 1)$, we get that the only order-embedding from E_n to E_m is $\alpha(1) = (n + 1)1$.

4.12. *The category \mathfrak{e}_∞ is the category whose objects are all elementary Cu-semigroups satisfying (O5) and (O6) with finitely many elements, and whose morphisms are all nonzero Cu-morphisms.*

Proposition 4.13. *The category \mathfrak{e}_∞ is Fraïssé.*

Proof. The category \mathfrak{e}_∞ contains countably many objects, and the morphisms between them are given by multiplication. It follows that \mathfrak{e}_∞ is separable.

Further, given $n_1, n_2 \in \mathbb{N}$, it follows from Lemma 4.11 that the maps $\alpha_1: E_{n_1} \rightarrow E_{n_1 n_2}$ and $\alpha_2: E_{n_2} \rightarrow E_{n_1 n_2}$ given by $1 \mapsto n_2 1$ and $1 \mapsto n_1 1$ respectively are order-embeddings (in particular, nonzero Cu-morphisms). This shows that the category \mathfrak{e}_∞ has the joint embedding property.

Finally, given any pair of nonzero Cu-morphisms $\alpha_1: E_n \rightarrow E_m$ and $\alpha_2: E_n \rightarrow E_m$, one can simply consider the map $\beta: E_m \rightarrow E_m$ given by $\beta(1) := \infty$. This Cu-morphism satisfies $\beta \circ \alpha_1 = \beta \circ \alpha_2$, which implies that the category \mathfrak{e}_∞ is Fraïssé. \square

Corollary 4.14. *The Fraïssé limit of \mathfrak{e}_∞ is $\{0, \infty\}$, that is, the Cuntz semigroup of any purely infinite simple C^* -algebra.*

Proof. Let $(S_i, \sigma_{i,j})_i$ be a Fraïssé sequence of \mathfrak{e}_∞ . For any $i \in \mathbb{N}$, let $\alpha: S_i \rightarrow \{0, \infty\}$ be the morphism that maps every nonzero element to ∞ . By definition, there exists $\beta: \{0, \infty\} \rightarrow S_j$ such that $\sigma_{i,j} = \beta \circ \alpha$. In other words, every element in S_i becomes idempotent eventually.

³ A quick proof of this fact goes as follows: Assume for the sake of contradiction that there exists A with $\text{Cu}(A)$ isomorphic to E_n for some n . Then, A must be simple and weakly purely infinite and, consequently, purely infinite. The Cuntz semigroup of any simple, purely infinite C^* -algebra is isomorphic to $\{0, \infty\}$, a contradiction.

Recall that a Cu-semigroup is called idempotent if $x = 2x$ for each element x . The only nonzero, simple, idempotent Cu-semigroup is $\{0, \infty\}$. Thus, since the morphisms in the sequence are nonzero, the Fraïssé limit of \mathfrak{e}_∞ is isomorphic to $\{0, \infty\}$. \square

Remark 4.15. The subcategory obtained by considering all order-embeddings instead of all nonzero Cu-morphisms is not Fraïssé, since it fails to satisfy the joint embedding property. For example, one can check that the morphisms $\alpha_1, \alpha_2: \{0, 1, \infty\} \rightarrow \{0, \dots, 6, \infty\}$ given by $\alpha_1(1) = 4 \cdot 1$ and $\alpha_2(1) = 5 \cdot 1$ cannot be amalgamated.

4.16. *The category \mathfrak{e}_n (for a fixed $n \geq 1$) is the category whose objects are all the elementary semigroups of the form E_{n^k} for some $k \in \mathbb{N}$, and whose morphisms are all the order-embeddings given by multiplication by a power of n .*

Note that not all maps given by powers of n are Cu-morphisms. As an example, set $n = 2$. Then, we know from Lemma 4.11 that multiplying by 2 does not give rise to a Cu-morphism $E_2 \rightarrow E_8$. In fact, it follows from Lemma 4.11 that for any given pair $k \leq s$ there exists a unique order-embedding $E_{n^k} \rightarrow E_{n^s}$ given by a power of n , namely $1 \mapsto n^{s-k}$.

Proposition 4.17. *The category \mathfrak{e}_n is Fraïssé for any $n \in \mathbb{N}$.*

Proof. That the category \mathfrak{e}_n is separable and satisfies the joint embedding property is proven similarly as in the proof of Proposition 4.13.

To see that the category \mathfrak{e}_n has amalgamation, let $\alpha_1: E_{n^{s_1}} \rightarrow E_{n^{s_2}}$ and $\alpha_2: E_{n^{s_1}} \rightarrow E_{n^{s_2}}$ be \mathfrak{e}_n -morphisms. From the joint embedding property, we may assume that $s_1 = s_2$. Now, we know from the comments above that these two maps must be the same. Thus, the category \mathfrak{e}_n has amalgamation. \square

Recall that a Cu-semigroup S is *stably finite* if $x \neq x + y$ for every nonzero $y \in S$ whenever $x \ll z$ for some $z \in S$. Also, extending the definition of [39], let us say that S is *(n-)weakly purely infinite* if $nx = 2(nx)$ for every $x \in S$.

Corollary 4.18. *The Fraïssé limit \mathcal{E}_n of \mathfrak{e}_n is a simple, non-stably finite, not weakly purely infinite Cu-semigroup whose order is total.*

Proof. The Fraïssé sequence of \mathfrak{e}_n consists of simple Cu-semigroups, so its limit must be simple. Further, since each Cu-semigroup has a total order, so does the limit.

A simple Cu-semigroup is not stably finite if and only if its greatest element, denoted by ∞ , satisfies $\infty \ll \infty$. The Fraïssé limit of \mathfrak{e}_n admits a nonzero Cu-morphism α from $\{0, 1, \dots, n, \infty\}$. In particular, $\alpha(\infty) = \infty$ and, since α preserves the \ll -relation, one has $\infty \ll \infty$ in the limit. This shows that the limit is not stably finite.

Finally, the limit cannot be weakly purely infinite. Indeed, for any $k \in \mathbb{N}$ we can use Theorem 3.27 to find an order-embedding α from $\{0, 1, \dots, n^k, \infty\}$ to the limit. Then,

$\alpha(1)$ satisfies $l\alpha(1) \neq (l-1)\alpha(1)$ for any $l \leq n^k$. This shows that there is no global bound on the idempotency of the elements, which implies that the limit is not weakly purely infinite. \square

Remark 4.19. Arguing similarly as in Theorem 4.2, we can deduce that, given a prime number p , the Fraïssé limit \mathcal{E}_p of \mathfrak{e}_p is a truncated version of the dimension semigroup of infinite type $S_p = \mathbb{N}[\frac{1}{p}] \sqcup (0, \infty]$. Explicitly, we have

$$\mathcal{E}_p \cong \left\{ x \in \mathbb{N} \left[\frac{1}{p} \right] \sqcup (0, 1] \mid x \leq 1 \right\} \cup \{\infty\}$$

where the order and sum between two elements x, y are defined as in S_p , with the exception that $x + y = \infty$ whenever this sum is (strictly) greater than 1_c in S_p .

Question 4.20. It can be checked that \mathcal{E}_p satisfies all the known extra axioms (O5)-(O8) that the Cuntz semigroup of any C^* -algebra satisfies. Thus, it is natural to ask: Is \mathcal{E}_p the Cuntz semigroup of some C^* -algebra A ?

Note that such a C^* -algebra A would be simple, not stably finite, and not purely infinite. However, A is not the C^* -algebra B constructed by Rørdam in [47], since $\text{Cu}(B)$ does not satisfy the Corona Factorization Property (see [11, Theorem 5.8]) but \mathcal{E}_p does.

D. The Cantor set and the pseudo-arc

Fraïssé Theory allows one to rewrite well-known topological spaces such as the Cantor set $2^{\mathbb{N}}$ and the pseudo-arc \mathbb{P} as Fraïssé limits and, in particular, to generically (re)prove some interesting facts about them, such as universality and homogeneity. We refer the reader to [9,35,40] for more details. Following these results, we show that the Cu -semigroups $\text{Lsc}(2^{\mathbb{N}}, \overline{\mathbb{N}})$ and $\text{Lsc}(\mathbb{P}, \overline{\mathbb{N}})$ are Fraïssé limits of well-chosen categories of Cuntz semigroups.

4.21 (Lower-semicontinuous functions). We begin by recalling some facts about monoids of lower-semicontinuous functions, which constitute a great source of example of abstract Cuntz semigroups. For instance, it is known that the monoid of lower-semicontinuous functions from a compact, metric (or, more generally, hereditarily Lindelöf, locally compact, and Hausdorff) space X to $\overline{\mathbb{N}}$, denoted by $\text{Lsc}(X, \overline{\mathbb{N}})$, is a Cu -semigroup when equipped with pointwise addition and order. (See [34, Proposition 1.16], and also [57, Corollary 4.22] and [4, Theorem 5.17].) In the specific case where X is a compact one-dimensional CW-complex, then $\text{Lsc}(X, \overline{\mathbb{N}})$ is in fact the concrete Cuntz semigroup of the C^* -algebra $C(X)$. (See [45,15] and also [4], [18] for other examples of concrete Cuntz semigroup of C^* -algebras that can be expressed as lower-semicontinuous functions.)

Further, as noted in [46] for the interval, and in [15] for compact one-dimensional CW-complexes (which include finite discrete sets, the interval and the circle), the set $\text{Hom}_{\text{Cu}}(\text{Lsc}(X, \overline{\mathbb{N}}), T)$ of Cu -morphisms can be equipped with the following Cu -metric

$$d_{\text{Cu}}(\alpha, \beta) := \inf \{r > 0 \mid \forall V \in \mathcal{O}(X), \alpha(1_V) \leq \beta(1_{V_r}) \text{ and } \beta(1_V) \leq \alpha(1_{V_r})\}$$

where $\mathcal{O}(X) := \{ \text{Open sets of } X \}$ and $V_r := \bigcup_{x \in V} B_r(x)$, that is, V_r is the set of points with distance less than r from V .

As proved implicitly in [58, Lemma 4.8] for the interval case and explicitly in [15] for compact one-dimensional CW-complexes, there is a strong link between the metric d_{Cu} and finite-set comparison:

Proposition 4.22 ([15, Proposition 5.6]). *Let X be a compact one-dimensional CW-complex and let $\{\overline{U}_k\}_1^n$ be a finite closed cover of X induced by an equidistant partition of size $1/n$.*

Let $F_n := \{f \in \text{Lsc}(X, \overline{\mathbb{N}}) \mid f|_{U_k} \text{ is constant for any } k \in \{1, \dots, n\}\}$. For any pair of morphisms $\alpha, \beta \in \text{Hom}_{\text{Cu}}(\text{Lsc}(X, \overline{\mathbb{N}}), T)$, we have

- (i) $\alpha \simeq_{F_n} \beta$ implies $d_{\text{Cu}}(\alpha, \beta) \leq 2/n$.
- (ii) $d_{\text{Cu}}(\alpha, \beta) \leq 1/n$ implies $\alpha \simeq_{F_n} \beta$.

As shown in [4, Lemma 5.16], any continuous map $f: Y \rightarrow X$ between second countable, compact, Hausdorff spaces induces a Cu-morphism $\text{Lsc}(f, \overline{\mathbb{N}}): \text{Lsc}(X, \overline{\mathbb{N}}) \rightarrow \text{Lsc}(Y, \overline{\mathbb{N}})$ given by $l \mapsto l \circ f$. In what follows, we prove that a weak converse of this result —akin to what happens for commutative C^* -algebras— also exists. These results might be well known to experts (for example, if X and Y are one-dimensional, they follow from [19] and standard facts about abelian C^* -algebras). However, since we have not found them in the literature with our generality, we provide a proof here for the convenience of the reader.

Proposition 4.23. *Let X, Y be compact, metric spaces and let $\alpha: \text{Lsc}(X, \overline{\mathbb{N}}) \rightarrow \text{Lsc}(Y, \overline{\mathbb{N}})$ be a Cu-morphism such that $\alpha(1) = 1$.*

Then there exists a continuous map $f_\alpha: Y \rightarrow X$ inducing α , in the sense that $\alpha = \text{Lsc}(f_\alpha, \overline{\mathbb{N}})$.

Proof. We begin our argument by proving the following claim.

Claim. Let $\lambda: \text{Lsc}(X, \overline{\mathbb{N}}) \rightarrow \overline{\mathbb{N}}$ be a Cu-morphism such that $\lambda(1) = 1$. Then, there exists $x \in X$ such that $\lambda = \text{ev}_x$, the evaluation at x .

Proof of the Claim. Let J be the family of open subsets U in X such that $\lambda(\mathbb{1}_U) = 1$. Given finitely many open sets $U_1, \dots, U_n \in J$, it follows from [57, Remark 4.4] that we can order the sum $\mathbb{1}_{U_1} + \dots + \mathbb{1}_{U_n}$ as follows

$$\mathbb{1}_{U_1} + \dots + \mathbb{1}_{U_n} = \mathbb{1}_{\cup_j U_j} + \mathbb{1}_{\cup_{i < j} (U_i \cap U_j)} + \dots + \mathbb{1}_{\cap_j U_j}$$

and, consequently,

$$n = \lambda(\mathbb{1}_{U_1}) + \dots + \lambda(\mathbb{1}_{U_n}) = \lambda(\mathbb{1}_{\cup_j U_j}) + \dots + \lambda(\mathbb{1}_{\cap_j U_j}).$$

This implies that $\lambda(\mathbb{1}_{\cap_j U_j}) = 1$ and, in particular, that $\cap_j U_j \neq \emptyset$. Thus, J has the finite intersection property and, since X is compact, we get that $\cap_{U \in J} \overline{U}$ is not empty.

To see that $\cap_{U \in J} \overline{U} = \{x\}$, assume for the sake of contradiction that there exist $x, y \in \cap_{U \in J} \overline{U}$ with $x \neq y$. Let B, C be closed disjoint balls of nonzero radius centred at x and y respectively. One has $1 \leq \mathbb{1}_{X-B} + \mathbb{1}_{X-C}$ and, consequently, that $\lambda(\mathbb{1}_{X-B})$ or $\lambda(\mathbb{1}_{X-C})$ is 1. Thus, we may assume without loss of generality that $\lambda(\mathbb{1}_{X-C}) = 1$ or, in other words, that $X - C \in J$. This is a contradiction, since $y \notin \overline{X - C}$. It follows that $\cap_{U \in J} \overline{U}$ contains a single point x .

Finally, to see that $\lambda = \text{ev}_x$, take any open subset U and let U' be a compactly contained open subset in U such that $\lambda(\mathbb{1}_U) = \lambda(\mathbb{1}_{U'})$. If $\lambda(\mathbb{1}_U) = 1$, then $x \in \overline{U'}$ and so $x \in U$. Conversely, if $x \in U$, take U' compactly contained in U such that $x \in U'$. Then, $\mathbb{1}_U + \mathbb{1}_{X-\overline{U'}} \geq 1$, which implies that either $\lambda(\mathbb{1}_U) = 1$ or $\lambda(\mathbb{1}_{X-\overline{U'}}) = 1$. The second equality cannot hold, since otherwise we would get $x \in X - \overline{U'}$, a contradiction. This proves the claim.

Now, for any $y \in Y$, the composition $\text{ev}_y \circ \alpha$ is a Cu-morphism such that $\lambda(1) = 1$. Using the claim, there exists $x \in X$ with $\text{ev}_y \circ \alpha = \text{ev}_x$. Let $f_\alpha : Y \rightarrow X$ be the map defined by $f_\alpha(y) := x$. To see that f_α is continuous, take an open subset U of X and let $V \subseteq Y$ be the open subset such that $\alpha(\mathbb{1}_V) = \mathbb{1}_U$. Then, we see that

$$f_\alpha^{-1}(U) = \{y \in Y \mid \text{ev}_{f_\alpha(y)}(\mathbb{1}_U) = 1\} = \{y \in Y \mid \text{ev}_y \circ \alpha(\mathbb{1}_U) = 1\} = V$$

which ends the proof. \square

Corollary 4.24. *Let X, Y be compact, metric spaces. Let $\alpha : \text{Lsc}(X, \overline{\mathbb{N}}) \rightarrow \text{Lsc}(Y, \overline{\mathbb{N}})$ be a Cu-morphism with $\alpha(1) = 1$. Let $f_\alpha : Y \rightarrow X$ be the continuous map obtained above.*

Then, α is an order-embedding if and only if f_α is surjective.

Proof. Assume for the sake of contradiction that α is an order-embedding and that there exists $x \in X \setminus f_\alpha(Y)$. Since $f_\alpha(Y)$ is compact, we can find an open neighbourhood U of x disjoint with $f_\alpha(Y)$. In particular, $f_\alpha^{-1}(U) = \emptyset$ and, therefore, $\alpha(\mathbb{1}_U) = \mathbb{1}_{f_\alpha^{-1}(U)} = 0$, a contradiction.

Conversely, assume now that $f_\alpha : Y \rightarrow X$ is surjective. The order in $\text{Lsc}(X, \overline{\mathbb{N}})$ is determined by the indicators $\mathbb{1}_U$ (see e.g. [15, Proposition 4.3] or [57]). Thus it is enough to prove that $\mathbb{1}_U \leq \mathbb{1}_V$ whenever $\alpha(\mathbb{1}_U) \leq \alpha(\mathbb{1}_V)$. Let $U, V \subseteq X$ be such that $\alpha(\mathbb{1}_U) \leq \alpha(\mathbb{1}_V)$. We have $\mathbb{1}_{f_\alpha^{-1}(U)} \leq \mathbb{1}_{f_\alpha^{-1}(V)}$. Consequently, $f_\alpha^{-1}(U) \subseteq f_\alpha^{-1}(V)$. By the surjectivity of f_α we deduce that $U \subseteq V$, as desired. \square

4.25 (The Cantor Set). Let us recall a characterization of the Cantor set. We use the language and formulations detailed in [9], even though this characterization had been obtained beforehand, e.g. in [12,40]. As mentioned in the discussion of [9, Example 4.55], the Cantor set $2^{\mathbb{N}}$ is the Fraïssé limit of the category of finite discrete sets and continuous

surjections in the category of zero-dimensional compacta and continuous surjections. Now, using the characterization of Fraïssé limits given in [9, Theorem 4.15], one obtains the following result.

Theorem 4.26 (Characterization of the Cantor set). *A zero-dimensional compactum C is the Cantor set if and only if C satisfies the following property:*

For any two finite discrete sets F, F' and any two continuous surjections $f: C \rightarrow F, g: F' \rightarrow F$, there exists a continuous surjection $h: C \rightarrow F'$ such that $h \circ g = f$.

4.27. *The category $\mathcal{K}_{2^{\mathbb{N}}}$ is the category whose objects are simplicial Cu-semigroups and whose morphisms are order-embeddings such that $1 \mapsto 1$.*

Theorem 4.28. *The category $\mathcal{K}_{2^{\mathbb{N}}}$ is Fraïssé.*

Proof. Throughout the proof, we will denote the r -tuple $(0, \dots, 1, \dots, 0)$ with value 1 at the i -th component and 0 everywhere else by δ_i^r . Note that $\{\delta_i^r\}_{i=1}^r$ generates $\overline{\mathbb{N}}^r$.

The category $\mathcal{K}_{2^{\mathbb{N}}}$ contains countably many objects and finitely many morphisms between two given objects. It follows that the category $\mathcal{K}_{2^{\mathbb{N}}}$ is separable.

Let $\overline{\mathbb{N}}^{r_1}, \overline{\mathbb{N}}^{r_2}$ be simplicial Cu-semigroups. We construct $\alpha_1: \overline{\mathbb{N}}^{r_1} \rightarrow \overline{\mathbb{N}}^{r_1} \oplus \overline{\mathbb{N}}^{r_2}$ that sends $\delta_1^{r_1} \mapsto \delta_1^{r_1} \oplus 1_{\overline{\mathbb{N}}^{r_2}}$ and $\delta_i^{r_1} \mapsto \delta_i^{r_1} \oplus 0_{\overline{\mathbb{N}}^{r_2}}$ for any $2 \leq i \leq r_1$. Similarly, we construct $\alpha_2: \overline{\mathbb{N}}^{r_2} \rightarrow \overline{\mathbb{N}}^{r_1} \oplus \overline{\mathbb{N}}^{r_2}$. It is readily checked that α_1, α_2 are $\mathcal{K}_{2^{\mathbb{N}}}$ -morphisms and, hence, the joint embedding property follows.

Let $\alpha_1: \overline{\mathbb{N}}^r \rightarrow \overline{\mathbb{N}}^{t_1}$ and $\alpha_2: \overline{\mathbb{N}}^r \rightarrow \overline{\mathbb{N}}^{t_2}$ be $\mathcal{K}_{2^{\mathbb{N}}}$ -morphisms. We know from Corollary 4.24 that $r \leq t_1, t_2$. Further, we may assume that $t_1 = t_2 = t$ and, upon a possible reindexing (i.e. composing with an isomorphism), we may also assume that the α_i 's are of the form $\text{id} \oplus \eta_i$ for some Cu-morphisms $\eta_i: \overline{\mathbb{N}}^r \rightarrow \overline{\mathbb{N}}^{t-r}$. Note that, since both α_1 and α_2 map 1_r to 1_t , the maps η_i also map 1_r to 1_{t-r} .

Let $\beta_i: \overline{\mathbb{N}}^r \oplus \overline{\mathbb{N}}^{t-r} \rightarrow \overline{\mathbb{N}}^r \oplus \overline{\mathbb{N}}^{t-r} \oplus \overline{\mathbb{N}}^{t-r}$ be the maps $\beta_1(x, y) = (x, y, \eta_2(x))$ and $\beta_2(x, y) = (x, \eta_1(x), y)$. Note that these order-embeddings map 1 to 1. By construction, we have $\beta_1 \circ \alpha_1 = \beta_2 \circ \alpha_2$, which shows that the category $\mathcal{K}_{2^{\mathbb{N}}}$ has the amalgamation property. \square

Corollary 4.29. *The Fraïssé limit of $\mathcal{K}_{2^{\mathbb{N}}}$ is $\text{Lsc}(2^{\mathbb{N}}, \overline{\mathbb{N}})$.*

Proof. First, note that we can identify any $\mathcal{K}_{2^{\mathbb{N}}}$ -object $\overline{\mathbb{N}}^r$ with $\text{Lsc}(X_r, \overline{\mathbb{N}})$, where X_r is any finite discrete set of cardinality r . Now, let $\alpha: \text{Lsc}(X_r, \overline{\mathbb{N}}) \rightarrow \text{Lsc}(X_t, \overline{\mathbb{N}})$ be a $\mathcal{K}_{2^{\mathbb{N}}}$ -morphism and consider the continuous surjective map $f_\alpha: X_t \rightarrow X_r$ obtained from Proposition 4.23. From the construction of f_α , we deduce that α can be identified with $\text{Lsc}(f_\alpha, \overline{\mathbb{N}}): \text{Lsc}(X_r, \overline{\mathbb{N}}) \rightarrow \text{Lsc}(X_t, \overline{\mathbb{N}})$ which sends $l \mapsto l \circ f_\alpha$.

On the other hand, we know that the Fraïssé limit is obtained from an inductive system in $\mathcal{K}_{2^{\mathbb{N}}}$ that we write $(\overline{\mathbb{N}}^{r_i}, \alpha_i)_i$. By the above, we identify the system with $(\text{Lsc}(X_{r_i}, \overline{\mathbb{N}}), \text{Lsc}(f_{\alpha_i}, \overline{\mathbb{N}}))_i$. Now combining Corollary 4.24 with Theorem 3.27 (and the fact that the category $\mathcal{K}_{2^{\mathbb{N}}}$ has exact amalgamation property), we get that $\lim_{\leftarrow} (X_{r_i}, f_{\alpha_i})$

is a zero-dimensional compactum satisfying the above characterization of the Cantor set. That is, $\varprojlim (X_{r_i}, f_{\alpha_i}) \cong 2^{\mathbb{N}}$ and the result follows from [4, Proposition 5.18]. \square

4.30 (The Pseudo-arc). Proceeding as before, we recall a characterization of the pseudo-arc in the language from [9], although this result had also been obtained in the past, e.g. in [35,40]. As shown in [9, Theorem 4.38], the pseudo-arc \mathbb{P} is the Fraïssé limit of the category consisting of a single object being the unit interval and continuous surjections in the category of arc-like continua and continuous surjections. Using [9, Theorem 4.15], one gets:

Theorem 4.31 (Characterization of the Pseudo-arc). *An arc-like continuum P is the pseudo-arc if and only if P satisfies the following property:*

For any two continuous surjections $f: P \rightarrow [0, 1]$ and $g: [0, 1] \rightarrow [0, 1]$ and any $\varepsilon > 0$, there exists a continuous surjection $h: P \rightarrow [0, 1]$ such that $\|h \circ g - f\| < \varepsilon$.

The following two lemmas will be needed in our proofs. The first is known as the Mountain Climbing Lemma (see [33]), while the second is readily obtained by generalizing the arguments in [58, Lemma 4.5].

Lemma 4.32 (Mountain Climbing Lemma). *Let $f_1, f_2: [0, 1] \rightarrow [0, 1]$ be continuous, piecewise linear maps that are not constant on any subinterval and such that $f_1(0) = 0 = f_2(0)$ and $f_1(1) = 1 = f_2(1)$. Then, there exist surjective, continuous maps $g_1, g_2: [0, 1] \rightarrow [0, 1]$ such that $f_1 \circ g_1 = f_2 \circ g_2$.*

Lemma 4.33. *Let X and Y be second countable, compact, Hausdorff spaces. Let $f, g: Y \rightarrow X$ be continuous surjective maps and consider their induced Cu-morphisms $\text{Lsc}(f, \overline{\mathbb{N}}), \text{Lsc}(g, \overline{\mathbb{N}})$ given by $l \mapsto l \circ f, l \circ g$ respectively. Then,*

- (i) *For any finite subset F of $\text{Lsc}(X, \overline{\mathbb{N}})$, there exists $\varepsilon_F > 0$ such that $\text{Lsc}(f, \overline{\mathbb{N}}) \simeq_F \text{Lsc}(g, \overline{\mathbb{N}})$ whenever $\|f - g\| < \varepsilon$.*
- (ii) *For any $\varepsilon > 0$, there exists a finite subset F_ε of $\text{Lsc}(X, \overline{\mathbb{N}})$ such that $\|f - g\| < \varepsilon$ whenever $\text{Lsc}(f, \overline{\mathbb{N}}) \simeq_{F_\varepsilon} \text{Lsc}(g, \overline{\mathbb{N}})$.*

Further, the ε_F only depends on F, X, Y , and F_ε only depends on ε, X, Y (not on f, g).

4.34. *The category $\mathcal{K}_{\mathbb{P}}$ is the category whose (single) object is $\text{Lsc}([0, 1], \overline{\mathbb{N}})$ and whose morphisms are order-embeddings such that $1 \mapsto 1$.*

Theorem 4.35. *The category $\mathcal{K}_{\mathbb{P}}$ is Fraïssé.*

Proof. The category $\mathcal{K}_{\mathbb{P}}$ contains only one object, so the joint embedding property is trivial.

To see that $\mathcal{K}_{\mathbb{P}}$ is separable, we consider the subcategory $\mathfrak{s} \subseteq \mathcal{K}_{\mathbb{P}}$ whose morphisms are $\mathcal{K}_{\mathbb{P}}$ -morphisms of the form $\text{Lsc}(h, \overline{\mathbb{N}}) : \text{Lsc}([0, 1], \overline{\mathbb{N}}) \rightarrow \text{Lsc}([0, 1], \overline{\mathbb{N}})$ where $h : [0, 1] \rightarrow [0, 1]$ is any piecewise linear, surjective map which is not constant on any subinterval and which has rational, finitely many peaks and valleys. We know from Corollary 4.24 that any $\mathcal{K}_{\mathbb{P}}$ -morphism α is of the form $\text{Lsc}(f_{\alpha}, \overline{\mathbb{N}})$ where $f_{\alpha} : [0, 1] \rightarrow [0, 1]$ is a continuous surjective map. Moreover, it is well-known that any such continuous surjective map can be approximated in norm by a piecewise linear map h which is not constant on any subinterval and which has rational, finitely many peaks and valleys. Therefore, it follows from Lemma 4.33 that \mathfrak{s} is a countable, dominating subcategory.

Finally, let us prove $\mathcal{K}_{\mathbb{P}}$ satisfies the near amalgamation property. Let α_1, α_2 be Cu-morphisms in $\mathcal{K}_{\mathbb{P}}$ and consider the continuous, surjective maps $f_{\alpha_1}, f_{\alpha_2} : [0, 1] \rightarrow [0, 1]$ obtained from Corollary 4.24. Also, let $F \subseteq \text{Lsc}([0, 1], \overline{\mathbb{N}})$ be a finite subset and consider the bound $\varepsilon_F > 0$ given by Lemma 4.33.

Without loss of generality, we can assume that $f_{\alpha_1}(0) = 0 = f_{\alpha_2}(0)$ and $f_{\alpha_1}(1) = 1 = f_{\alpha_2}(1)$. (If needed, we can precompose them with well-chosen continuous surjective maps.) As before, we can find piecewise linear maps $h_1, h_2 : [0, 1] \rightarrow [0, 1]$ which are not constant on any subinterval and which only have rational, finitely many peaks and valleys, at distance less than $\varepsilon/2$ from f_{α_1} and f_{α_2} respectively, and such that $h_1(0) = 0 = h_2(0)$ and $h_1(1) = 1 = h_2(1)$.

Applying the Mountain Climbing lemma (see Lemma 4.32), we obtain surjective continuous maps $g_1, g_2 : [0, 1] \rightarrow [0, 1]$ such that $h_1 \circ g_1 = h_2 \circ g_2$. By construction, we get that $f_{\alpha_1} \circ g_1$ and $f_{\alpha_2} \circ g_2$ are at distance at most ε . Let $\beta_1 := \text{Lsc}(g_1, \overline{\mathbb{N}})$ and $\beta_2 := \text{Lsc}(g_2, \overline{\mathbb{N}})$ be the $\mathcal{K}_{\mathbb{P}}$ -morphisms induced by g_1 and g_2 respectively. By Lemma 4.33, we obtain $\beta_1 \circ \alpha_1 \simeq_F \beta_2 \circ \alpha_2$, as desired. \square

Corollary 4.36. *The Fraïssé limit of $\mathcal{K}_{\mathbb{P}}$ is $\text{Lsc}(\mathbb{P}, \overline{\mathbb{N}})$.*

Proof. We know that the Fraïssé limit is obtained from an inductive system in $\mathcal{K}_{\mathbb{P}}$ that we write $(\text{Lsc}([0, 1], \overline{\mathbb{N}}), \alpha_i)_i$. Now combining Corollary 4.24 with Theorem 3.27, we get that $\varprojlim ([0, 1], f_{\alpha_i})$ is an arc-like continuum satisfying the above characterization of the pseudo-arc. We conclude that $\varprojlim ([0, 1], f_{\alpha_i}) \cong \mathbb{P}$ and the result follows from [4, Proposition 5.18]. \square

Remark 4.37. We have chosen to give self-contained proofs of the last two examples, but they could have alternatively been obtained as a combination of Remark 3.32 (ii), Corollary 4.24, [45] and known results in Fraïssé theory of C^* -algebras (for example, those in [56]). Indeed, since in both categories $\mathcal{K}_{2^{\mathbb{N}}}$ and $\mathcal{K}_{\mathbb{P}}$ the dimension of the underlying spaces is at most one, we know from [45] (see also [23] for the case of the interval) that all the objects are Cuntz semigroups of commutative C^* -algebras. Further, it follows from Corollary 4.24 that the Cu-morphisms in both categories are in correspondence with surjective continuous maps between the spaces, which in turn correspond to injective $*$ -homomorphisms between the commutative C^* -algebras.

In other words, this shows that both $\mathcal{K}_{2^{\mathbb{N}}}$ and $\mathcal{K}_{\mathbb{P}}$ can be written as $\text{Cu}(\mathfrak{c}_{2^{\mathbb{N}}})$ and $\text{Cu}(\mathfrak{c}_{\mathbb{P}})$ for well chosen subcategories of commutative C^* -algebras, which are well-known to be Fraïssé (e.g. [56, Theorem 3.4] for the case of the pseudo-arc). Using Remark 3.32 (ii), it follows that both $\mathcal{K}_{2^{\mathbb{N}}}$ and $\mathcal{K}_{\mathbb{P}}$ are Fraïssé, and that their limit coincides with the Cuntz semigroup of the limit of $\mathfrak{c}_{2^{\mathbb{N}}}$ and $\mathfrak{c}_{\mathbb{P}}$.

E. The Jiang-Su algebra

It is readily checked that all the results developed in this paper also hold for classes where the objects are pairs of the form (S, ρ) with $\rho: S \rightarrow [0, \infty]$ a generalized Cu-morphism (also known as a *functional*) and where the morphisms from a pair (S, ρ) to (T, δ) are simply Cu-morphisms $S \rightarrow T$ preserving the prescribed functional ρ . Namely, the statements still hold because a limit of functional-preserving Cu-morphisms is still functional-preserving.

Using Theorem 3.31 (and [43]) in conjunction with the fact that quasitraces on a C^* -algebra correspond to functionals on its Cuntz semigroup ([26]), one obtains the following result by using [42] (see also [25]).

4.38. *The category \mathcal{K}_Z is the category whose objects are pairs $(Z_{p,q}, \rho)$ with p, q coprime and ρ a faithful functional, and whose morphisms are functional-preserving order-embeddings. Here, $Z_{p,q}$ denotes the Cuntz semigroup of the prime dimension-drop algebra $\mathcal{Z}_{p,q}$. Using computations in [4], recall that we have*

$$\text{Cu}(\mathcal{Z}_{p,q}) \cong Z_{p,q} = \{f \in \text{Lsc}([0, 1], \overline{\mathbb{N}}) \mid \exists k_1, k_2 \text{ such that } f(0) = qk_1 \text{ and } f(1) = pk_2\}.$$

Theorem 4.39. *The class \mathcal{K}_Z is Fraïssé and its limit is the Cuntz semigroup Z of the Jiang-Su algebra \mathcal{Z} .*

5. Metrics on Hom_{Cu} -sets

In this last section we introduce a metric on any Hom_{Cu} -set building on the ideas of [15,20,46,58]. We also provide a number of examples and, in particular, we show that this notion generalizes the metrics introduced in [15, Definition 5.1]. We also study the relation between our proposed metric and finite-set comparison. Further, we prove that, in general, comparing Cu-morphisms via the metric is more restrictive than using finite-set comparison. As a consequence, when (re)formulating the notion of Cauchy sequences in terms of the metric, the limit we obtain might not behave as expected. Let us start by recalling an important fact about the category Cu.

5.1 (*A generator for Cu*). Let \mathbb{G} be the submonoid of $\text{Lsc}([0, 1], \overline{\mathbb{N}})$ defined as

$$\mathbb{G} = \{f \in \text{Lsc}([0, 1], \overline{\mathbb{N}}) \mid f(0) = 0, f \text{ increasing}\}.$$

It follows from [49, Section 5.2] that \mathbb{G} is a Cu-semigroup. Moreover, \mathbb{G} is a generator for the category Cu, in the sense that the functor $\text{Cu}(\mathbb{G}, -) : \text{Cu} \rightarrow \text{Set}$ is faithful.

Using terminology of [15], we can view \mathbb{G} as the sub-Cu-semigroup of $\text{Lsc}([0, 1], \overline{\mathbb{N}})$ chain-generated by $\Lambda_{\mathbb{G}} := \{\mathbb{1}_{(t,1]} \mid t \in (0, 1]\}$. Therefore, the order and the compact-containment relation in \mathbb{G} are completely determined by the ones in $\Lambda_{\mathbb{G}}$. (See [15, Section 4] for more details.)

5.2 (Thomsen semigroup of a Cu-semigroup). Let S be a Cu-semigroup. We define the *Thomsen semigroup of S* , in symbols $\text{Th}(S)$, to be the set of Cu-morphisms from the generator \mathbb{G} to S . In other words,

$$\text{Th}(S) := \text{Hom}_{\text{Cu}}(\mathbb{G}, S).$$

This construction is inspired by the C^* -case, where the *Thomsen semigroup of a C^* -algebra A* , in symbols $\mathcal{T}h(A)$, is the set of approximate unitary equivalence classes of $*$ -homomorphisms of the form $C_0((0, 1]) \rightarrow A \otimes \mathcal{K}$. (See [54].) Here, note that $C_0((0, 1])$ is a generator for the category C^* . (See e.g. [49].)

Therefore, the construction above is the natural Cu-analogue of the Thomsen semigroup for C^* -algebras. In fact, there exists a natural (monoid) map $\mathcal{T}h(A) \rightarrow \text{Th}(\text{Cu}(A))$, defined in the following proposition.

Proposition 5.3. *Let A be a C^* -algebra. The map $\iota : \mathcal{T}h(A) \rightarrow \text{Th}(\text{Cu}(A))$ given by $[\varphi] \mapsto \text{Cu}(\varphi)_{|\mathbb{G}}$ is a well-defined monoid morphism.*

If A has stable rank one, then ι is a bijection.

Proof. It is a well-known fact that any two approximate unitary equivalent $*$ -homomorphisms agree at level of Cu. Therefore, ι is a well-defined map for any C^* -algebra.

Now assume that A has stable rank one. Then, it follows from [48, Theorem 4.3, Lemma 7.2] that $\text{Cu}(A)$ is weakly cancellative and satisfies (O5).

Let $\tau \in \text{Th}(\text{Cu}(A))$. Proceeding as in the proof of [58, Proposition 3.4], we can construct a Cu-morphism $\tilde{\tau} : \text{Cu}(C_0((0, 1])) \rightarrow \text{Cu}(A)$ extending τ . Further, such a morphism is unique.

The functor Cu classifies $*$ -homomorphisms from $C_0((0, 1])$ to any C^* -algebra of stable rank one (See [19, 46, 43]). In particular, one can lift the extension $\tilde{\tau}$ to a $*$ -homomorphism $C_0((0, 1]) \rightarrow A$. This proves that ι is surjective.

Since the extension $\tilde{\tau}$ is unique, and the lift of any such τ is unique up to approximate unitary equivalence, ι is injective. \square

We will use the Thomsen semigroup of S to build a metric on any $\text{Hom}_{\text{Cu-set}} \text{Hom}_{\text{Cu}}(S, T)$. Let us start by equipping $\text{Th}(S)$ with the following metric, modelled after the distance in [19]. (See also [46] and Paragraph 4.21.)

Definition 5.4. Let S be a Cu-semigroup, and let $\alpha, \beta \in \text{Th}(S)$. We define

$$d_{\mathbb{G}}(\alpha, \beta) := \inf \{r > 0 \mid \forall t \in [0, 1], \alpha(\mathbb{1}_{(t+r,1]}) \leq \beta(\mathbb{1}_{(t,1]}) \text{ and } \beta(\mathbb{1}_{(t+r,1]}) \leq \alpha(\mathbb{1}_{(t,1]})\}.$$

Remark 5.5. Note that, by definition, $d_{\mathbb{G}}(\alpha, \beta) = 0$ precisely when $\alpha(\mathbb{1}_{(t,1]}) = \beta(\mathbb{1}_{(t,1]})$ for every t . Thus, since \mathbb{G} is generated (as a Cu-semigroup) by the elements $\mathbb{1}_{(t,1]}$, one gets $\alpha = \beta$. Consequently, $d_{\mathbb{G}}$ is a metric. This is in contrast to [19], where weak cancellation of the Cuntz semigroup is needed to ensure positivity.

5.6 (Sets with generating image). Let Λ be a subset of $\text{Th}(S)$. We say that Λ has a *generating image in S* if the submonoid generated by $\{\tau(\mathbb{1}_{(t,1]}) \mid t \in [0, 1], \tau \in \Lambda\}$ is sup-dense in S .

Equivalently, Λ has a generating image if for any $s', s \in S$ with $s' \ll s$, there exist τ_1, \dots, τ_n in Λ and $t_1, \dots, t_n \in [0, 1]$ such that $s' \ll \tau_1(\mathbb{1}_{(t_1,1]}) + \dots + \tau_n(\mathbb{1}_{(t_n,1]}) \ll s$.

For any Cu-semigroup S , there always exists a family (and, in fact, many) with a generating image. For instance, the following result shows that $\text{Th}(S)$ always has a generating image in S .

Proposition 5.7 ([49, Lemma 5.16]). *Let S be a Cu-semigroup and let $(x_n)_n$ be a \ll -increasing sequence in S . Then, there exists $\tau \in \text{Th}(S)$ such that $\tau(\mathbb{1}_{(\frac{1}{n},1]}) = x_n$.*

Proof. We give a brief argument for the convenience of the reader.

Using [6, Proposition 2.10], there exists a net $(y_t)_{t \in [0,1]}$, with $y_{\frac{1}{n}} = x_n$, such that $y_t \ll y_r$ whenever $r < t$ and $\sup_{t > r} y_t = y_r$. (This is achieved by an iterated application of (O2) and the use of (O1).) We let $\tau: \mathbb{G} \rightarrow S$ be the Cu-morphism defined by $\tau(\mathbb{1}_{(t,1]}) := y_t$ for each $t \in [0, 1]$. \square

Definition 5.8. Let $S, T \in \text{Cu}$ and let $\Lambda \subseteq \text{Th}(S)$ be a subset with a generating image in S . For any $\alpha, \beta \in \text{Hom}_{\text{Cu}}(S, T)$, we define

$$d_{\Lambda}(\alpha, \beta) := \sup_{\tau \in \Lambda} d_{\mathbb{G}}(\alpha \circ \tau, \beta \circ \tau).$$

Lemma 5.9. *Let $S, T \in \text{Cu}$ and let $\Lambda \subseteq \text{Th}(S)$ with a generating image in S . Then $d_{\Lambda}(\alpha, \beta)$ is a metric on $\text{Hom}_{\text{Cu}}(S, T)$.*

Proof. The symmetry, triangular inequality, and the fact that $d_{\Lambda}(\alpha, \beta) = 0$ whenever $\alpha = \beta$ are all immediate. We are left to show that $\alpha = \beta$ whenever $d_{\Lambda}(\alpha, \beta) = 0$.

Thus, assume that $d_{\Lambda}(\alpha, \beta) = 0$. Let $s', s \in S$ be such that $s' \ll s$. Since Λ has a generating image in S , there exist $\tau_1, \dots, \tau_n \in \text{Th}(S)$ and $t_1, \dots, t_n \in [0, 1]$ such that $s' \leq \tau_1(\mathbb{1}_{(t_1,1]}) + \dots + \tau_n(\mathbb{1}_{(t_n,1]}) \leq s$. Note that $d_{\mathbb{G}}(\alpha \circ \tau_i, \beta \circ \tau_i) = 0$ for any $i \leq n$. Therefore, we deduce that $\alpha \circ \tau_i = \beta \circ \tau_i$ for any $i \leq n$. Consequently, we have

$$\beta(s') \leq \beta(\tau_1(\mathbb{1}_{(t_1,1]}) + \dots + \beta(\tau_1(\mathbb{1}_{(t_n,1]}) = \alpha(\tau_1(\mathbb{1}_{(t_1,1]}) + \dots + \alpha(\tau_1(\mathbb{1}_{(t_n,1]}) \leq \alpha(s).$$

A symmetric argument gives us that $\alpha(s') \leq \beta(s)$ and Remark 3.3 give sus $\alpha = \beta$. \square

Example 5.10. Let S, T be Cu-semigroups. The metric $d_{\text{Th}(S)}$ is trivial, that is, $d_{\text{Th}(S)}(\alpha, \beta) = 1$ if and only if $\alpha \neq \beta$. We give a brief argument below.

Let $\alpha, \beta \in \text{Hom}_{\text{Cu}}(S, T)$ be such that $d_{\text{Th}(S)}(\alpha, \beta) < 1$ and let $\varepsilon > 0$ be such that $\varepsilon > d_{\text{Th}(S)}(\alpha, \beta)$. Now take any pair $x', x \in S$ such that $x' \ll x$. By Proposition 5.7, we can find $\tau \in \text{Th}(S)$ satisfying $\tau(\mathbb{1}_{(0,1]}) = x$ and $\tau(\mathbb{1}_{(\varepsilon,1]}) = x'$. Since $d_{\mathbb{G}}(\alpha \circ \tau, \beta \circ \tau) < \varepsilon$, we compute $\alpha(x') = \alpha(\tau(\mathbb{1}_{(\varepsilon,1]})) \leq \beta(\tau(\mathbb{1}_{(0,1]})) = \beta(x)$ and $\beta(x') \leq \alpha(x)$. We conclude that $\alpha = \beta$ by Remark 3.3.

The example above illustrates that, despite always having many sets with a generating image, they will only induce meaningful metrics as long as they are not too large. The following examples show that all the (meaningful) Cu-metrics considered in the past for specific Hom_{Cu} -sets can be recovered as d_{Λ} from well-chosen Λ 's with generating image.

Example 5.11. (i) The Cuntz semigroup of the Jacelon-Razak algebra \mathcal{W} can be identified with $[0, \infty]$; see [44]. Let $\tau: \mathbb{G} \rightarrow [0, \infty]$ be the Cu-morphism determined by $\mathbb{1}_{(t,1]} \mapsto 1 - t$.

The family $\Lambda = \{\tau\}$ has a generating image in $\text{Cu}(\mathcal{W})$. It can be computed that, for any $\alpha, \beta \in \text{Hom}_{\text{Cu}}(\text{Cu}(\mathcal{W}), T)$, we have

$$d_{\Lambda}(\alpha, \beta) = d_{\mathbb{G}}(\alpha \circ \tau, \beta \circ \tau) = \inf\{r > 0 \mid \forall t \in [0, 1], \alpha(t - r) \leq \beta(t) \text{ and } \beta(t - r) \leq \alpha(t)\}.$$

(ii) The Cuntz semigroup of the Jiang-Su algebra \mathcal{Z} can be identified with $(0, \infty] \sqcup \mathbb{N}$; see e.g. [30, Theorem 7.3]. Similarly, let $\tau: \mathbb{G} \rightarrow [0, \infty]$ be the Cu-morphism determined by $\mathbb{1}_{(t,1]} \mapsto 1 - t$ and let $c: \mathbb{1}_{(t,1]} \mapsto 1_c$ be the constant map.

The family $\Lambda = \{\tau, c\}$ has a generating image in $\text{Cu}(\mathcal{Z})$. One can show that, for any $\alpha, \beta \in \text{Hom}_{\text{Cu}}(\text{Cu}(\mathcal{Z}), T)$, we have

$$d_{\Lambda}(\alpha, \beta) = \begin{cases} 1, & \text{whenever } \alpha(1_c) \neq \beta(1_c). \\ \inf\{r > 0 \mid \forall t \in [0, 1], \alpha(t - r) \leq \beta(t) \text{ and } \beta(t - r) \leq \alpha(t)\}, & \text{otherwise.} \end{cases}$$

Example 5.12. Let A be a C^* -algebra and let $x \in \text{Cu}(A)$. Fix a contraction $a_x \in (A \otimes \mathcal{K})_+$ such that $x = [a_x]$. This element gives rise to the $*$ -homomorphism $\varphi_x: C_0((0, 1]) \rightarrow (A \otimes \mathcal{K})_+$ given by $\text{id}_{C_0(0,1]} \mapsto a_x$. Denote the canonical inclusion from \mathbb{G} to $\text{Cu}(C_0(0, 1])$ by ι .

The family $\Lambda = \{\text{Cu}(\varphi_x) \circ \iota \mid x \in \text{Cu}(A)\}$ has a generating image in $\text{Cu}(A)$. (In fact, $\Lambda(\mathbb{G}) = \text{Cu}(A)$). For any $\phi_1, \phi_2 \in \text{Hom}_{C^*}(A, B)$, we get

$$d_{\Lambda}(\text{Cu}(\phi_1), \text{Cu}(\phi_2)) \leq d_{C^*}(\phi_1, \phi_2).$$

Example 5.13. Let X be a compact, metric space. Note that, for any $x \in X$, we have

$$\bigcap_{t \in (0,1]} B(x, t \operatorname{diam}(X)) = \{x\} \quad \text{and} \quad \bigcup_{t \in (0,1]} B(x, t \operatorname{diam}(X)) = X.$$

Let $S := \operatorname{Lsc}(X, \overline{\mathbb{N}})$ and let $x \in X$. We define $x_0 := 0_S$ and $x_t := \mathbb{1}_{B(x, t \operatorname{diam}(X))}$ for any $t \in (0, 1]$. By Proposition 5.7, there is a Cu-morphism $\tau_x: \mathbb{G} \rightarrow S$ such that $\tau_x(\mathbb{1}_{(t,1]}) = x_{1-t}$ for any $t \in [0, 1]$.

The family $\Lambda = \{\tau_x \mid x \in X\}$ has a generating image in S . It can be computed that for any $\alpha, \beta \in \operatorname{Hom}_{\operatorname{Cu}}(S, T)$ their distance d_Λ is

$$d_\Lambda(\alpha, \beta) = \inf \{r > 0 \mid \forall V \in \mathcal{O}(X), \alpha(1_V) \leq \beta(1_{V_r}) \text{ and } \beta(1_V) \leq \alpha(1_{V_r})\}$$

where $\mathcal{O}(X) := \{ \text{Open sets of } X \}$ and $V_r := \bigcup_{x \in V} B_r(x)$. This generalizes the distance recalled in Paragraph 4.21 when X is a compact, one-dimensional CW-complex.

Example 5.14. We refer the reader to [15, Section 5.1] for details on finite uniform bases and induced Cu-semimetrics.

Let S be a uniformly-based Cu-semigroup with a finite uniform basis $\Lambda_f = (\Lambda_n, \varepsilon_n)_n$. Let $n \in \mathbb{N}$. Recall that Λ_n is finite and, in particular, that it has finitely many *chains*, i.e. finite \ll -increasing sequences. Let us denote the set of chains in Λ_n starting at 0_S by C_n .

Now let $c \in C_n$ and let l_c be the cardinal of c . From Proposition 5.7, we know that there exists a Cu-morphism $\tau_c: \mathbb{G} \rightarrow S$ such that $\tau_c(\mathbb{1}_{((l_c-k)/l_c, 1]}) = c(k)$ for any $0 \leq k \leq l_c$.

The family $\Lambda = \bigcup_{n \in \mathbb{N}} \{\tau_c \mid c \in C_n\}$ has a generating image in S . Following the ideas of [15, Proposition 5.5], one can show that the metric d_Λ is topologically equivalent to the Cu-semimetric $dd_{\operatorname{Cu}, \Lambda_f}$ induced by the finite uniform basis Λ_f .

We next expose relations between the above metrics and finite-set comparison for Cu-morphisms.

Lemma 5.15. *Let $S, T \in \operatorname{Cu}$ and let $\Lambda \subseteq \operatorname{Th}(S)$ with a generating image in S . Then*

- (i) *For any finite set $F \subseteq S$, there exists $\varepsilon_F > 0$ such that $\alpha \simeq_F \beta$ whenever $d_\Lambda(\alpha, \beta) < \varepsilon_F$.*

If moreover Λ is finite, then

- (ii) *For any $\varepsilon > 0$, there exists a finite set $F_\varepsilon \subseteq S$ such that $d_\Lambda(\alpha, \beta) < \varepsilon$ whenever $\alpha \simeq_{F_\varepsilon} \beta$.*

Proof. (i) Note that it is enough to prove the result for $F = \{x', x\}$ where $x', x \in S$ are such that $x' \ll x$. Thus, let $x', x \in S$ be such that $x' \ll x$. We know that there exist $n \in \mathbb{N}$, $\tau_1, \dots, \tau_n \in \Lambda$, and $t_1, \dots, t_n \in [0, 1]$ such that $x' \leq \tau_1(\mathbb{1}_{(t_1,1]}) + \dots + \tau_n(\mathbb{1}_{(t_n,1]}) \ll x$.

For each i , let $r_i > 0$ be such that $x' \ll \tau_1(\mathbb{1}_{(t_1+r_1,1]}) + \dots + \tau_n(\mathbb{1}_{(t_n+r_n,1]})$. Set $\varepsilon_F := \min_i r_i$, and let α, β be such that $d_\Lambda(\alpha, \beta) < \varepsilon_F$. We get

$$\alpha(x') \leq \alpha(\tau_1(\mathbb{1}_{(t_1+r_1,1]}) + \dots + \tau_n(\mathbb{1}_{(t_n+r_n,1]})) \leq \beta(\tau_1(\mathbb{1}_{(t_1,1]}) + \dots + \tau_n(\mathbb{1}_{(t_n,1]})) \leq \beta(x)$$

A symmetric argument gives $\beta(x') \leq \alpha(x)$, as required.

(ii) We now assume that Λ is finite. Let $\varepsilon > 0$ and let t_1, \dots, t_n be a partition of $[0, 1]$ such that $|t_i - t_{i+1}| < \varepsilon/2$. Let us define

$$F_\varepsilon := \{\tau(\mathbb{1}_{(t_i,1]}) \mid \tau \in \Lambda, i \leq n\} \cup \{\tau(\mathbb{1}_{(t_i+\varepsilon/2,1]}) \mid \tau \in \Lambda, i \leq n\}.$$

Let α, β be such that $\alpha \simeq_{F_\varepsilon} \beta$. By [58, Lemma 4.8], we have $d_{\mathbb{G}}(\alpha \circ \tau, \beta \circ \tau) < 1/n + \varepsilon/2 \leq \varepsilon$ for every $\tau \in \Lambda$. This implies $d_\Lambda(\alpha, \beta) < \varepsilon$, as desired. \square

Let S, T be Cu-semigroups and let $\Lambda \subseteq \text{Th}(S)$ be a set with a generating image in S . We will say that a sequence $(\alpha_i)_i$ in $\text{Hom}_{\text{Cu}}(S, T)$ is d_Λ -Cauchy if $\sum_i d_\Lambda(\alpha_i, \alpha_{i+1}) < \infty$. The proposition below shows that any d_Λ -Cauchy sequence has a unique limit α , in the sense of Definition 3.5. Nevertheless, an extra-assumption is needed (e.g. Λ is finite) in order for α to satisfy $d_\Lambda(\alpha_i, \alpha) \rightarrow 0$.

Proposition 5.16. *Let S, T be Cu-semigroups and let $\Lambda \subseteq \text{Th}(S)$ be a set with a generating image in S . Then any d_Λ -Cauchy sequence $(\alpha_i)_i$ in $\text{Hom}_{\text{Cu}}(S, T)$ has a (unique) limit.*

Proof. Let $F \subseteq S$ be a finite set and let ε_F be the bound given by Lemma 5.15 (i). Since $(\alpha_i)_i$ is d_Λ -Cauchy, there exists some $i_F \in \mathbb{N}$ such that $d_\Lambda(\alpha_j, \alpha_k) < \varepsilon_F$ whenever $j, k \geq i_F$. It follows from Lemma 5.15 (i) that $(\alpha_i)_i$ is Cauchy in the sense of Definition 3.4. Using Theorem 3.8, we get that $(\alpha_i)_i$ has a (unique) limit. \square

Proposition 5.17. *Let $S, T \in \text{Cu}$ and let $\Lambda \subseteq \text{Th}(S)$ with a generating image in S . Let $(\alpha_i)_i$ be a d_Λ -Cauchy sequence in $\text{Hom}_{\text{Cu}}(S, T)$ and let $\alpha \in \text{Hom}_{\text{Cu}}(S, T)$. Then the following are equivalent:*

- (i) α is the limit of a sequence $(\alpha_i)_i$.
- (ii) $d_{\mathbb{G}}(\alpha_i \circ \tau, \alpha \circ \tau) \rightarrow 0$ for any $\tau \in \Lambda$.

If moreover Λ is finite, then (i)-(ii) are in turn equivalent to

- (iii) $d_\Lambda(\alpha, \alpha_i) \rightarrow 0$.

Remark 5.18. (iii) always implies (ii). We exhibit an example where the converse does not hold.

For any $n \in \mathbb{N}$ we let $\tau_n \in \text{Th}(\mathbb{G})$ be the map given by $\mathbb{1}_{(t,1]} \mapsto \mathbb{1}_{(t+1/n,1]}$. It can be checked that $\text{id}_{\mathbb{G}}$ is the limit of the sequence $(\tau_n)_n$. Consider the piecewise linear functions $f_n: [0, 1] \rightarrow [0, 1]$ mapping $0 \mapsto 0$, $1/2 \mapsto 1/n$, and $1 \mapsto 1$. Let $\lambda_n \in \text{Th}(S)$ be the Cu-morphisms defined by $\lambda_n(\mathbb{1}_{(t,1]}) := \mathbb{1}_{(f_n(t),1]}$.

The family $\Lambda = \{\lambda_n\}_n$ has a generating image in S , and we get $d_{\mathbb{G}}(\tau_n \circ \lambda_n, \lambda_n) = 1/2$ for each n . This shows that $d_{\Lambda}(\tau_n, \text{id})$ is constantly $1/2$. In particular, the distance does not tend to 0.

Question 5.19. Let S be a Cu-semigroup. When does there exist $\Lambda \subseteq \text{Th}(S)$ such that $\sum_i d_{\Lambda}(\alpha_i, \alpha_{i+1}) < \infty$ implies $d_{\Lambda}(\alpha, \alpha_i) \rightarrow 0$?

Data availability

No data was used for the research described in the article.

References

- [1] R. Antoine, P. Ara, J. Bosa, F. Perera, E. Vilalta, The Cuntz semigroup of a ring, Preprint, arXiv:2307.07266, 2023.
- [2] R. Antoine, F. Perera, L. Robert, H. Thiel, Edwards' condition for quasitraces on C^* -algebras, Proc. R. Soc. Edinb., Sect. A 151 (2) (2021) 525–547.
- [3] R. Antoine, F. Perera, L. Robert, H. Thiel, C^* -algebras of stable rank one and their Cuntz semigroups, Duke Math. J. 171 (1) (2022) 33–99.
- [4] R. Antoine, F. Perera, L. Santiago, Pullbacks, $C(X)$ -algebras, and their Cuntz semigroup, J. Funct. Anal. 260 (10) (2011) 2844–2880.
- [5] R. Antoine, F. Perera, H. Thiel, Tensor products and regularity properties of Cuntz semigroups, Mem. Am. Math. Soc. 251 (1199) (2018) viii+191.
- [6] R. Antoine, F. Perera, H. Thiel, Abstract bivariant Cuntz semigroups, Int. Math. Res. Not. 17 (2020) 5342–5386.
- [7] Ramon Antoine, Francesc Perera, Hannes Thiel, Abstract bivariant Cuntz semigroups II, Forum Math. 32 (1) (2020) 45–62.
- [8] M. Ali Asadi-Vasfi, Hannes Thiel, Eduard Vilalta, Ranks of soft operators in nowhere scattered C^* -algebras, J. Inst. Math. Jussieu (2024), in press.
- [9] A. Bartos, W. Kubiś, Hereditarily indecomposable continua as generic mathematical structures, Preprint, arXiv:2208.06886, 2022.
- [10] D. Bartošová, A. Kwiatkowska, Lelek fan from a projective Fraïssé limit, Fundam. Math. 231 (1) (2015) 57–79.
- [11] J. Bosa, H. Petzka, Comparison properties of the Cuntz semigroup and applications to C^* -algebras, Can. J. Math. 70 (1) (2018) 26–52.
- [12] F. Cabello, On continuous surjections from Cantor set, Extr. Math. 19 (3) (2004) 335–337.
- [13] L. Cantier, A unitary Cuntz semigroup for C^* -algebras of stable rank one, J. Funct. Anal. 281 (9) (2021) 109175.
- [14] L. Cantier, Unitary Cuntz semigroups of ideals and quotients, Münster J. Math. 14 (2021) 585–606.
- [15] L. Cantier, Uniformly based Cuntz semigroups and approximate intertwinings, Int. J. Math. 33 (9) (2022) 2250062, 36.
- [16] L. Cantier, A systematic approach for invariants of C^* -algebras, arXiv:2304.08836 [math.OA], 2023.
- [17] L. Cantier, Towards a classification of unitary elements of C^* -algebras, preprint arXiv:2304.01761, 2023.
- [18] L. Cantier, The unitary Cuntz semigroup on the classification of non-simple C^* -algebras, J. Math. Anal. Appl. 522 (2) (2023) 127003.
- [19] A. Ciuperca, G.A. Elliott, A remark on invariants for C^* -algebras of stable rank one, Int. Math. Res. Not. 5 (158) (2008) 33.

- [20] A. Ciuperca, G.A. Elliott, L. Santiago, On inductive limits of type-I C^* -algebras with one-dimensional spectrum, *Int. Math. Res. Not.* 11 (2011) 2577–2615.
- [21] A. Ciuperca, L. Robert, L. Santiago, The Cuntz semigroup of ideals and quotients and a generalized Kasparov stabilization theorem, *J. Oper. Theory* 64 (1) (2010) 155–169.
- [22] K.T. Coward, G.A. Elliott, C. Ivanescu, The Cuntz semigroup as an invariant for C^* -algebras, *J. Reine Angew. Math.* 623 (2008) 161–193.
- [23] K.T. Coward, The Cuntz Semigroup as a Classification Functor for $C(\text{star})$ -Algebras, ProQuest LLC, Thesis (Ph.D.)– University of Toronto, Ann Arbor, MI, 2007, Canada.
- [24] J. Cuntz, Dimension functions on simple C^* -algebras, *Math. Ann.* 233 (2) (1978) 145–153.
- [25] C. Eagle, I. Farah, B. Hart, B. Kadets, V. Kalashnyk, M. Lupini, Fraïssé limits of C^* -algebras, *J. Symb. Log.* 81 (2016) 755.
- [26] G.A. Elliott, L. Robert, L. Santiago, The cone of lower semicontinuous traces on a C^* -algebra, *Am. J. Math.* 133 (4) (2011) 969–1005.
- [27] I. Farah, B. Hart, M. Lupini, L. Robert, A. Tikuisis, A. Vignati, W. Winter, Model theory of C^* -algebras, *Mem. Am. Math. Soc.* 271 (1324) (2021) viii+127.
- [28] I. Farah, B. Hart, D. Sherman, Model theory of operator algebras II: model theory, *Isr. J. Math.* 201 (1) (2014) 477–505.
- [29] R. Fraïssé, Sur l’extension aux relations de quelques propriétés des ordres, *Ann. Sci. Éc. Norm. Supér.* 71 (4) (1954) 363–388, 3e série.
- [30] E. Gardella, F. Perera, The modern theory of Cuntz semigroups of C^* -algebras., preprint arXiv: 2212.02290, 2022.
- [31] S. Ghasemi, Strongly self-absorbing C^* -algebras and Fraïssé limits, *Bull. Lond. Math. Soc.* 53 (3) (2021) 937–955.
- [32] S. Ghasemi, W. Kubiś, Universal AF-algebras, *J. Funct. Anal.* 279 (5) (2020) 108590, 32.
- [33] T. Homma, A theorem on continuous functions, *Kodai Math. Semin. Rep.* 4 (1952) 13–16.
- [34] J. Im, G.A. Elliott, Coloured isomorphism of classifiable C^* -algebras, *Canad. J. Math.* 76 (1) (2024) 216–245.
- [35] T. Irwin, S. Solecki, Projective Fraïssé limits and the pseudo-arc, *Trans. Am. Math. Soc.* 358 (7) (2006) 3077–3096.
- [36] B. Jacelon, A. Vignati, Stably projectionless Fraïssé limits, *Stud. Math.* 267 (2022) 161–199.
- [37] X. Jiang, H. Su, On a simple unital projectionless C^* -algebra, *Am. J. Math.* 121 (2) (1999) 359–413.
- [38] P.T. Johnstone, *Stone Spaces*, Cambridge Studies in Advanced Mathematics, vol. 3, Cambridge University Press, Cambridge, 1982.
- [39] E. Kirchberg, M. Rørdam, Non-simple purely infinite C^* -algebras, *Am. J. Math.* 122 (3) (2000) 637–666.
- [40] W. Kubiś, Metric-enriched categories and approximate Fraïssé limits, Preprint, arXiv:1210.6506, 2013.
- [41] H. Lin, Unitary groups and augmented Cuntz semigroups of separable simple \mathcal{Z} -stable C^* -algebras, *Int. J. Math.* 33 (02) (2022) 2250018.
- [42] S. Masumoto, The Jiang-Su algebra as a Fraïssé limit, *J. Symb. Log.* 82 (4) (2017) 1541–1559.
- [43] L. Robert, Classification of inductive limits of 1-dimensional NCCW complexes, *Adv. Math.* 231 (5) (2012) 2802–2836.
- [44] L. Robert, The cone of functionals on the Cuntz semigroup, *Math. Scand.* 113 (2) (2013) 161–186.
- [45] L. Robert, The Cuntz semigroup of some spaces of dimension at most two, *C. R. Math. Acad. Sci. Soc. R. Can.* 35 (1) (2013) 22–32.
- [46] L. Robert, L. Santiago, Classification of C^* -homomorphisms from $C_0(0, 1)$ to a C^* -algebra, *J. Funct. Anal.* 258 (3) (2010) 869–892.
- [47] M. Rørdam, A simple C^* -algebra with a finite and an infinite projection, *Acta Math.* 191 (1) (2003) 109–142.
- [48] M. Rørdam, W. Winter, The Jiang-Su algebra revisited, *J. Reine Angew. Math.* 642 (2010) 129–155.
- [49] C. Schons, *Categorical aspects of Cuntz semigroups*, Master thesis <http://hannesthiel.org/supervision>, 2018.
- [50] H. Thiel, E. Vilalta, Covering dimension of Cuntz semigroups II, *Int. J. Math.* 32 (14) (2021) 2150100, 27.
- [51] H. Thiel, E. Vilalta, Covering dimension of Cuntz semigroups, *Adv. Math.* 394 (2022) 108016, 44.
- [52] H. Thiel, E. Vilalta, Nowhere scattered C^* -algebras, *J. Noncommut. Geom.* 18 (1) (2024) 231–263.
- [53] H. Thiel, E. Vilalta, The global glimm property, *Trans. Am. Math. Soc.* 376 (2023) 4713–4744.
- [54] K. Thomsen, Inductive limits of interval algebras: unitary orbits of positive elements, *Math. Ann.* 293 (1) (1992) 47–63.

- [55] A.S. Toms, On the classification problem for nuclear C^* -algebras, *Ann. Math. (2)* 167 (3) (2008) 1029–1044.
- [56] A. Vignati, Fraïssé theory in operator algebras, in: Isaac Goldbring (Ed.), *Model Theory of Operator Algebras*, De Gruyter, Berlin, Boston, 2023.
- [57] E. Vilalta, The Cuntz semigroup of unital commutative AI-algebras, *Can. J. Math.* 75 (6) (2023) 1831–1868.
- [58] E. Vilalta, A local characterization for the Cuntz semigroup of AI-algebras, *J. Math. Anal. Appl.* 506 (1) (2022) 125612, 47.
- [59] E. Vilalta, Nowhere scattered multiplier algebras, *Proc. R. Soc. Edinb. A* (2024), in press, <https://doi.org/10.1017/prm.2023.123>.
- [60] I.B. Yaacov, Fraïssé limits of metric structures, *J. Symb. Log.* 80 (1) (2015) 100–115.