

# The volume of pseudoeffective line bundles and partial equilibrium

Downloaded from: https://research.chalmers.se, 2024-09-26 22:02 UTC

Citation for the original published paper (version of record):

Darvas, T., Xia, M. (2024). The volume of pseudoeffective line bundles and partial equilibrium. Geometry and Topology, 28(4): 1957-1993. http://dx.doi.org/10.2140/gt.2024.28.1957

N.B. When citing this work, cite the original published paper.

research.chalmers.se offers the possibility of retrieving research publications produced at Chalmers University of Technology. It covers all kind of research output: articles, dissertations, conference papers, reports etc. since 2004. research.chalmers.se is administrated and maintained by Chalmers Library



# Geometry & Topology

Volume 28 (2024)

The volume of pseudoeffective line bundles and partial equilibrium

Tamás Darvas Mingchen Xia





# The volume of pseudoeffective line bundles and partial equilibrium

#### TAMÁS DARVAS MINGCHEN XIA

Let  $(L, he^{-u})$  be a pseudoeffective line bundle on an *n*-dimensional compact Kähler manifold X. Let  $h^0(X, L^k \otimes \mathcal{I}(ku))$  be the dimension of the space of sections *s* of  $L^k$  such that  $h^k(s, s)e^{-ku}$  is integrable. We show that the limit of  $k^{-n}h^0(X, L^k \otimes \mathcal{I}(ku))$  exists, and equals the nonpluripolar volume of  $P[u]_{\mathcal{I}}$ , the  $\mathcal{I}$ -model potential associated to *u*. We give applications of this result to Kähler quantization: fixing a Bernstein–Markov measure v, we show that the partial Bergman measures of *u* converge weakly to the nonpluripolar Monge–Ampère measure of  $P[u]_{\mathcal{I}}$ , the partial equilibrium.

32W20; 53C55

1.	Introduction	1957
2.	Preliminaries	1963
3.	The closure of analytic singularity types in $\mathcal{G}(X, \theta)$	1968
4.	Proof of Theorem 1.1	1972
5.	Envelopes of singularity types with respect to compact sets	1978
6.	Quantization of partial equilibrium measures	1984
Re	ferences	1990

# **1** Introduction

A major theme in Kähler geometry has been the approximation/quantization of natural objects in the theory, going back to a problem of Yau [1987] and early work of Tian [1988]. Initial focus was on the quantization of smooth Kähler metrics, with asymptotic expansion results due to Tian [1990], Bouche [1990], Catlin [1999], Zelditch [1998], Lu [2000] and others. Later, Donaldson [2001] proposed to not just quantize Kähler metrics, but their infinite-dimensional geometry as well. This led to a flurry of activity helping to better understand notions of stability in Kähler geometry; see the work by Berndtsson [2018], Chen and Sun [2012], Phong and Sturm [2006], Song and Zelditch [2010], Darvas, Lu and Rubinstein [Darvas et al. 2020], Zhang [2023] to only mention a few works in a fast expanding literature. We refer to the excellent textbook by Ma and Marinescu [2007] for a detailed discussion and history of many classical results in this direction.

<sup>© 2024</sup> MSP (Mathematical Sciences Publishers). Distributed under the Creative Commons Attribution License 4.0 (CC BY). Open Access made possible by subscribing institutions via Subscribe to Open.

Our work fits into this broad context; however, we consider perhaps the most singular objects one can work with: positively curved metrics on a pseudoeffective line bundle. Despite the fact that potentials of these positively curved metrics are only integrable in general, we will be able to recover their volumes and partial equilibrium measure using quantization, significantly extending the scope of previous results in the literature.

The volume of a pseudoeffective line bundle We now describe our results. Let L be a holomorphic line bundle on a compact connected Kähler manifold  $(X, \omega)$  of dimension n. Let h be a smooth metric on L, and let  $\theta := c_1(L, h)$  denote the Chern form of h. Let  $(T, h_T)$  be an arbitrary Hermitian holomorphic vector bundle on X of rank r, which will be used to *twist* powers of L.

By PSH(*X*,  $\theta$ ) we will denote the space of quasi-plurisubharmonic (quasi-psh) functions v on X such that  $\theta + dd^c v = \theta + (i/2\pi)\partial\overline{\partial}v \ge 0$  in the sense of currents. Here  $d = \partial + \overline{\partial}$  and  $d^c = (i/4\pi)(-\partial + \overline{\partial})$ .

A priori  $PSH(X, \theta)$  may be empty, but if there exists  $u \in PSH(X, \theta)$ , then following terminology of Demailly, we say that the pair  $(L, he^{-u})$  is a pseudoeffective (psef) Hermitian line bundle. Moreover, to such u one can associate a nonpluripolar complex Monge–Ampère measure  $\theta_u^n$ , as introduced in [Boucksom et al. 2010; Guedj and Zeriahi 2007], following ideas by Cegrell [1998] and Bedford and Taylor [1976] in the local case; see Section 2.1 for more details.

We can associate to *u* the so-called  $\mathcal{I}$ -model potential/envelope  $P[u]_{\mathcal{I}} \in PSH(X, \theta)$ , defined by

(1) 
$$P[u]_{\mathcal{I}} := \sup\{w \in \mathsf{PSH}(X,\theta) : w \le 0, \ \mathcal{I}(tw) \subseteq \mathcal{I}(tu) \text{ for } t \ge 0\}.$$

Here  $\mathcal{I}(tu)$  is a multiplier ideal sheaf, locally generated by holomorphic functions f such that  $|f|^2 e^{-tu}$  is integrable. To our knowledge  $P[u]_{\mathcal{I}}$  was first considered in [Kim and Seo 2020], and we studied it in detail in [Darvas and Xia 2022, Section 2.4]; see also [Trusiani 2022].

Let  $H^0(X, T \otimes L^k \otimes \mathcal{I}(ku))$  be the space of global holomorphic sections *s* of  $T \otimes L^k$  satisfying  $\int_X h_T \otimes h^k(s, s) e^{-ku} \omega^n < \infty$ . We also introduce the notation

$$h^0(X, T \otimes L^k \otimes \mathcal{I}(ku)) := \dim_{\mathbb{C}} H^0(X, T \otimes L^k \otimes \mathcal{I}(ku)).$$

It was conjectured by Cao [2014, page 7] and Tsuji [2007, Section 4.4] that

$$\lim_{k \to \infty} \frac{1}{k^n} h^0(X, T \otimes L^k \otimes \mathcal{I}(ku))$$

always exists. We show that this is indeed the case, and we give a precise formula for the limit in terms of the nonpluripolar volume of  $P[u]_{\mathcal{G}}$ :

**Theorem 1.1** Let  $(L, he^{-u})$  be a pseudoeffective Hermitian line bundle on X, and let T be a holomorphic vector bundle of rank r on X. Then

(2) 
$$\lim_{k \to \infty} \frac{1}{k^n} h^0(X, T \otimes L^k \otimes \mathcal{I}(ku)) = \frac{r}{n!} \int_X \theta_{P[u]_{\mathcal{I}}}^n.$$

Geometry & Topology, Volume 28 (2024)

When *L* is ample and *T* is a line bundle, Theorem 1.2 was obtained using non-Archimedean methods in [Darvas and Xia 2022, Theorem 1.4]. As these techniques do not extend to the pseudoeffective case, we take a more elementary approach in this work. In addition, in Section 4.2 we show that the analogue of Theorem 1.1 holds for pseudoeffective  $\mathbb{R}$ -line bundles as well.

In the case that u has analytic singularity type with smooth remainder (see Section 2.2 for the definition), formula (2) is a well-known consequence of the Riemann–Roch theorem of Bonavero [1998, Théorème 1.1, Corollaire 1.2]; see [Darvas and Xia 2022, Theorem 2.26]. In this case, it is possible to apply a resolution of singularities to simplify/principalize the singularity locus of  $\mathscr{I}(u)$ , allowing for a precise asymptotic analysis. In addition, in this case one also has  $\int_X \theta_{P[u]_{\mathscr{I}}}^n = \int_X \theta_u^n$  [Darvas and Xia 2022, Proposition 2.20], simplifying the right-hand side of (2). However, for general  $u \in PSH(X, \theta)$ , one is forced to use the measures  $\theta_{P[u]_{\mathscr{I}}}^n$ , and this is one of the novelties of our work. Indeed, since  $u - \sup_X u \leq P[u]_{\mathscr{I}}$ , [Witt Nyström 2019, Theorem 1.1] gives that  $\int_X \theta_u^n \leq \int_X \theta_{P[u]_{\mathscr{I}}}^n$ , and strict inequality is possible, as pointed out in [Darvas and Xia 2022, Example 2.19].

Formula (2) is also known for  $u := V_{\theta} := \sup\{\varphi \in PSH(X, \theta) : \varphi \leq 0\}$ , the potential with minimal singularity type in PSH(X,  $\theta$ ); see [Boucksom et al. 2010, Proposition 1.18]. In this case we again have  $\int_X \theta_{P[V_{\theta}]_{\theta}}^n = \int_X \theta_{V_{\theta}}^n$ , recovering Boucksom's formula [Boucksom 2002b; Boucksom et al. 2010]:

$$\lim_{k \to \infty} \frac{1}{k^n} h^0(X, L^k) = \frac{1}{n!} \int_X \theta_{V_\theta}^n.$$

The above expression is called the volume of the line bundle L in the literature [Boucksom 2002a; Demailly 2012], justifying our terminology calling  $(1/n!) \int_X \theta_{P[u]_4}^n$  the volume of the pair  $(L, he^{-u})$ .

As T is allowed to be an arbitrary vector bundle in Theorem 1.1, one can hypothesize a version of this result with T being a coherent sheaf on X. This was pointed out to us by László Lempert.

At the slight expense of precision, we briefly describe the strategy behind the proof of Theorem 1.1. By [Witt Nyström 2019, Theorem 1.2], both the left and right sides of (2) only depend on the singularity type of the potential u. As a result, we can use the metric topology of singularity types introduced in [Darvas et al. 2021], and further developed in [Darvas and Xia 2022]. Let us very briefly recall the terminology. For  $v, w \in PSH(X, \theta)$  we say that

- v is more singular than w, and we write  $v \leq w$ , if there exists  $C \in \mathbb{R}$  such that  $v \leq w + C$ ;
- v has the same singularity type as w, and we write  $v \simeq w$ , if  $v \preceq w$  and  $w \preceq v$ .

The classes  $[v] \in \mathcal{G} := \text{PSH}(X, \theta)/\simeq$  of this latter equivalence relation are called *singularity types*. As pointed out in [Darvas et al. 2021], and recalled in Section 2.2,  $\mathcal{G}$  admits a natural pseudometric  $d_{\mathcal{G}}$ , making  $(\mathcal{G}, d_{\mathcal{G}})$  complete (in the presence of positive mass).

By [Darvas and Xia 2022, Proposition 2.20], we have

$$H^{0}(X, T \otimes L^{k} \otimes \mathcal{I}(ku)) = H^{0}(X, T \otimes L^{k} \otimes \mathcal{I}(kP[u]_{\mathcal{I}})) \quad \text{and} \quad P[u]_{\mathcal{I}} = P[P[u]_{\mathcal{I}}]_{\mathcal{I}},$$

ie  $u \to P[u]_{\mathscr{I}}$  is a projection. Hence, it is enough to prove (2) for potentials of the form  $P[u]_{\mathscr{I}}$ . In Section 3 we show that the singularity types  $[P[u]_{\mathscr{I}}] \in \mathscr{S}$  can be  $d_{\mathscr{I}}$ -approximated by analytic singularity types  $[u_j] \in \mathscr{S}$ . It is crucial to work with potentials of the form  $P[u]_{\mathscr{I}}$ , as the same property does not hold for general potentials u.

The proof is then completed by an approximation argument. We take a decreasing sequence  $u_j \in PSH(X, \theta)$  composed of potentials with analytic singularity types such that  $d_{\mathcal{P}}([u_j], [u]) \to 0$ . By Bonavero's theorem we know that (2) holds for each  $u_j$ . It is known that  $d_{\mathcal{P}}([u_j], [u]) \to 0$  implies  $\int_X \theta_{P[u_j]_g}^n \to \int_X \theta_{P[u_j]_g}^n$ , and we will prove a similar convergence result for the left-hand side of (2) as well, to finish the argument.

Let us mention applications of Theorem 1.1 that are treated elsewhere. By [Lazarsfeld and Mustață 2009; Kaveh and Khovanskii 2012] we can naturally assign a family of convex Okounkov bodies  $\Delta(L)$  to a given big line bundle L, depending only on the numerical class of L. Moreover, vol  $L = \text{vol } \Delta(L)$ . In [Xia 2021], based on Theorem 1.1, the second author extended this construction to Hermitian pseudoeffective line bundles: it is possible to define a natural family of convex bodies  $\Delta(L, \phi)$  associated with a given Hermitian pseudoeffective line bundle  $(L, \phi)$  such that vol  $\Delta(L, \phi) = \text{vol}(L, \phi)$ .

Another application concerns automorphic forms. Consider an automorphic line bundle L on a Shimura variety or mixed Shimura variety X. The global sections of  $L^k$  correspond to certain automorphic forms. It is a natural and important question in number theory to understand the asymptotic dimensions of these automorphic forms. In general, X is not compact, but it admits natural smooth compactifications [Ash et al. 2010]. Usually the smooth equivariant metrics on L only extends to singular metrics on a compactification. In this case, Theorem 1.1 can be naturally applied. In the special case of Siegel–Jacobi modular forms, this idea has been carried out concretely in the recent preprints [Botero et al. 2022a; 2022b]. Using a particular case of Theorem 1.1, they managed to prove that the ring of Siegel–Jacobi modular forms is not finitely generated, disproving a well-known claim by Runge [1995].

**Convergence of partial Bergman measures** As another application of Theorem 1.1, we give a very general convergence result for partial Bergman measures to the partial equilibrium, extending the scope of numerous results in the literature.

First we recall terminology introduced in [Berman and Boucksom 2010]. A *weighted subset* of X is a pair (K, v) consisting of a closed nonpluripolar subset  $K \subseteq X$  and a function  $v \in C^0(K)$ . Next, given  $u \in PSH(X, \theta)$ , we tailor the definition of  $\mathscr{I}$ -model envelope from (1) to the pair (K, v):

(3) 
$$P[u]_{\mathscr{I}}(v) := \operatorname{usc}(\sup\{w \in \operatorname{PSH}(X,\theta) : w|_{K} \le v \text{ and } \mathscr{I}(tw) \subseteq \mathscr{I}(tu), t \ge 0\}).$$

Here  $usc(\cdot)$  denotes the least upper semicontinuous envelope. In case that K = X,  $usc(\cdot)$  is unnecessary, moreover we have  $P[u]_{\mathcal{F}}(0) = P[u]_{\mathcal{F}}$ .

Geometry & Topology, Volume 28 (2024)

As a consequence of Corollary 5.7 below,  $\theta_{P[u]_{\mathfrak{f}}(v)}^{n}$  does not put mass on the set  $(X \setminus K) \cup \{P[u]_{\mathfrak{f}}(v) < v\}$ . What is more, when K = X and  $v \in C^{2}(X)$ , the main result of Di Nezza and Trapani [2021] implies that

$$\theta_{P[u]_{\mathscr{I}}(v)}^{n} = \mathbb{1}_{\{P[u]_{\mathscr{I}}(v)=v\}} \theta_{v}^{n}.$$

Analogous properties of equilibrium type measures in different contexts were obtained in [Shiffman and Zelditch 2003; Berman 2009; Ross and Witt Nyström 2017]. With this in mind, we will call the measure  $\theta_{P[u]_{\mathcal{F}}(v)}^{n}$  the *partial equilibrium (measure)* associated to *u* and (*K*, *v*). Theorem 1.2 will further justify this choice of terminology.

Let  $(T, h_T)$  be a Hermitian line bundle. Let  $\nu$  be a Borel probability measure on K. We consider the norms on  $H^0(X, L^k \otimes T)$  given by

$$N_{v,v}^{k}(s) := \left( \int_{K} h^{k} \otimes h_{T}(s,s) e^{-kv} \, \mathrm{d}v \right)^{1/2} \quad \text{and} \quad N_{v,K}^{k}(s) := \sup_{K} (h^{k} \otimes h_{T}(s,s) e^{-kv})^{1/2}.$$

Note that we always have  $N_{v,v}^k(s) \le N_{v,K}^k(s)$ . The measure v is a *Bernstein–Markov measure* with respect to (K, v) if for each  $\varepsilon > 0$ , there is a constant  $C_{\varepsilon} > 0$  such that

(4) 
$$N_{v,K}^{k}(s) \leq C_{\varepsilon} e^{\varepsilon k} N_{v,v}^{k}(s)$$

for any  $s \in H^0(X, L^k \otimes T)$ . A broad class of Bernstein–Markov measures are probability volume forms with respect to (X, v), where  $v \in C^{\infty}(X)$ . For more complicated examples we refer to [Berman et al. 2011, Section 1.2].

We introduce the associated *partial Bergman kernels*: for any  $k \in \mathbb{N}$ ,  $x \in K$ ,

$$B_{\nu,u,\nu}^k(x) := \sup\{h^k \otimes h_T(s,s)e^{-k\nu}(x) : N_{\nu,\nu}^k(s,s) \le 1, s \in H^0(X, L^k \otimes T \otimes \mathcal{I}(ku))\}.$$

The associated partial Bergman measures on X are identically zero on  $X \setminus K$  and on K are defined as

(5) 
$$\beta_{\nu,u,\nu}^k := \frac{n!}{k^n} B_{\nu,u,\nu}^k \,\mathrm{d}\nu.$$

Our next result states that the partial Bergman measures  $\beta_{v,u,v}^k$  quantize the nonpluripolar measure  $\theta_{P[u]_{x}(v)}^n$ , the partial equilibrium of this setting:

**Theorem 1.2** Let  $(L, he^{-u})$  be a pseudoeffective Hermitian holomorphic line bundle on X, and let  $(T, h_T)$  be a Hermitian line bundle. Suppose that v is a Bernstein–Markov measure with respect to a weighted subset (K, v). Then  $\beta_{u,v,v}^k \rightharpoonup \theta_{P[u]_{\psi}(v)}^n$  weakly as  $k \rightarrow \infty$ .

To our knowledge, this result is new even in the case when *L* is assumed to be ample. An important particular case is when *T* is trivial, K = X,  $v \equiv 0$  and  $\mu = \omega^n / \int_X \omega^n$ . In this case we simply put  $\beta_u^k := \beta_{u,0,\omega^n}^k$  and recall that  $P[u]_{\mathscr{I}} = P[u]_{\mathscr{I}}(0)$ . We have the following corollary:

**Corollary 1.3** For  $u \in \text{PSH}(X, \theta)$  we have that  $\beta_u^k \to \theta_{P[u]_{\theta}}^n$  weakly as  $k \to \infty$ .

When T is the trivial line bundle and u has minimal singularity, Theorem 1.2 recovers [Berman et al. 2011, Theorem B]. As part of our argument, in Sections 5 and 6 we also extend [Berman and Boucksom 2010, Theorems A and B] to our partial setting. We suspect that using our results one can now prove equidistribution theorems for (partial) Fekete point configurations, extending [Berman et al. 2011, Theorem A] to our context. However, to stay brief we omit this discussion here.

When T is the trivial line bundle, K = X,  $v \in C^2(X)$ ,  $\mu = \omega^n / \int_X \omega^n$  and u has minimal or exponentially continuous singularity type, we are essentially in the setting of [Berman 2009, Theorem 1.4] and [Ross and Witt Nyström 2017, Theorem 1.4]. Our Theorem 1.2 extends these results, to the extent that our singular setting allows. Indeed, as [u] is of  $\mathscr{I}$ -model type in these cases, we automatically get that  $P[u](v) = P[u]_{\mathscr{I}}(v)$ , where

$$P[u](v) := \operatorname{usc} \sup\{h \in \operatorname{PSH}(X, \theta) : h \le v, [h] \le [u]\}.$$

See Section 3 for more details. Hence, in this case the (partial) Bergman measures converge weakly to  $\theta_{P[u](v)}^n$ . Berman [2009] and Ross and Witt Nyström [2017] actually argue pointwise convergence of the density functions as well, on the locus where P[u](v) = v and  $\theta_v > 0$ . As our v in Theorem 1.2 is only continuous, it is not clear how to interpret the condition  $\theta_v > 0$  in our context.

Observe that  $\int_X \beta_{v,u,v}^k = n! k^{-n} h^0(X, L^k \otimes T \otimes \mathcal{I}(ku))$ . In particular, Theorem 1.2 recovers Theorem 1.1 after an integration. In fact, this plays a crucial role in the argument of Theorem 1.2. As all the measures  $\beta_{u,v,\mu}^k$  have uniformly bounded masses, they form a weakly compact family. The difficulty is to prove that each subsequential limit measure is dominated by  $\theta_{P[u]_{\mathcal{I}}(v)}^n$ . Then the argument is concluded by simply comparing total masses of the limit measures.

The literature on partial Bergman kernels/measures has been fast expanding in many directions. One particular line of study concerns partial Bergman kernels arising from sections vanishing along a smooth divisor V, with the vanishing order increasing in the large limit. As pointed out in numerous works mentioned below, this setup is closely related to ours, when one considers  $L^2$  integrable sections with respect to a weight that has logarithmic singularity along V. It would be interesting to study this connection in the future. One of the first works on this topic was that of Berman [2009], who proved  $L^1$  convergence of the volume densities of the partial Bergman measures. Ross and Singer [2017] and Zelditch and Zhou [2019b] considered this problem in the presence of an  $S^1$ -symmetry near the vanishing locus, identified the forbidden region in terms of the Hamiltonian action, and gave detailed asymptotic expansions. When symmetries are not present, Coman and Marinescu [2017] proved that the partial Bergman kernel has exponential decay near the vanishing locus. For recent extensions to smooth and singular subvarieties V, see [Coman et al. 2019; Sun 2020].

Applications of partial Bergman kernels related to test configurations and geodesic rays were explored in [Ross and Witt Nyström 2014; Darvas and Xia 2022].

In another line of study, Zelditch and Zhou [2019a] initiated the study of partial Bergman kernels that arise from spectral subspaces of the Toeplitz quantization of a smooth Hamiltonian. They showed that their

partial density of states also converges to an equilibrium type measure, suggesting possible connections with our Theorem 1.2. Specifically, given the Hamiltonian data (H, E) of [Zelditch and Zhou 2019a], we wonder if there exists  $v \in C^{\infty}(X)$  and  $u \in PSH(X, \theta)$  such that  $\{H(z) < E\} = \{P[u]_{\mathcal{I}}(v) = v\}$ . If the answer to this question is affirmative, then using the terminology of [Zelditch and Zhou 2019a, Main Theorem] we would obtain that  $\prod_{k,\mathcal{P}_k} \omega^n \rightharpoonup \theta_{P[u]_{\mathcal{I}}(v)}^n$ .

**Acknowledgements** We would like to thank Bo Berndtsson, Junyan Cao, Jakob Hultgren, László Lempert, Yaxiong Liu, Duc-Viet Vu and Steven Zelditch for discussions related to the topic of the paper. We thank the referee for suggesting many improvements. Darvas was partially supported by an Alfred P Sloan Fellowship and National Science Foundation grant DMS-1846942.

**Organization** In Section 2 we recall the relevant notions of envelopes, and adapt results in the literature about the metric topology of singularity types to our context. In Section 3 we characterize the closure of analytic singularity types in a big cohomology class. In Section 4 we prove Theorem 1.1. In Sections 5 and 6 we extend the related results of [Berman and Boucksom 2010; Berman et al. 2011] to our partial context, and prove Theorem 1.2.

# 2 Preliminaries

#### 2.1 Nonpluripolar products and singularity types

Let X be a compact Kähler manifold. Let  $\theta$  be a smooth real (1, 1)-form on X. Let  $PSH(X, \theta)$  be the set of  $\theta$ -plurisubharmonic ( $\theta$ -psh) functions on X. Assume that the cohomology class of  $\theta$  is pseudoeffective, ie that  $PSH(X, \theta)$  is nonempty.

Let  $V_{\theta} := \sup\{v \in PSH(X, \theta) : v \le 0\}$  be the potential with minimal singularity in  $PSH(X, \theta)$ . We recall the construction of nonpluripolar product associated to  $u_1, \ldots, u_n \in PSH(X, \theta)$  from [Boucksom et al. 2010].

Let  $k \in \mathbb{N}$ . Using Bedford–Taylor theory [1976], one can consider the following sequence of measures on *X*:

 $\mathbb{1}_{\bigcap_i \{\varphi_i > V_{\theta} - k\}} (\theta + \mathrm{dd}^{\mathrm{c}} \max(\varphi_1, V_{\theta} - k)) \wedge \cdots \wedge (\theta + \mathrm{dd}^{\mathrm{c}} \max(\varphi_n, V_{\theta} - k)).$ 

It has been shown in [Boucksom et al. 2010, Section 1] that these measures converge weakly to the so-called *nonpluripolar product*  $\theta_{\varphi_1} \wedge \cdots \wedge \theta_{\varphi_n}$  as  $k \to \infty$ . All complex Monge–Ampère measures will be interpreted in this sense in our work.

The resulting positive measure  $\theta_{\varphi_1} \wedge \cdots \wedge \theta_{\varphi_n}$  does not charge pluripolar sets. The particular case when  $u := u_1 = \cdots = u_n$  will yield  $\theta_u^n$ , the nonpluripolar complex Monge–Ampère measure of u.

For any  $u \in PSH(X, \theta)$ , let  $\mathscr{I}(u)$  denote Nadel's multiplier ideal sheaf of u, namely, the coherent ideal sheaf of holomorphic functions f, such that  $|f|^2 e^{-u}$  is integrable. These objects allow us to introduce

an algebraic refinement of the notion of singularity type from the introduction. For  $u, v \in PSH(X, \theta)$  we have the following relations:

- $u \leq_{\mathcal{F}} v$  (also written as  $[u] \leq_{\mathcal{F}} [v]$ ) if  $\mathcal{F}(tu) \subseteq \mathcal{F}(tv)$  for all t > 0.
- $u \simeq_{\mathcal{F}} v$  (also written as  $[u] \simeq_{\mathcal{F}} [v]$ ) if  $u \preceq_{\mathcal{F}} v$  and  $v \preceq_{\mathcal{F}} u$ .

The relation  $\simeq_{\mathscr{I}}$  induces equivalence classes called  $\mathscr{I}$ -singularity types  $[u]_{\mathscr{I}}$ , for any  $u \in PSH(X, \theta)$ . As pointed out in [Darvas and Xia 2022], [u] = [v] implies  $[u]_{\mathscr{I}} = [v]_{\mathscr{I}}$ , but not vice versa.

The different equivalence relations ( $\simeq$  and  $\simeq_{\mathcal{I}}$ ) admit two different envelope notions, as already alluded to in the introduction. Let us revisit them in a very general setup, that will be needed later. Let  $K \subseteq X$ compact and nonpluripolar, and let  $v: K \to [-\infty, \infty]$  measurable. To such v and  $u \in PSH(X, \theta)$  we associate the following notion of envelope:

$$P_{K}^{\theta}[u](v) := (\sup\{w \in \mathsf{PSH}(X, \theta) : [w] \leq [u], w|_{K} \leq v\}),$$
  
$$P_{K}^{\theta}[u]_{\mathscr{I}}(v) := \operatorname{usc}(\sup\{w \in \mathsf{PSH}(X, \theta) : [w] \leq_{\mathscr{I}} [u], w|_{K} \leq v\}).$$

Here and later  $usc(\cdot)$  denotes the upper semicontinuous regularization. We omit  $\theta$  and X from our notation when there is no risk of confusion. In addition, we will use the following shorthand notation, ubiquitous in the literature:

$$P[u] := P_X^{\theta}[u](0), \quad P[u]_{\mathcal{F}} := P_X^{\theta}[u]_{\mathcal{F}}(0).$$

A potential  $u \in PSH(X, \theta)$  is model if u = P[u], and it is  $\mathscr{I}$ -model if  $u = P[u]_{\mathscr{I}}$ .

For any usc function  $f: X \to [-\infty, \infty)$  we define

(6) 
$$P^{\theta}(f) := \operatorname{usc\,sup}\{\varphi \in \operatorname{PSH}(X,\theta) : \varphi \le f\}.$$

Building on the above, for usc functions  $f_1, \ldots, f_N$  we define a notion of rooftop envelope:

 $P^{\theta}(f_1,\ldots,f_N) := P^{\theta}(\min\{f_1,\ldots,f_N\}).$ 

The following lemma was essentially proved in [Darvas et al. 2021]. We recall the short proof as a courtesy to the reader:

**Lemma 2.1** Let  $u, v \in PSH(X, \theta)$  such that  $P^{\theta}(u, v) \in PSH(X, \theta)$ . If u, v are model (resp.  $\mathcal{I}$ -model), then  $P^{\theta}(u, v)$  is also model (resp.  $\mathcal{I}$ -model).

**Proof** Since  $P(u, v) \le \min(u, v)$ , we get that  $P[P(u, v)] \le P[u] = u$  and  $P[P(u, v)] \le P[v] = v$ , hence  $P[P(u, v)] \le P(u, v)$ . This implies P[P(u, v)] = P(u, v), as desired. The statement about  $\mathscr{I}$ -model potentials is proved in the same way.

For any  $x \in X$  and  $u \in PSH(X, \theta)$ , we denote by v(u, x) the Lelong number of  $\varphi$  at x. We recall the following result from [Boucksom et al. 2008], adapted to our context in [Darvas and Xia 2022, Corollary 2.16]:

#### **Proposition 2.2** Let $u, v \in PSH(X, \theta)$ . Then

- (i)  $[u] \leq_{\mathscr{I}} [v]$  if and only if  $v(\pi^*u, y) \geq v(\pi^*v, y)$  for any smooth modification  $\pi: Y \to X$  and any  $y \in Y$ .
- (ii)  $[u] \simeq_{\mathscr{I}} [v]$  if and only if  $v(\pi^*u, y) = v(\pi^*v, y)$  for any smooth modification  $\pi: Y \to X$  and any  $y \in Y$ .

**Corollary 2.3** Let  $u_0, u_1, v_0, v_1 \in \text{PSH}(X, \theta)$  with  $[u_0] \leq_{\mathscr{I}} [v_0]$  and  $[u_1] \leq_{\mathscr{I}} [v_1]$ . For any  $t \in [0, 1]$  we have  $[(1-t)u_0 + tu_1] \leq_{\mathscr{I}} [(1-t)v_0 + tv_1]$ .

**Proof** This follows from Proposition 2.2(i) and the additivity of Lelong numbers [Boucksom 2017, Corollary 2.10]. □

Lastly, we show concavity properties for the envelopes defined above:

**Proposition 2.4** Let  $v \in C^0(K)$  and  $u_0, u_1 \in PSH(X, \theta)$ . The following hold:

- (i) For any  $t \in [0, 1]$ , let  $u_t = tu_1 + (1 t)u_0$ . Then
- (7)  $tP_{K}[u_{1}]_{\mathfrak{F}}(v) + (1-t)P_{K}[u_{0}]_{\mathfrak{F}}(v) \le P_{K}[u_{t}]_{\mathfrak{F}}(v), \quad tP_{K}[u_{1}](v) + (1-t)P_{K}[u_{0}](v) \le P_{K}[u_{t}](v).$ 
  - (ii) If  $[u_0] \leq_{\mathscr{I}} [u_1]$  (resp.  $[u_0] \leq [u_1]$ ), then  $P_K[u_0]_{\mathscr{I}}(v) \leq P_K[u_1]_{\mathscr{I}}(v)$  (resp.  $P_K[u_0](v) \leq P_K[u_1](v)$ ).

**Proof** The proof of (ii) follows from the definitions. To prove (i), let  $h_0, h_1 \in PSH(X, \theta)$  be such that  $[h_i] \leq_{\mathscr{I}} [u_i]$  and  $h_i|_K \leq v$ . Then by Corollary 2.3,  $[th_1 + (1-t)h_0] \leq_{\mathscr{I}} [u_t]$ . It is clear that  $th_1|_K + (1-t)h_0|_K \leq v$ . Hence,  $th_1 + (1-t)h_0 \leq P_K[u_t]_{\mathscr{I}}(v)$ .

As  $h_1$  and  $h_0$  are arbitrary candidates, we conclude the first inequality in (7). The proof of the second inequality is similar.

#### 2.2 The metric topology of singularity types

Let  $\mathscr{G}(X,\theta)$  be the set of singularity types of  $\theta$ -psh functions:  $\mathscr{G}(X,\theta) := \text{PSH}(X,\theta)/\simeq$ . Let  $\mathscr{A}(X,\theta) \subseteq \mathscr{G}(X,\theta)$  be the set of *analytic singularity types*, namely, all singularity types [u] represented by an element  $u \in \text{PSH}(X,\theta)$  such that u is locally of the form

(8) 
$$u = c \log \sum_{i=1}^{N} |f_i|^2 + g$$

where  $c \in \mathbb{Q}^+$ ,  $f_1, \ldots, f_N$  are holomorphic functions and g is a bounded function. When g can be taken to be smooth, then following [Demailly 2018] we say that [u] is a *neat* analytic singularity type.

Darvas et al. [2021] constructed a pseudometric  $d_{\mathcal{G}}$  on  $\mathcal{G}(X, \theta)$ . As we will use the  $d_{\mathcal{G}}$  topology extensively in this work, we recall here a few basic facts, and refer to [Darvas et al. 2021] for a more complete picture.

The definition of  $d_{\mathcal{G}}$  involves embedding  $\mathcal{G}(X, \theta)$  into the space of  $L^1$  geodesic rays [Darvas et al. 2021, Section 3]. We do not recall the exact definition, but simply recall that there is a constant C > 0, depending only on n, such that for any  $[u], [v] \in \mathcal{G}(X, \theta)$  we have

$$(9) \quad d_{\mathcal{G}}([u],[v]) \leq \sum_{j=0}^{n} \left( 2 \int_{X} \theta_{V_{\theta}}^{j} \wedge \theta_{\max\{u,v\}}^{n-j} - \int_{X} \theta_{V_{\theta}}^{j} \wedge \theta_{u}^{n-j} - \int_{X} \theta_{V_{\theta}}^{j} \wedge \theta_{v}^{n-j} \right) \leq C d_{\mathcal{G}}([u],[v]).$$

Note that the term in the middle is independent of the choices of representatives u and v, as a consequence of [Darvas et al. 2018, Theorem 1.1].

**Theorem 2.5** [Darvas et al. 2021, Theorem 1.1] For any  $\delta > 0$ , the space

$$\mathscr{G}_{\delta}(X,\theta) := \left\{ [u] \in \mathscr{G}(X,\theta) \mid \int_{X} \theta_{u}^{n} \ge \delta \right\}$$

is  $d_{\mathcal{G}}$ -complete.

We paraphrase another result, to make it easily adaptable to our context:

**Lemma 2.6** [Darvas et al. 2021, Lemma 4.3] Let  $u, v \in PSH(X, \theta)$  be such that  $[u] \leq [v]$  and  $\int_X \theta_u^n > 0$ . For any

$$b \in \left(1, \left(\frac{\int_X \theta_v^n}{\int_X \theta_v^n - \int_X \theta_u^n}\right)^{1/n}\right),$$

there exists  $h \in PSH(X, \theta)$  such that  $h + (b-1)v \le bu$ . This allows to introduce:

(10)  $P(bu + (1-b)v) := \operatorname{usc} \sup\{h \in \operatorname{PSH}(X, \theta) : h + (b-1)v \le bu\} \in \operatorname{PSH}(X, \theta).$ 

To clarify, when  $\int_X \theta_v^n = \int_X \theta_u^n$  the condition on b in the above result is  $b \in (1, \infty)$ . In addition, by (10), we have that  $P(bu + (1-b)v) + (b-1)v \le u$  a.e. on X, hence this inequality holds globally, since both the left- and right-hand side are quasi-psh functions.

Next we prove continuity results for the envelopes defined above.

**Proposition 2.7** Let  $K \subseteq X$  be a compact and nonpluripolar subset. Let  $v \in C^0(K)$ . Suppose that  $u_j, u \in PSH(X, \theta)$  are such that  $d_{\mathcal{G}}([u_j], [u]) \to 0$  and  $\int_X \theta_u^n > 0$ . Then the following hold:

- (i) If  $u_j \searrow u$  then  $P_K[u_j]_{\mathscr{I}}(v) \searrow P_K[u]_{\mathscr{I}}(v)$  and  $P_K[u_j](v) \searrow P_K[u](v)$ .
- (ii) If  $u_j \nearrow u$  then  $P_K[u_j]_{\mathscr{I}}(v) \nearrow P_K[u]_{\mathscr{I}}(v)$  a.e. and  $P_K[u_j](v) \nearrow P_K[u](v)$  a.e.

The argument is very similar to that of [Darvas and Xia 2022, Lemma 2.21].

**Proof** We first prove (i). Since  $\int_X \theta_{u_j}^n \searrow \int_X \theta_u^n > 0$  [Darvas et al. 2021, Proposition 4.8], by Lemma 2.6, there exists  $\alpha_j \searrow 0$  and  $h_j := P((1/\alpha_j)u + (1-(1/\alpha_j))u_j) \in PSH(X, \theta)$  satisfying  $(1-\alpha_j)u_j + \alpha_j h_j \le u$ . By Proposition 2.4,

$$(1-\alpha_j)P_K[u_j]_{\mathscr{I}}(v) + \alpha_j P_K[h_j]_{\mathscr{I}}(v) \le P_K[(1-\alpha_j)u_j + \alpha_j h_j]_{\mathscr{I}}(v) \le P_K[u]_{\mathscr{I}}(v).$$

Since  $\{u_j\}_j$  is decreasing, so is  $\{P_K[u_j]_{\mathcal{F}}(v)\}_j$ , hence  $w := \lim_j P_K[u_j]_{\mathcal{F}}(v) \ge P[u]_{\mathcal{F}}(v)$  exists. Since  $\alpha_j \to 0$  and  $\sup_X P_K[h_j]_{\mathcal{F}}(v)$  is bounded, we can let  $j \to \infty$  in the above estimate to conclude that  $w = P_K[u]_{\mathcal{F}}(v)$ . The same ideas yield that  $P_K[u_j](v) \searrow P_K[u](v)$ .

Proving (ii) is similar. Since  $\int_X \omega_{u_j}^n \nearrow \int_X \omega_u^n > 0$  [Darvas et al. 2018, Theorem 2.3], by [Darvas et al. 2021, Lemma 4.3] there exists  $\alpha_j \searrow 0$  and  $h_j := P((1/\alpha_j)u_j + (1 - (1/\alpha_j))u) \in PSH(X, \theta)$  satisfying  $(1 - \alpha_j)u + \alpha_j h_j \le u_j$ . By Proposition 2.4,

$$(1-\alpha_j)P_K[u]_{\mathscr{I}}(v) + \alpha_j P_K[h_j]_{\mathscr{I}}(v) \le P_K[(1-\alpha_j)u + \alpha_j h_j]_{\mathscr{I}}(v) \le P_K[u_j]_{\mathscr{I}}(v)$$

Since  $\{u_j\}_j$  is increasing, so is  $\{P_K[u_j]_{\mathscr{I}}(v)\}_j$ , hence  $w := \operatorname{usc}\lim_j P_K[u_j]_{\mathscr{I}}(v) \leq P_K[u]_{\mathscr{I}}(v)$  exists. Since  $\alpha_j \to 0$  and  $\sup_X P_K[h_j]_{\mathscr{I}}(v)$  is bounded, we can let  $j \to \infty$  in the above estimate to conclude that  $w = P_K[u]_{\mathscr{I}}(v)$ . The same proof yields that  $P_K[u_j](v) \nearrow P_K[u](v)$  a.e.

#### 2.3 An approximation result of Demailly

Let X be a compact Kähler manifold of dimension n. Let  $\theta$  be a smooth representative of a pseudoeffective (1, 1)–class on X. Let  $\omega$  be a Kähler form on X.

Following the terminology of Cao [2014, Definition 2.3], we recall the existence of quasi-equisingular approximation for potentials in  $PSH(X, \theta)$ . As elaborated below, this result is implicit in the proof of [Demailly et al. 2001, Theorem 2.2.1; Demailly and Paun 2004, Theorem 3.2; Demailly 2015, Theorem 1.6].

**Theorem 2.8** Let  $u \in PSH(X, \theta)$ . Then there exists  $u_k^D \in PSH(X, \theta + \varepsilon_k \omega)$  with  $\varepsilon_k \searrow 0$  such that

(i)  $u_k^D \searrow u$ , (ii)  $[u_k^D] \in \mathcal{A}(X, \theta + \varepsilon_k \omega)$ , (iii)  $\mathcal{I}(s2^k/(2^k - s))u_k^D) \subseteq \mathcal{I}(su) \subseteq \mathcal{I}(su_k^D)$  for all s > 0.

**Proof** Parts (i) and (ii) follow from [Demailly 2015, Theorem 1.6]. The second inclusion of (iii) follows from  $u \le u_k^D$ , whereas the first inclusion of (iii) follows from [Demailly 2015, Corollary 1.12].

As shown in [Demailly 2012, page 135, formula (13.14)] (or [Demailly and Paun 2004, Theorem 3.2(iv)]), for each  $u_k^D$  in the above theorem, there exists a holomorphic modification  $\pi_k : Y_k \to X$ , a smooth closed (1, 1)–form  $\beta_k$ , and a  $\mathbb{Q}$ -divisor  $D_k$  with snc singularities on Y such that

(11) 
$$\theta_{u_k^D} = [D_k] + \beta_k.$$

In particular,  $u_k^D \circ \pi_k$  has neat analytic singularity type; recall (8).

When the pseudoeffective class is induced by a line bundle, we have a related approximation result:

**Remark 2.9** In the case that  $(L, h) \to X$  is a Hermitian line bundle with  $c_1(L, h) = \{\theta\}, (T, h_T) \to X$  is an arbitrary Hermitian line bundle, and  $\theta_u$  is a Kähler current with  $[u] \in \mathcal{A}(X, \theta)$ , it is possible to work with the following alternative approximating sequence:

(12) 
$$\widetilde{u}_{k}^{D} = \frac{1}{k} \log \sup_{\substack{s \in H^{0}(X, L^{k} \otimes T) \\ \int_{X} h^{k} \otimes h_{T}(s, s) e^{-ku} \omega^{n} \leq 1}} h^{k} \otimes h_{T}(s, s).$$

For k big enough, this sequence will satisfy  $\tilde{u}_k^D + C \log k/k \ge u$ , by the Ohsawa–Takegoshi theorem. However, it is not monotone in general. On the other hand, a stronger form of condition (iii) will hold in this case, namely  $[u] \le [\tilde{u}_k^D] \le [\alpha_k u]$  for some  $\alpha_k \nearrow 1$ .

**Proof** This is a known consequence of the Briancon–Skoda theorem [Demailly 2012], but as a courtesy to the reader we give a brief argument for the estimate  $[\tilde{u}_k^D] \leq [\alpha_k u]$ , the only part that needs to be proved. As we point out now, this actually follows from the arguments of [Demailly 2012, Remark 5.9].

Let  $\mathcal{J}$  be the coherent sheaf of holomorphic functions g satisfying  $|g| \leq De^{u/2c}$  with  $c \in \mathbb{Q}^+$ , as in (8), and D > 0 some positive constant. As pointed out in [Demailly 2012, Remark 5.9], we may assume that the  $f_j$  in (8) are local generators of  $\mathcal{J}$ .

Let  $\pi: Y \to X$  be a smooth modification such that  $\pi^{-1} \mathscr{G} \cdot \mathbb{O}_Y = \mathbb{O}(-D)$ , where  $D = \sum_j \lambda_j D_j$  is a normal crossing divisor on Y. The existence of such  $\pi$  follows from Hironaka desingularization.

Now suppose that  $s \in H^0(X, L^k \otimes T)$  satisfies  $\int_X h^k \otimes h_T(s, s) e^{-ku} \omega^n \leq 1$ . By pulling back we obtain

$$\int_Y h^k \otimes h_T(s \circ \pi, s \circ \pi) e^{-ku \circ \pi} (\pi^* \omega)^n \leq 1.$$

As  $u \circ \pi \simeq c \sum_j \lambda_j \log g_j$  for some local generators  $g_j$  of  $\mathbb{O}(-D_j)$ , by Fubini's theorem  $h^k \otimes h_T(s \circ \pi, s \circ \pi)$ vanishes to order at least  $\lfloor kc\lambda_j \rfloor + d$  along  $D_j$ , where d is an absolute constant, only dependent on  $\pi$ . In particular, one can find  $\alpha_k \nearrow 1$  such that  $h^k \otimes h_T(s \circ \pi, s \circ \pi)$  vanishes to order at least  $c\alpha_k k\lambda_j$  along  $D_j$ . Since  $u \circ \pi \simeq c \sum_j \lambda_j \log g_j$ , we obtain that  $[(1/k) \log h^k \otimes h_T(s \circ \pi, s \circ \pi)] \preceq [\alpha_k u \circ \pi]$ , which in turn gives  $[\tilde{u}_k^D \circ \pi] \preceq [\alpha_k u \circ \pi]$ , since  $H^0(X, L^k \otimes T \otimes \mathcal{I}(ku))$  is finite-dimensional. By pushing forward, we obtain that  $[\tilde{u}_k^D] \preceq [\alpha_k u]$ , as desired.

# **3** The closure of analytic singularity types in $\mathcal{G}(X, \theta)$

In this section we only assume that  $\theta$  is a smooth representative of a big (1, 1)–cohomology class on X. Our goal is to prove that the  $d_{\mathcal{G}}$ –closure of  $\mathcal{A}(X, \theta)$  is the space of  $\mathcal{I}$ –model singularity types, in the presence of positive mass. We start with an elementary lemma:

**Lemma 3.1** Let  $\pi: X' \to X$  be a smooth modification and  $u \in PSH(X, \theta)$ . Then we have

$$\pi^* P^{\theta}[u]_{\mathcal{Y}} = P^{\pi^* \theta}[\pi^* u]_{\mathcal{Y}}.$$

**Proof** Recall that

(13) 
$$P^{\theta}[u]_{\mathscr{Y}} = \sup\{v \in \mathrm{PSH}(X,\theta) : v \le 0, [v] \le_{\mathscr{Y}} [u]\}.$$

Let  $v \in \text{PSH}(X, \theta)$  be a candidate of the sup in (13). Then by Proposition 2.2, for any smooth modification  $p: Y \to X$  and any  $y \in Y$ ,  $v(p^*v, y) \ge v(p^*u, y)$ . In particular, for any smooth modification  $q: Z \to X'$  and any  $z \in Z$ , we have  $v(q^*\pi^*v, z) \ge v(q^*\pi^*u, z)$ . By Proposition 2.2 again,  $[\pi^*v] \preceq_{\mathscr{I}} [\pi^*u]$ . In particular,  $\pi^*v \le P^{\pi^*\theta}[\pi^*u]_{\mathscr{I}}$ . We arrive at the inequality

$$\pi^*(P^{\theta}[u]_{\mathscr{I}}) \le P^{\pi^*\theta}[\pi^*u]_{\mathscr{I}}.$$

It remains to prove the reverse inequality. There is a unique  $h \in PSH(X, \theta)$  such that  $\pi^* h = P^{\theta}[\pi^* u]_{\mathcal{F}}$ . We need to prove that  $P^{\theta}[u]_{\mathcal{F}} \ge h$ . It suffices to prove the following claim: for any k > 0,  $\mathcal{I}(ku) \supseteq \mathcal{I}(kh)$ . But we already know that  $\mathcal{I}(k\pi^* u) = \mathcal{I}(k\pi^* h)$ , while by [Demailly 2012, Proposition 5.8],

$$\mathfrak{P}(ku) = \pi_*(K_{X'/X} \otimes \mathfrak{P}(k\pi^*u)) \quad \text{and} \quad \mathfrak{P}(kh) = \pi_*(K_{X'/X} \otimes \mathfrak{P}(k\pi^*h)).$$

Hence, we conclude that  $\mathcal{I}(ku) = \mathcal{I}(kh)$ .

**Lemma 3.2** If  $u \in PSH(X, \theta)$  satisfies  $[u] \in \mathcal{A}(X, \theta)$ , then  $[u] = [P[u]] = [P[u]_{\mathcal{J}}]$ .

**Proof** Since  $u \sim_{\mathcal{J}} P[u]_{\mathcal{J}}$ , we get  $[u] = [P[u]_{\mathcal{J}}]$  from [Kim 2015, Theorem 4.3]. Since  $[u] \preceq [P[u]] \preceq [P[u]_{\mathcal{J}}]$ , it also follows that [u] = [P[u]].

**Proposition 3.3** Let  $u \in \text{PSH}(X, \theta)$ . Then  $P^{\theta + \varepsilon_j \omega}[u_j^D]_{\mathscr{I}} \searrow P^{\theta}[u]_{\mathscr{I}}$  as  $j \to \infty$ , where the sequence  $u_j^D \in \text{PSH}(X, \theta + \varepsilon_j \omega)$  is the approximating sequence of Theorem 2.8. Moreover, if  $\theta_u$  is a Kähler current, then  $P^{\theta}[u_j^D]_{\mathscr{I}} \searrow P^{\theta}[u]_{\mathscr{I}}$  as  $j \to \infty$ .

**Proof** We can suppose that  $u \leq 0$ . Since  $[u_j^D] \geq [u]$  we have that  $P^{\theta + \varepsilon_j \omega}[u_j^D]_{\mathscr{I}} \geq P^{\theta + \varepsilon_j \omega}[u]_{\mathscr{I}} \geq P^{\theta}[u]_{\mathscr{I}}$ . Since  $\{u_j^D\}_j$  is decreasing, we have that  $v := \lim_j P^{\theta + \varepsilon_j \omega}[u_j^D]_{\mathscr{I}} \in PSH(X, \theta)$  exists and  $u \leq v$ .

Observe that  $P^{\theta}[v]_{\mathscr{I}} = v$ , since any candidate  $h \in \text{PSH}(X, \theta)$  for  $P^{\theta}[v]_{\mathscr{I}}$  is also a candidate for each  $P^{\theta+\varepsilon_{j}\omega}[u_{j}^{D}]_{\mathscr{I}}$ . Hence, to finish the argument, it is enough to show that  $\mathscr{I}(tu) = \mathscr{I}(tv)$  for all t > 0. By Theorem 2.8, for any  $\delta > 1$  and t > 0 there exists  $k_{0}(\delta, t) > 0$  such that for all  $k \ge k_{0}$  we have  $\mathscr{I}(t\delta v) \subseteq \mathscr{I}(t\delta u_{k}^{D}) \subseteq \mathscr{I}(tu)$ . Letting  $\delta \searrow 1$ , the strong openness theorem of Guan and Zhou [2015] implies that  $\mathscr{I}(tv) \subseteq \mathscr{I}(tu)$ . Since the reverse inclusion is trivial, the proof of the first assertion is finished.

To prove the second assertion, assume that  $\theta_u$  is a Kähler current. Hence, for j large enough, it holds that  $u_j^D \in \text{PSH}(X, \theta)$ . On the other hand, observe that  $P^{\theta + \varepsilon_j \omega}[u_j^D]_{\mathscr{I}} \geq P^{\theta}[u_j^D]_{\mathscr{I}} \geq P^{\theta}[u]_{\mathscr{I}}$ , hence  $P^{\theta}[u_j^D]_{\mathscr{I}} \searrow P^{\theta}[u]_{\mathscr{I}}$  as  $j \to \infty$ .

We note the following important corollary of this result, which will be used numerous times in this work:

 $\Box$ 

Tamás Darvas and Mingchen Xia

**Corollary 3.4** Let  $u \in PSH(X, \theta)$ . Then

$$\int_X (\theta + \varepsilon_j \omega)_{u_j^D}^n = \int_X (\theta + \varepsilon_j \omega)_{P^{\theta + \varepsilon_j \omega}[u_j^D]_{\mathscr{I}}}^n \searrow \int_X \theta_{P^{\theta}[u]_{\mathscr{I}}}^n \quad \text{as } j \to \infty,$$

where  $u_i^D \in \text{PSH}(X, \theta + \varepsilon_j \omega)$  is the approximating sequence of Theorem 2.8.

**Proof** The equality follows from Lemma 3.2 and [Witt Nyström 2019, Theorem 1.1]. Since  $P^{\theta + \varepsilon_j \omega}[u_j^D]_{\mathscr{I}} \ge P^{\theta}[u]_{\mathscr{I}}$ , we can start with the following inequality:

$$\lim_{i \to \infty} \int_X (\theta + \varepsilon_j \omega)_{P^{\theta + \varepsilon_j \omega}[u_j^D]_{\mathscr{I}}}^n \ge \lim_{j \to \infty} \int_X (\theta + \varepsilon_j \omega)_{P^{\theta}[u]_{\mathscr{I}}}^n = \int_X \theta_{P^{\theta}[u]_{\mathscr{I}}}^n.$$

To finish the proof, we will argue that  $\overline{\lim}_{j} \int_{X} (\theta + \varepsilon_{j} \omega)_{P^{\theta + \varepsilon_{j} \omega}[u_{j}^{D}]_{j}}^{n} \leq \int_{X} \theta_{P^{\theta}[u]_{j}}^{n}$ . Indeed, fixing  $j_{0} \in \mathbb{N}$ ,

$$\overline{\lim_{j \to \infty}} \int_{X} (\theta + \varepsilon_{j}\omega)_{P^{\theta + \varepsilon_{j}\omega}[u_{j}^{D}]_{\delta}}^{n} = \overline{\lim_{j \to \infty}} \int_{\{P^{\theta + \varepsilon_{j}\omega}[u_{j}^{D}]_{\delta}=0\}} (\theta + \varepsilon_{j}\omega)_{P^{\theta + \varepsilon_{j}\omega}[u_{j}^{D}]_{\delta}}^{n} \\
\leq \overline{\lim_{j \to \infty}} \int_{\{P^{\theta + \varepsilon_{j}\omega}[u_{j}^{D}]_{\delta}=0\}} (\theta + \varepsilon_{j_{0}}\omega)_{P^{\theta + \varepsilon_{j}\omega}[u_{j}^{D}]_{\delta}}^{n} \\
\leq \int_{\{P^{\theta}[u]_{\delta}=0\}} (\theta + \varepsilon_{j_{0}}\omega)_{P^{\theta}[u]_{\delta}}^{n},$$

where in the first line we have used that  $P^{\theta+\varepsilon_j\omega}[u_j^D]_{\mathcal{I}} = P^{\theta+\varepsilon_j\omega}[u_j^D]$  from [Darvas and Xia 2022, Proposition 2.20] together with [Darvas et al. 2018, Theorem 3.8], and in the last line we have used Proposition 3.3 and [Darvas et al. 2021, Proposition 4.6]. Letting  $j_0 \to \infty$ , we arrive at the desired conclusion:

$$\overline{\lim_{j \to \infty}} \int_X (\theta + \varepsilon_j \omega)_{u_j^D}^n \leq \underline{\lim_{j_0 \to \infty}} \int_{\{P^{\theta}[u]_{\mathscr{I}} = 0\}} (\theta + \varepsilon_{j_0} \omega)_{P^{\theta}[u]_{\mathscr{I}}}^n = \int_{\{P^{\theta}[u]_{\mathscr{I}} = 0\}} \theta_{P^{\theta}[u]_{\mathscr{I}}}^n \leq \int_X \theta_{P^{\theta}[u]_{\mathscr{I}}}^n . \quad \Box$$

**Corollary 3.5** Let  $\psi \in PSH(X, \theta)$ . The following hold:

(i) 
$$\int_X (\theta + \varepsilon \omega + \mathrm{dd}^c P^{\theta + \varepsilon \omega} [\psi]_{\mathfrak{f}})^n \searrow \int_X \theta_{P^{\theta}[\psi]_{\mathfrak{f}}}^n as \varepsilon \searrow 0.$$
  
(ii) If  $\theta_{\psi}$  is a Kähler current, then  $\int_X (\theta - \varepsilon \omega + \mathrm{dd}^c P^{\theta - \varepsilon \omega} [\psi]_{\mathfrak{f}})^n \nearrow \int_X \theta_{P^{\theta}[\psi]_{\mathfrak{f}}}^n as \varepsilon \searrow 0.$ 

**Proof** We approximate  $\psi$  with  $\psi_j^D \in PSH(X, \theta + \varepsilon_j \omega)$  from Theorem 2.8. For  $\varepsilon > 0$ , applying Corollary 3.4 for  $\psi \in PSH(X, \theta + \varepsilon \omega)$  (for the same approximating sequence  $\psi_j^D \in PSH(X, \theta + (\varepsilon + \varepsilon_j)\omega)$  independent of  $\varepsilon$ ) we get that

$$\int_{X} (\theta + \varepsilon \omega + \mathrm{dd}^{\mathrm{c}} P^{\theta + \varepsilon \omega} [\psi]_{\mathscr{I}})^{n} = \lim_{j \to \infty} \int_{X} (\theta + (\varepsilon + \varepsilon_{j})\omega + \mathrm{dd}^{\mathrm{c}} \psi_{j}^{D})^{n},$$
$$\int_{X} (\theta + \mathrm{dd}^{\mathrm{c}} P^{\theta} [\psi]_{\mathscr{I}})^{n} = \lim_{j \to \infty} \int_{X} (\theta + \varepsilon_{j}\omega + \mathrm{dd}^{\mathrm{c}} \psi_{j}^{D})^{n}.$$

Using the multilinearity of nonpluripolar products, (i) follows. The proof of (ii) follows the same pattern and is left to the reader.  $\Box$ 

Geometry & Topology, Volume 28 (2024)

**Proposition 3.6** Let  $u \in PSH(X, \theta)$  be such that  $\int_X \theta_u^n > 0$ . Then there exists  $v \in PSH(X, \theta)$  such that  $u \ge v$  and  $\theta_v \ge \delta \omega$  for some  $\delta > 0$ .

**Proof** We may assume that  $u \le 0$ . Since  $u \le V_{\theta}$  and  $\int_X \theta_{V_{\theta}}^n \ge \int_X \theta_u^n > 0$ , by Lemma 2.6 there exists b > 0 such that  $h := P((1+b)u - bV_{\theta}) \in PSH(X, \theta)$  and

$$\frac{b}{b+1}V_{\theta} + \frac{1}{b+1}h \le u.$$

By [Boucksom 2002b], there exists  $w \in PSH(X, \theta)$  such that  $w \le 0$  and  $\theta_w \ge \delta' \omega$  for some  $\delta' > 0$ . Since  $w \leq V_{\theta}$ , we obtain that

$$v := \frac{b}{b+1}w + \frac{1}{b+1}h \le u$$

and  $\theta_v \geq (b\delta'/(b+1))\omega$ .

Next we extend [Darvas and Xia 2022, Theorem 2.24] to big cohomology classes.

**Lemma 3.7** Let  $u \in PSH(X, \theta)$ . Assume that  $\theta_u$  is a Kähler current. Let  $u_k^D$  be the approximation sequence in Theorem 2.8. Then

(14) 
$$d_{\mathcal{G}}([u_k^D], P^{\theta}[u]_{\mathcal{F}}) \to 0 \quad \text{as } k \to \infty$$

In particular,

(15) 
$$\lim_{k \to \infty} \int_X \theta_{u_k}^n = \int_X \theta_{P^{\theta}[u]_{\mathfrak{f}}}^n.$$

**Proof** First observe that  $u_k^D \in PSH(X, \theta)$  when k is large enough, so (14) indeed makes sense. The second assertion follows from the first and [Darvas et al. 2021, Lemma 3.7], so it suffices to prove the first. By Proposition 3.3,  $P^{\theta}[u_k^D]_{\mathcal{I}}$  decreases to  $P^{\theta}[u]_{\mathcal{I}}$  as  $k \to \infty$ .

Since the potentials  $P^{\theta}[u_k^D]_{\mathcal{I}}$  are model [Darvas and Xia 2022, Proposition 2.18(i)], by [Darvas et al. 2021, Lemma 3.6, Proposition 4.8] we obtain that  $d_{\mathcal{F}}([P^{\theta}[u]_{\mathscr{I}}], [P^{\theta}[u_k^D]]_{\mathscr{I}}) \to 0$  as  $k \to \infty$ . By Lemma 3.2 we conclude (14). 

**Theorem 3.8** Suppose that  $u \in \text{PSH}(X, \theta)$  is such that  $\int_X \theta_u^n > 0$ . Then  $[u] \in \overline{\mathcal{A}(X, \theta)}^{d_g}$  if and only if  $[P[u]] = [P[u]_{\mathcal{F}}]$ . Additionally, if  $[P[u]] = [P[u]_{\mathcal{F}}]$  and  $\theta_u$  is a Kähler current, then the regularization sequence  $\{[u_k^D]\}_k$  of Theorem 2.8  $d_{\mathcal{G}}$ -converges to [u].

Here the notation  $\overline{\mathcal{A}(X,\theta)}^{d_{\mathcal{G}}}$  means the closure of  $\mathcal{A}(X,\theta)$  in  $\mathcal{G}(X,\theta)$  with respect to the  $d_{\mathcal{G}}$ -metric.

**Proof** To begin, let  $v \in PSH(X, \theta)$  be such that  $v \le u$  and  $\theta_v \ge \delta \omega$  for some  $\delta > 0$ . Such v exists by Proposition 3.6. Let  $v_t := (1-t)v + tu$ , with  $t \in [0, 1]$ . Then  $\theta_{v_t}$  is a Kähler current for  $t \in [0, 1)$  and  $v_t \nearrow u$  a.e. as  $t \nearrow 1$ .

Assume first that  $[P[u]_{\mathcal{I}}] = [P[u]]$ . By replacing u with  $P[u]_{\mathcal{I}}$ , we can additionally assume that  $u = P[u]_{\mathcal{I}}$ . By [Darvas and Xia 2022, Lemma 2.21(iii)] we obtain that  $P[v_t]_{\mathcal{F}} \nearrow P[u]_{\mathcal{F}} = u$  a.e. as  $t \to 1$ . In particular, by [Darvas et al. 2021, Lemma 4.1] we obtain that  $d_{\mathcal{G}}(P[v_t]_{\mathcal{I}}, [u]) \to 0$  as  $t \to 1$ .

Let us fix  $t \in [0, 1)$ . By the above, it is enough to argue that  $[P[v_t]_{\mathcal{F}}] \in \overline{\mathcal{A}}^{d_{\mathcal{F}}}$ . For this we apply the regularization method of Theorem 2.8 to  $v_t$ , obtaining  $v_{t,k}^D \in \text{PSH}(X, \theta)$  such that  $[v_{t,k}^D] \in \mathcal{A}(X, \theta)$  (we used here that  $\theta_{v_t}$  is a Kähler current). By Lemma 3.7,  $d_{\mathcal{F}}([v_{t,k}^D], [v_t]) \to 0$  as  $k \to \infty$ . So  $[P[v_t]_{\mathcal{F}}] \in \overline{\mathcal{A}}^{d_{\mathcal{F}}}$ , and we conclude.

In the reverse direction, suppose there exists  $[v_j] \in \mathcal{A}(X, \theta)$  such that  $d_{\mathcal{G}}([v_j], [u]) \to 0$ . By Lemma 3.2, we can assume that  $v_j = P[v_j]_{\mathcal{F}} = P[v_j]$ . In addition, we can assume that u = P[u], since  $d_{\mathcal{G}}(u, P[u]) = 0$  [Darvas et al. 2021, Theorem 3.3]. Since  $\int_X \theta_u^n > 0$ , after possibly restricting to a subsequence of  $v_j$ , we can use [Darvas et al. 2021, Theorem 5.6] to conclude existence of an increasing sequence of model potentials  $\{w_j\} \in PSH(X, \theta)$  such that  $w_j \leq v_j$  and  $d_{\mathcal{G}}([w_j], [u]) \to 0$ . As pointed out after the statement of [Darvas et al. 2021, Theorem 5.6], after possibly taking a subsequence of the  $v_j$ , we can take

$$w_j := \lim_{k \to \infty} P(v_j, v_{j+1}, \dots, v_{j+k}).$$

Since all the  $v_i$  are  $\mathcal{I}$ -model, an iterated application of Lemma 2.1 implies that so is

$$h_{j,k} := P(v_j, v_{j+1}, \dots, v_{j+k}).$$

Moreover, since  $w_j$  is the decreasing limit of the  $h_{j,k}$ , then  $w_j$  is  $\mathscr{I}$ -model too [Darvas and Xia 2022, Lemma 2.21(i)]. Lastly, since u is the increasing limit of the  $w_j$ , then u is  $\mathscr{I}$ -model as well [Darvas and Xia 2022, Lemma 2.21(iii)].

# 4 **Proof of Theorem 1.1**

Let X be a connected compact Kähler manifold of dimension n. For this section, let T be an arbitrary holomorphic vector bundle on X, with rank r.

#### 4.1 The case of integral line bundles

Let *L* be a pseudoeffective line bundle on *X*. Let *h* be a smooth Hermitian metric on *L* such that  $\theta := c_1(L, h)$ . We fix a Kähler form  $\omega$  on *X* such that  $\omega - \theta$  is a Kähler form.

**Proposition 4.1** Suppose that  $u \in PSH(X, \theta)$ . Then

$$\lim_{k \to \infty} \frac{1}{k^n} h^0(X, T \otimes L^k \otimes \mathcal{I}(ku)) \leq \frac{r}{n!} \int_X \theta_{P[u]_{\mathcal{I}}}^n$$

**Proof** Since  $P[P[u]_{\mathcal{F}}] = P[u]_{\mathcal{F}}$  and  $\mathcal{F}(sP[u]_{\mathcal{F}}) = \mathcal{F}(su)$  for all s > 0 [Darvas and Xia 2022, Proposition 2.18(ii)], we can replace u with  $P[u]_{\mathcal{F}}$  to assume that u is  $\mathcal{F}$ -model.

Next we apply the regularization method of Theorem 2.8 to u, obtaining  $u_j^D \in \text{PSH}(X, \theta + \varepsilon_j \omega)$  such that  $[u_j^D] \in \mathcal{A}(X, \theta + \varepsilon_j \omega)$  and  $u_j^D \searrow u$ . Let  $\pi_k \colon Y_k \to X$  be the smooth resolution of singularities of (11).

By [Demailly 2012, Proposition 5.8] and [Bonavero 1998, Théorème 2.1] applied to q = 0 on  $Y_k$  (see also [Ma and Marinescu 2007, Theorem 2.3.18]), we obtain that

$$\begin{split} \overline{\lim_{k \to \infty} \frac{1}{k^n}} h^0(X, T \otimes L^k \otimes \mathscr{I}(ku)) &\leq \overline{\lim_{k \to \infty} \frac{1}{k^n}} h^0(X, T \otimes L^k \otimes \mathscr{I}(ku_j^D)) \\ &= \overline{\lim_{k \to \infty} \frac{1}{k^n}} h^0(Y, \pi_k^* T \otimes (\pi_k^* L)^k \otimes K_{Y/X} \otimes \mathscr{I}(ku_j^D \circ \pi_k)) \\ &\leq \frac{r}{n!} \int_{Y_k(0)} \pi_k^* \theta_{u_j^D}^n &= \frac{r}{n!} \int_{\pi_k(Y_k(0))} \theta_{u_j^D}^n \\ &\leq \frac{r}{n!} \int_{\pi_k(Y_k(0))} (\theta + \varepsilon_j \omega)_{u_j^D}^n \leq \frac{r}{n!} \int_X (\theta + \varepsilon_j \omega)_{u_j^D}^n, \end{split}$$

where  $Y_k(0) \subseteq Y_k$  is the set contained in the smooth locus of the (1, 1)-current  $\pi_k^* \theta_{u_j^D}$  where the eigenvalues of  $\pi_k^* \theta_{u_j^D}$  are all positive. By Corollary 3.4,  $\lim_{j\to\infty} \int_X (\theta + \varepsilon_j \omega)_{u_j^D}^n = \int_X \theta_u^n$ , finishing the argument.

**Lemma 4.2** Let  $u \in PSH(X, \theta)$  such that  $\theta_u$  is a Kähler current. Let  $\beta \in (0, 1)$ . Then there exists  $k_0 := k_0(u, \beta)$  such that for all  $k \ge k_0$  there exists  $v_{\beta,k} \in PSH(X, \theta)$  satisfying

(i)  $P[u]_{\mathscr{I}} \ge (1-\beta)u_k^D + \beta v_{\beta,k},$ (ii)  $\int_X \theta_{v_{\beta,k}}^n > 0.$ 

**Proof** Due to Lemma 3.7, we have that  $\int_X \theta_{u_k^D}^n \searrow \int_X \theta_{P[u]_{\#}}^n$ . In particular, there exists  $k_0 > 0$  such that

$$\frac{1}{\beta^n} < \frac{\int_X \theta_{u_k^D}^n}{\int_X \theta_{u_k^D}^n - \int_X \theta_{P[u]_{\mathcal{F}}}^n} \quad \text{for all } k \ge k_0.$$

By Lemma 2.6 we obtain that

$$v_{k,\beta} := P\left(\frac{1}{\beta}P[u]_{\mathscr{I}} - \frac{1-\beta}{\beta}u_k^D\right) \in \mathrm{PSH}(X,\theta) \quad \text{and} \quad P[u]_{\mathscr{I}} \ge (1-\beta)u_k^D + \beta v_{\beta,k}.$$

Now we show that  $v_{\beta,k}$  has positive mass. Pick  $\beta' \in (0,\beta)$  such that

$$\frac{1}{\beta'^n} < \frac{\int_X \theta_{u_k^D}^n}{\int_X \theta_{u_k^D}^n - \int_X \theta_{P[u]_{\mathcal{I}}}^n} \quad \text{for all } k \ge k_0.$$

Then

$$h := P\left(\frac{1}{\beta'}P[u]_{\mathcal{I}} - \frac{1-\beta'}{\beta'}u_k^D\right) \in \mathrm{PSH}(X,\theta)$$

is defined as well, and  $v_{k,\beta} \ge (\beta'/\beta)h + ((\beta - \beta')/\beta)u_k^D \in PSH(X, \theta)$ , implying that

$$\int_X \theta_{v_{k,\beta}}^n \ge \frac{(\beta - \beta')^n}{\beta^n} \int_X \theta_{u_k^D}^n \ge \frac{(\beta - \beta')^n}{\beta^n} \int_X \theta_u^n > 0,$$

where we applied [Witt Nyström 2019, Theorem 1.1] twice.

Geometry & Topology, Volume 28 (2024)

Tamás Darvas and Mingchen Xia

**Proposition 4.3** Suppose that  $u \in PSH(X, \theta)$  with  $\theta_u > \delta \omega$  for some  $\delta > 0$ . Then

$$\lim_{j\to\infty}\frac{1}{j^n}h^0(X,T\otimes L^j\otimes\mathfrak{I}(ju))\geq \frac{r}{n!}\int_X\theta_{P[u]_g}^n.$$

**Proof** To start, we fix a number  $\beta = p/q \in (0, \min(\delta, 1)) \cap \mathbb{Q}$ . It suffices to show that there is a constant C > 0, only dependent on r, n and  $\theta$ , such that

$$\lim_{j\to\infty}\frac{1}{j^n}h^0(X,T\otimes L^j\otimes \mathcal{I}(ju))\geq \frac{r}{n!}\int_X\theta_{P[u]_{\mathcal{I}}}^n-C\beta.$$

Writing j = aq + b for some b = 0, ..., q - 1, observe that

$$h^{0}(X, T \otimes L^{j} \otimes \mathcal{I}(ju)) \ge h^{0}(X, T \otimes L^{b-q} \otimes L^{(a+1)q} \otimes \mathcal{I}((a+1)qu)).$$

Absorbing  $L^{b-q}$  into T, and noticing that b-q can only take a finite number of values, we find that it suffices to prove that

(16) 
$$\lim_{j \to \infty} \frac{1}{j^n q^n} h^0(X, T \otimes L^{jq} \otimes \mathcal{I}(jqu)) \ge \frac{r}{n!} \int_X \theta_{P[u]_{\mathcal{I}}}^n - C\beta$$

for an arbitrary twisting bundle T.

By Lemma 4.2, there is  $k_0 > 0$  depending on  $\beta$  and u such that for  $k \ge k_0$ , there exists a potential  $v_{\beta,k} \in PSH(X, \theta)$  of positive mass such that

$$P[u]_{\mathscr{I}} \ge w_{\beta,k} := (1-\beta)u_k^D + \beta v_{\beta,k} \quad \text{for all } k \ge k_0.$$

For big enough  $k_0$  we also have  $\theta_{u_k^D} > \beta \omega \ge \beta \theta$  for all  $k \ge k_0$ . In particular,  $u_k^D \in \text{PSH}(X, (1-\beta)\theta)$ . We have  $H^0(X, T \otimes L^{jq} \otimes \mathcal{I}(jqu)) \supseteq H^0(X, T \otimes L^{jq} \otimes \mathcal{I}(jqw_{\beta,k}))$ , hence

(17) 
$$h^{0}(X, T \otimes L^{jq} \otimes \mathcal{I}(jqu)) \geq h^{0}(X, T \otimes L^{jq} \otimes \mathcal{I}(jqw_{\beta,k})).$$

For each fixed k > 0, we can take a resolution of singularities  $\pi: Y \to X$  such that  $\pi^* u_k^D$  has neat analytic singularities along a normal crossing  $\mathbb{Q}$ -divisor, as described in (11). By [Demailly 2012, Proposition 5.8] and the projection formula,

(18) 
$$h^{0}(X, T \otimes L^{jq} \otimes \mathcal{I}(jqw_{\beta,k})) = h^{0}(Y, \pi^{*}T \otimes K_{Y/X} \otimes (\pi^{*}L)^{jq} \otimes \mathcal{I}(jq\pi^{*}w_{\beta,k})).$$

Since  $\int_Y (\pi^*\theta + dd^c \pi^* v_{\beta,k})^n = \int_X \theta_{v_{\beta,k}}^n > 0$ , there exists a nonzero section

$$s_j \in H^0(Y, \pi^* L^{\beta j q} \otimes \mathcal{I}(\beta j q \pi^* v_{\beta,k})) = H^0(Y, \pi^* L^{j p} \otimes \mathcal{I}(j p \pi^* v_{\beta,k}))$$

for all *j* large enough, by Lemma 4.4. Hence applying Lemma 4.5 for  $T := \pi^* T \otimes K_{Y/X}$ ,  $E_1 = \pi^* L^{q-p}$ ,  $E_2 = \pi^* L^p$ ,  $\chi_1 := q \pi^* u_k^D$ ,  $\chi_2 := p \pi^* v_{\beta,k}$ ,  $s_j := s_j$  and  $\varepsilon := \beta$ , we find

(19) 
$$h^{0}(Y, \pi^{*}T \otimes K_{Y/X} \otimes \pi^{*}L^{jq} \otimes \mathcal{I}(jq\pi^{*}w_{k,\beta}))$$
$$= h^{0}(Y, \pi^{*}T \otimes K_{Y/X} \otimes \pi^{*}L^{(q-p)j} \otimes \pi^{*}L^{pj} \otimes \mathcal{I}((1-\beta)jq\pi^{*}u_{k}^{D} + jp\pi^{*}v_{\beta,k}))$$
$$\geq h^{0}(Y, \pi^{*}T \otimes K_{Y/X} \otimes \pi^{*}L^{(q-p)j} \otimes \mathcal{I}(jq\pi^{*}u_{k}^{D}))$$

for j large enough (depending on k).

Geometry & Topology, Volume 28 (2024)

Since  $\theta_{u_k^D} > \beta \omega \ge \beta \theta$ , we notice that  $q u_k^D \in PSH(X, \theta(q-p))$ . Hence, by [Bonavero 1998, Théorème 2.1, Corollaire 2.2] (see also [Darvas and Xia 2022, Theorem 2.26]), we have the estimates

$$(20) \quad \lim_{j \to \infty} \frac{1}{j^n q^n} h^0(Y, \pi^* T \otimes K_{Y/X} \otimes \pi^* L^{(1-\beta)qj} \otimes \mathscr{G}(jq\pi^* u_k^D)) = \lim_{j \to \infty} \frac{1}{j^n q^n} h^0(Y, \pi^* T \otimes K_{Y/X} \otimes \pi^* L^{(q-p)j} \otimes \mathscr{G}(jq\pi^* u_k^D)) = \frac{r}{q^n n!} \int_Y ((q-p)\pi^* \theta + q \mathrm{dd}^c \pi^* u_k^D)^n = \frac{r}{n!} \int_X ((1-\beta)\theta + \mathrm{dd}^c u_k^D)^n \geq \frac{r}{n!} \int_X \theta_{u_k^D}^n - C\beta,$$

where C > 0 depends only on r, n and  $\theta$ . Putting together (17)–(20), we obtain

$$\lim_{j \to \infty} \frac{1}{j^n} h^0(X, T \otimes L^j \otimes \mathcal{I}(ju)) \ge \frac{r}{n!} \int_X \theta_{u_k}^n - C\beta.$$

Letting  $k \to \infty$  and applying Lemma 3.7, we conclude (16).

**Lemma 4.4** Suppose that  $L \to X$  is a big line bundle, with smooth Hermitian metric h. Let  $\theta = c_1(L, h)$ . Let  $v \in PSH(X, \theta)$  with  $\int_X \theta_v^n > 0$ . Then for m big enough there exists an  $s \in H^0(X, L^m \otimes \mathcal{I}(mv))$  which is nonvanishing.

**Proof** By Proposition 3.6 there exists  $w \in PSH(X, \theta)$  such that  $w \le v$  and  $\theta_w \ge \delta \omega$ . By [Demailly 2012, Theorem 13.21], for *m* big enough there exists an  $s \in H^0(X, L^m \otimes \mathscr{I}(mw))$  which is nonzero. Since  $w \le v$ , we get that  $s \in H^0(X, L^m \otimes \mathscr{I}(mv))$ .

**Lemma 4.5** Suppose that  $E_1$ ,  $E_2$  and T are vector bundles over a connected complex manifold Y, with rank  $E_2 = 1$ , and that  $\chi_1$  and  $\chi_2$  are quasi-psh functions on Y, with  $\chi_1$  having normal crossing divisorial singularity type. Suppose that there exists a nonzero section  $s_j \in H^0(Y, E_2^{\otimes j} \otimes \mathcal{I}(j\chi_2))$  for all j big enough. Then for any  $\varepsilon \in (0, 1)$ , the map  $w \mapsto w \otimes s_j$  between the vector spaces

$$H^{0}(Y, T \otimes E_{1}^{\otimes j} \otimes \mathcal{I}(j\chi_{1})) \to H^{0}(Y, T \otimes E_{1}^{\otimes j} \otimes E_{2}^{\otimes j} \otimes \mathcal{I}(j(1-\varepsilon)\chi_{1}+j\chi_{2}))$$

is well defined and injective for all *j* big enough.

**Proof** Suppose that the singularity type of  $\chi_1$  is given by the effective normal crossing  $\mathbb{R}$ -divisor  $\sum_j \alpha_j D_j$  with  $\alpha_j > 0$ . By [Demailly 2012, Remark 5.9] we have that

$$\mathscr{I}(j\chi_1) = \mathbb{O}_Y\left(-\sum_m \lfloor \alpha_m j \rfloor D_j\right).$$

We obtain that  $we^{-j(1-\varepsilon)\chi_1}$  is bounded for any  $w \in H^0(Y, T \otimes E_1^j \otimes \mathcal{I}(j\chi_1))$  and j big enough. Since  $s_j \in H^0(Y, E_2^{\otimes j} \otimes \mathcal{I}(j\chi_2))$ , we obtain that

$$w \otimes s_j \in H^0(Y, T \otimes E_1^{\otimes j} \otimes E_2^{\otimes j} \otimes \mathcal{I}(j(1-\varepsilon)\chi_1 + j\chi_2))$$

Injectivity of  $w \mapsto w \otimes s_j$  follows from the identity theorem of holomorphic functions.

Geometry & Topology, Volume 28 (2024)

Tamás Darvas and Mingchen Xia

**Theorem 4.6** Suppose that  $u \in PSH(X, \theta)$ . Then

(21) 
$$\lim_{k \to \infty} \frac{1}{k^n} h^0(X, T \otimes L^k \otimes \mathcal{I}(ku)) = \frac{r}{n!} \int_X \theta_{P[u]_{\mathcal{I}}}^n.$$

**Proof** Since both sides of (21) only depend on  $P[u]_{\mathcal{I}}$ , we can assume that  $P[u]_{\mathcal{I}} = u$ . Proposition 4.1 implies (21) for  $\int_X \theta_u^n = 0$ , so we can also assume that  $\int_X \theta_u^n > 0$ . In particular, *L* is a big line bundle and *X* is projective. By Proposition 3.6, there exists  $v \le u$  such that  $\theta_v$  is a Kähler current. Let  $v_t := (1-t)v + tu$ . Then  $\theta_{v_t}$  is a Kähler current for  $t \in [0, 1)$ , so we can apply Proposition 4.3 to obtain that

$$\lim_{k \to \infty} \frac{1}{k^n} h^0(X, T \otimes L^k \otimes \mathscr{I}(ku)) \ge \lim_{k \to \infty} \frac{1}{k^n} h^0(X, T \otimes L^k \otimes \mathscr{I}(kv_t)) \ge \frac{r}{n!} \int_X \theta_{P[v_t]_{\mathscr{I}}}^n.$$

Letting  $t \rightarrow 0$  and using [Darvas and Xia 2022, Lemma 2.21(iii)], we obtain that

$$\lim_{k \to \infty} \frac{1}{k^n} h^0(X, T \otimes L^k \otimes \mathcal{I}(ku)) \ge \frac{r}{n!} \int_X \theta_{P[u]_{\mathcal{I}}}^n$$

The reverse inequality follows from Proposition 4.1.

#### **4.2** The case of $\mathbb{R}$ -line bundles

In this subsection we extend Theorem 4.6 to  $\mathbb{R}$ -line bundles. First we deal with the case of  $\mathbb{Q}$ -line bundles.

**Corollary 4.7** Let *L* be a pseudoeffective  $\mathbb{Q}$ -line bundle on *X*, represented by an effective  $\mathbb{Q}$ -divisor *D*. Let  $\theta$  be a smooth form representative of  $c_1(L)$ . Let  $u \in PSH(X, \theta)$ . Then

(22) 
$$\lim_{k \to \infty} \frac{1}{k^n} h^0(X, T \otimes \mathbb{O}_X(\lfloor kD \rfloor) \otimes \mathscr{I}(ku)) = \frac{r}{n!} \int_X \theta_{P[u]_{\mathscr{I}}}^n.$$

**Proof** We may assume that D' := aD is a line bundle L' for some  $a \in \mathbb{N}$ . For each  $k \in \mathbb{N}$ , write

 $k = k_0 a + k'$ , where  $k_0 \in \mathbb{N}, k' \in [0, a - 1)$ .

Note that the difference  $\lfloor kD \rfloor - k_0D'$  can represent only a finite number of different line bundles. Hence, in order to prove (22), it suffices to establish the following: for each fixed  $k' \in [0, a-1)$ ,

$$\lim_{k_0\to\infty}\frac{1}{k_0^n a^n}h^0(X,T\otimes L'^{k_0}\otimes \mathcal{I}(k_0au+k'u))=\frac{r}{n!}\int_X\theta_{P[u]_{\mathcal{I}}}^n.$$

Observe that  $\mathcal{I}(k_0au + k'u) \subseteq \mathcal{I}(k_0au)$ , so by Theorem 4.6 we have

$$\lim_{k_0\to\infty}\frac{1}{k_0^n a^n}h^0(X,T\otimes L'^{k_0}\otimes \mathcal{I}(k_0au+k'u))\leq \frac{r}{n!}\int_X\theta_{P[u]_{\mathcal{I}}}^n.$$

Geometry & Topology, Volume 28 (2024)

1976

On the other hand, as  $\mathcal{I}(k_0au + k'u) \supseteq \mathcal{I}((k_0 + 1)au)$ , we get

$$\begin{split} \lim_{k_0 \to \infty} \frac{1}{k_0^n a^n} h^0(X, T \otimes L'^{k_0} \otimes \mathcal{I}(k_0 a u + k' u)) \\ &\geq \underline{\lim} \frac{1}{k_0^n a^n} h^0(X, T \otimes L'^{k_0} \otimes \mathcal{I}((k_0 + 1) a u)) \\ &= \underline{\lim} \frac{1}{k_0 \to \infty} \frac{1}{((k_0 + 1)a)^n} h^0(X, T \otimes L'^* \otimes L'^{k_0 + 1} \otimes \mathcal{I}((k_0 + 1) a u)) \\ &= \frac{r}{n!} \int_X \theta_{P[u]_{\mathcal{I}}}^n, \end{split}$$

where in the last step we again used Theorem 4.6, finishing the proof.

**Corollary 4.8** Assume that X is projective. Let D be a big  $\mathbb{R}$ -divisor on X. Let  $\theta$  be a smooth form representing the cohomology class [D]. Let  $u \in PSH(X, \theta)$ . Then

(23) 
$$\lim_{k \to \infty} \frac{1}{k^n} h^0(X, T \otimes \mathbb{O}_X(\lfloor kD \rfloor) \otimes \mathscr{I}(ku)) = \frac{r}{n!} \int_X \theta_{P[u]_{\mathscr{I}}}^n.$$

**Proof** We first deal with the  $\leq$  direction in (23). Fix  $\delta > 0$ . Fix  $\varepsilon > 0$ , so that

$$\int_X (\theta + \varepsilon \omega + \mathrm{dd}^{\mathsf{c}} P^{\theta + \varepsilon \omega} [u]_{\mathscr{I}})^n < \int_X \theta_{P[u]_{\mathscr{I}}}^n + \delta.$$

This is possible by Corollary 3.5(i). Take a  $\mathbb{Q}$ -divisor  $D^{\delta}$  such that the cohomology class  $\{D^{\delta} - D\}$  has a smooth positive representative  $\theta^{\delta} \leq \varepsilon \omega$ . As a result,  $D^{\delta} - D$  is ample. This is possible as X is projective. We have  $u \in PSH(X, \theta + \theta^{\delta})$ . Then  $\mathbb{O}_X(\lfloor kD^{\delta} \rfloor - \lfloor kD \rfloor)$  has a nonzero global section s for k big enough. As a result, the map  $H^0(X, T \otimes \mathbb{O}_X(\lfloor kD \rfloor) \otimes \mathcal{I}(ku)) \to H^0(X, T \otimes \mathbb{O}_X(\lfloor kD^{\delta} \rfloor) \otimes \mathcal{I}(ku))$  given by  $s' \mapsto s' \otimes s$  is injective, allowing us to write the estimates

$$\begin{split} \overline{\lim}_{k \to \infty} \frac{1}{k^n} h^0(X, T \otimes \mathbb{O}_X(\lfloor kD \rfloor) \otimes \mathscr{I}(ku)) &\leq \overline{\lim}_{k \to \infty} \frac{1}{k^n} h^0(X, T \otimes \mathbb{O}_X(\lfloor kD^\delta \rfloor) \otimes \mathscr{I}(ku)) \\ &= \frac{r}{n!} \int_X (\theta + \theta^\delta + \mathrm{dd^c} \, P^{\theta + \theta^\delta}[u]_{\mathscr{I}})^n \\ &\leq \frac{r}{n!} \int_X (\theta + \varepsilon \omega + \mathrm{dd^c} \, P^{\theta + \varepsilon \omega}[u]_{\mathscr{I}})^n \\ &\leq \frac{r}{n!} \int_X \theta^n_{P[u]_{\mathscr{I}}} + \frac{r\delta}{n!}, \end{split}$$

where in the second line we have used Theorem 4.6. Letting  $\delta \to 0+$ , we conclude the  $\leq$  direction in (23). For the reverse direction, we can replace u by  $P[u]_{\mathcal{I}}$ , as in the proof of Theorem 4.6. Hence, we can assume that u is  $\mathcal{I}$ -model. If  $\int_X \theta_u^n = 0$ , we are done by the previous arguments, so we can assume that  $\int_X \theta_u^n > 0$ . We first treat the case where  $\theta_u > \varepsilon_0 \omega$  for some  $\varepsilon_0 > 0$ . Fix  $\delta > 0$ . Fix  $\varepsilon \in (0, \varepsilon_0)$ , so that

$$\int_{X} (\theta - \varepsilon \omega + \mathrm{dd}^{\mathrm{c}} P^{\theta - \varepsilon \omega} [u]_{\mathscr{I}})^{n} > \int_{X} \theta^{n}_{P[u]_{\mathscr{I}}} - \delta.$$

Geometry & Topology, Volume 28 (2024)

This is possible by Corollary 3.5(ii). Take a  $\mathbb{Q}$ -divisor  $D^{\delta}$  so that  $\{D - D^{\delta}\}$  has a smooth positive representative  $\theta^{\delta} \leq \varepsilon \omega$ . As a result,  $D - D^{\delta}$  is ample. Then we have  $u \in PSH(X, \theta - \theta^{\delta})$ . As before, we have the estimates

$$\begin{split} \lim_{k \to \infty} \frac{1}{k^n} h^0(X, T \otimes \mathbb{O}_X(\lfloor kD \rfloor) \otimes \mathscr{I}(ku)) &\geq \lim_{k \to \infty} \frac{1}{k^n} h^0(X, T \otimes \mathbb{O}_X(\lfloor kD^\delta \rfloor) \otimes \mathscr{I}(ku)) \\ &= \frac{r}{n!} \int_X (\theta - \theta^\delta + \mathrm{dd^c} P^{\theta - \theta^\delta}[u]_{\mathscr{I}})^n \\ &\geq \frac{r}{n!} \int_X (\theta - \varepsilon \omega + \mathrm{dd^c} P^{\theta - \varepsilon \omega}[u]_{\mathscr{I}})^n \\ &\geq \frac{r}{n!} \int_X \theta_{P[u]_{\mathscr{I}}}^n - \frac{r\delta}{n!}, \end{split}$$

where in the second line we have used Theorem 4.6. Letting  $\delta \rightarrow 0+$ , we conclude (23) in this case.

Finally, we treat the general case. By Proposition 3.6, there exists  $v \in PSH(X, \theta)$ , such that  $v \le u$  and  $\theta_v$  is a Kähler current. Set  $u_t := (1-t)u + tv$  for  $t \in [0, 1]$ . For  $t \in (0, 1]$ ,  $\theta_{u_t}$  is still a Kähler current. By the special case treated above, we get

(24) 
$$\lim_{k \to \infty} \frac{1}{k^n} h^0(X, T \otimes \mathbb{O}_X(\lfloor kD \rfloor) \otimes \mathscr{I}(ku)) \ge \lim_{k \to \infty} \frac{1}{k^n} h^0(X, T \otimes \mathbb{O}_X(\lfloor kD \rfloor) \otimes \mathscr{I}(ku_t))$$
$$= \frac{r}{n!} \int_X \theta_{P[u_t]_{\mathscr{I}}}^n$$

for  $t \in (0, 1]$ . As  $t \searrow 0$  we have  $u_t \nearrow u$ , hence  $P[u_t]_{\mathscr{I}} \nearrow P[u]_{\mathscr{I}}$  a.e. by Proposition 2.7(ii). By [Darvas et al. 2018, Theorem 2.3],  $\int_X \theta_{P[u_t]_{\mathscr{I}}}^n \nearrow \int_X \theta_{P[u]_{\mathscr{I}}}^n$ . Letting  $t \searrow 0$  in (24), we find the desired inequality

$$\lim_{k \to \infty} \frac{1}{k^n} h^0(X, T \otimes \mathbb{O}_X(\lfloor kD \rfloor) \otimes \mathscr{I}(ku)) \ge \frac{r}{n!} \int_X \theta_{P[u]_{\mathscr{I}}}^n.$$

# 5 Envelopes of singularity types with respect to compact sets

Let *X* be a connected compact Kähler manifold of dimension *n*. For this whole section, let  $K \subseteq X$  be a closed nonpluripolar set. Let  $\theta$  be a closed real (1, 1)–form on *X* representing a pseudoeffective cohomology class. Let  $u \in PSH(X, \theta)$ .

Let  $v: K \to [-\infty, \infty)$  be a function. We introduce the following *K*-relative envelopes and their regularizations, refining the definitions from Section 2.1:

$$\begin{split} E_{K}^{\theta}[u]_{\mathscr{I}}(v) &:= \sup\{h \in \mathrm{PSH}(X,\theta) : h|_{K} \leq v \text{ and } [h] \preceq_{\mathscr{I}} [u]\}, \quad P_{K}^{\theta}[u]_{\mathscr{I}}(v) := \mathrm{usc}(E_{K}^{\theta}[u]_{\mathscr{I}}(v)), \\ E_{K}^{\theta}[u](v) &:= \sup\{h \in \mathrm{PSH}(X,\theta) : h|_{K} \leq v \text{ and } [h] \preceq [u]\}, \qquad P_{K}^{\theta}[u](v) := \mathrm{usc}(E_{K}^{\theta}[u](v)). \end{split}$$

We omit  $\theta$  and K from the notation when there is no risk of confusion. When v is bounded, neither of the above candidate sets are empty: one can always take h = u - C for a large enough constant C.

We note the following maximum principles, that follow from the above definitions:

**Lemma 5.1** Let  $v \in C^0(K)$ . Let  $h \in PSH(X, \theta)$ . Assume that  $[h] \leq [u]$ , then

(25) 
$$\sup_{K} (h-v) = \sup_{X \setminus \{h=-\infty\}} (h-E_K[u](v)) = \sup_{X \setminus \{E_K[u](v)=-\infty\}} (h-E_K[u](v))$$

**Proof** We prove the first equality first. We write  $S = \{h = -\infty\}$ .

By definition,  $E_K[u](v)|_K \leq v$ , so

$$(h - E_K[u](v))|_{K \setminus S} \ge h|_{K \setminus S} - v|_{K \setminus S}.$$

This implies that  $\sup_{K}(h-v) \leq \sup_{X \setminus S}(h-E_{K}[u](v))$ .

Conversely, observe that  $\sup_K (h-v) > -\infty$  as K is nonpluripolar. Let  $h' := h - \sup_K (h-v)$ . Then h' is a candidate in the definition of  $E_K[u](v)$ , hence  $h' \leq E_K[u](v)$ , namely

$$h - \sup_{K} (h - v) \le E_{K}[u](v),$$

the latter implies that  $\sup_{K}(h-v) \ge \sup_{X \setminus S}(h-E_{K}[u](v))$ , finishing the proof of the first identity.

We have  $\{E_K[u](v) = -\infty\} \subseteq S$ , and we notice that points in  $S \setminus \{E_K[u](v) = -\infty\}$  do not contribute to the supremum of  $h - E_K[u](v)$  on  $X \setminus \{E_K[u](v) = -\infty\}$ , hence the last equality of (25) also follows.  $\Box$ 

Next we make the following observations about the singularity types of our envelopes:

**Lemma 5.2** For any  $v \in C^0(K)$  we have  $[P_K[u](v)] = [P[u]]$  and  $[P_K[u]_{\mathcal{F}}(v)] = [P[u]_{\mathcal{F}}]$ . In particular, if  $[u] \in \mathcal{A}(X, \theta)$ , then  $P_K[u](v) = P_K[u]_{\mathcal{F}}(v)$ .

**Proof** Let C > 0 such that  $-C \le v \le C$ . Then  $P[u] - C \le P_K[u](v)$ . Since K is nonpluripolar, for  $h \in PSH(X, \theta)$  the condition  $h|_K \le v \le C$  implies that  $h \le \tilde{C}$  on X for some  $\tilde{C} := \tilde{C}(C, K) > 0$ ; see [Guedj and Zeriahi 2007, Corollary 4.3]. This implies that  $P_K[u](v) \le P[u] + \tilde{C}$ , giving  $[P_K[u](v)] = [P[u]]$ . The exact same argument applies in case of the  $P[\cdot]_{\mathscr{I}}$  envelope as well. Finally, when  $[u] \in \mathscr{A}(X, \theta)$ , we have that  $[u] = [P[u]_{\mathscr{I}}] = [P[u]]$  (Lemma 3.2). We claim that for  $h \in PSH(X, \theta)$ ,  $[h] \le [u]$  if and only if  $[h] \le_{\mathscr{I}} [u]$ . This claim immediately gives  $P_K[u](v) = P_K[u]_{\mathscr{I}}(v)$ . The forward direction of the claim is trivial, so suppose  $[h] \le_{\mathscr{I}} [u]$ . We then have  $P[h]_{\mathscr{I}} \le P[u]_{\mathscr{I}}$ . This implies that  $[h] \le [P[h]_{\mathscr{I}}] \le [P[u]_{\mathscr{I}}] = [u]$ .  $\Box$ 

**Corollary 5.3** Let  $u \in \text{PSH}(X, \theta)$  and  $v \in C^0(X)$ . Then  $P_K[u]_{\mathscr{I}}(v) = P_K[P_K[u]_{\mathscr{I}}(v)]_{\mathscr{I}}(v)$ .

**Proof** By definition, the right-hand side is the usc regularization of

$$\sup\{h \in \mathsf{PSH}(X,\theta) : h|_K \le v, [h] \preceq_{\mathscr{I}} P_K[u]_{\mathscr{I}}(v)\}.$$

By Lemma 5.2 and [Darvas and Xia 2022, Proposition 2.18(ii)], this expression can be rewritten as

$$\sup\{h \in \mathsf{PSH}(X,\theta) : h|_K \le v, [h] \preceq_{\mathscr{I}} [u]\}.$$

The usc regularization of the latter expression is just  $P_K[u]_{\mathcal{I}}(v)$ .

Geometry & Topology, Volume 28 (2024)

**Lemma 5.4** Let  $u \in PSH(X, \theta)$  be a potential with positive mass. Let  $v \in C^0(K)$ . Let  $S \subseteq X$  be a pluripolar set. Let  $h \in PSH(X, \theta)$  satisfy  $[h] \leq [u]$ . Assume that h has positive mass and  $h|_{K \setminus S} \leq v|_{K \setminus S}$ . Then  $h \leq P_K[u](v)$ .

**Proof** By the global Josefson theorem [Guedj and Zeriahi 2005, Theorem 7.2], there is  $\chi \in PSH(X, \theta)$  such that  $S \subseteq \{\chi = -\infty\}$ . We claim that we can choose  $\chi$  so that  $\chi \leq h$ . In fact, since  $\int_X \theta_h^n > 0$ , fixing some  $\chi$  and  $\varepsilon > 0$  small enough, we have

$$\int_X \theta_{\varepsilon\chi+(1-\varepsilon)V_{\theta}}^n + \int_X \theta_h^n > \int_X \theta_{V_{\theta}}^n.$$

Thus, by [Darvas et al. 2021, Lemma 5.1], we have  $P(\varepsilon \chi + (1 - \varepsilon)V_{\theta}, h) \in PSH(X, \theta)$ . Since we have  $P(\varepsilon \chi + (1 - \varepsilon)V_{\theta}, h) \leq \varepsilon \chi$ , the claim is proved by replacing  $\chi$  with  $P(\varepsilon \chi + (1 - \varepsilon)V_{\theta}, h)$ .

Fix  $\chi \leq h$  as above. For any  $\delta \in (0, 1)$ , we have

$$(1-\delta)h|_{K} + \delta\chi|_{K} \le v$$
 and  $[(1-\delta)h + \delta\chi] \le [u].$ 

Hence,  $(1 - \delta)h + \delta \chi \leq P_K[u](v)$ . Letting  $\delta \searrow 0$ , we conclude that  $h \leq P_K[u](v)$ .

**Corollary 5.5** Let  $u \in PSH(X, \theta)$  be a potential with positive mass. Let  $v \in C^{0}(K)$ . Then

$$P_K[u](v) = P_X[u](P_K[V_\theta](v)).$$

**Proof** It is clear that  $P_K[u](v) \leq P_X[u](P_K[V_\theta](v))$ . For the reverse direction, it suffices to prove that any  $h \in PSH(X, \theta)$  such that  $[h] \leq [u], h \leq P_K[V_\theta](v)$  satisfies  $h \leq P_K[u](v)$ . As *u* has positive mass, we can assume that *h* has positive mass as well. Let  $S = \{P_K[V_\theta](v) > E_K[V_\theta](v)\}$ . By [Bedford and Taylor 1982, Theorem 7.1], *S* is a pluripolar set. Observe that  $h|_{K\setminus S} \leq v|_{K\setminus S}$ , hence by Lemma 5.4,  $h \leq P_K[u](v)$  and we conclude.

The next result motivates our terminology to call the measures  $\theta_{P_K[u](v)}^n$  the partial equilibrium measures of our context.

**Lemma 5.6** Let  $v \in C^0(K)$ . Let  $u \in PSH(X, \theta)$ . Then  $\theta_{P_K[u](v)}^n$  does not charge  $X \setminus K$ . Moreover,  $P_K[u](v)|_K = v$  a.e. with respect to  $\theta_{P_K[u](v)}^n$ . More precisely, we have

(26) 
$$\theta_{P_K[u](v)}^n \leq \mathbb{1}_{K \cap \{P_K[u](v) = P_K[V_\theta](v) = v\}} \theta_{P_K[V_\theta](v)}^n$$

**Proof** First we address the case when  $u = V_{\theta}$ .

Let  $S \subseteq X$  be a closed pluripolar set such that  $V_{\theta}$  is locally bounded on  $X \setminus S$ .

For the first assertion, it suffices to show that  $\theta_{P_K[V_\theta](v)}^n$  does not charge any open ball  $B \in X \setminus (S \cup K)$ . By Choquet's lemma, we can take an increasing sequence  $h_j \in \text{PSH}(X, \theta)$  converging to  $P_K[V_{\min}](v)$ a.e. and  $h_j|_K \leq v$ . By [Bedford and Taylor 1982, Proposition 9.1], we can find  $w_j \in \text{PSH}(X, \theta)$  such that  $(\theta + \text{dd}^c w_j|_B)^n = 0$  and  $w_j$  agrees with  $h_j$  outside B. Note that  $w_j$  is clearly increasing and  $w_j \geq h_j$ , along with  $w_j|_K \leq v$ . It follows that  $w_j$  converges to  $P_K[V_\theta](v)$  as well. By continuity of the

Monge–Ampère operator along increasing bounded sequences [Darvas et al. 2018, Theorem 2.3], we find that  $\theta_{P_K[V_a](v)}^n$  does not charge *B*, as desired.

For the second assertion, let  $x \in (X \setminus S) \cap K$  be a point such that  $P_K[V_\theta](v)(x) < v(x) - \varepsilon$  for some  $\varepsilon > 0$ . Let *B* be a ball centered at *x*, small enough so that  $\theta$  has a local potential on *B*, allowing us to identify  $\theta$ -psh functions with psh functions (on *B*). By shrinking *B*, we can further guarantee

(i) 
$$\overline{B} \subseteq X \setminus S$$
,

(ii) 
$$P_K[V_{\theta}](v)|_{\overline{B}} < v(x) - \varepsilon$$
,

(iii) 
$$v|_{\overline{B}\cap K} > v(x) - \varepsilon$$
.

Construct the sequences  $h_j$  and  $w_j$  as above. On B, by choosing a local potential of  $\theta$ , we may identify  $h_j$  and  $w_j$  with the corresponding psh functions in a neighborhood of  $\overline{B}$ . By 2. we have  $w_j \leq v(x) - \varepsilon$  on  $\partial B$ , hence by the comparison principle,  $w_j|_B \leq v(x) - \varepsilon$ . By (3) we have  $w_j|_{B\cap K} \leq v|_{B\cap K}$ . Thus, we conclude that  $\theta_{P_K[V_{\theta}](v)}^n$  does not charge B, as in the previous paragraph.

For the general case, we can assume  $\int_X \theta_u^n > 0$ . Indeed, due to Lemma 5.2, we have  $\int_X \theta_{P_K[u](v)}^n = \int_X \theta_u^n$ , hence there is nothing to prove if  $\int_X \theta_u^n = 0$ . By Corollary 5.5,  $P_K[u](v) = P_X[u](P_K[V_\theta](v))$ . Now [Darvas et al. 2018, Theorem 3.8] gives

$$\theta_{P_{K}[u](v)}^{n} \leq \mathbb{1}_{\{P_{K}[u](v)=P_{K}[V_{\theta}](v)\}} \theta_{P_{K}[V_{\theta}](v)}^{n} \leq \mathbb{1}_{\{P_{K}[u](v)=v\}} \theta_{P_{K}[V_{\theta}](v)}^{n}$$

where in the last inequality we have used the first part of the argument.

**Corollary 5.7** Let  $v \in C^0(K)$ . Let  $u \in PSH(X, \theta)$ . Then  $\theta_{P_K[u](v)}^n$  (resp.  $\theta_{P_K[u]_{\mathscr{I}}(v)}^n$ ) does not charge  $(X \setminus K) \cup \{P_K[u](v) < v\}$  (resp.  $(X \setminus K) \cup \{P_K[u]_{\mathscr{I}}(v) < v\}$ ).

**Proof** The first part of the corollary follows from Lemma 5.6. For the second part, we can assume that  $\int_X \theta_{P_K[u]_{\mathfrak{f}}(v)}^n > 0$ , otherwise there is nothing to prove. By definition, we have  $P_K[u]_{\mathfrak{f}}(v) = P_K[P[u]_{\mathfrak{f}}]_{\mathfrak{f}}(v)$ . Next we show that  $P_K[P[u]_{\mathfrak{f}}]_{\mathfrak{f}}(v) = P_K[P[u]_{\mathfrak{f}}](v)$ . The inequality  $P_K[P[u]_{\mathfrak{f}}]_{\mathfrak{f}}(v) \ge P_K[P[u]_{\mathfrak{f}}](v)$  is trivial. By Lemma 5.2 we get that  $[P_K[P[u]_{\mathfrak{f}}]_{\mathfrak{f}}(v)] = [P[u]_{\mathfrak{f}}]$ . Due to Choquet's lemma, we get that  $P_K[P[u]_{\mathfrak{f}}]_{\mathfrak{f}}(v) \le v$  on  $K \setminus S$ , where S is pluripolar. As a result, due to the nonvanishing mass assumption, Lemma 5.4 allows to conclude that  $P_K[P[u]_{\mathfrak{f}}]_{\mathfrak{f}}(v) \le P_K[P[u]_{\mathfrak{f}}]_{\mathfrak{f}}(v)$ .

Since  $P_K[P[u]_{\mathcal{J}}]_{\mathcal{J}}(v) = P_K[u]_{\mathcal{J}}(v)$ , we get that  $\theta_{P_K[u]_{\mathcal{J}}(v)}^n$  does not charge  $(X \setminus K) \cup \{P_K[u]_{\mathcal{J}}(v) < v\}$ , using the first part of the corollary.

**Proposition 5.8** Let  $u \in PSH(X, \theta)$  be a potential with positive mass. Let  $v \in C^0(K)$ . Then

(27) 
$$P_{K}[u](v) = P_{K}[P[u]](v).$$

In particular,  $P_K[u](v) = P_K[P_K[u](v)](v)$ .

**Proof** It is obvious that  $P_K[u](v) \le P_K[P[u]](v)$ . We to prove the reverse inequality. As  $P_K[u](v)$  and  $P_K[P[u]](v)$  have the same singularity types (Lemma 5.2), by the domination principle [Darvas et al. 2018, Corollary 3.10], it suffices to show that  $P_K[u](v) \ge P_K[P[u]](v)$  a.e. with respect to  $\theta_{P_K[u](v)}^n$ .

By (26),  $P_K[u](v) = P_K[V_\theta](v) = v$  a.e. with respect to  $\theta_{P_K[u](v)}^n$ . Hence,  $P_K[P[u]](v) = v$  a.e. with respect to  $\theta_{P_K[u](v)}^n$ . We conclude that  $P_K[u](v) = P_K[P[u]](v)$ .

Finally, that  $P_K[u](v) = P_K[P_K[u](v)](v)$  follows from Lemma 5.2 and (27).

**Lemma 5.9** Fix a Kähler form  $\omega$  on X. For  $v \in C^0(K)$  there exists an increasing bounded sequence  $\{v_j^-\}_j$  in  $C^\infty(X)$  and a decreasing bounded sequence  $\{v_j^+\}_j$  in  $C^\infty(X)$  such that for all  $u \in PSH(X, \theta)$  with  $\int_X \theta_u^n > 0$ , and  $\delta \in [0, 1]$ , we have

- (i)  $P_X^{\theta+\delta\omega}[u](v_j^+) \searrow P_K^{\theta+\delta\omega}[u](v),$
- (ii)  $P_X^{\theta+\delta\omega}[u](v_i^-) \nearrow P_K^{\theta+\delta\omega}[u](v)$  a.e.,
- (iii)  $\sup_X |v_j^-| \le C(\|v\|_{C^0(K)}, K, \theta + \omega) \text{ and } \sup_X |v_j^+| \le C(\|v\|_{C^0(K)}, K, \theta + \omega).$

**Proof** We fix  $\delta \in [0, 1]$ . First we prove the existence of  $\{v_i^-\}_j$ . Let

$$C_{K,v} := \sup \{ \sup_{X} w : w \in \mathsf{PSH}(X, \theta + \omega), w |_{K} \le v \}$$

Since *K* is nonpluripolar, we have that  $C_{K,v} \in \mathbb{R}$ . Now let  $\tilde{v}: X \to \mathbb{R}$  so that  $\tilde{v}|_K = v$  and  $\tilde{v}|_{X \setminus K} = C_{K,v} + 1$ . Since  $\tilde{v}$  is lsc, there exists an increasing and uniformly bounded sequence  $\{v_j^-\}_j$  in  $C^{\infty}(X)$  such that  $v_j^- \nearrow \tilde{v}$ .

Observe that  $P_X^{\theta+\delta\omega}[u](v_j^-)$  is increasing in j, and that  $P_X^{\theta+\delta\omega}[u](v_j^-) \leq P_K^{\theta+\delta\omega}[u](v)$ . To prove that  $P_X^{\theta+\delta\omega}[u](v_j^-) \nearrow P_K^{\theta+\delta\omega}[u](v)$  a.e., let w be a candidate for  $P_K^{\theta+\delta\omega}[u](v)$  such that  $\sup_K (w-v) < 0$ . Then, since w is use and  $w < \tilde{v}$ , by Dini's lemma there exists  $j_0$  such that  $w < v_j^-$  for  $j \ge j_0$ , ie  $w \le P_X^{\theta+\delta\omega}[u](v_j^-)$ , proving existence of  $\{v_j^-\}_j$ .

Next we prove the existence of  $\{v_j^+\}_j$ . Since  $h := \max(P_K^{\theta+\omega}[V_{\theta+\omega}](v), \inf_K v - 1)$  is usc, there exists a decreasing and uniformly bounded sequence  $\{v_j^+\}_j$  in  $C^{\infty}(X)$  such that  $v_j^+ \searrow h$ . Trivially,  $\chi := \lim_{j \to \infty} P_X^{\theta+\delta\omega}[u](v_j^+) \ge P_K^{\theta+\delta\omega}[u](v)$ . In particular,  $\chi$  has positive mass, since it has the same singularity types as  $P_K^{\theta+\delta\omega}[u](v)$  (Lemma 5.2). We introduce

$$S := \{ E_K^{\theta + \omega} [V_{\theta + \omega}](v) < P_K^{\theta + \omega} [V_{\theta + \omega}](v) \}.$$

By [Bedford and Taylor 1982, Theorem 7.1], *S* is a pluripolar set. Observe that  $P_X^{\theta+\delta\omega}[u](v_j^+) \le v_j^+$  for all *j*. Thus,  $\chi \le h$ . On the other hand,  $h \le v$  on  $K \setminus S$ . So in particular,  $\chi|_{K \setminus S} \le v|_{K \setminus S}$ . By Lemma 5.2 we also have that  $[\chi] = [P_K^{\theta+\delta\omega}[u](v)]$ . Hence, by Lemma 5.4,  $\chi \le P_K^{\theta+\delta\omega}[P_K^{\theta+\delta\omega}[u](v)](v) = P_K^{\theta+\delta\omega}[u](v)$ , where we also used the last statement of Proposition 5.8.

We recall the relative Monge–Ampère energy  $I_{[u]}^{\theta}$ :  $\mathscr{E}^1(X, \theta; P[u]) \to \mathbb{R}$  from [Darvas et al. 2018]:

(28) 
$$I_{[u]}^{\theta}(\varphi) := \frac{1}{n+1} \sum_{i=0}^{n} \int_{X} (\varphi - P[u]) \, \theta_{\varphi}^{i} \wedge \theta_{P[u]}^{n-i}$$

Using integration by parts (see [Xia 2019; Lu 2021, Theorem 1.2; Vu 2022, Theorem 2.6], and compare with [Boucksom et al. 2010, Theorem 1.14]), the argument of Berman and Boucksom [2010, Corollary 4.2] can be reproduced line by line to yield the following cocycle property: for  $\varphi_1, \varphi_2 \in \mathscr{E}^1(X, \theta; P[u])$  such that  $[\varphi_i] = [P[u]]$  for i = 1, 2, we have

(29) 
$$I_{[u]}^{\theta}(\varphi_1) - I_{[u]}^{\theta}(\varphi_2) = \frac{1}{n+1} \sum_{i=0}^n \int_X (\varphi_1 - \varphi_2) \, \theta_{\varphi_1}^i \wedge \theta_{\varphi_2}^{n-i}.$$

Following Berman and Boucksom [2010], we define the *partial equilibrium energy*  $\mathscr{I}^{\theta}_{[u],K}$  of  $v \in C^{0}(K)$ :

(30) 
$$\mathscr{I}^{\theta}_{[u],K}(v) := I^{\theta}_{[u]}(P_K[u](v)).$$

Berman and Boucksom [2010] used the symbol & for the above quantity. Due to potential confusion with the notation for (relative) full mass classes (that also uses the symbol  $\mathscr{E}$ ), we use the symbol  $\mathscr{I}$  instead.

Next we extend [Berman and Boucksom 2010, Theorem B] using the arguments of [Darvas 2019, Proposition 4.32], itself reproducing ideas from [Lu and Nguyen 2015]:

**Proposition 5.10** Let  $K \subseteq X$  be a closed nonpluripolar set, consider  $v, f \in C^0(K)$  and  $u \in PSH(X, \theta)$ satisfying  $\int_X \theta_u^n > 0$ . Then  $t \mapsto \mathcal{I}_{[u],K}^{\theta}(v+tf)$  for  $t \in \mathbb{R}$  is differentiable and

(31) 
$$\frac{\mathrm{d}}{\mathrm{d}t}\mathscr{I}_{[u],K}^{\theta}(v+tf) = \int_{K} f \,\theta_{P_{K}[u](v+tf)}^{n}$$

In this work, we will only need this result in the case that  $[u] \in \mathcal{A}(X, \theta)$ .

**Proof** It suffices to prove (31) at t = 0, which is equivalent to

(32) 
$$\lim_{t \to 0} \frac{I_{[u]}^{\theta}(P_K[u](v+tf)) - I_{[u]}^{\theta}(P_K[u](v))}{t} = \int_K f \,\theta_{P_K[u](v)}^n.$$

By switching f to -f, we may assume that t > 0 in the above limit.

By the comparison principle [Darvas et al. 2018, Proposition 3.5] and (29), we find that

$$\begin{aligned} \mathscr{I}^{\theta}_{[u],K}(v+tf) - \mathscr{I}^{\theta}_{[u],K}(v) &= \frac{1}{n+1} \sum_{i=0}^{n} \int_{X} \left( P_{K}[u](v+tf) - P_{K}[u](v) \right) \theta^{i}_{P_{K}[u](v+tf)} \wedge \theta^{n-i}_{P_{K}[u](v)} \\ &\leq \int_{X} \left( P_{K}[u](v+tf) - P_{K}[u](v) \right) \theta^{n}_{P_{K}[u](v)}. \end{aligned}$$
Lemma 5.6.

By Lemma 5.6,

$$\int_X (P_K[u](v+tf) - P_K[u](v)) \,\theta_{P_K[u](v)}^n \le t \int_K f \,\theta_{P_K[u](v)}^n$$

Thus, we get the inequality

$$\lim_{t \to 0+} \frac{I_{[u]}^{\theta}(P_K[u](v+tf)) - I_{[u]}^{\theta}(P_K[u](v))}{t} \le \int_K f \ \theta_{P_K[u](v)}^n$$

Similarly, we have

$$I_{[u]}^{\theta}(P_K[u](v+tf)) - I_{[u]}^{\theta}(P_K[u](v)) \ge \int_X (P_K[u](v+tf) - P_K[u](v)) \theta_{P_K[u](v+tf)}^n$$
$$\ge t \int_K f \theta_{P_K[u](v+tf)}^n.$$

Together with the above, this implies (32).

In the next lemma, we prove convergence results for the partial equilibrium energy:

**Lemma 5.11** Let  $v \in C^0(K)$  and  $u \in PSH(X, \theta)$  with  $\int_X \theta_u^n > 0$ . Let  $v_i^-$  and  $v_i^+$  be the sequences constructed in Lemma 5.9. Then

$$\lim_{j\to\infty} \mathscr{I}^{\theta}_{[u],X}(v_j^-) = \mathscr{I}^{\theta}_{[u],K}(v) \quad and \quad \lim_{j\to\infty} \mathscr{I}^{\theta}_{[u],X}(v_j^+) = \mathscr{I}^{\theta}_{[u],K}(v).$$

**Proof** This follows from Lemmas 5.2 and 5.9, and [Darvas et al. 2018, Theorem 2.3].

#### **Quantization of partial equilibrium measures** 6

In this section, we give a proof for Theorem 1.2. Throughout the section,  $L \to X$  is a pseudoeffective line bundle and h is a Hermitian metric on L such that  $\theta := c_1(L, h)$ . Let  $T \to X$  be a Hermitian line bundle on X with a smooth Hermitian metric  $h_T$ . We normalize the Kähler metric  $\omega$  on X so that

$$\int_X \omega^n = 1.$$

#### 6.1 Bernstein–Markov measures

Let  $K \subseteq X$  be a closed nonpluripolar subset. Let v be a measurable function on K and let v be a positive Borel probability measure on K. We introduce the following functions on  $H^0(X, L^k \otimes T)$ , with values possibly equaling  $\infty$ :

$$N_{v,v}^{k}(s) := \left( \int_{K} h^{k} \otimes h_{T}(s,s) e^{-kv} \, \mathrm{d}v \right)^{1/2} \quad \text{and} \quad N_{v,K}^{k}(s) := \sup_{K \setminus \{v = -\infty\}} (h^{k} \otimes h_{T}(s,s) e^{-kv})^{1/2}.$$

We start with the following elementary observation.

**Lemma 6.1** Let  $v_1 \le v_2$  be two measurable functions on *X*. Assume that  $\{v_1 = -\infty\} = \{v_2 = -\infty\}$ . Then for any  $s \in H^0(X, L^k \otimes T)$  and any k > 0, we have

$$N_{\boldsymbol{v}_1,\boldsymbol{K}}^{\boldsymbol{k}}(s) \ge N_{\boldsymbol{v}_2,\boldsymbol{K}}^{\boldsymbol{k}}(s).$$

If  $\nu$  puts no mass on  $\{v = -\infty\}$ , then we always have

$$(33) N_{v,v}^k(s) \le N_{v,K}^k(s).$$

We recall terminology introduced in [Berman and Boucksom 2010], providing a natural context in which the converse of (33) holds, with subexponential growth. A weighted subset of X is a pair (K, v) consisting of a closed nonpluripolar subset  $K \subseteq X$  and a function  $v \in C^0(K)$ .

Geometry & Topology, Volume 28 (2024)

1984

Let (K, v) be a weighted subset of X. A positive Borel probability measure v on K is *Bernstein–Markov* with respect to (K, v) if for each  $\varepsilon > 0$ , there is a constant  $C_{\varepsilon} > 0$  such that

(34) 
$$N_{v,K}^{k}(s) \le C_{\varepsilon} e^{\varepsilon k} N_{v,v}^{k}(s)$$

for any  $s \in H^0(X, L^k \otimes T)$  and any  $k \in \mathbb{N}$ . We write BM(K, v) for the set of Bernstein–Markov measures with respect to (K, v). As pointed out in [Berman et al. 2011], any volume form measure on X is Bernstein–Markov with respect to (X, v), with  $v \in C^{\infty}(X)$ .

**Proposition 6.2** Assume that (K, v) is a weighted subset of X. Then:

- (i)  $N_{n,K}^k$  is a norm on  $H^0(X, L^k \otimes T)$ .
- (ii) For any  $v \in BM(K, v)$ ,  $N_{v,v}^k$  is a norm on  $H^0(X, L^k \otimes T)$ .

**Proof** (i) As v is bounded,  $N_{v,K}^k$  is clearly finite on  $H^0(X, L^k \otimes T)$ . In order to show that it is a norm, it suffices to show that for any  $s \in H^0(X, L^k \otimes T)$ ,  $N_{v,K}^k(s) = 0$  implies that s = 0. In fact, we have  $s|_K = 0$ ; hence s = 0 by the connectedness of X.

(ii) As v is bounded, clearly  $N_{v,v}^k$  is finite and satisfies the triangle inequality. Nondegeneracy follows from the fact that  $N_{v,K}^k$  is a norm and (34).

#### 6.2 Partial Bergman kernels

In this section, with the terminology and context of the previous section, we fix a weighted subset (K, v) of X and  $v \in BM(K, v)$ . We introduce the associated partial Bergman kernels: for any  $k \in \mathbb{N}$  and  $x \in K$ ,

$$(35) \qquad B_{\nu,u,\nu}^k(x) := \sup\{h^k \otimes h_T(s,s)e^{-k\nu}(x) : N_{\nu,\nu}^k(s,s) \le 1, s \in H^0(X, L^k \otimes T \otimes \mathcal{I}(ku))\}.$$

The associated partial Bergman measures on X are identically zero on  $X \setminus K$ , and on K are defined as

(36) 
$$\beta_{\nu,u,\nu}^k := \frac{n!}{k^n} B_{\nu,u,\nu}^k \,\mathrm{d}\nu$$

Observe that

(37) 
$$\int_{K} \beta_{\nu,u,\nu}^{k} = \frac{n!}{k^{n}} h^{0}(X, L^{k} \otimes T \otimes \mathcal{I}(ku)).$$

Our aim is to show the following weak convergence result:

(38) 
$$\beta_{v,u,v}^{k} \rightharpoonup \theta_{P_{K}[u]_{\mathcal{J}}(v)}^{n}, \quad \text{as } k \to \infty.$$

We focus momentarily on the case when  $d\nu = \omega^n$  and K = X. That (38) holds in this particular case follows from [Ross and Witt Nyström 2017, Theorem 1.4]. Relying on the recent paper of Di Nezza and Trapani [2021], we give here a short proof of this result, borrowing ideas from Berman [2009] as well.

**Proposition 6.3** Let  $u \in \text{PSH}(X, \theta)$  be such that  $\theta_u$  is a Kähler current and  $[u] \in \mathcal{A}(X, \theta)$ . If  $v \in C^{\infty}(X)$ , then  $\beta_{v,u,\omega^n}^k \rightharpoonup \theta_{P_X[u]_{\mathcal{I}}(v)}^n = \theta_{P_X[u](v)}^n$  as  $k \to \infty$ .

Tamás Darvas and Mingchen Xia

**Proof** That  $\theta_{P_X[u]_{\delta}(v)}^n = \theta_{P_X[u](v)}^n$  follows from Lemma 5.2. We start with noticing that as  $k \to \infty$ ,  $\beta_{v,u,\omega^n}^k \le \beta_{v,V_{\theta},\omega^n}^k \rightharpoonup \theta_{P_X[V_{\theta}](v)}^n = \mathbb{1}_{\{v=P_X[V_{\theta}](v)\}} \theta_v^n$ ,

where the convergence follows from [Berman 2009, Theorem 1.2], and the last identity is due to [Di Nezza and Trapani 2021, Corollary 3.4]. Letting  $\mu$  be the weak limit of a subsequence of  $\beta_{v,u,\omega^n}^k$ , we obtain

(39) 
$$\mu \leq \lim_{k \to \infty} \beta_{v, V_{\theta}, \omega^n}^k = \mathbb{1}_{\{v = P_X[V_{\theta}](v)\}} \theta_v^n$$

Let  $s \in H^0(X, L^k \otimes T \otimes \mathcal{I}(ku))$  be a section such that  $N_{v,\omega^n}^k(s,s) \leq 1$ . Then by [Berman 2009, Lemma 4.1], there exists C > 0 such that  $h^k \otimes h_T(s,s)e^{-kv} \leq B_{v,u,\omega^n}^k \leq B_{v,V_\theta,\omega^n}^k \leq k^n C$ . This implies

$$\frac{1}{k}\log h^k \otimes h_T(s,s) \le v + \frac{\log C}{k} + n\frac{\log k}{k}$$

However, we also have that  $[(1/k) \log h^k \otimes h_T(s, s)] \leq [\tilde{u}_k^D] \leq \alpha_k[u]$ , where  $\tilde{u}_k^D$  is as defined in Remark 2.9, and  $\alpha_k \in (0, 1)$  is also from the notation of Remark 2.9. Let  $\varepsilon > 0$ . Observe that for all  $k \geq k_0(\varepsilon)$ , we have  $(1/k) \log h^k \otimes h_T(s, s) \in PSH(X, \theta + \varepsilon \omega)$ . In particular,

$$\frac{1}{k}\log h^k \otimes h_T(s,s) - \frac{\log C}{k} - n\frac{\log k}{k} \le P_X^{\theta + \varepsilon\omega}[\alpha_k u](v).$$

Now, taking the supremum over all candidates s, we obtain that

(40) 
$$B_{\nu,u,\omega^n}^k \le Ck^n e^{k(P_X^{\theta+\varepsilon\omega}[\alpha_k u](\nu)-\nu)} \quad \text{for } k \ge k_0.$$

We claim that  $\mu$  does not put mass on  $\{P_X^{\theta+\varepsilon\omega}[u](v) < v\}$  for any  $\varepsilon > 0$ . Since by Proposition 2.7  $P_X^{\theta+\varepsilon\omega}[\alpha_k u](v) \searrow P_X^{\theta+\varepsilon\omega}[u](v)$ , we get that  $\{P_X^{\theta+\varepsilon\omega}[\alpha_k u](v) < v\} \nearrow \{P_X^{\theta+\varepsilon\omega}[u](v) < v\}$ . As a result, to argue the claim, it suffices to show that  $\mu$  does not put mass on the set  $\{P_X^{\theta+\varepsilon\omega}[\alpha_k u](v) < v\}$  for any k. Note that the latter set is open, hence (40) implies our claim.

Since  $u \in \mathcal{A}(X,\theta)$ , we have that  $P_X^{\theta+\varepsilon\omega}[u](v) = [u]$  for all  $\varepsilon \ge 0$  by Lemma 5.2. As a result,  $P_X^{\theta+\varepsilon\omega}[u](v) \searrow P_X^{\theta}[u](v)$ . We can let  $\varepsilon \searrow 0$  to conclude that  $\mu$  does not put mass on  $\{P_X[u](v) < v\} = \bigcup_{\varepsilon>0} \{P_X^{\theta+\varepsilon\omega}[u](v) < v\}$ . Putting this together with (39), we obtain that

$$\mu \leq \mathbb{1}_{\{P_X[u](v)=v\}}\theta_v^n = \theta_{P_X[u](v)}^n,$$

where the last equality is due to [Di Nezza and Trapani 2021, Corollary 3.4]. Comparing total masses (via (37) and Theorem 1.1), we conclude that  $\mu = \theta_{P_X[u](v)}^n$ . As  $\mu$  is an arbitrary limit point of  $\beta_{v,u,\omega^n}^k$ , we conclude that  $\beta_{v,u,\omega^n}^k$  converges weakly to  $\theta_{P_X[u](v)}^n$  as  $k \to \infty$ .

Let Norm  $(H^0(X, L^k \otimes T \otimes \mathcal{I}(ku)))$  be the space of  $\mathbb{C}$ -norms on the vector space  $H^0(X, L^k \otimes T \otimes \mathcal{I}(ku))$ and let  $\mathcal{L}_{k,u}$ : Norm  $(H^0(X, L^k \otimes T \otimes \mathcal{I}(ku))) \to \mathbb{R}$  be the *partial Donaldson functional*, extending the definition from [Berman and Boucksom 2010],

$$\mathscr{L}_{k,u}(H) = \frac{n!}{k^{n+1}} \log \frac{\operatorname{vol}\{s : H(s) \le 1\}}{\operatorname{vol}\{s : N_{0,\omega^n}^k(s) \le 1\}},$$

where vol is simply the Euclidean volume.

Geometry & Topology, Volume 28 (2024)

**Proposition 6.4** Let  $w, w' \in C^0(X)$ . Suppose that  $u \in PSH(X, \theta)$  is such that  $\theta_u$  is a Kähler current and  $[u] \in \mathcal{A}(X, \theta)$ . Then

(41) 
$$\lim_{k \to \infty} \left( \mathscr{L}_{k,u}(N_{w,\omega^n}^k) - \mathscr{L}_{k,u}(N_{w',\omega^n}^k) \right) = \mathscr{I}_{[u],X}^{\theta}(w) - \mathscr{I}_{[u],X}^{\theta}(w').$$

In particular,

(42) 
$$\lim_{k \to \infty} \mathscr{L}_{k,u}(N_{w,\omega^n}^k) = \mathscr{I}_{[u],X}^{\theta}(w).$$

**Proof** First observe that by Proposition 6.2, for any k > 0, both  $N_{w,\omega^n}^k$  and  $N_{w',\omega^n}^k$  are norms, hence the expressions inside the limit in (41) make sense.

To start, we make the following classical observation:

$$\mathscr{L}_{k,u}(N_{w,\omega^{n}}^{k}) - \mathscr{L}_{k,u}(N_{w',\omega^{n}}^{k}) = \int_{0}^{1} \frac{\mathrm{d}}{\mathrm{d}t} \mathscr{L}_{k,u}(N_{w+t(w'-w),\omega^{n}}^{k}) \,\mathrm{d}t = \int_{0}^{1} \int_{X} (w'-w) \,\beta_{w+t(w'-w),u,\omega^{n}}^{k} \,\mathrm{d}t.$$

By Proposition 6.3, we have

$$\lim_{k \to \infty} \int_X (w' - w) \,\beta_{w+t(w'-w),u,\omega^n}^k = \int_X (w' - w) \,\theta_{P_X[u](w+t(w'-w))}^n$$

By Theorem 1.1 we have  $\left|\int_X (w'-w)\beta_{w+t(w'-w),u,\omega^n}^k\right| \le C \sup_X |w-w'|$ . Hence, by the dominated convergence theorem we obtain that

$$\lim_{k \to \infty} \left( \mathscr{L}_{k,u}(N_{w,\omega^n}^k) - \mathscr{L}_{k,u}(N_{w',\omega^n}^k) \right) = \int_0^1 \int_X (w' - w) \,\theta_{P_X[u](w + t(w' - w))}^n \,\mathrm{d}t = \mathscr{I}_{[u],X}^\theta(w) - \mathscr{I}_{[u],X}^\theta(w'),$$

where in the last equality we have used Proposition 5.10.

Finally, (42) is just a special case of (41) with w' = 0.

**Lemma 6.5** Let  $u \in PSH(X, \theta)$ . Let (K, v) be a weighted subset of X. Let  $v \in BM(K, v)$ . Then

(43) 
$$\lim_{k \to \infty} \left( \mathscr{L}_{k,u}(N_{v,K}^k) - \mathscr{L}_{k,u}(N_{v,v}^k) \right) = 0$$

This is a direct consequence of the definition of Bernstein–Markov measures (34).

**Corollary 6.6** Take  $w \in C^0(X)$  and  $u \in PSH(X, \theta)$  such that  $\theta_u$  is a Kähler current and  $[u] \in \mathcal{A}(X, \theta)$ . Then

$$\lim_{k \to \infty} \mathscr{L}_{k,u}(N_{w,X}^k) = \mathscr{I}_{[u],X}^{\theta}(w).$$

**Proof** This follows from Lemma 6.5 and Proposition 6.4 and the fact that  $\omega^n \in BM(X, 0)$ .

Using these preliminary facts we extend Proposition 6.4 for much less regular data, again relying on ideas from [Berman and Boucksom 2010]:

**Proposition 6.7** Let  $u \in PSH(X, \theta)$  be such that  $\theta_u$  is a Kähler current and  $[u] \in \mathcal{A}(X, \theta)$ . Let (K, v) and (K', v') be two weighted subsets of X. Then

(44) 
$$\lim_{k \to \infty} \left( \mathscr{L}_{k,u}(N_{v,K}^k) - \mathscr{L}_{k,u}(N_{v',K'}^k) \right) = \mathscr{I}_{[u],K}^{\theta}(v) - \mathscr{I}_{[u],K'}^{\theta}(v').$$

In particular,

(45) 
$$\lim_{k \to \infty} \mathscr{L}_{k,u}(N_{v,K}^k) = \mathscr{I}_{[u],K}^{\theta}(v).$$

**Proof** First observe that by Proposition 6.2, for any k > 0, both  $N_{v,K}^k$  and  $N_{v',K'}^k$  are norms, hence the expressions inside the limit in (44) make sense. Moreover, (45) is just a special case of (44) for K' = X and v' = 0.

To prove (44) it is enough to show that for any fixed  $w \in C^{\infty}(X)$  we have

(46) 
$$\lim_{k \to \infty} \left( \mathscr{L}_{k,u}(N_{v,K}^k) - \mathscr{L}_{k,u}(N_{w,\omega^n}^k) \right) = \mathscr{I}_{[u],K}^{\theta}(v) - \mathscr{I}_{[u],X}^{\theta}(w).$$

For  $\varepsilon \in (0, 1)$  small enough we have that  $\theta_{(1-\varepsilon)u}$  is still a Kähler current. Let us fix such  $\varepsilon$ , along with an arbitrary  $\varepsilon' \in (0, 1)$ .

Let  $\{v_j^-\}_j$  and  $\{v_j^+\}_j$  be the sequence of smooth global functions constructed in Lemma 5.9 for the data (K, v).

By Remark 2.9 there exists  $k_0(\varepsilon, \varepsilon') \in \mathbb{N}$  such that  $[(1/k) \log h^k \otimes h_T(s, s)] \leq [(1-\varepsilon)u]$ , as well as  $(1/k) \log h^k \otimes h_T(s, s) \in PSH(X, \theta + \varepsilon'\omega)$  for any  $s \in H^0(X, T \otimes L^k \otimes \mathfrak{I}(ku))$  and  $k \geq k_0(\varepsilon, \varepsilon')$ .

In particular, Lemma 5.1 gives that

$$N_{E_X^{\theta+\varepsilon'\omega}[(1-\varepsilon)u](v),X}^k(s) = N_{v,K}^k(s),$$
  

$$N_{E_X^{\theta+\varepsilon'\omega}[(1-\varepsilon)u](v_j^-),X}^k(s) = N_{v_j^-,X}^k(s),$$
  

$$N_{E_X^{\theta+\varepsilon'\omega}[(1-\varepsilon)u](v_j^+),X}^k(s) = N_{v_j^+,X}^k(s).$$

As  $E_X^{\theta+\varepsilon'\omega}[(1-\varepsilon)u](v_j^-) \le E_K^{\theta+\varepsilon'\omega}[(1-\varepsilon)u](v) \le E_X^{\theta+\varepsilon'\omega}[(1-\varepsilon)u](v_j^+)$ , by Lemma 6.1 we have  $N_{v_j^+,X}^k(s) \le N_{v,K}^k(s) \le N_{v_j^-,X}^k(s)$  for  $s \in H^0(X, T \otimes L^k \otimes \mathcal{I}(ku))$  and  $k \ge k_0(\varepsilon, \varepsilon')$ .

Composing with  $\mathcal{L}_{k,u}$ , we arrive at

$$\mathscr{L}_{k,u}(N_{v_j^-,X}^k) \le \mathscr{L}_{k,u}(N_{v,K}^k) \le \mathscr{L}_{k,u}(N_{v_j^+,X}^k) \quad \text{for } k \ge k_0(\varepsilon,\varepsilon').$$

For any j > 0, by Corollary 6.6 we get

$$\begin{aligned} \mathscr{I}^{\theta}_{[u],X}(v_{j}^{-}) - \mathscr{I}^{\theta}_{[u],X}(w) &= \lim_{k \to \infty} \left( \mathscr{L}_{k,u}(N_{v_{j}^{+},X}^{k}) - \mathscr{L}_{k,u}(N_{w,X}^{k}) \right) \\ &\leq \lim_{k \to \infty} \left( \mathscr{L}_{k,u}(N_{v,K}^{k}) - \mathscr{L}_{k,u}(N_{w,X}^{k}) \right) \leq \lim_{k \to \infty} \left( \mathscr{L}_{k,u}(N_{v,K}^{k}) - \mathscr{L}_{k,u}(N_{w,X}^{k}) \right) \\ &\leq \lim_{k \to \infty} \left( \mathscr{L}_{k,u}(N_{v_{j}^{-},X}^{k}) - \mathscr{L}_{k,u}(N_{w,X}^{k}) \right) = \mathscr{I}^{\theta}_{[u],X}(v_{j}^{+}) - \mathscr{I}^{\theta}_{[u],X}(w). \end{aligned}$$

Geometry & Topology, Volume 28 (2024)

Using Lemma 5.11, we can let  $j \to \infty$  to arrive at

$$\begin{aligned} \mathscr{I}^{\theta}_{[u],K}(v) - \mathscr{I}^{\theta}_{[u],K}(w) &\leq \underbrace{\lim_{k \to \infty}}_{k \to \infty} \left( \mathscr{L}_{k,u}(N^{k}_{v,K}) - \mathscr{L}_{k,u}(N^{k}_{w,X}) \right) \leq \underbrace{\lim_{k \to \infty}}_{k \to \infty} \left( \mathscr{L}_{k,u}(N^{k}_{v,K}) - \mathscr{L}_{k,u}(N^{k}_{w,X}) \right) \\ &\leq \mathscr{I}^{\theta}_{[u],K}(v) - \mathscr{I}^{\theta}_{[u],K}(w). \end{aligned}$$
ence, (46) follows.

Hence, (46) follows.

**Corollary 6.8** Let  $u \in PSH(X, \theta)$  such that  $\theta_u$  is a Kähler current and  $[u] \in \mathcal{A}(X, \theta)$ . Let (K, v) be a weighted subset of X. Assume that  $v \in BM(K, v)$ . Then

$$\lim_{k \to \infty} \mathscr{L}_{k,u}(N_{v,v}^k) = \mathscr{I}_{[u],K}^{\theta}(v)$$

**Proof** Our claim follows from Proposition 6.7 and Lemma 6.5.

**Proposition 6.9** Suppose that  $u \in PSH(X, \theta)$  with  $[u] \in \mathcal{A}(X, \theta)$ , and assume that  $\theta_u$  is a Kähler current. Let (K, v) be a weighted subset of X. Let  $v \in BM(K, v)$ . Then  $\beta_{v,u,v}^k \rightharpoonup \theta_{P_K[u]_{\mathscr{I}}(v)}^n = \theta_{P_K[u](v)}^n$  weakly as  $k \to \infty$ .

The following proof is similar to that of [Berman et al. 2011, Theorem B].

**Proof** For  $w \in C^0(X)$ , let

$$f_k(t) = \mathscr{L}_{k,u}(N_{v+tw,v}^k)$$
 and  $g(t) := \mathscr{I}_{[u],K}^{\theta}(v+tw).$ 

By Corollary 6.8  $\lim_{k\to\infty} f_k(t) = g(t)$ . Note that  $f_k$  is concave by Hölder's inequality (see [Berman et al. 2011, Proposition 2.4]), so by [Berman and Boucksom 2010, Lemma 7.6],  $\lim_{k\to\infty} f'_k(0) = g'(0)$ , which is equivalent to  $\beta_{v,u,v}^k \rightharpoonup \theta_{P_K[u](v)}^n$ , by Proposition 5.10. 

Next we deal with the case of Kähler currents:

**Proposition 6.10** Suppose that  $u \in PSH(X, \theta)$  such that  $\theta_u$  is a Kähler current. Let (K, v) be a weighted subset of X and  $v \in BM(K, v)$ . Then  $\beta_{u,v,v}^k \rightharpoonup \theta_{P_K[u]_{\mathscr{G}}(v)}^n$  as  $k \to \infty$ .

**Proof** Let  $\mu$  be the weak limit of a subsequence of  $\beta_{v,u,v}^k$ . We claim that

(47) 
$$\mu \le \theta_{P_K[u]_{\mathscr{I}}(v)}^n$$

Observe that this claim implies the conclusion. In fact, by Theorem 1.1, we have equality of the total masses, so equality holds in (47). As  $\mu$  is an arbitrary subsequential limit of the weak compact sequence  $\{\beta_{v,u,v}^k\}_k$ , we get that  $\beta_{v,u,v}^k \rightharpoonup \theta_{P_K[u]_{\emptyset}(v)}^n$  as  $k \to \infty$ .

We prove the claim. Let  $\{u_i^D\}_j$  be the approximation sequence of Theorem 2.8. By Lemmas 5.2 and 3.7, we know that  $d_{\mathcal{G}}([u_j^D], [P_K[u]_{\mathcal{F}}]) = d_{\mathcal{G}}([u_j^D], [P_K[u]_{\mathcal{F}}(v)]) \to 0$ . In particular,

(48) 
$$\lim_{j \to \infty} \int_X \theta_{P_K[u_j^D]_{\mathfrak{f}}(v)}^n = \int_X \theta_{P_K[u]_{\mathfrak{f}}(v)}^n.$$

We know that  $\theta_{u_i^D}$  are Kähler currents for high enough j. Since  $u \leq u_i^D$ , we trivially obtain that  $\beta_{v,u,v}^k \leq \beta_{v,u_j}^k$ , for any  $k \geq 1$ . As  $v \in BM(K, v)$ , by Proposition 6.9,  $\mu \leq \theta_{P_K[u_j^D]_j(v)}^n$  for any  $j \geq 1$ 

Geometry & Topology, Volume 28 (2024)

fixed. By Proposition 2.7,  $P_K[u_j^D]_{\mathscr{I}}(v) \searrow P_K[u]_{\mathscr{I}}(v)$  as  $j \to \infty$ . Hence, by (48) and [Darvas et al. 2018, Theorem 2.3], (47) follows.

Finally, the main result:

**Theorem 6.11** Suppose that  $u \in PSH(X, \theta)$ . Let (K, v) be a weighed subset of X, let  $v \in BM(K, v)$ . Then  $\beta_{v,u,v}^k \rightharpoonup \theta_{P_K[u]_{\ell}(v)}^n$  as  $k \to \infty$ .

Proof By Lemma 5.2 and [Darvas and Xia 2022, Proposition 2.18] we have that

$$H^{0}(X, L^{k} \otimes T \otimes \mathcal{I}(ku)) = H^{0}(X, L^{k} \otimes T \otimes \mathcal{I}(kP[u]_{\mathcal{I}})) = H^{0}(X, L^{k} \otimes T \otimes \mathcal{I}(kP[u]_{\mathcal{I}}(v))).$$

This allows us to replace u with  $P_K[u]_{\mathcal{I}}(v)$ . In addition, by Theorem 1.1 we can also assume that  $\int_X \theta_u^n > 0$ , otherwise there is nothing to prove.

By Proposition 3.6, there exists  $u_j \in PSH(X, \theta)$  such that  $u_j \nearrow u$  a.e. and  $\theta_{u_j}$  are Kähler currents. This gives  $\beta_{v,u_j,v}^k \leq \beta_{v,u,v}^k$ . Let  $\mu$  be the weak limit of a subsequence of  $\beta_{v,u,v}^k$ . Then by Proposition 6.10,  $\theta_{P_K[u_j]_{\mathfrak{f}}(v)}^n \leq \mu$ . By Proposition 2.7 and [Darvas et al. 2018, Theorem 2.3],  $\theta_{P_K[u_j]_{\mathfrak{f}}(v)}^n \nearrow \theta_{P_K[u]_{\mathfrak{f}}(v)}^n$ . Hence,

(49) 
$$\theta_{P_K[u]_4(v)}^n \le \mu.$$

A comparison of total masses ((37) and Theorem 1.1) gives that equality holds in (49). As  $\mu$  is an arbitrary subsequential limit of the weak compact sequence  $\{\beta_{v,u,\mu}^k\}_k$ , we obtain that  $\beta_{v,u,\nu}^k \rightharpoonup \theta_{P_K[u]_{\mathscr{I}}(v)}^n$  as  $k \rightarrow \infty$ .

# References

- [Ash et al. 2010] **A Ash, D Mumford, M Rapoport, Y-S Tai**, *Smooth compactifications of locally symmetric varieties*, 2nd edition, Cambridge Univ. Press (2010) MR Zbl
- [Bedford and Taylor 1976] **E Bedford**, **B A Taylor**, *The Dirichlet problem for a complex Monge–Ampère equation*, Invent. Math. 37 (1976) 1–44 MR Zbl
- [Bedford and Taylor 1982] E Bedford, B A Taylor, *A new capacity for plurisubharmonic functions*, Acta Math. 149 (1982) 1–40 MR Zbl
- [Berman 2009] **R J Berman**, *Bergman kernels and equilibrium measures for line bundles over projective manifolds*, Amer. J. Math. 131 (2009) 1485–1524 MR Zbl
- [Berman and Boucksom 2010] **R Berman**, **S Boucksom**, *Growth of balls of holomorphic sections and energy at equilibrium*, Invent. Math. 181 (2010) 337–394 MR Zbl
- [Berman et al. 2011] **R Berman, S Boucksom, D Witt Nyström,** *Fekete points and convergence towards equilibrium measures on complex manifolds*, Acta Math. 207 (2011) 1–27 MR Zbl
- [Berndtsson 2018] **B Berndtsson**, *Probability measures associated to geodesics in the space of Kähler metrics*, from "Algebraic and analytic microlocal analysis" (M Hitrik, D Tamarkin, B Tsygan, S Zelditch, editors), Springer Proc. Math. Stat. 269, Springer (2018) 395–419 MR Zbl

- [Bonavero 1998] L Bonavero, Inégalités de Morse holomorphes singulières, J. Geom. Anal. 8 (1998) 409–425 MR Zbl
- [Botero et al. 2022a] **A M Botero**, **J I Burgos Gil**, **D Holmes**, **R de Jong**, *Chern–Weil and Hilbert–Samuel formulae for singular Hermitian line bundles*, Doc. Math. 27 (2022) 2563–2624 MR Zbl
- [Botero et al. 2022b] A Botero, J I Burgos Gil, D Holmes, R de Jong, *Rings of Siegel–Jacobi forms of bounded relative index are not finitely generated* (2022) arXiv 2203.14583 To appear in Duke Math. J.
- [Bouche 1990] **T Bouche**, *Convergence de la métrique de Fubini–Study d'un fibré linéaire positif*, Ann. Inst. Fourier (Grenoble) 40 (1990) 117–130 MR Zbl
- [Boucksom 2002a] **S Boucksom**, *Cônes positifs des variétés complexes compactes*, PhD thesis, Université Joseph Fourier–Grenoble I (2002) Available at https://theses.hal.science/tel-00002268
- [Boucksom 2002b] S Boucksom, On the volume of a line bundle, Int. J. Math. 13 (2002) 1043–1063 MR Zbl
- [Boucksom 2017] **S Boucksom**, Singularities of plurisubharmonic functions and multiplier ideals, lecture notes (2017) Available at http://sebastien.boucksom.perso.math.cnrs.fr/notes/L2.pdf
- [Boucksom et al. 2008] **S Boucksom, C Favre, M Jonsson**, *Valuations and plurisubharmonic singularities*, Publ. Res. Inst. Math. Sci. 44 (2008) 449–494 MR Zbl
- [Boucksom et al. 2010] S Boucksom, P Eyssidieux, V Guedj, A Zeriahi, Monge–Ampère equations in big cohomology classes, Acta Math. 205 (2010) 199–262 MR Zbl
- [Cao 2014] J Cao, Numerical dimension and a Kawamata–Viehweg–Nadel-type vanishing theorem on compact Kähler manifolds, Compos. Math. 150 (2014) 1869–1902 MR Zbl
- [Catlin 1999] **D** Catlin, *The Bergman kernel and a theorem of Tian*, from "Analysis and geometry in several complex variables" (G Komatsu, M Kuranishi, editors), Birkhäuser, Boston (1999) 1–23 MR Zbl
- [Cegrell 1998] U Cegrell, Pluricomplex energy, Acta Math. 180 (1998) 187–217 MR Zbl
- [Chen and Sun 2012] X Chen, S Sun, Space of Kähler metrics, V: Kähler quantization, from "Metric and differential geometry" (X Dai, X Rong, editors), Progr. Math. 297, Birkhäuser, Basel (2012) 19–41 MR Zbl
- [Coman and Marinescu 2017] D Coman, G Marinescu, On the first order asymptotics of partial Bergman kernels, Ann. Fac. Sci. Toulouse Math. 26 (2017) 1193–1210 MR Zbl
- [Coman et al. 2019] **D Coman, G Marinescu, V-A Nguyên**, *Holomorphic sections of line bundles vanishing* along subvarieties, preprint (2019) arXiv 1909.00328
- [Darvas 2019] T Darvas, Geometric pluripotential theory on K\"ahler manifolds, from "Advances in complex geometry" (Y A Rubinstein, B Shiffman, editors), Contemp. Math. 735, Amer. Math. Soc., Providence, RI (2019) 1–104 MR Zbl
- [Darvas and Xia 2022] **T Darvas**, **M Xia**, *The closures of test configurations and algebraic singularity types*, Adv. Math. 397 (2022) art. id. 108198 MR Zbl
- [Darvas et al. 2018] **T Darvas**, **E Di Nezza**, **C H Lu**, *Monotonicity of nonpluripolar products and complex Monge–Ampère equations with prescribed singularity*, Anal. PDE 11 (2018) 2049–2087 MR Zbl
- [Darvas et al. 2020] **T Darvas, C H Lu, Y A Rubinstein**, *Quantization in geometric pluripotential theory*, Comm. Pure Appl. Math. 73 (2020) 1100–1138 MR Zbl
- [Darvas et al. 2021] T Darvas, E Di Nezza, H-C Lu, The metric geometry of singularity types, J. Reine Angew. Math. 771 (2021) 137–170 MR Zbl
- [Demailly 2012] **J-P Demailly**, *Analytic methods in algebraic geometry*, Surv. Modern Math. 1, International, Somerville, MA (2012) MR Zbl

- [Demailly 2015] J-P Demailly, On the cohomology of pseudoeffective line bundles, from "Complex geometry and dynamics" (JE Fornæss, M Irgens, E F Wold, editors), Abel Symp. 10, Springer (2015) 51–99 MR Zbl
- [Demailly 2018] **J-P Demailly**, *Extension of holomorphic functions and cohomology classes from non reduced analytic subvarieties*, from "Geometric complex analysis" (J Byun, HR Cho, S Y Kim, K-H Lee, J-D Park, editors), Springer Proc. Math. Stat. 246, Springer (2018) 97–113 MR Zbl
- [Demailly and Paun 2004] **J-P Demailly**, **M Paun**, *Numerical characterization of the Kähler cone of a compact Kähler manifold*, Ann. of Math. 159 (2004) 1247–1274 MR Zbl
- [Demailly et al. 2001] J-P Demailly, T Peternell, M Schneider, Pseudo-effective line bundles on compact Kähler manifolds, Int. J. Math. 12 (2001) 689–741 MR Zbl
- [Di Nezza and Trapani 2021] E Di Nezza, S Trapani, Monge–Ampère measures on contact sets, Math. Res. Lett. 28 (2021) 1337–1352 MR Zbl
- [Donaldson 2001] **S K Donaldson**, *Scalar curvature and projective embeddings*, *I*, J. Differential Geom. 59 (2001) 479–522 MR Zbl
- [Guan and Zhou 2015] **Q** Guan, X Zhou, A proof of Demailly's strong openness conjecture, Ann. of Math. 182 (2015) 605–616 MR Zbl
- [Guedj and Zeriahi 2005] V Guedj, A Zeriahi, Intrinsic capacities on compact Kähler manifolds, J. Geom. Anal. 15 (2005) 607–639 MR Zbl
- [Guedj and Zeriahi 2007] V Guedj, A Zeriahi, *The weighted Monge–Ampère energy of quasiplurisubharmonic functions*, J. Funct. Anal. 250 (2007) 442–482 MR Zbl
- [Kaveh and Khovanskii 2012] K Kaveh, A G Khovanskii, Newton–Okounkov bodies, semigroups of integral points, graded algebras and intersection theory, Ann. of Math. 176 (2012) 925–978 MR Zbl
- [Kim 2015] **D** Kim, Equivalence of plurisubharmonic singularities and Siu-type metrics, Monatsh. Math. 178 (2015) 85–95 MR Zbl
- [Kim and Seo 2020] **D** Kim, **H** Seo, *Jumping numbers of analytic multiplier ideals*, Ann. Polon. Math. 124 (2020) 257–280 MR Zbl
- [Lazarsfeld and Mustață 2009] R Lazarsfeld, M Mustață, Convex bodies associated to linear series, Ann. Sci. École Norm. Sup. 42 (2009) 783–835 MR Zbl
- [Lu 2000] Z Lu, On the lower order terms of the asymptotic expansion of Tian–Yau–Zelditch, Amer. J. Math. 122 (2000) 235–273 MR Zbl
- [Lu 2021] CHLu, Comparison of Monge-Ampère capacities, Ann. Polon. Math. 126 (2021) 31-53 MR Zbl
- [Lu and Nguyen 2015] CH Lu, V-D Nguyen, Degenerate complex Hessian equations on compact Kähler manifolds, Indiana Univ. Math. J. 64 (2015) 1721–1745 MR Zbl
- [Ma and Marinescu 2007] X Ma, G Marinescu, Holomorphic Morse inequalities and Bergman kernels, Progr. Math. 254, Birkhäuser, Basel (2007) MR Zbl
- [Phong and Sturm 2006] D H Phong, J Sturm, The Monge–Ampère operator and geodesics in the space of Kähler potentials, Invent. Math. 166 (2006) 125–149 MR Zbl
- [Ross and Singer 2017] J Ross, M Singer, Asymptotics of partial density functions for divisors, J. Geom. Anal. 27 (2017) 1803–1854 MR Zbl
- [Ross and Witt Nyström 2014] J Ross, D Witt Nyström, Analytic test configurations and geodesic rays, J. Symplectic Geom. 12 (2014) 125–169 MR Zbl

- [Ross and Witt Nyström 2017] J Ross, D Witt Nyström, *Envelopes of positive metrics with prescribed singularities*, Ann. Fac. Sci. Toulouse Math. 26 (2017) 687–728 MR Zbl
- [Runge 1995] B Runge, Theta functions and Siegel–Jacobi forms, Acta Math. 175 (1995) 165–196 MR Zbl
- [Shiffman and Zelditch 2003] **B Shiffman**, **S Zelditch**, *Equilibrium distribution of zeros of random polynomials*, Int. Math. Res. Not. 2003 (2003) 25–49 MR Zbl
- [Song and Zelditch 2010] J Song, S Zelditch, Bergman metrics and geodesics in the space of Kähler metrics on toric varieties, Anal. PDE 3 (2010) 295–358 MR Zbl
- [Sun 2020] J Sun, Asymptotics of partial density function vanishing along smooth subvariety, preprint (2020) arXiv 2010.04951
- [Tian 1988] **G Tian**, *Kähler metrics on algebraic manifolds*, PhD thesis, Harvard University (1988) Available at https://www.proquest.com/docview/303687845
- [Tian 1990] **G Tian**, On a set of polarized Kähler metrics on algebraic manifolds, J. Differential Geom. 32 (1990) 99–130 MR Zbl
- [Trusiani 2022] A Trusiani, Kähler–Einstein metrics with prescribed singularities on Fano manifolds, J. Reine Angew. Math. 793 (2022) 1–57 MR Zbl
- [Tsuji 2007] H Tsuji, Extension of log pluricanonical forms from subvarieties, preprint (2007) arXiv 0709.2710
- [Vu 2022] **D-V Vu**, *Convexity of the class of currents with finite relative energy*, Ann. Polon. Math. 128 (2022) 275–288 MR Zbl
- [Witt Nyström 2019] D Witt Nyström, Monotonicity of nonpluripolar Monge–Ampère masses, Indiana Univ. Math. J. 68 (2019) 579–591 MR Zbl
- [Xia 2019] M Xia, Integration by parts formula for non-pluripolar product, preprint (2019) arXiv 1907.06359
- [Xia 2021] **M Xia**, Partial Okounkov bodies and Duistermaat–Heckman measures of non-Archimedean metrics, preprint (2021) arXiv 2112.04290
- [Yau 1987] S-T Yau, Nonlinear analysis in geometry, Enseign. Math. 33 (1987) 109–158 MR Zbl
- [Zelditch 1998] **S Zelditch**, *Szegő kernels and a theorem of Tian*, Int. Math. Res. Not. 1998 (1998) 317–331 MR Zbl
- [Zelditch and Zhou 2019a] S Zelditch, P Zhou, Central limit theorem for spectral partial Bergman kernels, Geom. Topol. 23 (2019) 1961–2004 MR Zbl
- [Zelditch and Zhou 2019b] S Zelditch, P Zhou, Interface asymptotics of partial Bergman kernels on S<sup>1</sup>-symmetric Kähler manifolds, J. Symplectic Geom. 17 (2019) 793–856 MR Zbl
- [Zhang 2023] **K Zhang**, *A quantization proof of the uniform Yau–Tian–Donaldson conjecture*, J. Eur. Math. Soc. (online publication August 2023)

Department of Mathematics, University of Maryland

College Park, MD, United States

Department of Mathematics, Chalmers Tekniska Högskola

Göteborg, Sweden

Current address: Institut de mathématiques de Jussieu, Sorbonne Université

Paris, France

tdarvas@umd.edu, xia.mingchen@imj-prg.fr

Proposed: Gang Tian Seconded: Leonid Polterovich, John Lott Received: 29 May 2022 Revised: 29 September 2022

#### **GEOMETRY & TOPOLOGY**

msp.org/gt

#### MANAGING EDITOR

#### András I Stipsicz

cz Alfréd Rényi Institute of Mathematics stipsicz@renyi.hu

#### BOARD OF EDITORS

Mohammed Abouzaid	Stanford University abouzaid@stanford.edu	Mark Gross	University of Cambridge mgross@dpmms.cam.ac.uk
Dan Abramovich	Brown University dan_abramovich@brown.edu	Rob Kirby	University of California, Berkeley kirby@math.berkeley.edu
Ian Agol	University of California, Berkeley ianagol@math.berkeley.edu	Bruce Kleiner	NYU, Courant Institute bkleiner@cims.nyu.edu
Arend Bayer	University of Edinburgh arend.bayer@ed.ac.uk	Sándor Kovács	University of Washington skovacs@uw.edu
Mark Behrens	University of Notre Dame mbehren1@nd.edu	Urs Lang	ETH Zürich urs.lang@math.ethz.ch
Mladen Bestvina	University of Utah bestvina@math.utah.edu	Marc Levine	Universität Duisburg-Essen marc.levine@uni-due.de
Martin R Bridson	University of Oxford bridson@maths.ox.ac.uk	Ciprian Manolescu	University of California, Los Angeles cm@math.ucla.edu
Jim Bryan	University of British Columbia jbryan@math.ubc.ca	Haynes Miller	Massachusetts Institute of Technology hrm@math.mit.edu
Dmitri Burago	Pennsylvania State University burago@math.psu.edu	Tomasz Mrowka	Massachusetts Institute of Technology mrowka@math.mit.edu
Tobias H Colding	Massachusetts Institute of Technology colding@math.mit.edu	Aaron Naber	Northwestern University anaber@math.northwestern.edu
Simon Donaldson	Imperial College, London s.donaldson@ic.ac.uk	Peter Ozsváth	Princeton University petero@math.princeton.edu
Yasha Eliashberg	Stanford University eliash-gt@math.stanford.edu	Leonid Polterovich	Tel Aviv University polterov@post.tau.ac.il
Benson Farb	University of Chicago farb@math.uchicago.edu	Colin Rourke	University of Warwick gt@maths.warwick.ac.uk
David M Fisher	Rice University davidfisher@rice.edu	Roman Sauer	Karlsruhe Institute of Technology roman.sauer@kit.edu
Mike Freedman	Microsoft Research michaelf@microsoft.com	Stefan Schwede	Universität Bonn schwede@math.uni-bonn.de
David Gabai	Princeton University gabai@princeton.edu	Natasa Sesum	Rutgers University natasas@math.rutgers.edu
Stavros Garoufalidis	Southern U. of Sci. and Tech., China stavros@mpim-bonn.mpg.de	Gang Tian	Massachusetts Institute of Technology tian@math.mit.edu
Cameron Gordon	University of Texas gordon@math.utexas.edu	Ulrike Tillmann	Oxford University tillmann@maths.ox.ac.uk
Jesper Grodal	University of Copenhagen jg@math.ku.dk	Nathalie Wahl	University of Copenhagen wahl@math.ku.dk
Misha Gromov	IHÉS and NYU, Courant Institute gromov@ihes.fr	Anna Wienhard	Universität Heidelberg wienhard@mathi.uni-heidelberg.de

See inside back cover or msp.org/gt for submission instructions.

The subscription price for 2024 is US \$805/year for the electronic version, and \$1135/year (+\$70, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Geometry & Topology is indexed by Mathematical Reviews, Zentralblatt MATH, Current Mathematical Publications and the Science Citation Index.

Geometry & Topology (ISSN 1465-3060 printed, 1364-0380 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840.

GT peer review and production are managed by EditFLOW<sup>®</sup> from MSP.

PUBLISHED BY



# **GEOMETRY & TOPOLOGY**

Volume 28 Issue 4 (pages 1501–1993) 2024	
Localization in Khovanov homology	1501
MATTHEW STOFFREGEN and MELISSA ZHANG	
On definite lattices bounded by a homology 3–sphere and Yang–Mills instanton Floer theory	1587
CHRISTOPHER SCADUTO	
On automorphisms of high-dimensional solid tori	1629
MAURICIO BUSTAMANTE and OSCAR RANDAL-WILLIAMS	
On the subvarieties with nonsingular real loci of a real algebraic variety	1693
Olivier Benoist	
The triangulation complexity of fibred 3–manifolds	1727
MARC LACKENBY and JESSICA S PURCELL	
Sublinearly Morse boundary, II: Proper geodesic spaces	1829
YULAN QING, KASRA RAFI and GIULIO TIOZZO	
Small Dehn surgery and SU(2)	1891
JOHN A BALDWIN, ZHENKUN LI, STEVEN SIVEK and FAN YE	
Pseudo-Anosovs are exponentially generic in mapping class groups	1923
Іннуеок Сноі	
The volume of pseudoeffective line bundles and partial equilibrium	1957
TAMÁS DARVAS and MINGCHEN XIA	