

# **Rings and C\*-algebras generated by commutators**

Downloaded from: https://research.chalmers.se, 2024-11-22 21:35 UTC

Citation for the original published paper (version of record):

Gardella, E., Thiel, H. (2025). Rings and C\*-algebras generated by commutators. Journal of Algebra, 662: 214-241. http://dx.doi.org/10.1016/j.jalgebra.2024.08.020

N.B. When citing this work, cite the original published paper.

research.chalmers.se offers the possibility of retrieving research publications produced at Chalmers University of Technology. It covers all kind of research output: articles, dissertations, conference papers, reports etc. since 2004. research.chalmers.se is administrated and maintained by Chalmers Library

Contents lists available at ScienceDirect

Journal of Algebra

journal homepage: www.elsevier.com/locate/jalgebra



Research Paper

Rings and C\*-algebras generated by commutators  $\stackrel{\diamond}{\approx}$ 



ALGEBRA

Eusebio Gardella\*, Hannes Thiel

Department of Mathematical Sciences, Chalmers University of Technology and University of Gothenburg, Gothenburg SE-412 96, Sweden

#### ARTICLE INFO

Article history: Received 4 April 2024 Available online 30 August 2024 Communicated by Louis Rowen

MSC: primary 16N60, 46L05, 47B47 secondary 15A15, 15A23, 16S50, 19K14

Keywords: Commutators  $C^*$ -algebras Commutativity

#### ABSTRACT

We show that a unital ring is generated by its commutators as an ideal if and only if there exists a natural number N such that every element is a sum of N products of pairs of commutators. We show that one can take  $N \leq 2$  for matrix rings, and that one may choose  $N \leq 3$  for rings that contain a direct sum of matrix rings – this in particular applies to C\*-algebras that are properly infinite or have real rank zero. For Jiang-Su-stable C\*-algebras, we show that  $N \leq 6$  can be arranged.

For arbitrary rings, we show that every element in the commutator ideal admits a power that is a sum of products of commutators. Using that a C\*-algebra cannot be a radical extension over a proper ideal, we deduce that a C\*-algebra is generated by its commutators as a not necessarily closed ideal if and only if every element is a finite sum of products of pairs of commutators.

© 2024 The Author(s). Published by Elsevier Inc. This is an open access article under the CC BY license (http:// creativecommons.org/licenses/by/4.0/).

\* Corresponding author.

E-mail addresses: gardella@chalmers.se (E. Gardella), hannes.thiel@chalmers.se (H. Thiel).

URLs: https://www.math.chalmers.se/~gardella (E. Gardella), https://www.hannesthiel.org (H. Thiel).

#### https://doi.org/10.1016/j.jalgebra.2024.08.020

0021-8693/  $\odot 2024$  The Author(s). Published by Elsevier Inc. This is an open access article under the CC BY license (http://creativecommons.org/licenses/by/4.0/).

 $<sup>^{*}</sup>$  The first named author was partially supported by the Swedish Research Council Grant 2021-04561. The second named author was partially supported by the Knut and Alice Wallenberg Foundation (KAW 2021.0140).

## 1. Introduction

The commutator of two elements x and y in a ring is defined as [x, y] := xy - yx, and it is an elementary fact that the ideal generated by all commutators is the smallest ideal whose associated quotient ring is commutative. We study rings that are 'very noncommutative', in the sense that they are generated by their commutators as an ideal; equivalently, these are the rings that admit no nonzero ring homomorphism to a commutative ring.

A related property that has been studied is that of being *additively* generated by commutators [23,32]. This is however a much more restrictive condition, as for example the matrix algebras  $M_n(\mathbb{Z})$  for  $n \geq 2$  do not satisfy it, while they are easily seen to be generated by their commutators as an ideal.

For a unital ring, we show in Theorem 3.4 that being generated by its commutators as an ideal implies that every element is a sum of products of *pairs* of commutators, and there even is a uniform bound (depending only on the ring) on the number of summands. We reproduce a particular case here:

**Theorem A.** Let R be a unital ring. Then the following are equivalent:

- (1) The ring R is generated by its commutators as an ideal.
- (2) The ring R is generated by its commutators as a ring.
- (3) There exists  $N \in \mathbb{N}$  such that for every  $a \in R$  there exist  $b_j, c_j, d_j, e_j \in R$ , for  $j = 1, \ldots, N$ , such that

$$a = \sum_{j=1}^{N} [b_j, c_j] [d_j, e_j].$$

The fact that (1) implies (3) in the above theorem is most surprising, for two reasons: First, it states that double-products are sufficient to generate R. And second, it also shows that there exists a uniform bound on the number of summands needed.

A result of Baxter [1, Theorem 1] asserts that a simple ring is either a field, or every of its elements is a sum of products of pairs of commutators. Theorem A recovers Baxter's result in the unital case, and moreover shows that the minimal number of summands required is uniformly bounded.

Theorem A can be viewed as a Waring-type result, as we explain now. The classical Waring problem asks if for each  $k \geq 1$  there exists a uniform bound g(k) such that every integer can be written as a sum of at most g(k) many k-th powers of integers. This was solved positively by Hilbert in 1909, and since then numerous generalizations and variations of this problem have been considered. In the context of general rings, the Waring problem was introduced by Brešar in [6]; see also [30,10,11]. Here, given a unital ring R and a polynomial  $f \in \mathbb{Z}\langle x_1, \ldots, x_d \rangle$  in d noncommuting variables, one considers

the set  $f(R) := \{f(a_1, \ldots, a_d) : a_1, \ldots, a_d \in R\}$  and asks if there exists an integer N such that every element in R is a sum of at most N elements in f(R). The classical Waring problem is the case  $R = \mathbb{Z}$  and  $f(x) = x^k$ . Our results apply to the polynomial in four variables given by

$$f(a, b, c, d) = [a, b][c, d] = abcd - bacd - abdc + badc.$$

Theorem A shows that if a unital ring R is generated by f(R) then there is a constant N (depending only on R) such that every element is a sum of at most N elements in f(R).

The question of when a unital ring is generated by its commutators has also been studied in [14].

For not necessarily unital rings that are generated by their commutators as an ideal, we show in Theorem 3.5 that sums of products of commutators contain an ideal over which the ring is a radical extension. Here, a ring R is said to be a radical extension over a subring  $S \subseteq R$  if for every  $x \in R$  there exists  $n \ge 1$  with  $x^n \in S$ . The following is a particular case of Theorem 3.5.

**Theorem B.** Let R be a (not necessarily unital) ring that is generated by its commutators as an ideal. Then for every  $a \in R$ , there exist  $n, m \ge 1$  and  $b_j, c_j, d_j, e_j \in R$  for j = 1, ..., n such that

$$a^m = \sum_{j=1}^n [b_j, c_j] [d_j, e_j].$$

Of course, unital rings are never radical extensions over proper ideals. By [21, Proposition 2.2], C\*-algebras have this property as well. Using these facts, we obtain the following characterization of when a C\*-algebra is generated by its commutators; see Theorem 4.1.

**Theorem C.** Let A be a  $C^*$ -algebra. Then the following are equivalent:

- (1) The  $C^*$ -algebra A is generated by its commutators as an ideal.
- (2) The  $C^*$ -algebra A is generated by its commutators as a ring.
- (3) For every  $a \in A$  there exist  $n \in \mathbb{N}$  and  $b_j, c_j, d_j, e_j \in A$  for  $j = 1, \ldots, n$  such that

$$a = \sum_{j=1}^{n} [b_j, c_j] [d_j, e_j].$$

If a C\*-algebra is generated by its commutators as an ideal, then it admits no onedimensional irreducible representations. For unital C\*-algebras, the converse also holds, but there exist nonunital C\*-algebras that admit no one-dimensional irreducible representations which are not generated by their commutators as an ideal; see Example 4.4.

We note that, unless A is unital, it is not clear if there exists a uniform bound on the number of summands in part (3) of Theorem C; see Question 4.3.

Robert showed in [39, Theorem 3.2] that if a unital C\*-algebra A admits no onedimensional irreducible representations, then every element is a sum of commutators with a sum of products of commutators. Theorem A strengthens Robert's result by showing that it suffices to consider summands that are products of commutators; see Remarks 4.2. We point out that it is not clear that every commutator in a C\*-algebra (let alone in a ring) is a sum of products of pairs of commutators; see Question 7.1. This and related questions are discussed in Section 7.

Our proofs rely on the theory of Lie ideals in associative rings, as developed by Herstein [25] and others in the 1950s and 1960s. In the context of C\*-algebras, Lie ideals have been studied by Miers [33] and coauthors in the 1970s and 1980s, and more recently by Brešar, Kissin, and Shulman [9], and Robert [39].

In Section 5 we study the invariant  $\xi(R)$  that associates to a unital ring R generated by its commutators, the minimal N for which part (3) of Theorem A holds; see Definition 5.1. We obtain estimates for this invariant for matrix rings over arbitrary unital rings (Theorem 5.4), for division rings (Proposition 5.8), for semisimple<sup>1</sup> rings (Corollary 5.11), as well as for unital rings that contain a direct sum of matrix rings (Theorem 5.15).

In Section 6, we specialize to unital C\*-algebras and we obtain estimates for the invariant  $\xi$  for C\*-algebras that are properly infinite (Example 6.1), have real rank zero (Theorem 6.3), or contain a unital copy of the Jiang-Su algebra  $\mathcal{Z}$  (Theorem 6.4). The latter in particular applies to all C\*-algebras covered by the Elliott classification programme, as well as to the reduced group C\*-algebras  $C^*_{\lambda}(\mathbb{F}_n)$  of all nonabelian free groups.

In forthcoming work [22], we show that  $\xi(M) \leq 2$  for von Neumann algebras M that have no commutative summand, and that many such von Neumann algebras even satisfy the optimal value  $\xi(M) = 1$ . In [8], we show that  $\xi(M_n(D)) = 1$  for every division ring D with infinite center and  $n \geq 2$ , and we give an example of a commutative, unital ring R such that  $\xi(M_2(R)) = 2$ .

We have written this paper with both ring theorists and operator algebraists in mind. To make it accessible to both communities, we have included some basic definitions and statements of useful results along the way. We also sometimes mention  $L^p$ -operator algebras, for which we refer the reader to [15] for an introduction.

 $<sup>^{1}</sup>$  In ring theory, a semisimple ring is usually defined to be an *Artinian* ring with trivial Jacobson radical. We will follow this convention in this paper.

## A cknowledgments

The authors thank Matej Brešar, Tsiu-Kwen Lee and Leonel Robert for valuable comments on earlier versions of this paper. We also thank the referee for their valuable comments and suggestions.

## 2. A commutativity result

In this section, we study rings R satisfying  $[[R, R], [R, R]^2] = \{0\}$ , that is, rings where commutators commute with products of pairs of commutators. We show that a semiprime ring with this property is automatically commutative; see Theorem 2.3. For general rings, said condition implies that the commutator ideal is nil, meaning that each of its elements is nilpotent; see Corollary 2.4.

The results in this section are rather elementary and probably folklore in ring theory. Since we were not able to locate them in the literature, we give here complete proofs, but note that they could also be obtained by using either the theory of functional identities or the theory of polynomial identities.

Given elements x, y in a ring, we use [x, y] := xy - yx to denote the (additive) commutator. Following the customary convention in algebra, when X and Y are subsets of a ring, we use [X, Y] to denote the additive subgroup generated by the set of commutators [x, y] with  $x \in X$  and  $y \in Y$ . We sometimes consider just the set of commutators, and then we use the notation

$$[X,Y]_1 := \{ [x,y] \colon x \in X, y \in Y \}$$

for given subsets X and Y of a ring.

Similarly, we use XY to denote the additive subgroup generated by the set  $\{xy : x \in X, y \in Y\}$ , and we write  $X^2$  for XX. If we want to specify the set of products, we use the notation

$$X \bullet Y := \{xy \colon x \in X, y \in Y\}$$

for given subsets X and Y of a ring. Note that XY is therefore the additive subgroup generated by  $X \bullet Y$ .

An ideal  $I \subseteq R$  in a ring is said to be *prime* if  $I \neq R$  and whenever  $J_1, J_2 \subseteq R$  are ideals with  $J_1J_2 \subseteq I$ , then we have  $J_1 \subseteq I$  or  $J_2 \subseteq I$ . A ring R is *prime* if  $\{0\}$  is a prime ideal, that is, whenever  $I, J \subseteq R$  are ideals and  $IJ = \{0\}$ , then we have  $I = \{0\}$ or  $J = \{0\}$ . (Equivalently, whenever  $a, b \in R$  satisfy  $aRb = \{0\}$ , then either a = 0 or b = 0.) Further, a ring R is *semiprime* if the intersection of all prime ideals in R is  $\{0\}$ . (Equivalently, whenever  $a \in R$  satisfies  $aRa = \{0\}$ , then a = 0.) We refer to [28, Section 10] for details. An additive subgroup L of a ring R is said to be a Lie ideal if  $[R, L] \subseteq L$ , or equivalently  $[L, R] \subseteq L$ . Finally, recall that a ring R is said to have no 2-torsion if  $x \in R$  and 2x = 0 imply x = 0.

**Proposition 2.1.** Let R be a ring, and let  $L \subseteq R$  be a Lie ideal. Consider the following conditions:

- (1) We have  $[L, R] = \{0\}$ , that is, L is a subset of the center of R.
- (2) We have  $[L, L] = \{0\}.$
- (3) We have  $[L, L^2] = \{0\}.$

Then the implications  $(1) \Rightarrow (2) \Rightarrow (3)'$  hold. If R is semiprime, then the implication  $(3) \Rightarrow (2)'$  holds. Moreover, if R is semiprime and has no 2-torsion, then all conditions are equivalent.

**Proof.** It is clear that (1) implies (2). If [L, L] = 0, namely if every element of L commutes with L, then clearly also every element of L commutes with products of elements of L, so (2) implies (3).

Assume that R is semiprime. We will show that (3) implies (2), using an argument inspired by the proof of [29, Lemma 1]. We first establish the following:

Claim: For every  $x, y \in L$ , we have  $[x, y]^2 = 0$ . Using that y commutes with yx, we have yxy = yyx. Similarly, we have xxy = xyx. Using this at the last step, we get

$$[x, y]^{2} = xyxy - xyyx - yxxy + yxyx = x(yxy - yyx) - y(xxy - xyx) = 0,$$

which proves the claim.

Now, let  $x, y \in L$  and set u := [x, y]. We will show that u = 0. Let  $a \in R$ . Claim: We have  $(ua)^4 = 0$ . Applying the Jacobi identity, we have

$$[u, a] = [[x, y], a] = [x, [y, a]] + [y, [a, x]].$$
(2.1)

Since L is a Lie ideal, it follows that both [y, a] and [a, x] belong to L. Thus [x, [y, a]] and [y, [a, x]] belong to [L, L], and hence their squares are zero by the above claim. Further, [x, [y, a]] and [y, [a, x]] both belong to L (again because L is a Lie ideal), and they also belong to  $L^2$ . In particular, they commute. Set s = [x, [y, a]] and t = [y, [a, x]], and note that  $s^2 = t^2 = 0$  by the claim above. Using that s and t commute at the second step, it follows that

$$[u,a]^3 \stackrel{(2.1)}{=} \left( \left[ x, [y,a] \right] + \left[ y, [a,x] \right] \right)^3 = s^3 + 3s^2t + 3st^2 + t^3 = 0.$$

Using that  $u^2 = 0$  (by the claim) at the third step, one easily checks that

$$0 = [u, a]^3 ua = (ua - au)^3 ua = (ua)^4,$$

as desired.

It follows that  $\{u\} \cup uR$  is a right ideal of R that is nil of bounded index at most 4. By Herstein's generalization of Levitzki's theorem, [24, Lemma 1.1], it follows that u = 0; see also [28, Exercise 10.13].

Suppose that R is semiprime and has no 2-torsion. To show that (2) implies (1), assume that  $[L, L] = \{0\}$ . Let  $x \in L$  and  $a \in R$ . Since L is a Lie ideal, we have  $[x, a] \in L$ , and thus [x, [x, a]] = 0. Thus,  $[x, [x, R]] = \{0\}$ . By [25, Theorem 1], if M is a Lie ideal in a semiprime ring S without 2-torsion, and if  $y \in S$  satisfies  $[y, [y, M]] = \{0\}$ , then  $[y, M] = \{0\}$ . Applied to y = x and M = R, it follows that  $[x, R] = \{0\}$ . Thus,  $[L, R] = \{0\}$ , as desired.  $\Box$ 

In Proposition 2.1, condition (2) need not imply (1) in semiprime rings with 2-torsion, as we show in the following example.

**Example 2.2.** Let  $\mathbb{F}$  be a field of characteristic 2, set  $R := M_2(\mathbb{F})$ , the ring of 2-by-2 matrices over  $\mathbb{F}$ , and

$$L := \left\{ \begin{pmatrix} \mu & \lambda \\ \lambda & \mu \end{pmatrix} : \mu, \lambda \in \mathbb{F} \right\}.$$

Then R is semiprime and  $[L, L] = \{0\}$  (so that  $[L, L^2] = 0$  as well). Moreover, using that  $\mathbb{F}$  has characteristic 2, a direct computation shows that L is a Lie ideal in R. On the other hand, since the center of R consists only of scalar multiples of the identity, we see that L is not contained in the center and thus  $[L, R] \neq \{0\}$ .

We now specialize to the case that the Lie ideal is the commutator subgroup.

In preparation for the proof of the next result, let us recall that the characteristic of a ring R is defined as the smallest natural number  $n \in \mathbb{N}$  with  $n \geq 1$  such that  $nR = \{0\}$ , and as 0 if no such n exists. Note that a prime ring R has characteristic  $n \geq 1$  if and only if there exists  $a \in R \setminus \{0\}$  with na = 0. Indeed, given any  $b \in R$ , we have  $aR(nb) = (na)Rb = \{0\}$ . Using that R is prime, we have a = 0 or nb = 0, but  $a \neq 0$  and so nb = 0. In particular, a prime ring has characteristic 0 if and only if it has no torsion.

**Theorem 2.3.** Let R be a semiprime ring. Then the following are equivalent:

- (1) The ring R is commutative.
- (2) We have  $[[R, R], R] = \{0\}$ , that is, [R, R] is a subset of the center of R.
- (3) We have  $[[R, R], [R, R]^2] = \{0\}.$
- (4) We have  $[[R, R], [R, R]] = \{0\}.$

**Proof.** It is clear that (1) implies (2), and that (2) implies (3). By Proposition 2.1, (3) implies (4).

Let us show that (4) implies (1). To verify that R is commutative, let  $x, y \in R$ . We will show that [x, y] = 0. Since R is semiprime, the intersection of all of its prime ideals is  $\{0\}$ . Thus, to show that [x, y] = 0, it suffices to show that [x, y] belongs to every prime ideal of R. Let  $I \subseteq R$  be a prime ideal. Then the quotient ring Q := R/I is prime and satisfies  $[[Q, Q], [Q, Q]] = \{0\}$ . In order to show that [x, y] = 0, we will show that Q is commutative. In other words, and this is indeed what we will do, we may assume that R is prime.

Let Z denote the center of R. Using that [R, R] is a Lie ideal of R satisfying  $[[R, R], [R, R]] = \{0\}$ , it follows from [29, Theorem 4] that  $[R, R] \subseteq Z$ , unless we have the following exceptional case: R has characteristic 2,  $Z \neq \{0\}$ , and the localization  $RZ^{-1}$  of R at Z is isomorphic to the ring  $M_2(\mathbb{F})$  of 2-by-2 matrices over the quotient field  $\mathbb{F}$  of Z. We claim that the exceptional case cannot happen under our assumptions. Arguing by contradiction, assume that it does. Then  $\mathbb{F}$  is a field of characteristic 2. The assumption  $[[R, R], [R, R]] = \{0\}$  implies that  $[[RZ^{-1}, RZ^{-1}], [RZ^{-1}, RZ^{-1}]] = \{0\}$  as well. Set  $S := M_2(\mathbb{F})$  and

$$M := \left\{ \begin{pmatrix} e & f \\ g & e \end{pmatrix} : e, f, g \in \mathbb{F} \right\}.$$

Since

$$\begin{bmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix}, \begin{pmatrix} a' & b' \\ c' & d' \end{pmatrix} \end{bmatrix} = \begin{pmatrix} bc' - cb' & * \\ * & cb' - bc' \end{pmatrix}$$

and  $\mathbb{F}$  has characteristic 2, it follows that every commutator in S belongs to M. Further, for  $e, f, g \in \mathbb{F}$ , we have

$$\begin{bmatrix} \begin{pmatrix} e & 0 \\ g & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} = \begin{pmatrix} g & e \\ 0 & g \end{pmatrix}, \text{ and } \begin{bmatrix} \begin{pmatrix} e & f \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \end{bmatrix} = \begin{pmatrix} f & 0 \\ e & f \end{pmatrix}$$

Thus, M is the commutator subgroup of S. In particular, the matrices  $\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$  and  $\begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$  belong to M, but their commutator does not vanish. Thus,  $[M, M] \neq \{0\}$ . This shows that R is not of the exceptional case.

Thus, we have  $[R, R] \subseteq Z$ . We conclude that  $R \subseteq Z$  by [29, Lemma 7], and hence R is commutative.  $\Box$ 

**Corollary 2.4.** Let R be a ring such that  $[[R, R], [R, R]^2] = \{0\}$ . Then the commutator ideal of R is nil.

**Proof.** Let *P* denote the lower nilradical (or prime radical) of *R*. Then *P* is a nil ideal and R/P is semiprime; see [28, Definition 10.13]. It follows from Theorem 2.3 that R/P is commutative. Hence, the commutator ideal of *R* is contained in *P* and therefore nil.  $\Box$ 

#### 3. Rings generated by commutators

In this section, we show that a unital ring is generated by its commutators as an ideal if and only if every element can be written as a sum of products of pairs of commutators. Moreover, we show that there is a universal bound on the number of summands required; see Theorem 3.4. In the general case of a not necessarily unital ring, we show that every element in the commutator ideal has a power that is a sum of products of pairs of commutators; see Theorem 3.5.

We actually work in the more general setting of algebras over rings, so we fix a unital commutative ring K and will consider K-algebras. The case of rings is obtained by specializing to  $K = \mathbb{Z}$ .

Given a subset  $X \subseteq A$  in a K-algebra A and  $n \in \mathbb{N}$ , we use

$$\sum^{n} X := \{x_1 + x_2 + \ldots + x_n \in A \colon x_1, \ldots, x_n \in X\}$$

to denote the collection of elements in A that can be written as a sum of n elements from X.

A Lie ideal in a K-algebra A is a K-linear subspace  $L \subseteq A$  such that  $[A, L] \subseteq L$ . Given Lie ideals  $L, M \subseteq A$ , it is easy to show, using the Jacobi identity, that the subspace [L, M] is a Lie ideal as well.

Given subsets  $X, Y \subseteq A$ , recall from the beginning of Section 2 that

$$[X, Y]_1 := \{ [x, y] : x \in X, y \in Y \}$$
 and  $X \bullet Y := \{ xy : x \in X, y \in Y \}.$ 

We further set  $X^{\bullet 2} := X \bullet X$ .

**Lemma 3.1.** Let L, M be Lie ideals in a K-algebra A. Then

$$A[L, M]A \subseteq A[L, M].$$

Moreover, if  $L_0 \subseteq L$  and  $M_0 \subseteq M$  are subsets, and  $m, n \in \mathbb{N}$  satisfy

$$[A, L_0]_1 \subseteq \sum^n L_0, \text{ and } [A, M_0]_1 \subseteq \sum^m M_0,$$

then  $A \bullet [L_0, M_0]_1 \bullet A \subseteq \sum^{1+n+m} A \bullet [L_0, M_0]_1$ .

**Proof.** We only prove the quantitative statement since the proof of the non-quantitative statement is analogous. Let  $a, b \in A$ , let  $x \in L_0$ , and let  $y \in M_0$ . It follows from the Jacobi identity that

$$[[x, y], b] = -[[y, b], x] - [[b, x], y] = [[b, y], x] + [[x, b], y],$$

and therefore

$$a[x,y]b = a[[x,y],b] + ab[x,y]$$
  
=  $a[[b,y],x] + a[[x,b],y] + ab[x,y] \in \sum_{n+1}^{n+m} A \bullet [L_0, M_0]_1,$ 

as desired.  $\hfill\square$ 

It is a folklore result from Lie theory that if L is a Lie ideal in an algebra, then the ideal generated by [L, L] is contained in  $L + L^2$ ; see, for example [39, Lemma 1.1]. We include the result together with a quantitative statement for future reference.

Lemma 3.2. Let L be a Lie ideal in a K-algebra A. Then

$$A[L,L]A \subseteq A[L,L] \subseteq L+L^2.$$

Moreover, if  $L_0 \subseteq L$  and  $n \in \mathbb{N}$  satisfy  $[A, L_0]_1, [L_0, A]_1 \subseteq \sum^n L_0$ , then

$$A \bullet [L_0, L_0]_1 \subseteq \sum^n (L_0 + L_0^{\bullet 2}), \text{ and } A \bullet [L_0, L_0]_1 \bullet A \subseteq \sum^{n(1+2n)} (L_0 + L_0^{\bullet 2}).$$

**Proof.** We only prove the quantitative statement since the proof of the non-quantitative statement is analogous. Given  $a \in A$  and  $x, y \in L_0$ , we have

$$a[x,y] = [ax,y] + [y,a]x \in [A, L_0]_1 + [L_0, A]_1 \bullet L_0 \subseteq \sum^n (L_0 + L_0^{\bullet 2}).$$

Applying Lemma 3.1 at the first step with  $M_0 = L_0$  and m = n, we obtain

$$A \bullet [L_0, L_0]_1 \bullet A \subseteq \sum^{1+2n} A \bullet [L_0, L_0]_1 \subseteq \sum^{n(1+2n)} (L_0 + L_0^{\bullet 2}),$$

as desired.  $\Box$ 

The next result is a variation of Lemma 3.2 that allows us to strengthen containment in  $L + L^2$  to containment in  $L^2$ .

**Lemma 3.3.** Let L be a Lie ideal in a K-algebra A such that  $[A, L^2] \subseteq L$ . (For example, this is automatically the case if L contains [A, A].) Then:

$$A[L, L^2]A \subseteq A[L, L^2] \subseteq L^2.$$

Moreover, if  $L_0 \subseteq L$  and  $m, n \in \mathbb{N}$  satisfy  $[A, L_0]_1 \subseteq \sum^n L_0$  and  $[L_0^{\bullet 2}, A]_1 \subseteq \sum^m L_0$ , then

$$A \bullet [L_0, L_0^{\bullet 2}]_1 \subseteq \sum^{(2n+m)} L_0^{\bullet 2}, \quad and \quad A \bullet [L_0, L_0^{\bullet 2}]_1 \bullet A \subseteq \sum^{(2n+m)(1+3n)} L_0^{\bullet 2}.$$

**Proof.** Again we only prove the quantitative statement. For  $a \in A$  and  $x, y \in L_0$ , we have

E. Gardella, H. Thiel / Journal of Algebra 662 (2025) 214-241

$$[a, xy] = [a, x]y + x[a, y] \in \left(\sum^{n} L_{0}\right) \bullet L_{0} + L_{0} \bullet \left(\sum^{n} L_{0}\right) = \sum^{2n} L_{0}^{\bullet 2}$$

Thus, setting  $M_0 = L_0^{\bullet 2}$ , we get  $[A, M_0]_1 \subseteq \sum^{2n} M_0$ . Since  $[A, L_0]_1 \subseteq \sum^n L_0$ , it follows from Lemma 3.1 that

$$A \bullet [L_0, L_0^{\bullet 2}]_1 \bullet A \subseteq \sum^{(1+3n)} A \bullet [L_0, L_0^{\bullet 2}]_1.$$
(3.1)

Given  $a \in A$  and  $x, y, z \in L_0$ , using that  $[A, L_0^{\bullet 2}]_1 \subseteq \sum^{2n} L_0^{\bullet 2}$  and  $[A, L_0^{\bullet 2}]_1 \bullet L_0 \subseteq \sum^m L_0^{\bullet 2}$ , we have

$$a[x, yz] = [ax, yz] + [yz, a]x \in [A, L_0^{\bullet 2}]_1 + [L_0^{\bullet 2}, A]_1 \bullet L_0 \subseteq \sum_{n=1}^{\infty} (2n+m) L_0^{\bullet 2}.$$

Thus  $A \bullet [L_0, L_0^{\bullet 2}]_1 \subseteq \sum_{n=1}^{\infty} (2n+m) L_0^{\bullet 2}$ . Using this at the second step, we get

$$A \bullet [L_0, L_0^{\bullet 2}]_1 \bullet A \stackrel{(3.1)}{\subseteq} \sum {}^{(1+3n)} A \bullet [L_0, L_0^{\bullet 2}]_1 \subseteq \sum {}^{(2n+m)(1+3n)} L_0^{\bullet 2},$$

as desired.  $\Box$ 

**Theorem 3.4.** Let A be a unital K-algebra. Then the following are equivalent:

- (1) We have A = A[A, A]A, that is, A agrees with its commutator ideal.
- (2) The algebra A is generated by its commutators as a K-algebra.
- (3) There exists  $N \in \mathbb{N}$  such that for every  $a \in A$  there exist  $b_j, c_j, d_j, e_j \in A$  for  $j = 1, \ldots, N$  such that

$$a = \sum_{j=1}^{N} [b_j, c_j] [d_j, e_j].$$

**Proof.** It is clear that (3) implies (2), which in turn implies (1). To show that (1) implies (3), assume that A = A[A, A]A. Set

$$L := [A, A], \text{ and } I := A[L, L^2]A.$$

We first show that I = A. To reach a contradiction, assume that  $I \neq A$ . Using that A is unital, choose a maximal ideal  $J \subseteq A$  with  $I \subseteq J$ . Set B := A/J, which is a simple and unital K-algebra, and therefore a prime ring.

We claim that  $[[B, B], [B, B]^2] = \{0\}$ . To see this, let  $b_1, \ldots, b_6 \in B$ . We want to show that  $[[b_1, b_2], [b_3, b_4][b_5, b_6]] = 0$ . Let  $\pi: A \to B$  denote the quotient map, and find  $a_1, \ldots, a_6 \in A$  such that  $\pi(a_j) = b_j$  for all  $j = 1, \ldots, 6$ . Then  $[[a_1, a_2], [a_3, a_4][a_5, a_6]]$  belongs to I by definition (and since A is unital), and thus

$$\left[ [b_1, b_2], [b_3, b_4] [b_5, b_6] \right] = \pi \left( \left[ [a_1, a_2], [a_3, a_4] [a_5, a_6] \right] \right) = 0,$$

224

as desired.

Thus, the prime ring B satisfies  $[[B, B], [B, B]^2] = \{0\}$ . Applying Theorem 2.3, we deduce that B is commutative. Since, by assumption, the algebra A is generated by its commutators as an ideal, the same is true for B. This is a contradiction, which shows that I = A.

Set  $L_0 := [A, A]_1$ . It follows that A is the linear span of  $A \bullet [L_0, L_0^2]_1 \bullet A$ , and we obtain  $d \in \mathbb{N}$  and  $a_j, b_j \in A$  and  $x_j, y_j, z_j \in L_0$ , for  $j = 1, \ldots, d$ , such that

$$1 = \sum_{j=1}^d a_j [x_j, y_j z_j] b_j.$$

We claim that every element in A is a sum of at most 12d products of pairs of commutators. To show this, note that  $[A, L_0]_1 \subseteq L_0$  and  $[A, L_0^2]_1 \subseteq L_0$ . Applying Lemma 3.3 with m = n = 1 gives

$$A \bullet [L_0, L_0^2]_1 \bullet A \subseteq \sum{}^{12} L_0^{\bullet 2}.$$

$$(3.2)$$

Let  $a \in A$ . Then

$$a = \sum_{j=1}^{d} a a_j [x_j, y_j z_j] b_j \in \sum^{d} A \bullet [L_0, L_0^{\bullet 2}]_1 \bullet A \stackrel{(3.2)}{\subseteq} \sum^{12d} L_0^{\bullet 2}$$

This verifies (3) for N := 12d, and finishes the proof.  $\Box$ 

In Sections 5 and 6, we will study the minimal N for which (3) in Theorem 3.4 is satisfied.

We now turn to not necessarily unital rings and K-algebras. In this case, we show that every element in the commutator ideal has a power that is a sum of products of pairs of commutators.

Given a non-unital K-algebra A, we let  $\widetilde{A}$  denote the unitization, which is given by  $\widetilde{A} = K \times A$  with coordinatewise addition and scalar multiplication, and multiplication defined as  $(\lambda, x)(\mu, y) = (\lambda \mu, \mu x + \lambda y + xy)$  for  $\lambda, \mu \in K$  and  $x, y \in A$ . Then  $\widetilde{A}$  is a unital K-algebra, with unit (1,0), and the canonical map  $A \to \widetilde{A}$  identifies A with an ideal in  $\widetilde{A}$ . If A is unital, we set  $\widetilde{A} := A$ . See [5, Section 2.3] for details. The minimal unitization of a C\*-algebra A agrees with this construction when A is viewed as a  $\mathbb{C}$ -algebra.

Given a K-algebra A, the ideal generated by a subspace L is L + AL + LA + ALA, which agrees with  $\widetilde{A}L\widetilde{A}$  viewed inside  $\widetilde{A}$ . Note that ALA is also an ideal in A, which, however, may not contain L if A is not unital.

**Theorem 3.5.** Let A be a K-algebra and consider the ideals

$$I := \widetilde{A} \big[ [A, A], [A, A]^2 \big] \widetilde{A} \quad and \quad J := \widetilde{A} [A, A] \widetilde{A}.$$

Then  $I \subseteq [A, A]^2 \subseteq J$ , and J/I is nil. Thus, for every  $a, d \in \widetilde{A}$  and  $b, c \in A$ , there exist  $n, m \ge 1$  and  $w_j, x_j, y_j, z_j \in A$  for  $j = 1, \ldots, n$  such that

$$(a[b,c]d)^m = \sum_{j=1}^n [w_j, x_j][y_j, z_j].$$

In particular, if  $A = \widetilde{A}[A, A]\widetilde{A}$ , then for every  $a \in A$ , there exist  $n, m \geq 1$  and  $b_j, c_j, d_j, e_j \in A$  for  $j = 1, \ldots, n$ , such that

$$a^m = \sum_{j=1}^n [b_j, c_j] [d_j, e_j].$$

**Proof.** That I is contained in  $[A, A]^2$  follows from Lemma 3.3, while the inclusion  $[A, A]^2 \subseteq J$  is clear. Note that the quotient B := J/I satisfies  $[[B, B], [B, B]^2] = \{0\}$  and is generated by its commutators as an ideal. It thus follows from Corollary 2.4 that B is nil, as desired.  $\Box$ 

## 4. C\*-algebras generated by commutators

In this section, we prove that if a C\*-algebra is generated by its commutators as a (not necessarily closed) ideal, then every element is a sum of products of pairs of commutators; see Theorem 4.1.

We use  $\widetilde{A}$  to denote the minimal unitization of a C\*-algebra A; see the comments before Theorem 3.5. The (not necessarily closed) ideal generated by a subspace  $L \subseteq A$  is  $\widetilde{A}L\widetilde{A}$ . While in general it is not clear if  $ALA = \widetilde{A}L\widetilde{A}$  since L may not be a subset of ALA, for the special case L = [A, A] we showed in [21, Proposition 3.2] that  $[A, A] \subseteq A[A, A]A$ and therefore  $A[A, A]A = \widetilde{A}[A, A]\widetilde{A}$ .

A character on a C\*-algebra A is a one-dimensional, irreducible representation. (Equivalently, a nonzero homomorphism  $A \to \mathbb{C}$ .)

**Theorem 4.1.** Let A be a  $C^*$ -algebra. Then the following are equivalent:

- (1) We have  $A = \widetilde{A}[A, A]\widetilde{A}$ , that is, A is generated by its commutators as a (not necessarily closed) ideal.
- (2) We have A = A[A, A]A, that is, for every  $a \in A$  there exist  $n \in \mathbb{N}$  and  $b_j, c_j, d_j, e_j \in A$ , for  $j = 1, \ldots, n$ , such that

$$a = \sum_{j=1}^{n} b_j [c_j, d_j] e_j.$$

(3) We have A = A[[A.A], [A.A]]A.

- (4) The  $C^*$ -algebra A is generated by its commutators as a ring.
- (5) We have  $A = [A, A]^2$ , that is, for every  $a \in A$  there exist  $n \in \mathbb{N}$  and  $b_j, c_j, d_j, e_j \in A$ , for  $j = 1, \ldots, n$ , such that

$$a = \sum_{j=1}^{n} [b_j, c_j] [d_j, e_j].$$

Moreover, these conditions imply:

(6) The  $C^*$ -algebra A admits no character.

Moreover, (6) implies that A[A, A]A is dense in A, and thus all conditions are equivalent if A is unital.

**Proof.** By [21, Theorem 3.3], (1), (2) and (3) are equivalent. It is clear that (5) implies (4), which in turn implies (1).

Let us show that (1) implies (5). Thus, assuming that  $A = \widetilde{A}[A, A]\widetilde{A}$ , consider the ideal

$$I := \widetilde{A}[[A, A], [A, A]^2]\widetilde{A}.$$

Applying Theorem 3.5, we obtain that

$$I \subseteq [A, A]^2 \subseteq A,$$

and A/I is nilradical. Thus A is a radical extension over I. Since C\*-algebras are never radical extension over proper ideals ([21, Proposition 2.2]), we get I = A, and hence  $A = [A, A]^2$ , as desired.

We have seen that (1)-(5) are equivalent. It is easy to see that every character vanishes on commutators, which shows that (4) implies (6).

We will now show that (6) implies A[A, A]A = A. To reach a contradiction, assume that  $J := \overline{A[A, A]A}$  is a proper subset of A, and thus a proper (closed, two-sided) ideal. The quotient A/J is a commutative C\*-algebra and therefore admits a character. Composed with the quotient map  $A \to A/J$ , we obtain a character for A, which is the desired contradiction.

The last claim follows from the fact that the only dense ideal in a unital C\*-algebra is the C\*-algebra itself, and hence (6) implies (2) in this case.  $\Box$ 

**Remarks 4.2.** (1) It was shown in [7, Lemma 2.5] that a unital  $C^*$ -algebra is generated by its commutators as an ideal if and only if it is generated by its commutators as an algebra, and if and only if it admits no characters. Theorem 4.1 refines and generalizes this result. (2) Let A be a unital C\*-algebra that admits no characters. Robert showed in [39, Theorem 3.2] that every element in A can be written as

$$\sum_{j=1}^{N} [a_j, b_j] + \sum_{j=1}^{N} [c_j, d_j] [e_j, f_j]$$

for some  $N \in \mathbb{N}$  and elements  $a_j, b_j, c_j, d_j, e_j, f_j \in A$ , for  $j = 1, \ldots, N$ . Theorem 4.1 shows that the summands  $[a_j, b_j]$  are not necessary, and every element can be written as a sum of products of commutators.

It is not known if every commutator in a C\*-algebra is a linear combination of products of pairs of commutators; see Question 7.1(c).

**Question 4.3.** Let A be a nonunital C\*-algebra that is generated by its commutators as an ideal. Is there a uniform upper bound on the number of summands necessary in statement (3) in Theorem 4.1?

The following example was provided by Ozawa, who kindly allowed us to include it here. It shows that (6) does not imply (1)-(5) in Theorem 4.1 for nonunital C\*-algebras. In the example, we use the C\*-algebraic product of C\*-algebras  $A_n$  for  $n \in \mathbb{N}$ , which is defined as

$$\prod_{n \in \mathbb{N}} A_n = \Big\{ (a_n)_{n \in \mathbb{N}} \colon a_n \in A_n, \sup_{n \in \mathbb{N}} \|a_n\| < \infty \Big\}.$$

**Example 4.4.** By [40, Corollary 8.6], there exists a sequence  $(A_n)_{n \in \mathbb{N}}$  of unital, simple, infinite-dimensional C\*-algebras  $A_n$  such that the product  $\prod_{n \in \mathbb{N}} A_n$  has a character. We claim that for every  $m \geq 1$  and C > 1, there exists n = n(m, C) such that in  $A_n$  the following holds: If  $1 = \sum_{j=1}^m a_j[b_j, c_j]d_j$  for  $a_j, b_j, c_j, d_j \in A_n$ , then  $\sum_{j=1}^m \|a_j\| \|b_j\| \|c_j\| \|d_j\| > C$ .

Indeed, if this is not the case, then there exist  $m \ge 1$  and C > 1 such that for every  $n \in \mathbb{N}$  there are  $a_{n,j}, b_{n,j}, c_{n,j}, d_{n,j} \in A_n$  such that

$$1 = \sum_{j=1}^{m} a_{n,j} [b_{n,j}, c_{n,j}] d_{n,j}, \text{ and } \sum_{j=1}^{m} \|a_{n,j}\| \|b_{n,j}\| \|c_{n,j}\| \|d_{n,j}\| \le C.$$

One may arrange that  $||a_{n,j}|| = ||b_{n,j}|| = ||c_{n,j}|| = ||d_{n,j}||$  for each n and j. It follows that

$$a_j := (a_{n,j})_n, \quad b_j := (b_{n,j})_n, \quad c_j := (c_{n,j})_n, \text{ and } d_j := (d_{n,j})_n$$

are bounded sequences and thus belong to  $\prod_{n \in \mathbb{N}} A_n$ . Then  $1 = \sum_{j=1}^m a_j [b_j, c_j] d_j$  in  $\prod_{n \in \mathbb{N}} A_n$ , which contradicts that  $\prod_{n \in \mathbb{N}} A_n$  has a one-dimensional irreducible representation.

For  $m \in \mathbb{N}$ , consider the natural number  $n(m, m^3)$  as in the claim for m = m and  $C = m^3$ . Let us consider the direct sum  $B := \bigoplus_m A_{n(m,m^3)}$ . Since each summand  $A_{n(m,m^3)}$  has no character, neither does B. On the other hand, we claim that B does not agree with its commutator ideal. To see this, assume that the element  $e := (\frac{1}{m})_{m \in \mathbb{N}}$  belongs to the commutator ideal. Then there exist  $r \in \mathbb{N}$  and sequences  $a_k = (a_{m,k})_{m \in \mathbb{N}}$ ,  $b_k = (b_{m,k})_{m \in \mathbb{N}}$ ,  $c_k = (c_{m,k})_{m \in \mathbb{N}}$ ,  $d_k = (d_{m,k})_{m \in \mathbb{N}} \in B$  with  $e = \sum_{k=1}^r a_k [b_k, c_k] d_k$ . Choose m such that  $m \ge r$  and  $m^2 \ge \sum_{k=1}^r \|a_k\| \|b_k\| \|c_k\| \|d_k\|$ . In the quotient  $A_{n(m,m^3)}$  of B, we have

$$\frac{1}{m} = \sum_{k=1}^{r} a_{m,k} [b_{m,k}, c_{m,k}] d_{m,k}, \text{ and } \sum_{k=1}^{r} \|a_{m,k}\| \|b_{m,k}\| \|c_{m,k}\| \|d_{m,k}\| \le m^2,$$

a contradiction to the choice of  $n(m, m^3)$ .

Given a C\*-algebra A, we use  $A_0$  to denote the set of self-adjoint elements in A that vanish under every tracial state on A. A *self-commutator* in A is an element of the form  $[x^*, x]$  for some  $x \in A$ . Clearly, every self-commutator belongs to  $A_0$ , and consequently every norm-convergent sum  $\sum_{j=1}^{\infty} [x_j^*, x_j]$  belongs to  $A_0$ . It was shown by Cuntz and Pedersen, [13], that all elements of  $A_0$  arise this way.

Thus, every element in  $A_0$  can be approximated in norm by finite sums of selfcommutators. However, the following example shows that there is no uniform bound on the number of summands needed – even if A is unital. This should be contrasted with Theorem 3.4.

**Example 4.5.** Based on examples by Pedersen and Petersen [34], Bice and Farah showed in [2, Theorem 2.1] that for each m there exist a unital C\*-algebra B and an element  $b \in B_0$  that is not a limit of sums of m self-commutators.

Robert showed that this phenomenon can even be accomplished in a simple C\*-algebra for all m simultaneously: By [38, Theorem 1.4], there exists a simple, separable, unital C\*-algebra A with a unique tracial state  $\tau$  that contains contractive elements  $a_m \in A_0$ , for  $m \ge 1$ , satisfying  $||a_m - \sum_{j=1}^m [x_j^*, x_j]|| \ge 1$  for all  $x_1, \ldots, x_m \in A$ . Let us see that dist $(a_{4m}, \sum_{j=1}^m [A, A]_1) = 1$ . Since  $a_{4m}$  is contractive, it suffices to

Let us see that  $\operatorname{dist}(a_{4m}, \sum^{m}[A, A]_1) = 1$ . Since  $a_{4m}$  is contractive, it suffices to show that  $\operatorname{dist}(a_{4m}, \sum^{m}[A, A]_1) \geq 1$ . To reach a contradiction, assume that there is  $c \in \sum^{m}[A, A]_1$  with  $||a_{4m} - c|| < 1$ . Then  $||a_{4m} - \frac{1}{2}(c + c^*)|| < 1$ . Note that  $\frac{1}{2}(c + c^*)$  is a self-adjoint element in  $\sum^{2m}[A, A]_1$ . It follows from [31, Theorem 2.4] (and its proof), that every self-adjoint element in  $\sum^{k}[A, A]_1$  is a sum of at most 2k self-commutators, for every  $k \in \mathbb{N}$ . Thus,  $a_{4m}$  has distance strictly less than 1 to a sum of at most 4mself-commutators, which is the desired contradiction.

It follows that  $A_0$  is not contained in the closure of  $\sum_{i=1}^{m} [A, A]_1$  for any  $m \in \mathbb{N}$ . On the other hand, since A is unital, by Theorem 3.4 there exists  $N \in \mathbb{N}$  such that  $A = \sum_{i=1}^{N} [A, A]_1^{\bullet 2}$ .

## 5. Bounds on the number of commutators for rings

In Theorem 3.4 we have seen that if a unital ring R is generated by its (additive) commutators, then there exists  $N \in \mathbb{N}$  such that  $R = \sum^{N} [R, R]_{1}^{\bullet 2}$ , where  $[R, R]_{1}$  denotes the set of commutators in R, and  $[R, R]_{1}^{\bullet 2}$  denotes the set of products of pairs of commutators. We define the invariant  $\xi(R)$  as the minimal N with this property; see Definition 5.1. We show that  $\xi(M_n(S)) \leq 2$  for the ring  $M_n(S)$  of *n*-by-*n* matrices over an arbitrary unital ring S and  $n \geq 2$ ; see Theorem 5.4. For matrix algebras over a field  $\mathbb{F}$ , we prove that  $\xi(M_n(\mathbb{F})) = 1$ , which amounts to showing that every matrix is a product of two matrices of trace zero; see Example 5.5.

We then show that every noncommutative division ring D satisfies  $\xi(D) \leq 2$  (Proposition 5.8), and deduce that  $\xi(R) \leq 2$  for every semisimple ring R that has no commutative summand (Corollary 5.11).

In Lemma 5.13, we develop a method to obtain good bounds for  $\xi(R)$  for a unital ring R from the existence of special elements in a unital subring of R. Using this, we prove that  $\xi(R) \leq 3$  as soon as R admits a unital ring homomorphism from a direct sum of matrix rings; see Theorem 5.15.

**Definition 5.1.** Given a unital ring R generated by its commutators, we set

$$\xi(R) := \min\left\{N \in \mathbb{N} : R = \sum{}^{N} [R, R]_{1}^{\bullet 2}\right\}.$$

**Remarks 5.2.** (1) In [37], Pop shows that a unital C\*-algebra A admits no tracial states if and only if there exists n such that  $A = \sum^{n} [A, A]_{1}$ , and in this case he defines the invariant  $\nu(A)$  as the minimal number n such that  $A = \sum^{n} [A, A]_{1}$ . Our invariant  $\xi$  is defined analogously.

(2) For a K-algebra A over a unital, commutative ring K, it would arguably be more natural to define  $\xi(A)$  as the smallest N such that every element of A is a K-linear combination of (at most) N elements in  $[A, A]_1^{\bullet 2}$ . Since the set  $[A, A]_1^{\bullet 2}$  is K-invariant, this alternative definition agrees with the one given above.

(3) In [39, Example 3.11], Robert shows that for every  $N \in \mathbb{N}$  there exists a unital C\*-algebra A such that not every element of A can be expressed as a sum of N elements in  $[A, A]_{1}^{\bullet 2}$ , that is, such that  $\xi(A) > N$ .

**Question 5.3.** What are the possible values  $\xi(R)$  for unital rings R? What are the possible values for unital, simple rings?

The rest of this work is concerned with finding upper bounds for  $\xi(R)$  for different classes of rings R. We begin with matrix rings. In [32, Theorem 15], Mesyan showed that for every unital ring S and  $n \geq 2$ , every matrix in  $M_n(S)$  with trace zero is a sum of two commutators. Based on his ideas, we show that every matrix in  $M_n(S)$ , regardless of its trace, is a sum of two products of pairs of commutators. **Theorem 5.4.** Let S be a unital ring and let  $n \ge 2$ . Then  $\xi(M_n(S)) \le 2$ .

**Proof.** Assume first that n = 2. Given  $x, y \in S$ , we have

$$\left[ \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & x \\ -y & 0 \end{pmatrix} \right] = \begin{pmatrix} 0 & x \\ y & 0 \end{pmatrix} \in [M_2(S), M_2(S)]_1.$$

Further, we have

$$\begin{bmatrix} \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \in [M_2(S), M_2(S)]_1.$$
(5.1)

Therefore, given any  $a, b, c, d \in S$ , we have

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} 0 & b \\ -c & 0 \end{pmatrix} + \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & d \\ a & 0 \end{pmatrix} \in \sum^2 [M_2(S), M_2(S)]_1^{\bullet 2}.$$

Assume now that  $n \geq 3$ . For j, k = 1, ..., n, let  $e_{j,k} \in M_n(\{0,1\}) \subseteq M_n(S)$  denote the standard matrix unit. Let  $a \in M_n(S)$  and set

$$x := \sum_{j=1}^{n-1} e_{j+1,j}, \quad y := \sum_{j=1}^{n-1} e_{j,j+1}, \text{ and } c := a + xay + \ldots + x^{n-1}ay^{n-1}.$$

As shown in the proof of [32, Theorem 15], one can readily check that

$$[cy, x] = a - ce_{n,n}.$$

Note that the entries of the matrix  $a - ce_{n,n}$  agree with those of a in the first n - 1 columns.

Using block-diagonal matrices with blocks  $\binom{0\ 1}{0\ 0}$  and  $\binom{0\ 0}{1\ 0}$  and using the identity in (5.1), we see that the diagonal matrix d with entries  $1, -1, \ldots, 1, -1, 0$  (for n odd) and with entries  $1, -1, \ldots, 1, -1, 0$  (for n odd) and with entries  $1, -1, \ldots, 1, -1, 0, 0$  (for n even) is a commutator in  $M_n(S)$ . Moreover, we have [cy, x]d = ad, so ad is a product of two commutators. Set  $p_n = 1$  if n is odd and  $p_n = 2$  if n is even. Note that ad is an arbitrary matrix in  $M_n(S)$  whose last  $p_n$ -many columns are zero.

We have shown that every matrix whose last  $p_n$ -many columns are zero is a product of two commutators in  $M_n(S)$ . Analogously, every matrix whose first  $p_n$ -many columns are zero is also a product of two commutators. Since  $n \ge 3$ , every matrix is a sum of two such matrices, thus concluding the proof.  $\Box$ 

For matrix algebras over a field  $\mathbb{F}$ , we have the optimal estimate  $\xi(M_n(\mathbb{F})) = 1$ .

**Example 5.5.** Let  $\mathbb{F}$  be a field, and let  $n \geq 2$ . For characteristic 0, it was shown by Wu in [46, Theorem 5.12] that every matrix in  $M_n(\mathbb{F})$  is a product of two (additive) commutators. Botha showed in [4, Theorem 4.1] that the same holds in positive characteristic.

Thus, we have  $\xi(M_n(\mathbb{F})) = 1$ . This means that the image of the polynomial f(a, b, c, d) = [a, b][c, d] in  $M_n(\mathbb{F})$  is all of  $M_n(\mathbb{F})$ . In particular, this verifies the L'vov-Kaplansky conjecture for this polynomial, for all fields, and for all  $n \geq 2$ . We refer the reader to the survey [27] for background on the L'vov-Kaplansky conjecture.

**Example 5.6.** Let R be a unital ring that admits a surjective inner derivation, that is, we have  $R = \{[a, x] : x \in R\}$  for some element  $a \in R$ . This clearly implies that every element in R is a product of two commutators and thus  $\xi(R) = 1$ . Examples of such rings include the algebra of endomorphisms on an infinite-dimensional vector space, and the Weyl algebras over fields of characteristic 0; see [44, Examples 1.1].

This means that the polynomial f(a, b, c, d) = [a, b][c, d] is surjective on unital rings admitting a surjective inner derivation. Much more generally, Vitas showed in [44, Theorem 1.2] that A = f(A) for every nonzero multilinear polynomial f on a unital algebra A over a field such that A admits a surjective inner derivation.

Let S be a unital ring and let  $n \ge 2$ . While every matrix of trace zero in  $M_n(S)$  is a sum of two commutators by [32, Theorem 15], it was shown by Rosset and Rosset in [42] that not every matrix of trace zero in  $M_n(S)$  is necessarily a commutator. In forthcoming work [8], we show that the answer to the following question is 'Yes'.

**Question 5.7.** Do there exist a unital ring S and  $n \ge 2$  such that  $\xi(M_n(S)) = 2$ ?

Recall that a division ring, being simple, is generated by its commutators as an ideal if and only if it is not commutative (that is, not a field).

**Proposition 5.8.** Let D be a noncommutative division ring. Then

$$D = [D, D]_1 + [D, D]_1 \bullet [D, D]_1 = [D, D]_1 \bullet [D, D]_1 + [D, D]_1 \bullet [D, D]_1,$$

and in particular  $\xi(D) \leq 2$ .

**Proof.** It is clear that  $[D, [D, D]_1]_1 \subseteq [D, D]_1$ . We claim that also  $[D, [D, D]_1]_1 \subseteq [D, D]_1^{\bullet 2}$ . To see this, let  $r, s, t \in D$ . If [s, t] = 0, then  $[r, [s, t]] = 0 \in [D, D]_1^{\bullet 2}$ . If  $[s, t] \neq 0$ , then

$$[r, [s, t]] = [r[s, t]^{-1}, [s, t]][s, t] \in [D, D]_1^{\bullet 2}.$$

By Theorem 2.3, we have  $[[D, D], [D, D]] \neq \{0\}$ . Since every nonzero element in D is invertible, this shows that there exist  $a \in D$  and  $v, w \in [D, D]_1$  such that 1 = a[v, w]. Now, given any  $d \in D$ , we have

$$d = da[v, w] = [dav, w] + [w, da]v \in [D, [D, D]_1]_1 + [D, D]_1 \bullet [D, D]_1.$$

By the previous comments, the above is contained both in  $[D, D]_1 + [D, D]_1 \bullet [D, D]_1$ and in  $[D, D]_1 \bullet [D, D]_1 + [D, D]_1 \bullet [D, D]_1$ , as desired.  $\Box$ 

It was shown by Harris in [23] that there exist division rings D with  $D = [D, D]_1$ , hence also  $D = [D, D]_1^{*2}$ . In that case, we have  $\xi(D) = 1$ .

**Question 5.9.** Do we have  $\xi(D) = 1$  for every noncommutative division ring D?

**Question 5.10.** Do we have  $\xi(M_n(D)) = 1$  for every  $n \ge 2$  and every (noncommutative) division ring D?

Recall that a ring is said to be *semisimple* if it is Artinian (namely, if it has no infinite descending sequence of left or right ideals), and its Jacobson radical is trivial.

**Corollary 5.11.** Let R be a semisimple ring which is generated by its commutators as an ideal. Then  $\xi(R) \leq 2$ . If R is additionally finite-dimensional over an algebraically closed field, then  $\xi(R) = 1$ .

**Proof.** Since R is semisimple, by the Artin–Wedderburn Theorem there exist  $m, n_1, \ldots, n_m \in \mathbb{N}$  and division rings  $D_1, \ldots, D_m$  such that  $R \cong \bigoplus_{j=1}^m M_{n_j}(D_j)$ . Since  $M_{n_j}(D_j)$  is simple for all  $j = 1, \ldots, m$ , it follows that R is generated by its commutators as an ideal if and only if for all j with  $n_j = 1$ , the division ring  $D_j$  is not commutative, that is, if and only if no summand  $M_{n_j}(D_j)$  is a field. Applying Theorem 5.4 for the summands with  $n_j \ge 2$ , and Proposition 5.8 for the summands with  $n_j = 1$ , we deduce that  $\xi(R) \le 2$ .

If R is additionally finite-dimensional over an algebraically closed field  $\mathbb{F}$ , then  $R \cong \bigoplus_{j=1}^{m} M_{n_j}(\mathbb{F})$  for some integers  $n_j \ge 1$ . In this case, R is generated by its commutators if and only if  $n_j \ge 2$  for every j, in which case  $\xi(R) = 1$  by Example 5.5.  $\Box$ 

We now turn to estimates relating the invariant  $\xi$  of a ring and a subring. The following is a first crude estimate that can probably be improved.

**Proposition 5.12.** Let R be a unital ring, and let  $S \subseteq R$  be a subring with  $1 \in S$ . Then  $\xi(R) \leq 15\xi(S)^3$ .

**Proof.** Set  $M_0 := [S, S]_1$  and  $N := \xi(S)$ . We clearly have  $[M_0, S]_1 \subseteq M_0$ . Moreover, we claim that  $[M_0^{\bullet 2}, S]_1 \subseteq \sum^2 M_0^{\bullet 2}$ . To see this, let  $s \in S$  and  $x, y \in M_0$ . Then

$$[s, xy] = [s, x]y + x[s, y] \in M_0^{\bullet 2} + M_0^{\bullet 2},$$

as desired. Using the above, it follows from Lemma 3.1 that

$$S \bullet [M_0, M_0^{\bullet 2}]_1 \bullet S \subseteq \sum {}^4S \bullet [M_0, M_0^{\bullet 2}]_1.$$

$$(5.2)$$

Note also that  $S = \sum^{N} M_0^{\bullet 2}$  immediately implies that

$$S \bullet [S, S]_1 = \sum^{N^2} S \bullet [M_0^{\bullet 2}, M_0^{\bullet 2}]_1.$$
 (5.3)

Using unitality of S and the identity [ab, cd] = a[b, cd] + [a, cd]b at the fourth step, we get

$$S = \sum^{N} [S, S]_{1} \bullet [S, S]_{1} \subseteq \sum^{N} S \bullet [S, S]_{1} \stackrel{(5.3)}{=} \sum^{N^{3}} S \bullet [M_{0}^{\bullet 2}, M_{0}^{\bullet 2}]_{1}$$
(5.4)  
= 
$$\sum^{N^{3}} \left( S \bullet [M_{0}, M_{0}^{\bullet 2}]_{1} + S \bullet [M_{0}, M_{0}^{\bullet 2}]_{1} \bullet S \right) \stackrel{(5.2)}{\subseteq} \sum^{5N^{3}} S \bullet [M_{0}, M_{0}^{\bullet 2}]_{1}.$$

Set  $L_0 := [R, R]_1$ . We have  $[R, L_0]_1 \subseteq L_0$  and  $[L_0^{\bullet 2}, R]_1 \subseteq L_0$ , and therefore  $R \bullet [L_0, L_0^{\bullet 2}]_1 \subseteq \sum^3 L_0^{\bullet 2}$  by Lemma 3.3. Using that  $M_0 \subseteq L_0$  and using the above at the last step, we get

$$R = R \bullet S \stackrel{(5,4)}{\subseteq} \sum {}^{5N^3} R \bullet S \bullet [M_0, M_0^{\bullet 2}]_1 \subseteq \sum {}^{5N^3} R \bullet [L_0, L_0^{\bullet 2}]_1 \subseteq \sum {}^{15N^3} L_0^{\bullet 2}.$$
  
Thus,  $R = \sum {}^{15N^3} [R, R]_1 \bullet [R, R]_1$  and therefore  $\xi(R) \le 15\xi(S)^3$ .  $\Box$ 

If R is a unital ring that contains a unital subring of the form  $\bigoplus_{j=1}^{m} M_{n_j}(S_j)$  for unital rings  $S_j$  and  $n_j \ge 2$ , then Proposition 5.12 and Theorem 5.4 show that  $\xi(R) \le 120$ . We now present a method that can be used to obtain a much better bound, and which we will use in Theorem 5.15 to verify that  $\xi(R) \le 3$ .

**Lemma 5.13.** Let R be a unital ring, let  $S \subseteq R$  be a subring with  $1 \in S$ , and let  $m, n \in \mathbb{N}$ .

- (1) If  $1 \in \sum^{m} S \bullet [[S,S]_{1}, [S,S]_{1}]_{1}$ , then  $R = \sum^{m} ([R,R]_{1} + [R,R]_{1} \bullet [R,R]_{1})$ . In particular, if there exist  $a \in S$  and  $u, v \in [S,S]_{1}$  with 1 = a[u,v], then  $R = [R,R]_{1} + [R,R]_{1} \bullet [R,R]_{1}$ .
- (2) If  $1 \in \sum^{n} S \bullet [[S,S]_1, [S,S]_1^{\bullet 2}]$ , then  $\xi(R) \leq 3n$ . In particular, if there exist  $a \in S$  and  $u, v, w \in [S,S]_1$  such that 1 = a[u,vw], then  $\xi(R) \leq 3$ .
- **Proof.** Set  $L_0 := [R, R]_1$ , and note that  $[R, L_0]_1, [L_0, R]_1 \subseteq L_0$  and  $[L_0^{\bullet 2}, R]_1 \subseteq L_0$ .
  - (1). Using the assumption at the first step, and Lemma 3.2 at the second step, we get

$$R = \sum^{m} R \bullet [L_0, L_0]_1 \subseteq \sum^{m} (L_0 + L_0^{\bullet 2}) = \sum^{m} ([R, R]_1 + [R, R]_1 \bullet [R, R]_1).$$

(2). Using the assumption at the first step, and Lemma 3.3 at the second step, we get

$$R = \sum^{n} R \bullet [L_0, L_0^{\bullet 2}]_1 \subseteq \sum^{3n} L_0^{\bullet 2} = \sum^{3n} [R, R]_1 \bullet [R, R]_1. \quad \Box$$

We will need to know that matrix algebras over  $\mathbb{Z}$  contain certain special commutators.

**Lemma 5.14.** Let  $n \ge 2$ . Then the integral matrix ring  $M_n(\mathbb{Z})$  contains matrices  $u, v, w \in [M_n(\mathbb{Z}), M_n(\mathbb{Z})]_1$  such that [u, v] is invertible and v = vw.

**Proof.** Suppose n = 2. Set

$$u := \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} = \begin{bmatrix} \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \end{bmatrix} \quad v := \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} = \begin{bmatrix} \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \end{bmatrix}.$$

One checks that

$$[u,v] = \left[ \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \right] = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix},$$

which is invertible. Since one may also readily check that v = v[u, v], we may take w = [u, v].

Suppose n = 3. Consider

$$u := \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} = \begin{bmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \end{bmatrix},$$
$$v := \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} = \begin{bmatrix} \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}, \begin{pmatrix} 2 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix} \end{bmatrix}.$$

Then

$$[u,v] = \begin{bmatrix} \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix} = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & -1 \\ -1 & 0 & 0 \end{pmatrix}$$

is readily seen to be invertible. Moreover, setting

$$w := \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -2 \end{pmatrix} = \begin{bmatrix} \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 2 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \end{bmatrix},$$

one checks that vw = v as well.

Suppose that  $n \geq 4$ . Find  $k, \ell \in \mathbb{N}$  such that  $n = 2k + 3\ell$ . One can find the desired matrices in  $M_n(\mathbb{Z})$  as block-diagonal matrices using k blocks in  $M_2(\mathbb{Z})$  and  $\ell$  blocks in  $M_3(\mathbb{Z})$ . We omit the details.  $\Box$ 

**Theorem 5.15.** Let R be a unital ring, and suppose that there exist unital rings  $S_1, \ldots, S_m$ and natural numbers  $n_1, \ldots, n_m \ge 2$  such that  $\bigoplus_{j=1}^m M_{n_j}(S_j)$  embeds unitally into R. Then

$$R = [R, R]_1 + [R, R]_1 \bullet [R, R]_1, \quad and \quad R = \sum{}^3 [R, R]_1 \bullet [R, R]_1.$$

In particular,  $\xi(R) \leq 3$ .

**Proof.** Set  $S := \bigoplus_{j=1}^{m} M_{n_j}(S_j)$ . Without loss of generality, we assume that S is a subring of R. Applying Lemma 5.14 in each summand  $M_{n_j}(S_j)$  and adding them, we find  $a \in S$  and  $u, v, w \in [S, S]_1$  such that 1 = a[u, v] = a[u, vw]. The result now follows from Lemma 5.13.  $\Box$ 

**Question 5.16.** In the setting of Theorem 5.15, does one have  $\xi(R) \leq 2$ ?

**Remark 5.17.** Theorem 5.15 applies in particular to matrix rings and shows that for every unital ring S and  $n \ge 2$ , every element in  $M_n(S)$  is a sum of a commutator with a product of two commutators. Similarly, one can show that  $R = [R, R]_1 + [R, R]_1 \cdot [R, R]_1$  whenever R is a noncommutative division ring, or a semisimple ring without commutative summands.

## 6. Bounds on the number of commutators for $C^*$ -algebras

In this section, we prove estimates for the invariant  $\xi(A)$  in the case that A is a unital C\*-algebra. We show that  $\xi(A) \leq 3$  if A is properly infinite (Example 6.1), or has real rank zero (Theorem 6.3). We prove the estimate  $\xi(A) \leq 6$  whenever the Jiang-Su algebra  $\mathcal{Z}$  embeds unitally into A, in particular for all  $\mathcal{Z}$ -stable C\*-algebras; see Theorem 6.4.

A unital C\*-algebra A is said to be properly infinite if there exist mutually orthogonal projections  $p, q \in A$  and isometries  $s, t \in A$  satisfying  $ss^* = p$  and  $tt^* = q$ . Equivalently, and with  $\mathcal{O}_{\infty}$  denoting the infinite Cuntz algebra, there is a unital embedding  $\mathcal{O}_{\infty} \to A$ ; see [3, Proposition III.1.3.3].

**Example 6.1.** Let A be a unital, properly infinite  $C^*$ -algebra. Then

$$A = \sum^{2} [A, A]_{1} = [A, A]_{1} + [A, A]_{1} \bullet [A, A]_{1} = \sum^{3} [A, A]_{1} \bullet [A, A]_{1},$$

so in particular  $\xi(A) \leq 3$ . Indeed, the first equality was shown by Pop; see [37, Remark 3]. The two last equalities follow from Theorem 5.15 using that A admits a unital embedding of the Cuntz algebra  $\mathcal{O}_{\infty}$ , which in turn admits a unital embedding of  $M_2(\mathbb{C}) \oplus M_3(\mathbb{C})$ .

It would be interesting to compute  $\xi(A)$  for some particular cases of properly (or even purely) infinite C\*-algebras. For example:

Question 6.2. What is  $\xi(\mathcal{O}_{\infty})$ ?

For Cuntz algebras  $\mathcal{O}_n$  with finite n, a smaller upper bound can be given. Indeed, for  $n \geq 2$  it is well-known that  $\mathcal{O}_n$  contains a unital copy of  $M_n(\mathbb{C})$ . (Indeed, if  $s_1, \ldots, s_n$  denote the canonical isometries generating  $\mathcal{O}_n$ , one can readily check that  $\{s_j s_k^* : j, k = 1, \ldots, n\}$  is a system of matrix units, so it generates a (unital) copy of  $M_n(\mathbb{C})$ .) It follows that  $\mathcal{O}_n$  is isomorphic to the ring of *n*-by-*n* matrices over a unital ring, and therefore  $\xi(\mathcal{O}_n) \leq 2$  by Theorem 5.4.

The same argument applies to the  $L^p$ -versions  $\mathcal{O}_n^p$  of the Cuntz algebras introduced in [36] and studied in [17,12], thus giving  $\xi(\mathcal{O}_n^p) \leq 2$  for every  $p \in [1, \infty)$  and every natural number  $n \geq 2$ . We do not know whether  $\xi(\mathcal{O}_n^p) = 1$ .

A unital C\*-algebra is said to have *real rank zero* if the self-adjoint elements with finite spectrum are dense among all self-adjoint elements. See [3, Section V.3.2] for details.

**Theorem 6.3.** Let A be a unital C\*-algebra of real rank zero that admits no characters. Then  $\xi(A) \leq 3$  and

$$A = [A, A]_1 + [A, A]_1 \bullet [A, A]_1.$$

**Proof.** Set  $B := M_2(\mathbb{C}) \oplus M_3(\mathbb{C})$ . By [35, Proposition 5.7], the unit of A is weakly divisible of degree 2, that is, there exists a unital homomorphism  $B \to A$ ; see [35, Page 164, Lines 10ff]. It follows that A contains a unital copy of B, or one of its unital quotients  $M_2(\mathbb{C})$  or  $M_3(\mathbb{C})$ . Now the result follows from Theorem 5.15.  $\Box$ 

We now turn to C\*-algebras that admit an embedding of the Jiang-Su algebra  $\mathcal{Z}$ , which is a unital, simple, separable, nuclear C\*-algebra with unique tracial state and without nontrivial idempotents; see [41, Example 3.4.5]. There are plenty of C\*-algebras that contain  $\mathcal{Z}$  unitally but which do not contain any matrix subalgebras, so that Theorem 5.15 does not apply to them. One such example is given by the reduced group C\*-algebra  $C^*_{\lambda}(\mathbb{F}_n)$  of the free group; see Example 6.5.

We will need the so-called *dimension drop algebra*  $Z_{2,3}$ , which is given by

$$Z_{2,3} = \left\{ f \colon [0,1] \to M_2(\mathbb{C}) \otimes M_3(\mathbb{C}) \text{ continuous:} \begin{array}{l} f(0) \in M_2(\mathbb{C}) \otimes 1, \\ f(1) \in 1 \otimes M_3(\mathbb{C}) \end{array} \right\}.$$

It is known that  $\mathcal{Z}$  contains  $Z_{2,3}$  as a unital subalgebra.

A unital C\*-algebra is said to be Z-stable if A is \*-isomorphic to the C\*-algebraic tensor product  $A \otimes Z$ . Such algebras clearly admit a unital embedding of Z, and therefore of  $Z_{2,3}$ .

The Jiang-Su algebra and  $\mathcal{Z}$ -stable C\*-algebras play a crucial role in Elliott's classification program of simple, nuclear C\*-algebras. For a general overview, as well as the statement of the recent classification theorem and corresponding references, we refer the reader to Winter's ICM proceedings [45].

**Theorem 6.4.** Let A be a unital C\*-algebra that admits a unital \*-homomorphism  $Z_{2,3} \rightarrow A$ . Then  $\xi(A) \leq 6$ . This applies in particular to all unital,  $\mathcal{Z}$ -stable C\*-algebras.

**Proof.** We claim that there exist  $a, b \in Z_{2,3}$  and  $q, r, s, x, y, z \in [Z_{2,3}, Z_{2,3}]_1$  such that

$$1 = a[q, rs] + b[x, yz].$$

Once we prove this, the fact that  $\xi(A) \leq 6$  will follow from part (2) of Lemma 5.13 with n = 2.

Use Lemma 5.14 with n = 3 to find  $u, v, w \in [M_3(\mathbb{Z}), M_3(\mathbb{Z})]_1$  and  $e \in M_3(\mathbb{Z})$  satisfying  $1_3 = e[u, vw]$ . Define functions  $a, q, r, s \colon [0, 1] \to M_2(\mathbb{C}) \otimes M_3(\mathbb{C})$  by

$$a(t) = t^{\frac{1}{2}}(1_2 \otimes e), \quad q(t) = t^{\frac{1}{4}}(1_2 \otimes u), \quad r(t) = t^{\frac{1}{8}}(1_2 \otimes v), \text{ and } s(t) = t^{\frac{1}{8}}(1_2 \otimes w).$$

Denote by  $h \in \mathbb{Z}_{2,3}$  the function given by  $h(t) = t \mathbb{1}_6$ . One readily checks that a, q, r, s belong to  $\mathbb{Z}_{2,3}$ ; that q, r, s are commutators in  $\mathbb{Z}_{2,3}$ , and that h = a[q, rs].

Similarly, using Lemma 5.14 with n = 2, we find  $u', v', w' \in [M_2(\mathbb{Z}), M_2(\mathbb{Z})]_1$  and  $e' \in M_2(\mathbb{Z})$  satisfying  $1_2 = e'[u', v'w']$ . Define  $b, x, y, z \colon [0, 1] \to M_2(\mathbb{C}) \otimes M_3(\mathbb{C})$  by

$$b(t) = (1-t)^{\frac{1}{2}} (e' \otimes 1_3), \quad x(t) = (1-t)^{\frac{1}{4}} (u' \otimes 1_3),$$
  
$$y(t) = (1-t)^{\frac{1}{8}} (v' \otimes 1_3), \text{ and } z(t) = (1-t)^{\frac{1}{8}} (w' \otimes 1_3)$$

One readily checks that b, x, y, z belong to  $Z_{2,3}$ ; that x, y, z are commutators in  $Z_{2,3}$ , and that 1 - h = b[x, yz]. It follows that 1 = a[q, rs] + b[x, yz], as desired.  $\Box$ 

Although the primary interest in the theorem above is to  $\mathcal{Z}$ -stable C\*-algebras, it also gives information for certain reduced group C\*-algebras:

**Example 6.5.** Let G be a discrete group containing a nonabelian free group. Then  $\xi(C^*_{\lambda}(G)) \leq 6$ . Indeed, it was shown in [43, Proposition 4.2] that  $C^*_{\lambda}(\mathbb{F}_2)$  contains a unital copy of  $\mathcal{Z}$ , and the assumptions on G guarantee that  $C^*_{\lambda}(\mathbb{F}_2)$  embeds unitally into  $C^*_{\lambda}(G)$ . Now the assertion follows from Theorem 6.4.

The C\*-algebras here considered often contain no nontrivial idempotents, and hence no matrix subalgebras, for example for  $G = \mathbb{F}_n$ . In such situations, Theorem 5.15 is not applicable.

The methods we have developed to show that  $\xi(C^*_{\lambda}(\mathbb{F}_n)) \leq 6$  do not seem to give any information about the  $L^p$ -versions  $F^p_{\lambda}(\mathbb{F}_n)$  of these group algebras introduced by Herz in [26] and further studied, among others, in [18–20]. It would be interesting to find explicit upper bounds for  $\xi(F^p_{\lambda}(\mathbb{F}_n))$ .

## 7. Outlook

We end this paper with some questions for future work. Given a C\*-algebra A, consider the following properties:

- (1) We have A = A[A, A]A.
- (2) We have  $A = [A, A]^2$ .
- (3) We have  $[A, A] = [[A, A], [A, A]^2].$

- (4) We have  $A[A, A]A = [A, A]^2$ .
- (5) We have  $[A, A] \subseteq [A, A]^2$ .

Then the implications  $(1) \Leftrightarrow (2) \Rightarrow (3) \Rightarrow (4) \Rightarrow (5)$ ' hold. Indeed, (1) and (2) are equivalent by Theorem 4.1. To show that (2) implies (3), assume that  $A = [A, A]^2$ . Given  $a, b, c \in A$ , we have [ab, c] = [a, bc] + [b, ca] and thus

$$[A, A] = [[A, A]^2, A] \subseteq [[A, A], [A, A]A] + [[A, A], A[A, A]]$$
$$\subseteq [[A, A], A] \subseteq [[A, A], [A, A]^2].$$

To show that (3) implies (4), first note that the inclusion  $[A, A]^2 \subseteq A[A, A]A$  holds in general, since the ideal A[A, A]A contains [A.A] by [21, Proposition 3.2]. Further, assuming that  $[A, A] = [[A, A], [A, A]^2]$ , using Lemma 3.3 at the second step, we get

$$A[A, A]A = A[[A, A], [A, A]^2]A \subseteq [A, A]^2.$$

Finally, we see that (4) implies (5), again using that  $[A, A] \subseteq A[A, A]A$ .

If A is commutative, then (2) does not hold, while (3) is satisfied. Thus, (3) need not imply (2), but it remains unclear if (3)-(5) are equivalent. In fact, it is possible that (3)-(5) are always true.

Question 7.1. Let A be a C\*-algebra.

- (a) Do we have  $[A, A] = [[A, A], [A, A]^2]$ ?
- (b) Do we have  $A[A, A]A = [A, A]^2$ ?
- (c) Do we have  $[A, A] \subseteq [A, A]^2$ ?

For C\*-algebras that are generated by their commutators as an ideal (or more generally, C\*-algebras for which the commutator ideal is closed), all subquestions of Question 7.1 have a positive answer. Further, by passing to the closed ideal generated by commutators, one may assume that the C\*-algebra has no character. Thus, Question 7.1 is only unclear for C\*-algebras that have no character but are not generated by their commutators as an ideal, such as the one described in Example 4.4.

**Remark 7.2.** Assume that a C\*-algebra A satisfies  $[A, A] \subseteq [A, A]^2$ . We claim that the commutator ideal  $I := \widetilde{A}[A, A]\widetilde{A}$  is semiprime, that is, an intersection of prime ideals. To see this, we note that I is idempotent, that is,  $I = I^2$ . Indeed, the inclusion  $I^2 \subseteq I$  is clear, and conversely we have

$$I = \widetilde{A}[A, A]\widetilde{A} \subseteq \widetilde{A}[A, A]^2 \widetilde{A} \subseteq \widetilde{A}[A, A]\widetilde{A}[\widetilde{A}A, A]\widetilde{A} = I^2.$$

Since an ideal in a C\*-algebra is idempotent if and only if it is semiprime by [16, Theorem A], the claim follows.

### Data availability

No data was used for the research described in the article.

## References

- W.E. Baxter, Concerning the commutator subgroup of a ring, Proc. Am. Math. Soc. 16 (1965) 803–805.
- [2] T. Bice, I. Farah, Traces, ultrapowers and the Pedersen-Petersen C\*-algebras, Houst. J. Math. 41 (2015) 1175–1190.
- [3] B. Blackadar, Operator Algebras: Theory of C\*-Algebras and von Neumann Algebras, Operator Algebras and Non-commutative Geometry, III, Encyclopaedia of Mathematical Sciences, vol. 122, Springer-Verlag, Berlin, 2006.
- [4] J.D. Botha, Products of matrices with prescribed nullities and traces, Linear Algebra Appl. 252 (1997) 173–198.
- [5] M. Brešar, Introduction to Noncommutative Algebra, Universitext, Springer, Cham, 2014.
- [6] M. Brešar, Commutators and images of noncommutative polynomials, Adv. Math. 374 (2020) 107346.
- [7] M. Brešar, A. Fošner, M. Fošner, Jordan ideals revisited, Monatshefte Math. 145 (2005) 1–10.
- [8] M. Brešar, E. Gardella, H. Thiel, Products of commutators in matrix rings, Can. Math. Bull. (2024), in press.
- [9] M. Brešar, E. Kissin, V.S. Shulman, Lie ideals: from pure algebra to C\*-algebras, J. Reine Angew. Math. 623 (2008) 73–121.
- [10] M. Brešar, P. Šemrl, The Waring problem for matrix algebras, Isr. J. Math. 253 (2023) 381–405.
- [11] M. Brešar, P. Šemrl, The Waring problem for matrix algebras, II, Bull. Lond. Math. Soc. 55 (2023) 1880–1889.
- [12] Y. Choi, E. Gardella, H. Thiel, Rigidity results for L<sup>p</sup>-operator algebras and applications, Adv. Math. 452 (2024) 109747.
- [13] J. Cuntz, G.K. Pedersen, Equivalence and traces on C\*-algebras, J. Funct. Anal. 33 (1979) 135–164.
- [14] M.P.n. Eroğlu, On the subring generated by commutators, J. Algebra Appl. 21 (2022) 2250059.
- [15] E. Gardella, A modern look at algebras of operators on  $L^p$ -spaces, Expo. Math. 39 (2021) 420–453.
- [16] E. Gardella, K. Kitamura, H. Thiel, Semiprime ideals in C\*-algebras, preprint, arXiv:2311.17480 [math.OA], 2023.
- [17] E. Gardella, M. Lupini, Representations of étale groupoids on L<sup>p</sup>-spaces, Adv. Math. 318 (2017) 233–278.
- [18] E. Gardella, H. Thiel, Group algebras acting on L<sup>p</sup>-spaces, J. Fourier Anal. Appl. 21 (2015) 1310–1343.
- [19] E. Gardella, H. Thiel, Representations of p-convolution algebras on L<sup>q</sup>-spaces, Trans. Am. Math. Soc. 371 (2019) 2207–2236.
- [20] E. Gardella, H. Thiel, Isomorphisms of algebras of convolution operators, Ann. Sci. Éc. Norm. Supér. (4) 55 (2022) 1433–1471.
- [21] E. Gardella, H. Thiel, Prime ideals in C\*-algebras and applications to Lie theory, Proc. Am. Math. Soc. 152 (2024) 3647–3656.
- [22] E. Gardella, H. Thiel, Products of additive commutators in von Neumann algebras, 2024, in preparation.
- [23] B. Harris, Commutators in division rings, Proc. Am. Math. Soc. 9 (1958) 628-630.
- [24] I.N. Herstein, Topics in Ring Theory, The University of Chicago Press, Chicago, Ill.-London, 1969.
- [25] I.N. Herstein, On the Lie structure of an associative ring, J. Algebra 14 (1970) 561–571.
- [26] C. Herz, Harmonic synthesis for subgroups, Ann. Inst. Fourier (Grenoble) 23 (1973) 91–123.
- [27] A. Kanel-Belov, S. Malev, L. Rowen, R. Yavich, Evaluations of noncommutative polynomials on algebras: methods and problems, and the L'vov-Kaplansky conjecture, SIGMA 16 (2020) 071.
- [28] T.Y. Lam, A First Course in Noncommutative Rings, second ed., Graduate Texts in Mathematics, vol. 131, Springer-Verlag, New York, 2001.
- [29] C. Lanski, S. Montgomery, Lie structure of prime rings of characteristic 2, Pac. J. Math. 42 (1972) 117–136.
- [30] T.-K. Lee, Additive subgroups generated by noncommutative polynomials, Monatshefte Math. 199 (2022) 149–165.

- [31] L.W. Marcoux, Sums of small number of commutators, J. Oper. Theory 56 (2006) 111-142.
- [32] Z. Mesyan, Commutator rings, Bull. Aust. Math. Soc. 74 (2006) 279–288.
- [33] C.R. Miers, Closed Lie ideals in operator algebras, Can. J. Math. 33 (1981) 1271–1278.
- [34] G.K. Pedersen, N.H. Petersen, Ideals in a C\*-algebra, Math. Scand. 27 (1970) 193–204 (1971).
- [35] F. Perera, M. Rørdam, AF-embeddings into C\*-algebras of real rank zero, J. Funct. Anal. 217 (2004) 142–170.
- [36] N.C. Phillips, Analogs of Cuntz algebras on L<sup>p</sup> spaces, preprint, arXiv:1201.4196 [math.FA], 2012.
- [37] C. Pop, Finite sums of commutators, Proc. Am. Math. Soc. 130 (2002) 3039-3041.
- [38] L. Robert, Nuclear dimension and sums of commutators, Indiana Univ. Math. J. 64 (2015) 559–576.
- [39] L. Robert, On the Lie ideals of C\*-algebras, J. Oper. Theory 75 (2016) 387–408.
- [40] L. Robert, M. Rørdam, Divisibility properties for C\*-algebras, Proc. Lond. Math. Soc. (3) 106 (2013) 1330–1370.
- [41] M. Rørdam, Classification of nuclear, simple C\*-algebras, in: Classification of Nuclear C\*-Algebras. Entropy in Operator Algebras, in: Encyclopaedia Math. Sci., vol. 126, Springer, Berlin, 2002, pp. 1–145.
- [42] M. Rosset, S. Rosset, Elements of trace zero that are not commutators, Commun. Algebra 28 (2000) 3059–3072.
- [43] H. Thiel, W. Winter, The generator problem for Z-stable C\*-algebras, Trans. Am. Math. Soc. 366 (2014) 2327–2343.
- [44] D. Vitas, Multilinear polynomials are surjective on algebras with surjective inner derivations, J. Algebra 565 (2021) 255–281.
- [45] W. Winter, Structure of nuclear C\*-algebras: from quasidiagonality to classification and back again, in: Proceedings of the International Congress of Mathematicians—Rio de Janeiro 2018. Vol. III. Invited Lectures, World Sci. Publ., Hackensack, NJ, 2018, pp. 1801–1823.
- [46] P.Y. Wu, The operator factorization problems, Linear Algebra Appl. 117 (1989) 35–63.