



Splitting integrators for linear Vlasov equations with stochastic perturbations

Downloaded from: <https://research.chalmers.se>, 2025-12-08 23:25 UTC

Citation for the original published paper (version of record):

Bréhier, C., Cohen, D. (2024). Splitting integrators for linear Vlasov equations with stochastic perturbations. *Journal of Computational Dynamics*, 11(4): 494-532.
<http://dx.doi.org/10.3934/jcd.2024014>

N.B. When citing this work, cite the original published paper.



SPLITTING INTEGRATORS FOR LINEAR VLASOV EQUATIONS WITH STOCHASTIC PERTURBATIONS

CHARLES-EDOUARD BRÉHIER^{✉1} AND DAVID COHEN^{✉*2}

¹Universite de Pau et des Pays de l'Adour,
 E2S UPPA, CNRS, LMAP Pau, France

²Department of Mathematical Sciences Chalmers University of Technology,
 and University of Gothenburg, Gothenburg, Sweden

ABSTRACT. We consider a class of linear Vlasov partial differential equations driven by Wiener noise. Different types of stochastic perturbations are treated: additive noise, multiplicative Itô and Stratonovich noise, and transport noise. We propose to employ splitting integrators for the temporal discretization of these stochastic partial differential equations. These integrators are designed in order to preserve qualitative properties of the exact solutions depending on the stochastic perturbation, such as preservation of norms or positivity of the solutions. We provide numerical experiments in order to illustrate the properties of the proposed integrators and investigate mean-square rates of convergence.

1. Introduction. In this article, we are interested in stochastic perturbations of the linear Vlasov equation

$$\begin{cases} \partial_t f(t, x, v) + v \cdot \nabla_x f(t, x, v) + E(x) \cdot \nabla_v f(t, x, v) = 0, \\ f(0, x, v) = f_0(x, v), \end{cases}$$

see the next sections for the precise setting. Our objective is to identify some qualitative properties of the solutions and to propose temporal discretization schemes which are able to preserve those properties.

The Vlasov equation has been introduced in the literature in the middle of the 20th century [62], and has become a fundamental tool for the mathematical description of (collisionless) plasma in astrophysics and in plasma physics. It is common to interpret $f(t, x, v)$ as the density of particles having position x (assumed to take values in a torus) and velocity v at time t , starting from an initial configuration $f_0(x, v)$, and where the particles are transported by the ordinary differential equation

$$\begin{cases} \dot{x}_t = v_t, \\ \dot{v}_t = E(x_t). \end{cases}$$

The linear model above is a simple version, where the vector field E is imposed and is independent of time. In the Vlasov–Maxwell or the Vlasov–Poisson equations, for

2020 *Mathematics Subject Classification.* 35Q83. 60-08. 60H15. 60H35. 65C30. 65J08.

Key words and phrases. Stochastic partial differential equations, stochastic Vlasov equation, splitting scheme, trace formula, preservation properties, positivity-preserving scheme.

* Corresponding author: David Cohen.

example, E is not fixed and depends on the solution f , see for instance [6, 23, 54]. We refer to [61] for a review of kinetic models which take into account collisional effects, which may be linear (for instance Vlasov–Fokker–Planck, Bhatnagar–Gross–Krook equations) or nonlinear (Boltzmann equation) and to the monograph [42] for the analysis of kinetic partial differential equations (PDEs), including the Vlasov equation. Note that kinetic PDEs have also been used as popular models in mathematical biology in the recent years.

Let us mention the main qualitative properties satisfied by the linear Vlasov equations, we refer to Section 3 for details. If the initial value is nonnegative, then at all times the solution is also nonnegative. In addition, all L^p norms are preserved.

Stochastic versions of Vlasov kinetic equations have been considered in some recent works, for instance to model random injection or removal of particles in the system or to model the influence of stochastic (space-time) perturbations of the external vector field E . In [40] the authors prove regularization by noise results for a class of linear Vlasov equations with transport noise, and similar results are proved for nonlinear Vlasov–Poisson(–Fokker–Planck) equations in [4]. See also the article [34], where it is shown that introducing a stochastic (space-time) perturbation of the vector field E can prevent collapse in Vlasov–Poisson systems. Variational integrators are tested for some stochastic versions of the Vlasov equation in the article [47]. Finally, numerical methods are applied to some linear collisional kinetic equations with stochastic perturbation in the diffusion limit in the article [2]. We refer to [36] for a review of numerical methods applied to kinetic PDEs. The article [28] promotes the application of particle methods to approximate solutions of Vlasov equations. See also [41] for a description of various numerical approaches.

Our motivation in this work is driven by the perspective of geometric numerical integration and the desire to illustrate how the qualitative behavior of the solutions and of well-chosen integrators are modified under stochastic perturbations of various types. In addition, in this work we do not provide rigorous convergence analysis for the proposed integrators. To the best of our knowledge, the numerical schemes constructed below have not been studied previously in the literature.

The numerical schemes considered in this article combine two techniques which have been extensively studied for the discretization of the Vlasov equation: the semi-Lagrangian approach [59, 7, 38, 31, 8, 20] and the splitting technique [19, 39, 30, 56]. The lists of references above are not exhaustive. The semi-Lagrangian approach exploits characteristic curves to express the numerical solution at time t_{n+1} at grid points in terms of the numerical solution at time t_n , with the application of an interpolation procedure. The splitting approach consists in decomposing on each time interval the dynamics into subsystems which can be solved exactly. We briefly review semi-Lagrangian and splitting techniques when applied to the deterministic linear Vlasov equation in Section 3. In the sequel, we focus on the temporal discretization and we thus only deal with semi-discrete splitting methods.

Various types of stochastic perturbations of the linear Vlasov equation are considered. We illustrate below how the properties of the exact solutions differ for each type of perturbation, and how this needs to be taken into account in the construction of the splitting integrators. Basic concepts and results on splitting schemes, when applied to deterministic differential equations, are found for instance in the monographs [43, 10] and in the review articles [51, 11], see also references therein. For applications of splitting schemes to the temporal discretization of stochastic

partial differential equations, we refer the interested reader to the following (non-exhaustive) list: [50, 29, 3, 32, 53, 16, 17, 5, 13, 12, 14, 15]. In all the considered stochastic perturbations of the linear Vlasov equation, a similar strategy is applied to define the splitting schemes: the contributions of the deterministic and stochastic parts in the evolution are treated separately. In addition, the deterministic part is always treated by the same Lie–Trotter splitting scheme. For each type of noise the exact solution for the stochastic contribution is known exactly. Considering splitting schemes is appealing in our context since this leads to effective explicit numerical schemes which are able to preserve some qualitative properties of the exact solutions. This is justified theoretically and demonstrated numerically for all the stochastic versions of the linear Vlasov equation considered in this work. Let us now describe these versions and the main results. We refer to Section 2 below for the notation.

- *Additive noise perturbation:* in Section 4 we consider the following stochastic PDE

$$\begin{cases} df^{\text{add}}(t, x, v) + v \cdot \nabla_x f^{\text{add}}(t, x, v) dt \\ \quad + E(x) \cdot \nabla_v f^{\text{add}}(t, x, v) dt = dW(t, x, v) , \\ f^{\text{add}}(0, x, v) = f_0(x, v) , \end{cases}$$

see equation (15). The splitting scheme in this case is given by (19). Since the noise is additive, the solution is a Gaussian process, thus the positivity or the L^p norm of the initial value f_0 cannot be preserved. Instead, we prove trace formulas: the second-order moment of the L^2 norm of the exact solution grows linearly with time (see Proposition 1). This linear evolution is preserved for the numerical solution computed using the proposed splitting scheme (see Proposition 2).

- *Multiplicative noise perturbation with Itô interpretation:* in Section 5.1 we consider the following stochastic PDE

$$\begin{cases} df^{\text{mI}}(t, x, v) + v \cdot \nabla_x f^{\text{mI}}(t, x, v) dt + E(x) \cdot \nabla_v f^{\text{mI}}(t, x, v) dt \\ \quad = f^{\text{mI}}(t, x, v) dW(t, x, v) , \\ f^{\text{mI}}(0, x, v) = f_0(x, v) , \end{cases}$$

see equation (27). The splitting scheme in this case is given by (31). Proposition 3 provides the qualitative properties for the exact solution. First, one has almost sure preservation of the positivity of the initial value. Second, the mass of the initial value is preserved in expectation. Finally, under an appropriate condition on the diffusion coefficients, the second-order moment of the L^2 norm is shown to satisfy a simple evolution law. All those properties are shown to be preserved by the splitting scheme (31), see Proposition 5.

- *Multiplicative noise perturbation with Stratonovich interpretation:* in Section 5.2 we consider the following stochastic PDE

$$\begin{cases} df^{\text{mS}}(t, x, v) + v \cdot \nabla_x f^{\text{mS}}(t, x, v) dt \\ \quad + E(x) \cdot \nabla_v f^{\text{mS}}(t, x, v) dt = f^{\text{mS}}(t, x, v) \circ dW(t, x, v) , \\ f^{\text{mS}}(0, x, v) = f_0(x, v) , \end{cases}$$

see equation (33). The splitting scheme in this case is given by (38). Proposition 8 provides the qualitative properties for the exact solution. First, one has almost sure preservation of the positivity of the initial value. Moreover,

under an appropriate condition on the diffusion coefficients, the second-order moment of the L^2 norm is shown to satisfy a simple evolution law. All those properties are shown to be preserved by the splitting scheme (38), see Proposition 9.

- *Transport noise perturbation:* in Section 6 we consider the following stochastic PDE

$$\begin{cases} df^{\text{tr}}(t, x, v) + v \cdot \nabla_x f^{\text{tr}}(t, x, v) dt \\ \quad + E(x) \cdot \nabla_v f^{\text{tr}}(t, x, v) dt + \nabla_v f^{\text{tr}}(t, x, v) \odot d\mathbf{W}(t, x) = 0, \\ f^{\text{tr}}(0, x, v) = f_0(x, v), \end{cases}$$

see equation (39). The splitting scheme in this case is given by (44). Almost preservation of positivity and L^p norms are stated in Proposition 11 for the exact solution. We show that those properties are also preserved by the splitting scheme (44), see Proposition 12.

All the theoretical results mentioned above are illustrated by numerical experiments for each stochastic partial differential equation (SPDE). Snapshots of the numerical solutions also illustrate the influence of the stochastic perturbation on the behavior of the solutions. Even if we do not provide a rigorous convergence analysis, we conjecture that the proposed splitting schemes are consistent. Moreover, we report below numerical experiments to investigate mean-square rates of convergence and identify that in all cases convergence with order 1 is expected to hold. We leave a convergence analysis and other questions such as treating nonlinear models or constructing higher order integrators for future works. Our implementation is based on the code from [46] and is available under <https://doi.org/10.5281/zenodo.10495233>.

Sections 4, 5 and 6 below are organized similarly, this allows us to exhibit the main common or different features of the considered SPDEs and their numerical discretizations.

2. Notation. The dimension $d \in \mathbb{N}$ is an arbitrary integer. Let $\mathbb{T}^d = (\mathbb{R}/\mathbb{Z})^d$ denote the d -dimensional torus. For any differentiable mapping $f : \mathbb{T}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$, let $\nabla_x f = (\partial_{x_1} f, \dots, \partial_{x_d} f)$ and $\nabla_v f = (\partial_{v_1} f, \dots, \partial_{v_d} f)$ denote the gradients of f with respect to the spatial variable $x \in \mathbb{T}^d$ and the velocity $v \in \mathbb{R}^d$ respectively.

For any real number $p \in [1, \infty)$, and any measurable mapping $f : \mathbb{T}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$, define

$$\|f\|_{L^p_{x,v}} = \left(\iint |f(x, v)|^p dx dv \right)^{\frac{1}{p}} := \left(\int_{\mathbb{R}^d} \int_{\mathbb{T}^d} |f(x, v)|^p dx dv \right)^{\frac{1}{p}}$$

and set $L^p_{x,v} = \{f; \|f\|_{L^p_{x,v}} < \infty\}$. Below, to simplify the notation we will not write the domains of the integrals. In addition, for any measurable mapping $f : \mathbb{T}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$, define

$$\|f\|_{L^\infty_{x,v}} = \operatorname{ess\,sup}_{(x,v) \in \mathbb{T}^d \times \mathbb{R}^d} |f(x, v)|$$

and set $L^\infty_{x,v} = \{f; \|f\|_{L^\infty_{x,v}} < \infty\}$.

Given a function $f : (t, x, v) \in \mathbb{R}^+ \times \mathbb{T}^d \times \mathbb{R}^d \mapsto f(t, x, v) \in \mathbb{R}$, the notation $f(t)$ for $t \geq 0$ is frequently used in the sequel to denote the mapping $(x, v) \in \mathbb{T}^d \times \mathbb{R}^d \mapsto f(t, x, v)$. The expression $f(t)(x, v) = f(t, x, v)$ is also often employed below.

To describe the considered class of (stochastic) partial differential equations, let us denote by $E : x \in \mathbb{T}^d \mapsto E(x) \in \mathbb{R}^d$ a vector field which is assumed to be of class C^∞ . In addition, the initial value is denoted by $f_0 : (x, v) \in \mathbb{T}^d \times \mathbb{R}^d \mapsto f_0(x, v) \in \mathbb{R}$

in the sequel. Below, it is assumed that f_0 is non-random. Precise regularity and integrability conditions on f_0 are imposed below.

It remains to describe the setting for the considered stochastic perturbation of linear Vlasov equations. Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space and $(\mathcal{F}_t)_{t \geq 0}$ be a filtration satisfying the usual conditions. The expectation operator is denoted by $\mathbb{E}[\cdot]$. Let $K \in \mathbb{N}$ be an integer, and let $(\beta_k)_{1 \leq k \leq K}$ be a family of independent standard real-valued Wiener processes, adapted to the filtration.

In Sections 4 and 5, the stochastic perturbation is of additive or multiplicative type, and is written as a real-valued Wiener process defined by

$$W(t, x, v) = \sum_{k=1}^K \beta_k(t) \sigma_k(x, v), \quad t \geq 0, x \in \mathbb{T}^d, v \in \mathbb{R}^d, \quad (1)$$

where $\sigma_k : (x, v) \in \mathbb{T}^d \times \mathbb{R}^d \mapsto \sigma_k(x, v) \in \mathbb{R}$ are real-valued mappings, for $1 \leq k \leq K$.

In Section 6, the stochastic perturbation is of transport type and it does not depend on of the variable v . It is written as a \mathbb{R}^d -valued Wiener process defined by

$$\mathbf{W}(t, x) = \sum_{k=1}^K \beta_k(t) \boldsymbol{\sigma}_k(x), \quad t \geq 0, x \in \mathbb{T}^d, \quad (2)$$

where $\sigma_{j,k} : x \in \mathbb{T}^d \mapsto \sigma_{j,k}(x) \in \mathbb{R}$ are real-valued mappings, and for all $x \in \mathbb{T}^d$ one has $\boldsymbol{\sigma}_k(x) = (\sigma_{1,k}(x), \dots, \sigma_{d,k}(x)) \in \mathbb{R}^d$.

The notation $W(t)$ is also used below for the random mapping $(x, v) \in \mathbb{T}^d \times \mathbb{R}^d \mapsto W(t, x, v)$, associated with (1) while the notation $\mathbf{W}(t)$ stands for the random mapping $x \in \mathbb{T}^d \mapsto \mathbf{W}(t, x)$, associated with (2) respectively, for all $t \geq 0$. We refer for instance to [33, Chapter 4] and [49, Chapter 10] for details on Wiener processes and stochastic integrals with values in Hilbert spaces.

It is assumed that the mappings σ_k and $\sigma_{j,k}$, $1 \leq j \leq d$, $1 \leq k \leq K$ are of class \mathcal{C}^∞ . Further growth or integrability conditions on these mappings are imposed below depending on the considered class of problems.

Finally, the numerical methods considered in this paper use the following notation. For the temporal discretization, the time-step size is denoted by τ . Without loss of generality, it is assumed that $\tau \in (0, 1)$. For any nonnegative integer $n \geq 0$, set $t_n = n\tau$, and introduce Wiener increments

$$\delta\beta_{k,n} = \beta_k(t_{n+1}) - \beta_k(t_n), \quad 1 \leq k \leq K. \quad (3)$$

The random variables $(\tau^{-\frac{1}{2}} \delta\beta_{n,k})_{n \geq 0, 1 \leq k \leq K}$ are independent standard real-valued Gaussian random variables (with mean equal to 0 and variance equal to 1).

Increments of the Wiener processes defined by (1) and (2) are given for any nonnegative integer $n \geq 0$ by

$$\begin{aligned} \delta W_n(x, v) &= W(t_{n+1}, x, v) - W(t_n, x, v) \\ &= \sum_{k=1}^K \delta\beta_{k,n} \sigma_k(x, v), \quad (x, v) \in \mathbb{T}^d \times \mathbb{R}^d, \end{aligned} \quad (4)$$

$$\delta \mathbf{W}_n(x) = \mathbf{W}(t_{n+1}, x) - \mathbf{W}(t_n, x) = \sum_{k=1}^K \delta\beta_{k,n} \boldsymbol{\sigma}_k(x), \quad x \in \mathbb{T}^d. \quad (5)$$

In this work, we focus on the temporal discretization of stochastic linear Vlasov equations. However, practical implementation requires some discretization procedure with respect to the spatial and velocity variables $x \in \mathbb{T}^d$ and $v \in \mathbb{R}^d$. In

the numerical experiments below, a semi-Lagrangian approach is employed, see for instance [59, 7, 38] or [37, Chapter 6]. The semi-Lagrangian approach exploits characteristic curves to express the numerical solution at time t_{n+1} at grid points in terms of the numerical solution at time t_n . Since the characteristic curves in general do not hit grid points, an interpolation procedure is employed to the numerical solution at time t_n . Note that, in order to preserve the positivity property of solutions to the considered SPDEs (see below), we use a linear interpolation in the implementation of the studied numerical schemes. However the analysis of semi-Lagrangian discretization is out of the scope of this work. The mesh sizes with respect to the spatial and velocity domains are denoted by δx and δv respectively. Since the velocity space \mathbb{R}^d is unbounded, in practice a truncation procedure at large velocities v is imposed, whereas for the spatial variable x , periodic boundary conditions are used. Besides, the verification of trace formulas and evolution laws for some moments of the solutions and the study of mean-square convergence require a Monte Carlo averaging procedure in order to compute approximate values of expectations.

3. Preliminaries on the deterministic linear Vlasov equation. The objective of this section is to provide basic and well-known background on the deterministic version of the linear Vlasov equation, before considering stochastically perturbed versions in the next sections. Some notation and fundamental properties introduced below are employed in the sequel. We study the linear Vlasov equation

$$\begin{cases} \partial_t f^{\text{det}}(t, x, v) + v \cdot \nabla_x f^{\text{det}}(t, x, v) + E(x) \cdot \nabla_v f^{\text{det}}(t, x, v) = 0, \\ t \geq 0, x \in \mathbb{T}^d, v \in \mathbb{R}^d, \\ f^{\text{det}}(0, x, v) = f_0(x, v), \quad x \in \mathbb{T}^d, v \in \mathbb{R}^d, \end{cases} \quad (6)$$

where the unknown is a mapping $f^{\text{det}} : (t, x, v) \in \mathbb{R}^+ \times \mathbb{T}^d \times \mathbb{R}^d \mapsto f^{\text{det}}(t, x, v) \in \mathbb{R}$. Under appropriate regularity and integrability conditions, given any initial value f_0 , the partial differential equation (6) admits a unique solution.

3.1. Analysis and properties of the problem. The solution of (6) can be constructed using solutions of the associated ordinary differential equation

$$\begin{cases} \dot{x}_t = v_t \\ \dot{v}_t = E(x_t). \end{cases} \quad (7)$$

Let $(\phi_t)_{t \in \mathbb{R}}$ denote the associated flow of this differential equation. Recall that this means that for any initial value $(x_0, v_0) \in \mathbb{T}^d \times \mathbb{R}^d$, the solution at time $t \geq 0$ is given by $(x_t, v_t) = \phi_t(x_0, v_0)$. In addition, the flow map property $\phi_{t+s} = \phi_t \circ \phi_s$ is satisfied for all $t, s \in \mathbb{R}$. Finally, for all $t \in \mathbb{R}$, the mapping ϕ_t is a smooth diffeomorphism and one has $(\phi_t)^{-1} = \phi_{-t}$. Since the vector field $(x, v) : \mathbb{T}^d \times \mathbb{R}^d \mapsto (v, E(x)) \in \mathbb{R}^d \times \mathbb{R}^d$ does not depend on the time variable, $(\phi_t^{-1})_{t \in \mathbb{R}} = (\phi_{-t})_{t \in \mathbb{R}}$ is obtained by reverting time in the dynamics (7), i. e. by considering the flow of the ordinary differential equation

$$\begin{cases} \dot{x}_t = -v_t \\ \dot{v}_t = -E(x_t). \end{cases}$$

The flow $(\phi_t)_{t \in \mathbb{R}}$ associated with (7) satisfies a remarkable property: it preserves the volume in $\mathbb{T}^d \times \mathbb{R}^d$. This can be written as follows: for any integrable function

$\varphi : \mathbb{T}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$, one has

$$\iint \varphi(\phi_t(x, v)) \, dx \, dv = \iint \varphi(x, v) \, dx \, dv.$$

We are now in position to recall the links between the Vlasov PDE (6) and the ODE system (7). On the one hand, assume that $(f^{\det}(t))_{t \geq 0}$ is solution of the PDE (6), then for any solution $(x_t, v_t)_{t \geq 0}$ of the ODE system (7), applying the chain rule, one has

$$\frac{d f^{\det}(t, x_t, v_t)}{dt} = 0.$$

As a consequence, for all $t \geq 0$, $x_0 \in \mathbb{T}^d$ and $v_0 \in \mathbb{R}^d$, one has

$$f^{\det}(t, \phi_t(x_0, v_0)) = f^{\det}(t, x_t, v_t) = f^{\det}(0, x_0, v_0) = f_0(x_0, v_0).$$

Therefore the ODE system (7) provides characteristic curves for the Vlasov equation (6). This gives a strategy to solve (6) by the method of lines: the solution $f(t)$ at any time $t \geq 0$ is given by

$$f^{\det}(t, x, v) = f_0(\phi_t^{-1}(x, v)), \quad x \in \mathbb{T}^d, v \in \mathbb{R}^d. \quad (8)$$

On the other hand, under sufficient regularity conditions, one can check that the mapping $f^{\det} : (t, x, v) \in \mathbb{R}^+ \times \mathbb{T}^d \times \mathbb{R}^d \mapsto f^{\det}(t, x, v) \in \mathbb{R}$ defined by the expression (8) solves the Vlasov equation (6).

The expression (8) leads to define a group of linear operators $(S(t))_{t \geq 0}$ as follows: for all $t \in \mathbb{R}$ and any measurable mapping $f : \mathbb{T}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$, set

$$S(t)f = f(\phi_t^{-1}(\cdot)). \quad (9)$$

The group property $S(t+s) = S(t)S(s)$, $t, s \in \mathbb{R}$, follows from the group property of the flow $(\phi_t)_{t \in \mathbb{R}}$.

Let us describe some remarkable properties of solutions of the Vlasov equation (6), which are considered below for stochastic perturbations of this PDE and for its numerical discretization. Let $f^{\det}(t) = S(t)f_0$ for all $t \geq 0$.

- *Preservation of positivity.* Assume that $f_0(x, v) \geq 0$ for all $(x, v) \in \mathbb{T}^d \times \mathbb{R}^d$. Then one has $f^{\det}(t, x, v) \geq 0$ for all $t \geq 0$ and $(x, v) \in \mathbb{T}^d \times \mathbb{R}^d$.
- *Preservation of integrals.* Let $\Phi : \mathbb{R} \rightarrow \mathbb{R}^+$ be a real-valued measurable mapping. Then for all $t \geq 0$ one has

$$\iint \Phi(f^{\det}(t, x, v)) \, dx \, dv = \iint \Phi(f_0(x, v)) \, dx \, dv.$$

- *Isometry property.* For all $p \in [1, \infty]$ and $t \geq 0$, the linear operator $S(t) : L_{x,v}^p \rightarrow L_{x,v}^p$ is an isometry: if $f_0 \in L_{x,v}^p$, then $f^{\det}(t) = S(t)f_0 \in L_{x,v}^p$ for all $t \geq 0$ and

$$\|f^{\det}(t)\|_{L_{x,v}^p} = \|S(t)f_0\|_{L_{x,v}^p} = \|f_0\|_{L_{x,v}^p}.$$

The preservation of positivity and the preservation of the $L_{x,v}^\infty$ norm are straightforward consequences of the expression (8). The preservation of integrals property follows from the preservation of volume by the flow $(\phi_t)_{t \in \mathbb{R}}$ and implies the isometry property above when $p \in [1, \infty)$ by choosing $\Phi = |\cdot|^p$.

Finally, the preservation of volume by the flow $(\phi_t)_{t \geq 0}$ also implies the following result. Assume that $f_0 \in L_{x,v}^1$, then for any bounded and continuous function $\varphi : \mathbb{T}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$ and for all $t \geq 0$, one has

$$\iint f^{\det}(t, x, v) \varphi(x, v) \, dx \, dv = \iint f(0, x, v) \varphi(\phi_t(x, v)) \, dx \, dv.$$

The properties above provide a possible probabilistic interpretation of solutions to the Vlasov PDE (6). If the initial value f_0 is a probability density function, in particular, this requires the conditions $f_0 \geq 0$ and $\|f_0\|_{L^1_{x,v}} = 1$, then for all $t \geq 0$, the mapping $f^{\det}(t) = S(t)f_0$ is a probability density function. If $(\mathcal{X}_0, \mathcal{V}_0)$ is a $\mathbb{T}^d \times \mathbb{R}^d$ -valued random variable having density f_0 with respect to the Lebesgue measure, then $f^{\det}(t)$ is the density with respect to the Lebesgue measure of the random variable $(\mathcal{X}_t, \mathcal{V}_t) = \phi_t(\mathcal{X}_0, \mathcal{V}_0)$. In other words, $f^{\det}(t)$ is the probability density function associated with the transport by the dynamics (7) of the distribution f_0 of the initial values (x_0, v_0) . In some contexts, in particular for Hamiltonian dynamics ($E = -\nabla V$ where $V : \mathbb{T}^d \rightarrow \mathbb{R}$ is a smooth mapping), the Vlasov equation (6) is referred to as the Liouville equation associated with the dynamics (7).

3.2. Numerical approximation. Let us describe numerical integrators applied to the linear Vlasov equation (6). As already mentioned, we focus on temporal discretization only in this work. To approximate solutions and preserve the properties mentioned above, it is natural to rely on splitting integrators. We only present Lie–Trotter versions, for simplicity and motivated by the fact that stochastic perturbations usually lead to numerical methods which are of strong order less than 2 (the usual order of Strang splitting for deterministic problems, see for instance [22, 59, 39, 19]). Let us mention that high-order numerical schemes for the deterministic part of the problem (for instance using a Strang splitting instead of a Lie–Trotter splitting) could however be of interest for stochastic problems with small noise, see for instance [52, Chapter 4].

The principle of splitting integrators is to combine solutions of subsystems of evolution equations which can be solved exactly. For the deterministic linear Vlasov equation (6) it is natural to decompose the problem into the two subsystems

$$\partial_t f^1(t, x, v) + v \cdot \nabla_x f^1(t, x, v) = 0, \quad t \geq 0, x \in \mathbb{T}^d, v \in \mathbb{R}^d \quad (10a)$$

$$\partial_t f^2(t, x, v) + E(x) \cdot \nabla_v f^2(t, x, v) = 0, \quad t \geq 0, x \in \mathbb{T}^d, v \in \mathbb{R}^d. \quad (10b)$$

Let $(S^1(t))_{t \in \mathbb{R}}$ and $(S^2(t))_{t \in \mathbb{R}}$ be the associated groups of linear operators: the solutions at time $t \geq 0$ of (10a) and (10b) are respectively $f^1(t) = S^1(t)f_0$ and $f^2(t) = S^2(t)f_0$, where for all $t \in \mathbb{R}$, $x \in \mathbb{T}^d$, $v \in \mathbb{R}^d$ and any measurable mapping $f : \mathbb{T}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$, one has

$$S^1(t)f(x, v) = f(x - tv, v)$$

$$S^2(t)f(x, v) = f(x, v - tE(x)).$$

Note that the partial differential equations (10a) and (10b) are associated with the ordinary differential equations

$$\begin{cases} \dot{x}_t^1 = v_t^1 \\ \dot{v}_t^1 = 0, \end{cases} \quad \text{and} \quad \begin{cases} \dot{x}_t^2 = 0 \\ \dot{v}_t^2 = E(x_t^2), \end{cases}$$

for which expressions of the exact solutions are known: for all $t \geq 0$, one has

$$\begin{aligned} (x_t^1, v_t^1) &= \phi_t^1(x_0^1, v_0^1) = (x_0^1 + tv_0^1, v_0^1) \\ (x_t^2, v_t^2) &= \phi_t^2(x_0^2, v_0^2) = (x_0^2, v_0^2 + tE(x_0^2)). \end{aligned}$$

The link between the Vlasov equation (6) and the dynamics (7) is retrieved for the subsystems (10a) and (10b) and the associated dynamics: for all $t \geq 0$ and any

measurable mapping $f : \mathbb{T}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$ one has

$$S^1(t)f = f((\phi_t^1)^{-1}(\cdot)) , \quad S^2(t)f = f((\phi_t^2)^{-1}(\cdot)) \quad (11)$$

where $(\phi_t^1)_{t \in \mathbb{R}}$ and $(\phi_t^2)_{t \in \mathbb{R}}$ are the flows associated with the dynamics introduced above.

As a consequence, it is straightforward to check that the properties stated above for solutions of the Vlasov equation (6) (preservation of positivity, preservation of integrals and isometry property) and the interpretation of the solution as the probability density functions associated with the transport by dynamics of the distribution of initial value, persist for the solutions of the subsystems (10a) and (10b).

Applying the Lie–Trotter splitting method yields to the definition of the following numerical method for the deterministic linear Vlasov equation (6): set $f_0^{\det} = f_0$ and for any nonnegative integer $n \geq 0$, set

$$f_{n+1}^{\det} = S^2(\tau)S^1(\tau)f_n^{\det}. \quad (12)$$

Using the definitions (11) of $S^1(t)$ and $S^2(t)$ above, one has for all $n \geq 0$, $x \in \mathbb{T}^d$ and $v \in \mathbb{R}^d$

$$\begin{aligned} f_{n+1}^{\det}(x, v) &= (S^2(\tau)S^1(\tau)f_n^{\det})(x, v) = (S^1(\tau)f_n^{\det})(x, v - \tau E(x)) \\ &= f_n^{\det}(x - \tau v - \tau^2 E(x), v - \tau E(x)). \end{aligned}$$

With that expression, it appears that the scheme (12) can be interpreted as a discrete version of the expression (8) for the exact solution of (6), where the flow ϕ_t is approximated using a splitting integrator based on the two subsystems of ordinary differential equations above.

It is straightforward to check that the properties stated above for the exact solution of (6) also hold for the Lie–Trotter splitting scheme (12), for any value of the time step size τ . In particular, one has the following results.

- *Preservation of positivity.* Assume that $f_0(x, v) \geq 0$ for all $(x, v) \in \mathbb{T}^d \times \mathbb{R}^d$. Then for all $n \in \mathbb{N}$ one has $f_n^{\det}(x, v) \geq 0$ for all $(x, v) \in \mathbb{T}^d \times \mathbb{R}^d$.
- *Isometry property.* Let $p \in [1, \infty]$ and assume that $f_0 \in L_{x,v}^p$. Then for all $n \geq 0$ one has $f_n^{\det} \in L_{x,v}^p$ and

$$\|f_n^{\det}\|_{L_{x,v}^p} = \|f_0\|_{L_{x,v}^p}.$$

We conclude these preliminaries on the deterministic linear Vlasov equation (6) with a numerical experiment with the goal to illustrate the behavior of the solution to the linear Vlasov PDE (6), in dimension $d = 1$. In Figure 1 below, snapshots at times $\{0, 0.5, 1, 1.5, 2, 2.5\}$ of the numerical solution computed using the Lie–Trotter splitting scheme 12 are displayed. We consider a standard two-stream instability test case: the initial value f_0 is given by

$$f_0(x, v) = \frac{e^{-v^2/2}}{\sqrt{2\pi}} (1 + 0.05 \cos(2\pi x)) v^2, \quad x \in \mathbb{T}, v \in \mathbb{R}, \quad (13)$$

The two-stream instability has been extensively used to illustrate wave-particle interactions since the seminal paper [35], see also for instance [9, 45, 22]. In addition, the vector field E is given by

$$E(x) = \cos(2\pi x), \quad x \in \mathbb{T}. \quad (14)$$

The discretization parameters for Figure 1 are $\delta x = \frac{1}{500}$, $\delta v = \frac{4\pi}{500}$, and $\tau = 0.1$.




FIGURE 1. Snapshots: approximation of the solution of the deterministic PDE (6) with initial value f_0 given by (13), at times $\{0, 0.5, 1, 1.5, 2, 2.5\}$, using the Lie-Trotter splitting scheme (12) with time-step size $\tau = 0.1$.

4. The stochastic linear Vlasov equation perturbed by additive noise. In this section, we consider a version of the linear Vlasov equation (6) perturbed by additive noise of the type (1): for $t \geq 0$, $x \in \mathbb{T}^d$, $v \in \mathbb{R}^d$

$$\begin{cases} df^{\text{add}}(t, x, v) + v \cdot \nabla_x f^{\text{add}}(t, x, v) dt + E(x) \cdot \nabla_v f^{\text{add}}(t, x, v) dt = dW(t, x, v), \\ f^{\text{add}}(0, x, v) = f_0(x, v). \end{cases} \quad (15)$$

4.1. Analysis and properties of the problem. We recall that the initial value f_0 is assumed to be non-random. The solution of the stochastic partial differential equation (15) can be interpreted in different ways, as explained below. Those interpretations are equivalent under appropriate regularity and integrability conditions.

On the one hand, using the group $(S(t))_{t \in \mathbb{R}}$ of linear operators given by (9) in Section 3 and the definition of the noise (1), one can consider mild solutions in the sense of [33, Chapter 5], see also [49, Section 10.4] for instance: for all $t \geq 0$, one has

$$f^{\text{add}}(t) = S(t)f_0 + \int_0^t S(t-s) dW(s) = S(t)f_0 + \sum_{k=1}^K \int_0^t S(t-s) \sigma_k d\beta_k(s), \quad (16)$$

where $\int_0^t S(t-s) dW(s)$ is considered as a stochastic integral with values in the Hilbert space $L^2_{x,v}$, see [33, Chapter 4]. On the other hand, using the expression (9) for the linear operator $S(t)$, one has for all $t \geq 0$, $x \in \mathbb{T}^d$ and $v \in \mathbb{R}^d$

$$f^{\text{add}}(t, x, v) = f_0(\phi_t^{-1}(x, v)) + \sum_{k=1}^K \int_0^t \sigma_k(\phi_{t-s}^{-1}(x, v)) d\beta_k(s), \quad (17)$$

where $(\phi_t)_{t \in \mathbb{R}}$ denotes the flow of (7).

Finally note that the expression above can be retrieved by applying the Itô–Wentzell formula (see [Appendix A](#) for the statement): if $t \mapsto (x_t, v_t) = \phi_t(x_0, v_0)$ is the solution of the ordinary differential equation (7) with arbitrary initial value $(x_0, v_0) \in \mathbb{T}^d \times \mathbb{R}^d$, and if the solution of (15) is sufficiently regular, then the stochastic process $t \in \mathbb{R}^+ \mapsto f^{\text{add}}(t, x_t, v_t)$ satisfies

$$\begin{aligned} df^{\text{add}}(t, x_t, v_t) &= \sum_{k=1}^K \sigma_k(x_t, v_t) d\beta_k(t) \\ f^{\text{add}}(t, x_t, v_t) &= f_0(x_0, v_0) + \sum_{k=1}^K \int_0^t \sigma_k(x_s, v_s) d\beta_k(s). \end{aligned}$$

Let us now study properties of the solution to the SPDE (15). First, note that the random field $(t, x, v) \in \mathbb{R}^+ \times \mathbb{T}^d \times \mathbb{R}^d \mapsto f^{\text{add}}(t, x, v)$ is Gaussian. Owing to (17) one has $\mathbb{E}[f^{\text{add}}(t, x, v)] = f_0(\phi_t^{-1}(x, v)) = f^{\text{det}}(t, x, v)$ for all $t \geq 0$, $x \in \mathbb{T}^d$ and $v \in \mathbb{R}^d$, where $(f^{\text{det}}(t))_{t \geq 0}$ is the solution of deterministic linear Vlasov equation (6) with initial value f_0 . The covariance structure of the Gaussian process $\mathbb{E}[f^{\text{add}}(t_1, x_1, v_1) f^{\text{add}}(t_2, x_2, v_2)]$ is computed using Itô’s isometry formula applied to (17): for all $t_1, t_2 \geq 0$, $x_1, x_2 \in \mathbb{T}^d$ and $v_1, v_2 \in \mathbb{R}^d$ one has

$$\begin{aligned} \mathbb{E}[f^{\text{add}}(t_1, x_1, v_1) f^{\text{add}}(t_2, x_2, v_2)] &= f_0(\phi_{t_1}^{-1}(x_1, v_1)) f_0(\phi_{t_2}^{-1}(x_2, v_2)) \\ &+ \sum_{k=1}^K \int_0^{\min(t_1, t_2)} \sigma_k(\phi_{t_1-s}^{-1}(x_1, v_1)) \sigma_k(\phi_{t_2-s}^{-1}(x_2, v_2)) ds. \end{aligned}$$

Since the solution to the SPDE (15) is a Gaussian process, the preservation properties satisfied in the deterministic case cannot be satisfied. However, one has the following remarkable property, which is often called a *trace formula* in the literature, see for instance [60, 44, 18, 58, 57, 26, 1, 25, 21, 27, 13].

Proposition 1. *Assume that $\sigma_k \in L^2_{x,v}$ for all $1 \leq k \leq K$, and that $f_0 \in L^2_{x,v}$. Let $(f^{\text{add}}(t))_{t \geq 0}$ be the solution of the SPDE (15) driven by the additive noise (1). Then for all $t \geq 0$ one has $f^{\text{add}}(t) \in L^2(\Omega, L^2_{x,v})$ and*

$$\mathbb{E}[\|f^{\text{add}}(t)\|_{L^2_{x,v}}^2] = \|f_0\|_{L^2_{x,v}}^2 + t \sum_{k=1}^K \|\sigma_k\|_{L^2_{x,v}}^2. \quad (18)$$

The trace formula (18) is proved below using the two formulations (16) and (17).

First proof of Proposition 1. Choosing $t_1 = t_2 = t$, $x_1 = x_2 = x$ and $v_1 = v_2 = v$ in the expression of the covariance above, and integrating with respect to the variables x and v , one obtains

$$\begin{aligned} \mathbb{E}[\|f^{\text{add}}(t)\|_{L^2_{x,v}}^2] &= \iint \mathbb{E}[f^{\text{add}}(t, x, v)^2] dx dv \\ &= \iint f_0(\phi_t^{-1}(x, v))^2 dx dv + \sum_{k=1}^K \int_0^t \iint \mathbb{E}[\sigma_k(\phi_{t-s}^{-1}(x, v))^2] dx dv ds \end{aligned}$$

$$\begin{aligned}
&= \iint f_0(x, v)^2 \, dx \, dv + \sum_{k=1}^K \int_0^t \iint \mathbb{E}[\sigma_k(x, v)^2] \, dx \, dv \, ds \\
&= \|f_0\|_{L_{x,v}}^2 + t \sum_{k=1}^K \|\sigma_k\|_{L_{x,v}^2}^2
\end{aligned}$$

using the fact that for all $t \geq 0$ the mapping ϕ_t preserves volume in $\mathbb{T}^d \times \mathbb{R}^d$, see Section 3. \square

Second proof of Proposition 1. Using the mild formulation (16) of the solution $f(t)$ of (15), the Itô isometry formula in the Hilbert space $L_{x,v}^2$ (see for instance [49, Theorem 10.16]), and the isometry property for the linear operators $S(t) : L_{x,v}^2 \rightarrow L_{x,v}^2$ (see Section 3), for all $t \geq 0$, one has

$$\begin{aligned}
\mathbb{E}[\|f^{\text{add}}(t)\|_{L_{x,v}^2}^2] &= \mathbb{E}[\|S(t)f_0\|_{L_{x,v}^2}^2] + \sum_{k=1}^K \int_0^t \|S(t-s)\sigma_k\|_{L_{x,v}^2}^2 \, ds \\
&= \mathbb{E}[\|f_0\|_{L_{x,v}^2}^2] + \sum_{k=1}^K \int_0^t \|\sigma_k\|_{L_{x,v}^2}^2 \, ds \\
&= \mathbb{E}[\|f_0\|_{L_{x,v}^2}^2] + t \sum_{k=1}^K \|\sigma_k\|_{L_{x,v}^2}^2.
\end{aligned}$$

\square

4.2. Splitting scheme. Let us now describe the proposed numerical scheme for the temporal discretization of the SPDE (15). The strategy is to use a splitting method in order to treat first the deterministic part, second the stochastic part. Using the Lie–Trotter integrator (12) for the deterministic part yields the following scheme: given the initial value f_0 and the time-step size $\tau \in (0, 1)$, set $f_0^{\text{add}} = f_0$ and for any nonnegative integer $n \geq 0$ set

$$f_{n+1}^{\text{add}} = S^2(\tau)S^1(\tau)f_n^{\text{add}} + \delta W_n = S^2(\tau)S^1(\tau)f_n^{\text{add}} + \sum_{k=1}^K \delta\beta_{n,k}\sigma_k, \quad (19)$$

where the Wiener increments δW_n and $\delta\beta_{n,k}$ are given by (4) and (3) respectively, see Section 2. The scheme (19) can also be written as follows: for all $n \geq 0$, $x \in \mathbb{T}^d$ and $v \in \mathbb{R}^d$ one has

$$f_{n+1}^{\text{add}}(x, v) = f_n^{\text{add}}(x - \tau v - \tau^2 E(x), v - \tau E(x)) + \sum_{k=1}^K \sigma_k(x, v) \delta\beta_{n,k}. \quad (20)$$

The main result of this section states that the Lie–Trotter splitting scheme (19) preserves the trace formula from Proposition 1 for all times and for any value of the time-step size τ .

Proposition 2. Assume that $\sigma_k \in L_{x,v}^2$ for all $1 \leq k \leq K$, and that $f_0 \in L_{x,v}^2$. Let $(f_n^{\text{add}})_{n \geq 0}$ be given by the Lie–Trotter splitting scheme (19) with time-step size $\tau \in (0, 1)$. Then for any nonnegative integer $n \geq 0$, one has $f_n^{\text{add}} \in L^2(\Omega, L_{x,v}^2)$ and

$$\mathbb{E}[\|f_n^{\text{add}}\|_{L_{x,v}^2}^2] = \|f_0\|_{L_{x,v}^2}^2 + t_n \sum_{k=1}^K \|\sigma_k\|_{L_{x,v}^2}^2, \quad (21)$$

where $t_n = n\tau$.

To obtain the trace formula (21), it suffices to prove that for any nonnegative integer $n \geq 0$ one has

$$\mathbb{E}[\|f_{n+1}^{\text{add}}\|_{L_{x,v}^2}^2] = \mathbb{E}[\|f_n^{\text{add}}\|_{L_{x,v}^2}^2] + \tau \sum_{k=1}^K \|\sigma_k\|_{L_{x,v}^2}^2.$$

Like for Proposition 1, two proofs are given below.

First proof of Proposition 2. Observe that for all $x \in \mathbb{T}^d$ and $v \in \mathbb{R}^d$ the random variable $f_n^{\text{add}}(x - \tau v - \tau^2 E(x), v - \tau E(x))$ and the Gaussian random variable $(\delta\beta_{n,k})_{1 \leq k \leq K}$ are independent. Using the expression (20), one obtains, for all $(x, v) \in \mathbb{T}^d \times \mathbb{R}^d$, the identity

$$\mathbb{E}[f_{n+1}^{\text{add}}(x, v)^2] = \mathbb{E}[f_n^{\text{add}}(x - tv - t^2 E(x), v - tE(x))^2] + \tau \sum_{k=1}^K \sigma_k(x, v)^2.$$

Note that applying twice a change of variables formulas one has

$$\begin{aligned} \iint \mathbb{E}[f_n^{\text{add}}(x - \tau v - \tau^2 E(x), v - \tau E(x))^2] dx dv &= \iint \mathbb{E}[f_n^{\text{add}}(x - \tau v, v)^2] dx dv \\ &= \iint \mathbb{E}[f_n^{\text{add}}(x, v)^2] dx dv. \end{aligned}$$

Integrating with respect to the variables x and v , one then obtains

$$\begin{aligned} \mathbb{E}[\|f_{n+1}^{\text{add}}\|_{L_{x,v}^2}^2] &= \iint \mathbb{E}[f_{n+1}^{\text{add}}(x, v)^2] dx dv \\ &= \iint \mathbb{E}[f_n^{\text{add}}(x, v)^2] dx dv + \tau \sum_{k=1}^K \iint \sigma_k(x, v)^2 dx dv \\ &= \mathbb{E}[\|f_n^{\text{add}}\|_{L_{x,v}^2}^2] + \tau \sum_{k=1}^K \|\sigma_k\|_{L_{x,v}^2}^2. \end{aligned}$$

□

Second proof of Proposition 2. Observe that the random mapping f_n^{add} and the Gaussian random variable $(\delta\beta_{n,k})_{1 \leq k \leq K}$ are independent. Using the isometry property for the linear operators $S^1(\tau), S^2(\tau) : L_{x,v}^2 \rightarrow L_{x,v}^2$, one then obtains

$$\begin{aligned} \mathbb{E}[\|f_{n+1}^{\text{add}}\|_{L_{x,v}^2}^2] &= \mathbb{E}[\|S^2(\tau)S^1(\tau)f_n^{\text{add}}\|_{L_{x,v}^2}^2] + \tau \sum_{k=1}^K \|\sigma_k\|_{L_{x,v}^2}^2 \\ &= \mathbb{E}[\|f_n^{\text{add}}\|_{L_{x,v}^2}^2] + \tau \sum_{k=1}^K \|\sigma_k\|_{L_{x,v}^2}^2. \end{aligned}$$

□

4.3. Numerical experiments. We begin the numerical experiments by illustrating the behavior of the linear Vlasov equation perturbed by additive noise (15), in dimension $d = 1$. Like for Figure 1 in the deterministic case (see Section 3), the initial value f_0 is given by (13) and the vector field E is given by (14).

The noise perturbation is given either by

$$\sigma_1(x, v) = \cos(v)\mathbf{1}_{|v| \leq 3}, \quad x \in \mathbb{T}, v \in \mathbb{R}, \quad (22)$$

or by

$$\sigma_1(x, v) = \sin(v) \mathbf{1}_{|v| \leq 3}, \quad x \in \mathbb{T}, v \in \mathbb{R}, \quad (23)$$

with $K = 1$ in both cases. In Figures 2 and 3 below, snapshots at the times $\{0, 0.5, 1, 1.5, 2, 2.5\}$ of the numerical solution computed using the Lie–Trotter splitting scheme 19 are displayed, with σ_1 given by (22) and (23) respectively. The discretization parameters are given by $\delta x = \frac{1}{500}$, $\delta v = \frac{4\pi}{500}$, and $\tau = 0.1$. One observes that the solutions behave differently from the deterministic case displayed in Figure 1. One also observes major differences between Figures 2 and 3 which are due to imposing a noise perturbation which is either symmetric or skew-symmetric with respect to the velocity variable v . Recall that the average value $\mathbb{E}[f^{\text{add}}(t, x, v)] = f^{\text{det}}(t, x, v)$ is solution of the deterministic PDE (6), which justifies the persistence of the deterministic behavior and of the influence of the initial condition in the snapshots.




FIGURE 2. Snapshots: approximation of the solution of the stochastic PDE with additive noise (15) with initial value f_0 given by (13), with σ_1 given by (22) at times $\{0, 0.5, 1, 1.5, 2, 2.5\}$, using the Lie–Trotter splitting scheme (19) with time-step size $\tau = 0.1$.

We continue these numerical experiments with Figure 4 in order to illustrate the trace formula (21) from Proposition 2 for the $L^2_{x,v}$ -norm of the Lie–Trotter splitting scheme (19). Let $d = 1$ and $T = 1$, the initial value f_0 is given by (13) and the vector field E is given by (14). For the noise perturbation, one has either $K = 1$ and

$$\sigma_1(x, v) = 0.5 \sin(v) \mathbf{1}_{|v| \leq 3}, \quad (24)$$

or $K = 2$ and

$$\sigma_1(x, v) = 0.5e^{-v^2/2} \cos(2\pi x), \quad \sigma_2(x, v) = 0.5e^{-v^2/2} \sin(2\pi x), \quad (25)$$

for all $x \in \mathbb{T}$ and $v \in \mathbb{R}$. The discretization parameters are given by $\delta x = \frac{1}{200}$, $\delta v = \frac{4\pi}{400}$, and $\tau = 0.1$. The expectation in the trace formula (21) is approximated




FIGURE 3. Snapshots: approximation of the solution of the stochastic PDE with additive noise (15) with initial value f_0 given by (13), with σ_1 given by (23) at times $\{0, 0.5, 1, 1.5, 2, 2.5\}$, using the Lie–Trotter splitting scheme (19) with time-step size $\tau = 0.1$.

by a standard Monte Carlo averaging procedure over 10^6 independent samples. The exact line in Figure 4 corresponds to the trace formula (18) from Proposition 1 satisfied by the exact solution. Even if Proposition 2 states that the Lie–Trotter splitting scheme (19) preserves the trace formula at all times $t_n = n\tau$, some error is visible. This may be due to the discretization procedure with respect to the spatial and velocity variables x and v , in particular since a truncation procedure for large v is applied. We have verified that increasing the Monte Carlo sample size does not seem to reduce the error visible in Figure 4.




FIGURE 4. Trace formula: illustration of Proposition 2 when applying the Lie–Trotter splitting scheme (19) to the SPDE with additive noise (15) with time-step size $\tau = 0.1$.

The final experiment in the additive noise case is devoted to investigate the mean-square order of convergence of the Lie–Trotter splitting scheme (19). In Figure 5, a loglog plot displays how the error

$$\sqrt{\sup_{x,v} \mathbb{E}[|f^{\text{add}}(x,v) - f^{\text{add,ref}}(x,v)|^2]}$$

converges to 0 when τ decreases, where x, v are grid points, the reference solution $f^{\text{add,ref}}$ is computed using the splitting scheme with time-step size $\tau_{\text{ref}} = 2^{-14}$. The time-step size τ takes values in $\{2^{-7}, \dots, 2^{-13}\}$, and the expectation is computed using a Monte Carlo averaging procedure over 500 independent samples. Note that the sample size used to illustrate the behaviour of the mean-square error is much smaller than the sample size used for the illustration of the trace formula in Figure 4 above. This is due to the fact that the variance also decreases when τ decreases. We have verified that increasing the Monte Carlo sample size does not significantly modify the behaviour of the mean-square error observed below. The discretization parameters are $\delta x = \frac{1}{100}$, $\delta v = \frac{4\pi}{200}$. The final time is given by $T = 0.5$, whereas f_0 and E are again given by (13) and (14) respectively. Like for Figure 4 above, the noise is given by (24) or (25). Based on these numerical experiments, we conjecture that the order of mean-square convergence of the scheme (19) is equal to 1. We leave the rigorous verification of this conjecture for future works.




FIGURE 5. Mean-square errors: Lie–Trotter scheme (19) applied to the SPDE with additive noise (15) driven by one-dimensional noise ($K = 1$, left) and two-dimensional noises ($K = 2$, right).

5. The stochastic linear Vlasov equation perturbed by multiplicative noise. In this section, we consider stochastic perturbations of the Vlasov equation (6) where the noise is multiplicative. In the analysis and applications of stochastic (partial) differential equations, it is well-known that several interpretations of multiplicative noise perturbations are possible. The Itô interpretation of the noise is considered in Section 5.1, then the Stratonovich interpretation is considered in Section 5.2. The objective of this section is to explain how to construct numerical schemes which are consistent with the two possible interpretations of the multiplicative noise, and to investigate which properties of the exact solution can be preserved at the discrete level.

In this section, the following condition is imposed: for all $1 \leq k \leq K$, the mapping $(x, v) \mapsto \sigma_k(x, v)$ is bounded. Some arguments below hold assuming in addition that there exists a real number σ such that the mapping $\sum_{k=1}^K \sigma_k^2$ is constant equal to σ^2 :

$$\sum_{k=1}^K \sigma_k(x, v)^2 =: \sigma^2, \quad \forall (x, v) \in \mathbb{T}^d \times \mathbb{R}^d. \quad (26)$$

For instance the condition (26) is satisfied if $d = 1$, $K = 2$, $\sigma_1(x, v) = \cos(2\pi x)$ and $\sigma_2(x, v) = \sin(2\pi x)$, with $\sigma = 1$. Note that this two-dimensional noise leads to a different dynamics for the SPDE than the case $K = 1$ and $\sigma_1(x, v) = \sigma = 1$.

5.1. Itô interpretation. We consider the following linear Vlasov equation driven by multiplicative noise interpreted in the Itô sense: for $t \geq 0$, $x \in \mathbb{T}^d$, $v \in \mathbb{R}^d$

$$\begin{cases} df^{\text{mI}}(t, x, v) + v \cdot \nabla_x f^{\text{mI}}(t, x, v) dt + E(x) \cdot \nabla_v f^{\text{mI}}(t, x, v) dt \\ = f^{\text{mI}}(t, x, v) dW(t, x, v), \\ f^{\text{mI}}(0, x, v) = f_0(x, v), \end{cases} \quad (27)$$

where the noise is defined by (1).

5.1.1. Analysis and properties of the problem. Like in the additive noise case (Section 4), let us consider several ways to define and deal with solutions of the SPDE (27).

On the one hand, using the group $(S(t))_{t \in \mathbb{R}}$ of linear operators given by (9) in Section 3, one can consider mild solutions: for all $t \geq 0$, one has

$$\begin{aligned} f^{\text{mI}}(t) &= S(t)f_0 + \int_0^t S(t-s)f^{\text{mI}}(s) dW(s) \\ &= S(t)f_0 + \sum_{k=1}^K \int_0^t S(t-s)(f^{\text{mI}}(s)\sigma_k) d\beta_k(s). \end{aligned} \quad (28)$$

On the other hand, using the expression (9) for the linear operator $S(t)$, one has for all $t \geq 0$, $x \in \mathbb{T}^d$ and $v \in \mathbb{R}^d$

$$f^{\text{mI}}(t, x, v) = f_0(\phi_t^{-1}(x, v)) + \sum_{k=1}^K \int_0^t f^{\text{mI}}(s, \phi_{t-s}^{-1}(x, v)) \sigma_k(\phi_{t-s}^{-1}(x, v)) d\beta_k(s). \quad (29)$$

Finally, the connection with the ordinary differential equation (7) can also be seen by applying the Itô–Wentzell formula (see Appendix A): if $t \mapsto (x_t, v_t) = \phi_t(x_0, v_0)$ is the solution of the ordinary differential equation (7) with arbitrary initial value $(x_0, v_0) \in \mathbb{T}^d \times \mathbb{R}^d$, and if the solution of (27) is sufficiently regular, then the stochastic process $t \geq 0 \mapsto f^{\text{mI}}(t, x_t, v_t)$ satisfies

$$df^{\text{mI}}(t, x_t, v_t) = \sum_{k=1}^K f^{\text{mI}}(t, x_t, v_t) \sigma_k(x_t, v_t) d\beta_k(t). \quad (30)$$

The formula (30) allows to retrieve the expression (29) of $f^{\text{mI}}(t, x, v)$ above by writing $(x_t, v_t) = \phi_t(x, v)$.

Let us now describe the properties of the solutions of the SPDE (27).

Proposition 3. *Let $(f^{\text{mI}}(t))_{t \geq 0}$ be the solution of the SPDE (27) with (non-random) initial value f_0 . One has the following properties.*

- Preservation of positivity. Assume that $f_0(x, v) \geq 0$ for all $(x, v) \in \mathbb{T}^d \times \mathbb{R}^d$. Then one has $f^{\text{mI}}(t, x, v) \geq 0$ almost surely for all $t \geq 0$ and $(x, v) \in \mathbb{T}^d \times \mathbb{R}^d$.
- Preservation of the expected mass. Assume that $f_0 \in L^1_{x,v}$. Then almost surely one has $f^{\text{mI}}(t) \in L^1_{x,v}$ for all $t \geq 0$, and

$$\iint \mathbb{E}[f^{\text{mI}}(t, x, v)] dx dv = \iint f_0(x, v) dx dv.$$

- Evolution law for the L^2 norm. Assume that $f_0 \in L^2_{x,v}$ and that the condition (26) is satisfied. Then one has $f^{\text{mI}}(t) \in L^2(\Omega, L^2_{x,v})$ for all $t \geq 0$, and

$$\mathbb{E}[\|f^{\text{mI}}(t)\|_{L^2_{x,v}}^2] = e^{\sigma^2 t} \|f_0\|_{L^2_{x,v}}^2.$$

Proof of Proposition 3.

- Owing to the expression (30) of the solution and applying a comparison principle for solutions of stochastic differential equations, one has $f^{\text{mI}}(t, x_t, v_t) \geq 0$ almost surely for all $t \geq 0$, and for any arbitrary initial value $(x_0, v_0) \in \mathbb{T}^d \times \mathbb{R}^d$. Choosing $(x_0, v_0) = \phi_t^{-1}(x, v)$ then yields $f^{\text{mI}}(t, x, v) \geq 0$ almost surely for all $(t, x, v) \in \mathbb{R}^+ \times \mathbb{T}^d \times \mathbb{R}^d$.
- Owing to the expression (29) of the solution using the flow $(\phi_t)_{t \in \mathbb{R}}$, one has for all $t \geq 0$

$$\begin{aligned} & \iint \mathbb{E}[f^{\text{mI}}(t, x, v)] dx dv \\ &= \iint f_0(\phi_t^{-1}(x, v)) dx dv \\ &+ \sum_{k=1}^K \iint \mathbb{E}\left[\int_0^t \sigma_k(\phi_{t-s}^{-1}(x, v)) f^{\text{mI}}(s, \phi_{t-s}^{-1}(x, v)) d\beta_k(s)\right] dx dv \\ &= \iint f_0(x, v) dx dv, \end{aligned}$$

since the expectation of the stochastic Itô integral vanishes and since ϕ_t preserves the volume in $\mathbb{T}^d \times \mathbb{R}^d$ for all $t \geq 0$.

- Owing to the expression (29) of the solution using the flow $(\phi_t)_{t \in \mathbb{R}}$, applying Itô's isometry formula yields

$$\begin{aligned} & \iint \mathbb{E}[f^{\text{mI}}(t, x, v)^2] dx dv = \iint f_0(\phi_t^{-1}(x, v))^2 dx dv \\ &+ \sum_{k=1}^K \int_0^t \iint \mathbb{E}[\sigma_k(\phi_{t-s}^{-1}(x, v))^2 f^{\text{mI}}(s, \phi_{t-s}^{-1}(x, v))^2] dx dv ds \\ &= \iint f_0(x, v)^2 dx dv + \sum_{k=1}^K \int_0^t \iint \mathbb{E}[\sigma_k(x, v)^2 f^{\text{mI}}(s, x, v)^2] dx dv ds, \end{aligned}$$

since the mapping ϕ_{t-s} preserves the volume in $\mathbb{T}^d \times \mathbb{R}^d$ for all $t \geq s \geq 0$. Using the condition (26) then yields the identity

$$\begin{aligned} \iint \mathbb{E}[f^{\text{mI}}(t, x, v)^2] dx dv &= \iint f_0(\phi_t^{-1}(x, v))^2 dx dv \\ &+ \sigma^2 \int_0^t \iint \mathbb{E}[f^{\text{mI}}(s, x, v)^2] dx dv ds. \end{aligned}$$

This implies that for all $t \geq 0$ one has

$$\mathbb{E}[\|f^{\text{mI}}(t)\|_{L^2_{x,v}}^2] = e^{\sigma^2 t} \|f_0\|_{L^2_{x,v}}^2.$$

□

Remark 4. The evolution law for the L^2 norm in Proposition 3 can be proved by an alternative approach: using the expression (28) of the mild solution, and using the Itô isometry formula in the Hilbert space $L^2_{x,v}$, one has

$$\begin{aligned} \mathbb{E}[\|f^{\text{mI}}(t)\|_{L^2_{x,v}}^2] &= \|S(t)f_0\|_{L^2_{x,v}}^2 + \sum_{k=1}^K \int_0^t \mathbb{E}[\|S(t-s)(f^{\text{mI}}(s)\sigma_k)\|_{L^2_{x,v}}^2] ds \\ &= \|f_0\|_{L^2_{x,v}}^2 + \sum_{k=1}^K \int_0^t \mathbb{E}[\|f^{\text{mI}}(s)\sigma_k\|_{L^2_{x,v}}^2] ds, \end{aligned}$$

since the linear operator $S(t-s) : L^2_{x,v} \rightarrow L^2_{x,v}$ is an isometry, for all $t \geq s \geq 0$. When the condition (26) is satisfied, one has

$$\sum_{k=1}^K \|f^{\text{mI}}(s)\sigma_k\|_{L^2_{x,v}}^2 = \sigma^2 \|f^{\text{mI}}(s)\|_{L^2_{x,v}}^2$$

and the conclusion is obtained as in the proof above.

5.1.2. Splitting scheme. Let us now describe the proposed numerical scheme for the temporal discretization of the SPDE (27) driven by multiplicative Itô noise. Like in the additive noise case presented in Section 4, a Lie–Trotter splitting strategy is applied. The treatment of the deterministic part is not modified. The auxiliary stochastic subsystem with multiplicative Itô noise

$$df(t, x, v) = f(t, x, v) dW(t, x, v), \quad (t, x, v) \in \mathbb{R}^+ \times \mathbb{T}^d \times \mathbb{R}^d$$

is solved exactly: for all $t \geq s \geq 0$, one has

$$f(t, x, v) = e^{\sum_{k=1}^K \sigma_k(\beta_k(t) - \beta_k(s)) - \frac{(t-s) \sum_{k=1}^K \sigma_k(x,v)^2}{2}} f(s, x, v), \quad \forall (x, v) \in \mathbb{T}^d \times \mathbb{R}^d.$$

Using the Lie–Trotter integrator (12) for the deterministic part and combining the discretizations of the deterministic and stochastic parts yields the following scheme: given the initial value f_0 and the time-step size $\tau \in (0, 1)$, set $f_0^{\text{mI}} = f_0$ and for any nonnegative integer $n \geq 0$ set

$$\begin{cases} \hat{f}_{n+1}^{\text{mI}} = S^2(\tau) S^1(\tau) f_n^{\text{mI}} \\ f_{n+1}^{\text{mI}}(x, v) = e^{\sum_{k=1}^K \sigma_k(x,v) \delta \beta_{n,k} - \frac{\tau \sum_{k=1}^K \sigma_k(x,v)^2}{2}} \hat{f}_{n+1}^{\text{mI}}(x, v), \quad x \in \mathbb{T}^d, v \in \mathbb{R}^d, \end{cases} \quad (31)$$

where the Wiener increments $\delta \beta_{n,k}$ are given by (3) in Section 2.

The Lie–Trotter splitting scheme (31) satisfies the same properties as the exact solution stated in Proposition 3.

Proposition 5. Let $(f_n^{\text{mI}})_{n \geq 0}$ be the solution of the Lie–Trotter splitting scheme (31) with initial value f_0 . One then has the following properties.

- Preservation of positivity. Assume that $f_0(x, v) \geq 0$ for all $(x, v) \in \mathbb{T}^d \times \mathbb{R}^d$. Then for any time-step size $\tau \in (0, 1)$, one has $f_n^{\text{mI}}(x, v) \geq 0$ almost surely for any nonnegative integer $n \geq 0$ and all $(x, v) \in \mathbb{T}^d \times \mathbb{R}^d$.

- Preservation of the expected mass. Assume that $f_0 \in L^1_{x,v}$. Then almost surely one has $f_n^{\text{mI}} \in L^1_{x,v}$ for all $n \geq 0$, and

$$\iint \mathbb{E}[f_n^{\text{mI}}(x, v)] dx dv = \iint f_0(x, v) dx dv.$$

- Evolution law for the L^2 norm. Assume that $f_0 \in L^2_{x,v}$ and that the condition (26) is satisfied. Then one has $f_n^{\text{mI}} \in L^2(\Omega, L^2_{x,v})$ for all $n \geq 0$, and

$$\mathbb{E}[\|f_n^{\text{mI}}\|_{L^2_{x,v}}^2] = e^{\sigma^2 t_n} \|f_0\|_{L^2_{x,v}}^2,$$

where $t_n = n\tau$.

Concerning the positivity-preserving property, a similar splitting scheme has been proposed in the recent work [15] for another class of SPDEs.

Proof of Proposition 5.

- The proof of the positivity preserving property is performed by recursion. It is satisfied if $n = 0$ since $f_0^{\text{mI}} = f_0$. Assume that $f_n^{\text{mI}}(x, v) \geq 0$ almost surely for all $(x, v) \in \mathbb{T}^d \times \mathbb{R}^d$. Owing to the positivity preserving property for the deterministic problem (see Section 3), one has for all $(x, v) \in \mathbb{T}^d \times \mathbb{R}^d$

$$\hat{f}_{n+1}^{\text{mI}}(x, v) = (S^2(\tau)S^1(\tau)f_n^{\text{mI}})(x, v) = f_n^{\text{mI}}(x - tv - t^2E(x), v - tE(x)) \geq 0$$

$$f_{n+1}^{\text{mI}}(x, v) = e^{\sum_{k=1}^K \sigma_k \delta \beta_{n,k} - \frac{(t-s) \sum_{k=1}^K \sigma_k (x,v)^2}{2}} \hat{f}_{n+1}^{\text{mI}}(x, v) \geq 0.$$

Therefore one has $f_{n+1}^{\text{mI}}(x, v) \geq 0$ almost surely for all $(x, v) \in \mathbb{T}^d \times \mathbb{R}^d$.

- Observe that the random variables $\hat{f}_{n+1}^{\text{mI}}(x, v)$ and $(\delta \beta_{n,k})_{1 \leq k \leq K}$ are independent. As a result, using the well-known expression for the exponential moments of Gaussian random variables, one has

$$\begin{aligned} \iint \mathbb{E}[f_{n+1}^{\text{mI}}(x, v)] dx dv &= \iint \mathbb{E}[e^{\sum_{k=1}^K \delta \beta_{n,k} \sigma_k(x,v) - \frac{\tau}{2} \sum_{k=1}^K \sigma_k(x,v)^2}] \mathbb{E}[\hat{f}_{n+1}^{\text{mI}}(x, v)] dx dv \\ &= \iint \mathbb{E}[\hat{f}_{n+1}^{\text{mI}}(x, v)] dx dv. \end{aligned}$$

Finally, one has $\hat{f}_{n+1}^{\text{mI}} = S^2(\tau)S^1(\tau)f_n^{\text{mI}}$ owing to (31), where the linear operators $S^1(\tau)$ and $S^2(\tau)$ preserve the integral, thus one obtains the identity

$$\iint \mathbb{E}[f_{n+1}^{\text{mI}}(x, v)] dx dv = \iint \mathbb{E}[\hat{f}_{n+1}^{\text{mI}}(x, v)] dx dv = \iint \mathbb{E}[f_n^{\text{mI}}(x, v)] dx dv.$$

- Applying the same arguments as above, one obtains

$$\begin{aligned} \mathbb{E}[\|f_{n+1}^{\text{mI}}\|_{L^2_{x,v}}^2] &= \iint \mathbb{E}[e^{2 \sum_{k=1}^K \sigma_k(x,v) \delta \beta_{n,k} - \tau \sum_{k=1}^K \sigma_k(x,v)^2}] \mathbb{E}[\hat{f}_{n+1}^{\text{mI}}(x, v)^2] dx dv \\ &= \iint e^{\tau \sum_{k=1}^K \sigma_k(x,v)^2} \mathbb{E}[\hat{f}_{n+1}^{\text{mI}}(x, v)^2] dx dv. \end{aligned}$$

Then using the condition (26) and the expression $\hat{f}_{n+1}^{\text{mI}} = S^2(\tau)S^1(\tau)f_n^{\text{mI}}$ gives

$$\mathbb{E}[\|f_{n+1}^{\text{mI}}\|_{L^2_{x,v}}^2] = e^{\sigma^2 \tau} \mathbb{E}[\|(S^2(\tau)S^1(\tau))f_n^{\text{mI}}\|_{L^2_{x,v}}^2] = e^{\sigma^2 \tau} \mathbb{E}[\|f_n^{\text{mI}}\|_{L^2_{x,v}}^2],$$

using the fact that $S^1(\tau)$ and $S^2(\tau)$ are isometries from $L^2_{x,v}$ to $L^2_{x,v}$. The evolution law then follows from a straightforward recursion argument. \square

Remark 6. If the noise in the SPDE (27) is a purely temporal Wiener process, i.e. $W(t, x, v) = \beta(t)$ for all $(t, x, v) \in \mathbb{R}^+ \times \mathbb{T}^d \times \mathbb{R}^d$ where $(\beta(t))_{t \geq 0}$ is a standard real-valued Wiener process, then the exact solution of (27) and the numerical solution given by (31) can be written

$$f^{\text{mI}}(t) = e^{\beta(t) - \frac{t}{2}} f^{\text{det}}(t), \quad t \geq 0; \quad f_n^{\text{mI}} = e^{\beta(t_n) - \frac{t_n}{2}} f_n^{\text{det}}, \quad n \geq 0,$$

where $(f^{\text{det}}(t))_{t \geq 0}$ is the exact solution of the deterministic equation (6), given by (8), and $(f_n^{\text{det}})_{n \geq 0}$ is given by the deterministic Lie–Trotter splitting scheme (12).

In that situation, Propositions 3 and 5 are straightforward consequences of the results described for the deterministic problem in Section 3.

Remark 7. Applying the standard Euler–Maruyama scheme to treat the stochastic part of (27) provides the scheme

$$f_{n+1}^{\text{mIEM}} = S^2(\tau) S^1(\tau) f_n^{\text{mIEM}} + \sum_{k=1}^K \delta \beta_{n,k} \sigma_k f_n^{\text{mIEM}}.$$

That scheme does not satisfy the positivity preserving property and the evolution law for the L^2 norm stated in Proposition 5 for the proposed Lie–Trotter scheme (31).

5.1.3. Numerical experiments. We begin the numerical experiments by illustrating the behavior of the linear Vlasov equation perturbed by multiplicative Itô noise (27), in dimension $d = 1$. In all the experiments below, the initial value f_0 is given by (13) and the vector field E is given by (14). The discretization parameters are given by $\delta x = \frac{1}{400}$, $\delta v = \frac{4\pi}{800}$, and $\tau = 0.1$. The snapshots of the numerical solution at times $\{0, 0.5, 1, 1.5, 2, 2.5\}$ computed using the splitting scheme (31) are provided in Figures 6 and 7, with $K = 1$ and diffusion coefficient σ_1 given by (22) and (23) respectively. In both experiments, one observes that the solution remains nonnegative, which illustrates the positivity preserving property stated in Proposition 5 on the considered realization.

Let us now check the almost sure positivity preserving property for the Lie–Trotter splitting scheme (31) in a more rigorous way: we have run 2.10^4 independent samples on the time interval $[0, 1]$, with initial value (13), with the same discretization parameters as above, and with different choices of the diffusion coefficients: either $K = 1$ and σ_1 given by (23), or $K = 2$ and σ_1 and σ_2 give by

$$\sigma_1(x, v) = \cos(2\pi x), \quad \sigma_2(x, v) = \sin(2\pi x). \quad (32)$$

All the samples only take nonnegative values, which confirms the positivity preserving property stated in Proposition 5.

Next, we illustrate the preservation of the expected mass and the evolution law of the L^2 norm stated in Proposition 5. In these experiments, one has $d = 1$, $T = 1$, $\delta x = \frac{1}{200}$, $\delta v = \frac{4\pi}{400}$ and $\tau = 0.1$. The expectations are computed using an averaging procedure over 5.10^5 samples. Since the solution is nonnegative, the mass is in fact equal to the L^1 norm of the solution. The results are presented in Figure 8, with different choices of the diffusion coefficients: $K = 1$ with $\sigma_1(x, v) = 1$, $K = 1$ with σ_1 given by (23), and $K = 2$ with σ_1 and σ_2 given by (32), respectively. Note that the condition (26) is satisfied in the first and in the third case. We observe a good agreement with the theoretical results given in Proposition 5.

We conclude these numerical experiments in the multiplicative Itô noise case by investigating the mean-square order of convergence of the Lie–Trotter splitting




FIGURE 6. Snapshots: approximation of the solution of the stochastic PDE with multiplicative Itô noise (27) with initial value f_0 given by (13), with σ_1 given by (22) at times $\{0, 0.5, 1, 1.5, 2, 2.5\}$, using the Lie–Trotter splitting scheme (31) with time-step size $\tau = 0.1$.

scheme (31). The same procedure as in the additive noise case (Section 4) is applied. A reference solution is computed using the splitting scheme with time-step size $\tau_{\text{ref}} = 2^{-14}$, and the errors are computed when the time-step size τ takes values in $\{2^{-7}, \dots, 2^{-13}\}$. The expectation is computed using a Monte Carlo averaging procedure over 500 independent samples. The discretization parameters are $\delta x = \frac{1}{100}$, $\delta v = \frac{4\pi}{200}$. The final time is $T = 0.5$. The noise is given by (23) or by (32). The results are presented in a loglog plot in Figure 9. We observe a mean-square convergence order equal to 1.

5.2. Stratonovich interpretation. Let us now consider the linear Vlasov equation driven by a multiplicative noise interpreted in the Stratonovich sense, for $t \geq 0$, $x \in \mathbb{T}^d$ and $v \in \mathbb{R}^d$:

$$\begin{cases} df^{\text{mS}}(t, x, v) + v \cdot \nabla_x f^{\text{mS}}(t, x, v) dt + E(x) \cdot \nabla_v f^{\text{mS}}(t, x, v) dt \\ = f^{\text{mS}}(t, x, v) \circ dW(t, x, v), \\ f^{\text{mS}}(0, x, v) = f_0(x, v), \end{cases} \quad (33)$$

where we recall that the noise is defined by (1) (Section 2) and that the symbol \circ denotes the Stratonovich product. The stochastic partial differential equation (33) has the equivalent Itô formulation, for $t \geq 0$, $x \in \mathbb{T}^d$, $v \in \mathbb{R}^d$:

$$\begin{cases} df^{\text{mS}}(t, x, v) + v \cdot \nabla_x f^{\text{mS}}(t, x, v) dt + E(x) \cdot \nabla_v f^{\text{mS}}(t, x, v) dt \\ = f^{\text{mS}}(t, x, v) dW(t, x, v) + \frac{1}{2} \sum_{k=1}^K \sigma_k(x, v)^2 f^{\text{mS}}(t, x, v) dt, \\ f^{\text{mS}}(0, x, v) = f_0(x, v). \end{cases} \quad (34)$$




FIGURE 7. Snapshots: approximation of the solution of the stochastic PDE with multiplicative Itô noise (27) with initial value f_0 given by (13), with σ_1 given by (23) at times $\{0, 0.5, 1, 1.5, 2, 2.5\}$, using the Lie–Trotter splitting scheme (31) with time-step size $\tau = 0.1$.

5.2.1. *Analysis and properties of the problem.* Using the tools described in Section 3 and like in the multiplicative Itô noise case studied in Section 5.1, solutions of (33) can be written in different ways.


On the one hand, using the group $(S(t))_{t \in \mathbb{R}}$ of linear operators given by (9) in Section 3, one can consider mild solutions of (34): for all $t \geq 0$, one has

$$\begin{aligned} f^{\text{mS}}(t) = & S(t)f_0 + \sum_{k=1}^K \int_0^t S(t-s)(f^{\text{mS}}(s)\sigma_k) d\beta_k(s) \\ & + \frac{1}{2} \sum_{k=1}^K \int_0^t S(t-s)(f^{\text{mS}}(s)\sigma_k^2) ds. \end{aligned} \quad (35)$$


On the other hand, using the expression (9) for the linear operator $S(t)$, one has for all $t \geq 0$, $x \in \mathbb{T}^d$ and $v \in \mathbb{R}^d$

$$\begin{aligned} f^{\text{mS}}(t, x, v) = & f_0(\phi_t^{-1}(x, v)) + \sum_{k=1}^K \int_0^t f^{\text{mS}}(s, \phi_{t-s}^{-1}(x, v))\sigma_k(\phi_{t-s}^{-1}(x, v)) d\beta_k(s) \\ & + \frac{1}{2} \sum_{k=1}^K \int_0^t f^{\text{mS}}(s, \phi_{t-s}^{-1}(x, v))\sigma_k^2(\phi_{t-s}^{-1}(x, v)) ds. \end{aligned} \quad (36)$$

Finally, the connection with the ordinary differential equation (7) can also be seen by applying the Itô–Wentzell formula (see Appendix A): if $t \mapsto (x_t, v_t) = \phi_t(x_0, v_0)$ is the solution of the ordinary differential equation (7) with arbitrary initial value $(x_0, v_0) \in \mathbb{T}^d \times \mathbb{R}^d$, and if the solution of (33) is sufficiently regular, then the




(A) Noise given by $\sigma_1(x, v) = 1$, $K = 1$. (B) Noise given by (23), $K = 1$.



(C) Noise given by (32), $K = 2$.

FIGURE 8. Preservation of the expected mass and evolution law for the $L^2_{x,v}$ norm: illustration of Proposition 5 when applying the Lie–Trotter scheme (31) with $\tau = 0.1$ to the SPDE with multiplicative Itô noise (27) with time-step size $\tau = 0.1$.



(A) Noise given by (23).

(B) Noise given by (32).

FIGURE 9. Mean-square errors: Mean-square errors: Lie–Trotter scheme (31) applied to the SPDE with multiplicative Itô noise (27) driven by one-dimensional noise ($K = 1$, left) and by two-dimensional noise ($K = 2$, right).

stochastic process $t \geq 0 \mapsto f^{\text{mS}}(t, x_t, v_t)$ satisfies

$$\begin{aligned} df^{\text{mS}}(t, x_t, v_t) &= \sum_{k=1}^K f^{\text{mS}}(t, x_t, v_t) \sigma_k(x_t, v_t) d\beta_k(t) + \frac{1}{2} \sum_{k=1}^K f^{\text{mS}}(t, x_t, v_t) \sigma_k(x_t, v_t)^2 dt \\ &= \sum_{k=1}^K f^{\text{mS}}(t, x_t, v_t) \sigma_k(x_t, v_t) \circ d\beta_k(t). \end{aligned} \tag{37}$$

The formula (37) allows to retrieve the expression (36) of $f^{\text{mI}}(t, x, v)$ above by writing $(x_t, v_t) = \phi_t(x, v)$.

Let us now describe the properties of the solutions of the SPDE (33).

Proposition 8. *Let $(f^{\text{mS}}(t))_{t \geq 0}$ be the solution of the SPDE (33) with initial value f_0 . One has the following properties.*

- Preservation of positivity. Assume that $f_0(x, v) \geq 0$ for all $(x, v) \in \mathbb{T}^d \times \mathbb{R}^d$. Then, one has $f^{\text{mS}}(t, x, v) \geq 0$ almost surely for all $t \geq 0$ and $(x, v) \in \mathbb{T}^d \times \mathbb{R}^d$.
- Evolution law for the L^2 norm. Assume that $f_0 \in L^2_{x,v}$ and that the condition (26) is satisfied. Then one has $f(t) \in L^2(\Omega, L^2_{x,v})$ for all $t \geq 0$, and

$$\mathbb{E}[\|f^{\text{mS}}(t)\|_{L^2_{x,v}}^2] = e^{2\sigma^2 t} \|f_0\|_{L^2_{x,v}}^2.$$

Proof of Proposition 8.

- Owing to the expression (37) of the solution and applying a comparison principle for solutions of stochastic differential equations, one has $f^{\text{mS}}(t, x_t, v_t) \geq 0$ almost surely for all $t \geq 0$, and for any arbitrary initial value $(x_0, v_0) \in \mathbb{T}^d \times \mathbb{R}^d$. Choosing $(x_0, v_0) = \phi_t^{-1}(x, v)$ then yields $f^{\text{mS}}(t, x, v) \geq 0$ almost surely for all $(t, x, v) \in \mathbb{R}^+ \times \mathbb{T}^d \times \mathbb{R}^d$.
- Owing to the expression (35) of the mild solution, using Itô's formula in the Hilbert space $L^2_{x,v}$ and the isometry property of the group $(S(t))_{t \geq 0}$ (see Section 3), one obtains for all $t \geq 0$

$$\frac{1}{2} \frac{d\mathbb{E}[\|f^{\text{mS}}(t)\|_{L^2_{x,v}}^2]}{dt} = \sum_{k=1}^K \mathbb{E}[\|\sigma_k f^{\text{mS}}(t)\|_{L^2_{x,v}}^2].$$

Using the condition (26) and integrating then yields the identity

$$\mathbb{E}[\|f^{\text{mS}}(t)\|_{L^2_{x,v}}^2] = e^{2\sigma^2 t} \|f_0\|_{L^2_{x,v}}^2.$$

□

5.2.2. Splitting scheme. Let us now describe the proposed numerical scheme for the temporal discretization of the SPDE (33) driven by multiplicative Itô noise. Like in the multiplicative Itô noise case presented in Section 5.1, a Lie–Trotter splitting strategy is applied. The treatment of the deterministic part is not modified. Compared with Section 5.1, the auxiliary stochastic subsystem needs to be considered with Stratonovich interpretation of the noise:

$$df(t, x, v) = f(t, x, v) \circ dW(t, x, v), \quad (t, x, v) \in \mathbb{R}^+ \times \mathbb{T}^d \times \mathbb{R}^d$$

The auxiliary stochastic subsystem above is solved exactly: for all $t \geq s \geq 0$ one has

$$f(t, x, v) = e^{\sum_{k=1}^K \sigma_k(\beta_k(t) - \beta_k(s))} f(s, x, v), \quad \forall (x, v) \in \mathbb{T}^d \times \mathbb{R}^d.$$

Using the Lie–Trotter integrator (12) for the deterministic part and combining the discretizations of the deterministic and stochastic parts yields the following scheme: given the initial value f_0 and the time-step size $\tau \in (0, 1)$, set $f_0^{\text{mS}} = f_0$ and for any nonnegative integer $n \geq 0$ set

$$\begin{cases} \hat{f}_{n+1}^{\text{mS}}(x, v) = S^2(\tau)S^1(\tau)f_n^{\text{mS}}(x, v) \\ f_{n+1}^{\text{mS}}(x, v) = e^{\sum_{k=1}^K \sigma_k(x, v)\delta\beta_{n,k}} \hat{f}_{n+1}^{\text{mS}}(x, v), \end{cases} \quad x \in \mathbb{T}^d, v \in \mathbb{R}^d, \quad (38)$$

where we recall that the Wiener increments $\delta\beta_{n,k}$ are given by (3), see Section 2.

The Lie–Trotter splitting scheme (38) satisfies the same properties as the exact solution stated in Proposition 8.

Proposition 9. *Let $(f_n^{\text{mS}})_{n \geq 0}$ be the solution of the Lie–Trotter splitting scheme (38) with initial value f_0 . One then has the following properties.*

- Preservation of positivity. Assume that $f_0(x, v) \geq 0$ for all $(x, v) \in \mathbb{T}^d \times \mathbb{R}^d$. Then, for any time-step size $\tau \in (0, 1)$, one has $f_n^{\text{mS}}(x, v) \geq 0$ almost surely for any nonnegative integer $n \in \mathbb{N}$ and all $(x, v) \in \mathbb{T}^d \times \mathbb{R}^d$.
- Evolution law for the L^2 norm. Assume that $f_0 \in L^2_{x,v}$ and that the condition (26) is satisfied. Then, one has $f_n^{\text{mS}} \in L^2(\Omega, L^2_{x,v})$, for all $n \geq 0$, and

$$\mathbb{E}[\|f_n^{\text{mS}}\|_{L^2_{x,v}}^2] = e^{2\sigma^2 t_n} \|f_0\|_{L^2_{x,v}}^2,$$

where $t_n = n\tau$.

The proof is similar to the proof of Proposition 3 from Section 5.1.

Proof of Proposition 9.

- The proof of the positivity preserving property is performed by recursion. It is satisfied if $n = 0$ since $f_0^{\text{mS}} = f_0$. Assume that $f_n^{\text{mS}}(x, v) \geq 0$ almost surely for all $(x, v) \in \mathbb{T}^d \times \mathbb{R}^d$. Owing to the positivity preserving property for the deterministic problem (see Section 3), one has for all $(x, v) \in \mathbb{T}^d \times \mathbb{R}^d$

$$\begin{aligned} \hat{f}_{n+1}^{\text{mS}}(x, v) &= (S^2(\tau)S^1(\tau)f_n^{\text{mS}})(x, v) = f_n^{\text{mS}}(x - tv - t^2E(x), v - tE(x)) \geq 0 \\ f_{n+1}^{\text{mS}}(x, v) &= e^{\sum_{k=1}^K \sigma_k(x, v)\delta\beta_{n,k}} \hat{f}_{n+1}^{\text{mS}}(x, v) \geq 0. \end{aligned}$$

Therefore one has $f_{n+1}^{\text{mS}}(x, v) \geq 0$ almost surely for all $(x, v) \in \mathbb{T}^d \times \mathbb{R}^d$.

- Observe that the random variables $\hat{f}_{n+1}^{\text{mS}}(x, v)$ and $(\delta\beta_{n,k})_{1 \leq k \leq K}$ are independent. As a result, using the well-known expression for the exponential moments of Gaussian random variables, one has

$$\begin{aligned} \mathbb{E}[\|f_{n+1}^{\text{mS}}\|_{L^2_{x,v}}^2] &= \iint \mathbb{E}[e^{2\sum_{k=1}^K \sigma_k(x, v)\delta\beta_{n,k}}] \mathbb{E}[\hat{f}_{n+1}^{\text{mS}}(x, v)^2] dx dv \\ &= \iint e^{2\tau \sum_{k=1}^K \sigma_k(x, v)^2} \mathbb{E}[\hat{f}_{n+1}^{\text{mS}}(x, v)^2] dx dv. \end{aligned}$$

Then using the condition (26) and the expression $\hat{f}_{n+1}^{\text{mS}} = S^2(\tau)S^1(\tau)f_n^{\text{mS}}$ gives

$$\mathbb{E}[\|f_{n+1}^{\text{mS}}\|_{L^2_{x,v}}^2] = e^{2\sigma^2 \tau} \mathbb{E}[\|(S^2(\tau)S^1(\tau))f_n^{\text{mS}}\|_{L^2_{x,v}}^2] = e^{\sigma^2 \tau} \mathbb{E}[\|f_n^{\text{mS}}\|_{L^2_{x,v}}^2],$$

using the fact that $S^1(\tau)$ and $S^2(\tau)$ are isometries from $L^2_{x,v}$ to $L^2_{x,v}$. The evolution law then follows by a straightforward recursion argument. \square

Remark 10 below is a discussion in the Stratonovich noise case of the situation described in Remark 6 in the Itô noise case above.

Remark 10. If the noise in the SPDE (33) is a purely temporal Wiener process, i.e. $W(t, x, v) = \beta(t)$ for all $(t, x, v) \in \mathbb{R}^+ \times \mathbb{T}^d \times \mathbb{R}^d$ where $(\beta(t))_{t \geq 0}$ is a standard real-valued Wiener process, then the exact solution of (33) and the numerical solution given by (38) can be written

$$f^{\text{mS}}(t) = e^{\beta(t)} f^{\text{det}}(t), \quad t \geq 0; \quad f_n^{\text{mS}} = e^{\beta(t_n)} f_n^{\text{det}}, \quad n \geq 0,$$

where $(f^{\text{det}}(t))_{t \geq 0}$ is the exact solution of the deterministic equation (6), given by (8), and $(f_n^{\text{det}})_{n \geq 0}$ is given by the deterministic Lie–Trotter splitting scheme (12).

In that situation, Propositions 8 and 9 are straightforward consequences of the results described for the deterministic problem in Section 3.

5.2.3. Numerical experiments. We begin the numerical experiments by illustrating the behavior of the linear Vlasov equation perturbed by multiplicative Stratonovich noise (33), in dimension $d = 1$. In all the experiments below, the initial value f_0 is given by (13) and the vector field E is given by (14). The discretization parameters are given by $\delta x = \frac{1}{400}$, $\delta v = \frac{4\pi}{800}$, and $\tau = 0.1$. The snapshots of the numerical solution at times $\{0, 0.5, 1, 1.5, 2, 2.5\}$ computed using the splitting scheme (38) are provided in Figures 10 and 11, with $K = 1$ and diffusion coefficient σ_1 given by (22) and (23) respectively. In both experiments, one observes that the solution remains nonnegative, which illustrates the positivity preserving property stated in Proposition 9 on the considered realization.




FIGURE 10. Snapshots: approximation of the solution of the stochastic PDE with multiplicative Stratonovich noise (33) with initial value f_0 given by (13), with σ_1 given by (22) at times $\{0, 0.5, 1, 1.5, 2, 2.5\}$, using the Lie–Trotter splitting scheme (38) with time-step size $\tau = 0.1$.




FIGURE 11. Snapshots: approximation of the solution of the stochastic PDE with multiplicative Stratonovich noise (33) with initial value f_0 given by (13), with σ_1 given by (23) at times $\{0, 0.5, 1, 1.5, 2, 2.5\}$, using the Lie–Trotter splitting scheme (38) with time-step size $\tau = 0.1$.

Next, we illustrate the evolution law of the L^2 norm stated in Proposition 9. In these experiments, one has $d = 1$, $T = 1$, $\delta x = \frac{1}{200}$, $\delta v = \frac{4\pi}{400}$ and $\tau = 0.1$. The expectations are computed using an averaging procedure over $5 \cdot 10^5$ samples. The results are presented in Figure 12, with different choices of the diffusion coefficients: $K = 1$ with $\sigma_1(x, v) = 1$, $K = 1$ with σ_1 given by (23), and $K = 2$ with σ_1 and σ_2 given by (32), respectively. Note that the condition (26) is satisfied in the first and in the third case. We observe a good agreement with the theoretical results given in Proposition 9.

We conclude these numerical experiments in the multiplicative Stratonovich noise case by investigating the mean-square order of convergence of the Lie–Trotter splitting scheme (38). The same procedure as in the multiplicative Itô noise case (Section 5.1) is applied. A reference solution is computed using the splitting scheme with time-step size $\tau_{\text{ref}} = 2^{-14}$, and the errors are computed when the time-step size τ takes values in $\{2^{-7}, \dots, 2^{-13}\}$. The expectation is computed using a Monte Carlo averaging procedure over 500 independent samples. The discretization parameters are $\delta x = \frac{1}{100}$, $\delta v = \frac{4\pi}{200}$. The final time is $T = 0.5$. The noise is given by (23) or by (32). The results are presented in a loglog plot in Figure 13. We observe a mean-square convergence order equal to 1.

6. The stochastic linear Vlasov equation perturbed by transport noise. In

this section, we consider a stochastic perturbation of the Vlasov partial differential equation (6) where the noise is of transport type: for $t \geq 0, x \in \mathbb{T}^d, v \in \mathbb{R}^d$

$$\begin{cases} df^{\text{tr}}(t, x, v) + v \cdot \nabla_x f^{\text{tr}}(t, x, v) dt + E(x) \cdot \nabla_v f^{\text{tr}}(t, x, v) dt \\ \quad + \nabla_v f^{\text{tr}}(t, x, v) \odot dW(t, x) = 0, \\ f^{\text{tr}}(0, x, v) = f_0(x, v), \end{cases} \quad (39)$$





FIGURE 12. Evolution law for the $L^2_{x,v}$ norm: illustration of Proposition 9 when applying the Lie-Trotter scheme (38) with $\tau = 0.1$ to the SPDE with multiplicative Stratonovich noise (33) with time-step size $\tau = 0.1$.

where the noise $\mathbf{W}(t, x)$ is given by (2) and the notation

$$\begin{aligned} \nabla_v f^{\text{tr}}(t, x, v) \odot d\mathbf{W}(t, x) &= \sum_{k=1}^K \nabla_v f^{\text{tr}}(t, x, v) \cdot \sigma_k(x) \odot d\beta_k(t) \\ &= \sum_{j=1}^d \sum_{k=1}^K \partial_{v_j} f^{\text{tr}}(t, x, v) \sigma_{j,k}(x) \odot d\beta_k(t), \end{aligned}$$

is used, with \odot denoting the Stratonovich product and \odot denoting the combination of the Stratonovich product \odot and of the inner product \cdot in the Euclidean space \mathbb{R}^d .

To identify the equivalent Itô formulation of the stochastic partial differential equation (39), it is convenient to set for all $x \in \mathbb{T}^d$ and $1 \leq i, j \leq d$

$$a_{i,j}(x) = \sum_{k=1}^K \sigma_{i,k}(x) \sigma_{j,k}(x).$$




FIGURE 13. Mean-square errors: Lie–Trotter scheme (38) applied to the SPDE with multiplicative Stratonovich noise (33) driven by one-dimensional noise ($K = 1$, left) and by two-dimensional noise ($K = 2$, right).

The Itô formulation of (39) is then the following: for $t \geq 0, x \in \mathbb{T}^d, v \in \mathbb{R}^d$

$$\begin{cases} df^{\text{tr}}(t, x, v) + v \cdot \nabla_x f^{\text{tr}}(t, x, v) dt \\ \quad + E(x) \cdot \nabla_v f^{\text{tr}}(t, x, v) dt + \nabla_v f^{\text{tr}}(t, x, v) \cdot d\mathbf{W}(t, x) \\ \quad = \frac{1}{2} \sum_{k=1}^K (\sigma_k(x) \cdot \nabla_v)^2 f^{\text{tr}}(t, x, v) dt, \\ f^{\text{tr}}(0, x, v) = f_0(x, v), \end{cases} \quad (40)$$

with the notation

$$\begin{aligned} \nabla_v f^{\text{tr}}(t, x, v) \cdot d\mathbf{W}(t, x) &= \sum_{j=1}^d \sum_{k=1}^K \partial_{v_j} f^{\text{tr}}(t, x, v) \sigma_{j,k}(x) d\beta_k(t) \\ \sum_{k=1}^K (\sigma_k(x, v) \cdot \nabla_v)^2 f^{\text{tr}}(t, x, v) &= \sum_{i,j=1}^d \sum_{k=1}^K \sigma_{i,k}(x) \sigma_{j,k}(x) \partial_{v_i} \partial_{v_j} f^{\text{tr}}(t, x, v) \\ &= \sum_{i,j=1}^d a_{i,j}(x) \partial_{v_i} \partial_{v_j} f^{\text{tr}}(t, x, v). \end{aligned}$$

6.1. Analysis and properties of the problem. Recall that the deterministic linear Vlasov equation (6) is connected with the ordinary differential equation (7), see Section 3. This connection is explored in Sections 4 and 5 for some stochastic perturbations of (6). Contrary to those situations, for the stochastic linear Vlasov equation driven by transport noise (39), the connection requires to introduce the stochastic differential equation

$$\begin{cases} dX_t = V_t \\ dV_t = E(X_t) dt + \sum_{k=1}^K \sigma_k(X_t) d\beta_k(t), \end{cases} \quad (41)$$

for all $t \geq 0$, instead of the ordinary differential equation (7). In the second line of (41), one may use the notation $d\mathbf{W}(t, X_t)$ to refer to $\sum_{k=1}^K \sigma_k(X_t) d\beta_k(t)$.

Assume that $(f^{\text{tr}}(t))_{t \geq 0}$ is a sufficiently regular solution of the SPDE (39) (or of its equivalent formulation (40)), then for any solution $(X_t, V_t)_{t \geq 0}$ of the SDE system (41), applying the Itô–Wentzell formula (see Appendix A) one obtains the identity

$$df^{\text{tr}}(t, X_t, V_t) = 0.$$

This means that the SDE (41) provides characteristic curves for the stochastic Vlasov equation (39) driven by transport noise. This gives a strategy to solve the SPDE (39) by the method of lines. Instead of using the flow $(\phi_t)_{t \in \mathbb{R}}$ associated with the ODE (7), in the present case this strategy is based on the notion of stochastic flow of diffeomorphisms, see for instance the monograph [48]. We use the notation $(\psi_t)_{t \geq 0}$ to denote the stochastic flow of diffeomorphisms associated with the SDE (41). In particular, for any (deterministic) initial values $(X_0, V_0) \in \mathbb{T}^d \times \mathbb{R}^d$, the unique solution of the SDE (41) at any time $t \geq 0$ is given by $(X_t, V_t) = \psi_t(X_0, V_0)$. In addition, almost surely the mapping ψ_t preserves the volume in $\mathbb{T}^d \times \mathbb{R}^d$ for all $t \geq 0$.

It results from the identity above that, for all $t \geq 0$ and all (deterministic) $X_0 \in \mathbb{T}^d$ and $V_0 \in \mathbb{R}^d$, one has

$$f^{\text{tr}}(t, \psi_t(X_0, V_0)) = f^{\text{tr}}(t, X_t, V_t) = f^{\text{tr}}(0, X_0, V_0) = f_0(X_0, V_0).$$

Finally one obtains the expression of the solution of (39) using characteristic curves: for all $t \geq 0$, $x \in \mathbb{T}^d$, $v \in \mathbb{R}^d$ one has

$$f^{\text{tr}}(t, x, v) = f_0(\psi_t^{-1}(x, v)). \quad (42)$$

Conversely, if $(t, x, v) \in \mathbb{R}^+ \times \mathbb{T}^d \times \mathbb{R}^d \mapsto f^{\text{tr}}(t, x, v)$ is defined by (42) and if the initial value f_0 is of class \mathcal{C}^2 , then the stochastic process $(f^{\text{tr}}(t))_{t \geq 0}$ defined by (42) is a weak solution of the SPDE (40) (the Itô formulation is considered): for any smooth compactly supported function $\varphi : \mathbb{T}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$, one has

$$\begin{aligned} d\left(\iint \varphi(x, v) f(t, x, v) dx dv\right) &= \iint v \cdot \nabla_x \varphi(x, v) f(t, x, v) dx dv dt \\ &\quad + \iint E(x) \cdot \nabla_v \varphi(x, v) f(t, x, v) dx dv dt \\ &\quad + \iint \sum_{k=1}^K \sigma_k(x) \cdot \nabla_v \varphi(x, v) f(t, x, v) dx dv d\beta_k(t) \\ &\quad + \frac{1}{2} \iint \sum_{i,j=1}^d a_{i,j}(x) \partial_{v_i} \partial_{v_j} \varphi(x, v) f(t, x, v) dx dv dt. \end{aligned} \quad (43)$$

The proof of the identity (43) combines two arguments. First, since the diffeomorphism ψ_t preserves the volume of $\mathbb{T}^d \times \mathbb{R}^d$ for all $t \geq 0$, using (42) one has the identity

$$\begin{aligned} \iint \varphi(x, v) f(t, x, v) dx dv &= \iint \varphi(\psi_t(x, v)) f(0, x, v) dx dv \\ &= \iint \varphi(X_t^{x,v}, V_t^{x,v}) f(0, x, v) dx dv, \end{aligned}$$

where $(X_t^{x,v}, V_t^{x,v}) = \psi_t(x, v)$ is the solution at time $t \geq 0$ of (41) with initial values $X_0^{x,v} = x$ and $V_0^{x,v} = v$. It then remains to apply the standard Itô formula to $t \mapsto \varphi(X_t^{x,v}, V_t^{x,v})$ and to use integration by parts arguments combined with the identity above obtained by the first argument to obtain (43). The details are omitted. We refer to [24] for instance where this type of arguments are employed in the analysis of a class of nonlinear transport equations with transport noise.

Using the expression (42) of the solution of (39), it is straightforward to check that the preservation properties satisfied in the deterministic case by the solution $(f^{\text{det}}(t))_{t \geq 0}$ of the deterministic linear Vlasov equation (6) are also satisfied when a transport noise perturbation is applied. The proof is omitted.

Proposition 11. *Let $(f^{\text{tr}}(t))_{t \geq 0}$ be the solution of the SPDE (39) with (non-random) initial value f_0 . One has the following properties.*

- Preservation of positivity. Assume that $f_0(x, v) \geq 0$ for all $(x, v) \in \mathbb{T}^d \times \mathbb{R}^d$. Then one has $f^{\text{tr}}(t, x, v) \geq 0$ almost surely for all $t \geq 0$ and $(x, v) \in \mathbb{T}^d \times \mathbb{R}^d$.
- Preservation of integrals. Let $\Phi : \mathbb{R} \rightarrow \mathbb{R}^+$ be a real-valued measurable mapping. Then for all $t \geq 0$ one has almost surely

$$\iint \Phi(f^{\text{tr}}(t, x, v)) \, dx \, dv = \iint \Phi(f_0(x, v)) \, dx \, dv.$$

In particular, if $p \in [1, \infty)$ and if $f_0 \in L_{x,v}^p$, then for all $t \geq 0$ one has $f^{\text{tr}}(t) \in L_{x,v}^p$ almost surely and

$$\|f^{\text{tr}}(t)\|_{L_{x,v}^p} = \|f_0\|_{L_{x,v}^p}.$$

6.2. Splitting scheme. We are now in position to introduce the proposed numerical scheme for the discretization of the SPDE with transport noise (39). Like in the previous sections, a Lie–Trotter splitting strategy is applied. To deal with the stochastic perturbation in (39), one needs to consider the auxiliary stochastic subsystem

$$df(t, x, v) + \nabla_v f(t, x, v) \odot d\mathbf{W}(t, x) = 0, \quad (t, x, v) \in \mathbb{R}^+ \times \mathbb{T}^d \times \mathbb{R}^d.$$

This auxiliary stochastic subsystem can be solved exactly: for all $t \geq s \geq 0$ and all $(x, v) \in \mathbb{T}^d \times \mathbb{R}^d$ one has

$$f(t, x, v) = f(s, x, v - (\mathbf{W}(t, x) - \mathbf{W}(s, x))).$$

Alternative notation for solving the auxiliary stochastic subsystem above can be used: for all $t \geq s \geq 0$ one has

$$f(t) = T(\mathbf{W}(t) - \mathbf{W}(s))f(s) = T^d(\beta_d(t) - \beta_d(s)) \dots T^1(\beta_1(t) - \beta_1(s))f(s),$$

where for any mapping $\sigma : x \in \mathbb{T}^d \mapsto \sigma(x) \in \mathbb{R}^d$ one has

$$T(\sigma)f(x, v) = f(x, v - \sigma(x)), \quad x \in \mathbb{T}^d, v \in \mathbb{R}^d$$

and where the auxiliary linear operators $T^k(y)$, with $y \in \mathbb{R}$ and $1 \leq k \leq K$, are defined by

$$T^k(y)f(x, v) = f(x, v - y\sigma_k(x)), \quad x \in \mathbb{T}^d, v \in \mathbb{R}^d.$$

Using the Lie–Trotter integrator (12) for the deterministic part and combining the discretizations of the deterministic and stochastic parts yields the following scheme: given the initial value f_0 and the time-step size $\tau \in (0, 1)$, set $f_0^{\text{tr}} = f_0$ and for any nonnegative integer $n \geq 0$ set

$$f_{n+1}^{\text{tr}} = T(\delta\mathbf{W}_n)S^2(\tau)S^1(\tau)f_n^{\text{tr}} = T^d(\delta\beta_{n,d}) \circ \dots \circ T^1(\delta\beta_{n,1}) \circ S^2(\tau)S^1(\tau)f_n^{\text{tr}}, \quad n \geq 0. \quad (44)$$

where the Wiener increments $\delta\beta_{n,k}$ and $\delta\mathbf{W}_n$ are given by (3) and (5) respectively, see Section 2.

The Lie–Trotter splitting scheme (44) is consistent with Stratonovich interpretation of the noise in the SPDE (39), and satisfies the same properties as the exact solution stated in Proposition 11.

Proposition 12. *Let $(f_n^{\text{tr}})_{n \geq 0}$ be the solution of the Lie–Trotter splitting scheme (44) with initial value f_0 . One then has the following properties.*

- Preservation of positivity. *Assume that $f_0(x, v) \geq 0$ for all $(x, v) \in \mathbb{T}^d \times \mathbb{R}^d$. Then one has $f_n^{\text{tr}}(x, v) \geq 0$ almost surely for any nonnegative integer $n \geq 0$ and all $(x, v) \in \mathbb{T}^d \times \mathbb{R}^d$.*
- Preservation of integrals. *Let $\Phi : \mathbb{R} \rightarrow \mathbb{R}^+$ be a real-valued measurable mapping. Then for all $n \geq 0$ one has almost surely*

$$\iint \Phi(f_n^{\text{tr}}(x, v)) \, dx \, dv = \iint \Phi(f_0(x, v)) \, dx \, dv.$$

In particular, if $p \in [1, \infty)$ and if $f_0 \in L_{x,v}^p$, then for all $n \geq 0$ one has $f_n^{\text{tr}} \in L_{x,v}^p$ almost surely and

$$\|f_n^{\text{tr}}\|_{L_{x,v}^p} = \|f_0\|_{L_{x,v}^p}$$

almost surely.

Proof of Proposition 12. All the results are proved by recursion, using the fact that the operators $T(\delta\mathbf{W}_n)$, $S^2(\tau)$ and $S^1(\tau)$ satisfy the considered properties. \square

Like for the splitting scheme (12) defined in Section 3 for the discretization of the deterministic linear Vlasov equation (6), the Lie–Trotter splitting scheme (44) can be interpreted as a discrete version of the expression (42) for the exact solution of (39), where the stochastic flow ψ_t is approximated using a splitting integrator applied to the SDE system (41).

Remark 13. If the noise in the SPDE (39) is a purely temporal Wiener process, i.e. $\mathbf{W}(t, x) = \mathbf{W}(t) = \sum_{k=1}^K \beta_k(t) \boldsymbol{\sigma}_k$ for all $(t, x) \in \mathbb{R}^+ \times \mathbb{T}^d$ where $\boldsymbol{\sigma}_1, \dots, \boldsymbol{\sigma}_K$ are elements of \mathbb{R}^d , then the exact solution of (39) and the numerical solution given by (44) can be written

$$f^{\text{tr}}(t) = T(\mathbf{W}(t)) f^{\text{det}}(t), \quad t \geq 0; \quad f_n^{\text{tr}} = T(\mathbf{W}(t_n)) f_n^{\text{det}}, \quad n \geq 0,$$

where the auxiliary continuous-time process $(f^{\text{det}}(t))_{t \geq 0}$ is the solution of the deterministic PDE (6) and the auxiliary continuous-time process $(f_n^{\text{det}})_{n \geq 0}$ is given by the deterministic Lie–Trotter splitting scheme (12).

In that situation, Propositions 11 and 12 are straightforward consequences of the results described for the deterministic problem in Section 3.

6.3. Numerical experiments. We begin the numerical experiments by illustrating the behavior of the linear Vlasov equation perturbed by transport noise (39), in dimension $d = 1$. In all the experiments below, the initial value f_0 is given by (13) and the vector field E is given by (14). The discretization parameters are given by $\delta x = \frac{1}{400}$, $\delta v = \frac{4\pi}{800}$, and $\tau = 0.1$. The snapshots of the numerical solution at times $\{0, 0.5, 1, 1.5, 2, 2.5\}$ computed using the splitting scheme (44) are provided in Figure 14, with $K = 1$ and diffusion coefficient $\sigma_1(x, v) = 0.5$. One observes that the solution remains nonnegative, which illustrates the positivity preserving property stated in Proposition 12 on the considered realization.




FIGURE 14. Snapshots: approximation of the solution of the stochastic PDE with transport noise (39) with initial value f_0 given by (13), with $\sigma_1(x, v) = 0.5$ at times $\{0, 0.5, 1, 1.5, 2, 2.5\}$, using the Lie–Trotter splitting scheme (44) with time-step size $\tau = 0.1$.

Next, in Figure 15, we illustrate the almost sure preservation of the L^1 , L^3 and L^{55} norms when the Lie–Trotter splitting scheme (44) is applied to the SPDE (39). This is performed for three independent realizations. The discretization parameters are chosen to be: $\delta x = \frac{1}{3000}$, $\delta v = \frac{4\pi}{3000}$, and $\tau = 0.1$. The final time is $T = 1$. In this figure, we observe the preservation of the norms, which illustrates Proposition 12.




FIGURE 15. Evolution of the $L^1_{x,v}$, $L^3_{x,v}$ and $L^{55}_{x,v}$ norms: illustration of Proposition 12 when applying the Lie–Trotter scheme (44) to the SPDE with transport noise (39), with time-step size $\tau = 0.1$.

We conclude these numerical experiments in the transport noise case by investigating the mean-square order of convergence of the Lie–Trotter splitting scheme (44).

The same procedure as in the previous cases is applied. A reference solution is computed using the splitting scheme with time-step size $\tau_{\text{ref}} = 2^{-14}$, and the errors are computed when the time-step size τ takes values in $\{2^{-7}, \dots, 2^{-13}\}$. The expectation is computed using a Monte Carlo averaging procedure over 500 independent samples. The discretization parameters are $\delta x = \frac{1}{100}$, $\delta v = \frac{4\pi}{200}$. The final time is $T = 0.5$. The results are presented in a loglog plot in Figure 16. We observe a mean-square convergence order of at least $1/2$.




FIGURE 16. Mean-square errors: Lie–Trotter scheme (44) applied to the SPDE (39) driven by one noise ($K = 1$, $\sigma_1(x, v) = 0.5$).

7. Conclusion and perspectives. In this paper, we numerically investigate the behaviour of solutions of linear Vlasov partial differential equations perturbed by Wiener noise, which may be additive noise, multiplicative Itô and Stratonovich noise, and transport noise. For this purpose, we apply Lie–Trotter splitting integrators for the temporal discretization of these SPDEs. We show that these time integrators are able to preserve key qualitative properties of the exact solutions, such as preserving norms or ensuring positivity. Several promising theoretical results are provided and illustrated numerically.

We leave some fundamental questions open for future work. Let us mention a few of them.

1. Based on the numerical experiments of this paper, we conjecture that the order of mean-square convergence of the Lie–Trotter splitting scheme is equal to 1 for all the considered types of noise. Thus the main perspective for a possible future work is to prove mean-square error bounds.
2. We have only considered the behavior of the temporal discretization error. It will be interesting to study the full discretization error, and in particular whether some conditions on the time-step size τ and the mesh sizes δx and δv need to be imposed for stability or accuracy reasons.
3. Constructing higher-order methods could also be an interesting question. For instance, in the context of SPDEs with small noise, it may be possible to apply a Strang splitting strategy to deal with the deterministic subsystems and obtain better rates of convergence.

4. We have only considered some simple linear Vlasov equations perturbed by additive or linear noise. It may be interesting to study more complex problems, for instance with nonlinear terms.

Appendix A. The Itô–Wentzell formula. Let $D, K \in \mathbb{N}$ be two integers. Let β_1, \dots, β_K be independent standard real-valued Wiener processes, and let $b_j : \mathbb{R}^D \rightarrow \mathbb{R}$ and $\varsigma_{j,k} : \mathbb{R}^D \rightarrow \mathbb{R}$, $1 \leq j \leq D$, $1 \leq k \leq K$, be Lipschitz continuous mappings.

Assume that the \mathbb{R}^D -valued stochastic process $(\xi(t))_{t \geq 0}$ is solution of the stochastic differential equation

$$d\xi_j(t) = b_j(\xi(t)) dt + \sum_{k=1}^K \varsigma_{j,k}(\xi(t)) d\beta_k(t), \quad 1 \leq j \leq D, \quad t \geq 0.$$

Let $G : (t, \xi) \in \mathbb{R}^+ \times \mathbb{R}^D \mapsto G(t, \xi) \in \mathbb{R}$ be a stochastic process, such that for all $\xi \in \mathbb{R}^D$ and $t \geq 0$ one has

$$dG(t, \xi) = J(t, \xi) dt + \sum_{k=1}^K H_k(t, \xi) d\beta_k(t),$$

where $J : (t, \xi) \in \mathbb{R}^+ \times \mathbb{R}^D \mapsto J(t, \xi) \in \mathbb{R}$ and $H_k : (t, \xi) \in \mathbb{R}^+ \times \mathbb{R}^D \mapsto H_k(t, \xi) \in \mathbb{R}$, $1 \leq k \leq K$, are stochastic processes. Assume that G is almost surely of class $\mathcal{C}^{0,2}$.

Then the real-valued stochastic processes $t \mapsto G(t, \xi(t))$ is solution of the stochastic differential equation

$$\begin{aligned} dG(t, \xi(t)) &= J(t, \xi(t)) dt + \sum_{k=1}^K H_k(t, \xi(t)) d\beta_k(t) \\ &+ \sum_{j=1}^d b_j(\xi(t)) \partial_{\xi_j} G(t, \xi(t)) dt + \sum_{j=1}^d \sum_{k=1}^K \varsigma_{j,k}(t, \xi(t)) \partial_{x_j} G(t, \xi(t)) d\beta_k(t) \\ &+ \frac{1}{2} \sum_{i,j=1}^d \sum_{k=1}^K \varsigma_{i,k}(t, \xi(t)) \varsigma_{j,k}(t, \xi(t)) \partial_{x_i} \partial_{x_j} G(t, \xi(t)) dt \\ &+ \sum_{j=1}^d \sum_{k=1}^K \varsigma_{j,k}(t, \xi(t)) \partial_{\xi_j} H_k(t, \xi(t)) dt. \end{aligned}$$

We refer to [55, Theorem 1.17] for a statement of the formula above.

Acknowledgments. We thank the referees and the associated editor for helpful comments and suggestions on the initial version of this work.

The numerical experiments have been performed by adapting the code from [46] to take into account the considered stochastic perturbations.

We thank Simone Calogero and Lukas Einkemmer for interesting discussions. The work of CEB is partially supported by the project SIMALIN (ANR-19-CE40-0016) operated by the French National Research Agency. The work of DC is partially supported by the Swedish Research Council (VR) (project no. 2018-04443). The computations were performed on resources provided by the National Academic Infrastructure for Supercomputing in Sweden (NAISS) at UPPMAX, Uppsala University partially funded by the Swedish Research Council through grant agreement no. 2022-06725.

REFERENCES

- [1] R. Anton and D. Cohen, [Exponential integrators for stochastic Schrödinger equations driven by Itô noise](#), *J. Comput. Math.*, **36** (2018), 276-309,
- [2] N. Ayi and E. Faou, [Analysis of an asymptotic preserving scheme for stochastic linear kinetic equations in the diffusion limit](#), *SIAM/ASA J. Uncertain. Quantif.*, **7** (2019), 760-785,
- [3] C. Bayer and H. Oberhauser, Splitting methods for SPDEs: From robustness to financial engineering, optimal control, and nonlinear filtering, in *Splitting Methods in Communication, Imaging, Science, and Engineering*, Sci. Comput., Springer, Cham, (2016), 499-539.
- [4] J. Bedrossian and S. Papathanasiou, *The Vlasov-Poisson and Vlasov-Poisson-Fokker-Planck, Systems in Stochastic Electromagnetic Fields: Local Well-Posedness*, 2022.
- [5] A. Berg, D. Cohen and G. Dujardin, [Lie-Trotter splitting for the nonlinear stochastic Manakov system](#), *J. Sci. Comput.*, **88** (2021), Paper No. 6, 31 pp.
- [6] P. Bertrand, M. Albrecht-Marc, T. Réveillé and A. Ghizzo, [Vlasov models for laser-plasma interaction](#), *Transport Theory and Statistical Physics*, **34** (2005), 103-126,
- [7] N. Besse, [Convergence of a semi-Lagrangian scheme for the one-dimensional Vlasov-Poisson system](#), *SIAM J. Numer. Anal.*, **42** (2004), 350-382.
- [8] N. Besse and M. Mehrenberger, [Convergence of classes of high-order semi-Lagrangian schemes for the Vlasov-Poisson system](#), *Math. Comp.*, **77** (2008), 93-123,
- [9] J. A. Bittencourt, *Fundamentals of Plasma Physics*, Pergamon Press, Oxford, 1986.
- [10] S. Blanes and F. Casas, *A Concise Introduction to Geometric Numerical Integration*, Monographs and Research Notes in Mathematics, CRC Press, Boca Raton, FL, 2016.
- [11] S. Blanes, F. Casas and A. Murua, *Splitting Methods for Differential Equations*, 2024.
- [12] C.-E. Bréhier and D. Cohen, [Strong rates of convergence of a splitting scheme for Schrödinger equations with nonlocal interaction cubic nonlinearity and white noise dispersion](#), *SIAM/ASA J. Uncertain. Quantif.*, **10** (2022), 453-480,
- [13] C.-E. Bréhier and D. Cohen, [Analysis of a splitting scheme for a class of nonlinear stochastic Schrödinger equations](#), *Appl. Numer. Math.*, **186** (2023), 57-83.
- [14] C.-E. Bréhier, D. Cohen and G. Giordano, *Splitting Schemes for FitzHugh-Nagumo Stochastic Partial Differential Equations*, 2023.
- [15] C.-E. Bréhier, D. Cohen and J. Ulander, Analysis of a positivity-preserving splitting scheme for some nonlinear stochastic heat equations, 2023.
- [16] C.-E. Bréhier, J. Cui and J. Hong, [Strong convergence rates of semidiscrete splitting approximations for the stochastic Allen-Cahn equation](#), *IMA J. Numer. Anal.*, **39** (2019), 2096-2134,
- [17] C.-E. Bréhier and L. Goudenège, [Weak convergence rates of splitting schemes for the stochastic Allen-Cahn equation](#), *BIT*, **60** (2020), 543-582.
- [18] P. M. Burrage and K. Burrage, [Structure-preserving Runge-Kutta methods for stochastic Hamiltonian equations with additive noise](#), *Numer. Algorithms*, **65** (2014), 519-532.
- [19] F. Casas, N. Crouseilles, E. Faou and M. Mehrenberger, [High-order Hamiltonian splitting for the Vlasov-Poisson equations](#), *Numer. Math.*, **135** (2017), 769-801.
- [20] F. Charles, B. Després and M. Mehrenberger, [Enhanced convergence estimates for semi-Lagrangian schemes: application to the Vlasov-Poisson equation](#), *SIAM J. Numer. Anal.*, **51** (2013), 840-863.
- [21] C. Chen, D. Cohen, R. D'Ambrosio and A. Lang, [Drift-preserving numerical integrators for stochastic Hamiltonian systems](#), *Adv. Comput. Math.*, **46** (2020), Paper No. 27, 22.
- [22] C. Z. Cheng and G. Knorr, The integration of the Vlasov equation in configuration space, *Journal of Computational Physics*, **22** (1976), 330-351, <https://www.sciencedirect.com/science/article/pii/002199917690053X>.
- [23] L. Cheung and W. Horton, Equilibrium and electrostatic stability theory of tokamaks from the drift-kinetic equation, *Annals of Physics*, **81** (1973), 201-230, <https://www.sciencedirect.com/science/article/pii/0003491673904867>.
- [24] M. Coghi and F. Flandoli, [Propagation of chaos for interacting particles subject to environmental noise](#), *Ann. Appl. Probab.*, **26** (2016), 1407-1442.
- [25] D. Cohen, J. Cui, J. Hong and L. Sun, [Exponential integrators for stochastic Maxwell's equations driven by Itô noise](#), *J. Comput. Phys.*, **410** (2020), 109382, 21 pp.
- [26] D. Cohen, S. Larsson and M. Sigg, [A trigonometric method for the linear stochastic wave equation](#), *SIAM J. Numer. Anal.*, **51** (2013), 204-222.
- [27] D. Cohen and G. Vilmart, [Drift-preserving numerical integrators for stochastic Poisson systems](#), *Int. J. Comput. Math.*, **99** (2022), 4-20.

- [28] G.-H. Cottet and P.-A. Raviart, [Particle methods for the one-dimensional Vlasov-Poisson equations](#), *SIAM J. Numer. Anal.*, **21** (1984), 52-76.
- [29] S. Cox and J. van Neerven, [Convergence rates of the splitting scheme for parabolic linear stochastic Cauchy problems](#), *SIAM J. Numer. Anal.*, **48** (2010), 428-451.
- [30] N. Crouseilles, L. Einkemmer and E. Faou, [Hamiltonian splitting for the Vlasov-Maxwell equations](#), *J. Comput. Phys.*, **283** (2015), 224-240.
- [31] N. Crouseilles, M. Mehrenberger and E. Sonnendrücker, [Conservative semi-Lagrangian schemes for Vlasov equations](#), *J. Comput. Phys.*, **229** (2010), 1927-1953.
- [32] J. Cui, J. Hong, Z. Liu and W. Zhou, [Strong convergence rate of splitting schemes for stochastic nonlinear Schrödinger equations](#), *J. Differential Equations*, **266** (2019), 5625-5663.
- [33] G. Da Prato and J. Zabczyk, *Stochastic Equations in Infinite Dimensions*, 2nd edition, Cambridge University Press, Cambridge, 2014.
- [34] F. Delarue, F. Flandoli and D. Vincenzi, [Noise prevents collapse of Vlasov-Poisson point charges](#), *Comm. Pure Appl. Math.*, **67** (2014), 1700-1736.
- [35] J. Denavit and W. L. Kruer, [Comparison of numerical solutions of the vlasov equation with particle simulations of collisionless plasmas](#), *The Physics of Fluids*, **14** (1971), 1782-1791.
- [36] G. Dimarco and L. Pareschi, [Numerical methods for kinetic equations](#), *Acta Numer.*, **23** (2014), 369-520.
- [37] D. R. Durran, *Numerical Methods for Wave Equations in Geophysical Fluid Dynamics*, vol. 32 of Texts in Applied Mathematics, Springer-Verlag, New York, 1999.
- [38] L. Einkemmer, [Semi-Lagrangian Vlasov simulation on GPUs](#), *Comput. Phys. Commun.*, **254** (2020), 107351, 12 pp.
- [39] L. Einkemmer and A. Ostermann, [Convergence analysis of Strang splitting for Vlasov-type equations](#), *SIAM J. Numer. Anal.*, **52** (2014), 140-155.
- [40] E. Fedrizzi, F. Flandoli, E. Priola and J. Vovelle, [Regularity of stochastic kinetic equations](#), *Electron. J. Probab.*, **22** (2017), Paper No. 48, 42.
- [41] F. Filbet and E. Sonnendrücker, [Comparison of Eulerian Vlasov solvers](#), *Comput. Phys. Comm.*, **150** (2003), 247-266.
- [42] R. T. Glassey, *The Cauchy Problem in Kinetic Theory*, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 1996.
- [43] E. Hairer, C. Lubich and G. Wanner, *Geometric Numerical Integration*, vol. 31 of Springer Series in Computational Mathematics, Springer Ser. Comput. Math., 31 Springer, Heidelberg, 2010.
- [44] J. Hong, R. Scherer and L. Wang, [Midpoint rule for a linear stochastic oscillator with additive noise](#), *Neural Parallel Sci. Comput.*, **14** (2006), 1-12.
- [45] G. Knorr, Plasma simulation with few particles, *Journal of Computational Physics*, **13** (1973), 165-180, <https://www.sciencedirect.com/science/article/pii/0021999173900211>.
- [46] V. Kozhevnikov, *ChengKnorrSonnerdrucker*, MATLAB Central File Exchange, 2024, https://www.mathworks.com/matlabcentral/fileexchange/50487-cheng_knorr_sonnerdrucker.
- [47] M. Kraus and T. M. Tyranowski, [Variational integrators for stochastic dissipative Hamiltonian systems](#), *IMA J. Numer. Anal.*, **41** (2021), 1318-1367.
- [48] H. Kunita, *Stochastic Flows and Stochastic Differential Equations*, vol. 24 of Cambridge Studies in Advanced Mathematics, Cambridge University Press, Cambridge, 1997.
- [49] G. J. Lord, C. E. Powell and T. Shardlow, *An Introduction to Computational Stochastic PDEs*, Cambridge Texts in Applied Mathematics, Cambridge University Press, New York, 2014.
- [50] R. Marty, [On a splitting scheme for the nonlinear Schrödinger equation in a random medium](#), *Commun. Math. Sci.*, **4** (2006), 679-705.
- [51] R. I. McLachlan and G. R. W. Quispel, [Splitting methods](#), *Acta Numer.*, **11** (2002), 341-434.
- [52] G. N. Milstein and M. V. Tretyakov, *Stochastic Numerics for Mathematical Physics*, 2nd edition, Scientific Computation, Springer, Cham, 2021.
- [53] J. L. Padgett and Q. Sheng, [Convergence of an operator splitting scheme for abstract stochastic evolution equations](#), in *Advances in Mathematical Methods and High Performance Computing*, **41** (2019), 163-179.
- [54] G. Rein, [Collisionless kinetic equations from astrophysics—the Vlasov-Poisson system](#), in *Handb. Differ. Equ. Elsevier/North-Holland, Amsterdam*, **3** (2007), 383-476.
- [55] B. L. Rozovsky and S. V. Lototsky, *Stochastic Evolution Systems*, 2nd edition, Probab. Theory Stoch. Model., 89 Springer, Cham, 2018.

- [56] J. Schaeffer, [Higher order time splitting for the linear Vlasov equation](#), *SIAM J. Numer. Anal.*, **47** (2009), 2203-2223.
- [57] H. Schurz, [Analysis and discretization of semi-linear stochastic wave equations with cubic nonlinearity and additive space-time noise](#), *Discrete Contin. Dyn. Syst. Ser. S*, **1** (2008), 353-363.
- [58] M. J. Senosiain and A. Tocino, [A review on numerical schemes for solving a linear stochastic oscillator](#), *BIT*, **55** (2015), 515-529.
- [59] E. Sonnendrücker, J. Roche, P. Bertrand and A. Ghizzo, [The semi-Lagrangian method for the numerical resolution of the Vlasov equation](#), *J. Comput. Phys.*, **149** (1999), 201-220.
- [60] A. H. Strømmen Melbø and D. J. Higham, [Numerical simulation of a linear stochastic oscillator with additive noise](#), *Appl. Numer. Math.*, **51** (2004), 89-99.
- [61] C. Villani, [A review of mathematical topics in collisional kinetic theory](#), in *Handbook of Mathematical Fluid Dynamics*, **1** (2002), 71-305.
- [62] A. A. Vlasov, *Many-Particle Theory and Its Application to Plasma*, Probab. Theory Stoch. Model., 89 Springer, Cham, 2018.

Received July 2023; revised January 2024; early access March 2024.