

Wilson Lines in the Abelian Lattice Higgs Model

Downloaded from: https://research.chalmers.se, 2024-12-22 17:10 UTC

Citation for the original published paper (version of record):

Palö Forsström, M. (2024). Wilson Lines in the Abelian Lattice Higgs Model. Communications in Mathematical Physics, 405(11). http://dx.doi.org/10.1007/s00220-024-05128-x

N.B. When citing this work, cite the original published paper.

research.chalmers.se offers the possibility of retrieving research publications produced at Chalmers University of Technology. It covers all kind of research output: articles, dissertations, conference papers, reports etc. since 2004. research.chalmers.se is administrated and maintained by Chalmers Library



Communications in

Mathematical

Wilson Lines in the Abelian Lattice Higgs Model

Malin P. Forsström

Mathematical Sciences, Chalmers University of Technology and University of Gothenburg, 41296 Gothenburg, Sweden. E-mail: palo@chalmers.se

Received: 18 November 2021 / Accepted: 21 August 2024 Published online: 5 November 2024 – © The Author(s) 2024

Abstract: Lattice gauge theories are lattice approximations of the Yang–Mills theory in physics. The abelian lattice Higgs model is one of the simplest examples of a lattice gauge theory interacting with an external field. In a previous paper (Forsström et al. in Math Phys 4(2):257–329, 2023), we calculated the leading order term of the expected value of Wilson loop observables in the low-temperature regime of the abelian lattice Higgs model on \mathbb{Z}^4 , with structure group $G = \mathbb{Z}_n$ for some $n \ge 2$. In the absence of a Higgs field, these are important observables since they exhibit a phase transition which can be interpreted as distinguishing between regions with and without quark confinement. However, in the presence of a Higgs field, this is no longer the case, and a more relevant family of observables are so-called open Wilson lines. In this paper, we extend and refine the ideas introduced in Forsström et al. (Math Phys 4(2):257–329, 2023) to calculate the leading order term of the expected value of the more general Wilson line observables. Using our main result, we then calculate the leading order term of several natural ratios of expected values and confirm the behavior predicted by physicists.

1. Introduction

1.1. Background. Lattice gauge theories are spin models which describe the interaction of elementary particles. These were first introduced by Wilson [32] as lattice approximations of the quantum field theories that appear in the standard model, known as Yang-Mills theory. Since then, lattice gauge theories have been successfully used to understand the corresponding continuous models, and several of the predictions made using these lattice approximations have been verified experimentally. At about the same time as lattice gauge theories were introduced in the physics literature by Wilson, Wegner [31] introduced what he then called *generalized Ising models* as an example of a family of models with a phase transition without a local order parameter. In special cases, these generalized Ising models are lattice gauge theories, and as such, they have

been used extensively as toy models for the lattice gauge theories that are more relevant for physics.

In the last couple of years, there has been a renewed interest in the rigorous analysis of four-dimensional lattice gauge theories in the mathematical community, see, e.g., [7–9,16,17,21]. Most relevant for this work are the papers [7,9,17], in which the leading order term for the expectation of Wilson loop observables was computed for lattice gauge theories with Wilson action and finite structure groups.

Pure gauge theories model only the gauge field itself, and to advance towards physically relevant theories; it is necessary also to understand models that include external fields interacting with the gauge field, see, e.g., [19, 30]. In this paper, we consider a lattice gauge theory that models a gauge field coupled to a scalar Bosonic field with a quartic Higgs potential. The resulting model is called the *lattice Higgs model*. This model has received significant attention in the physics community. Some examples are the works [1–3], where calculations to obtain critical parameters of these models were performed, and [27,28], in which phase diagrams were sketched. For further background, as well as more references, we refer the reader to [19] and [30].

In a recent paper [18], we extended the theory developed in [7,9,17] in order to describe the leading order term for the expectation of Wilson loop observables in the fixed length and low-temperature regime of the abelian Higgs model. Wilson loop expectations are natural observables in lattice gauge theories and were introduced by Wilson as a means to detect whether quark confinement occurs, see [32]. In lattice gauge theories without matter fields, one can show that the expected value of large Wilson loops undergo a phase transition, where it changes from following a so-called area law to following a so-called perimeter law. However, as discussed in, e.g., [29], in gauge theories with matter fields, the Wilson loop observable obeys a perimeter law for all parameters, and hence one cannot see a relevant phase transition using only the Wilson loop observable. For this reason, alternative observables have been suggested for studying the lattice Higgs model. One such observable is the open Wilson line observable, in which the loop in the Wilson loop observable is replaced by an open path that is saturated at the end-points by the Higgs field. This type of observable has been relatively well studied in the physics literature (see, e.g., [5,6,13,15,23,24,29,30]). Moreover, the asymptotic behavior of such observables has been argued to be related to, e.g., the absence of bound states of the charged particle in the presence of an external source [13], confinement versus deconfinement in lattice gauge theories with matter fields [6], and binding versus unbinding of dynamical quarks in the field of a static color source [6]. Hence the Wilson line observables are of physical relevance.

1.2. Preliminary notation. For $m \ge 2$, the graph naturally associated to \mathbb{Z}^m has a vertex at each point $x \in \mathbb{Z}^m$ with integer coordinates and a non-oriented edge between nearest neighbors. We will work with oriented edges throughout this paper, and for this reason we associate to each non-oriented edge \bar{e} two oriented edges e_1 and $e_2 = -e_1$ with the same endpoints as \bar{e} and opposite orientations.

Let $d\mathbf{e}_1 := (1, 0, 0, ..., 0)$, $d\mathbf{e}_2 := (0, 1, 0, ..., 0)$, ..., $d\mathbf{e}_m := (0, ..., 0, 1)$ be oriented edges corresponding to the unit vectors in \mathbb{Z}^m . We say that an oriented edge e is *positively oriented* if it is equal to a translation of one of these unit vectors, i.e., if there is a $v \in \mathbb{Z}^m$ and a $j \in \{1, 2, ..., m\}$ such that $e = v + d\mathbf{e}_j$. If $v \in \mathbb{Z}^m$ and $j_1 < j_2$, then $p = (v + d\mathbf{e}_{j_1}) \land (v + d\mathbf{e}_{j_2})$ is a positively oriented 2-cell, also known as a *positively oriented plaquette*. We let $C_0(\mathbb{Z}^4)$, $C_1(\mathbb{Z}^4)$, and $C_2(\mathbb{Z}^4)$ denote the sets of oriented vertices, edges, and plaquettes. Next, we let B_N denote the set $[-N, N]^m \subseteq \mathbb{Z}^m$, and we let $C_0(B_N)$, $C_1(B_N)$, and $C_2(B_N)$ denote the sets of oriented vertices, edges, and plaquettes, respectively, whose end-points are all in B_N .

Whenever we talk about a lattice gauge theory we do so with respect to some (abelian) group (G, +), referred to as the *structure group*. We also fix a unitary and faithful representation ρ of (G, +). In this paper, we will always assume that $G = \mathbb{Z}_n$ for some $n \ge 2$ with the group operation + given by standard addition modulo n. Also, we will assume that ρ is a one-dimensional representation of G. We note that a natural such representation is given by $j \mapsto e^{j \cdot 2\pi i/n}$.

Now assume that a structure group (G, +), a one-dimensional unitary representation ρ of (G, +), and an integer $N \ge 1$ are given. We let $\Omega^1(B_N, G)$ denote the set of all *G*-valued 1-forms σ on $C_1(B_N)$, i.e., the set of all *G*-valued functions $\sigma : e \mapsto \sigma(e)$ on $C_1(B_N)$ such that $\sigma(e) = -\sigma(-e)$ for all $e \in C_1(B_N)$. Similarly, we let $\Omega^0(B_N, G)$ denote the set of all *G*-valued functions $\phi : x \mapsto \phi(x)$ on $C_0(B_N)$ which are such that $\phi(x) = -\phi(-x)$ for all $x \in C_1(B_N)$. When $\sigma \in \Omega^1(B_N, G)$ and $p \in C_2(B_N)$, we let ∂p denote the formal sum of the four edges e_1 , e_2 , e_3 , and e_4 in the oriented boundary of p (see Sect. 2.1.5), and define

$$d\sigma(p) := \sigma(\partial p) := \sum_{e \in \partial p} \sigma(e) := \sigma(e_1) + \sigma(e_2) + \sigma(e_3) + \sigma(e_4).$$

Similarly, when $\phi \in \Omega^0(B_N, G)$ and $e \in C_1(B_N)$ is an edge from x_1 to x_2 , we let ∂e denote the formal sum $x_2 - x_1$, and define $d\phi(e) := \phi(\partial e) := \phi(x_2) - \phi(x_1)$.

1.3. The abelian lattice Higgs model. Given $\beta, \kappa, \zeta \ge 0$, the action $S_{N,\beta,\kappa,\zeta}$ for lattice gauge theory with Wilson action coupled to a Higgs field on B_N is, for $\sigma \in \Omega^1(E_N, G)$, $\phi \in \Omega^0(B_N, G)$, and a symmetric function $r: C_0(B_N) \to \mathbb{R}_+$, defined by

$$S_{N,\beta,\kappa,\zeta}(\sigma,\phi,r) := -\beta \sum_{p \in C_2(B_N)} \rho(d\sigma(p)) - \kappa \sum_{\substack{e \in C_1(B_N):\\\partial e = y - x}} r(x)r(y)\rho(\sigma(e) - \phi(\partial e)) + \zeta \sum_{x \in C_0(B_N)} (r(x)^2 - 1)^2 + \sum_{x \in C_0(B_N)} r(x)^2.$$

$$(1.1)$$

The first term on the right hand side of (1.1) is referred to as the *Wilson action functional* for pure gauge theory (see, e.g., [32]), the second term on the right hand side of (1.1) is referred to as the interaction term, and the third and fourth term on the right hand side of (1.1) together are referred to as a *sombrero potential*. Since $\phi \in \Omega^0(B_N, G)$ and $\sigma \in \Omega^1(B_N, G)$, the action $S_{N,\beta,\kappa,\zeta}(\sigma, \phi, r)$ is real for all σ, ϕ , and r. Elements $\sigma \in \Omega^1(B_N, G)$ will be referred to as *gauge field configurations*, and pairs (ϕ, r) , with $\psi \in \Omega^0(B_N, G)$ and $r : C_0(B_N) \to \mathbb{R}_+$ symmetric, will be referred to as *Higgs field configurations*. The quantity β is known as the *gauge coupling constant*, κ is known as the *hopping parameter*, and ζ is known as the *quartic Higgs self coupling*.

The Gibbs measure corresponding to the action $S_{N,\beta,\kappa,\zeta}$ is given by

$$d\mu_{N,\beta,\kappa,\zeta}(\sigma,\phi,r) = Z_{N,\beta,\kappa,\zeta}^{-1} e^{-S_{N,\beta,\kappa,\zeta}(\sigma,\phi,r)} \prod_{e \in C_1(B_N)^+} d\mu_G(\sigma(e))$$
$$\prod_{x \in C_0(B_N)^+} d\mu_G(\phi(x)) d\mu_{\mathbb{R}_+}(r(x)),$$

where $C_1(B_N)^+$ denotes the set of positively oriented edges in $C_1(B_N)$, $d\mu_G$ is the uniform measure on G, and $\mu_{\mathbb{R}_+}$ is the Lebesgue measure on \mathbb{R}_+ . We refer to this lattice gauge theory as the *abelian lattice Higgs model*.

We will work with the model obtained from this action in the *fixed length limit* $\zeta \to \infty$, in which the radial component of the Higgs field concentrates at one. In the physics literature, this is sometimes called the *London limit*. We do not discuss the limit of the Gibbs measure corresponding to $S_{N,\beta,\kappa,\zeta}$ as $\zeta \to \infty$ here, but simply from the outset adopt the action resulting from only considering $r: C_0(B_N) \to \mathbb{R}_+$ with r(x) = 1 for all $x \in C_0(B_N)$. In this case, for $\sigma \in \Omega^1(B_N, G)$, and $\phi \in \Omega^0(B_N, G)$, we obtain the action

$$S_{N,\beta,\kappa,\infty}(\sigma,\phi) := -\beta \sum_{p \in C_2(B_N)} \rho(d\sigma(p)) - \kappa \sum_{\substack{e \in C_1(B_N):\\ \partial e = y - x}} \rho(\sigma(e) - \phi(\partial e))$$

We then consider a corresponding probability measure $\mu_{N,\beta,\kappa,\infty}$ on $\Omega^1(B_N, G) \times \Omega^0(B_N, G)$ given by

$$\mu_{N,\beta,\kappa,\infty}(\sigma,\phi) := Z_{N,\beta,\kappa,\infty}^{-1} e^{-S_{N,\beta,\kappa,\infty}(\sigma,\phi)}, \qquad \sigma \in \Omega^1(B_N,G), \ \phi \in \Omega^0(B_N,G),$$

where $Z_{N,\beta,\kappa,\infty}$ is a normalizing constant. This is the *fixed length lattice Higgs model*. We let $\mathbb{E}_{N,\beta,\kappa,\infty}$ denote the corresponding expectation. Whenever $f: \Omega^1(B_M, G) \times \Omega^0(B_M, G) \to \mathbb{R}$ for some $M \ge 1$, then, as a consequence of the Ginibre inequalities (see Sect. 2.6), the infinite volume limit

$$\left\langle f(\sigma,\phi) \right\rangle_{\beta,\kappa,\infty} := \lim_{N \to \infty} \mathbb{E}_{N,\beta,\kappa,\infty} \left[f(\sigma,\phi) \right]$$

exists, and it is this limit that we will use in our main result.

1.4. Wilson loops and Wilson lines. For $k \in \{0, 1, ..., m, a k$ -chain is a formal sum of positively oriented k-cells with integer coefficients, see Sect. 2.1.4 below. The support of a 1-chain γ , written supp γ , is the set of directed edges with non-zero coefficient in γ . We say that a 1-chain with finite support is a *generalized loop* if it has coefficients in $\{-1, 0, 1\}$ and empty boundary, see Definition 2.13. Roughly speaking, this means that a generalized loop is a disjoint union of a finite number of closed loops, where each closed loop is a nearest-neighbor path in the graph \mathbb{Z}^4 starting and ending at the same vertex. For example, any rectangular loop, as well as any finite disjoint union of such loops, is a generalized loop. We say that a 1-chain with finite support is an *open path* from $x_1 \in \Omega_0^+(B_N)$ to $x_2 \in \Omega_2^+(B_N)$ if it has coefficients in $\{-1, 0, 1\}$ and boundary $\partial \gamma := x_2 - x_1$. If γ is either an open path or a generalized loop, we refer to γ as a *path*.

Given a path γ , the Wilson line observable $L_{\gamma}(\sigma, \phi)$ is defined by

 $L_{\gamma}(\sigma,\phi) := \rho \big(\sigma(\gamma) - \phi(\partial \gamma) \big), \qquad \sigma \in \Omega^{1}(B_{N},G), \ \phi \in \Omega^{0}(B_{N},G),$

where $\sigma(\gamma) := \sum_{e \in \gamma} \sigma(e)$, and $\phi(\partial \gamma) = \phi(x_2) - \phi(x_1)$ if γ is an open path from x_1 to x_2 , and $\phi(\partial \gamma) = 0$ if the boundary of γ is empty. If γ is a generalized loop, then $W_{\gamma}(\sigma) := L_{\gamma}(\sigma, \phi)$ is referred to as a *Wilson loop observable*.

1.5. Main results.

Theorem 1.1. Consider the fixed length lattice Higgs model on \mathbb{Z}^4 , with structure group $G = \mathbb{Z}_2$, and representation $\rho: G \to \mathbb{C}$ given by $\rho(0) = 1$ and $\rho(1) = -1$.

Let $\beta, \kappa \geq 0$ be such that $18^2 e^{-4\kappa} (2 + e^{-4\kappa}) < 1$ and $6\beta > \kappa$. Further, let γ be a path along the boundary of a rectangle with side lengths $\ell_1, \ell_2 \geq 8$, and assume that $|\operatorname{supp} \gamma| \geq 24$. Finally, let $e \in C_1(\mathbb{Z}^4)$ be arbitrary.

Then

$$\left| \left\langle L_{\gamma}(\sigma,\phi) \right\rangle_{\beta,\kappa,\infty} - \Theta_{\beta,\kappa}'(\gamma) H_{\kappa}(\gamma) \right| \le K_0 \left(e^{-4(\beta+\kappa/6)} + |\operatorname{supp}\gamma|^{-1/2} \right)^{\frac{1}{4}}, \quad (1.2)$$

where

$$\begin{split} \Theta_{\beta,\kappa}'(\gamma) &:= e^{-2|\operatorname{supp}\gamma|e^{-24\beta-4\kappa}\left(1+(e^{8\kappa}-1)\langle L_e(\sigma,\phi)\rangle_{\infty,\kappa,\infty}\right)}, \\ H_{\kappa}(\gamma) &:= \left\langle L_{\gamma}(\sigma,\phi)\right\rangle_{\infty,\kappa,\infty}, \end{split}$$

and $K_0 = K_0(\kappa, \beta, \ell_1, \ell_2, \gamma)$ is a non-negative function with

$$K_0 \le 2 \cdot 18^3 + |\operatorname{supp} \gamma|^{1/2} e^{-4\kappa} \left(18^2 (2 + e^{-4\kappa}) \right)^{\min(\ell_1, \ell_2)} + o_{\kappa}(1).$$

An exact expression for K_0 is given in (10.42).

Remark 1.2. We later show, in Corollary 2.18, that if γ is an open path, then the function $H_{\kappa}(\gamma)$ is exactly equal to the spin-spin-correlation of the spins at the endpoints of γ in the Ising model with coupling parameter κ . By the same argument, the term $\langle L_e(\sigma, \phi) \rangle_{\infty,\kappa,\infty}$ in the function $\Theta'_{\beta,\kappa}(\gamma)$ will be equal to the spin-spin-correlation of the spins at the endpoints of the (arbitrary) edge *e*.

It is well known (see, e.g., [12]) that when κ is larger than the critical parameter for the Ising model, then $H_{\kappa}(\gamma)$ is uniformly bounded from below for all γ . At the same time, by standard arguments, we have $\langle L_e(\sigma, \phi) \rangle_{\infty,\kappa,\infty} = e^{-4 \cdot 8\kappa} + o_{\kappa}(1)$.

Remark 1.3. Using the previous remark, we now interpret our main theorem. To this, end, assume that γ is a loop along the boundary of a rectangle *R*. Assume further that the two sides of *R* are of the same order, so that K_0 is bounded from above, and that β and $|\sup \gamma|$ are both very large. Then, by Theorem 1.1, the following holds. If $|\sup \gamma|e^{-24\beta-4\kappa}$ is very large, then $\langle L_{\gamma}(\sigma, \phi) \rangle_{\beta,\kappa,\infty}$ is very close to zero, and if $|\operatorname{supp} \gamma|e^{-24\beta-4\kappa}$ is bounded from above, then $\langle L_{\gamma}(\sigma, \phi) \rangle_{\beta,\kappa,\infty}$ will be non-trivial.

Remark 1.4. The assumption that $18^2 e^{-4\kappa_0}(2 + e^{-4\kappa_0}) < 1$ guarantees that the clusters formed by the edges in unitary gauge (see Sect. 2.5) are finite almost surely, and this is one of the main properties of the model which we use in the proof of Theorem 1.1. The assumptions that $6\beta \ge \kappa$ and that the path γ is along the boundary of a rectangle is used only to simplify $\Theta'_{\beta,\kappa}(\gamma)$ and K_0 , and is not needed for any of the main ideas of the proof. In particular, the strategy used to do this also works for more general classes of loops, as long as their shape is not too rough.

Remark 1.5. If γ is a generalized loop, then $H_{\kappa}(\gamma) = 1$, and hence, in this case, we essentially recover Theorem 1.1 [18].

In Sect. 10, we state a more general version of Theorem 1.1 (Theorem 10.1). While this result is stated for cyclic groups, with minor changes, this paper's arguments should also work in a more general setting. In particular, the proof strategy should work for all finite abelian groups. Finally, we also mention that alternative versions of our main result, with different error bounds, are given by Proposition 7.1 and Proposition 10.18.

1.6. Applications. We now apply our main result to a few different Wilson lines, and ratios of Wilson lines, which has been considered in the physics literature. In all of these examples, we will work under the assumptions of Theorem 1.1. We note that, when these hold, if γ is a loop along a rectangle with side lengths of the same order, then, by Theorem 1.1, we have

$$\left\langle L_{\gamma}(\sigma,\phi)\right\rangle_{\beta,\kappa,\infty} = \Theta_{\beta,\kappa}'(\gamma)H_{\kappa}(\gamma) + o_{\beta}(1) + o_{|\operatorname{supp}\gamma|}(1).$$

Example 1.6. In [5], Bricmont and Frölich consider Wilson line observables $L_{\gamma}(\sigma, \phi)$ for axis parallel paths γ which are a shortest path between two points x_1 and x_2 (see Fig. 1).

The authors argue that the expectation $\langle L_{\gamma}(\sigma, \phi) \rangle_{\beta,\kappa}$ should exhibit a phase transition, corresponding to binding versus unbinding of dynamical quarks in the field of a static colour source.

In detail, they argue that $\langle L_{\gamma}(\sigma, \phi) \rangle_{\beta,\kappa,\infty}$ should have exponential decay with polynomial corrections if β is large and κ is small, and exponential decay if either β is large and κ is not too small.

Since, under assumption 3, $H_{\kappa}(\gamma)$ is uniformly bounded from below for all γ , and $\Theta'_{\beta,\kappa}(\gamma)$ has exponential decay in $|\operatorname{supp} \gamma|$, we see that $\langle L_{\gamma}(\sigma, \phi) \rangle_{\beta,\kappa,\infty}$ indeed has exponential decay in $|\operatorname{supp} \gamma|$ when β is large and κ is not too small.

Example 1.7 [The Marcu-Fredenhagen parameter].

Let γ and γ' be as in Fig. 2. In [15,29], they consider the ratio

$$\frac{\langle L_{\gamma'}(\sigma,\phi)\rangle_{\beta,\kappa,\infty}\langle L_{\gamma-\gamma'}(\sigma,\phi)\rangle_{\beta,\kappa,\infty}}{\langle W_{\gamma}(\sigma)\rangle_{\beta,\kappa,\infty}}.$$
(1.3)

$$x_1 \longrightarrow x_2$$

Fig. 1. The open path γ . Note that for any $\ell_1 \ge |x_2 - x_1|$ and $\ell_2 \ge 0$ there is a rectangle *R* with side lengths ℓ_1 and ℓ_2 so that γ is a path along the boundary of *R*, and hence γ satisfies the assumptions of Theorem 1.1





The limit of this ratio, when $|x_2 - x_1|$ is proportional to h and $h \to \infty$, is often referred to as the *Marcu-Fredenhagen order parameter*.

If this limit is zero, the model in argued to have charged states, and no confinement, whereas if the limit is non-zero, then there should be no charged states and confinement.

We mention that this ratio is also studied in, e.g., [6,20,25,29,30].

As an immediate consequence of our Theorem 1.1, if κ is not too small and β , κ , and γ are such that $|\operatorname{supp} \gamma| e^{-24\beta - 4\kappa}$ is bounded away from infinity, then the right hand side of (1.3) is equal to $H_{\kappa}(\gamma')^2 + o_{|\operatorname{supp} \gamma|}(1) + o_{\beta}(1)$.

However, since letting $| \operatorname{supp} \gamma |$ tend to infinity while keeping β and κ fixed violates that assumption that $| \operatorname{supp} \gamma | e^{-24\beta - 4\kappa}$ is bounded away from infinity, we cannot use this approximate equation to make conclusions about the Marcu-Fredenhagen parameter itself.

Example 1.8. Let γ and γ' be as in Fig. 3.

In [23], Gliozzi considers the ratio

$$\frac{\langle L_{\gamma'}(\sigma,\phi)\rangle_{\beta,\kappa,\infty}L_{\gamma-\gamma'}(\sigma,\phi)\rangle_{\beta,\kappa,\infty}}{\langle W_{\gamma}(\sigma)\rangle_{\beta,\kappa,\infty}},$$
(1.4)

and note that it, asymptotically, seem to only depend on the distance $|x_2 - x_1|$. Indeed, from Theorem 1.1, it follows that if $| \operatorname{supp} \gamma | e^{-24\beta - 4\kappa}$ is bounded away from infinity and κ is not too small, then the expression in (1.8) is equal to $H_{\kappa}(\gamma')^2 + o_{|\operatorname{supp} \gamma|}(1) + o_{\beta}(1)$. Using Remark 1.2 to recognise $H_{\kappa}(\gamma')$ as the spin-spin-correlation function for the Ising model, evaluated at the end-points of γ' , this confirms the observation made in [23].

Example 1.9 [Almost closed Wilson lines]. Let γ and γ' be as in Fig. 4, and let *r* be the distance between the endpoints of γ' . In [23], when *r* is much smaller that $|\operatorname{supp} \gamma|$, the path γ' is referred to as an *almost closed Wilson line*, and it was argued that the following functional equation should hold.

$$\left\langle L_{\gamma'}(\sigma,\phi)\right\rangle_{\beta,\kappa,\infty}\left\langle L_{\gamma-\gamma'}(\sigma,\phi)\right\rangle_{\beta,\kappa,\infty}\simeq\left\langle W_{\gamma}(\sigma)\right\rangle_{\beta,\kappa,\infty}f(r)$$
(1.5)

for some function f(r) that should neither depend on γ nor on the placement of the open path $\gamma - \gamma'$ on γ .

Using our main result, it indeed see that if κ is not too small, then (assuming that the side lengths of the rectangle are proportional to | supp γ | is large), we have

$$\left\langle L_{\gamma'}(\sigma,\phi)\right\rangle_{\beta,\kappa,\infty}\left\langle L_{\gamma-\gamma'}(\sigma,\phi)\right\rangle_{\beta,\kappa,\infty} = \left\langle W_{\gamma}(\sigma)\right\rangle_{\beta,\kappa,\infty}H_{\kappa}(\gamma')^{2} + o_{|\operatorname{supp}\gamma|}(1) + o_{\beta+6\kappa}(1).$$



(A) The open path γ' . (B) The generalized loop γ .

Fig. 3. The open path γ' and generalized loop γ considered in Example 1.8



Fig. 4. The open path γ' and the loop γ considered considered in Example 1.9

In particular, using Remark 1.2, this shows that the functional equation in (1.5) indeed hold when $|\operatorname{supp} \gamma|$ and β are both large, and with f(r) given by the spin-spin-correlation function evaluated at the endpoints of γ' .

1.7. Relation to other work. Many of the ideas used in this paper are refined versions of analogue ideas used in [18], which in turn build upon the works [7,9,17]. However, since this paper deals with general paths γ , and not only generalized loops as in [7,9,17,18], the first main idea in these papers, which is to pass from a generalized loop to an oriented surface, does not work. One of the main contributions of this paper thus consists in dealing with this obstacle. Even in the case when the path γ in Theorem 1.1 is a generalized loop, our proof is different from that in [18], and we hence provides an alternative proof in this case. In addition, when γ is a generalized loop, we express the leading-order term in a more transparent way than in [18].

We mention that although the recent paper [21] also calculate the first order term of Wilson loop observables in an abelian lattice gauge theory, they work with a continuous structure group, and thus their methods are fundamentally different from the ideas used here.

1.8. Structure of paper. In Sect. 2, we give a brief introduction to the cell complex of \mathbb{Z}^m and the discrete exterior calculus on this cell complex. We also define vortices and recall some of their properties from [17] to [18]. Moreover, we recall the definition of generalized loops and oriented surfaces from [18], explain unitary gauge and define a corresponding measure, and discuss the existence of the infinite volume limit $\mathbb{E}_{\beta,\kappa,\infty}[L_{\gamma}(\sigma,\phi)]$. In Sect. 3, we introduce additional notation which will be useful throughout the paper. In Sect. 4, we recall the notion of activity of gauge field configurations from [18]. In Sect. 5, we describe a useful edge graph, and introduce two couplings, one between the abelian lattice Higgs model and a \mathbb{Z}_n -model, and one between two \mathbb{Z}_n models. These will be important in the proof of our main result. In Sect. 6, using the edge graph from Sect. 5, we give upper bounds on a number of events related to the couplings introduced in Sect. 5. Next, in Sect. 7, we show how one of the couplings introduced in Sect. 5 can be used to obtain a first version of our main result, which is useful when $|\operatorname{supp} \gamma| e^{-4(\kappa+6\beta)}$ is small. This result is not needed for the proof of Theorem 1.1, but illustrates the usefulness of the coupling. In Sect. 8, we introduce a spin decomposition of two coupled configurations. In Sect.9, we describe how different 1-forms affect the Wilson line observable. Finally, in Sect. 10, we use the setup from the earlier sections to give a proof of our main result.

2. Preliminaries

2.1. The cell complex. In this section, we introduce notation for the cell complexes of the lattices \mathbb{Z}^m and $B_N := [-N, N]^m \cap \mathbb{Z}^m$ for $m, N \ge 1$. This section will closely follow the corresponding section in [17], where we refer the reader for further details.

To simplify notation, we define $e_1 := (1, 0, ..., 0), e_2 := (0, 1, 0, ..., 0), ..., e_m := (0, ..., 0, 1).$

2.1.1. Boxes and cubes. A set B of the form $([a_1, b_1] \times \cdots \times [a_m, b_m]) \cap \mathbb{Z}^m$ where, for each $j \in \{1, 2, \dots, m\}, \{a_j, b_j\} \subset \mathbb{Z}$ satisfies $a_j < b_j$, will be referred to as a box. If all the intervals $[a_j, b_j], 1 \leq j \leq m$, have the same length, then the set $([a_1, b_1] \times \cdots \times [a_m, b_m]) \cap \mathbb{Z}^m$ will be referred to as a cube.

2.1.2. Non-oriented cells. When $a \in \mathbb{Z}^m$, $k \in \{0, 1, \ldots, m\}$, and $\{j_1, \ldots, j_k\} \subseteq \{1, 2, \ldots, m\}$, we say that the set

$$(a; e_{j_1}, \dots, e_{j_k}) := \left\{ x \in \mathbb{R}^m : \exists b_1, \dots, b_k \in [0, 1] \text{ such that } x = a + \sum_{i=1}^k b_i e_{j_i} \right\}$$

is a *non-oriented k-cell*. Note that if σ is a permutation, then $(a; e_{j_1}, \ldots, e_{j_k})$ and $(a; \sigma(e_{j_1}, \ldots, e_{j_k}))$ represent the same non-oriented *k*-cell.

2.1.3. Oriented cells. To each non-oriented *k*-cell $(a; e_{j_1}, \ldots, e_{j_k})$ with $a \in \mathbb{Z}^m, k \ge 1$, and $1 \le j_1 < \cdots < j_k \le m$, we associate two oriented *k*-cells, denoted $\frac{\partial}{\partial x^{j_1}}\Big|_a \land \cdots \land \frac{\partial}{\partial x^{j_k}}\Big|_a$ and $-\frac{\partial}{\partial x^{j_1}}\Big|_a \land \cdots \land \frac{\partial}{\partial x^{j_k}}\Big|_a$, with opposite orientation. When $a \in \mathbb{Z}^m$, $1 \le j_1 < \cdots < j_k \le m$, and σ is a permutation of $\{1, 2, \ldots, k\}$, we define

$$\frac{\partial}{\partial x^{j_{\sigma(1)}}}\Big|_a \wedge \dots \wedge \frac{\partial}{\partial x^{j_{\sigma(k)}}}\Big|_a := \operatorname{sgn}(\sigma) \frac{\partial}{\partial x^{j_1}}\Big|_a \wedge \dots \wedge \frac{\partial}{\partial x^{j_k}}\Big|_a$$

If $\operatorname{sgn}(\sigma) = 1$, then $\frac{\partial}{\partial x^{j_{\sigma(1)}}}\Big|_a \wedge \cdots \wedge \frac{\partial}{\partial x^{j_{\sigma(k)}}}\Big|_a$ is said to be *positively oriented*, and if $\operatorname{sgn}(\sigma) = -1$, then $\frac{\partial}{\partial x^{j_{\sigma(1)}}}\Big|_a \wedge \cdots \wedge \frac{\partial}{\partial x^{j_{\sigma(k)}}}\Big|_a$ is said to be *negatively oriented*. Analogously, we define

$$-\frac{\partial}{\partial x^{j_{\sigma(1)}}}\bigg|_{a} \wedge \cdots \wedge \frac{\partial}{\partial x^{j_{\sigma(k)}}}\bigg|_{a} := -\operatorname{sgn}(\sigma) \left.\frac{\partial}{\partial x^{j_{1}}}\right|_{a} \wedge \cdots \wedge \frac{\partial}{\partial x^{j_{k}}}\bigg|_{a},$$

and say that $-\frac{\partial}{\partial x^{j_{\sigma(1)}}}\Big|_{a} \wedge \cdots \wedge \frac{\partial}{\partial x^{j_{\sigma(k)}}}\Big|_{a}$ is positively oriented if $-\operatorname{sgn}(\sigma) = 1$, and negatively oriented if $-\operatorname{sgn}(\sigma) = -1$.

Let $\mathcal{L} = \mathbb{Z}^m$ or $\mathcal{L} = B_N \subseteq \mathbb{Z}^m$. An oriented cell $\frac{\partial}{\partial x^{j_1}}\Big|_a \wedge \cdots \wedge \frac{\partial}{\partial x^{j_k}}\Big|_a$ is said to be in \mathcal{L} if all corners of $(a; e_{j_1}, \ldots, e_{j_k})$ belong to \mathcal{L} ; otherwise it is said to be *outside* \mathcal{L} . The set of all oriented *k*-cells in \mathcal{L} will be denoted by $C_k(\mathcal{L})$. The set of all positively and negatively oriented cells in $C_k(\mathcal{L})$ will be denoted by $C_k^+(\mathcal{L})$ and $C_k^-(\mathcal{L})$, respectively. A set $C \subseteq C_k(\mathcal{L})$ is said to be *symmetric* if for each $c \in C$ we have $-c \in C$.

A non-oriented 0-cell $a \in \mathbb{Z}^m$ is simply a point, and to each point we associate two oriented 0-cells a^+ and a^- with opposite orientation. We let $C_0(\mathcal{L})$ denote the set of all oriented 0-cells.

Oriented 1-cells will be referred to as *edges*, and oriented 2-cells will be referred to as *plaquettes*.

2.1.4. *k*-chains. The space of finite formal sums of positively oriented *k*-cells with integer coefficients will be denoted by $C_k(\mathcal{L}, \mathbb{Z})$. Elements of $C_k(\mathcal{L}, \mathbb{Z})$ will be referred to as *k*-chains. If $q \in C_k(\mathcal{L}, \mathbb{Z})$ and $c \in C_k^+(\mathcal{L})$, we let q[c] denote the coefficient of c in q. If $c \in C_k^-(\mathcal{L})$, we let q[c] := -q[-c]. For $q, q' \in C_k(\mathcal{L}, \mathbb{Z})$, we define

$$q+q' := \sum_{c \in C_k^+(\mathcal{L})} (q[c]+q'[c])c.$$

Using this operation, $C_k(\mathcal{L}, \mathbb{Z})$ becomes a group.

When $q \in C_k(\mathcal{L}, G)$, we let the *support* of q be defined by

$$\operatorname{supp} q := \left\{ c \in C_k^+(\mathcal{L}) \colon q[c] \neq 0 \right\}.$$

To simplify notation, when $q \in C_k(\mathcal{L}, G)$ and $c \in C_k(\mathcal{L})$, we write $c \in q$ if either (1) $c \in C_k^+(\mathcal{L})$ and q[c] > 0, or (2) $c \in C_k^-(\mathcal{L})$ and q[-c] < 0.

2.1.5. The boundary of a cell When $k \ge 2$, we define the boundary $\partial c \in C_{k-1}(\mathcal{L}, \mathbb{Z})$ of $c = \frac{\partial}{\partial x^{j_1}}\Big|_a \land \cdots \land \frac{\partial}{\partial x^{j_k}}\Big|_a \in C_k(\mathcal{L})$ by

$$\partial c := \sum_{k' \in \{1, \dots, k\}} \left((-1)^{k'} \frac{\partial}{\partial x^{j_1}} \bigg|_a \wedge \dots \wedge \frac{\partial}{\partial x^{j_{k'-1}}} \bigg|_a \wedge \frac{\partial}{\partial x^{j_{k'+1}}} \bigg|_a \wedge \dots \wedge \frac{\partial}{\partial x^{j_k}} \bigg|_a + (-1)^{k'+1} \frac{\partial}{\partial x^{j_1}} \bigg|_{a+e_{j_{k'}}} \wedge \dots \wedge \frac{\partial}{\partial x^{j_{k'-1}}} \bigg|_{a+e_{j_{k'}}} \wedge \frac{\partial}{\partial x^{j_{k'+1}}} \bigg|_{a+e_{j_{k'}}} \wedge \dots \wedge \frac{\partial}{\partial x^{j_k}} \bigg|_{a+e_{j_{k'}}} \right).$$

$$(2.1)$$

When $c := \frac{\partial}{\partial x^{j_1}} \Big|_a \in C_1(\mathcal{L})$ we define the boundary $\partial c \in C_0(\mathcal{L}, \mathbb{Z})$ by

$$\partial c = (-1)^1 a^+ + (-1)^{1+1} (a + e_{j_1})^+ = (a + e_{j_1})^+ - a^+.$$

We extend the definition of ∂ to *k*-chains $q \in C_k(\mathcal{L}, \mathbb{Z})$ by linearity. One verifies, as an immediate consequence of this definition, that if $k \in \{2, 3, ..., m\}$, then $\partial \partial c = 0$ for any $c \in \Omega_k(\mathcal{L})$.

2.1.6. The coboundary of an oriented cell If $k \in \{0, 1, ..., n-1\}$ and $c \in C_k(\mathcal{L})$ is an oriented k-cell, we define the *coboundary* $\hat{\partial} c \in C_{k+1}(\mathcal{L})$ of c as the (k + 1)-chain

$$\hat{\partial}c := \sum_{c' \in C_{k+1}(\mathcal{L})} (\partial c'[c])c'.$$

Note in particular that if $c' \in C_{k+1}(\mathcal{L})$, then $\hat{\partial}c[c'] = \partial c'[c]$. We extend the definition of $\hat{\partial}$ to *k*-chains $q \in C_k(\mathcal{L}, \mathbb{Z})$ by linearity.

2.1.7. The boundary of a box An oriented k-cell $c = \frac{\partial}{\partial x^{j_1}}\Big|_a \wedge \cdots \wedge \frac{\partial}{\partial x^{j_k}}\Big|_a \in C_k(B_N)$ is said to be a boundary cell of a box $B = ([a_1, b_1] \times \cdots \times [a_m, b_m]) \cap \mathbb{Z}^m \subseteq B_N$, or equivalently to be in the boundary of B, if the non-oriented cell $(a; e_{j_1}, \ldots, e_{j_k})$ is a subset of the boundary of $[a_1, b_1] \times \cdots \times [a_m, b_m]$.

When $k \in C_k(B_N)$, we let $\partial C_k(B_N)$ denote the set cells in $C_k(B_N)$ which are boundary cells of B_N .

2.2. Discrete exterior calculus. In what follows, we give a brief overview of discrete exterior calculus on the cell complexes of \mathbb{Z}^m and $B = [a_1, b_1] \times \cdots \times [a_m, b_m] \cap \mathbb{Z}^m$ for $m \ge 1$. As with the previous section, this section will closely follow the corresponding section in [17], where we refer the reader for further details and proofs.

All of the results in this subsection are obtained under the assumption that an abelian group G, which is not necessarily finite, has been given. In particular, they all hold for both $G = \mathbb{Z}_n$ and $G = \mathbb{Z}$.

2.2.1. Discrete differential forms A homomorphism from the group $C_k(\mathcal{L}, \mathbb{Z})$ to the group *G* is called a *k*-form. The set of all such *k*-forms will be denoted by $\Omega^k(\mathcal{L}, G)$. This set becomes an abelian group if we add two homomorphisms by adding their values in *G*.

The set $C_k^+(\mathcal{L})$ of positively oriented *k*-cells is naturally embedded in $C_k(\mathcal{L}, \mathbb{Z})$ via the map $c \mapsto 1 \cdot c$, and we will frequently identify $c \in C_k^+(\mathcal{L})$ with the *k*-chain $1 \cdot c$ using this embedding. Similarly, we will identify a negatively oriented *k*-cell $c \in C_k^-(\mathcal{L})$ with the *k*-chain $(-1) \cdot (-c)$. In this way, a *k*-form ω can be viewed as a *G*-valued function on $C_k(\mathcal{L})$ with the property that $\omega(c) = -\omega(-c)$ for all $c \in C_k(\mathcal{L})$. Indeed, if $\omega \in \Omega^k(\mathcal{L}, G)$ and $q = \sum a_i c_i \in C_k(\mathcal{L}, \mathbb{Z})$, we have

$$\omega(q) = \omega \left(\sum a_i c_i \right) = \sum a_i \omega(c_i),$$

and hence a k-form is uniquely determined by its values on positively oriented k-cells.

If ω is a k-form, it is useful to represent it by the formal expression

$$\sum_{1\leq j_1<\cdots< j_k\leq m}\omega_{j_1\ldots j_k}dx^{j_1}\wedge\cdots\wedge dx^{j_k}.$$

where $\omega_{j_1...j_k}$ is a *G*-valued function on the set of all $a \in \mathbb{Z}^m$ such that $\frac{\partial}{\partial x^{j_1}}\Big|_a \wedge \cdots \wedge \frac{\partial}{\partial x^{j_k}}\Big|_a \in C_k(\mathcal{L})$, defined by

$$\omega_{j_1\dots j_k}(a) = \omega \left(\frac{\partial}{\partial x^{j_1}} \bigg|_a \wedge \dots \wedge \frac{\partial}{\partial x^{j_k}} \bigg|_a \right).$$

If $1 \le j_1 < \cdots < j_k \le m$ and σ is a permutation of $\{1, 2, \ldots, k\}$, we define

$$dx^{j_{\sigma(1)}} \wedge \cdots \wedge dx^{j_{\sigma(k)}} := \operatorname{sgn}(\sigma) dx^{j_1} \wedge \cdots \wedge dx^{j_k},$$

and if $1 \le j_1, \ldots, j_k \le n$ are such that $j_i = j_{i'}$ for some $1 \le i < i' \le k$, then we let

$$dx^{j_1}\wedge\cdots\wedge dx^{j_k}:=0.$$

Given a k-form ω , we let supp ω denote the support of ω , i.e., the set of all oriented k-cells c such that $\omega(c) \neq 0$. Note that supp ω always contains an even number of elements.

2.2.2. The exterior derivative Given $h: \mathbb{Z}^m \to G, a \in \mathbb{Z}^m$, and $i \in \{1, 2, ..., m\}$, we let

$$\partial_i h(a) := h(a + e_i) - h(a).$$

If $k \in \{0, 1, 2, ..., m\}$ and $\omega \in \Omega^k(\mathcal{L}, G)$, we define the (k+1)-form $d\omega \in \Omega^{k+1}(\mathcal{L}, G)$ by

$$d\omega = \sum_{1 \le j_1 < \cdots < j_k \le m} \sum_{i=1}^m \partial_i \omega_{j_1, \dots, j_k} \, dx^i \wedge (dx^{j_1} \wedge \cdots \wedge dx^{j_k}).$$

The operator *d* is called the *exterior derivative*. Using (2.1), one can show that $\omega \in \Omega^k(\mathcal{L}, G)$ and $c \in C_k(\mathcal{L}, \mathbb{Z})$, we have $d\omega(c) = \omega(\partial c)$. This equality is known as the *discrete Stokes' theorem*. Recalling that when $k \in \{2, 3, ..., m-2\}$ and $c \in C_{k+2}(\mathcal{L})$, then $\partial \partial c = 0$, it follows from the discrete Stokes theorem that for any $\omega \in \Omega^k(\mathcal{L}, G)$, we have $dd\omega = 0$.

2.2.3. Closed forms and the Poincaré lemma For $k \in \{0, ..., m\}$, we say that a k-form $\omega \in \Omega^k(\mathcal{L}, G)$ is closed if $d\omega(c) = 0$ for all $c \in C_{k+1}(\mathcal{L})$. The set of all closed forms in $\Omega^k(\mathcal{L}, G)$ will be denoted by $\Omega_0^k(\mathcal{L}, G)$.

Lemma 2.1 [The Poincaré lemma, Lemma 2.2 in [9]]. Let $k \in \{1, ..., m\}$ and let *B* be a box in \mathbb{Z}^m . Then the exterior derivative *d* is a surjective map from the set $\Omega^{k-1}(B \cap \mathbb{Z}^m, G)$ to $\Omega_0^k(B \cap \mathbb{Z}^m, G)$. Moreover, if *G* is finite, then this map is an $|\Omega_0^{k-1}(B \cap \mathbb{Z}^m, G)|$ -to-1 correspondence.

Lastly, if $k \in \{1, 2, ..., m - 1\}$ and $\omega \in \Omega_0^k(B \cap \mathbb{Z}^m, G)$ vanishes on the boundary of B, then there is a (k - 1)-form $\omega' \in \Omega^{k-1}(B \cap \mathbb{Z}^m, G)$ that also vanishes on the boundary of B and satisfies $d\omega' = \omega$.

2.2.4. Non-trivial forms. We say that a k-form $\omega \in \Omega^k(\mathcal{L}, G)$ is non-trivial if there is at least one k-cell $c \in C_k(\mathcal{L})$ such that $\omega(c) \neq 0$.

2.2.5. Restrictions of forms. If $\omega \in \Omega^k(\mathcal{L}, G)$, $C \subseteq C_k(\mathcal{L})$ is symmetric, and $c \in C$, we define

$$\omega|_C(c) := \begin{cases} \omega(c) & \text{if } c \in C, \\ 0 & \text{else.} \end{cases}$$

2.2.6. A partial ordering of $\Omega^k(\mathcal{L}, G)$. We now recall the partial ordering on differential forms, which was introduced in [18].

Definition 2.2 [Definition 2.6 in [18]]. When $k \in \{0, 1, ..., m\}$ and $\omega, \omega' \in \Omega^k(\mathcal{L}, G)$, we write $\omega' \leq \omega$ if

(i) $\omega' = \omega|_{\text{supp }\omega'}$, and (ii) $d\omega' = (d\omega)|_{\text{supp }d\omega'}$.

If $\omega' \neq \omega$ and $\omega' \leq \omega$, we write $\omega' < \omega$.

The following lemma from [18] collects some basic facts about the relation \leq on $\Omega^k(\mathcal{L}, G)$, and shows that \leq is a partial order on $\Omega^k(\mathcal{L}, G)$.

Lemma 2.3 [Lemma 2.7 in [18]]. Let $k \in \{0, 1, ..., m\}$ and $\omega, \omega', \omega'' \in \Omega^k(\mathcal{L}, G)$. The relation \leq on $\Omega^k(\mathcal{L}, G)$ has the following properties.

- (*i*) *Reflexivity*: $\omega \leq \omega$.
- (*i*) Antisymmetry: If $\omega' \leq \omega$ and $\omega \leq \omega'$, then $\omega = \omega'$.
- (iii) Transitivity: If $\omega'' \leq \omega'$ and $\omega' \leq \omega$, then $\omega'' \leq \omega$.
- (iv) If $\omega' \leq \omega$, then $\omega \omega' = \omega|_{C_1(B_N) \smallsetminus (\operatorname{supp} \omega')} \leq \omega$.
- (v) If $\omega' \leq \omega$, then supp $d\omega'$ and supp $d(\omega \omega')$ are disjoint.

The next lemma guarantees the existence of minimal elements satisfying certain constraints.

Lemma 2.4 [Lemma 2.8 in [18]]. Let $k \in \{0, 1, ..., m\}$, let $\Omega \subseteq \Omega^k(\mathcal{L}, G)$, and let $\omega \in \Omega$. Then there is $\omega' \leq \omega$ such that

(i) $\omega' \in \Omega$, and (ii) There is no $\omega'' < \omega'$ such that $\omega'' \in \Omega$.

2.2.7. *Irreducible forms* The partial ordering given in Definition 2.2 allows us to introduce a notion of irreducibility.

Definition 2.5 [Definition 2.9 in [18]]. When $k \in \{0, 1, ..., m - 1\}$, a k-form $\omega \in \Omega^k(\mathcal{L}, G)$ is said to be *irreducible* if there is no non-trivial k-form $\omega' \in \Omega^k(\mathcal{L}, G)$ such that $\omega' < \omega$.

Equivalently, $\omega \in \Omega^k(\mathcal{L}, G)$ is irreducible if there is no non-empty set $S \subsetneq \operatorname{supp} \omega$ such that $\operatorname{supp} d(\omega|_S)$ and $\operatorname{supp} d(\omega|_{S^c})$ are disjoint. Note that if $\omega \in \Omega^k(\mathcal{L}, G)$ satisfies $d\omega = 0$, then ω is irreducible if and only if there is no non-empty set $S \subsetneq \operatorname{supp} \omega$ such that $d(\omega|_S) = d(\omega|_{S^c}) = 0$.

Lemma 2.6 [Lemma 2.10 in [18]]. Let $k \in \{0, 1, ..., m-1\}$, and let $\omega \in \Omega^k(\mathcal{L}, G)$ be non-trivial and have finite support.

Then there is an integer $j \ge 1$ and k-forms $\omega_1, \ldots, \omega_j \in \Omega^k(\mathcal{L}, G)$ such that

- (i) For each $i \in \{1, 2, ..., j\}$, ω_i is non-trivial and irreducible,
- (*ii*) For each $i \in \{1, 2, \ldots, j\}$, $\omega_i \leq \omega$,
- (iii) $\omega_1, \ldots, \omega_j$ have disjoint supports,
- (*iv*) $\omega = \omega_1 + \cdots + \omega_j$, and
- (v) $d\omega_1, \ldots, d\omega_i$ have disjoint supports.

A set $\Omega := \{\omega_1, \ldots, \omega_j\} \subseteq \Omega^k(\mathcal{L}, G)$ such that $\omega_1, \ldots, \omega_j$ satisfies (2.6) –(2.6) of Lemma 2.6 will be referred to as a *decomposition* of $\omega \in \Omega^k(\mathcal{L}, G)$.

We note that as an immediate consequence of the previous lemma, if $\omega \in \Omega_0^2(\mathcal{L}, G)$ has finite support, then there is a set $\Omega \subseteq \Omega_0^2(\mathcal{L}, G)$ which is a decomposition of ω (see also Lemma 2.12 in [18]).

2.2.8. *Minimal forms*. In this section, we recall three lemmas from [18] which gives lower bounds on the size of the support of differential forms. Throughout this section, we assume that m = 4. In other words, we assume that we are working on the \mathbb{Z}^4 -lattice.

Lemma 2.7 [Lemma 2.16 in [18]]. Let $\sigma \in \Omega^1(\mathcal{L}, G)$. Then

$$|\operatorname{supp} \sigma| \geq |\operatorname{supp} d\sigma|/6.$$

Lemma 2.8. Let $\omega \in \Omega_0^2(\mathcal{L}, G)$ be non-trivial and have finite support, and assume that there is a plaquette $p \in \text{supp } \omega$ such that supp ∂p contains no boundary edges of B_N .

Then $|(\sup \omega)^+| \ge 6$, and if $|(\sup \omega)^+| = 6$, then there is an edge $e_0 \in C_1(B_N)$ such that $\operatorname{supp} \nu = \operatorname{supp} \hat{\partial} e_0 \cup \operatorname{supp} \hat{\partial} (-e_0)$.

For a proof of Lemma 2.8, see, e.g., Lemma 3.4.6 in [7].

Lemma 2.9 [Lemma 2.19 in [18]]. Let $\sigma \in \Omega_0^1(B_N, G)$ be non-trivial, and assume that there is an edge $e \in \text{supp } \sigma$ such that the support of $\hat{\partial} e$ contains no boundary cells of B_N . Then $|(\text{supp } \sigma)^+| \ge 8$.

2.3. Vortices. In this section, we use the notion of irreducibility introduced in Sect. 2.2.7 to define what we refer to as vortices. We mention that the definition of a vortex given in Definition 2.10 below is identical to the definitions used in [17, 18], but is different from the corresponding definitions in [7,9].

Definition 2.10 [Vortex]. Let $\sigma \in \Omega^1(B_N, G)$. A non-trivial and irreducible 2-form $\nu \in \Omega^2_0(B_N, G)$ is said to be a *vortex* in σ if $\nu \leq d\sigma$, i.e., if $d\sigma(p) = \nu(p)$ for all $p \in \text{supp } \nu$.

We say that $\sigma \in \Omega^1(B_N, G)$ has a vortex at $V \subseteq C_2(B_N)$ if $(d\sigma)|_V$ is a vortex in σ .

Lemma 2.11 [Lemma 3.6 in [18]]. Let $\sigma', \sigma \in \Omega^1(B_N, G)$ be such that $\sigma' \leq \sigma$, and let $v \in \Omega^2_0(B_N, G)$ be a vortex in σ' . Then v is a vortex in σ .

With Lemma 2.8 in mind, we say that a vortex v such that no plaquette in supp v is a boundary plaquette of \mathcal{L} is a *minimal vortex* if $|\operatorname{supp} v| = 12$.

Lemma 2.12 [Lemma 3.2 in [18]]. Let $\sigma \in \Omega^1(B_N, G)$, and let $v \in \Omega^2_0(B_N, G)$ be a minimal vortex in σ . Then there is an edge $\partial x_j \in C_1(B_N)$ and a group element $g \in G \setminus \{0\}$ such that

$$\nu = d(g \, dx_i). \tag{2.2}$$

In particular, $d\sigma(p) = v(p) = g$ whenever $p \in \hat{\partial} e_0$.

If $\sigma \in \Omega^1(B_N, G)$ and $\nu \in \Omega_0^2(B_N, G)$ is a minimal vortex in σ which can be written as in (2.2) for some $e_0 \in C_1(B_N)$ and $g \in G \setminus \{0\}$, then we say that ν is a *minimal vortex centered at* e_0 .

2.4. Generalized loops and oriented surfaces. In this section, we recall the definitions of generalized loops and oriented surfaces from [17], and outline their connection.

Definition 2.13 [Definition 2.6 in [17]]. A 1-chain $\gamma \in C_1(\mathcal{L}, \mathbb{Z})$ with finite support is a *generalized loop* if

(1) for all $e \in \Omega^1(\mathcal{L})$, we have $\gamma[e] \in \{-1, 0, 1\}$, and (2) $\partial \gamma = 0$.

Definition 2.14 [Definition 2.7 in [17]]. Let $\gamma \in C_1(\mathcal{L}, \mathbb{Z})$ be a generalized loop. A 2-chain $q \in C_2(\mathcal{L}, \mathbb{Z})$ is an *oriented surface* with *boundary* γ if $\partial q = \gamma$.

We recall that by Stokes' theorem (see Sect. 2.2.2), for any $q \in C_2(\mathcal{L}, G)$ and any $\sigma \in \Omega^1(\mathcal{L}, G)$, we have

$$\sigma(\partial q) = d\sigma(q).$$

The following lemma gives a connection between generalized loops and oriented surfaces.

Lemma 2.15 [Lemma 2.8 in [17]]. Let $\gamma \in C_1(\mathcal{L}, \mathbb{Z})$ be a generalized loop, and let $B \subseteq \mathcal{L}$ be a box containing the support of γ . Then there is an oriented surface $q \in C_2(\mathcal{L}, \mathbb{Z})$ with support contained in B such that γ is the boundary of q.

2.5. Unitary gauge. In this section, we introduce gauge transforms, and the describe how these can be used to rewrite the Wilson line expectation as an expectation with respect to a slightly simpler probability measure.

Before we can state the main results of this section, we need to briefly discuss gauge transformations. To this end, for $\eta \in \Omega_0(B_N, G)$, consider the bijection $\tau := \tau_\eta := \tau_\eta^{(1)} \times \tau_\eta^{(2)}$: $\Omega^1(B_N, G) \times \Omega^0(B_N, G) \to \Omega_1(B_N, G) \times \Omega^0(B_N, G)$, defined by

$$\begin{aligned} \sigma(e) &\mapsto -\eta(x) + \sigma(e) + \eta(y), \quad e = (x, y) \in C_1(B_N), \\ \phi(x) &\mapsto \phi(x) + \eta(x), \qquad x \in C_0(B_N). \end{aligned}$$
 (2.3)

Any mapping τ of this form is called a *gauge transformation*. Any mapping τ of this form is called a *gauge transformation*, and functions $f : \Omega^1(B_N, G) \times \Omega^0(B_N, G) \to \mathbb{C}$ which are invariant under such mappings in the sense that $f = f \circ \tau$ are said to be *gauge invariant*.

For $\beta, \kappa \geq 0$ and $\sigma \in \Omega^1(B_N, G)$, define

$$\mu_{N,\beta,\kappa}(\sigma) := Z_{N,\beta,\kappa}^{-1} \exp\Big(\beta \sum_{p \in C_2(B_N)} \rho\big(d\sigma(p)\big) + \kappa \sum_{e \in C_1(B_N)} \rho\big(\sigma(e)\big)\Big), \quad (2.4)$$

where $Z_{N,\beta,\kappa}^{-1}$ is a normalizing constant which ensures that $\mu_{N,\beta,\kappa}$ is a probability measure. We let $\mathbb{E}_{N,\beta,\kappa}$ denote the corresponding expectation.

The main reason that gauge transformations are useful to us is the following result.

Proposition 2.16 [Proposition 2.21 in [18]]. Let $\beta, \kappa \geq 0$, and let and assume that the function $f: \Omega^1(B_N, G) \times \Omega^0(B_N, G) \to \mathbb{C}$ is gauge invariant. Then

$$\mathbb{E}_{N,\beta,\kappa,\infty}[f(\sigma,\phi)] = \mathbb{E}_{N,\beta,\kappa}[f(\sigma,1)].$$

The main idea of the proof of Proposition 2.16 is to perform a change of variables, where we for each pair (σ, ϕ) apply the gauge transformation $\tau_{-\phi}$, thus mapping ϕ to 0. After having applied this gauge transformation, we are said to be working in *unitary* gauge.

Noting that for any path γ , the function $(\sigma, \phi) \mapsto L_{\gamma}(\sigma, \phi)$ is gauge invariant, we obtain the following result as an immediate corollary of Proposition 2.16.

Corollary 2.17. Let $\beta \in [0, \infty]$, $\kappa \ge 0$, and let γ be a path in $C_1(B_N)$. Then

$$\mathbb{E}_{N,\beta,\kappa,\infty} \big[L_{\gamma}(\sigma,\phi) \big] = \mathbb{E}_{N,\beta,\kappa} \big[L_{\gamma}(\sigma,1) \big] = \mathbb{E}_{N,\beta,\kappa} \big[\rho \big(\sigma(\gamma) \big) \big]$$

Results analogous to Proposition 2.16 are considered well-known in the physics literature.

By combining the previous result with Lemma 2.1, we obtain the following result, which will help us interpret our main result.

Corollary 2.18. Let $\kappa \ge 0$, and let γ be an open path from $x_1 \in C_0(B_N)$ to $x_2 \in C_0(B_N)$. Then

$$H_{\kappa}(\gamma) = \lim_{N \to \infty} Z_{N,\kappa}^{-1} \sum_{\eta \in \Omega^{0}(B_{N},G)} \rho(\eta(-x_{1})) \rho(\eta(x_{2})) e^{-\kappa \sum_{e \in C_{1}(B_{N})} \rho(\eta(\partial e))}$$

where

$$Z_{N,\kappa} := \sum_{\eta \in \Omega^0(B_N,G)} e^{\kappa \sum_{e \in C_1(B_N)} \rho(\eta(\partial e))}.$$

If particular, if $G = \mathbb{Z}_2$, then $H_{\kappa}(\gamma)$ is the spin-spin-correlation between for the spins at the endpoints of γ for the Ising model on B_N with coupling constant κ .

Proof. By Corollary 2.17, we have

$$H_{\kappa}(\gamma) = \left\langle L_{\gamma}(\sigma, \phi) \right\rangle_{\infty, \kappa, \infty} = \lim_{N \to \infty} \left\langle L_{\gamma}(\sigma, \phi) \right\rangle_{N, \infty, \kappa, \infty}$$

=
$$\lim_{N \to \infty} \mathbb{E}_{N, \infty, \kappa, \infty} \left[L_{\gamma}(\sigma, \phi) \right] = \lim_{N \to \infty} \mathbb{E}_{N, \infty, \kappa} \left[L_{\gamma}(\sigma, 1) \right]$$

=
$$\lim_{N \to \infty} Z_{N, \infty, \kappa, \infty}^{-1} \sum_{\sigma \in \Omega_{0}^{1}(B_{N}, G)} \rho(\sigma(\gamma)) e^{\kappa \sum_{e \in C_{1}(B_{N})} \rho(\sigma(e))}.$$

Since $\beta = \infty$, we only need to sum over $\sigma \in \Omega_0^1(B_N, G)$. Now recall that by Lemma 2.1, for each $\sigma \in \Omega_0^1(B_N, G)$ there is $\eta \in \Omega^0(B_N, G)$ such that $d\eta = \sigma$. Moreover, the mapping $\eta \mapsto d\eta$ is a $|\Omega_0^0(B_N, G)|$ -to-1 correspondence. From this the desired conclusion immediately follows.

With the current section in mind, we will work with $\sigma \sim \mu_{N,\beta,\kappa}$ rather than $(\sigma, \phi) \sim \mu_{N,\beta,\kappa,\infty}$ throughout the rest of this paper, together with the observable

$$L_{\gamma}(\sigma) := L_{\gamma}(\sigma, 1) = \prod_{e \in \gamma} \rho(\sigma(e)) = \rho(\sigma(\gamma)).$$

2.6. Existence of the infinite volume limit. In this section, we recall a result which shows existence and translation invariance of the infinite volume limit $\langle L_{\gamma}(\sigma, \phi) \rangle_{\beta,\kappa}$ defined in the introduction. This result is well-known, and is often mentioned in the literature as a direct consequence of the Ginibre inequalities. A full proof of this result in the special case $\kappa = 0$ was included in [17], and the general case can be proven completely analogously, hence we omit the proof here.

Proposition 2.19. Let $G = \mathbb{Z}_n$, $M \ge 1$, and let $f : \Omega^1(B_M, G) \to \mathbb{R}$.

For $M' \ge M$, we abuse notation and let f denote the natural extension of f to $C_1(B_{M'})$, i.e., the unique function such that $f(\sigma) = f(\sigma|_{C_1(B_M)})$ for all $\sigma \in \Omega^1(B_{M'}, G)$.

Further, let $\beta \in [0, \infty]$ *and* $\kappa \ge 0$ *. Then the following hold.*

- (*i*) The limit $\lim_{N\to\infty} \mathbb{E}_{N,\beta,\kappa}[f(\sigma)]$ exists.
- (ii) For any translation τ of \mathbb{Z}^m , we have $\lim_{N\to\infty} \mathbb{E}_{N,\beta,\kappa} [f \circ \tau(\sigma)] = \lim_{N\to\infty} \mathbb{E}_{N,\beta,\kappa} [f(\sigma)].$

3. Additional Notation and Standing Assumptions

Throughout the rest of this paper, we will assume that $N \ge 1$ is given, and that $G = \mathbb{Z}_n$ for some $n \ge 2$.

To simplify the notation, we now introduce some additional notation.

For $r \ge 0$ and $g \in G$, we define

$$\varphi_r(g) := e^{r\Re(\rho(g) - \rho(0))}.$$
(3.1)

We extend this notation to $r = \infty$ by letting

$$\varphi_{\infty}(g) := \begin{cases} 1 & \text{if } g = 0, \\ 0 & \text{if } g \in G \smallsetminus \{0\} \end{cases}$$

Next, for $\hat{g} \in G$ and $\beta, \kappa \ge 0$, we define

$$\theta_{\beta,\kappa}(\hat{g}) := \frac{\sum_{g \in G} \rho(g) \varphi_{\beta}(g)^{12} \varphi_{\kappa}(g + \hat{g})^2}{\sum_{g \in G} \varphi_{\beta}(g)^{12} \varphi_{\kappa}(g + \hat{g})^2}.$$
(3.2)

When γ is a path, or when $E \subseteq C_1(B_N)$ is a finite set, we define

$$\Theta_{N,\beta,\kappa}(\gamma) := \mathbb{E}_{N,\infty,\kappa} \Big[\prod_{e \in \gamma} \theta_{\beta,\kappa} \big(\sigma(e) \big) \Big] \quad \text{and} \quad \Theta_{N,\beta,\kappa}(E) := \mathbb{E}_{N,\infty,\kappa} \Big[\prod_{e \in E} \theta_{\beta,\kappa} \big(\sigma(e) \big) \Big].$$
(3.3)

We next define a number of functions which will be used as error bounds. To this end, for $r \ge 0$, let

$$\alpha_0(r) := \sum_{g \in G \smallsetminus \{0\}} \varphi_r(g)^2 \text{ and } \alpha_1(r) := \max_{g \in G \smallsetminus \{0\}} \varphi_r(g)^2.$$
(3.4)

Next, for $\beta, \kappa \ge 0$, define

$$\alpha_{2}(\beta,\kappa) := \alpha_{0}(\beta)\alpha_{0}(\kappa)^{1/6}, \quad \alpha_{3}(\beta,\kappa) := \left|1 - \theta_{\beta,\kappa}(0)\right|, \quad \alpha_{4}(\beta,\kappa)$$
$$:= \max_{g \in G} \left|\theta_{\beta,\kappa}(g) - \theta_{\beta,\kappa}(0)\right|, \qquad (3.5)$$
$$\alpha_{5}(\beta,\kappa) := \min_{g_{1},g_{2},\dots,g_{6} \in G} \left(1 - \left|\frac{\sum_{g \in G} \rho(g) \left(\prod_{k=1}^{6} \varphi_{\beta}(g+g_{k})^{2}\right) \varphi_{\kappa}(g)^{2}}{\sum_{g \in G} \left(\prod_{k=1}^{6} \varphi_{\beta}(g+g_{k})^{2}\right) \varphi_{\kappa}(g)^{2}}\right|\right)$$

and

$$\alpha_6(\beta,\kappa) := \max_{g \in G} \left| 1 - \theta_{\beta,\kappa}(g) \right|. \tag{3.7}$$

When γ is a path, an edge $e \in \operatorname{supp} \gamma$ is said to be a corner edge in γ if there is another edge $e' \in \gamma$ and a plaquette $p \in \hat{\partial} e$ such that $p \in \pm \hat{\partial} e'$. We define the 1-form $\gamma_c \in C_1(\mathcal{L}, \mathbb{Z})$ for $c' \in C_1(B_N)$ by

$$\gamma_c[c'] := \begin{cases} \gamma[c'] & \text{if } c' \text{ is a corner edge of } \gamma, \\ 0 & \text{else.} \end{cases}$$
(3.8)

In the rest of this paper, we will often work under the following assumption. [A] $18^2 \alpha_0(\kappa)(2 + \alpha_0(\kappa)) < 1$.

In essence, the purpose of this assumption is to guarantee that we are in the sub-critical regime of the model, where certain edge clusters are finite almost surely.

(3.6)

4. Activity of Gauge Field Configurations

In this section, we recall the useful notion of the activity of a gauge field configurations from [18]. To this end, recall the definition of φ_r from the previous section. Since ρ is a unitary representation of *G*, for any $g \in G$ we have $\rho(g) = \overline{\rho(-g)}$, and hence $\Re \rho(g) = \Re \rho(-g)$. In particular, this implies that for any $g \in G$ and any $r \ge 0$, we have

$$\varphi_r(g) = e^{r(\Re\rho(g) - \rho(0))} = e^{r\beta(\Re\rho(-g) - \rho(0))} = \varphi_r(-g).$$
(4.1)

Clearly, we also have $\varphi_{\infty}(g) = \varphi_{\infty}(-g)$ for all $g \in G$. Moreover, if $a \ge 0$ and $r \ge 0$, then

$$\varphi_r(g)^a = \varphi_{ar}(g).$$

Abusing notation, for $\sigma \in \Omega^1(B_N, G)$ and $r \in [0, \infty]$, we define

$$\varphi_r(\sigma) := \prod_{e \in C_1(B_N)} \varphi_r(\sigma(e)),$$

and for $\omega \in \Omega_0^2(B_N, G)$, we define

$$\varphi_r(\omega) := \prod_{p \in C_2(B_N)} \varphi_r(\omega(p)).$$

For $\beta \in [0, \infty]$ and $\kappa \ge 0$, we define the *activity* of $\sigma \in \Omega^1(B_N, G)$ by

$$\varphi_{\beta,\kappa}(\sigma) := \varphi_{\kappa}(\sigma)\varphi_{\beta}(d\sigma).$$

Note that with this notation, for $\sigma \in \Omega^1(B_N, G)$, $\beta \in [0, \infty]$, and $\kappa \ge 0$, we have

$$\mu_{N,\beta,\kappa}(\sigma) = \frac{\varphi_{\beta,\kappa}(\sigma)}{\sum_{\sigma' \in \Omega^1(B_N,G)} \varphi_{\beta,\kappa}(\sigma')}.$$
(4.2)

Before ending this section, we recall two results from [18] about the activity of gauge field configurations, which will be useful to us.

Lemma 4.1 [Lemma 4.1 in [18]]. Let $\sigma, \sigma' \in \Omega^1(B_N, G)$ be such that $\sigma' \leq \sigma$, let $\beta \in [0, \infty]$, and let $\kappa \geq 0$. Then

$$\varphi_{\beta,\kappa}(\sigma) = \varphi_{\beta,\kappa}(\sigma')\varphi_{\beta,\kappa}(\sigma - \sigma'). \tag{4.3}$$

Proposition 4.2 [Proposition 5.1 in [18]]. Let $\sigma' \in \Omega^1(B_N, G)$, let $\beta \in [0, \infty]$, and let $\kappa \ge 0$. Then

$$\mu_{N,\beta,\kappa}(\left\{\sigma\in\Omega^1(B_N,G)\colon\sigma'\leq\sigma\right\})\leq\varphi_{\beta,\kappa}(\sigma').$$

5. Two Couplings

The main purpose of this section is to introduce two couplings which will be useful to us throughout this paper. Both of these couplings use ideas from disagreement percolation, and will be constructed so that the two coupled configurations agree as often as possible, given certain constraints. Before we introduce the two couplings, we will recall the definition of a certain edge graph from [16], and state and prove some of its properties, and introduce a set $E_{E_0,\hat{\sigma},\hat{\sigma}'}$ which will be used for the definitions of the two couplings.

5.1. A useful edge graph.

Definition 5.1. Given $\sigma, \sigma' \in \Omega^1(B_N, G)$, let $\mathcal{G}(\sigma, \sigma')$ be the graph with vertex set $C_1(B_N)$, and with an edge between two distinct vertices $e, e' \in C_1(B_N)$ if either

(i) e' = -e, or

(ii) $e, e' \in \operatorname{supp} \sigma \cup \operatorname{supp} \sigma'$, and either supp $\hat{\partial} e \cap \operatorname{supp} \hat{\partial} e' \neq \emptyset$ or supp $\hat{\partial} e \cap \operatorname{supp} \hat{\partial} (-e') \neq \emptyset$.

Given $\sigma \in \Omega^1(B_N)$, we let $\mathcal{G}(\sigma) := \mathcal{G}(\sigma, 0)$.

Given $\sigma, \sigma' \in \Omega^1(B_N)$, $\mathcal{G} := \mathcal{G}(\sigma, \sigma')$, and $e \in C_1(B_N)$, we let $\mathcal{C}_{\mathcal{G}}(e)$ be set of all edges $e' \in C_1(B_N)$ which belong to the same connected component as e in \mathcal{G} . For $E \subseteq C_1(B_N)$, we let $\mathcal{C}_{\mathcal{G}}(E) := \bigcup_{e \in E} \mathcal{C}_{\mathcal{G}}(e)$.

We now state and prove a number of lemmas, which describe different properties of the sets $C_{\mathcal{G}(\hat{\sigma}, \hat{\sigma}')}(E)$.

Lemma 5.2 [Lemma 7.2 in [18]]. Let $\sigma, \sigma' \in \Omega^1(B_N, G)$, $E \subseteq C_1(B_N)$, and $E' := C_{\mathcal{G}(\sigma,\sigma')}(E)$. Then

(i) $\sigma|_{E'} \leq \sigma$, (ii) $\sigma|_{C_1(B_N) \setminus E'} \leq \sigma$, (iii) $\sigma'|_{E'} \leq \sigma'$, and (iv) $\sigma'|_{C_1(B_N) \setminus E'} \leq \sigma'$.

Lemma 5.3. Let $\sigma \in \Omega^1(B_N, G)$ be nontrivial and irreducible. Then the support of σ is a connected set in $\mathcal{G}(\sigma)$.

Proof. Let $e \in \text{supp } \sigma$, and define $\sigma' := \sigma |_{\mathcal{C}_{\mathcal{G}(\sigma)}(e)}$. Then, by definition, σ' is non-trivial, and by Lemma 5.2, we have $\sigma' \leq \sigma$. Since σ is irreducible, it follows that $\sigma = \sigma'$, and hence the desired conclusion follows.

Lemma 5.4. Let $\sigma, \sigma' \in \Omega^1(B_N, G)$. Assume that $\sigma'' \leq \sigma$ is nontrivial and irreducible, and let $e \in \text{supp } \sigma''$. Then

$$\sigma'' \le \sigma |_{\mathcal{C}_{\mathcal{G}(\sigma,\sigma')}(e)}.$$
(5.1)

Proof. Since σ'' is irreducible, by Lemma 5.3, the support of σ'' is a connected set $\mathcal{G}(\sigma'', 0)$.

Since $\sigma'' \leq \sigma$, we have $\sigma|_{\text{supp }\sigma''} = \sigma''$, and hence it follows that the support of σ'' is a connected set in $\mathcal{G}(\sigma, \sigma')$.

Consequently, since $e \in \operatorname{supp} \sigma''$, we have

$$\operatorname{supp} \sigma'' \subseteq \mathcal{C}_{\mathcal{G}(\sigma,\sigma')}(e), \tag{5.2}$$

and thus, since $\sigma|_{\text{supp }\sigma''} = \sigma''$, it follows that

$$(\sigma|_{\mathcal{C}_{\mathcal{G}}(\sigma,\sigma')}(e)})|_{\operatorname{supp}\sigma''} = \sigma|_{\operatorname{supp}\sigma''} = \sigma''.$$

For (5.1) to follow, it thus remains to show that

$$\left(d(\sigma|_{\mathcal{C}_{\mathcal{G}(\sigma,\sigma')}(e)})\right)\Big|_{\operatorname{supp} d\sigma''} = d\sigma''.$$
(5.3)

If $d\sigma'' = 0$, then this immediately follows. Hence, assume that $d\sigma'' \neq 0$, and let $p \in \text{supp } d\sigma''$. Since $p \in \text{supp } d\sigma''$, there must exist at least one $e' \in \partial p$ with $\sigma''(e') \neq 0$.

Since $e' \in \text{supp } \sigma''$, it follows from (5.2) that $e' \in C_{\mathcal{G}(\sigma,\sigma')}(e)$. Since $e' \in \partial p$, it follows from the definition of $C_{\mathcal{G}(\sigma,\sigma')}(e)$ that any edge $e'' \in \partial p$ with $\sigma(e'') \neq 0$ is also a member of $C_{\mathcal{G}(\sigma,\sigma')}(e)$. Consequently, we must have $\sigma(e'') = \sigma|_{\mathcal{C}_{\mathcal{G}(\sigma,\sigma')}(e)}(e'')$ for all $e'' \in \partial p$, and hence

$$d\sigma|_{\mathcal{C}_{\mathcal{C}}(\sigma,\sigma')}(e)}(p) = d\sigma(p).$$

Since $\sigma'' \leq \sigma$ and $p \in \operatorname{supp} d\sigma''$, we also have $d\sigma''(p) = d\sigma(p)$, and hence we conclude that

$$d\sigma|_{\mathcal{C}_{\mathcal{G}(\sigma,\sigma')}(e)}(p) = d\sigma''(p).$$

Since this holds for any $p \in \text{supp } d\sigma''$, we obtain (5.3).

This concludes the proof.

Lemma 5.5. Let $\sigma, \sigma' \in \Omega^1(B_N, G)$, and let $E \subseteq C_1(B_N)$. Assume that $\sigma'' \leq \sigma$ is nontrivial and irreducible, and that $\operatorname{supp} \sigma'' \cap C_{\mathcal{G}(\sigma,\sigma')}(E) \neq \emptyset$. Then $\sigma'' \leq \sigma|_{\mathcal{C}_{\mathcal{G}(\sigma,\sigma')}(E)}$.

Proof. Fix some $e \in \operatorname{supp} \sigma'' \cap C_{\mathcal{G}(\sigma,\sigma')}(E)$.

Since σ'' is irreducible and $e \in \operatorname{supp} \sigma''$, it follows from Lemma 5.4 that $\sigma'' \leq \sigma|_{\mathcal{C}_{G(\sigma,\sigma')}(e)}$.

Next, since $e \in C_{\mathcal{G}(\sigma,\sigma')}(E)$, we have $C_{\mathcal{G}(\sigma,\sigma')}(e) \subseteq C_{\mathcal{G}(\sigma,\sigma')}(E)$, and hence, by Lemma 5.2, it follows that

$$\sigma|_{\mathcal{C}_{\mathcal{G}(\sigma,\sigma')}(e)} = (\sigma|_{\mathcal{C}_{\mathcal{G}(\sigma,\sigma')}(E)})|_{\mathcal{C}_{\mathcal{G}(\sigma,\sigma')}(e)} \le \sigma|_{\mathcal{C}_{\mathcal{G}(\sigma,\sigma')}(E)}.$$

Since $\sigma'' \leq \sigma|_{\mathcal{C}_{\mathcal{G}(\sigma,\sigma')}(E)}$, using Lemma 2.3 (2.3), we thus conclude that $\sigma'' \leq \sigma|_{\mathcal{C}_{\mathcal{G}(\sigma,\sigma')}(E)}$.

Lemma 5.6. Let $\sigma, \sigma' \in \Omega^1(B_N, G)$, let $E \subseteq C_1(B_N)$, and assume that $\sigma'' \leq \sigma$ is nontrivial and irreducible. Then either $\sigma'' \leq \sigma|_{\mathcal{C}_{\mathcal{G}(\sigma,\sigma')}(E)}$, or $\sigma'' \leq \sigma|_{C_1(B_N) \setminus \mathcal{C}_{\mathcal{G}(\sigma,\sigma')}(E)}$.

Proof. If supp $\sigma'' \cap C_{\mathcal{G}(\sigma,\sigma')}(E) \neq \emptyset$, then, by Lemma 5.5, we have $\sigma'' \leq \sigma|_{\mathcal{C}_{\mathcal{G}(\sigma,\sigma')}(E)}$, and hence the desired conclusion holds in this case.

Now instead assume that $\operatorname{supp} \sigma'' \cap C_{\mathcal{G}(\sigma,\sigma')}(E) = \emptyset$, and note that this implies that $\operatorname{supp} \sigma'' \subseteq C_1(B_N) \smallsetminus C_{\mathcal{G}(\sigma,\sigma')}(E)$.

Define

$$E' := \operatorname{supp} \sigma|_{C_1(B_N) \smallsetminus \mathcal{C}_{\mathcal{G}(\sigma,\sigma')}(E)} \cup \operatorname{supp} \sigma'|_{C_1(B_N) \smallsetminus \mathcal{C}_{\mathcal{G}(\sigma,\sigma')}(E)}.$$

Then, since $\sigma'' \leq \sigma$ and $\operatorname{supp} \sigma'' \subseteq C_1(B_N) \setminus C_{\mathcal{G}(\sigma,\sigma')}(E)$, we have $\operatorname{supp} \sigma'' \subseteq E'$. Consequently, by Lemma 5.5, we have $\sigma'' \leq \sigma |_{\mathcal{C}_{\mathcal{G}(\sigma,\sigma')}(E')}$. Since

$$\sigma|_{C_1(B_N)\smallsetminus \mathcal{C}_{\mathcal{G}(\sigma,\sigma')}(E)} = \sigma|_{\mathcal{C}_{\mathcal{G}(\sigma,\sigma')}(E')},$$

we obtain $\sigma'' \leq \sigma|_{C_1(B_N) \setminus C_{\mathcal{G}(\sigma,\sigma')}(E)}$, and hence the desired conclusion holds also in this case.

This completes the proof.

M. P. Forsström

5.2. The set $E_{E_0,\hat{\sigma},\hat{\sigma}'}$. We now define a set which we will need for the definitions of the couplings in Sects. 5.3 and 5.4

Definition 5.7. For σ , $\sigma' \in \Omega^1(B_N, G)$ and $E_0 \subseteq C_1(B_N)$, define

$$E_{E_0,\sigma,\sigma'} := \mathcal{C}_{\mathcal{G}(\sigma,\sigma')} \Big(E_0 \cup \{ e \in \operatorname{supp} \sigma : d\sigma |_{\pm \operatorname{supp} \hat{\partial} e} \neq 0 \} \\ \cup \{ e \in \operatorname{supp} \sigma' : d\sigma' |_{\pm \operatorname{supp} \hat{\partial} e} \neq 0 \} \Big).$$
(5.4)

Lemma 5.8. Let $\sigma, \sigma' \in \Omega^1(B_N, G)$, and let $E_0 \subseteq C_1(B_N)$.

Then

(i) $d(\sigma|_{E_{E_0,\sigma,\sigma'}}) = d\sigma$, and

(*ii*) $d(\sigma|_{C_1(B_N) \smallsetminus E_{E_0,\sigma,\sigma'}}) = 0.$

Proof. To simplify notation, let $E := E_{E_0,\sigma,\sigma'}$.

By Lemma 5.2, applied with σ , σ' , and E, we then have $\sigma|_E \leq \sigma$, $\sigma'|_E \leq \sigma'$, $\sigma|_{C_1(B_N) \setminus E} \leq \sigma$ and $\sigma'|_{C_1(B_N) \setminus E} \leq \sigma'$.

We now show that $d(\sigma|_E) = d\sigma$. Since $\sigma|_E \leq \sigma$, it suffices to show that $d(\sigma|_E)(p) \neq 0$ whenever $d\sigma(p) \neq 0$. To this end, assume that $d\sigma(p) \neq 0$. Then the set supp $\partial p \cap$ supp σ must be non-empty. Fix one edge $e \in \text{supp } \partial p \cap \text{supp } \sigma$. Recalling the definition of E, we see that $e \in E$, and hence any edge $e' \in \text{supp } \partial p \setminus \{e\}$ must satisfy either $\sigma'(e') = \sigma(e') = 0$, or $e' \in E$.

Consequently, $\sigma|_E(\partial p) = \sigma(\partial p)$, and hence

$$d\sigma|_E(p) = \sigma|_E(\partial p) = \sigma(\partial p) = d\sigma(p)$$

as desired. This concludes the proof of 5.8.

To see that (5.8) holds, note simply that, using 5.8, we have

$$d(\sigma|_{C_1(B_N)\smallsetminus E}) = d(\sigma - \sigma|_E) = d\sigma - d(\sigma|_E) = d\sigma - d\sigma = 0,$$

and hence (5.8) holds. This concludes the proof.

Lemma 5.9. Let $\hat{\sigma}, \hat{\sigma}' \in \Omega^1(B_N, G)$, and let $E_0 \subseteq C_1(B_N)$. Further, either let

$$\begin{cases} \sigma := \hat{\sigma}|_{E_{E_0,\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}'|_{C_1(B_N) \smallsetminus E_{E_0,\hat{\sigma},\hat{\sigma}'}} \\ \sigma' := \hat{\sigma}'. \end{cases}$$

or let

$$\begin{aligned} \sigma &:= \hat{\sigma}|_{E_{E_0,\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}'|_{C_1(B_N) \smallsetminus E_{E_0,\hat{\sigma},\hat{\sigma}'}} \\ \sigma' &:= \hat{\sigma}'|_{E_{E_0,\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}|_{C_1(B_N) \smallsetminus E_{E_0,\hat{\sigma},\hat{\sigma}'}}. \end{aligned}$$

Then $E_{E_0,\sigma,\sigma'} = E_{E_0,\hat{\sigma},\hat{\sigma}'}$.

Proof. By Lemma 5.8, we have $d\hat{\sigma}|_{E_{E_0,\sigma,\sigma'}} = d\hat{\sigma}$, and $d\hat{\sigma}'|_{C_1(B_N) \smallsetminus E_{E_0,\hat{\sigma},\hat{\sigma}'}} = 0$ and hence

$$d\sigma = d(\hat{\sigma}|_{E_{E_0,\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}'|_{C_1(B_N) \smallsetminus E_{E_0,\sigma,\sigma'}}) = d(\hat{\sigma}|_{E_{E_0,\hat{\sigma},\hat{\sigma}'}}) + d(\hat{\sigma}'|_{C_1(B_N) \smallsetminus E_{E_0,\sigma,\sigma'}})$$
$$= d\hat{\sigma} + 0 = d\hat{\sigma}.$$

If $e \in \operatorname{supp} \hat{\sigma}$ is such that $d\hat{\sigma}|_{\pm \operatorname{supp} \hat{\partial} e} \neq 0$, then $e \in E_{E_0, \hat{\sigma}, \hat{\sigma}'}$, and thus $\sigma(e) = \hat{\sigma}(e)$, implying in particular that $e \in \operatorname{supp} \sigma$. Since $d\sigma = d\hat{\sigma}$, it follows that $d\sigma|_{\pm \operatorname{supp} \hat{\partial} e} \neq 0$, and hence

$$\{e \in \operatorname{supp} \hat{\sigma} : d\hat{\sigma}|_{\pm \operatorname{supp} \hat{\partial} e} \neq 0\} \subseteq \{e \in \operatorname{supp} \sigma : d\sigma|_{\pm \operatorname{supp} \hat{\partial} e} \neq 0\}.$$

Analogously, we obtain

$$\{e \in \operatorname{supp} \hat{\sigma}' \colon d\hat{\sigma}'|_{\pm \operatorname{supp} \hat{\partial}_e} \neq 0\} \subseteq \{e \in \operatorname{supp} \sigma' \colon d\sigma'|_{\pm \operatorname{supp} \hat{\partial}_e} \neq 0\}.$$

Noting that $\mathcal{G}(\hat{\sigma}, \hat{\sigma}') = \mathcal{G}(\sigma, \sigma')$, we thus obtain

$$\begin{split} E_{E_{0},\hat{\sigma},\hat{\sigma}'} &= \mathcal{C}_{\mathcal{G}(\hat{\sigma},\hat{\sigma}')} \Big(E_{0} \cup \{ e \in \operatorname{supp} \hat{\sigma} : d\hat{\sigma} \mid_{\pm \operatorname{supp} \partial e} \neq 0 \} \cup \{ e \in \operatorname{supp} \hat{\sigma}' : d\hat{\sigma}' \mid_{\pm \operatorname{supp} \partial e} \neq 0 \} \Big) \\ &\subseteq \mathcal{C}_{\mathcal{G}(\sigma,\sigma')} \Big(E_{0} \cup \{ e \in \operatorname{supp} \sigma : d\sigma \mid_{\pm \operatorname{supp} \partial e} \neq 0 \} \cup \\ &\{ e \in \operatorname{supp} \sigma' : d\sigma' \mid_{\pm \operatorname{supp} \partial e} \neq 0 \} \Big) = E_{E_{0},\sigma,\sigma'}. \end{split}$$

For the other direction, assume that $e \in \operatorname{supp} \sigma$ is such that $d\sigma|_{\pm \operatorname{supp} \hat{\partial} e} \neq 0$. Then $\sigma(e) \neq 0$, and there must exist $p \in \hat{\partial} e$ such that $d\sigma(p) \neq 0$.

Since $d\hat{\sigma} = d\sigma$, it follows that $d\hat{\sigma}(p) \neq 0$. Consequently, there must exist $e' \in \partial p$ such that $\hat{\sigma}(e') \neq 0$. For any such edge e', we have $d\hat{\sigma}|_{\pm \operatorname{supp} \hat{\partial} e'} \neq 0$, and hence $e' \in E_{E_0,\hat{\sigma},\hat{\sigma}'}$. In particular, this implies that $\sigma(e') = \hat{\sigma}(e') \neq 0$, and hence $e \in E_{E_0,\hat{\sigma},\hat{\sigma}'}$. Consequently,

$$\{e \in \operatorname{supp} \sigma \colon d\sigma|_{\pm \operatorname{supp} \hat{\partial}_e} \neq 0\} \subseteq E_{E_0, \hat{\sigma}, \hat{\sigma}'}.$$

Analogously, we also obtain

$$\{e \in \operatorname{supp} \sigma' \colon d\sigma'|_{\pm \operatorname{supp} \hat{\partial}_e} \neq 0\} \subseteq E_{E_0, \hat{\sigma}, \hat{\sigma}'}.$$

Again recalling that $\mathcal{G}(\hat{\sigma}, \hat{\sigma}') = \mathcal{G}(\sigma, \sigma')$, we thus obtain

$$E_{E_{0},\sigma,\sigma'} = C_{\mathcal{G}(\sigma,\sigma')} (E_{0} \cup \{e \in \operatorname{supp} \sigma : d\sigma|_{\pm \operatorname{supp} \partial e} \neq 0\}$$
$$\cup \{e \in \operatorname{supp} \sigma' : d\sigma'|_{\pm \operatorname{supp} \partial e} \neq 0\})$$
$$\subseteq C_{\mathcal{G}(\hat{\sigma},\hat{\sigma}')} (E_{0} \cup E_{E_{0},\hat{\sigma},\hat{\sigma}'} \cup \{e \in \operatorname{supp} \hat{\sigma}' : d\hat{\sigma}'|_{\pm \operatorname{supp} \partial e} \neq 0\}) = E_{E_{0},\hat{\sigma},\hat{\sigma}'}.$$

This concludes the proof.

Lemma 5.10. Let $\sigma, \sigma' \in \Omega^1(B_N, G)$, let $E_0 \subseteq C_1(B_N)$, and let $e \in C_1(B_N)$. Then $e \in E_{E_0,\sigma,\sigma'}$ if and only if one of the following holds.

$$\begin{array}{l} (i) \ d(\sigma|_{\mathcal{C}_{\mathcal{G}(\sigma,\sigma')}(e)}) \neq 0\\ (ii) \ d(\sigma'|_{\mathcal{C}_{\mathcal{G}(\sigma,\sigma')}(e)}) \neq 0\\ (iii) \ \mathcal{C}_{\mathcal{G}(\sigma,\sigma')}(e) \cap E_0 \neq \ell \end{array}$$

Proof. Suppose first that $e \in E_{E_0,\sigma,\sigma'}$. By the definition of $C_{\mathcal{G}(\sigma,\sigma')}(e)$, there exists an edge $e' \in C_{\mathcal{G}(\sigma,\sigma')}(e)$ such that

$$e' \in E_0 \cup \{e'' \in \operatorname{supp} \sigma : d\sigma|_{\pm \operatorname{supp} \hat{\partial} e''} \neq 0\} \cup \{e'' \in \operatorname{supp} \sigma' : d\sigma'|_{\pm \operatorname{supp} \hat{\partial} e''} \neq 0\}.$$

If $e' \in E_0$, then $e' \in C_{\mathcal{G}(\sigma,\sigma')}(e) \cap E_0$, and hence $C_{\mathcal{G}(\sigma,\sigma')}(e) \cap E_0 \neq \emptyset$. If $e' \notin E_0$, then, by symmetry, we can assume that $e' \in \{e'' \in \operatorname{supp} \sigma : d\sigma|_{\pm \operatorname{supp}} \hat{\partial}e'' \neq 0\}$.

In this case, we infer that there exists a plaquette $p \in \hat{\partial} e'$ such that $d\sigma(p) \neq 0$. Since $e' \in C_{\mathcal{G}(\sigma,\sigma')}(e)$, we have $\operatorname{supp} \sigma \cap \operatorname{supp} \partial p \subseteq C_{\mathcal{G}(\sigma,\sigma')}(e)$, and so $d(\sigma|_{\mathcal{C}_{\mathcal{G}(\sigma,\sigma')}(e)})$ $(p) = d\sigma(p) \neq 0$.

For the other direction, assume first that $C_{\mathcal{G}(\sigma,\sigma')}(e) \cap E_0 \neq \emptyset$. Then there is $e' \in E_0$ such that $e' \in C_{\mathcal{G}(\sigma,\sigma')}(e)$. Since $e' \in C_{\mathcal{G}(\sigma,\sigma')}(e)$ we must also have $e \in C_{\mathcal{G}(\sigma,\sigma')}(e')$, which is a subset of $E_{E_0,\sigma,\sigma'}$ since $e' \in E_0$. Next, assume instead that there is a plaquette $p \in C_2(B_N)$ such that $d(\sigma|_{\mathcal{C}_{\mathcal{G}(\sigma,\sigma')}(e)})(p) \neq 0$. Then there exists an edge $e' \in \partial p$ with $\sigma(e') \neq 0$ and $e' \in C_{\mathcal{G}(\sigma,\sigma')}(e)$. Thus $\sup p \sigma \cap \sup p \partial p \subseteq C_{\mathcal{G}(\sigma,\sigma')}(e)$, and so $d\sigma(p) = d(\sigma|_{\mathcal{C}_{\mathcal{G}(\sigma,\sigma')}(e)})(p) \neq 0$. In particular, it follows that $e' \in \sup p \partial p \subseteq \{e'' \in \sup p \sigma : d\sigma|_{\pm \sup p \partial e''} \neq 0\}$. Consequently, we must have $e' \in \{e'' \in \sup p \sigma : d\sigma|_{\pm \sup p \partial e''} \neq 0\}$, and hence $e' \in E_{\sigma,\sigma'}$. Since $e' \in C_{\mathcal{G}_{\sigma,\sigma'}(e)}$, we thus have $e \in C_{\mathcal{G}_{\sigma,\sigma'}(e')} \subseteq E_{E_0,\sigma,\sigma'}$ as desired. Using symmetry, this concludes the proof.

Lemma 5.11. Let $\beta_1, \beta_2 \in [0, \infty]$, $\kappa \ge 0$, $E_0 \subseteq C_1(B_N)$, and $\sigma, \hat{\sigma}, \hat{\sigma}' \in \Omega^1(B_N, G)$. *Then*

$$\begin{split} \varphi_{\beta_{1,\kappa}}(\hat{\sigma})\varphi_{\beta_{2,\kappa}}(\hat{\sigma}') \cdot \mathbb{1}\left(\hat{\sigma}|_{E_{E_{0},\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}'|_{C_{1}(B_{N}) \smallsetminus E_{E_{0},\hat{\sigma},\hat{\sigma}'}} = \sigma\right) \\ &= \varphi_{\beta_{1,\kappa}}(\sigma) \sum_{\sigma' \in \Omega^{1}(B_{N},G)} \varphi_{\beta_{2,\kappa}}(\sigma') \cdot \mathbb{1}\left(\hat{\sigma} = \sigma|_{E_{E_{0},\sigma,\sigma'}} + \sigma'|_{C_{1}(B_{N}) \smallsetminus E_{E_{0},\sigma,\sigma'}}\right) \\ &\quad \cdot \mathbb{1}\left(\hat{\sigma}' = \sigma'|_{E_{E_{0},\sigma,\sigma'}} + \sigma|_{C_{1}(B_{N}) \smallsetminus E_{E_{0},\sigma,\sigma'}}\right). \end{split}$$
(5.5)

Proof. By Lemma 5.2, we have $\hat{\sigma}|_{E_{E_0,\hat{\sigma},\hat{\sigma}'}} \leq \hat{\sigma}$ and $\hat{\sigma}'|_{E_{E_0,\hat{\sigma},\hat{\sigma}'}} \leq \hat{\sigma}'$ and hence, by Lemma 4.1,

$$\varphi_{\beta_{1},\kappa}(\hat{\sigma}) = \varphi_{\beta_{1},\kappa}\left(\hat{\sigma}|_{E_{E_{0},\hat{\sigma},\hat{\sigma}'}}\right)\varphi_{\beta_{1},\kappa}\left(\hat{\sigma}|_{C_{1}(B_{N})\smallsetminus E_{E_{0},\hat{\sigma},\hat{\sigma}'}}\right)$$
(5.6)

and

$$\varphi_{\beta_{2},\kappa}(\hat{\sigma}') = \varphi_{\beta_{2},\kappa}\left(\hat{\sigma}'|_{E_{E_{0},\hat{\sigma},\hat{\sigma}'}}\right)\varphi_{\beta_{2},\kappa}\left(\hat{\sigma}'|_{C_{1}(B_{N})\smallsetminus E_{E_{0},\hat{\sigma},\hat{\sigma}'}}\right).$$
(5.7)

Next, by Lemma 5.8, we have

$$d(\hat{\sigma}|_{C_1(B_N) \smallsetminus E_{E_0,\hat{\sigma},\hat{\sigma}'}}) = 0$$

and hence

$$\varphi_{\beta_1,\kappa}\left(\hat{\sigma}|_{C_1(B_N)\smallsetminus E_{E_0,\hat{\sigma},\hat{\sigma}'}}\right) = \varphi_{\kappa}\left(\hat{\sigma}|_{C_1(B_N)\smallsetminus E_{E_0,\hat{\sigma},\hat{\sigma}'}}\right) = \varphi_{\beta_2,\kappa}\left(\hat{\sigma}|_{C_1(B_N)\smallsetminus E_{E_0,\hat{\sigma},\hat{\sigma}'}}\right).$$

Since $\hat{\sigma}'|_{E_{E_0,\hat{\sigma},\hat{\sigma}'}}$ and $\hat{\sigma}|_{C_1(B_N) \setminus E_{E_0,\hat{\sigma},\hat{\sigma}'}}$ have disjoint supports, it also follows that

$$\hat{\sigma}'|_{E_{E_0,\hat{\sigma},\hat{\sigma}'}} \leq \hat{\sigma}'|_{E_{E_0,\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}|_{C_1(B_N) \smallsetminus E_{E_0,\hat{\sigma},\hat{\sigma}'}}.$$

Thus, by Lemma 4.1, it follows that

$$\varphi_{\beta_{2,\kappa}}(\hat{\sigma}'|_{E_{E_{0},\hat{\sigma},\hat{\sigma}'}})\varphi_{\beta_{1,\kappa}}(\hat{\sigma}|_{C_{1}(B_{N})\smallsetminus E_{E_{0},\hat{\sigma},\hat{\sigma}'}}) = \varphi_{\beta_{2,\kappa}}(\hat{\sigma}'|_{E_{E_{0},\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}|_{C_{1}(B_{N})\smallsetminus E_{E_{0},\hat{\sigma},\hat{\sigma}'}}).$$
(5.8)

By symmetry, we also have

$$\varphi_{\beta_1,\kappa}\left(\hat{\sigma}|_{E_{E_0,\hat{\sigma},\hat{\sigma}'}}\right)\varphi_{\beta_2,\kappa}\left(\hat{\sigma}'|_{C_1(B_N)\smallsetminus E_{E_0,\hat{\sigma},\hat{\sigma}'}}\right) = \varphi_{\beta_1,\kappa}\left(\hat{\sigma}|_{E_{E_0,\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}'|_{C_1(B_N)\smallsetminus E_{E_0,\hat{\sigma},\hat{\sigma}'}}\right).$$

Combining (5.6), (5.7), (5.8), and (5.9), it follows that

$$\begin{split} \varphi_{\beta_{1,\kappa}}(\hat{\sigma})\varphi_{\beta_{2,\kappa}}(\hat{\sigma}') & \mathbb{1}\Big(\hat{\sigma}|_{E_{E_{0},\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}'|_{C_{1}(B_{N}) \smallsetminus E_{E_{0},\hat{\sigma},\hat{\sigma}'}} = \sigma\Big) \\ &= \varphi_{\beta_{1,\kappa}}\Big(\hat{\sigma}|_{E_{E_{0},\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}'|_{C_{1}(B_{N}) \smallsetminus E_{E_{0},\hat{\sigma},\hat{\sigma}'}}\Big)\varphi_{\beta_{2,\kappa}}\Big(\hat{\sigma}'|_{E_{E_{0},\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}|_{C_{1}(B_{N}) \smallsetminus E_{E_{0},\hat{\sigma},\hat{\sigma}'}}\Big) \\ & \cdot \mathbb{1}\Big(\hat{\sigma}|_{E_{E_{0},\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}'|_{C_{1}(B_{N}) \smallsetminus E_{E_{0},\hat{\sigma},\hat{\sigma}'}} = \sigma\Big) \\ &= \varphi_{\beta_{1,\kappa}}(\sigma)\varphi_{\beta_{2,\kappa}}\Big(\hat{\sigma}'|_{E_{E_{0},\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}|_{C_{1}(B_{N}) \smallsetminus E_{E_{0},\hat{\sigma},\hat{\sigma}'}}\Big) \cdot \mathbb{1}\Big(\hat{\sigma}|_{E_{E_{0},\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}'|_{C_{1}(B_{N}) \smallsetminus E_{E_{0},\hat{\sigma},\hat{\sigma}'}} = \sigma\Big) \\ &= \varphi_{\beta_{1,\kappa}}(\sigma)\sum_{\sigma'\in\Omega^{1}(B_{N},G)}\varphi_{\beta_{2,\kappa}}(\sigma') \cdot \mathbb{1}\Big(\hat{\sigma}'|_{E_{E_{0},\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}|_{C_{1}(B_{N}) \smallsetminus E_{E_{0},\hat{\sigma},\hat{\sigma}'}} = \sigma'\Big) \\ & \cdot \mathbb{1}\Big(\hat{\sigma}|_{E_{E_{0},\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}'|_{C_{1}(B_{N}) \smallsetminus E_{E_{0},\hat{\sigma},\hat{\sigma}'}} = \sigma\Big). \end{split}$$

$$(5.10)$$

Now fix $\sigma' \in \Omega^1(B_N, G)$ and assume that

$$\begin{cases} \sigma = \hat{\sigma}|_{E_{E_0,\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}'|_{C_1(B_N) \smallsetminus E_{E_0,\hat{\sigma},\hat{\sigma}'}} \\ \sigma' = \hat{\sigma}'|_{E_{E_0,\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}|_{C_1(B_N) \smallsetminus E_{E_0,\hat{\sigma},\hat{\sigma}'}} \end{cases}$$

By Lemma 5.9, we have $E_{E_0,\hat{\sigma},\hat{\sigma}'} = E_{E_0,\sigma,\sigma'}$. Since $\sigma|_{E_{E_0,\hat{\sigma},\hat{\sigma}'}} = \hat{\sigma}|_{E_{E_0,\hat{\sigma},\hat{\sigma}'}}$ and $\sigma'|_{C_1(B_N) \smallsetminus E_{E_0,\hat{\sigma},\hat{\sigma}'}} = \hat{\sigma}|_{C_1(B_N) \smallsetminus E_{E_0,\hat{\sigma},\hat{\sigma}'}}$, it follows that

$$\hat{\sigma} = \sigma|_{E_{E_0,\hat{\sigma},\hat{\sigma}'}} + \sigma'|_{C_1(B_N) \smallsetminus E_{E_0,\hat{\sigma},\hat{\sigma}'}} = \sigma|_{E_{E_0,\sigma,\sigma'}} + \sigma'|_{C_1(B_N) \smallsetminus E_{E_0,\sigma,\sigma'}}.$$

Analogously, since $\sigma'|_{E_{E_0,\hat{\sigma},\hat{\sigma}'}} = \hat{\sigma}'|_{E_{E_0,\hat{\sigma},\hat{\sigma}'}}$ and $\sigma|_{C_1(B_N) \smallsetminus E_{E_0,\hat{\sigma},\hat{\sigma}'}} = \hat{\sigma}'|_{C_1(B_N) \smallsetminus E_{E_0,\hat{\sigma},\hat{\sigma}'}}$, it follows that

$$\hat{\sigma}' = \sigma'|_{E_{E_0,\hat{\sigma},\hat{\sigma}'}} + \sigma|_{C_1(B_N) \smallsetminus E_{E_0,\hat{\sigma},\hat{\sigma}'}} = \sigma'|_{E_{E_0,\sigma,\sigma'}} + \sigma|_{C_1(B_N) \smallsetminus E_{E_0,\sigma,\sigma'}}.$$

This shows that for any $\sigma' \in \Omega^1(B_N, G)$, we have

$$\mathbb{1}\left(\hat{\sigma}'|_{E_{E_{0},\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}|_{C_{1}(B_{N}) \smallsetminus E_{E_{0},\hat{\sigma},\hat{\sigma}'}} = \sigma'\right) \cdot \mathbb{1}\left(\hat{\sigma}|_{E_{E_{0},\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}'|_{C_{1}(B_{N}) \smallsetminus E_{E_{0},\hat{\sigma},\hat{\sigma}'}} = \sigma\right) \\
= \mathbb{1}\left(\hat{\sigma} = \sigma|_{E_{E_{0},\sigma,\sigma'}} + \sigma'|_{C_{1}(B_{N}) \smallsetminus E_{E_{0},\sigma,\sigma'}}\right) \cdot \mathbb{1}\left(\hat{\sigma}' = \sigma'|_{E_{E_{0},\sigma,\sigma'}} + \sigma|_{C_{1}(B_{N}) \smallsetminus E_{E_{0},\sigma,\sigma'}}\right). \tag{5.11}$$

Combining (5.10) and (5.11), we obtain (5.5) as desired.

5.3. A coupling between two \mathbb{Z}_n -models. In this section, we define a coupling between two copies of $\mu_{N,\infty,\kappa}$, constructed to always agree on a given set $E_0 \subseteq C_1(B_N)$

Definition 5.12 [A coupling of two \mathbb{Z}_n -models]. For $\kappa \ge 0$, σ , $\sigma' \in \Omega_0^1(B_N, G)$, $E_0 \subseteq C_1(B_N)$, and $E_{E_0,\sigma,\sigma'} = C_{\mathcal{G}_{\sigma,\sigma'}}(E_0)$, we define

$$\mu_{N,(\infty,\kappa),(\infty,\kappa)}^{E_0}(\sigma,\sigma') := \mu_{N,\infty,\kappa} \times \mu_{N,\infty,\kappa} \left(\left\{ (\hat{\sigma},\hat{\sigma}') \in \Omega_0^1(B_N,G) \times \Omega_0^1(B_N,G) : \sigma = \hat{\sigma} |_{E_{E_0,\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}' |_{C_1(B_N) \smallsetminus E_{E_0,\hat{\sigma},\hat{\sigma}'}} \text{ and } \sigma' = \hat{\sigma}' \right\} \right).$$

We let $\mathbb{E}_{N,(\infty,\kappa),(\infty,\kappa)}^{E_0}$ denote the corresponding expectation.



support of $\hat{\sigma} \in \Omega_0^1(B_N, \mathbb{Z}_2)$.

(A) Red edges correspond to the (B) Blue edges correspond to the support of $\hat{\sigma}' \in \Omega_0^1(B_N, \mathbb{Z}_2)$.

(C) Red edges correspond to the support of $\hat{\sigma}$, and blue edges correspond to the support of $\hat{\sigma}'$.



(D) Red edges correspond to the $\,$ (E) Blue edges correspond to the support of $\hat{\sigma}|_{E_{E_0,\hat{\sigma},\hat{\sigma}'}}$, and blue support of $\hat{\sigma}'|_{C_1(B_N)\smallsetminus E_{\hat{\sigma},\hat{\sigma}'}}$. edges correspond to the support of $\hat{\sigma}'|_{E_{E_0,\hat{\sigma},\hat{\sigma}'}}$.

(F) Red edges correspond to the support of $\hat{\sigma}|_{E_{E_0,\hat{\sigma},\hat{\sigma}'}}$, and purple edges correspond to the support of $\hat{\sigma}'|_{E_N \smallsetminus E_{\hat{\sigma},\hat{\sigma}'}}$.

Fig. 5. Illustration of the coupling $(\sigma, \sigma') \sim \mu_{N,(\infty,\beta),(\infty,\beta)}^{E_0}$ defined in Definition 5.18, simulated on a 2-dimensional lattice, with $G = \mathbb{Z}_2$, and with $E_0 = C_1(B_{N/4})$.

Remark 5.13. When $\sigma, \sigma' \in \Omega_0^1(B_N, G)$, then $d\sigma = d\sigma' = 0$, and hence the definition of $E_{E_0,\sigma,\sigma'}$ in Definition 5.12 in consistent with (5.4).

Remark 5.14. By definition, if $\hat{\sigma}$, $\hat{\sigma}' \sim \mu_{N,\infty,\kappa}$ are independent, and we let $\sigma := \hat{\sigma}|_{E_{E_{\alpha},\hat{\sigma},\hat{\sigma}'}}$ $\hat{\sigma}'|_{C_1(B_N)\smallsetminus E_{E_0,\hat{\sigma},\hat{\sigma}'}}$ and $\sigma':=\hat{\sigma}'$, then $(\sigma,\sigma')\sim \mu_{N,(\infty,\kappa),(\infty,\kappa)}^{E_0}$.

The next result shows that the measure introduced in Definition 5.12 is indeed a coupling.

Proposition 5.15. Let $\kappa \geq 0$, and let $E_0 \subseteq C_1(B_N)$. Then $\mu_{N,(\infty,\kappa),(\infty,\kappa)}^{E_0}$ is a coupling of $\mu_{N,\infty,\kappa}$ and $\mu_{N,\infty,\kappa}$.

Proof. It is immediate from the definition that if $(\sigma, \sigma') \sim \mu_{N,(\infty,\kappa),(\infty,\kappa)}^{E_0}$, then $\sigma' \sim$ $\mu_{N,\infty,\kappa}$, and it is hence sufficient to show that $\sigma \sim \mu_{N,\infty,\kappa}$.

To this end, fix some $\sigma \in \Omega_0^1(B_N)$. We need to show that

$$\begin{split} \mu_{N,\infty,\kappa} &\times \mu_{N,\infty,\kappa} \left(\left\{ (\hat{\sigma}, \hat{\sigma}') \in \Omega_0^1(B_N) \times \Omega_0^1(B_N) : \hat{\sigma}|_{E_{E_0,\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}'|_{C_1(B_N) \smallsetminus E_{E_0,\hat{\sigma},\hat{\sigma}'}} \right. \\ &= \sigma \right\} \right) = \mu_{N,\infty,\kappa}(\sigma), \end{split}$$

or equivalently, that

$$\sum_{\substack{\hat{\sigma} \in \Omega_0^1(B_N,G),\\ \hat{\sigma}' \in \Omega_0^1(B_N,G)}} \varphi_{\kappa}(\hat{\sigma}') \cdot \mathbb{1}\left(\hat{\sigma}|_{E_{E_0,\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}'|_{C_1(B_N) \smallsetminus E_{E_0,\hat{\sigma},\hat{\sigma}'}} = \sigma\right)$$
$$= \varphi_{\kappa}(\sigma) \sum_{\substack{\sigma' \in \Omega_0^1(B_N,G)}} \varphi_{\kappa}(\sigma').$$
(5.12)

We now rewrite the left-hand side of (5.12) in order to see that this equality indeed holds.

To this end, note first that by Lemma 5.11, applied with $\beta_1 = \beta_2 = \infty$, we have

$$\sum_{\substack{\hat{\sigma} \in \Omega_0^1(B_N,G),\\ \hat{\sigma}' \in \Omega_0^1(B_N,G)}} \varphi_{\kappa}(\hat{\sigma}') \mathbb{1}\left(\hat{\sigma}|_{E_{E_0,\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}'|_{C_1(B_N) \smallsetminus E_{E_0,\hat{\sigma},\hat{\sigma}'}} = \sigma\right)$$

$$= \varphi_{\kappa}(\sigma) \sum_{\sigma' \in \Omega_{0}^{1}(B_{N},G)} \varphi_{\kappa}(\sigma') \cdot \sum_{\substack{\hat{\sigma} \in \Omega_{0}^{1}(B_{N},G),\\ \hat{\sigma}' \in \Omega_{0}^{1}(B_{N},G)}} \mathbb{1}\left(\hat{\sigma} = \sigma'|_{E_{E_{0},\sigma,\sigma'}} + \sigma|_{C_{1}(B_{N}) \smallsetminus E_{E_{0},\sigma,\sigma'}}\right)$$

$$(5.13)$$

Since $\sigma, \sigma' \in \Omega_0^1(B_N)$, we can apply Lemma 5.8 to see that $\sigma|_{E_{E_0,\sigma,\sigma'}} + \sigma'|_{C_1(B_N) \smallsetminus E_{E_0,\sigma,\sigma'}} \in \Omega_0^1(B_N, G)$ and $\sigma'|_{E_{E_0,\sigma,\sigma'}} + \sigma|_{C_1(B_N) \smallsetminus E_{E_0,\sigma,\sigma'}} \in \Omega_0^1(B_N, G)$.

From this it follows that the double sum on the right-hand side of (5.13) is equal to 1, and hence we obtain (5.12) as desired. This completes the proof.

Lemma 5.16. Let $\beta, \kappa \geq 0$, let $E_0 \subseteq C_1(B_N)$, and let $(\sigma, \sigma') \in \Omega_0^1(B_N, G) \times \Omega_0^1(B_N, G)$ be such that $\mu_{N,(\infty,\kappa),(\infty,\kappa)}^{E_0}(\sigma, \sigma') \neq 0$. Then $\sigma(e) = \sigma'(e)$ for all $e \in C_1(B_N) \setminus E_{E_0,\sigma,\sigma'}$.

Proof. Since $\mu_{N,(\infty,\kappa),(\infty,\kappa)}^{E_0}(\sigma,\sigma') \neq 0$, by definition, there is $(\hat{\sigma}, \hat{\sigma}') \in \Omega_0^1(B_N, G) \times \Omega_0^1(B_N, G)$ such that $\sigma = \hat{\sigma}|_{E_{E_0,\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}'|_{C_1(B_N) \smallsetminus E_{E_0,\hat{\sigma},\hat{\sigma}'}}$ and $\sigma' = \hat{\sigma}'$. Using Lemma 5.9, we immediately obtain the desired conclusion.

One application of the coupling introduced in Definition 5.12, which will be particularly useful to us, is the following proposition.

Proposition 5.17. Let $\kappa \geq 0$, and let $E_0, E_1 \subseteq C^1(B_N, \mathbb{Z})$ have disjoint supports. Further, let $f_0, f_1: \Omega^1(B_N, G) \to \mathbb{R}$ be such that $f_0(\sigma) = f(\sigma|_{E_0})$ and $f_1(\sigma) = f_1(\sigma|_{E_1})$ for all $\sigma \in \Omega^1(B_N)$.

Then

$$\begin{aligned} \left| \mathbb{E}_{N,\infty,\kappa} \left[f_0(\sigma) f_1(\sigma) \right] - \mathbb{E}_{N,\infty,\kappa} \left[f_0(\sigma) \right] \mathbb{E}_{N,\infty,\kappa} \left[f_1(\sigma) \right] \right| \\ &\leq 2 \| f_0 \|_{\infty} \| f_1 \|_{\infty} \sum_{e \in E_1} \mu^{E_0}_{N,(\infty,\kappa),(\infty,\kappa)} \left(\left\{ (\hat{\sigma}, \hat{\sigma}') \in \Omega^1_0(B_N, G) \times \Omega^1_0(B_N, G) \right\} \right) \\ &: e \in E_{E_0,\hat{\sigma},\hat{\sigma}'} \right\} \right). \end{aligned}$$

$$(5.14)$$

We provide an upper bound on the right hand side of (5.14) in Proposition 6.4.

Proof of Proposition 5.17. To simplify notation, for $\sigma \in \Omega^1(B_N, G)$, let $F(\sigma) := f_1(\sigma) f_2(\sigma)$. Let $\hat{\sigma}, \hat{\sigma}' \sim \mu_{N,\infty,\kappa}$. Note that when $d\hat{\sigma} = d\hat{\sigma}' = 0$, we have $E_{E_0,\hat{\sigma},\hat{\sigma}'} = C_{\mathcal{G}(\hat{\sigma},\hat{\sigma}')}(E_0)$.

Define

$$\begin{cases} \sigma := \hat{\sigma}|_{E_{E_0,\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}'|_{C_1(B_N) \smallsetminus E_{E_0,\hat{\sigma},\hat{\sigma}'}} \\ \sigma' := \hat{\sigma}'. \end{cases}$$

Then $(\sigma, \sigma') \sim \mu_{N,(\infty,\kappa),(\infty,\kappa)}^{E_0}$, and hence $\sigma, \sigma' \sim \mu_{N,\infty,\kappa}$. Consequently, we have

$$\mathbb{E}_{N,\infty,\kappa} \left[F(\sigma) \right] = \mathbb{E}_{N,(\infty,\kappa),(\infty,\kappa)}^{L_0} \left[F(\sigma) \right]$$

$$= \mathbb{E}_{N,\infty,\kappa} \times \mathbb{E}_{N,\infty,\kappa} \left[F(\hat{\sigma}|_{E_{E_0,\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}'|_{C_1(B_N) \smallsetminus E_{E_0,\hat{\sigma},\hat{\sigma}'}} \right]$$

$$= \mathbb{E}_{N,\infty,\kappa} \times \mathbb{E}_{N,\infty,\kappa} \left[F(\hat{\sigma}|_{E_{E_0,\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}'|_{C_1(B_N) \smallsetminus E_{E_0,\hat{\sigma},\hat{\sigma}'}} \right) \cdot$$

$$\mathbb{1}(E_1 \cap E_{E_0,\hat{\sigma},\hat{\sigma}'}) = \emptyset) \right]$$

$$+ \mathbb{E}_{N,\infty,\kappa} \times \mathbb{E}_{N,\infty,\kappa} \left[F(\hat{\sigma}|_{E_{E_0,\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}'|_{C_1(B_N) \smallsetminus E_{E_0,\hat{\sigma},\hat{\sigma}'}} \right) \cdot$$

$$\mathbb{1}(E_1 \cap E_{E_0,\hat{\sigma},\hat{\sigma}'}) \neq \emptyset) \right]. \tag{5.15}$$

Since $E_{E_0,\hat{\sigma},\hat{\sigma}'} = C_{\mathcal{G}(\hat{\sigma},\hat{\sigma}')}(E_0)$, we have $(\operatorname{supp} \hat{\sigma} \cup \operatorname{supp} \hat{\sigma}') \cap E_0 \subseteq E_{E_0,\hat{\sigma},\hat{\sigma}'}$. This implies in particular that

 $(\hat{\sigma}|_{E_{E_0,\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}'|_{C_1(B_N) \smallsetminus E_{E_0,\hat{\sigma},\hat{\sigma}'}})|_{E_0} = (\hat{\sigma}|_{E_{E_0,\hat{\sigma},\hat{\sigma}'}})|_{E_0} + (\hat{\sigma}'|_{C_1(B_N) \smallsetminus E_{E_0,\hat{\sigma},\hat{\sigma}'}})|_{E_0} = \hat{\sigma}|_{E_0},$ and hence

$$f_0(\hat{\sigma}|_{E_{E_0,\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}'|_{C_1(B_N) \smallsetminus E_{E_0,\hat{\sigma},\hat{\sigma}'}}) = f_0(\hat{\sigma}).$$

At the same time, on the event $E_1 \cap E_{E_0,\hat{\sigma},\hat{\sigma}'} = \emptyset$, for all $e \in E_1$ we have

$$\begin{aligned} (\hat{\sigma}|_{E_{E_0,\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}'|_{C_1(B_N) \smallsetminus E_{E_0,\hat{\sigma},\hat{\sigma}'}})|_{E_1} &= (\hat{\sigma}|_{E_{E_0,\hat{\sigma},\hat{\sigma}'}})|_{E_1} + (\hat{\sigma}'|_{C_1(B_N) \smallsetminus E_{E_0,\hat{\sigma},\hat{\sigma}'}})|_{E_1} \\ &= 0 + \hat{\sigma}'|_{E_1}, \end{aligned}$$

and hence

$$f_1(\hat{\sigma}|_{E_{E_0,\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}'|_{C_1(B_N) \smallsetminus E_{E_0,\hat{\sigma},\hat{\sigma}'}}) = f_1(\hat{\sigma}').$$

Consequently, on the event $E_1 \cap E_{E_0,\sigma,\sigma'} = \emptyset$, we have

$$F(\hat{\sigma}|_{E_{E_0,\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}'|_{C_1(B_N) \smallsetminus E_{E_0,\hat{\sigma},\hat{\sigma}'}}) = f_0(\hat{\sigma}) f_1(\hat{\sigma}'),$$

and hence

$$\begin{split} \mathbb{E}_{N,\infty,\kappa} \times \mathbb{E}_{N,\infty,\kappa} \Big[F(\hat{\sigma}|_{E_{E_0,\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}'|_{C_1(B_N) \smallsetminus E_{E_0,\hat{\sigma},\hat{\sigma}'}}) \cdot \mathbb{1}(E_1 \cap E_{E_0,\hat{\sigma},\hat{\sigma}'} = \emptyset) \Big] \\ &= \mathbb{E}_{N,\infty,\kappa} \times \mathbb{E}_{N,\infty,\kappa} \Big[f_0(\hat{\sigma}) f_1(\hat{\sigma}') \cdot \mathbb{1}(E_1 \cap E_{E_0,\hat{\sigma},\hat{\sigma}'} = \emptyset) \Big] \\ &= \mathbb{E}_{N,\infty,\kappa} \times \mathbb{E}_{N,\infty,\kappa} \Big[f_0(\hat{\sigma}) f_1(\hat{\sigma}') \Big] - \mathbb{E}_{N,\infty,\kappa} \times \mathbb{E}_{N,\infty,\kappa} \Big[f_0(\hat{\sigma}) f_1(\hat{\sigma}') \\ \cdot \mathbb{1}(E_1 \cap E_{E_0,\hat{\sigma},\hat{\sigma}'} \neq \emptyset) \Big] \\ &= \mathbb{E}_{N,\infty,\kappa} \Big[f_0(\hat{\sigma}) \Big] \mathbb{E}_{N,\infty,\kappa} \Big[f_1(\hat{\sigma}') \Big] - \mathbb{E}_{N,\infty,\kappa} \times \mathbb{E}_{N,\infty,\kappa} \Big[f_0(\hat{\sigma}) f_1(\hat{\sigma}') \\ \cdot \mathbb{1}(E_1 \cap E_{E_0,\hat{\sigma},\hat{\sigma}'} \neq \emptyset) \Big]. \end{split}$$

Inserting this into (5.15), we see that

$$\begin{split} & \mathbb{E}_{N,\infty,\kappa} \Big[f_0(\sigma) f_1(\sigma) \Big] \\ &= \mathbb{E}_{N,\infty,\kappa} \Big[f_0(\hat{\sigma}) \Big] \mathbb{E}_{N,\infty,\kappa} \Big[f_1(\hat{\sigma}') \Big] - \mathbb{E}_{N,\infty,\kappa} \times \mathbb{E}_{N,\infty,\kappa} \Big[f_0(\hat{\sigma}) f_1(\hat{\sigma}') \\ & \cdot \mathbb{1} (E_1 \cap E_{E_0,\hat{\sigma},\hat{\sigma}'}) \neq \emptyset) \Big] \\ & + \mathbb{E}_{N,\infty,\kappa} \times \mathbb{E}_{N,\infty,\kappa} \Big[F(\hat{\sigma}'|_{E_{E_0,\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}|_{C_1(B_N) \smallsetminus E_{E_0,\hat{\sigma},\hat{\sigma}'}}) \cdot \mathbb{1} (E_1 \cap E_{E_0,\hat{\sigma},\hat{\sigma}'} \neq \emptyset) \Big]. \end{split}$$

In particular, this implies that

$$\begin{split} \left| \mathbb{E}_{N,\infty,\kappa} \Big[f_0(\sigma) f_1(\sigma) \Big] - \mathbb{E}_{N,\infty,\kappa} \Big[f_0(\hat{\sigma}) \Big] \mathbb{E}_{N,\infty,\kappa} \Big[f_1(\hat{\sigma}') \Big] \right| \\ &\leq 2 \| f_0 \|_{\infty} \| f_1 \|_{\infty} \mu_{N,\infty,\kappa} \times \mu_{N,\infty,\kappa} \Big(E_1 \cap E_{E_0,\hat{\sigma},\hat{\sigma}'} \neq \emptyset \Big) \\ &\leq 2 \| f_0 \|_{\infty} \| f_1 \|_{\infty} \sum_{e \in E_1} \mu_{N,\infty,\kappa} \times \mu_{N,\infty,\kappa} \Big(e \in E_{E_0,\hat{\sigma},\hat{\sigma}'} \neq \emptyset \Big). \end{split}$$

To obtain the desired conclusion, we note that by Lemma 5.9, we have $E_{E_0,\hat{\sigma},\hat{\sigma}'} = E_{E_0,\sigma,\sigma'}$. This concludes the proof.

5.4. A coupling between the Abelian Higgs model and the \mathbb{Z}_n -model. In this section, we recall the coupling between $\mu_{N,\beta,\kappa}$ and $\mu_{N,\infty,\kappa}$ introduced in [18].

Definition 5.18 [The coupling to the \mathbb{Z}_n model]. For $\beta, \kappa \ge 0, \sigma \in \Omega^1(B_N, G)$, and $\sigma' \in \Omega_0^1(B_N, G)$, let let

$$E_{\sigma,\sigma'} := E_{\emptyset,\sigma,\sigma'} = \mathcal{C}_{\mathcal{G}(\sigma,\sigma')} \big(\{ e \in \operatorname{supp} \sigma : d\sigma |_{\pm \operatorname{supp} \hat{\partial} e} \neq 0 \} \big).$$
(5.16)

and define

$$\mu_{N,(\beta,\kappa),(\infty,\kappa)}(\sigma,\sigma') := \mu_{N,\beta,\kappa} \times \mu_{N,\infty,\kappa} \left(\left\{ (\hat{\sigma},\hat{\sigma}') \in \Omega^1(B_N,G) \times \Omega^1_0(B_N,G) : \sigma = \hat{\sigma} |_{E_{\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}' |_{C_1(B_N) \smallsetminus E_{\hat{\sigma},\hat{\sigma}'}} \text{ and } \sigma' = \hat{\sigma}' \right\} \right).$$

We let $\mathbb{E}_{N,(\beta,\kappa),(\infty,\kappa)}$ denote the corresponding expectation.

Remark 5.19. If [18], the measure $\mu_{N,(\beta,\kappa),(\infty,\kappa)}$ in Definition 5.18 above was defined slightly differently, but using the Proposition 5.20 below, one easily shows that they are equivalent.

The next result shows that this is indeed a coupling.

Proposition 5.20. Let β , $\kappa \geq 0$. Then $\mu_{N,(\beta,\kappa),(\infty,\kappa)}$ is a coupling of $\mu_{N,\beta,\kappa}$ and $\mu_{N,\infty,\kappa}$.

Proof. It is immediate from the definition that if $(\sigma, \sigma') \sim \mu_{N,(\beta,\kappa),(\infty,\kappa)}$, then $\sigma' \sim \mu_{N,\infty,\kappa}$, and it is hence sufficient to show that $\sigma \sim \mu_{N,\beta,\kappa}$. This is exactly equivalent to, for each $\sigma \in \Omega^1(B_N, G)$, showing that

$$\mu_{N,\beta,\kappa} \times \mu_{N,\infty,\kappa} \left(\left\{ (\hat{\sigma}, \hat{\sigma}') \in \Omega^1(B_N) \times \Omega^1_0(B_N) \colon \hat{\sigma}|_{E_{\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}'|_{C_1(B_N) \smallsetminus E_{\hat{\sigma},\hat{\sigma}'}} = \sigma \right\} \right)$$

= $\mu_{N,\beta,\kappa}(\sigma).$





 $d\hat{\sigma}$.

(A) Red edges correspond to the (B) Blue edges correspond to the (C) Red edges correspond to the $d\hat{\sigma}' = 0.$



support of $\hat{\sigma}|_{E_{\hat{\sigma},\hat{\sigma}'}}$, blue edges cor-support of $\hat{\sigma}'|_{C_1(B_N)\smallsetminus E_{\hat{\sigma},\hat{\sigma}'}}$. respond to the support of $\hat{\sigma}'|_{E_{\hat{\sigma},\hat{\sigma}'}}$, and black squares correspond to the support of $d\hat{\sigma}$.





support of $\hat{\sigma} \in \Omega^1(B_N, \mathbb{Z}_2)$, and support of $\hat{\sigma}' \in \Omega^1(B_N, \mathbb{Z}_2)$. In support of $\hat{\sigma}$, blue edges correblack squares to the support of this case we automatically have spond to the support of $\hat{\sigma}'$, and black squares to the support of $d\hat{\sigma}$.



(D) Red edges correspond to the (E) Blue edges correspond to the (F) Red edges correspond to the support of $\hat{\sigma}|_{E_{\hat{\sigma},\hat{\sigma}'}}$, blue edges correspond to the support of $\hat{\sigma}'|_{C_1(B_N)\smallsetminus E_{\hat{\sigma},\hat{\sigma}'}}$, and black squares correspond to the support of $d\hat{\sigma}$.

Fig. 6. Illustration of the coupling $(\sigma, \sigma') \sim \mu_{N,(\beta,\kappa),(\infty,\kappa)}$ defined in Definition 5.18 (simulated on a 2-dimensional lattice, with $G = \mathbb{Z}_2$)

or, equivalently, that

$$\sum_{\substack{\hat{\sigma} \in \Omega^{1}(B_{N},G),\\ \hat{\sigma}' \in \Omega^{1}_{0}(B_{N},G)}} \varphi_{\beta,\kappa}(\hat{\sigma})\varphi_{\kappa}(\hat{\sigma}') \cdot \mathbb{1}\left(\hat{\sigma}|_{E_{\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}'|_{C_{1}(B_{N}) \smallsetminus E_{\hat{\sigma},\hat{\sigma}'}}\right)$$
$$= \sigma = \varphi_{\beta,\kappa}(\sigma) \sum_{\substack{\sigma' \in \Omega^{1}_{0}(B_{N},G)}} \varphi_{\kappa}(\sigma').$$
(5.17)

We now show that (5.17) holds. To this end, fix some $\sigma \in \Omega^1(B_N, G)$. By Lemma 5.11, applied with $\beta_1 = \beta$ and $\beta_2 = \infty$, we have

$$\sum_{\substack{\hat{\sigma} \in \Omega^{1}(B_{N},G),\\ \hat{\sigma}' \in \Omega^{1}_{0}(B_{N},G)}} \varphi_{\beta,\kappa}(\sigma) \varphi_{\kappa}(\sigma') \mathbb{1} \left(\hat{\sigma} |_{E_{\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}' |_{C_{1}(B_{N}) \smallsetminus E_{\hat{\sigma},\hat{\sigma}'}} = \sigma \right)$$

$$= \varphi_{\kappa}(\sigma) \sum_{\sigma' \in \Omega^{1}(B_{N},G)} \varphi_{\beta,\kappa}(\sigma') \sum_{\substack{\hat{\sigma} \in \Omega^{1}(B_{N},G),\\\hat{\sigma}' \in \Omega^{1}(B_{N},G)}} \mathbb{1}\left(\hat{\sigma} = \sigma|_{E_{\sigma,\sigma'}} + \sigma'|_{C_{1}(B_{N}) \smallsetminus E_{\sigma,\sigma'}}\right)$$
$$= \varphi_{\kappa}(\sigma) \sum_{\sigma' \in \Omega^{1}(B_{N},G)} \varphi_{\beta,\kappa}(\sigma') \mathbb{1}\left(\sigma'|_{E_{\sigma,\sigma'}} + \sigma|_{C_{1}(B_{N}) \smallsetminus E_{\sigma,\sigma'}} \in \Omega^{1}_{0}(B_{N},G)\right)$$
$$= \varphi_{\kappa}(\sigma) \sum_{\sigma' \in \Omega^{1}(B_{N},G)} \varphi_{\beta,\kappa}(\sigma') \mathbb{1}\left(d\left(\sigma'|_{E_{\sigma,\sigma'}} + \sigma|_{C_{1}(B_{N}) \smallsetminus E_{\sigma,\sigma'}}\right) = 0\right).$$
(5.18)

By Lemma 5.8, we have

$$d(\sigma'|_{E_{\sigma,\sigma'}} + \sigma|_{C_1(B_N) \smallsetminus E_{\sigma,\sigma'}}) = d(\sigma'|_{E_{\sigma,\sigma'}}) + d(\sigma|_{C_1(B_N) \smallsetminus E_{\sigma,\sigma'}}) = d(\sigma'|_{E_{\sigma,\sigma'}}) + 0$$

= $d\sigma'$,

and hence

$$\varphi_{\kappa}(\sigma) \sum_{\sigma' \in \Omega^{1}(B_{N},G)} \varphi_{\beta,\kappa}(\sigma') \mathbb{1} \left(d\left(\sigma'|_{E_{\sigma,\sigma'}} + \sigma|_{C_{1}(B_{N}) \smallsetminus E_{\sigma,\sigma'}}\right) = 0 \right)$$

$$= \varphi_{\kappa}(\sigma) \sum_{\sigma' \in \Omega^{1}(B_{N},G)} \varphi_{\beta,\kappa}(\sigma') \mathbb{1} (d\sigma' = 0) = \varphi_{\kappa}(\sigma) \sum_{\sigma' \in \Omega^{1}_{0}(B_{N},G)} \varphi_{\beta,\kappa}(\sigma').$$
(5.19)

Combining (5.18) and (5.19), we obtain (5.17) as desired. This concludes the proof. \Box

Lemma 5.21. Let $\beta, \kappa \geq 0$, let $E_0 \subseteq C_1(B_N)$, and let $(\sigma, \sigma') \in \Omega^1(B_N, G) \times \Omega_0^1(B_N, G)$ be such that $\mu_{N,(\beta,\kappa),(\infty,\kappa)}(\sigma, \sigma') \neq 0$. Then $\sigma(e) = \sigma'(e)$ for all $e \in C_1(B_N) \setminus E_{\sigma,\sigma'}$.

Proof. Since $\mu_{N,(\beta,\kappa),(\infty,\kappa)}(\sigma,\sigma') \neq 0$, by definition, there is $(\hat{\sigma}, \hat{\sigma}') \in \Omega^1(B_N, G) \times \Omega^1_0(B_N, G)$ such that $\sigma = \hat{\sigma}|_{E_{\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}'|_{C_1(B_N) \smallsetminus E_{\hat{\sigma},\hat{\sigma}'}}$ and $\sigma' = \hat{\sigma}'$. Using Lemma 5.9, we immediately obtain the desired conclusion.

6. Distribution of Vortices and Edge Configurations

In this section, we use the edge graph defined in Sect. 5.1 to give upper bounds on several useful events. Throughout this section, constants K_1, K_2, \ldots, K_{15} will be introduced. We use distinct names for these to make it possible to find explicit upper bounds, but stress that under the assumptions of the main results these are all bounded from above, and will thus not affect the decay rate of the upper bounds obtained throughout this section.

For $E_0 \subseteq C_1(B_N)$ and $e \in C_1(B_N)$, we define

$$dist_1(e, E_0) := \frac{1}{2} \min \left\{ \left| \mathcal{C}_{\mathcal{G}_{\hat{\sigma}, \hat{\sigma}'}}(e) \right| : \sigma, \sigma' \in \Omega^1(B_N, G), \ \mathcal{C}_{\mathcal{G}_{\hat{\sigma}, \hat{\sigma}'}}(e) \cap E_0 \neq \emptyset \right\} \\ = \frac{1}{2} \min \left\{ \left| \mathcal{C}_{\mathcal{G}_{\hat{\sigma}}}(e) \right| : \sigma \in \Omega^1(B_N, G), \ \mathcal{C}_{\mathcal{G}_{\hat{\sigma}}}(e) \cap E_0 \neq \emptyset \right\}, \ e \in C_1(B_N),$$

and

$$\operatorname{dist}_{0}(e, E_{0}) := \frac{1}{2} \min \left\{ \left| \mathcal{C}_{\mathcal{G}_{\hat{\sigma}, \hat{\sigma}'}}(e) \right| : \sigma, \sigma' \in \Omega_{0}^{1}(B_{N}, G), \ \mathcal{C}_{\mathcal{G}_{\hat{\sigma}, \hat{\sigma}'}}(e) \cap E_{0} \neq \emptyset \right\},\ e \in C_{1}(B_{N}).$$

Note that, by Lemma 2.9, if $e \notin E$, then dist $_0(e, E) \ge 8$. We extend this definition to sets $E \subseteq C_1(B_N)$ by letting dist $_1(E, E_0) := \min_{e \in E} \text{dist}_1(e, E_0)$ and dist $_0(E, E_0) := \min_{e \in E} \text{dist}_0(e, E_0)$.

In this section, we will state and prove the following three propositions.

Proposition 6.1. Let β , κ_1 , $\kappa_2 \in [0, \infty]$ be such that $18^2(\alpha_0(\kappa_1) + \alpha_0(\kappa_2) + \alpha_0(\kappa_1)\alpha_0(\kappa_2))$ < 1, let $e \in C_1(B_N)$ be such that $dist_0(e, \partial C_1(B_N)) \ge 8$, and let $M \ge 1$ and $M' \ge 0$. Then

$$\begin{split} &\mu_{N,\beta,\kappa_{1}} \times \mu_{N,\infty,\kappa_{2}} \Big(\Big\{ (\hat{\sigma},\hat{\sigma}') \in \Omega^{1}(B_{N},G) \times \Omega^{1}_{0}(B_{N},G) : \\ &|\mathcal{C}_{\mathcal{G}(\hat{\sigma},\hat{\sigma}')}(e)| \geq 2M, \text{ and } |\text{supp } d(\hat{\sigma}|_{\mathcal{C}_{\mathcal{G}(\hat{\sigma},\hat{\sigma}')}(e)}) \Big| \geq 2M' \Big\} \Big) \\ &\leq \mathbb{1}_{M'>0} \cdot \mathbb{1}_{M=1} \cdot \Big(18^{2} \big(\alpha_{0}(\kappa_{1}) + \alpha_{0}(\kappa_{2}) + \alpha_{0}(\kappa_{1})\alpha_{0}(\kappa_{2}) \big) \Big) \alpha_{1}(\beta)^{\max(6,M')} \\ &+ \mathbb{1}_{M'>0} \cdot K_{1} \Big(18^{2} \big(\alpha_{0}(\kappa_{1}) + \alpha_{0}(\kappa_{2}) + \alpha_{0}(\kappa_{1})\alpha_{0}(\kappa_{2}) \big) \Big)^{\max(M,2)} \alpha_{1}(\beta)^{\max(6,M')} \\ &+ \mathbb{1}_{M'\in\{1,2,3,4,5\}} \cdot K_{1} \Big(18^{2} \big(\alpha_{0}(\kappa_{1}) + \alpha_{0}(\kappa_{2}) + \alpha_{0}(\kappa_{1})\alpha_{0}(\kappa_{2}) \big) \Big)^{\dim(4,8)} \alpha_{1}(\beta) \\ &+ \mathbb{1}_{M'=0} \cdot K_{1} \Big(18^{2} \big(\alpha_{0}(\kappa_{1}) + \alpha_{0}(\kappa_{2}) + \alpha_{0}(\kappa_{1})\alpha_{0}(\kappa_{2}) \big) \Big)^{\max(M,8)}, \end{split}$$

where

$$K_1 = K_1(\kappa_1, \kappa_2) := 18^{-3} \left(1 - 18^2 \left(\alpha_0(\kappa_1) + \alpha_0(\kappa_2) + \alpha_0(\kappa_1)\alpha_0(\kappa_2) \right) \right)^{-1}.$$
 (6.2)

Remark 6.2. If $\kappa_1 = \kappa_2 =: \kappa$, then the assumption on κ_1 and κ_2 in Proposition (6.1) is equivalent to 3.

Proposition 6.3. Let $\beta, \kappa \ge 0$ be such that 3 holds, and assume that $p \in C_2(B_N)$ is such that dist₀(supp $\partial e, \partial C_1(B_N)$)) ≥ 8 . Then

$$\mu_{N,\beta,\kappa}\left(\left\{\hat{\sigma}\in\Omega^1(B_N,G)\colon d\hat{\sigma}(p)\neq 0\right\}\right)\leq K_2\alpha_2(\beta,\kappa),\tag{6.3}$$

where

$$K_2 := 4 \left(18^2 + 18\alpha_0(\kappa) \left(1 - 18^2 \alpha_0(\kappa) \right)^{-1} \right) \frac{\alpha_1(\beta)^6}{\alpha_0(\beta)^6}.$$
 (6.4)

Proposition 6.4. Let $\kappa \ge 0$ be such that 3 holds, let $E_0 \subseteq C_1(B_N)$ be non-empty, and let $e \in C_1(B_N)$ be such that $dist_0(e, \partial C_1(B_N))) \ge 8$.

Then

$$\mu_{N,(\infty,\kappa),(\infty,\kappa)}^{E_0}\left(\left\{(\sigma,\sigma')\in\Omega_0^1(B_N,G)\times\Omega_0^1(B_N,G)\colon e\in E_{E_0,\sigma,\sigma'}\right\}\right)$$

$$\leq K_3\left(K_4\alpha_0(\kappa)\right)^{\operatorname{dist}_0(e,E_0)}$$
(6.5)

where

$$K_3 := 18^{-3} (1 - 18^2 (2 + \alpha_0(\kappa)) \alpha_0(\kappa))^{-1}, \quad and \quad K_4 := 18^2 (2 + \alpha_0(\kappa)).$$
(6.6)

Proposition 6.5. Let β , $\kappa \ge 0$ be such that 3 hold, let $e \in C_1(B_N)$ be such that for all $p \in \hat{\partial}e$ we have dist₀(supp ∂p , $\partial C_1(B_N)$) ≥ 8 , and let $E_0 := \{e' \in C_1(B_N) : \hat{\partial}e' \cap \hat{\partial}e \neq \emptyset\}$. Then

$$\mu_{N,\beta,\kappa} \left(\left\{ \sigma \in \Omega^1(B_N, G) : |\mathcal{C}_{\mathcal{G}(\sigma)}(E_0)| \ge 2M \text{ and } \left| \operatorname{supp} d(\sigma|_{\mathcal{C}_{\mathcal{G}(\sigma)}(E_0)}) \right| \ge 2M' \right\} \right) \\ \le K_5 \left(18^2 \alpha_0(\kappa) \right)^M \alpha_1(\beta)^{M'}.$$

where

$$K_5 := (18^{-3} + 18^{-1})(1 - 18^2 \alpha_0(\kappa))^{-1}.$$
(6.7)

Before we give proofs of the above propositions, we introduce some additional notation and prove two useful lemmas. To this end, we first define a graph $\overline{\mathcal{G}}$ as follows. Fix some $g \in G \setminus \{0\}$ and define $\overline{\sigma} \in \Omega^1(B_N, G)$ by letting $\sigma(e) = g$ for all $e \in C_1(B_N)^+$. Let $\overline{\mathcal{G}} := \mathcal{G}(\overline{\sigma}, 0)$ and note that $\overline{\mathcal{G}}$ does not depend on the choice of g. Note also that if $\sigma, \sigma' \in \Omega^1(B_N, G)$, then $\mathcal{G}(\sigma, \sigma')$ is a subgraph of $\overline{\mathcal{G}}$.

Lemma 6.6. [See also Lemma 7.15 and Lemma 7.16 in [18]] Let $e \in C_1^+(B_N)$, and let $m \ge 1$. Then

$$\left|\left\{E \subseteq C_1^+(B_N) \colon e \in E, |E| = m, \text{ and } \overline{\mathcal{G}}|_E \text{ is connected}\right\}\right| \le 18^{\max(0,2m-3)}.$$

Proof. Since the case m = 1 is trivial, we can assume that $m \ge 2$.

Fix some set $E \subseteq C_1^+(B_N)$ such that $e \in E$, |E| = m, and $\overline{\mathcal{G}}|_E$ is connected.

Since the graph $\overline{\mathcal{G}}|_E$ is connected, it has a spanning tree. Let \mathcal{T} be such a spanning tree. By definition, \mathcal{T} must contain exactly m-1 edges. Since any spanning tree is connected, \mathcal{T} must have a spanning walk which uses each edge in \mathcal{T} exactly twice, and starts and ends at the same vertex. This walk must have length 2(m-1) = 2m-2. By removing one of the edges adjacent to the vertex e, we obtain a spanning walk of $\overline{\mathcal{G}}|_E$ which has length 2m-3, starts at the vertex e and visits every vertex of $\overline{\mathcal{G}}|_E$ at least once.

Since for each $e' \in E_N$, we have $|\{e'' \in C_1(B_N) \setminus \{e'\}: \hat{\partial}e'' \cap \hat{\partial}e' \neq \emptyset\}| = 6 \cdot 3 = 18$, there can exists at most 18^{2m-3} walks in $\overline{\mathcal{G}}$ which starts at e and has length 2m - 3, and hence the desired conclusion follows.

Lemma 6.7. Let $\kappa_1, \kappa_2 \ge 0$, and let $E \subseteq C_1^+(B_N)$. Then

$$\sum_{\substack{\hat{\hat{\sigma}} \in \Omega^{1}(B_{N},G), \, \hat{\hat{\sigma}}' \in \Omega^{1}(B_{N},G):\\(\operatorname{supp} \hat{\hat{\sigma}} \cup \operatorname{supp} \hat{\hat{\sigma}}')^{+} = E}} \varphi_{\kappa_{1}}(\hat{\hat{\sigma}})\varphi_{\kappa_{2}}(\hat{\hat{\sigma}}') \leq (\alpha_{0}(\kappa_{1}) + \alpha_{0}(\kappa_{2}) + \alpha_{0}(\kappa_{1})\alpha_{0}(\kappa_{2}))^{|E|}.$$
(6.8)

Proof. If $\hat{\sigma} \in \Omega^1(B_N, G)$ and $e' \notin \operatorname{supp} \hat{\sigma}$, then, for any $\kappa \ge 0$, we have $\varphi_{\kappa}(\hat{\sigma}(e')) = \varphi_{\kappa}(0) = 1$. Also, if $\hat{\sigma}, \hat{\sigma}' \in \Omega^1(B_N, G)$ and $e' \in (\operatorname{supp} \hat{\sigma} \cup \operatorname{supp} \hat{\sigma}')^+ = E$, then either $\hat{\sigma}(e') \neq 0$ and $\hat{\sigma}'(e') = 0$, $\hat{\sigma}(e') = 0$ and $\hat{\sigma}'(e') \neq 0$, or $\hat{\sigma}(e'), \hat{\sigma}'(e') \neq 0$.

Combining these observations, we find that

$$\begin{split} &\sum_{\hat{\hat{\sigma}} \in \Omega^{1}(B_{N},G), \, \hat{\hat{\sigma}}' \in \Omega^{1}(B_{N},G):} \prod_{e' \in C_{1}(B_{N})^{+}} \varphi_{\kappa_{1}}(\hat{\hat{\sigma}}(e'))^{2} \prod_{e'' \in C_{1}(B_{N})^{+}} \varphi_{\kappa_{2}}(\hat{\hat{\sigma}}'(e'')) \\ &(\operatorname{supp} \hat{\hat{\sigma}} \cup \operatorname{supp} \hat{\hat{\sigma}}')^{+} = E \end{split} \\ &\leq \prod_{e' \in E} \left\{ \varphi_{\kappa_{1}}(0)^{2} \bigg(\sum_{\hat{\hat{\sigma}}'(e') \in G \smallsetminus \{0\}} \varphi_{\kappa_{2}}(\hat{\hat{\sigma}}'(e'))^{2} \bigg) + \bigg(\sum_{\hat{\hat{\sigma}}(e') \in G \smallsetminus \{0\}} \varphi_{\kappa_{1}}(\hat{\hat{\sigma}}(e'))^{2} \bigg) \varphi_{\kappa_{2}}(0)^{2} \\ &+ \bigg(\sum_{\hat{\hat{\sigma}}(e') \in G \smallsetminus \{0\}} \varphi_{\kappa_{1}}(\hat{\hat{\sigma}}(e'))^{2} \bigg) \bigg(\sum_{\hat{\hat{\sigma}}'(e') \in G \smallsetminus \{0\}} \varphi_{\kappa_{2}}(\hat{\hat{\sigma}}'(e'))^{2} \bigg) \bigg\} \\ &= \prod_{e' \in E} \bigg(\alpha_{0}(\kappa_{1}) + \alpha_{0}(\kappa_{2}) + \alpha_{0}(\kappa_{1})\alpha_{0}(\kappa_{2}) \bigg) = \big(\alpha_{0}(\kappa_{1}) + \alpha_{0}(\kappa_{2}) + \alpha_{0}(\kappa_{1})\alpha_{0}(\kappa_{2}) \big)^{|E|}. \end{split}$$

This concludes the proof.

Proof of Proposition 6.1. Since $C_{\mathcal{G}(\hat{\sigma},\hat{\sigma}')}(e)$ is symmetric, induces a connected subgraph in $\overline{\mathcal{G}}$, and contains *e* if it non-empty, we have

$$\begin{split} & \mu_{N,\beta,\kappa_{1}} \times \mu_{N,\infty,\kappa_{2}} \Big(\Big\{ (\hat{\sigma},\hat{\sigma}') \in \Omega^{1}(B_{N},G) \times \Omega^{1}_{0}(B_{N},G) : \\ & |\mathcal{C}_{\mathcal{G}(\hat{\sigma},\hat{\sigma}')}(e)| \geq 2M, \text{ and } \left| \operatorname{supp} d \left(\hat{\sigma} |_{\mathcal{C}_{\mathcal{G}(\hat{\sigma},\hat{\sigma}')}(e)} \right) \right| \geq 2M' \Big\} \Big) \\ &= \sum_{\substack{E \subseteq C_{1}^{+}(B_{N}):\\ e \in E, |E| \geq M,\\ \bar{\mathcal{G}}|_{E} \text{ is connected}} \\ & \mathcal{C}_{\mathcal{G}(\hat{\sigma},\hat{\sigma}')}(e)^{+} = E, \text{ and } \left| \operatorname{supp} d \left(\hat{\sigma} |_{\mathcal{C}_{\mathcal{G}(\hat{\sigma},\hat{\sigma}')}(e)} \right) \right| \geq 2M' \Big\} \Big) \end{split}$$

$$(6.9)$$

Given $(\hat{\sigma}, \hat{\sigma}') \in \Omega^1(B_N, G) \times \Omega^1_0(B_N, G)$, if we let $\hat{\hat{\sigma}} := \hat{\sigma}|_{\mathcal{C}_{G(\hat{\sigma}, \hat{\sigma}')}(e)}$ and $\hat{\hat{\sigma}}' := \hat{\sigma}'|_{\mathcal{C}_{G(\hat{\sigma}, \hat{\sigma}')}(e)}$, then the following statements hold.

(1) By Lemma 5.8, we have $\hat{\sigma}' \in \Omega_0^1(B_N, G)$.

(2) If $|\operatorname{supp} d(\hat{\sigma}|_{\mathcal{C}_{\mathcal{G}}(\hat{\sigma},\hat{\sigma}')}(e))| \ge 2M'$, then, by definition, $|\operatorname{supp} d\hat{\sigma}| \ge 2M'$. (3) If $|\mathcal{C}_{\mathcal{G}}(\hat{\sigma},\hat{\sigma}')(e)| \ge 2$, then we have $e \in \operatorname{supp} \hat{\sigma} \cup \operatorname{supp} \hat{\sigma}'$, and thus $(\mathcal{C}_{\mathcal{G}}(\hat{\sigma},\hat{\sigma}')(e))^+ =$ $(\operatorname{supp} \hat{\hat{\sigma}} \cup \operatorname{supp} \hat{\hat{\sigma}}')^+$. Consequently, $(\mathcal{C}_{\mathcal{G}(\hat{\sigma}, \hat{\sigma}')}(e))^+ = E$ if and only if $(\operatorname{supp} \hat{\hat{\sigma}} \cup e)^+$ $\operatorname{supp} \hat{\hat{\sigma}}')^+ = E.$

As a consequence, (6.9) can be bounded from above by

$$\sum_{\substack{E \subseteq C_1^+(B_N): \\ e \in E, |E| \ge M, \\ \tilde{\mathcal{G}}|_E \text{ is connected}}} \sum_{\substack{\hat{\sigma} \in \Omega^1(B_N, G), \hat{\sigma}' \in \Omega_0^1(B_N, G): \\ (\sup \hat{\sigma} \cup \sup \hat{\sigma}')^+ = E, \\ |\sup \hat{\sigma} d\hat{\sigma}| \ge 2M'}} \mu_{N, \beta, \kappa_1} \times \mu_{N, \infty, \kappa_2} \big\{ \{(\hat{\sigma}, \hat{\sigma}') \in \Omega^1(B_N, G) \times \Omega_0^1(B_N, G): \\ \hat{\sigma}|_{\mathcal{C}_{\mathcal{G}}(\hat{\sigma}, \hat{\sigma}')}(e) = \hat{\sigma}, \text{ and } \hat{\sigma}'|_{\mathcal{C}_{\mathcal{G}}(\hat{\sigma}, \hat{\sigma}'(e)} = \hat{\sigma}'\} \big).$$

For any $\hat{\sigma}$, and $\hat{\sigma}'$ as in the sum above, by Lemma 5.2, we have

$$\begin{split} \mu_{N,\beta,\kappa_{1}} &\times \mu_{N,\infty,\kappa_{2}} \big(\{ (\hat{\sigma}, \hat{\sigma}') \in \Omega^{1}(B_{N}, G) \times \Omega^{1}_{0}(B_{N}, G) \colon \hat{\sigma} |_{\mathcal{C}_{\mathcal{G}(\hat{\sigma}, \hat{\sigma}')}(e)} \\ &= \hat{\sigma} \text{ and } \hat{\sigma}' |_{\mathcal{C}_{\mathcal{G}(\hat{\sigma}, \hat{\sigma}')}(e)} = \hat{\sigma}' \} \big) \\ &\leq \mu_{N,\beta,\kappa_{1}} \times \mu_{N,\infty,\kappa_{2}} \big(\{ (\hat{\sigma}, \hat{\sigma}') \in \Omega^{1}(B_{N}, G) \times \Omega^{1}_{0}(B_{N}, G) \colon \hat{\sigma} \leq \hat{\sigma} \text{ and } \hat{\sigma}' \leq \hat{\sigma}' \} \big) \\ &= \mu_{N,\beta,\kappa_{1}} \big(\{ \hat{\sigma} \in \Omega^{1}(B_{N}, G) \colon \hat{\sigma} \leq \hat{\sigma} \} \big) \mu_{N,\infty,\kappa_{2}} \big(\{ \hat{\sigma}' \in \Omega^{1}_{0}(B_{N}, G) \colon \hat{\sigma}' \leq \hat{\sigma}' \} \big) \\ &\leq \varphi_{\beta,\kappa_{1}}(\hat{\sigma}) \varphi_{\infty,\kappa_{2}}(\hat{\sigma}'), \end{split}$$

where the last inequality follows by applying Proposition 4.2 twice.

Taken together, the above equations thus show that

$$\mu_{N,\beta,\kappa_{1}} \times \mu_{N,\infty,\kappa_{2}} \Big(\Big\{ (\hat{\sigma}, \hat{\sigma}') \in \Omega^{1}(B_{N}, G) \times \Omega^{1}_{0}(B_{N}, G) : \\ |\mathcal{C}_{\mathcal{G}}(\hat{\sigma}, \hat{\sigma}')(e)| \geq 2M \text{ and } |\operatorname{supp} d \left(\hat{\sigma} |_{\mathcal{C}_{\mathcal{G}}(\hat{\sigma}, \hat{\sigma}')(e)} \right) \Big| \geq 2M' \Big\} \Big) \\ \leq \sum_{\substack{E \subseteq C_{1}^{+}(B_{N}): \\ e \in E, |E| \geq M, \\ \tilde{\mathcal{G}}|_{E} \text{ is connected}}} J_{\beta,\kappa_{1},\kappa_{2}}(E),$$

$$(6.10)$$

where

$$J_{\beta,\kappa_1,\kappa_2}(E) := \sum_{\substack{\hat{\hat{\sigma}} \in \Omega^1(B_N,G), \, \hat{\hat{\sigma}}' \in \Omega^1_0(B_N,G):\\(\operatorname{supp} \hat{\hat{\sigma}} \cup \operatorname{supp} \hat{\hat{\sigma}}')^* = E,\\|\operatorname{supp} d\sigma| \ge 2M'}} \varphi_{\beta,\kappa_1}(\hat{\hat{\sigma}}) \varphi_{\infty,\kappa_2}(\hat{\hat{\sigma}}').$$

Fix some set $E \subseteq C_1^+(B_N)$. Then

$$J_{\beta,\kappa_1,\kappa_2}(E) = \sum_{\substack{\hat{\hat{\sigma}} \in \Omega^1(B_N,G), \, \hat{\hat{\sigma}}' \in \Omega^1_0(B_N,G):\\(\operatorname{supp} \hat{\hat{\sigma}} \cup \operatorname{supp} \hat{\hat{\sigma}}')^+ = E,\\|\operatorname{supp} d\hat{\hat{\sigma}}| \ge 2M'}} \varphi_{\kappa_1}(\hat{\hat{\sigma}})\varphi_{\kappa_2}(\hat{\hat{\sigma}}') \prod_{p \in C_2(B_N)} \varphi_{\beta}(d\hat{\hat{\sigma}}(p)).$$

Now recall that for any $r \ge 0$ and $g \in G$, we have $\varphi_r(0) = 1$ and $\varphi_r(g) = e^{r\Re(\rho(g)-1)} \in (0, 1]$. If $g \ne 0$, then $\varphi_r(g) < 1$ and hence $\varphi_\beta(g)^2 \le \alpha_1(\beta) < 1$.

If $\hat{\sigma}$, $\hat{\sigma}'$ and *E* are as above, then we must be in one of the following three cases.

(1) If $|(\operatorname{supp} d\hat{\hat{\sigma}})^+| \ge 6$, then

$$\prod_{p \in C_2(B_N)} \varphi_\beta \left(d\hat{\hat{\sigma}}(p) \right) = \prod_{p \in C_2(B_N, G)^+} \varphi_\beta \left((d\hat{\hat{\sigma}})_p \right)^2 \le \alpha_1(\beta)^{\max(M', 6)}.$$

(2) If $|(\sup p d\hat{\sigma})^+| \in \{1, 2, 3, 4, 5\}$, by Lemma 2.8, $\hat{\sigma}$ must support a vortex with support at the boundary of B_N , and hence we must have $|E| \ge \operatorname{dist}_1(e, \partial C_1(B_N))$. At the same time, by definition, we also have $\prod_{p \in C_1(B_N,G)} \varphi_\beta(d\hat{\sigma}(p)) \le \alpha_1(\beta)$.

(3) If $|(\operatorname{supp} d\hat{\hat{\sigma}})^+| = 0$, then $\hat{\hat{\sigma}} \in \Omega_0^1(B_N, G)$. Since $|\operatorname{supp}(\hat{\hat{\sigma}})^+ \cup \operatorname{supp}(\hat{\hat{\sigma}}')^+| > 0$, it follows from Lemma 2.9 that $|E| \ge \min(M, 8, \operatorname{dist}_0(e, \partial C_1(B_N)))$. Moreover, we have $\prod_{p \in C_1(B_N, G)} \varphi_\beta(d\hat{\hat{\sigma}}(p)) = 1$.

Consequently, we have

$$J_{\beta,\kappa_{1},\kappa_{2}}(E) \leq \left(\alpha_{1}(\beta)^{\max(6,M')} + \mathbb{1}_{M' \in \{1,2,3,4,5\}, |E| \geq \text{dist}_{1}(e,\partial C_{1}(B_{N}))} \cdot \alpha_{1}(\beta) \right.$$

+ $\left.\mathbb{1}_{M'=0, |E| \geq \max(M,\min(8,\text{dist}_{0}(e,\partial C_{1}(B_{N}))))}\right)$
$$\sum_{\hat{\hat{\sigma}} \in \Omega^{1}(B_{N},G), \hat{\hat{\sigma}}' \in \Omega^{1}_{0}(B_{N},G):} \varphi_{\kappa_{1}}(\hat{\hat{\sigma}})\varphi_{\kappa_{2}}(\hat{\hat{\sigma}}').$$

$$\left.(\sup_{(\sup_{\hat{\sigma}}) \in \sup_{\hat{\sigma}})^{+}=E,} |\sup_{(\sup_{\hat{\sigma}}) \geq 2M'}$$

By dropping the condition $|\operatorname{supp} d\hat{\hat{\sigma}}| \geq 2M'$, and replacing the condition $\hat{\hat{\sigma}}' \in \Omega_0^1(B_N, G)$ with the condition that $\hat{\hat{\sigma}}' \in \Omega^1(B_N, G)$, we make the sum larger. Hence

$$\sum_{\substack{\hat{\hat{\sigma}} \in \Omega^{1}(B_{N},G), \, \hat{\hat{\sigma}}' \in \Omega^{1}_{0}(B_{N},G):\\ (\operatorname{supp} \hat{\hat{\sigma}} \cup \operatorname{supp} \hat{\hat{\sigma}}')^{*} = E,\\ |\operatorname{supp} d\hat{\hat{\sigma}}| \geq 2M'} \varphi_{\kappa_{1}}(\hat{\hat{\sigma}}) \varphi_{\kappa_{2}}(\hat{\hat{\sigma}}') \leq \sum_{\substack{\hat{\hat{\sigma}} \in \Omega^{1}(B_{N},G), \, \hat{\hat{\sigma}}' \in \Omega^{1}(B_{N},G):\\ (\operatorname{supp} \hat{\hat{\sigma}} \cup \operatorname{supp} \hat{\hat{\sigma}}')^{*} = E}} \varphi_{\kappa_{1}}(\hat{\hat{\sigma}}) \varphi_{\kappa_{2}}(\hat{\hat{\sigma}}').$$

Using Lemma 6.7, we thus obtain

$$J_{\beta,\kappa_{1},\kappa_{2}}(E) \leq \left(\alpha_{1}(\beta)^{\max(6,M')} + \mathbb{1}_{M' \in \{1,2,3,4,5\}, |E| \geq \operatorname{dist}_{1}(e,\partial C_{1}(B_{N}))} \cdot \alpha_{1}(\beta) + \mathbb{1}_{M'=0, |E| \geq \max(M,\min(8,\operatorname{dist}_{0}(e,\partial C_{1}(B_{N})))))} \right)$$

$$\cdot \left(\left(\alpha_{0}(\kappa_{1}) + \alpha_{0}(\kappa_{2}) + \alpha_{0}(\kappa_{1})\alpha_{0}(\kappa_{2})\right)^{|E|} \right).$$
(6.11)

Combining (6.10) and (6.11) and applying Lemma 6.6, we now finally obtain

$$\begin{split} & \mu_{N,\beta,\kappa} \times \mu_{N,\infty,\kappa} \Big(\Big\{ (\hat{\sigma}, \hat{\sigma}') \in \Omega^{1}(B_{N}, G) \times \Omega^{1}_{0}(B_{N}, G) : \\ & |\mathcal{C}_{\mathcal{G}(\hat{\sigma}, \hat{\sigma}')}(e)| \geq 2M \text{ and } | \text{supp } d \left(\hat{\sigma} |_{\mathcal{C}_{\mathcal{G}(\hat{\sigma}, \hat{\sigma}')}(e)} \right) | \geq 2M' \Big\} \Big) \\ & \leq \sum_{m=M}^{\infty} 18^{\max(0, 2m-3)} \Big(\alpha_{0}(\kappa_{1}) + \alpha_{0}(\kappa_{2}) + \alpha_{0}(\kappa_{1})\alpha_{0}(\kappa_{2}) \Big)^{m} \alpha_{1}(\beta)^{\max(6, M')} \\ & + \mathbbm{1}_{M' \in \{1, 2, 3, 4, 5\}} \sum_{m=\text{dist}_{1}(e, \partial C_{1}(B_{N}))}^{\infty} 18^{\max(0, 2m-3)} \Big(\alpha_{0}(\kappa_{1}) + \alpha_{0}(\kappa_{2}) + \alpha_{0}(\kappa_{1})\alpha_{0}(\kappa_{2}) \Big)^{m} \alpha_{1}(\beta) \\ & + \mathbbm{1}_{M'=0} \sum_{m=\max(M, \min(8, \text{dist}_{0}(e, \partial C_{1}(B_{N}))))}^{\infty} 18^{\max(0, 2m-3)} \Big(\alpha_{0}(\kappa_{1}) + \alpha_{0}(\kappa_{2}) + \alpha_{0}(\kappa_{1})\alpha_{0}(\kappa_{2}) \Big)^{m}. \end{split}$$

Computing the above geometric sums, we obtain (6.3) as desired.

Proof of Proposition 6.3. If $\hat{\sigma} \in \Omega^1(B_N, G)$ satisfies $d\hat{\sigma}(p) \neq 0$, then there must exist $e \in \partial p$ such that $\sigma(e) \neq 0$. For any such e, we must have $|(\mathcal{C}_{\mathcal{G}(\hat{\sigma},0)}(e))^+| \geq 1$. Moreover, since $\sigma(e) \neq 0$, for any $e' \in \partial p$ such that $\sigma(e') \neq 0$, by definition, we have $e' \in \mathcal{C}_{\mathcal{G}(\hat{\sigma},0)}(e)$. Consequently, we must have $d(\hat{\sigma}|_{\mathcal{C}_{\mathcal{G}(\hat{\sigma},0)}(e)})(p) = d\hat{\sigma}(p) \neq 0$. Using Lemma 2.8, it follows that $|(\sup d(\hat{\sigma}|_{\mathcal{C}_{\mathcal{G}}(\hat{\sigma},0)}(e)))^+| \ge 6$. Combining these observations with a union bound, it follows that

$$\mu_{N,\beta,\kappa} \left(\left\{ \hat{\sigma} \in \Omega^1(B_N, G) \colon d\hat{\sigma}(p) \neq 0 \right\} \right)$$

$$\leq \sum_{e \in \partial p} \mu_{N,\beta,\kappa} \left(\left\{ \hat{\sigma} \in \Omega^1(B_N, G) \colon |(\mathcal{C}_{\mathcal{G}(\hat{\sigma})}(e))^+| \geq 1 \text{ and } |(\operatorname{supp} d(\hat{\sigma}|_{\mathcal{C}_{\mathcal{G}(\hat{\sigma})}(e)}))^+| \geq 6 \right\} \right)$$

Applying Proposition 6.1 with $\kappa_1 = \kappa$, $\kappa_2 = \infty$, M = 1, and M' = 6, we obtain (6.3) as desired.

Proof of Proposition 6.4. Recall first that by the definition of $\mu_{N,(\infty,\kappa),(\infty,\kappa)}^{E_0}$, using Lemma 5.9, we have

$$\mu_{N,(\infty,\kappa),(\infty,\kappa)}^{E_0} \left(\left\{ (\sigma,\sigma') \in \Omega_0^1(B_N,G) \times \Omega_0^1(B_N,G) \colon e \in E_{E_0,\sigma,\sigma'} \right\} \right)$$

= $\mu_{N,\infty,\kappa} \times \mu_{N,\infty,\kappa} \left(\left\{ (\hat{\sigma},\hat{\sigma}') \in \Omega_0^1(B_N,G) \times \Omega_0^1(B_N,G) \colon e \in E_{\hat{\sigma},\hat{\sigma}'} \right\} \right).$

Next, since $\hat{\sigma}, \hat{\sigma}' \in \Omega_0^1(B_N, G)$, we have $d\hat{\sigma} = d\hat{\sigma}' = 0$. Consequently,

$$e \in E_{E_0,\hat{\sigma},\hat{\sigma}'} \Leftrightarrow e \in \mathcal{C}_{\mathcal{G}(\hat{\sigma},\hat{\sigma}')}(E_0) \Leftrightarrow E_0 \cap \mathcal{C}_{\mathcal{G}(\hat{\sigma},\hat{\sigma}')}(e) \neq \emptyset.$$

Finally, note that if $E_0 \cap \mathcal{C}_{\mathcal{G}(\hat{\sigma},\hat{\sigma}')}(e) \neq \emptyset$, then, by definition, we must have $|(\mathcal{C}_{\mathcal{G}(\hat{\sigma},\hat{\sigma}')}(e))^+| \ge \text{dist}_0(e, E_0)$.

Combining these observations, it follows that

$$\mu_{N,(\infty,\kappa),(\infty,\kappa)}^{E_0} \big(\big\{ (\sigma,\sigma') \in \Omega_0^1(B_N,G) \times \Omega_0^1(B_N,G) \colon e \in E_{E_0,\sigma,\sigma'} \big\} \big)$$

$$\leq \mu_{N,\infty,\kappa} \times \mu_{N,\infty,\kappa} \big(\big\{ (\hat{\sigma},\hat{\sigma}') \in \Omega_0^1(B_N,G) \times \Omega_0^1(B_N,G) \colon |\mathcal{C}_{\mathcal{G}(\hat{\sigma},\hat{\sigma}')}(e)|$$

$$\geq \operatorname{dist}_0(e,E_0) \big\} \big).$$

Applying Proposition 6.1 with $\kappa_1 = \kappa_2 = \kappa$, $\beta = \infty$, $M = \text{dist}_0(e, E_0)$, and M' = 0, we obtain (6.5) as desired.

Proof of Proposition 6.5. Without loss of generality, we can assume that $e \in C_1(B_N)^+$. To simplify notation, let

$$\mathcal{E} := \left\{ \sigma \in \Omega^1(B_N, G) \colon |\mathcal{C}_{\mathcal{G}(\sigma)}(E_0)| \ge 2M, \text{ and } \left| \operatorname{supp} d\left(\sigma|_{\mathcal{C}_{\mathcal{G}(\sigma)}(E_0)}\right) \right| \ge 2M' \right\}.$$

Now note that $C_{\mathcal{G}(\hat{\sigma},\hat{\sigma}')}(E_0)$ is symmetric, and that the set $C_{\mathcal{G}(\hat{\sigma},\hat{\sigma}')}(E_0) \cup \{e, -e\}$ induces a connected set in $\overline{\mathcal{G}}$. Consequently, we have

$$\mu_{N,\beta,\kappa}(\mathcal{E}) = \sum_{\substack{E \subseteq C_1^+(B_N): |E| \ge M, \\ \bar{\mathcal{G}}|_{E \cup \{e\}} \text{ is connected}}} \mu_{N,\beta,\kappa} \Big(\Big\{ \sigma \in \Omega^1(B_N,G): \mathcal{C}_{\mathcal{G}(\sigma)}(E_0)^{-1} \\ = E, \ \Big| \text{supp} \, d \big(\sigma |_{\mathcal{C}_{\mathcal{G}(\sigma)}(E_0)} \big) \Big| \ge 2M' \Big\} \Big).$$

Given $\sigma \in \Omega^1(B_N, G)$, if we let $\hat{\sigma} := \sigma|_{\mathcal{C}_{\mathcal{G}(\sigma)}(E_0)}$, then the following statements hold.

(1) If $|\operatorname{supp} d(\sigma|_{\mathcal{C}_{\mathcal{G}(\sigma,0)}(E_0)})| \ge 2M'$, then, by definition, $|\operatorname{supp} d\hat{\hat{\sigma}}| \ge 2M'$.

(2) If $d(\sigma|_{\mathcal{C}_{\mathcal{G}(\sigma,0)}(E_0)}) \neq 0$, then $E_0 \cap \operatorname{supp} \sigma \neq \emptyset$ and thus $(\mathcal{C}_{\mathcal{G}(\sigma)}(E_0))^+ = (\operatorname{supp} \hat{\hat{\sigma}})^+$.

As a consequence, the expression in the previous equation is bounded from above by

$$\sum_{\substack{E \subseteq C_1^+(B_N): |E| \ge M, \ \hat{\sigma} \in \Omega^1(B_N, G): \\ \tilde{\mathcal{G}}|_{E \cup \{e\}} \text{ is connected }}} \sum_{\substack{\hat{\sigma} \in \Omega^1(B_N, G): \\ (\operatorname{supp} \hat{\sigma})^+ = E, \\ | \operatorname{supp} d\hat{\hat{\sigma}} | \ge 2M'}} \mu_{N,\beta,\kappa} \Big(\Big\{ \sigma \in \Omega^1(B_N, G): \hat{\sigma}|_{\mathcal{C}_{\mathcal{G}}(\hat{\sigma}, 0)}(E_0) = \hat{\sigma} \Big\} \Big).$$
(6.12)

For any $\hat{\sigma}$ as in the sum above, by applying first Lemma 5.2, and then Proposition 4.2, we have

$$\mu_{N,\beta,\kappa} \left\{ \{ \hat{\sigma} \in \Omega^1(B_N, G) : \hat{\sigma}|_{\mathcal{C}_{\mathcal{G}}(\hat{\sigma}, 0)}(E_0)} = \hat{\hat{\sigma}} \} \right\} \leq \mu_{N,\beta,\kappa} \left\{ \{ \hat{\sigma} \in \Omega^1(B_N, G) : \hat{\hat{\sigma}} \leq \hat{\sigma} \} \right\}$$
$$\leq \varphi_{\beta,\kappa}(\hat{\hat{\sigma}}).$$

Taken together, the above equations show that

$$\mu_{N,\beta,\kappa}(\mathcal{E}) \leq \sum_{\substack{E \subseteq C_1^+(B_N):\\e \in E, |E| \geq M,\\ \bar{\mathcal{G}}|_E \text{ is connected}}} J_{\beta,\kappa}(E),$$
(6.13)

where

$$J_{\beta,\kappa}(E) := \sum_{\substack{\hat{\hat{\sigma}} \in \Omega^1(B_N,G):\\(\sup \hat{\hat{\sigma}})^+ = \sup p E,\\|\sup p d\hat{\hat{\sigma}}| \ge 2M'}} \varphi_{\beta,\kappa}(\hat{\hat{\sigma}}).$$

Now recall that

$$\varphi_{\beta,\kappa}(\hat{\hat{\sigma}}) = \prod_{e' \in C_1(B_N)} \varphi_{\kappa}(\hat{\hat{\sigma}}(e')) \prod_{p \in C_2(B_N)} \varphi_{\beta}(d\hat{\hat{\sigma}}(p)).$$

Also, recall that for any $r \ge 0$ and $g \in G$, we have $\varphi_r(0) = 1$ and $\varphi_r(g) = e^{r\Re(\rho(g)-1)} \in (0, 1]$. If $g \ne 0$, then $\varphi_r(g) < 1$ and hence $\varphi_\beta(g)^2 \le \alpha_1(\beta) < 1$.

If $\hat{\sigma}$ is as above, then $|\operatorname{supp} d\hat{\sigma}| \ge 2 M'$, and hence

$$\prod_{p \in C_2(B_N,G)^+} \varphi_{\beta} \left((d\hat{\hat{\sigma}})_p \right)^2 \leq \alpha_1(\beta)^{M'}.$$

Consequently, if E is as above, then

$$J_{\beta,\kappa}(E) \leq \alpha_1(\beta)^{M'} \sum_{\substack{\hat{\hat{\sigma}} \in \Omega^1(B_N,G): \\ (\operatorname{supp} \hat{\hat{\sigma}})^+ = E, \\ |\operatorname{supp} d\hat{\hat{\sigma}}| \geq 2M'}} \prod_{\substack{e' \in C_1(B_N)^+ \\ |\operatorname{supp} d\hat{\hat{\sigma}}| \geq 2M'}} \varphi_{\kappa} \left(\hat{\hat{\sigma}}(e')\right)^2.$$

By dropping the condition $|\operatorname{supp} d\hat{\hat{\sigma}}| > 2 M'$ we make the sum larger. Hence

$$J_{\beta,\kappa}(E) \le \alpha_1(\beta)^{M'} \sum_{\substack{\hat{\sigma} \in \Omega^1(B_N,G):\\(\sup p\hat{\sigma})^+ = \operatorname{supp} W}} \prod_{\substack{e' \in C_1(B_N)^+}} \varphi_{\kappa} \left(\hat{\sigma}(e')\right)^2.$$

If $\hat{\sigma} \in \Omega^1(B_N, G)$ and $e' \notin \operatorname{supp} \hat{\sigma}$, then $\varphi_{\kappa}(\hat{\sigma}(e')) = \varphi_{\kappa}(0) = 1$. Also, if $\hat{\sigma} \in \Omega^1(B_N, G)$ $\Omega^1(B_N, G)$ and $e' \in (\operatorname{supp} \hat{\hat{\sigma}})^+$, then $\hat{\hat{\sigma}}(e') \neq 0$.

Using this observation, we obtain

$$\sum_{\substack{\hat{\hat{\sigma}} \in \Omega^{1}(B_{N},G):\\(\sup p \,\hat{\hat{\sigma}}')^{+} = E\\}} \prod_{e' \in C_{1}(B_{N})^{+}} \varphi_{\kappa} \left(\hat{\hat{\sigma}}(e') \right)^{2} \leq \prod_{e' \in E} \sum_{\hat{\hat{\sigma}}'(e') \in G \smallsetminus \{0\}} \varphi_{\kappa} (\hat{\hat{\sigma}}'(e'))^{2} = \prod_{e' \in E} \alpha_{0}(\kappa)$$

$$= \alpha_{0}(\kappa)^{|E|}, \qquad (6.14)$$

We thus have

$$J_{\beta,\kappa}(E) \le \alpha_1(\beta)^{M'} \alpha_0(\kappa)^{|E|}.$$
(6.15)

Now note that, by Lemma 6.6, for any m > M, we have

$$\begin{split} \left| \left\{ E \subseteq C_1^+(B_N) \colon |E| = m, \ \bar{\mathcal{G}}|_{E \cup \{e\}} \text{ is connected} \right\} \right| \\ &= \left| \left\{ E \subseteq C_1^+(B_N) \colon |E| = m, \ e \in E, \ \bar{\mathcal{G}}|_E \text{ is connected} \right\} \right| \\ &+ \left| \left\{ E \subseteq C_1^+(B_N) \colon |E| = m+1, \ e \in E, \ \bar{\mathcal{G}}|_E \text{ is connected} \right\} \right| \\ &< 18^{2m-3} + 18^{2(m+1)-3}. \end{split}$$

Combining this with (6.13) and (6.11), we thus find that

$$\mu_{N,\beta,\kappa}(\mathcal{E}) \leq \sum_{m=M}^{\infty} (18^{2m-3} + 18^{2(m+1)-3}) \alpha_0(\kappa)^m \alpha_1(\beta)^{M'}.$$

Computing the above geometric sum, we obtain (6.3).

7. A First Version of Our Main Result

In this section, we present a first application of the coupling introduced in Sect. 5.4, by giving a first version of Theorem 10.1. This result provides an upper bound on $(L_{\nu}(\sigma, \phi))$ which is good when the probability is small that there is a cluster in $\mathcal{G}(\hat{\sigma}, \hat{\sigma}')$ which both intersects supp γ and supports a vortex. We later present a strengthening of this result in Proposition 10.18.

Proposition 7.1. Let $\beta, \kappa \geq 0$ be such that 3 holds, and let γ be a path with finite support. Then

$$\left| \left\langle L_{\gamma}(\sigma,\phi) \right\rangle_{\beta,\kappa,\infty} - \left\langle L_{\gamma}(\sigma,\phi) \right\rangle_{\infty,\kappa,\infty} \right| \le 2K_4 \left(1 + K_3 K_4 \alpha_0(\kappa) \right) |\operatorname{supp} \gamma | \alpha_0(\kappa) \alpha_1(\beta)^6,$$
(7.1)

where K_3 and K_4 are defined by (6.6).

Proof. Let $N \ge 1$ be large enough so that supp $\gamma \subseteq C_1(B_N)$ and dist₀(supp γ , $\partial C_1(B_N)$)) ≥ 8 .

Then, by definition, we have

$$\mathbb{E}_{N,\beta,\kappa} \big[L_{\gamma}(\sigma) \big] = \mathbb{E}_{N,(\beta,\kappa),(\infty,\kappa)} \big[L_{\gamma}(\sigma) \big] = \mu_{N,\beta,\kappa} \times \mu_{N,\infty,\kappa} \big[L_{\gamma}(\hat{\sigma}|_{E_{\hat{\sigma},\hat{\sigma}'}}) \\ + \hat{\sigma}'|_{C_1(B_N) \smallsetminus E_{\hat{\sigma},\hat{\sigma}'}}) \big].$$

On the event supp $\gamma \cap E_{\hat{\sigma},\hat{\sigma}'} = \emptyset$, we have

$$\begin{split} L_{\gamma}(\hat{\sigma}|_{E_{\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}'|_{C_1(B_N) \smallsetminus E_{\hat{\sigma},\hat{\sigma}'}}) &= L_{\gamma}(0 + \hat{\sigma}'|_{C_1(B_N) \smallsetminus E_{\hat{\sigma},\hat{\sigma}'}}) = L_{\gamma}(\hat{\sigma}'|_{E_{\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}'|_{C_1(B_N) \smallsetminus E_{\hat{\sigma},\hat{\sigma}'}}) \\ &= L_{\gamma}(\hat{\sigma}'). \end{split}$$

As a consequence,

$$\begin{split} & \mu_{N,\beta,\kappa} \times \mu_{N,\infty,\kappa} \Big[L_{\gamma}(\hat{\sigma}|_{E_{\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}'|_{C_{1}(B_{N}) \smallsetminus E_{\hat{\sigma},\hat{\sigma}'}}) \Big] \\ &= \mu_{N,\beta,\kappa} \times \mu_{N,\infty,\kappa} \Big[L_{\gamma}(\hat{\sigma}') \cdot \mathbb{1}_{\operatorname{supp}\gamma \cap E_{\hat{\sigma},\hat{\sigma}'}} = \emptyset \Big] \\ &+ \mu_{N,\beta,\kappa} \times \mu_{N,\infty,\kappa} \Big[L_{\gamma}(\hat{\sigma}|_{E_{\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}'|_{C_{1}(B_{N}) \smallsetminus E_{\hat{\sigma},\hat{\sigma}'}}) \cdot \mathbb{1}_{\operatorname{supp}\gamma \cap E_{\hat{\sigma},\hat{\sigma}'} \neq \emptyset} \Big] \\ &= \mu_{N,\beta,\kappa} \times \mu_{N,\infty,\kappa} \Big[L_{\gamma}(\hat{\sigma}') \Big] - \mu_{N,\beta,\kappa} \times \mu_{N,\infty,\kappa} \Big[L_{\gamma}(\hat{\sigma}') \cdot \mathbb{1}_{\operatorname{supp}\gamma \cap E_{\hat{\sigma},\hat{\sigma}'} \neq \emptyset} \Big] \\ &+ \mu_{N,\beta,\kappa} \times \mu_{N,\infty,\kappa} \Big[L_{\gamma}(\hat{\sigma}|_{E_{\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}'|_{C_{1}(B_{N}) \smallsetminus E_{\hat{\sigma},\hat{\sigma}'}}) \cdot \mathbb{1}_{\operatorname{supp}\gamma \cap E_{\hat{\sigma},\hat{\sigma}'} \neq \emptyset} \Big] \end{split}$$

Since ρ is unitary and $L_{\gamma}(\sigma) = \rho(\sigma(\gamma))$ for any $\sigma \in \Omega^1(B_N, G)$, it follows that

$$\begin{split} \left| \mathbb{E}_{N,\beta,\kappa} \big[L_{\gamma}(\sigma) \big] - \mathbb{E}_{N,\infty,\kappa} \big[L_{\gamma}(\hat{\sigma}') \big] \right| \\ &\leq 2\mu_{N,\beta,\kappa} \times \mu_{N,\infty,\kappa} \Big(\big\{ (\hat{\sigma}, \hat{\sigma}') \in \Omega^{1}(B_{N}, G) \times \Omega^{1}_{0}(B_{N}, G) \colon \operatorname{supp} \gamma \cap E_{\hat{\sigma}, \hat{\sigma}'} \neq \emptyset \big\} \Big) \\ &\leq 2 \sum_{e \in \gamma} \mu_{N,\beta,\kappa} \times \mu_{N,\infty,\kappa} \Big(\big\{ (\hat{\sigma}, \hat{\sigma}') \in \Omega^{1}(B_{N}, G) \times \Omega^{1}_{0}(B_{N}, G) \colon e \in E_{\hat{\sigma}, \hat{\sigma}'} \neq \emptyset \big\} \Big) \end{split}$$

By Lemma 5.10, if $e \in C_1(B_N)$, $\hat{\sigma} \in \Omega^1(B_N, G)$ and $\hat{\sigma}' \in \Omega^1_0(B_N, G)$, then $e \in E_{\hat{\sigma},\hat{\sigma}'}$ if and only if $d(\hat{\sigma}|_{\mathcal{C}_{\mathcal{G}(\hat{\sigma},\hat{\sigma}')}(e)}) \neq 0$. On the other hand, if $d(\hat{\sigma}|_{\mathcal{C}_{\mathcal{G}(\hat{\sigma},\hat{\sigma}')}(e)}) \neq 0$, then we must have $e \in \operatorname{supp} \hat{\sigma}|_{\mathcal{C}_{\mathcal{G}(\hat{\sigma},\hat{\sigma}')}(e)}$, implying in particular that $-e \in \mathcal{C}_{\mathcal{G}(\hat{\sigma},\hat{\sigma}')}(e)$, and hence $|\mathcal{C}_{\mathcal{G}(\hat{\sigma},\hat{\sigma}')}(e)| \geq 2$. Applying Proposition 6.1 with M = M' = 1 and $\kappa_1 = \kappa_2 = \kappa$, we thus obtain

$$\mu_{N,\beta,\kappa} \times \mu_{N,\infty,\kappa} \Big(\Big\{ (\hat{\sigma}, \hat{\sigma}') \in \Omega^1(B_N, G) \times \Omega^1_0(B_N, G) \colon e \in E_{\hat{\sigma},\hat{\sigma}'} \Big\} \Big)$$

$$\leq K_4 \Big(1 + K_3 K_4 \alpha_0(\kappa) \Big) \alpha_0(\kappa) \alpha_1(\beta)^6 + K_3 \Big(K_4 \alpha_0(\kappa) \Big)^{\operatorname{dist}_1(e,\partial C_1(B_N))} \alpha_1(\beta).$$

Combining the above equations and letting $N \to \infty$ (using Proposition 2.19 and Corollary 2.17), we obtain (7.1) as desired.

8. A Decomposition of the Coupled Spin Configuration

The main result in this section is the following proposition, which gives a decomposition of $\sigma := \hat{\sigma}|_{E_{\sigma,\sigma'}} + \hat{\sigma}'|_{C_1(B_N) \smallsetminus E_{\hat{\sigma},\hat{\sigma}'}}$ in terms of decompositions of $\hat{\sigma}$ and $\hat{\sigma}'$.

Proposition 8.1. Let $\hat{\sigma} \in \Omega^1(B_N, G)$ and $\hat{\sigma}' \in \Omega^1_0(B_N, G)$, let $\hat{\Sigma}$ be a decomposition of $\hat{\sigma}$ and $\hat{\Sigma}'$ be a decomposition of $\hat{\sigma}'$ (these are guaranteed to exist by Lemma 2.6), and define

$$\Sigma := \{ \hat{\sigma} \in \hat{\Sigma} : \operatorname{supp} \hat{\sigma} \subseteq E_{\hat{\sigma}, \hat{\sigma}'} \}$$

and

$$\Sigma' := \{ \hat{\sigma} \in \hat{\Sigma}' \colon \operatorname{supp} \hat{\sigma} \subseteq C_1(B_N) \smallsetminus E_{\hat{\sigma}, \hat{\sigma}'} \}.$$

Then $\Sigma \cup \Sigma'$ is a decomposition of $\sigma := \hat{\sigma}|_{E_{\sigma,\sigma'}} + \hat{\sigma}'|_{C_1(B_N) \smallsetminus E_{\hat{\sigma},\hat{\sigma}'}}$.

Proof. We need to show that (2.6)-(2.6) of Lemma 2.6 holds, i.e. that

- (i) If $\hat{\sigma} \in \Sigma \cup \Sigma'$, then $\hat{\sigma}$ is non-trivial and irreducible,
- (ii) If $\hat{\sigma} \in \Sigma \cup \Sigma'$, then $\hat{\sigma} \leq \sigma$,
- (iii) If $\hat{\sigma}_1, \hat{\sigma}_2 \in \Sigma \cup \Sigma'$, then $\hat{\sigma}_1$ and $\hat{\sigma}_2$ have disjoint supports,
- (iv) $\sigma = \sum_{\hat{\sigma} \in \Sigma \cup \Sigma'} \hat{\sigma}$, and
- (v) if $\hat{\sigma}_1, \hat{\sigma}_2 \in \Sigma \cup \Sigma'$, then $d\hat{\sigma}_1$ and $d\hat{\sigma}_2$ have disjoint supports,

We now show that (i)–(v) holds.

- (i) Since Σ̂ and Σ̂' are decompositions of ô and ô' respectively, (8) holds with Σ̂ ∪ Σ̂' replaced with Σ ∪ Σ'. Since Σ ∪ Σ' ⊆ Σ̂ ∪ Σ̂', the desired conclusion follows.
- (ii) Fix some $\hat{\sigma} \in \Sigma$. By the definition of Σ , we have $\operatorname{supp} \hat{\sigma} \subseteq E_{\hat{\sigma},\hat{\sigma}'}$, and hence $\hat{\sigma} = \hat{\sigma}|_{E_{\hat{\sigma},\hat{\sigma}'}}$. At the same time, since $\Sigma \subseteq \hat{\Sigma}$ and $\hat{\Sigma}$ is a decomposition of $\hat{\sigma}$, we have $\hat{\sigma} \leq \hat{\sigma}$. Finally, note that, by Lemma 5.9, we have $E_{\hat{\sigma},\hat{\sigma}'} = E_{\sigma,\hat{\sigma}'}$. By applying Lemma 5.2 twice, we obtain

$$\hat{\sigma} = \hat{\sigma}|_{E_{\hat{\sigma},\hat{\sigma}'}} \leq \hat{\sigma}|_{E_{\hat{\sigma},\hat{\sigma}'}} = \sigma|_{E_{\hat{\sigma},\hat{\sigma}'}} = \sigma|_{E_{\sigma,\hat{\sigma}'}} \leq \sigma,$$

and hence $\hat{\sigma} \leq \sigma$. Since proof in the case $\hat{\sigma} \in \Sigma'$ is analogous, we omit it here.

- (iii) Since Σ̂ is a decomposition of ô, for any distinct ô₁, ô₂ ∈ Σ ⊆ Σ̂, ô₁ and ô₂ have disjoint supports. Analogously, since Σ̂' is a decomposition of ô', for any distinct ô₁, ô₂ ∈ Σ' ⊆ Σ̂', ô₁ and ô₂ have disjoint supports. Finally, if ô₁ ∈ Σ and ô₂ ∈ Σ'', then, since supp ô₁ ⊆ E_{σ,σ'}, supp ô₂ ⊆ C₁(B_N) \ E_{σ,σ'}, and the sets E_{σ,σ'} and C₁(B_N) \ E_{σ,σ'} are disjoint, it follows that ô₁ and ô₂ have disjoint supports. This concludes the proof of (8).
- (iv) Since $\hat{\Sigma}$ is a spin decomposition of $\hat{\sigma}$, each $\hat{\sigma} \in \hat{\Sigma}$ is non-trivial and irreducible. Consequently, using Lemma 5.6, it follows that for each $\hat{\sigma} \in \hat{\Sigma}$, we have either $\sup \hat{\sigma} \subseteq E_{\hat{\sigma},\hat{\sigma}'}$ or $\sup \hat{\sigma} \subseteq C_1(B_N) \setminus E_{\hat{\sigma},\hat{\sigma}'}$, and hence

$$\hat{\sigma}|_{E_{\hat{\sigma},\hat{\sigma}'}} = \left(\sum_{\hat{\sigma}\in\hat{\Sigma}}\hat{\sigma}\right)\Big|_{E_{\hat{\sigma},\hat{\sigma}'}} = \sum_{\hat{\sigma}\in\hat{\Sigma}}\hat{\sigma}|_{E_{\hat{\sigma},\hat{\sigma}'}} = \sum_{\hat{\sigma}\in\Sigma}\hat{\sigma}.$$

Completely analogously, we find that

$$\hat{\sigma}'|_{C_1(B_N)\smallsetminus E_{\hat{\sigma},\hat{\sigma}'}} = \left(\sum_{\hat{\sigma}\in\hat{\Sigma}'}\hat{\sigma}\right)\Big|_{C_1(B_N)\smallsetminus E_{\hat{\sigma},\hat{\sigma}'}} = \sum_{\hat{\sigma}\in\hat{\Sigma}'}\hat{\sigma}|_{E_{\hat{\sigma},\hat{\sigma}'}} = \sum_{\hat{\sigma}\in\Sigma'}\hat{\sigma}.$$

Combining the previous equations and using the definition of σ , we obtain (8).

(v) If *σ̂* ∈ Σ', then, since Σ' ⊆ Σ̂' and Σ̂' is a decomposition of *σ̂*', we have *σ̂* ≤ *σ̂*'. Since *σ̂*' ∈ Ω₀¹(*B_N*, *G*), we have *dσ̂*' = 0, and hence *dσ̂* = 0. Consequently, the desired conclusion will follow if we can show that (8) holds with Σ ∪ Σ' replaced with Σ. To see that this holds, let *σ̂*₁, *σ̂*₂ ∈ Σ. Then, since Σ ⊆ Σ̂, we also have *σ̂*₁, *σ̂*₂ ∈ Σ̂. Since Σ̂ is a decomposition of *σ̂*, the 2-forms *dσ*₁ and *dσ̂*₂ must have disjoint support. This concludes the proof of (8).

9. Disturbing 1-Forms

The main purpose of this section is to introduce the following definition.

Definition 9.1. Let $\sigma \in \Omega^1(B_N, G)$, and let $\gamma \in C^1(B_N)$ be a path. If there is no path $\hat{\gamma} \in C^1(B_N)$ with $\partial \hat{\gamma} = -\partial \gamma$ and 1-form $\hat{\sigma} \in \Omega^1(B_N, G)$ such that

- (i) $d\hat{\sigma} \leq d\sigma$
- (ii) $\sigma(\hat{\gamma}) = 0$,
- (iii) $\hat{\sigma}(\gamma + \hat{\gamma}) = 0$,
- (iv) Any vortex ν in $\sigma \hat{\sigma}$ is a minimal vortex centered around an edge in $\gamma \gamma_c$ (see (3.8) for a definition of γ_c), and
- (v) If $d\sigma(p) = d\sigma(p')$ for all $p, p' \in \hat{\partial}e$, then $d\hat{\sigma}(p) = 0$ for all $p \in \hat{\partial}e$,

then we say that σ disturbs γ .

Note that if $\gamma \in C^1(B_N)$ is a generalized loop and $\sigma \in \Omega^1(B_N, G)$, then we can pick $\hat{\gamma} = 0$ in Definition 9.1, and hence, in this case, (ii) automatically holds.

The main reason for introducing the previous definition is Lemma 9.2 below. To simplify the notation in this lemma, we define

$$\gamma'[e] := (\gamma - \gamma_c)[e] \cdot \mathbb{1}(\exists p, p' \in \hat{\partial}e : d\sigma(p) \neq d\sigma(p')), \quad e \in C_1^+(B_N).$$
(9.1)

Lemma 9.2. Let $\sigma \in \Omega^1(B_N, G)$ and let $\gamma \in C^1(B_N)$ be a path. For each $e \in \gamma$, fix one plaquette $p_e \in \hat{\partial}e$.

Then, if σ does not disturb γ , we have

$$\sigma(\gamma) = \sum_{e \in (\gamma - \gamma_c) - \gamma'} d\sigma(p_e).$$

Proof. Assume that σ does not disturb γ . Then, by definition, there is $\hat{\gamma}$ and $\hat{\sigma}$ which satisfies (9.1)–(9.1) of Definition 9.1.

To simplify notation, define $\bar{\sigma} := \sigma - \hat{\sigma}$. Then

$$\begin{aligned} \sigma(\gamma) &= \sigma(\gamma) + 0 \stackrel{(9,1)}{=} \sigma(\gamma) + \sigma(\hat{\gamma}) \\ &= \sigma(\gamma + \hat{\gamma}) = (\sigma - \hat{\sigma} + \hat{\sigma})(\gamma + \hat{\gamma}) = (\sigma - \hat{\sigma})(\gamma + \hat{\gamma}) + \hat{\sigma}(\gamma + \hat{\gamma}) \\ \stackrel{(9,1)}{=} (\sigma - \hat{\sigma})(\gamma + \hat{\gamma}) + 0 = (\sigma - \hat{\sigma})(\gamma + \hat{\gamma}) = \bar{\sigma}(\gamma + \hat{\gamma}). \end{aligned}$$

Since $\partial(\gamma + \hat{\gamma}) = \partial\gamma + \partial\hat{\gamma} = \partial\gamma - \partial\gamma = 0$, $\gamma + \hat{\gamma}$ is a generalized loop. Let *B* be a cube of width $|\operatorname{supp}(\gamma + \hat{\gamma})|$ which contains $\gamma + \hat{\gamma}$. Since $\gamma + \hat{\gamma} \subseteq C^1(B_N)$, such a cube exists. Next, let *q* be an oriented surface inside *B* such that $\gamma + \hat{\gamma}$ is the boundary of *q*. The existence of such a surface is guaranteed by Lemma 2.15.

By Lemma 2.6, there is a set $\Omega \subseteq \Omega_2^0(B_N, G)$ which is a decomposition of $d\bar{\sigma}$. Fix such a set Ω , and note that, by definition, each $\omega \in \Omega$ is a vortex in $\bar{\sigma}$. Let Ω^q be the set of all $\omega \in \Omega$ with $\omega(q) \neq 0$. Then, by the discrete Stokes' theorem, we have

$$\bar{\sigma}(\gamma + \hat{\gamma}) = d\bar{\sigma}(q) = \sum_{\omega \in \Omega^q} \omega(q).$$

Now fix some $\omega \in \Omega^q$. Since $\omega(q) \neq 0$, by (9.1) and Lemma 2.12, there must exist $e := \frac{\partial}{\partial x^j} \Big|_a \in \Omega_1^+(B_N)$ and $g \in G \setminus \{0\}$ such that $\gamma[e] = 1$ and $\omega = d(g \mathbb{1}_a dx_j)$. Then, by definition, we have $\omega(p_e) = g$, and since $\omega \leq d\bar{\sigma}$ and $g \neq 0$, it follows that $d\bar{\sigma}(p_e) = \omega(p_e) = g$. Since q is an oriented surface with boundary γ , we thus have

$$\omega(q) = d(g\mathbb{1}_a \, dx_i)(q) = (g\mathbb{1}_a \, dx_i)(\gamma) = g = \omega(p_e) = d\bar{\sigma}(p_e).$$

Define

 $\gamma_5[e] := (\gamma - \gamma_c)[e] \cdot \mathbb{1} \big(\exists \omega \in \Omega^q \text{ such that } \operatorname{supp} \omega = \hat{\partial} e \cup \hat{\partial} (-e) \big), \quad e \in C_1^+(B_N).$

Then, since minimal vortices around distinct edges in $\gamma - \gamma_c$ have disjoint supports, it follows that

$$\sum_{\omega\in\Omega^q}\omega(q)=\sum_{e\in\gamma_5}d\bar{\sigma}(p_e).$$

Since, by assumption, we have $d\hat{\sigma} \leq d\sigma$, and $d\bar{\sigma} = d\sigma - d\hat{\sigma}$, it follows from Lemma 2.3 (2.3) that $d\bar{\sigma} \leq d\sigma$.

Using the definition of γ_5 , it follows that for any $e \in \gamma_5$, we have $d\bar{\sigma}(p_e) = d\sigma(p_e)$. Consequently,

$$\sum_{e \in \gamma_5} d\bar{\sigma}(p_e) = \sum_{e \in \gamma_5} d\sigma(p_e).$$

Now note that by the definition of γ' , we have

$$((\gamma - \gamma_c) - \gamma')[e] = (\gamma - \gamma_c)[e] \cdot \mathbb{1} (d\sigma(p) = d\sigma(p') \text{ for all } p, p' \in \hat{\partial}e), \\ e \in C_1^+(B_N).$$

Since $d\bar{\sigma} \leq d\sigma$, it follows that if $e \in \gamma_5$ then $e \in (\gamma - \gamma_c) - \gamma'$. Finally, we note that if $e \in (\gamma - \gamma_c) - \gamma'$, then $d\hat{\sigma}(p_e) = 0$, and hence $d\bar{\sigma}(p_e) = d\sigma(p_e) = 0$. As a consequence,

$$\sum_{e \in \gamma_5} d\sigma(p_e) = \sum_{e \in (\gamma - \gamma_c) - \gamma'} d\sigma(p_e).$$

By combining the previous equations, we obtain the desired conclusion.

10. Proof of the Main Result

In this section, we will first give a proof of the following result, which is more general that Theorem 1.1, and then show how this proof, with very small adjustments, implies Theorem 1.1.

Theorem 10.1. Let $G = \mathbb{Z}_n$ for some $n \ge 2$, let β , $\kappa \ge 0$ satisfy 3, let γ be a path, and let $\gamma_0 \in C^1(B_N)$ be any path with $\partial \gamma_0 = -\partial \gamma$. Then

$$\begin{split} \left| \left\langle L_{\gamma}(\sigma,\phi) \right\rangle_{\beta,\kappa,\infty} &- \Theta_{\beta,\kappa}(\gamma) H_{\kappa}(\gamma) \right| \\ &\leq K_6 \left(\alpha_2(\beta,\kappa) + \sqrt{\frac{\max(1,|\operatorname{supp}\gamma_c|)}{|\operatorname{supp}\gamma|}} \right)^{|\operatorname{supp}(\gamma-\gamma_c)|/(|\operatorname{supp}(\gamma-\gamma_c)|+2|\operatorname{supp}\gamma|)}, \\ &(10.1) \end{split}$$

where

$$\begin{split} \Theta_{\beta,\kappa}(\gamma) &:= \left\langle \prod_{e \in \gamma} \theta_{\beta,\kappa} \left(\sigma(e) - \phi(\partial e) \right) \right\rangle_{\infty,\kappa,\infty}, \\ H_{\kappa}(\gamma) &:= \left\langle L_{\gamma}(\sigma,\phi) \right\rangle_{\infty,\kappa,\infty}, \\ K_{6} &:= 2^{2|\sup p \gamma|/(2|\sup p \gamma|+|\sup p(\gamma-\gamma_{c})|)} \\ &\cdot \left[\mathbbm{1}(\partial \gamma \neq 0) \cdot 2K_{3}K_{4}^{8}\alpha_{0}(\kappa)^{7} \sum_{e \in \gamma} \left(18^{2} \left(2 + \alpha_{0}(\kappa) \right) \alpha_{0}(\kappa) \right)^{\max(0,\operatorname{dist}_{0}(e,\sup p \gamma_{0}) - 8)} \\ &\cdot \left(\frac{\alpha_{1}(\beta)}{\alpha_{0}(\beta)} \right)^{6} \cdot \frac{\alpha_{2}(\beta,\kappa)^{6}}{\alpha_{5}(\beta,\kappa)} \\ &+ K_{2} \cdot \frac{\alpha_{2}(\beta,\kappa)^{6}}{\alpha_{5}(\beta,\kappa)} + K_{3}K_{4}^{2}\alpha_{0}(\kappa)^{5/6} \cdot \left(\frac{\alpha_{1}(\beta)}{\alpha_{0}(\beta)} \right)^{7} \cdot \frac{\alpha_{2}(\beta,\kappa)^{6}}{\alpha_{5}(\beta,\kappa)} \\ &+ \frac{18^{4}K_{5}\alpha_{2}(\beta,\kappa)^{5}}{2} \cdot \left(\frac{\alpha_{1}(\beta)}{\alpha_{0}(\beta)} \right)^{12} \cdot \frac{\alpha_{2}(\beta,\kappa)^{6}}{\alpha_{5}(\beta,\kappa)} \\ &+ \sqrt{\frac{2K_{8}\alpha_{0}(\kappa)^{5}\alpha_{4}(\beta,\kappa) \max(\alpha_{0}(\kappa),\alpha_{1}(\beta)^{6})}{\alpha_{5}(\beta,\kappa)}} \cdot \sqrt{\left(\frac{\alpha_{1}(\beta)}{\alpha_{0}(\beta)} \right)^{6}} \cdot \sqrt{\frac{\alpha_{2}(\beta,\kappa)^{6}}{\alpha_{5}(\beta,\kappa)}} \\ &+ \sqrt{12K_{2}} \cdot \sqrt{\frac{\alpha_{2}(\beta,\kappa)^{6}}{\alpha_{5}(\beta,\kappa)}} \cdot \sqrt{\frac{\alpha_{3}(\beta,\kappa)}{\alpha_{5}(\beta,\kappa)}} + \sqrt{\frac{K_{10}\alpha_{0}(\kappa)^{8}\alpha_{4}(\beta,\kappa)}{\alpha_{5}(\beta,\kappa)}} \\ &+ \sqrt{\frac{\alpha_{3}(\beta,\kappa)}{\alpha_{5}(\beta,\kappa)}} \right]^{|\sup p(\gamma-\gamma_{c})|/(|\sup p(\gamma-\gamma_{c})|+2|\sup p\gamma|)}, \end{split}$$
(10.2)

where K_2 is given by (6.4), where K_3 and K_4 are given by (6.6), K_5 is given by (6.7), K_7 is given by (10.21), K_8 is given by (10.22), and K_{10} is given by (10.24).

Remark 10.2. Using the equations in the beginning of Sect. 10.9, together with (10.39), one easily shows that if $G = \mathbb{Z}_2$, then

$$\begin{split} K_{6} &= 2^{2|\operatorname{supp}\gamma|/(2|\operatorname{supp}\gamma|+|\operatorname{supp}(\gamma-\gamma_{c})|)} \\ &\cdot \bigg[2K_{3}K_{4}^{8}\alpha_{0}(\kappa)^{7}\sum_{e\in\gamma} \big(K_{4}\alpha_{0}(\kappa)\big)^{\max(0,\operatorname{dist}_{0}(e,\gamma_{0})-8)} \\ &+ K_{2} + K_{3}K_{4}^{2} + \sqrt{K_{7}} + \sqrt{K_{8}} + \sqrt{K_{10}} + \sqrt{12K_{2}} + 1 \bigg]^{|\operatorname{supp}(\gamma-\gamma_{c})|/(|\operatorname{supp}(\gamma-\gamma_{c})|+2|\operatorname{supp}\gamma|)}, \end{split}$$

10.1. A first application of the coupling. In this section, we split the expected value we are interested in into two parts, later corresponding to the two functions $\Theta_{\beta,\kappa}(\gamma)$ and $H_{\kappa}(\gamma)$ in Theorem 10.1. In order to do this, we first define three useful events;

$$\mathcal{E}_{1} := \left\{ (\hat{\sigma}, \hat{\sigma}') \in \Omega^{1}(B_{N}, G) \times \Omega^{1}_{0}(B_{N}, G) : \exists \text{ irreducible } \bar{\sigma} \leq \hat{\sigma}'|_{E_{\hat{\sigma}, \hat{\sigma}'}} \right.$$

$$\text{that disturbs } \gamma \right\}, \tag{10.3}$$

$$\mathcal{E}_{2} := \left\{ (\hat{\sigma}, \hat{\sigma}') \in \Omega^{1}(B_{N}, G) \times \Omega^{1}_{0}(B_{N}, G) : \exists \text{ irreducible } \bar{\sigma} \leq \hat{\sigma}|_{E_{\hat{\sigma}, \hat{\sigma}'}} \right.$$

$$\text{that disturbs } \gamma \right\}, \tag{10.4}$$

and

$$\mathcal{E}_{3} := \left\{ \sigma \in \Omega^{1}(B_{N}, G) : \exists e \in \gamma, \ \tilde{\sigma} \leq \sigma, \ \tilde{\sigma}' \leq \sigma - \tilde{\sigma} \text{ s.t. } d\tilde{\sigma}|_{\pm \operatorname{supp} \hat{\partial} e} \neq 0 \right.$$

$$\operatorname{and} d\tilde{\sigma}'|_{\pm \operatorname{supp} \hat{\partial} e} \neq 0 \right\}.$$
(10.5)

We provide upper bounds of the probabilities of these events occurring in Sect. 10.4.

Using this notation, we have the following result, which is the main result of this section.

Proposition 10.3. Let $\beta, \kappa \geq 0$, and let $\gamma \in C^1(B_N)$ be a path. For each $e \in \gamma$, let $p_e \in \hat{\partial}e$. Then

$$\begin{aligned} \left| \mathbb{E}_{N,\beta,\kappa} \big[L_{\gamma}(\sigma) \big] - \mathbb{E}_{N,\infty,\kappa} \big[L_{\gamma}(\sigma) \big] \mathbb{E}_{N,\beta,\kappa} \Big[\prod_{e \in (\gamma - \gamma_{c}) - \gamma'} \rho \big(d\sigma(p_{e}) \big) \Big] \right| \\ &\leq 2\mu_{N,(\beta,\kappa),(\infty,\kappa)}(\mathcal{E}_{1}) + 2\mu_{N,(\beta,\kappa),(\infty,\kappa)}(\mathcal{E}_{2}) + 2\mu_{N,(\beta,\kappa),(\infty,\kappa)}(\mathcal{E}_{3}). \end{aligned}$$

Proof. Let $\hat{\sigma} \in \Omega^1(B_N, G)$ and $\hat{\sigma}' \in \Omega^1_0(B_N, G)$, and let

$$\sigma := \hat{\sigma}|_{E_{\hat{\sigma},\hat{\sigma}'}} + \hat{\sigma}'|_{C_1(B_N) \smallsetminus E_{\hat{\sigma},\hat{\sigma}'}}.$$

Let $\hat{\Sigma}$ be a decomposition of $\hat{\sigma}$,

let $\hat{\Sigma}'$ be a decomposition of $\hat{\sigma}'$, and define

$$\begin{split} \Sigma &:= \{ \hat{\sigma} \in \hat{\Sigma} \colon \operatorname{supp} \hat{\sigma} \subseteq E_{\hat{\sigma}, \hat{\sigma}'} \}, \\ \Sigma' &:= \{ \hat{\sigma} \in \hat{\Sigma}' \colon \operatorname{supp} \hat{\sigma} \subseteq C_1(B_N) \smallsetminus E_{\hat{\sigma}, \hat{\sigma}'} \}, \\ \hat{\Sigma}_{bad} &:= \{ \hat{\sigma} \in \hat{\Sigma} \colon \hat{\sigma} \text{ disturbs } \gamma \}, \end{split}$$

and

$$\hat{\Sigma}_{bad}' := \{ \hat{\sigma} \in \hat{\Sigma}' : \hat{\sigma} \text{ disturbs } \gamma \}.$$

Note that these sets depend on $\hat{\sigma}$ and $\hat{\sigma}'$. Note also that if $(\hat{\Sigma}' \setminus \Sigma') \cap \hat{\Sigma}'_{bad} \neq \emptyset$, then $(\hat{\sigma}, \hat{\sigma}') \in \mathcal{E}_1$, and if $\Sigma \cap \hat{\Sigma}_{bad} = \emptyset$, then $(\hat{\sigma}, \hat{\sigma}') \in \mathcal{E}_2$. By Proposition 8.1, $\Sigma \cup \Sigma'$ is a decomposition of σ . This implies in particular that the 1-forms in $\Sigma \cup \Sigma'$ have disjoint supports, and hence

$$L_{\gamma}(\sigma) = L_{\gamma} \Big(\sum_{\hat{\sigma} \in \Sigma} \hat{\sigma} + \sum_{\hat{\sigma}' \in \Sigma'} \hat{\sigma}' \Big) = L_{\gamma} \Big(\sum_{\hat{\sigma} \in \Sigma} \hat{\sigma} \Big) L_{\gamma} \Big(\sum_{\hat{\sigma}' \in \Sigma'} \hat{\sigma}' \Big).$$
(10.6)

If $(\hat{\sigma}, \hat{\sigma}') \notin \mathcal{E}_1$, then $(\hat{\Sigma}' \setminus \Sigma') \cap \hat{\Sigma}'_{bad} = \emptyset$. Since $d\hat{\sigma} = 0$ for all $\hat{\sigma} \in \hat{\Sigma}'$, and hence, using Lemma 9.2, it follows that, on this event, we have

$$L_{\gamma}\Big(\sum_{\hat{\sigma}\in\hat{\Sigma}'\smallsetminus\Sigma'}\hat{\sigma}\Big) = \rho\Big(\Big(\sum_{\hat{\sigma}\in\hat{\Sigma}'\smallsetminus\Sigma'}\hat{\sigma}\Big)(\gamma)\Big) = \rho\Big(\sum_{\hat{\sigma}\in\hat{\Sigma}'\smallsetminus\Sigma'}\hat{\sigma}(\gamma)\Big) = \rho\Big(\sum_{\hat{\sigma}\in\hat{\Sigma}'\smallsetminus\Sigma'}0\Big) = 1,$$

and hence

$$L_{\gamma}\left(\sum_{\hat{\sigma}\in\Sigma'}\hat{\sigma}\right) = L_{\gamma}\left(\sum_{\hat{\sigma}\in\Sigma'}\hat{\sigma}\right) \cdot L_{\gamma}\left(\sum_{\hat{\sigma}\in\hat{\Sigma}'\smallsetminus\Sigma'}\hat{\sigma}\right) = L_{\gamma}\left(\sum_{\hat{\sigma}\in\Sigma'}\hat{\sigma} + \sum_{\hat{\sigma}\in\hat{\Sigma}'\smallsetminus\Sigma'}\hat{\sigma}\right) = L_{\gamma}(\hat{\sigma}').$$

In particular, this shows that

$$\mathbb{E}_{N,(\beta,\kappa),(\infty,\kappa)}\left(\left|L_{\gamma}\left(\sum_{\hat{\sigma}\in\Sigma'}\hat{\sigma}\right)-L_{\gamma}(\hat{\sigma}')\right|\right)\leq 2\mu_{N,(\beta,\kappa),(\infty,\kappa)}(\mathcal{E}_{1}).$$
(10.7)

Next, note that since the 1-forms in Σ have disjoint supports, we have

$$L_{\gamma}\left(\sum_{\hat{\sigma}\in\Sigma}\hat{\sigma}\right) = \rho\left(\left(\sum_{\hat{\sigma}\in\Sigma}\hat{\sigma}\right)(\gamma)\right) = \rho\left(\sum_{\hat{\sigma}\in\Sigma}\hat{\sigma}(\gamma)\right) = \prod_{\hat{\sigma}\in\Sigma}\rho\left(\hat{\sigma}(\gamma)\right)$$

For $\hat{\sigma} \in \Omega^1(B_N, G)$, define

$$\gamma_{\hat{\sigma}}'[e] := (\gamma - \gamma_c)[e] \cdot \mathbb{1} \big(\exists p, p' \in \hat{\partial} e \colon d\hat{\sigma}(p) \neq d\hat{\sigma}(p') \big), \quad e \in C_1^+(B_N).$$

If $(\hat{\sigma}, \hat{\sigma}') \notin \mathcal{E}_2$, then we have $\Sigma \cap \hat{\Sigma}_{bad} = \emptyset$. Consequently, for any $\hat{\hat{\sigma}} \in \Sigma$ we can apply Lemma 9.2 to obtain

$$\rho(\hat{\sigma}(\gamma)) = \prod_{e \in (\gamma - \gamma_c) - \gamma'_{\hat{\sigma}}} \rho(d\hat{\sigma}(p_e)).$$

If $\hat{\sigma} \in \hat{\Sigma} \setminus \Sigma$, then $d\hat{\sigma} = 0$. Consequently, if $(\hat{\sigma}, \hat{\sigma}') \notin \mathcal{E}_2$, then

$$\prod_{\overset{\circ}{\sigma}\in\Sigma}\rho(\overset{\circ}{\sigma}(\gamma))=\prod_{\overset{\circ}{\sigma}\in\Sigma}\prod_{e\in(\gamma-\gamma_c)-\gamma'_{\overset{\circ}{\sigma}}}\rho(d\overset{\circ}{\sigma}(p_e))=\prod_{\overset{\circ}{\sigma}\in\hat{\Sigma}}\prod_{e\in(\gamma-\gamma_c)-\gamma'_{\overset{\circ}{\sigma}}}\rho(d\overset{\circ}{\sigma}(p_e)).$$

We now make a few observations.

- If $\hat{\sigma} \in \hat{\Sigma}$ satisfies $d\hat{\sigma}(p) \neq 0$ for some $e \in (\gamma \gamma_c) \gamma'_{\hat{\sigma}}$ and $p \in \hat{\partial}e$, then, since $\hat{\Sigma}$ is a decomposition of $\hat{\sigma}$, we must have $d\hat{\sigma}'(p) = 0$ for all $\hat{\sigma}' \in \hat{\Sigma} \setminus \{\hat{\sigma}\}$.
- If $\hat{\sigma} \in \hat{\Sigma}$, then, since $\hat{\Sigma}$ is a decomposition of $\hat{\sigma}$, we must have $\hat{\sigma} \leq \hat{\sigma}$. Consequently, if $d\hat{\sigma}(p) \neq 0$ for some $p \in C_2(B_N)$, then $d\hat{\sigma}(p) = d\hat{\sigma}(p)$.
- If $\hat{\sigma}, \hat{\sigma}' \in \hat{\Sigma}$ are distinct and $e \in \gamma'_{\hat{\sigma}}$, then either $e \in \gamma'_{\hat{\sigma}'}$ or $d\hat{\sigma}(p) = 0$ for all $p \in \hat{\partial} e$.

Define

$$\gamma'''[e] := (\gamma - \gamma_c)[e] \cdot \mathbb{1} \big(\exists p, p' \in \hat{\partial} e, \, \hat{\sigma} \in \hat{\Sigma} : \, d\hat{\sigma}(p) \neq d\hat{\sigma}(p') \big), \quad e \in C_1^+(B_N).$$

Combining these observations, it follows that

$$\prod_{\hat{\sigma}\in\hat{\Sigma}}\prod_{e\in(\gamma-\gamma_c)-\gamma'_{\hat{\sigma}}}\rho\left(d\hat{\sigma}(p_e)\right) = \prod_{\hat{\sigma}\in\hat{\Sigma}}\prod_{e\in(\gamma-\gamma_c)-\gamma'_{\hat{\sigma}}:}\rho\left(d\hat{\sigma}(p_e)\right)$$
$$=\prod_{\hat{\sigma}\in\hat{\Sigma}}\prod_{e\in(\gamma-\gamma_c)-\gamma'_{\hat{\sigma}}:}\rho\left(d\hat{\sigma}(p_e)\right) = \prod_{e\in(\gamma-\gamma_c)-\gamma'''}\rho\left(d\hat{\sigma}(p_e)\right).$$
$$\stackrel{(d\hat{\sigma}(p_e)\neq 0}{=}$$

Combining the previous equations, it follows that if $(\hat{\sigma}, \hat{\sigma}') \notin \mathcal{E}_2$, we have

$$L_{\gamma}\Big(\sum_{\hat{\sigma}\in\Sigma}\hat{\sigma}\Big)=\prod_{e\in(\gamma-\gamma_c)-\gamma'''}\rho\Big(d\hat{\sigma}(p_e)\Big),$$

and hence

$$\mathbb{E}_{N,(\beta,\kappa),(\infty,\kappa)}\left(\left|L_{\gamma}\left(\sum_{\hat{\sigma}\in\Sigma}\hat{\sigma}\right)-\prod_{e\in(\gamma-\gamma_{c})-\gamma'''}\rho\left(d\hat{\sigma}(p_{e})\right)\right|\right)\leq 2\mu_{N,(\beta,\kappa),(\infty,\kappa)}(\mathcal{E}_{2}).$$
(10.8)

We now argue that if $\gamma'_{\hat{\sigma}} \neq \gamma'''$, then the event \mathcal{E}_3 must happen. To this end, first assume that $e \in \gamma'_{\hat{\sigma}}$. Then there is $p, p' \in \hat{\partial} e$ with $\hat{\sigma}(p) \neq \hat{\sigma}(p')$. Without loss of generality, we can assume that $\hat{\sigma}(p) \neq 0$. Since $\hat{\Sigma}$ is a decomposition of $\hat{\sigma}$, there is $\hat{\sigma} \in \hat{\Sigma}$ with $\hat{\sigma} \leq \hat{\sigma}$ such that $d\hat{\sigma}(p) = d\hat{\sigma}(p)$. Since $\hat{\sigma} \leq \hat{\sigma}$, we must have either $d\hat{\sigma}(p') = d\hat{\sigma}(p')$ or $d\hat{\sigma}(p') = 0$. Using the assumption that $d\hat{\sigma}(p) \neq 0$, it follows that $d\hat{\sigma}(p) \neq d\hat{\sigma}(p')$, and hence $e \in \gamma'''$.

Now, instead assume that $e \in \gamma''' - \gamma'$. Then, since $e \in \gamma'''$, there must exist $p, p' \in \hat{\partial} e$ and $\hat{\sigma} \in \hat{\Sigma}$ such that $d\hat{\sigma}(p) \neq d\hat{\sigma}(p')$. Without loss of generality, we can assume that $d\hat{\sigma}(p) \neq 0$. Since $\hat{\sigma} \in \hat{\Sigma}$, we have $\hat{\sigma} \leq \hat{\sigma}$, and hence, since $d\hat{\sigma}(p) \neq 0$, it follows that $d\hat{\sigma}(p) = d\hat{\sigma}(p) \neq 0$.

Since $\hat{\sigma} \leq \hat{\sigma}$, we must have either $\hat{\sigma}(p') = \hat{\sigma}(p')$ or $\hat{\sigma}(p') = 0$. Since $e \in \gamma'$, we have $d\hat{\sigma}(p) = d\hat{\sigma}(p')$, and hence, since $d\hat{\sigma}(p) = d\hat{\sigma}(p)$ and $d\hat{\sigma}(p) \neq d\hat{\sigma}(p')$, we conclude that $d\hat{\sigma}(p') = 0$.

Since $d\hat{\sigma}(p') \neq 0$ and $d\hat{\sigma}(p') = 0$, there must exist $\hat{\sigma}' \in \hat{\Sigma} \setminus \{\hat{\sigma}\}$ such that $d\hat{\sigma}'(p') = d\hat{\sigma}(p') \neq 0$.

Consequently,

$$\mathbb{E}_{N,(\beta,\kappa),(\infty,\kappa)}\left(\left|\prod_{e\in(\gamma-\gamma_c)-\gamma'''}\rho\left(d\hat{\sigma}(p_e)\right)-\prod_{e\in(\gamma-\gamma_c)-\gamma'}\rho\left(d\hat{\sigma}(p_e)\right)\right|\right) \le 2\mu_{N,\beta,\kappa}(\mathcal{E}_3).$$
(10.9)

Combining (10.6), (10.7), (10.8), and 10.9 we obtain

$$\begin{split} \left| \mathbb{E}_{N,\beta,\kappa} \left[L_{\gamma}(\sigma) \right] - \mathbb{E}_{N,\infty,\kappa} \left[L_{\gamma}(\sigma) \right] \mathbb{E}_{N,\beta,\kappa} \left[\prod_{e \in (\gamma - \gamma_{c}) - \gamma'} \rho(d\sigma(p_{e})) \right] \right| \\ &= \left| \mathbb{E}_{N,(\beta,\kappa),(\infty,\kappa)} \left[L_{\gamma} \left(\sum_{\hat{\sigma}' \in \Sigma'} \hat{\sigma}' \right) L_{\gamma} \left(\sum_{\hat{\sigma} \in \Sigma} \hat{\sigma} \right) - L_{\gamma}(\hat{\sigma}') \prod_{e \in (\gamma - \gamma_{c}) - \gamma''} \rho(d\hat{\sigma}(p_{e})) \right] \right| \\ &\leq \mathbb{E}_{N,(\beta,\kappa),(\infty,\kappa)} \left[\left| L_{\gamma} \left(\sum_{\hat{\sigma}' \in \Sigma'} \hat{\sigma}' \right) - L_{\gamma}(\hat{\sigma}') \right| \right] \\ &\leq \mathbb{E}_{N,(\beta,\kappa),(\infty,\kappa)} \left[\left| L_{\gamma} \left(\sum_{\hat{\sigma}' \in \Sigma'} \hat{\sigma}' \right) - L_{\gamma}(\hat{\sigma}') \right| \right] \\ &+ \mathbb{E}_{N,(\beta,\kappa),(\infty,\kappa)} \left[\left| L_{\gamma} \left(\sum_{\hat{\sigma} \in \Sigma} \hat{\sigma} \right) - \prod_{e \in (\gamma - \gamma_{c}) - \gamma'''} \rho(d\hat{\sigma}(p_{e})) \right| \right] \\ &+ \mathbb{E}_{N,(\beta,\kappa),(\infty,\kappa)} \left[\left| \prod_{e \in (\gamma - \gamma_{c}) - \gamma'''} \rho(d\hat{\sigma}(p_{e})) - \prod_{e \in (\gamma - \gamma_{c}) - \gamma'} \rho(d\hat{\sigma}(p_{e})) \right| \right] \\ &\leq 2\mu_{N,(\beta,\kappa),(\infty,\kappa)} (\mathcal{E}_{1}) + 2\mu_{N,(\beta,\kappa),(\infty,\kappa)} (\mathcal{E}_{2}) + 2\mu_{N,\beta,\kappa} (\mathcal{E}_{3}). \end{split}$$

This concludes the proof.

10.2. A resampling trick. Recall that given a path γ and $\sigma \in \Omega^1(B_N, G)$, we have let

$$\gamma'[e] = (\gamma - \gamma_c)[e] \cdot \mathbb{1}(\exists p, p' \in \hat{\partial}e : d\sigma(p) \neq d\sigma(p')), \quad e \in C_1(B_N)$$

In this section, we describe a resampling trick, first introduced (in a different setting) in [9].

Proposition 10.4 [Proposition 10.1 in [18]]. Let $\beta, \kappa \ge 0$, and let $\gamma \in C^1(B_N)$ be a path such that $\operatorname{dist}_0(\gamma, \partial C_1(B_N)) \ge 8$. For each $e \in \gamma$, fix one plaquette $p_e \in \hat{\partial}e$. Then

$$\mathbb{E}_{N,\beta,\kappa}\left[\rho\left(\sum_{e\in(\gamma-\gamma_c)-\gamma'}d\sigma(p_e)\right)\right] = \mathbb{E}_{N,\beta,\kappa}\left[\prod_{e\in(\gamma-\gamma_c)-\gamma'}\theta_{\beta,\kappa}\left(\sigma(e) - d\sigma(p_e)\right)\right].$$
(10.10)

For a proof of Proposition 10.4, we refer the reader to [18, Proposition 10.1]. In [18, Proposition 10.1], γ is assumed to be a generalized loop rather than a path as in Proposition 10.4. However, since the proofs in the two cases are identical, we do not include a proof here.

10.3. A second application of the coupling. In this section, we take the next step towards the proof of Theorem 10.1, by giving an upper bound on the distance between the right hand side of (10.10) and $\Theta'_{N,\beta,\kappa}(\gamma)$. To this end, using the notation of Sect. 5, we now introduce a few additional useful events which will be used to express the upper bound in the Proposition 10.5 below.

Given an edge $e \in C_1(B_N)$, let

$$\mathcal{E}_{4}(e) := \left\{ (\sigma, \sigma') \in \Omega^{1}(B_{N}, G) \times \Omega^{1}_{0}(B_{N}, G) : e \in E_{\sigma, \sigma'} \text{ and } \sigma'(e) \neq 0 \right\}, \quad (10.11)$$

$$\mathcal{E}_{5}(e) := \left\{ (\sigma, \sigma') \in \Omega^{1}(B_{N}, G) \times \Omega^{1}_{0}(B_{N}, G) : \exists e' \in E_{\sigma, \sigma'} \text{ s.t. } \hat{\partial}e' \cap \hat{\partial}e \neq \emptyset \right\}$$

and
$$\exists g \in G \setminus \{0\}$$
 s.t. $\sigma(e) - d\sigma(p) = g \forall p \in \hat{\partial}e\},$ (10.12)

$$\mathcal{E}_6(e) := \{ \sigma' \in \Omega_0^1(B_N, G) : \, \sigma'(e) \neq 0 \},$$
(10.13)

$$\mathcal{E}_{7}(e) := \{ (\sigma, \sigma') \in \Omega^{1}(B_{N}, G) \times \Omega^{1}_{0}(B_{N}, G) : \exists p, p' \in \hat{\partial} e \text{ s.t. } d\sigma(p) \neq d\sigma(p') \}.$$
(10.14)

Proposition 10.5. Let $\beta, \kappa \geq 0$, and let $\gamma \in C^1(B_N)$ be a path such that for all $e \in \gamma$, $p \in \hat{\partial}e$ and $p' \in \partial C_2(B_N)$, we have $\operatorname{supp} \partial p \cap \operatorname{supp} \hat{\partial}p' = \emptyset$.

Then

$$\begin{aligned} \left| \mathbb{E}_{N,\beta,\kappa} \left[\prod_{e \in (\gamma - \gamma_c) - \gamma'} \theta_{\beta,\kappa} \left(\sigma(e) - d\sigma(p_e) \right) \right] - \Theta_{N,\beta,\kappa}(\gamma) \right| \\ &\leq 2 \sqrt{2\alpha_4(\beta,\kappa)} \sum_{e \in \gamma} \mu_{N,(\beta,\kappa),(\infty,\kappa)} \left(\mathcal{E}_5(e) \right) + 4 \sqrt{2\alpha_4(\beta,\kappa)} \sum_{e \in \gamma} \mu_{N,(\beta,\kappa),(\infty,\kappa)} \left(\mathcal{E}_4(e) \right) \\ &+ 2 \sqrt{2\alpha_4(\beta,\kappa)} \sum_{e \in \gamma_c} \mu_{N,\infty,\kappa} \left(\mathcal{E}_6(e) \right) + 2 \sqrt{2|\operatorname{supp} \gamma_c|\alpha_3(\beta,\kappa)} \\ &+ 2 \sqrt{2\alpha_3(\beta,\kappa)} \sum_{e \in \gamma} \mu_{N,(\beta,\kappa),(\infty,\kappa)} \left(\mathcal{E}_7(e) \right). \end{aligned}$$
(10.15)

For the proof of Proposition 10.5 we need a two lemmas from [18], which we now recall.

Lemma 10.6 [Lemma 11.2 in [18]]. Assume that $z_1, z_2, z'_1, z'_2 \in \mathbb{C}$ are such that $|z_1|, |z_2|, |z'_1|, |z'_2| \leq 1$. Then

$$|z_1z_2 - z_1'z_2'| \le |z_1 - z_1'| + |z_2 - z_2'|.$$

Lemma 10.7 [Lemma 11.3 in [18]]. Let a, b > 0. Assume that $A \subseteq C_1(B_N)$ is a random set with $\mathbb{E}[|A|] \leq a$, and that

(i) $X_e \in \mathbb{C}$ and $|X_e| \le 1$ for all $e \in C_1(B_N)$, and (ii) there exists a $c \in [-1, 1]$ such that $|X_e - c| \le b$ for all $e \in C_1(B_N)$.

Then

$$\mathbb{E}\left[\left|\prod_{e\in A} c - \prod_{e\in A} X_e\right|\right] \le 2\sqrt{2ab}.$$

Proof of Proposition 10.5. Recall the coupling $(\sigma, \sigma') \sim \mu_{N,(\beta,\kappa),(\infty,\kappa)}$ between $\sigma \sim \mu_{N,\beta,\kappa}$ and $\sigma' \sim \mu_{N,\infty,\kappa}$ described in Sect. 5.4, and the set $E_{\sigma,\sigma'}$ defined in (5.16). Since $\mu_{N,(\beta,\kappa),(\infty,\kappa)}$ is a coupling of $\mu_{N,\beta,\kappa}$ and $\mu_{N,\infty,\kappa}$, we have

$$\mathbb{E}_{N,\beta,\kappa} \Big[\prod_{e \in (\gamma - \gamma_c) - \gamma'} \theta_{\beta,\kappa} \big(\sigma(e) - d\sigma(p_e) \big) \Big] - \mathbb{E}_{N,\infty,\kappa} \Big[\prod_{e \in \gamma} \theta_{\beta,\kappa} \big(\sigma(e) \big) \Big] \\= \mathbb{E}_{N,(\beta,\kappa),(\infty,\kappa)} \Big[\prod_{e \in (\gamma - \gamma_c) - \gamma'} \theta_{\beta,\kappa} \big(\sigma(e) - d\sigma(p_e) \big) - \prod_{e \in \gamma} \theta_{\beta,\kappa} \big(\sigma'(e) \big) \Big].$$

Given $(\sigma, \sigma') \in \Omega^1(B_N, G) \times \Omega^1_0(B_N, G)$, define

$$\gamma''[e] := (\gamma - \gamma_c)[e] \cdot \mathbb{1} \big(\exists e' \in E_{\sigma,\sigma'} \colon \hat{\partial} e' \cap \hat{\partial} e \neq \emptyset \big), \quad e \in C_1(B_N)^+.$$

In other words, γ' is the indicator function for all edges is $\gamma - \gamma_c$ that is adjacent to some edge in $E_{\sigma,\sigma'}$. By Lemma 5.21, if $e \in (\gamma - \gamma_c) - \gamma''$, then $\sigma(e') = \sigma'(e')$ whenever $e' \in \partial p$ for some $p \in \hat{\partial} e$, and hence $d\sigma(p_e) = d\sigma'(p_e) = 0$. In particular, this implies that if $e \in (\gamma - \gamma_c) - \gamma''$, then

$$\sigma(e) - d\sigma(p_e) = \sigma'(e) - d\sigma'(p_e) = \sigma'(e) - 0 = \sigma'(e).$$
(10.16)

By the definition of γ' , if $e' \in \gamma'$ then there exists $p' \in \hat{\partial} e'$ and $e'' \in \partial p'$ such that $e'' \in \{e''' \in \sup p \sigma : d\sigma|_{\pm \sup p \hat{\partial} e'''} \neq 0\} \subseteq E_{\sigma,\sigma'}$. Consequently, there is $e'' \in E_{\sigma,\sigma'}$ such that $\hat{\partial} e'' \cap \hat{\partial} e \neq \emptyset$.

Hence, if $e \in \gamma'$ then $e \in \gamma''$, and it follows that

$$\prod_{e \in (\gamma - \gamma_c) - \gamma'} \theta_{\beta,\kappa} (\sigma(e) - d\sigma(p_e)) = \prod_{e \in (\gamma - \gamma_c) - \gamma''} \theta_{\beta,\kappa} (\sigma(e) - d\sigma(p_e))$$
$$\prod_{e \in \gamma'' - \gamma'} \theta_{\beta,\kappa} (\sigma(e) - d\sigma(p_e))$$
$$\stackrel{(10.16)}{=} \prod_{e \in (\gamma - \gamma_c) - \gamma'} \theta_{\beta,\kappa} (\sigma'(e)) \prod_{e \in \gamma'' - \gamma'} \theta_{\beta,\kappa} (\sigma(e) - d\sigma(p_e)).$$

Consequently, using Lemma 5.21, we have

$$\begin{split} &\prod_{e \in (\gamma - \gamma_c) - \gamma'} \theta_{\beta,\kappa} \left(\sigma(e) - d\sigma(p_e) \right) - \prod_{e \in \gamma} \theta_{\beta,\kappa} \left(\sigma'(e) \right) \\ &= \prod_{e \in (\gamma - \gamma_c) - \gamma''} \theta_{\beta,\kappa} \left(\sigma(e) \right) \prod_{e \in \gamma'' - \gamma'} \theta_{\beta,\kappa} \left(\sigma(e) - d\sigma(p_e) \right) \\ &- \prod_{e \in (\gamma - \gamma_c) - \gamma''} \theta_{\beta,\kappa} \left(\sigma'(e) \right) \prod_{e \in \gamma'' - \gamma'} \theta_{\beta,\kappa} \left(\sigma'(e) \right) \prod_{e \in \gamma_c + \gamma'} \theta_{\beta,\kappa} \left(\sigma'(e) \right) \\ &= \prod_{e \in (\gamma - \gamma_c) - \gamma''} \theta_{\beta,\kappa} \left(\sigma'(e) \right) \left(\prod_{e \in \gamma'' - \gamma'} \theta_{\beta,\kappa} \left(\sigma(e) - d\sigma(p_e) \right) - \prod_{e \in \gamma'' - \gamma'} \theta_{\beta,\kappa} \left(\sigma'(e) \right) \right) \\ &+ \prod_{e \in (\gamma - \gamma_c) - \gamma'} \theta_{\beta,\kappa} \left(\sigma'(e) \right) \left(1 - \prod_{e \in \gamma_c} \theta_{\beta,\kappa} \left(\sigma'(e) \right) \right) \end{split}$$

$$+\prod_{e\in(\gamma-\gamma_c)-\gamma'}\theta_{\beta,\kappa}\big(\sigma'(e)\big)\prod_{e\in\gamma_c}\theta_{\beta,\kappa}\big(\sigma'(e)\big)\Big(1-\prod_{e\in\gamma'}\theta_{\beta,\kappa}\big(\sigma'(e)\big)\Big).$$

Combining the above equations, we find that

$$\begin{split} \mathbb{E}_{N,\beta,\kappa} \bigg[\prod_{e \in (\gamma - \gamma_c) - \gamma'} \theta_{\beta,\kappa} \big(\sigma(e) - d\sigma(p_e) \big) \bigg] - \mathbb{E}_{N,\infty,\kappa} \bigg[\prod_{e \in \gamma} \theta_{\beta,\kappa} \big(\sigma(e) \big) \bigg] \\ &= \mathbb{E}_{N,(\beta,\kappa),(\infty,\kappa)} \bigg[\prod_{e \in (\gamma - \gamma_c) - \gamma''} \theta_{\beta,\kappa} (\sigma'(e)) \bigg(\prod_{e \in \gamma'' - \gamma'} \theta_{\beta,\kappa} \big(\sigma(e) - d\sigma(p_e) \big) \\ &- \prod_{e \in \gamma'' - \gamma'} \theta_{\beta,\kappa} (\sigma'(e)) \bigg) \bigg] \\ &+ \mathbb{E}_{N,(\beta,\kappa),(\infty,\kappa)} \bigg[\prod_{e \in (\gamma - \gamma_c) - \gamma'} \theta_{\beta,\kappa} (\sigma'(e)) \Big(1 - \prod_{e \in \gamma_c} \theta_{\beta,\kappa} (\sigma'(e)) \Big) \bigg] \\ &+ \mathbb{E}_{N,(\beta,\kappa),(\infty,\kappa)} \bigg[\prod_{e \in (\gamma - \gamma_c) - \gamma'} \theta_{\beta,\kappa} (\sigma'_e) \prod_{e \in \gamma_c} \theta_{\beta,\kappa} (\sigma'(e)) \Big(1 - \prod_{e \in \gamma'} \theta_{\beta,\kappa} (\sigma'(e)) \Big) \bigg] . \end{split}$$

Now note that

$$\begin{split} &\prod_{e \in \gamma'' - \gamma'} \theta_{\beta,\kappa} \left(\sigma(e) - d\sigma(p_e) \right) - \prod_{e \in \gamma'' - \gamma'} \theta_{\beta,\kappa}(\sigma'(e)) \\ &= \prod_{\substack{e \in \gamma'' - \gamma' : \\ \sigma(e) - d\sigma(p_e) = 0}} \theta_{\beta,\kappa}(0) \left(\prod_{\substack{e \in \gamma'' - \gamma' : \\ \sigma(e) - d\sigma(p_e) \neq 0}} \theta_{\beta,\kappa}(\sigma(e) - d\sigma(p_e)) - \prod_{\substack{e \in \gamma'' - \gamma' : \\ \sigma(e) - d\sigma(p_e) \neq 0}} \theta_{\beta,\kappa}(0) \right) \\ &+ \prod_{\substack{e \in \gamma'' - \gamma' : \\ \sigma'(e) = 0}} \theta_{\beta,\kappa}(0) \left(\prod_{\substack{e \in \gamma'' - \gamma' : \\ \sigma'(e) \neq 0}} \theta_{\beta,\kappa}(\sigma'(e)) \right), \end{split}$$

and similarly, that

$$1 - \prod_{e \in \gamma_{c}} \theta_{\beta,\kappa}(\sigma'(e)) = \left(1 - \theta_{\beta,\kappa}(0)^{|\operatorname{supp}\gamma_{c}|}\right) + \prod_{\substack{e \in \gamma_{c}:\\\sigma'(e)=0}} \theta_{\beta,\kappa}(0) \left(\prod_{\substack{e \in \gamma_{c}:\\\sigma'(e)\neq 0}} \theta_{\beta,\kappa}(0) - \prod_{\substack{e \in \gamma_{c}:\\\sigma'(e)\neq 0}} \theta_{\beta,\kappa}(\sigma'(e))\right)$$

and

$$1 - \prod_{e \in \gamma'} \theta_{\beta,\kappa}(\sigma'(e)) = \left(1 - \theta_{\beta,\kappa}(0)^{|\operatorname{supp}\gamma'|}\right) + \prod_{e \in \gamma'\sigma'(e)=0} \theta_{\beta,\kappa}(0) \left(\prod_{e \in \gamma'\sigma'(e)\neq 0} \theta_{\beta,\kappa}(0) - \prod_{e \in \gamma'\sigma'(e)\neq 0} \theta_{\beta,\kappa}(\sigma'(e))\right).$$

Consequently, by applying the triangle inequality and noting that, since ρ is unitary, we have $|\theta_{\beta,\kappa}(g)| \leq 1$ for all $g \in G$, we obtain

$$\begin{split} \mathbb{E}_{N,\beta,\kappa} \bigg[\prod_{e \in (\gamma - \gamma_{c}) - \gamma'} \theta_{\beta,\kappa} \big(\sigma(e) - d\sigma(p_{e}) \big) \big] - \mathbb{E}_{N,\infty,\kappa} \bigg[\prod_{e \in \gamma} \theta_{\beta,\kappa} \big(\sigma(e) \big) \bigg] \bigg| \\ &\leq \mathbb{E}_{N,(\beta,\kappa),(\infty,\kappa)} \bigg[\bigg| \prod_{\substack{e \in \gamma'' - \gamma': \\ \sigma(e) - d\sigma(p_{e}) \neq 0}} \theta_{\beta,\kappa} \big(\sigma(e) - d\sigma(p_{e}) \big) - \prod_{\substack{e \in \gamma'' - \gamma': \\ \sigma(e) - d\sigma(p_{e}) \neq 0}} \theta_{\beta,\kappa}(0) \bigg| \bigg] \\ &+ \mathbb{E}_{N,(\beta,\kappa),(\infty,\kappa)} \bigg[\bigg| \prod_{\substack{e \in \gamma'' - \gamma': \\ \sigma'(e) \neq 0}} \theta_{\beta,\kappa}(0) - \prod_{\substack{e \in \gamma'' - \gamma': \\ \sigma'(e) \neq 0}} \theta_{\beta,\kappa} \big(\sigma'(e) \big) \bigg| \bigg] \\ &+ \mathbb{E}_{N,(\beta,\kappa),(\infty,\kappa)} \bigg[\bigg| \prod_{\substack{e \in \gamma': \\ \sigma'(e) \neq 0}} \theta_{\beta,\kappa}(0) - \prod_{\substack{e \in \gamma': \\ \sigma'(e) \neq 0}} \theta_{\beta,\kappa} \big(\sigma'(e) \big) \bigg| \bigg] + \big(1 - \theta_{\beta,\kappa}(0)^{|\operatorname{supp}\gamma'|} \big) \\ &+ \mathbb{E}_{N,(\beta,\kappa),(\infty,\kappa)} \bigg[\bigg| \prod_{\substack{e \in \gamma': \\ \sigma'(e) \neq 0}} \theta_{\beta,\kappa}(0) - \prod_{\substack{e \in \gamma': \\ \sigma'(e) \neq 0}} \theta_{\beta,\kappa} \big(\sigma'(e) \big) \bigg| \bigg] + \big(1 - \theta_{\beta,\kappa}(0)^{|\operatorname{supp}\gamma'|} \big). \end{split}$$

We now use Lemma 10.7 to obtain upper bounds for each of the terms on the righthand side of the previous equation.

Claim 10.8.

$$\mathbb{E}_{N,(\beta,\kappa),(\infty,\kappa)}\Big[\big|\{e\in\gamma''\colon\sigma'(e)\neq0\}\big|\Big]\leq\sum_{e\in\gamma}\mu_{N,(\beta,\kappa),(\infty,\kappa)}\big(\mathcal{E}_4(e)\big).$$
 (10.18)

Proof of claim. For any $e \in \gamma''$, by definition, there is at least one $e' \in E_{\sigma,\sigma'}$ such that $\hat{\partial}e' \cap \hat{\partial}e \neq \emptyset$. Consequently, by the definition of $E_{\sigma,\sigma'}$, if $\sigma'(e) \neq 0$, we $e \in E_{\sigma,\sigma'}$.

From this it follows that

$$\begin{split} \mathbb{E}_{N,(\beta,\kappa),(\infty,\kappa)} \Big[\big| \{e \in \gamma'' : \sigma'(e) \neq 0\} \big| \Big] \\ &= \sum_{e \in \gamma''} \mu_{N,(\beta,\kappa),(\infty,\kappa)} \Big(\{(\sigma,\sigma') \in \Omega^1(B_N,G) \times \Omega^1_0(B_N,G) : \sigma'(e) \neq 0\} \Big) \\ &= \sum_{e \in \gamma''} \mu_{N,(\beta,\kappa),(\infty,\kappa)} \Big(\{(\sigma,\sigma') \in \Omega^1(B_N,G) \times \Omega^1_0(B_N,G) : e \in E_{\sigma,\sigma'} \\ &\text{and } \sigma'(e) \neq 0\} \Big) \\ &= \sum_{e \in \gamma''} \mu_{N,(\beta,\kappa),(\infty,\kappa)} \Big(\mathcal{E}_4(e) \Big) \leq \sum_{e \in \gamma} \mu_{N,(\beta,\kappa),(\infty,\kappa)} \Big(\mathcal{E}_4(e) \Big), \end{split}$$

which is the desired conclusion.

Claim 10.9.

$$\mathbb{E}_{N,(\beta,\kappa),(\infty,\kappa)}\Big[\big|\{e \in \gamma'' - \gamma' \colon \sigma(e) - d\sigma(p_e) \neq 0\}\big|\Big] \le \sum_{e \in \gamma} \mu_{N,(\beta,\kappa),(\infty,\kappa)}\big(\mathcal{E}_5(e)\big).$$
(10.19)

Proof of claim. Note fist that by definition, for any $e \in \gamma'' - \gamma'$, we have

$$\left\{(\sigma,\sigma')\in\Omega^1(B_N,G)\times\Omega^1_0(B_N,G)\colon\sigma(e)-d\sigma(p_e)\neq0\right\}=\mathcal{E}_5(e).$$

Consequently,

$$\begin{split} \mathbb{E}_{N,(\beta,\kappa),(\infty,\kappa)} \Big[\big| \{ e \in \gamma'' - \gamma' : \sigma(e) - d\sigma(p_e) \neq 0 \} \big| \Big] \\ &= \sum_{e \in \gamma'' - \gamma'} \mu_{N,(\beta,\kappa),(\infty,\kappa)} \Big(\{ (\sigma, \sigma') \in \Omega^1(B_N, G) \times \Omega^1_0(B_N, G) : \\ \sigma(e) - d\sigma(p_e) \neq 0 \} \Big) \\ &= \sum_{e \in \gamma'' - \gamma'} \mu_{N,(\beta,\kappa),(\infty,\kappa)} \Big(\mathcal{E}_5^4 \Big) \leq \sum_{e \in \gamma} \mu_{N,(\beta,\kappa),(\infty,\kappa)} \big(\mathcal{E}_5(e) \big), \end{split}$$

which is the desired conclusion.

Next, by definition, we have

$$\mathbb{E}_{N,(\beta,\kappa),(\infty,\kappa)}\Big[\big|\{e\in\gamma_c\colon\sigma'(e)\neq0\}\big|\Big]=\sum_{e\in\gamma_c}\mu_{N,\infty,\kappa}\big(\mathcal{E}_6(e)\big)$$

and

$$\begin{split} \mathbb{E}_{N,\beta,\kappa} \big[|\operatorname{supp} \gamma'| \big] \\ &= \sum_{e \in \gamma \smallsetminus \gamma_c} \mu_{N,(\beta,\kappa),(\infty,\kappa)} \Big(\big\{ (\sigma,\sigma') \in \Omega^1(B_N,G) \times \Omega^1_0(B_N,G) \colon \exists p, p' \in \hat{\partial} e \\ \text{s.t.} \ d\sigma(p) \neq d\sigma(p') \big\} \Big) \\ &= \sum_{e \in \gamma \smallsetminus \gamma_c} \mu_{N,(\beta,\kappa),(\infty,\kappa)} \big(\mathcal{E}_7(e) \big) \le \sum_{e \in \gamma} \mu_{N,(\beta,\kappa),(\infty,\kappa)} \big(\mathcal{E}_7(e) \big). \end{split}$$

Finally, since for any $e \in \gamma'$ we also have $e \in \gamma''$, we have

$$\mathbb{E}_{N,(\beta,\kappa),(\infty,\kappa)}\Big[\big|\{e\in\gamma'\colon\sigma'(e)\neq0\}\big|\Big]\leq\mathbb{E}_{N,(\beta,\kappa),(\infty,\kappa)}\Big[\big|\{e\in\gamma''\colon\sigma'(e)\neq0\}\big|\Big],$$

the right-hand side of which we have given an upper bound for in (10.18),

Applying Lemma 10.7 to the terms in (10.17), we thus obtain

$$\begin{aligned} \left| \mathbb{E}_{N,\beta,\kappa} \Big[\prod_{e \in (\gamma - \gamma_c) - \gamma'} \theta_{\beta,\kappa} \big(\sigma(e) - d\sigma(p_e) \big) \Big] - \mathbb{E}_{N,\infty,\kappa} \Big[\prod_{e \in \gamma} \theta_{\beta,\kappa} \big(\sigma(e) \big) \Big] \right| \\ &\leq 2 \sqrt{2 \max_{g \in G} \left| \theta_{\beta,\kappa}(g) - \theta_{\beta,\kappa}(0) \right| \sum_{e \in \gamma} \mu_{N,(\beta,\kappa),(\infty,\kappa)} \big(\mathcal{E}_{5}(e) \big)} \\ &+ 4 \sqrt{2 \max_{g \in G} \left| \theta_{\beta,\kappa}(g) - \theta_{\beta,\kappa}(0) \right| \sum_{e \in \gamma} \mu_{N,(\beta,\kappa),(\infty,\kappa)} \big(\mathcal{E}_{4}(e) \big)} \\ &+ 2 \sqrt{2 \max_{g \in G} \left| \theta_{\beta,\kappa}(g) - \theta_{\beta,\kappa}(0) \right| \sum_{e \in \gamma_c} \mu_{N,\infty,\kappa} \big(\mathcal{E}_{6}(e) \big)} \\ &+ 2 \sqrt{2 \left| \sup \gamma_c \right| \left| 1 - \theta_{\beta,\kappa}(0) \right|} + 2 \sqrt{2 \left| 1 - \theta_{\beta,\kappa}(0) \right| \sum_{e \in \gamma} \mu_{N,(\beta,\kappa),(\infty,\kappa)} \big(\mathcal{E}_{7}(e) \big)}. \end{aligned}$$

Recalling the definitions of $\alpha_3(\beta, \kappa)$ and $\alpha_4(\beta, \kappa)$, we obtain the desired conclusion. \Box

10.4. Upper bounds on events. In this section we provide upper bounds on the events \mathcal{E}_1 , \mathcal{E}_2 , and \mathcal{E}_3 , defined in Sect. 10.1, and the events $\mathcal{E}_4(e)$, $\mathcal{E}_5(e)$, $\mathcal{E}_6(e)$, and $\mathcal{E}_7(e)$, from Sect. 10.3.

Proposition 10.10. Let β , $\kappa \geq 0$ be such that 3 hold, let $\gamma \in C^1(B_N)$ be a path with $\operatorname{dist}_0(\operatorname{supp} \gamma, \partial C_1(B_N)) \geq 8$, let $\gamma_0 \in C^1(B_N)$ be any path such that $\partial \gamma_0 = -\partial \gamma$, and let \mathcal{E}_1 be given by (10.3). Then

$$\begin{split} \mu_{N,(\beta,\kappa),(\infty,\kappa)}(\mathcal{E}_{1}) &\leq \mathbb{1}(\partial\gamma \neq 0) \Big(K_{3} K_{4}^{8} \alpha_{0}(\kappa)^{8} \alpha_{1}(\beta)^{6} \sum_{e \in \gamma} (K_{4} \alpha_{0}(\kappa))^{\max(0,\operatorname{dist}_{0}(e,\operatorname{supp}\gamma_{0})-8)} \\ &+ K_{3} |\operatorname{supp}\gamma| \big(K_{4} \alpha_{0}(\kappa) \big)^{\operatorname{dist}_{1}(\operatorname{supp}\gamma,\partial C_{1}(B_{N}))} \Big), \end{split}$$

where K_3 and K_4 are given in (6.6).

Proposition 10.11. Let β , $\kappa \ge 0$ be such that 3 hold, let $\gamma \in C^1(B_N)$ be a path such that $e \in \gamma$ and $p \in \hat{\partial}e$ we have dist₀(supp ∂p , $\partial C_1(B_N)$)) ≥ 8 , let $\gamma_0 \in C^1(B_N)$ be any path such that $\partial \gamma_0 = -\partial \gamma$, and let \mathcal{E}_2 be given by (10.4). Then

$$\begin{split} & \mu_{N,(\beta,\kappa),(\infty,\kappa)}(\mathcal{E}_{2}) \\ & \leq \mathbb{1}(\partial \gamma \neq 0) \cdot K_{3}K_{4}^{8}\alpha_{0}(\kappa)^{8}\alpha_{1}(\beta)^{6}\sum_{e \in \gamma} (K_{4}\alpha_{0}(\kappa))^{\max(0,\operatorname{dist}_{0}(e,\operatorname{supp}\gamma_{0})-8)} \\ & + K_{2}|\operatorname{supp}\gamma_{c}|\alpha_{2}(\beta,\kappa)^{6} + K_{3}K_{4}^{2}|\operatorname{supp}\gamma|\alpha_{0}(\kappa)^{2}\alpha_{1}(\beta)^{7} \\ & + 4K_{3}|\operatorname{supp}\gamma|(K_{4}\alpha_{0}(\kappa))^{\operatorname{dist}_{1}(\operatorname{supp}\gamma,\partial C_{1}(B_{N}))}, \end{split}$$

where K_2 is defined in (6.4), and K_3 and K_4 are given in (6.6).

Proposition 10.12. Let $\beta, \kappa \ge 0$ be such that 3 hold, let $\gamma \in C^1(B_N)$ be a path such for all $e \in \gamma$ and $p \in \hat{\partial}e$ we have $\operatorname{dist}_0(\operatorname{supp} \partial p, \partial C_1(B_N)) \ge 8$, and let \mathcal{E}_3 be defined by (10.5). Then

$$\mu_{N,\beta,\kappa}(\mathcal{E}_3) \le 18^4 K_5 |\operatorname{supp} \gamma | \alpha_0(\kappa)^2 \alpha_1(\beta)^{12}$$
(10.20)

where K_5 is given by (6.7).

Proposition 10.13 [Proposition 7.10 in [18]]. Let $\beta, \kappa \geq 0$ be such that 3 holds, let $e \in C_1(B_N)$ be such that the support of $\hat{\partial} e$ contains no boundary plaquettes of B_N , and let $\mathcal{E}_4(e)$ be given by (10.11).

Then

$$\mu_{N,(\beta,\kappa),(\infty,\kappa)}\left(\mathcal{E}_4(e)\right) \le K_7 \alpha_0(\kappa)^9 \alpha_1(\beta)^6 + K_3 \left(K_4 \alpha_0(\kappa)\right)^{\operatorname{dist}_1(e,\partial C_1(B_N))}$$

where

$$K_7 := 18^{10} K_3 \Big(2^8 \alpha_0(\kappa)^{-1} \Big(\Big(1 + \alpha_0(\kappa)/2 \Big)^8 - 1 \Big) + 2^8 K_4 \Big), \tag{10.21}$$

and K_3 and K_4 are given by (6.6).

Proposition 10.14 [Proposition 7.12 in [18]]. Let $\beta, \kappa \ge 0$ be such that 3 holds. Next, let $e \in C_1(B_N)$ be such that the support of $\hat{\partial} e$ contains no boundary plaquettes of B_N , and let $\mathcal{E}_5(e)$ be given by (10.12).

Then

$$\mu_{N,(\beta,\kappa),(\infty,\kappa)}(\mathcal{E}_{5}(e)) \leq K_{8}\alpha_{1}(\beta)^{6}\alpha_{0}(\kappa)^{6}\max\left(\alpha_{0}(\kappa),\alpha_{1}(\beta)^{6}\right) \\ +K_{9}\left(K_{4}\alpha_{0}(\kappa)\right)^{\operatorname{dist}_{1}(e,\partial C_{1}(B_{N}))},$$

where

$$K_8 := 2K_3 \cdot 18^8 (18^2 + 1) (2 + \alpha_0(\kappa))^7, \quad K_9 := K_3 (1 + ((2 + \alpha_0(\kappa)) \alpha_0(\kappa))^{-1}),$$
(10.22)

and K_3 and K_4 are given by (6.6).

Proposition 10.15 [Proposition 7.14 in [18]]. Let $\beta, \kappa \geq 0$ be such that 3 holds, let $e \in C_1(B_N)$ be such that $dist_0(supp \partial p, \partial C_1(B_N)) \geq 8$ for all $p \in \hat{\partial}e$, and let $\mathcal{E}_6(e)$ be given by (10.13).

Then

$$\mu_{N,\infty,\kappa}\left(\mathcal{E}_6(e)\right) \le K_{10}\alpha_0(\kappa)^8,\tag{10.23}$$

where

$$K_{10} := 18^{13} (1 - 18^2 \alpha_0(\kappa))^{-1}.$$
(10.24)

Proposition 10.16. Let $\beta, \kappa \ge 0$ be such that 3 holds, let $e \in C_1(B_N)$ be such that for all $p \in \hat{\partial}e$, the support of ∂p contains no boundary edges of B_N , and let $\mathcal{E}_7(e)$ be given by (10.14). Then

 $\mu_{N,(\beta,\kappa),(\infty,\kappa)}\big(\mathcal{E}_7(e)\big) \leq 6K_2 \,\alpha_2(\beta,\kappa)^6,$

where K_2 is given by (6.4).

Before we provide proofs of Propositions 10.10, 10.11, 10.12, and 10.16, we state and prove the following lemma, which will be useful in these proofs.

Lemma 10.17. Let $\gamma \in C^1(B_N)$ be a path with $\partial \gamma \neq 0$ and dist₀(supp $\gamma, \partial C_1(B_N)$)) ≥ 8 , let $\sigma \in \Omega_0^1(B_N, G)$ be such that the support of any path $\hat{\gamma} \in C^1(B_N)$ with $\partial \hat{\gamma} = -\partial \gamma$ intersects supp σ . Then, for any $e \in \text{supp } \sigma$ and any path $\gamma_0 \in C^1(B_N)$ such that $\partial \gamma_0 = -\partial \gamma$, we have

$$|(\operatorname{supp} \sigma)^+| \ge \max(\operatorname{dist}_0(e, \operatorname{supp} \gamma_0), 8).$$

Proof. Since $\partial \gamma \neq 0$ and the support of any path $\hat{\gamma} \in C^1(B_N)$ with $\partial \hat{\gamma} = -\partial \gamma$ intersects supp σ , we must have supp $\sigma \neq \emptyset$. Since $\sigma \in \Omega^1_0(B_N, G)$, it thus follows from Lemma 2.9 that

$$\left| (\operatorname{supp} \sigma)^+ \right| \ge 8.$$

Using the definition of γ_0 , the desired conclusion now follows.

By Lemma 5.2, we have $\hat{\sigma}'|_{E_{\hat{\sigma},\hat{\sigma}'}} \leq \hat{\sigma}'$, and hence, since $\hat{\sigma}' \leq \hat{\sigma}'|_{E_{\hat{\sigma},\hat{\sigma}'}}$, it follows from Lemma 2.3 (2.3) that $\hat{\sigma}' \leq \hat{\sigma}'$. Since $\hat{\sigma}' \in \Omega_0^1(B_N, G)$, we have $d\hat{\sigma}' = 0$, and hence we conclude that $d\hat{\sigma}' = 0$.

Since $\hat{\sigma}'$ disturbs γ , the set supp $\gamma \cap$ supp $\hat{\sigma}'$ must be non-empty (since otherwise, we could let $\hat{\gamma} = -\gamma$ and $\bar{\sigma} = \hat{\sigma}'$ in Definition 9.1). Fix some edge $e \in$ supp $\gamma \cap$ supp $\hat{\sigma}'$.

Since $\hat{\sigma}' \leq \hat{\sigma}'|_{E_{\hat{\sigma},\hat{\sigma}'}}$, we have $\operatorname{supp} \hat{\sigma}' \subseteq E_{\hat{\sigma},\hat{\sigma}'}$. Using the definition of $E_{\hat{\sigma},\hat{\sigma}'}$, we conclude that $d(\hat{\sigma}|_{\mathcal{C}_{\mathcal{G}(\hat{\sigma},\hat{\sigma}')}(\operatorname{supp} \hat{\sigma}')}) \neq 0$. Since $\hat{\sigma}'$ is irreducible and $e \in \operatorname{supp} \hat{\sigma}'$, Lemma 5.3 implies that $\mathcal{C}_{\mathcal{G}(\hat{\sigma},\hat{\sigma}')}(\operatorname{supp} \hat{\sigma}') = \mathcal{C}_{\mathcal{G}(\hat{\sigma},\hat{\sigma}')}(e)$, and hence $d(\hat{\sigma}|_{\mathcal{C}_{\mathcal{G}(\hat{\sigma},\hat{\sigma}')}(e)}) \neq 0$. Since $\hat{\sigma}'$ is irreducible, satisfies $d\hat{\sigma}' = 0$, and disturbs γ , the support of any path $\hat{\gamma} \in C^1(B_N)$ with $\partial \hat{\gamma} = -\partial \gamma$ must intersect the support of $\hat{\sigma}'$. This implies in particular that we must have $\partial \gamma \neq 0$. Applying Lemma 10.17, we thus obtain $|(\operatorname{supp} \hat{\sigma}')^+| \geq \max(\operatorname{dist}_0(e, \operatorname{supp} \gamma_0), 8)$, and consequently $|\mathcal{G}_{\mathcal{G}(\hat{\sigma},\hat{\sigma}')}(e)| \geq 2 \max(\operatorname{dist}_0(e, \operatorname{supp} \gamma_0), 8)$. To sum up, we have showed that

$$\begin{split} &\mu_{N,(\beta,\kappa),((\infty,\kappa))}(\mathcal{E}_{1})\\ &\leq \mathbb{1}(\partial\gamma\neq 0)\,\mu_{N,\beta,\kappa}\times\mu_{N,\infty,\kappa}\Big(\Big\{(\hat{\sigma},\hat{\sigma}')\in\Omega^{1}(B_{N},G)\times\Omega^{1}_{0}(B_{N},G):\\ &\exists e\in\gamma \text{ such that}\\ &d\big(\hat{\sigma}|_{\mathcal{C}_{\mathcal{G}(\hat{\sigma},\hat{\sigma}')}(e)}\big)\neq 0 \text{ and }|\mathcal{C}_{\mathcal{G}(\hat{\sigma},\hat{\sigma}')}(e)|\geq 2\max\big(\text{dist}_{0}(e,\operatorname{supp}\gamma_{0}),8\big)\Big\}\Big)\\ &\leq \mathbb{1}(\partial\gamma\neq 0)\,\sum_{e\in\gamma}\mu_{N,\beta,\kappa}\times\mu_{N,\infty,\kappa}\Big(\Big\{(\hat{\sigma},\hat{\sigma}')\in\Omega^{1}(B_{N},G)\times\Omega^{1}_{0}(B_{N},G):\\ &d\big(\hat{\sigma}|_{\mathcal{C}_{\mathcal{G}(\hat{\sigma},\hat{\sigma}')}(e)}\big)\neq 0 \text{ and }|\mathcal{C}_{\mathcal{G}(\hat{\sigma},\hat{\sigma}')}(e)|\geq 2\max\big(\text{dist}_{0}(e,\operatorname{supp}\gamma_{0}),8\big)\Big\}\Big)\end{split}$$

Applying Proposition 6.1 with $M = \max(\text{dist}_0(e, \text{supp } \gamma_0), 8)$ and M' = 1 for each $e \in \gamma$, we obtain (10.10) as desired.

Proof of Proposition 10.11. Assume first that $(\hat{\sigma}, \hat{\sigma}') \in \mathcal{E}_2$. Then there exists an irreducible 1-form $\hat{\sigma}' \leq \hat{\sigma}|_{E_{\hat{\sigma},\hat{\sigma}'}}$ that disturbs γ . Since $\hat{\sigma}'$ disturbs γ , the set supp $\gamma \cap \text{supp } \hat{\sigma}'$ must be non-empty (since otherwise, we could let $\hat{\gamma} = -\gamma$ and $\bar{\sigma} = \hat{\sigma}'$ in Definition 9.1). Fix some edge $e \in \text{supp } \gamma \cap \text{supp } \hat{\sigma}'$.

By definition, we must have $\operatorname{supp} \hat{\sigma}' \subseteq E_{\hat{\sigma},\hat{\sigma}'}$. Using the definition of $E_{\hat{\sigma},\hat{\sigma}'}$, we see that $d(\hat{\sigma}|_{\mathcal{C}_{\mathcal{G}(\hat{\sigma},\hat{\sigma}')}(\operatorname{supp}\hat{\sigma}')}) \neq 0$. Since $e \in \operatorname{supp} \hat{\sigma}'$ and $\hat{\sigma}'$ is irreducible, it follows from Lemma 5.3 that $\mathcal{C}_{\mathcal{G}(\hat{\sigma},\hat{\sigma}')}(\operatorname{supp} \hat{\sigma}') = \mathcal{C}_{\mathcal{G}(\hat{\sigma},\hat{\sigma}')}(e)$, and hence $d(\hat{\sigma}|_{\mathcal{C}_{\mathcal{G}(\hat{\sigma},\hat{\sigma}')}(e)}) \neq 0$.

Since $\hat{\sigma}'$ is irreducible and disturbs γ , we must be in one of the following four cases.

- (1) $\partial \gamma \neq 0$, and all paths $\hat{\gamma} \in C^1(B_N)$ with $\partial \hat{\gamma} = -\partial \gamma$ intersects the support of $\hat{\sigma}'$. In this case, by Lemma 10.17, we must have $|\operatorname{supp} \hat{\sigma}'| \geq 2 \max(\operatorname{dist}_0(e, \operatorname{supp} \gamma_0), 8)$, and consequently, $|\mathcal{G}_{\mathcal{G}}(\hat{\sigma}, \hat{\sigma}')(e)| \geq 2 \max(\operatorname{dist}_0(e, \operatorname{supp} \gamma_0), 8)$.
- (2) $\hat{\sigma}'$ contains a minimal vortex centered around some edge $e' \in \gamma_c$. Since $\hat{\sigma}' \leq \hat{\sigma}|_{E_{\hat{\sigma},\hat{\sigma}'}}$ by definition, and $\hat{\sigma}|_{E_{\hat{\sigma},\hat{\sigma}'}} \leq \hat{\sigma}$ by Lemma 5.2, is follows from Lemma 2.11, applied twice, that $\hat{\sigma}$ also contains a minimal vortex centered around some edge $e' \in \gamma_c$.

- (3) $|(\sup p d\hat{\sigma}')^+| > 6$. In this case, by the same argument as above, we must have both $|\sup p d\hat{\sigma}|_{\mathcal{C}_{\mathcal{G}}(\hat{\sigma}, \hat{\sigma}')}(e)| \ge 2 \cdot 7$ and $|\mathcal{C}_{\mathcal{G}}(\hat{\sigma}, \hat{\sigma}')(e)| \ge 2 \cdot 2$.
- (4) $\hat{\sigma}'$ supports a vortex ν with support at the boundary of B_N . In this case, by the same argument as above, we must have $|\mathcal{C}_{\mathcal{G}}(\hat{\sigma},\hat{\sigma}')(e)| \ge 2 \cdot \text{dist}_1(e, \partial C_1(B_N))$.

Consequently, we have showed that

$$\begin{split} &\mu_{N,(\beta,\kappa),(\infty,\kappa)}(\mathcal{E}_{1}) \\ &\leq \mathbb{1}(\partial\gamma \neq 0) \ \mu_{N,\beta,\kappa} \times \mu_{N,\infty,\kappa} \Big(\Big\{ (\hat{\sigma}, \hat{\sigma}') \in \Omega^{1}(B_{N}, G) \times \Omega^{1}_{0}(B_{N}, G) \colon \exists e \in \gamma \text{ such that} \\ & d\big(\hat{\sigma}|_{\mathcal{C}_{\mathcal{G}}(\hat{\sigma}, \hat{\sigma}')}(e) \big) \neq 0 \text{ and } |\mathcal{C}_{\mathcal{G}}(\hat{\sigma}, \hat{\sigma}')(e)| \geq 2 \max \big(\operatorname{dist}_{0}(e, \operatorname{supp} \gamma_{0}), 8 \big) \Big\} \Big) \\ &+ \mu_{N,\beta,\kappa} \Big(\Big\{ \hat{\sigma} \in \Omega^{1}(B_{N}, G) \colon \exists e' \in \gamma_{c} \text{ and } \nu \leq \hat{\sigma} \text{ s.t. } \nu \text{ is a minimal vortex around} e' \Big\} \Big) \\ &+ \mu_{N,\beta,\kappa} \times \mu_{N,\infty,\kappa} \Big(\Big\{ (\hat{\sigma}, \hat{\sigma}') \in \Omega^{1}(B_{N}, G) \times \Omega^{1}_{0}(B_{N}, G) \colon \exists e \in \gamma \text{ such that} \\ & \left| d\big(\hat{\sigma}|_{\mathcal{C}_{\mathcal{G}}(\hat{\sigma}, \hat{\sigma}')}(e) \big| \geq 2 \cdot 7 \text{ and } |\mathcal{C}_{\mathcal{G}}(\hat{\sigma}, \hat{\sigma}')(e)| \geq 2 \cdot 2 \Big\} \Big) \\ &+ \mu_{N,\beta,\kappa} \times \mu_{N,\infty,\kappa} \Big(\Big\{ (\hat{\sigma}, \hat{\sigma}') \in \Omega^{1}(B_{N}, G) \times \Omega^{1}_{0}(B_{N}, G) \colon \exists e \in \gamma \text{ such that} \\ & d\big(\hat{\sigma}|_{\mathcal{C}_{\mathcal{G}}(\hat{\sigma}, \hat{\sigma}')}(e) \big) \neq 0 \text{ and } |\mathcal{C}_{\mathcal{G}}(\hat{\sigma}, \hat{\sigma}')(e)| \geq 2 \operatorname{dist}_{1}(e, \partial C_{1}(B_{N})) \Big\} \Big). \end{split}$$

By first applying union bounds to all terms, and then using Proposition 6.1 to upper bound the first, third, and fourth term and Proposition 6.3 to upper bound the second term, we obtain

$$\begin{split} & \mu_{N,(\beta,\kappa),(\infty,\kappa)}(\mathcal{E}_{2}) \\ & \leq \mathbb{1}(\partial \gamma \neq 0) \cdot K_{3}K_{4}^{8}\alpha_{0}(\kappa)^{8}\alpha_{1}(\beta)^{6}\sum_{e \in \gamma} (K_{4}\alpha_{0}(\kappa))^{\max(0,\operatorname{dist}_{0}(e,\operatorname{supp}\gamma_{0})-8)} \\ & + K_{3}|\operatorname{supp}\gamma|(K_{4}\alpha_{0}(\kappa))^{\operatorname{dist}_{0}(\operatorname{supp}\gamma,\partial C_{1}(B_{N}))} + K_{2}|\operatorname{supp}\gamma_{c}|\alpha_{2}(\beta,\kappa)^{6} \\ & + K_{3}K_{4}^{2}|\operatorname{supp}\gamma|\alpha_{0}(\kappa)^{2}\alpha_{1}(\beta)^{7} \\ & + K_{3}|\operatorname{supp}\gamma|(K_{4}\alpha_{0}(\kappa))^{\operatorname{dist}_{1}(\operatorname{supp}\gamma,\partial C_{1}(B_{N}))}\alpha_{1}(\beta)^{6} \\ & + 2K_{3}|\operatorname{supp}\gamma|(K_{4}\alpha_{0}(\kappa))^{\operatorname{dist}_{1}(\operatorname{supp}\gamma,\partial C_{1}(B_{N}))}. \end{split}$$

Simplifying this expression, and noting that by definition, we have $\alpha_1(\beta) \leq 1$, we obtain (10.11) as desired. This concludes the proof.

Proof of Proposition 10.12. Assume first that $\sigma \in \mathcal{E}_3$, and let Σ be a decomposition of σ . Further, let $e \in \gamma$, $p, p' \in \hat{\partial}e$, and $\hat{\sigma}, \hat{\sigma}' \in \Sigma$ be such that $\hat{\sigma} \neq \hat{\sigma}', d\hat{\sigma}(p) \neq 0$, and $d\hat{\sigma}'(p') \neq 0$. Without loss of generality we can assume that $e \in C_1(B_N)^+$. By Lemma 2.8, we must then have $|(\operatorname{supp} d\hat{\sigma})^+| \ge 6$ and $|(\operatorname{supp} d\hat{\sigma}')^+| \ge 6$, and hence $|(\operatorname{supp} d(\hat{\sigma} + \hat{\sigma}'))^+| \ge 12$.

Define $E_e := \{ e' \in C_1(B_N) : \hat{\partial} e' \cap \hat{\partial} e \neq \emptyset \}.$

Since $d\hat{\sigma}(p) \neq 0$, there must exist $e' \in \partial p$ such that $\hat{\sigma}(e') \neq 0$. Since $\hat{\sigma} \leq \sigma$, it follows that $\sigma(e') \neq 0$. Since $e' \in \partial p \subseteq E_e$, it follows that $C_{\mathcal{G}(\sigma,0)}(E_e)$ is nonempty. Moreover, since $\hat{\sigma}$ is irreducible, using Lemma 5.5, it follows that $\sup p \hat{\sigma} \subseteq C_{\mathcal{G}(\sigma,0)}(E_e)$. Completely analogously, we also obtain $\sup p \hat{\sigma}' \subseteq C_{\mathcal{G}(\sigma,0)}(E_e)$. Since $d\hat{\sigma}(p), d\hat{\sigma}'(p') \neq 0$, by Lemma (2.8), we must have $|(\sup p d\hat{\sigma})^+|, |(\sup p d\hat{\sigma}')^+| \geq 6$, and hence $|\sup d\sigma|_{\mathcal{C}_{\mathcal{G}(\sigma,0)}(E_e)})^+| \geq 2 \cdot 6 = 12$. Using Lemma 2.7, it follows that $|(\mathcal{C}_{\mathcal{G}(\sigma,0)}(E_e))^+| \ge 12/6 = 2.$

Combining these observations and using a union bound, we see that

$$\mu_{N,\beta,\kappa}(\mathcal{E}_{3}) \leq \sum_{e \in \mathcal{V}} \mu_{N,\beta,\kappa} \Big(\Big\{ \sigma \in \Omega^{1}(B_{N},G) \colon |\mathcal{C}_{\mathcal{G}(\sigma,0)}(E_{e})| \geq 2 \cdot 2, \text{ and } \Big| \text{supp} \, d\big(\sigma|_{\mathcal{C}_{\mathcal{G}(\sigma,0)}(E_{e})}\big) \Big| \\ \geq 2 \cdot 12 \Big\} \Big).$$

Applying Proposition 6.5 with M = 2 and M' = 12, we obtain (10.20) as desired.

Proof of Proposition 10.16. Recall that

$$\mathcal{E}_7(e) = \left\{ (\sigma, \sigma') \in \Omega^1(B_N, G) \times \Omega^1_0(B_N, G) \colon \exists p, p' \in \hat{\partial}e \text{ s.t. } d\sigma(p) \neq d\sigma(p') \right\}.$$

On this event, there must exist some $p \in \hat{\partial} e$ with $d\sigma(p) \neq 0$. Since $|\hat{\partial} e| = 6$, together with a union bound, the desired conclusion follows from Proposition 6.3. п

10.5. A second version of our main result. In this section, we prove a second version of our main result, by giving a refinement of Proposition 7.1. While the error term in Proposition 7.1 corresponds to the probability of the event that no cluster in $\mathcal{G}(\hat{\sigma}, \hat{\sigma}')$ both intersects γ and at the same time supports a vortex, the error term in Proposition 10.18 below essentially corresponds to the probability that no cluster in $\mathcal{G}(\hat{\sigma}, \hat{\sigma}')$ both intersects γ and at the same time supports a non-minimal vortex.

Proposition 10.18. Let β , $\kappa > 0$ be such that 3 hold, let $\gamma \in C^1(B_N)$ be a path such that dist₀(supp γ , $\partial C_1(B_N)$) > 8 and such that for each $e \in \gamma$ the support of $\hat{\partial} e$ contains no boundary plaquettes of B_N , and let $\gamma_0 \in C^1(B_N)$ be any path such that $\partial \gamma_0 = -\partial \gamma$. Then

$$\left| \mathbb{E}_{N,\beta,\kappa} \left[L_{\gamma}(\sigma) \right] - \mathbb{E}_{N,\infty,\kappa} \left[L_{\gamma}(\sigma) \right] \Theta_{N,\beta,\kappa}(\gamma) \right| \\ \leq 2K_{11} |\operatorname{supp} \gamma | \alpha_2(\beta,\kappa)^6 + 2K_{12} \sqrt{2|\operatorname{supp} \gamma | \alpha_2(\beta,\kappa)^6},$$
(10.25)

where

$$K_{11} := \mathbb{1}(\partial \gamma \neq 0) \cdot \frac{2K_3 K_4^8 \alpha_0(\kappa)^8 \alpha_1(\beta)^6 \sum_{e \in \gamma} (K_4 \alpha_0(\kappa))^{\max(0, \operatorname{dist}_0(e, \operatorname{supp} \gamma_0) - 8)}}{|\operatorname{supp} \gamma | \alpha_2(\beta, \kappa)^6} + \frac{4K_3 (K_4 \alpha_0(\kappa))^{\operatorname{dist}_1(\operatorname{supp} \gamma, \partial C_1(B_N))}}{\alpha_2(\beta, \kappa)^6} + \frac{K_2 |\operatorname{supp} \gamma_c|}{|\operatorname{supp} \gamma|} + \frac{K_3 K_4^2 \alpha_0(\kappa)^2 \alpha_1(\beta)^7}{\alpha_2(\beta, \kappa)^6} + \frac{18^4 K_5 \alpha_0(\kappa)^2 \alpha_1(\beta)^{12}}{2\alpha_2(\beta, \kappa)^6} + \sqrt{\frac{2K_8 \alpha_0(\kappa)^6 \alpha_1(\beta)^6 \alpha_4(\beta, \kappa) \max(\alpha_0(\kappa), \alpha_1(\beta)^6)}{|\operatorname{supp} \gamma | \alpha_2(\beta, \kappa)^{12}}}$$

$$+\sqrt{\frac{2K_{9}\alpha_{4}(\beta,\kappa)(K_{3}\alpha_{0}(\kappa))^{\operatorname{dist}_{1}(\operatorname{supp}\gamma,\partial C_{1}(B_{N}))}{|\operatorname{supp}\gamma|\alpha_{2}(\beta,\kappa)^{12}}}$$

$$+2\sqrt{\frac{2K_{7}\alpha_{0}(\kappa)^{9}\alpha_{1}(\beta)^{6}\alpha_{4}(\beta,\kappa)}{|\operatorname{supp}\gamma|\alpha_{2}(\beta,\kappa)^{12}}}$$

$$+2\sqrt{\frac{2K_{3}\alpha_{4}(\beta,\kappa)(K_{4}\alpha_{0}(\kappa))^{\operatorname{dist}_{1}(\operatorname{supp}\gamma,\partial C_{1}(B_{N}))}}{|\operatorname{supp}\gamma|\alpha_{2}(\beta,\kappa)^{12}}}$$

$$+\sqrt{\frac{12K_{2}\alpha_{2}(\beta,\kappa)^{6}\alpha_{3}(\beta,\kappa)}{|\operatorname{supp}\gamma|\alpha_{2}(\beta,\kappa)^{12}}},$$
(10.26)
$$=\sqrt{\frac{K_{10}|\operatorname{supp}\gamma_{c}|\alpha_{0}(\kappa)^{8}\alpha_{4}(\beta,\kappa)}{|\operatorname{supp}\gamma_{c}|\alpha_{3}(\beta,\kappa)}} + \sqrt{\frac{|\operatorname{supp}\gamma_{c}|\alpha_{3}(\beta,\kappa)}{|\operatorname{supp}\gamma_{c}|\alpha_{3}(\beta,\kappa)}}$$
(10.27)

$$K_{12} := \sqrt{\frac{K_{10} | \operatorname{supp} \gamma_c | \alpha_0(\kappa)^{\circ} \alpha_4(\beta, \kappa)}{| \operatorname{supp} \gamma | \alpha_2(\beta, \kappa)^6}} + \sqrt{\frac{| \operatorname{supp} \gamma_c | \alpha_3(\beta, \kappa)}{| \operatorname{supp} \gamma | \alpha_2(\beta, \kappa)^6}}.$$
(10.27)

 K_2 is given by (6.4), K_3 and K_4 are given by (6.6), K_5 is given by (6.7), K_7 is given by (10.21), K_8 and K_9 are given by (10.22), and K_{10} is given by (10.24).

Proof. By using first the definition of $\Theta_{N,\beta,\kappa}(\gamma)$, and then the triangle inequality, we see that

$$\begin{split} & \left| \mathbb{E}_{N,\beta,\kappa} \left[L_{\gamma}(\sigma) \right] - \mathbb{E}_{N,\infty,\kappa} \left[L_{\gamma}(\sigma) \right] \Theta_{N,\beta,\kappa}(\gamma) \right| \\ &= \left| \mathbb{E}_{N,\beta,\kappa} \left[L_{\gamma}(\sigma) \right] - \mathbb{E}_{N,\infty,\kappa} \left[L_{\gamma}(\sigma) \right] \mathbb{E}_{N,\infty,\kappa} \left[\prod_{e \in \gamma} \theta_{\beta,\kappa}(\sigma_{e}) \right] \right| \\ &\leq \left| \mathbb{E}_{N,\beta,\kappa} \left[L_{\gamma}(\sigma) \right] - \mathbb{E}_{N,\infty,\kappa} \left[L_{\gamma}(\sigma) \right] \mathbb{E}_{N,\beta,\kappa} \left[\prod_{e \in (\gamma - \gamma_{c}) - \gamma'} \rho(d\sigma(p_{e})) \right] \right| \\ &+ \left| \mathbb{E}_{N,\infty,\kappa} \left[L_{\gamma}(\sigma) \right] \right| \cdot \left| \mathbb{E}_{N,\beta,\kappa} \left[\prod_{e \in (\gamma - \gamma_{c}) - \gamma'} \rho(d\sigma(p_{e})) \right] \right| \\ &- \mathbb{E}_{N,\beta,\kappa} \left[\prod_{e \in (\gamma - \gamma_{c}) - \gamma'} \theta_{\beta,\kappa}(\sigma(e) - d\sigma(p_{e})) \right] \right| \\ &+ \left| \mathbb{E}_{N,\infty,\kappa} \left[L_{\gamma}(\sigma) \right] \right| \cdot \left| \mathbb{E}_{N,\beta,\kappa} \left[\prod_{e \in (\gamma - \gamma_{c}) - \gamma'} \theta_{\beta,\kappa}(\sigma(e) - d\sigma(p_{e})) \right] \right| \\ &- \mathbb{E}_{N,\infty,\kappa} \left[\prod_{e \in \gamma} \theta_{\beta,\kappa}(\sigma(e)) \right] \right|. \end{split}$$

Since $|L_{\gamma}(\sigma)| \le 1$, we can apply Proposition 10.3, Proposition 10.4 and Proposition 10.5 in order to obtain

$$\begin{split} \left| \mathbb{E}_{N,\beta,\kappa} \big[L_{\gamma}(\sigma) \big] - \mathbb{E}_{N,\infty,\kappa} \big[L_{\gamma}(\sigma) \big] \mathbb{E}_{N,\infty,\kappa} \Big[\prod_{e \in \gamma} \theta_{\beta,\kappa}(\sigma_{e}) \Big] \right| \\ &\leq 2\mu_{N,(\beta,\kappa),(\infty,\kappa)}(\mathcal{E}_{1}) + 2\mu_{N,(\beta,\kappa),(\infty,\kappa)}(\mathcal{E}_{2}) + 2\mu_{N,\beta,\kappa}(\mathcal{E}_{3}) \\ &+ 2\sqrt{2\alpha_{4}(\beta,\kappa)} \sum_{e \in \gamma} \mu_{N,(\beta,\kappa),(\infty,\kappa)} \big(\mathcal{E}_{5}(e) \big) + 4\sqrt{2\alpha_{4}(\beta,\kappa)} \sum_{e \in \gamma} \mu_{N,(\beta,\kappa),(\infty,\kappa)} \big(\mathcal{E}_{4}(e) \big) \end{split}$$

$$+ 2 \sqrt{2\alpha_4(\beta,\kappa) \sum_{e \in \gamma_c} \mu_{N,\infty,\kappa} (\mathcal{E}_6(e)) + 2\sqrt{2|\operatorname{supp} \gamma_c|\alpha_3(\beta,\kappa)}} + 2 \sqrt{2\alpha_3(\beta,\kappa) \sum_{e \in \gamma} \mu_{N,(\beta,\kappa),(\infty,\kappa)} (\mathcal{E}_7(e))}.$$

Inserting the upper bounds from Proposition 10.10, Proposition 10.11, Proposition 10.12, Proposition 10.13, Proposition 10.14, Proposition 10.15, and Proposition 10.16, and using the inequality $\sqrt{a+b} \le \sqrt{a} + \sqrt{b}$, we obtain (10.25) as desired.

10.6. An upper bound. The following result generalizes [9, Lemma 7.12] and [17, Lemma 3.3], and is completely analogous to Lemma 12.3 in [18].

Proposition 10.19 [Lemma 12.3 in [18]]. Let $\beta, \kappa \ge 0$, and let $\gamma \in C^1(B_N)$ be a path such that no edge in γ is in the boundary of B_N . Then

$$\left|\mathbb{E}_{N,\beta,\kappa}\left[L_{\gamma}(\sigma)\right]\right| \leq \exp\left(-|\operatorname{supp}(\gamma - \gamma_{c})| \alpha_{5}(\beta,\kappa)\right).$$

Remark 10.20. In Lemma 12.3 in [18], γ is assumed to be a generalized loop, rather than a path. However, since the proof is identical in the two cases, we do not include a proof here.

10.7. A proof of Theorem 10.1. In this section, we give a proof of Theorem 10.1. Before we give this proof, we recall the following lemma from [18].

Lemma 10.21 [Lemma 8.2 in [18]]. Let $\beta, \kappa \ge 0$, and for each $g \in G$, let $j_g > 0$ be given. Further, let $j := \sum_{g \in G} j_g$. Then

$$\left|\prod_{g\in G}\theta_{\beta,\kappa}(g)^{j_g}\right| \leq e^{-j\alpha_5(\beta,\kappa)}.$$

Proof of Theorem 10.1. Let *N* be sufficiently large, so that $dist_0(supp \gamma, \partial C_1(B_N)) \ge 8$, and so that for each $e \in \gamma$, the support of $\hat{\partial} e$ contains no boundary plaquettes of B_N .

Then the assumptions of Proposition 10.18 holds, and hence

$$\begin{aligned} \left| \mathbb{E}_{N,\beta,\kappa} \left[L_{\gamma}(\sigma) \right] - \mathbb{E}_{N,\infty,\kappa} \left[L_{\gamma}(\sigma) \right] \Theta_{N,\beta,\kappa}(\gamma) \right| \\ &\leq B \cdot 2 |\operatorname{supp} \gamma | \alpha_{5}(\beta,\kappa) + B' \cdot 2\sqrt{2} |\operatorname{supp} \gamma | \alpha_{5}(\beta,\kappa) \end{aligned}$$

where

$$B := K_{11}\alpha_2(\beta,\kappa)^6 / \alpha_5(\beta,\kappa) \quad \text{and} \quad B' := K_{12} \sqrt{\alpha_2(\beta,\kappa)^6 / \alpha_5(\beta,\kappa)}, \quad (10.28)$$

where K_{11} and K_{12} are given in (10.26) and (10.26) respectively.

Using that for x > 0, we have $x \le e^x$, and $2\sqrt{x} \le e^x$, it follows that

$$\left|\mathbb{E}_{N,\beta,\kappa}\left[L_{\gamma}(\sigma)\right] - \mathbb{E}_{N,\infty,\kappa}\left[L_{\gamma}(\sigma)\right]\Theta_{N,\beta,\kappa}(\gamma)\right| \le (B+B')e^{2|\operatorname{supp}\gamma|\alpha_{5}(\beta,\kappa)}.$$
(10.29)

Now recall that, by Proposition 10.19, we have $\left|\mathbb{E}_{N,\beta,\kappa}[L_{\gamma}(\sigma)]\right| \leq e^{-|\operatorname{supp}\gamma - \gamma_{c}|\alpha_{5}(\beta,\kappa)}$. By using the triangle inequality and applying Lemma 10.21, it follows that

$$\begin{aligned} \left| \mathbb{E}_{N,\beta,\kappa} \left[L_{\gamma}(\sigma) \right] - \mathbb{E}_{N,\infty,\kappa} \left[L_{\gamma}(\sigma) \right] \Theta_{N,\beta,\kappa}(\gamma) \right| \\ &\leq \left| \mathbb{E}_{N,\beta,\kappa} \left[L_{\gamma}(\sigma) \right] \right| + \left| \mathbb{E}_{N,\infty,\kappa} \left[L_{\gamma}(\sigma) \right] \right| \cdot \left| \Theta_{N,\beta,\kappa}(\gamma) \right| \\ &\leq e^{-|\operatorname{supp}\gamma - \gamma_{c}|\alpha_{5}(\beta,\kappa)} + 1 \cdot e^{-|\operatorname{supp}\gamma|\alpha_{5}(\beta,\kappa)} \leq 2e^{-|\operatorname{supp}\gamma - \gamma_{c}|\alpha_{5}(\beta,\kappa)}. \end{aligned}$$
(10.30)

Combining (10.29) and (10.30), we obtain

$$\left| \mathbb{E}_{N,\beta,\kappa} [L_{\gamma}(\sigma)] - \mathbb{E}_{N,\infty,\kappa} [L_{\gamma}(\sigma)] \Theta_{N,\beta,\kappa}(\gamma) \right|^{1+2|\operatorname{supp}\gamma|/|\operatorname{supp}(\gamma-\gamma_{c})|}$$

$$\leq 2^{2|\operatorname{supp}\gamma|/|\operatorname{supp}(\gamma-\gamma_{c})|} (B+B').$$
(10.31)

Recalling Proposition 2.16 and Proposition 2.19, and letting $N \to \infty$, the desired conclusion thus follows from (10.31) after simplification.

10.8. Simplifications for rectangular paths and $G = \mathbb{Z}_2$. The purpose of this section is to establish the tools we need in order make the small adjustments to the proof of Theorem 10.1 needed to instead obtain Theorem 1.1.

In order to simplify notations, for β , $\kappa \ge 0$ and a path γ , we define

$$\Theta_{N,\beta,\kappa}'(\gamma) := e^{-2|\operatorname{supp}\gamma|e^{-24\beta-4\kappa}\left(1+(e^{8\kappa}-1)|\operatorname{supp}\gamma|^{-1}\sum_{e\in\gamma}\mathbb{E}_{N,\infty,\kappa}[\mathbb{1}_{\sigma(e)=1}]\right)}.$$

The main result in this section is the following proposition.

Proposition 10.22. Let $\beta, \kappa \geq 0$ be such that 3 and $6\beta > \kappa$ both hold, let γ be a path along the boundary of a rectangle with side lengths ℓ_1 and ℓ_2 which is such that $\operatorname{dist}_0(\operatorname{supp} \gamma, \partial C_1(B_N)) \geq 8$, and let $G = \mathbb{Z}_2$.

Then

$$\left|\Theta_{N,\beta,\kappa}(\gamma) - \Theta_{N,\beta,\kappa}'(\gamma)\right| \le 2\sqrt[3]{\frac{K_{13}\alpha_2(\beta,\kappa)}{|\operatorname{supp}\gamma|}} + K_{14}|\operatorname{supp}\gamma|\alpha_2(\beta,\kappa)^{12}, \quad (10.32)$$

where

$$K_{13} := \left(4 + 4K_1 K_4^8 \alpha_0(\kappa)^4 + 4K_3 K_4^4 \alpha_2(\beta, \kappa)^6 \cdot |\operatorname{supp} \gamma| (K_4 \alpha_0(\kappa))^{\min(\ell_1, \ell_2) - 4} + 4K_3 K_4^4 \alpha_2(\beta, \kappa)^6 \cdot 32(K_4 \alpha_0(\kappa))^4 + 4K_3 K_4^4 \alpha_2(\beta, \kappa)^6 \cdot \frac{4(K_4 \alpha_0(\kappa))^5}{1 - K_4 \alpha_0(\kappa)}\right),$$
(10.33)

$$K_{14} := 4 \left(1 + K_1(\infty, \kappa) K_4^8 (\alpha_0(\kappa))^4\right),$$
(10.34)

 K_1 is given by (6.2), K_3 and K_4 are given by (6.6).

The second result which we will state and prove in this section is the following proposition, which will be used to simplify the error term in the proof of Theorem 1.1.

Proposition 10.23. Let β , $\kappa \ge 0$ be such that 3 holds, and let γ be an open path along the boundary of a rectangle with side lengths ℓ_1 and ℓ_2 . Then there is a path $\gamma_0 \in C^1(B_N)$ such that $\partial \gamma_0 = -\partial \gamma$ and

$$\sum_{e \in \gamma} \left(K_4 \alpha_0(\kappa) \right)^{\max(0, \operatorname{dist}_0(e, \operatorname{supp} \gamma_0) - 8)} \leq 16 + \frac{2K_4 \alpha_0(\kappa)}{1 - K_4 \alpha_0(\kappa)} + \frac{|\operatorname{supp} \gamma|}{2} \left(K_4 \alpha_0(\kappa) \right)^{\max(0, \min(\ell_1, \ell_2) - 7)},$$

where K_4 is given by (6.6).

Before we give a proofs of Proposition 10.22 and Proposition 10.23, we will state and prove a few useful lemmas. For these lemmas, it will be useful to note that when $G = \mathbb{Z}_2$ and $\rho(G) = \{1, -1\}$, then

$$\theta_{\beta,\kappa}(0) = \frac{1 - e^{-24\beta - 4\kappa}}{1 + e^{-24\beta - 4\kappa}} \quad \text{and} \quad \theta_{\beta,\kappa}(1) = \frac{1 - e^{-24\beta + 4\kappa}}{1 + e^{-24\beta + 4\kappa}}.$$
 (10.35)

From this, it in particular follows that when $6\beta > \kappa$, then $\theta_{\beta,\kappa}(0)$, $\theta_{\beta,\kappa} > 0$. Next, we recall from Section 12.2 in [18], that when $G = \mathbb{Z}_2$ and $\rho(G) = \{1, -1\}$, we have

$$\begin{aligned} \alpha_{0}(r) &= \alpha_{1}(r) = \varphi_{r}(1)^{2} = e^{-4r}, \qquad \alpha_{2}(\beta,\kappa) = e^{-4(\beta+\kappa/6)}, \\ \alpha_{3}(\beta,\kappa) &= 1 - \theta_{\beta,\kappa}(0) = \frac{2e^{-24\beta-4\kappa}}{1 + e^{-24\beta-4\kappa}}, \qquad \alpha_{5}(\beta,\kappa) = 1 - \theta_{\beta,\kappa}(0) \\ &= \frac{2e^{-24\beta-4\kappa}}{1 + e^{-24\beta-4\kappa}}, \\ \alpha_{4}(\beta,\kappa) &= \theta_{\beta,\kappa}(0) - \theta_{\beta,\kappa}(1) = \frac{2e^{-24\beta}(e^{4\kappa} - e^{-4\kappa})}{(1 + e^{-24\beta-4\kappa})(1 + e^{-24\beta+4\kappa})}. \end{aligned}$$
(10.36)

Proof of Proposition 10.23. Choose γ_0 so that $\gamma + \gamma_0$ is a generalized loop along the boundary of a rectangle with side lengths $\ell_1, \ell_2 \ge 2$.

Let $e_1, e_2, \ldots, e_{|\operatorname{supp} \gamma|}$ be the edges in γ , labelled according to their order in the path γ . Then, for any $j \in \{1, 2, \ldots, |\operatorname{supp} \gamma|\}$, one verifies that (see Fig. 7)

dist₀(*e*, supp
$$\gamma_0$$
) $\geq \max(8, \operatorname{dist}_1(e, \operatorname{supp} \gamma_0)) \geq \max(8, \min(j, |\operatorname{supp} \gamma| - j + 1, \ell_1, \ell_2) + 1).$

Using this inequality, we obtain

$$\begin{split} \sum_{e \in \gamma} & \left(K_4 \alpha_0(\kappa) \right)^{\max(0, \operatorname{dist}_0(e, \operatorname{supp} \gamma_0) - 8)} \leq \sum_{e \in \gamma} \left(K_4 \alpha_0(\kappa) \right)^{\max(0, \min(j, |\operatorname{supp} \gamma| - j + 1, \ell_1, \ell_2) + 1 - 8)} \\ & \leq 16 + 2 \sum_{j=9}^{\infty} \left(K_4 \alpha_0(\kappa) \right)^{j-8} + \frac{|\operatorname{supp} \gamma|}{2} \left(K_4 \alpha_0(\kappa) \right)^{\max(0, \min(\ell_1, \ell_2) - 7)}. \end{split}$$

Evaluating the geometric sum above, we obtain the desired conclusion.



(C) Given an edge $e_j \in \gamma$, we draw the support of a 1-form σ with $\operatorname{supp} \gamma_0 \cap C_{\mathcal{G}(\sigma)}(e_j) \neq \emptyset$ which minimizes $|\operatorname{supp} \sigma|$.

(D) Given an edge $e_{j'} \in \gamma$, we draw the support of a 1-form σ with $\operatorname{supp} \gamma_0 \cap \mathcal{C}_{\mathcal{G}(\sigma)}(e_{j'}) \neq \emptyset$ which minimizes $|\operatorname{supp} \sigma|$.

Fig. 7. In the figures above, we illustrate the setting of the proof of Proposition 10.23. Note in particular that for the edge e_j in 7a, we have dist₁(e_j , supp γ_0) = $| \operatorname{supp} \gamma | - j + 2$, and for the edge $e_{j'}$ in 7b, we have dist₁($e_{j'}$, supp γ_0) = $\min(\ell_1, \ell_2) + 1$

We now proceed to the proof of Proposition 10.22. Before we give this proof, we will state and prove a few lemmas. To simplify the notation in these lemmas, when $6\beta > \kappa$ and $\sigma \sim \mu_{N,\infty,\kappa}$, we define the following random variable.

$$\Upsilon_{\beta,\kappa}(\gamma) := |\operatorname{supp} \gamma|^{-1} \sum_{e \in \gamma} \log \theta_{\beta,\kappa} \big(\sigma(e) \big).$$
(10.37)

Lemma 10.24. Let $\beta, \kappa \ge 0$ be such that 3 and $6\beta > \kappa$ both hold, let γ be path along a rectangle with side lengths ℓ_1 and ℓ_2 and such that $\operatorname{dist}_0(\gamma, \partial C_1(B_N)) \ge 8$, and let $G = \mathbb{Z}_2$.

Then, for any $\varepsilon > 0$ *, we have*

$$\mu_{N,\infty,\kappa}\Big(\Big|\Upsilon_{\beta,\kappa}(\gamma) - \mathbb{E}_{N,\infty,\kappa}\big[\Upsilon_{\beta,\kappa}(\gamma)\big]\Big| \ge \varepsilon\Big) \le \frac{K_{13}\alpha_2(\beta,\kappa)}{\varepsilon^2|\operatorname{supp}\gamma|},$$

where K_{13} is given by (10.33).

Remark 10.25. The idea of the proof of Lemma 10.24 is essentially to use the weak law of large numbers for correlated random variables with exponential decay. For this to approach to work, we need the loop to be "smooth" enough for the sum of the covariances of all pairs of edges in γ to be finite. The reason for working with rectangular loops is that in this case, it is relatively easy to show that this holds. However, with small modifications, the conclusion of this lemma holds for more general classes of loops as well, as long as the path γ do not have too many corners.

Proof of Lemma 10.24. Fix some $\varepsilon > 0$.

By Chebyshev's inequality, we have

$$\varepsilon^{2} \mu_{N,\infty,\kappa} \Big(\Big| \Upsilon_{\beta,\kappa}(\gamma) - \mathbb{E}_{N,\infty,\kappa} \big[\Upsilon_{\beta,\kappa}(\gamma) \big] \Big| \ge \varepsilon \Big) \le \operatorname{Var}_{N,\infty,\kappa} \big[\Upsilon_{\beta,\kappa}(\gamma) \big]$$
$$= \sum_{e,e'\in\gamma} \operatorname{Cov}_{N,\infty,\kappa} \big(\Upsilon_{\beta,\kappa}(e), \Upsilon_{\beta,\kappa}(e') \big)$$
$$= |\operatorname{supp} \gamma|^{-2} \sum_{e,e'\in\gamma} \operatorname{Cov}_{N,\infty,\kappa} \Big(\log \theta_{\beta,\kappa} \big(\sigma(e) \big), \log \theta_{\beta,\kappa} \big(\sigma(e') \big) \Big).$$

By combining Proposition 5.17, applied with $f_0 = \log \theta_{\beta,\kappa}(\sigma(e))$ and $f_1 = \log \theta_{\beta,\kappa}(\sigma(e'))$, and Proposition 6.4, it follows that

$$\sum_{e,e'\in\gamma} \operatorname{Cov}_{N,\infty,\kappa} \left(\log \theta_{\beta,\kappa} (\sigma(e)), \log \theta_{\beta,\kappa} (\sigma(e')) \right)$$

$$\leq \sum_{e\in\operatorname{supp}\gamma} \operatorname{Var}_{N,\beta,\kappa} \left(\log \theta_{\beta,\kappa} (\sigma(e)) \right) + 2 \left\| \log \theta_{\beta,\kappa} \right\|_{\infty}^{2}$$

$$\sum_{e,e'\in\operatorname{supp}\gamma: e\neq e'} K_{3}(K_{4}\alpha_{0}(\kappa))^{\operatorname{dist}_{0}(e,e')}.$$

Since $0 \le \theta_{\beta,\kappa}(1) \le \theta_{\beta,\kappa}(0) \le 1$ for all $\beta, \kappa \ge 0$, we have

$$\begin{aligned} \left\| \log \theta_{\beta,\kappa} \right\|_{\infty} &\leq \left| \log \theta_{\beta,\kappa}(1) \right| \leq \left| \log \frac{1 - \varphi_{\beta}(1)^{12} \varphi_{\kappa}(1)^{-2}}{1 + \varphi_{\beta}(1)^{12} \varphi_{\kappa}(1)^{-2}} \right| \leq \left| \log e^{-2\varphi_{\beta}(1)^{12} \varphi_{\kappa}(1)^{-2}} \right| \\ &\leq 2\varphi_{\beta}(1)^{12} \varphi_{\kappa}(1)^{-2}. \end{aligned}$$

Next, recall that, by Proposition (6.1), applied with M = 1, M' = 0, $\beta = \kappa_1 = \infty$, and $\kappa_2 = \kappa$, for any edge $e \in \gamma$, we have

$$\mu_{N,\infty,\kappa}\Big(\big\{\hat{\sigma}\in\Omega^1_0(B_N,G)\colon \sigma(e)\neq 0\big\}\Big)\leq K_1(\infty,\kappa)\big(K_4\alpha_0(\kappa)\big)^8.$$

Consequently, for any $e \in \gamma$, we have

$$\begin{aligned} \operatorname{Var}_{N,\infty,\kappa} \left(\log \theta_{\beta,\kappa} \left(\sigma(e) \right) \right) &\leq \mathbb{E}_{N,\infty,\kappa} \left[\left(\log \theta_{\beta,\kappa} \left(\sigma(e) \right) \right)^2 \right] \\ &\leq \left(\log \theta_{\beta,\kappa} \left(0 \right) \right)^2 + K_1 \left(K_4 \alpha_0(\kappa) \right)^8 \left(\log \theta_{\beta,\kappa} \left(1 \right) \right)^2 \\ &\leq 4\varphi_\beta (1)^{24} \varphi_\kappa (1)^4 + K_1 \left(K_4 \alpha_0(\kappa) \right)^8 \cdot 4\varphi_\beta (1)^{24} \varphi_\kappa (1)^{-4} \\ &\leq \varphi_\beta (1)^{12} \varphi_\kappa (1)^2 \left(4 + 4K_1 K_4^8 \alpha_0(\kappa)^4 \right). \end{aligned}$$

Finally, note that

$$\sum_{e,e'\in\operatorname{supp}\gamma:\ e\neq e'} (K_4\alpha_0(\kappa))^{\operatorname{dist}_0(e,e')} \leq \sum_{e,e'\in\operatorname{supp}\gamma_R:\ e\neq e'} (K_4\alpha_0(\kappa))^{\operatorname{dist}_0(e,e')}$$
$$\leq |\operatorname{supp}\gamma| \Big(|\operatorname{supp}\gamma|/2 \cdot (K_4\alpha_0(\kappa))^{\min(\ell_1,\ell_2)} + 2\sum_{j=1}^{\infty} (K_4\alpha_0(\kappa))^{\min(j,8)} \Big)$$
$$\leq |\operatorname{supp}\gamma| \Big(|\operatorname{supp}\gamma|/2 \cdot (K_4\alpha_0(\kappa))^{\min(\ell_1,\ell_2)} + 16(K_4\alpha_0(\kappa))^8 + \frac{2(K_4\alpha_0(\kappa))^9}{1 - K_4\alpha_0(\kappa)} \Big).$$

Combining the above equations and recalling that when $G = \mathbb{Z}_2$, we have $\alpha_0(\kappa) = \varphi_{\kappa}(1)^2$ and $\alpha_2(\beta, \kappa)^6 = \alpha_0(\beta)^6 \alpha_0(\kappa)^6 = \varphi_{\beta}(1)^{12} \varphi_{\kappa}(1)^2$, we finally obtain

$$\begin{aligned} \varepsilon^{2} |\operatorname{supp} \gamma | \mu_{N,\infty,\kappa} \Big(\Big| \Upsilon_{\beta,\kappa}(\gamma) - \mathbb{E}_{N,\infty,\kappa} \big[\Upsilon_{\beta,\kappa}(\gamma) \big] \Big| \ge \varepsilon \Big) \\ &\le \alpha_{2}(\beta,\kappa)^{6} \Big(4 + 4K_{1}K_{4}^{8}\alpha_{0}(\kappa)^{4} \Big) \\ &+ 4K_{3}K_{4}^{4}\alpha_{2}(\beta,\kappa)^{12} \Big(|\operatorname{supp} \gamma | (K_{4}\alpha_{0}(\kappa))^{\min(\ell_{1},\ell_{2})-4} + 32(K_{4}\alpha_{0}(\kappa))^{4} + \frac{4(K_{4}\alpha_{0}(\kappa))^{5}}{1 - K_{4}\alpha_{0}(\kappa)} \Big). \end{aligned}$$

Rearranging this equation, the desired conclusion now immediately follows. This concludes the proof. $\hfill \Box$

Lemma 10.26. Let $\beta, \kappa \ge 0$ be such that 3 and $6\beta > \kappa$ both hold, let γ be path along the boundary of a rectangle with side lengths ℓ_1 and ℓ_2 , and let $G = \mathbb{Z}_2$. Then

$$\left|\Theta_{N,\beta,\kappa}(\gamma)-e^{|\operatorname{supp}\gamma|\mathbb{E}_{N,\infty,\kappa}[\Upsilon_{\beta,\kappa}(\gamma)]}\right| \leq 2\sqrt[3]{\frac{K_{13}\alpha_{2}(\beta,\kappa)}{|\operatorname{supp}\gamma|}},$$

where $K_{13} = K_{13}(\ell_1, \ell_2)$ is given by (10.33).

Proof. Recall the definition of $\Upsilon_{\beta,\kappa}(\gamma)$ from (10.37), and note that

$$\Theta_{N,\beta,\kappa}(\gamma) = \mathbb{E}_{N,\infty,\kappa} \Big[\prod_{e \in \gamma} \theta_{\beta,\kappa} \big(\sigma(e) \big) \Big] = \mathbb{E}_{N,\infty,\kappa} \Big[e^{\sum_{e \in \gamma} \log \theta_{\beta,\kappa} \big(\sigma(e) \big)} \Big]$$
$$= \mathbb{E}_{N,\infty,\kappa} \Big[e^{|\operatorname{supp} \gamma| \Upsilon_{\beta,\kappa}(\gamma)} \Big].$$

Consequently,

$$\begin{aligned} \left| Theta_{N,\beta,\kappa}(\gamma) - e^{|\operatorname{supp}\gamma|\mathbb{E}_{N,\infty,\kappa}[\Upsilon_{\beta,\kappa}(\gamma)]} \right| \\ &= \left| \mathbb{E}_{N,\infty,\kappa} \Big[e^{|\operatorname{supp}\gamma|\Upsilon_{\beta,\kappa}(\gamma)} - e^{|\operatorname{supp}\gamma|\mathbb{E}_{N,\infty,\kappa}[\Upsilon_{\beta,\kappa}(\gamma)]} \Big] \right| \\ &\leq \mathbb{E}_{N,\infty,\kappa} \Big[\left| e^{|\operatorname{supp}\gamma|\Upsilon_{\beta,\kappa}(\gamma)} - e^{|\operatorname{supp}\gamma|\mathbb{E}_{N,\infty,\kappa}[\Upsilon_{\beta,\kappa}(\gamma)]} \right| \Big] \end{aligned}$$

Next, note that since ρ is unitary, we have $|\theta_{\beta,\kappa}(g)| \leq 1$ for all $g \in G$, and hence $\Upsilon_{\beta,\kappa}(\gamma) \leq 0$. Now fix some $\varepsilon > 0$.

On the event $|\Upsilon_{\beta,\kappa}(\gamma) - \mathbb{E}_{N,\beta,\kappa}[\Upsilon_{\beta,\kappa}(\gamma)]| \ge \varepsilon$, since $\Upsilon_{\beta,\kappa}(\gamma) \le 0$, we must have

$$\left|e^{|\operatorname{supp}\gamma|\Upsilon_{\beta,\kappa}(\gamma)}-e^{|\operatorname{supp}\gamma|\mathbb{E}_{N,\infty,\kappa}[\Upsilon_{\beta,\kappa}(\gamma)]}\right|\leq 1.$$

On the other hand, on the event $|\Upsilon_{\beta,\kappa}(\gamma) - \mathbb{E}_{N,\beta,\kappa}[\Upsilon_{\beta,\kappa}(\gamma)]| < \varepsilon$, since $\Upsilon_{\beta,\kappa}(\gamma) \le 0$, we have

$$\left|e^{|\operatorname{supp}\gamma|\Upsilon_{\beta,\kappa}(\gamma)}-e^{|\operatorname{supp}\gamma|\mathbb{E}_{N,\infty,\kappa}[\Upsilon_{\beta,\kappa}(\gamma)]}\right|\leq \left|\Upsilon_{\beta,\kappa}(\gamma)-\mathbb{E}_{N,\infty,\kappa}[\Upsilon_{\beta,\kappa}(\gamma)]\right|<\varepsilon.$$

Using Lemma 10.24 with $\varepsilon = (K_{13}\alpha_2(\beta, \kappa)|\operatorname{supp} \gamma|^{-1})^{1/3}$, we obtain the desired conclusion.

Lemma 10.27. Let $\beta, \kappa \geq 0$ be such that $6\beta > \kappa$, and let $G = \mathbb{Z}_2$. Then

$$\begin{aligned} \left| \theta_{\beta,\kappa}(0) - e^{-2e^{-24\beta - 4\kappa}} \right| &\leq 4 \left(e^{-24\beta - 4\kappa} \right)^2 \quad and \quad \left| \theta_{\beta,\kappa}(1) - e^{-2e^{-24\beta + 4\kappa}} \right| \\ &\leq 4 \left(e^{-24\beta + 4\kappa} \right)^2. \end{aligned}$$

Proof. For the first inequality, note that

$$\begin{aligned} \left| \theta_{\beta,\kappa}(0) - e^{-2e^{-24\beta - 4\kappa}} \right| &= \left| \frac{1 - e^{-24\beta - 4\kappa}}{1 + e^{-24\beta - 4\kappa}} - e^{-2e^{-24\beta - 4\kappa}} \right| \\ &\leq \left| \frac{1 - e^{-24\beta - 4\kappa}}{1 + e^{-24\beta - 4\kappa}} - (1 - 2e^{-24\beta - 4\kappa}) \right| + \left| (1 - 2e^{-24\beta - 4\kappa}) - e^{-2e^{-24\beta - 4\kappa}} \right) \right| \\ &\leq 2(e^{-24\beta - 4\kappa})^2 + \left(2e^{-24\beta - 4\kappa} \right)^2 / 2 = 4(e^{-24\beta - 4\kappa})^2. \end{aligned}$$

The second inequality follows analogously.

Proof of Proposition 10.22. By definition, we have

$$\mathbb{E}_{N,\infty,\kappa}[\Upsilon_{\beta,\kappa}(\gamma)] = \mathbb{E}_{N,\infty,\kappa}\Big[|\operatorname{supp} \gamma|^{-1} \sum_{e \in \gamma} \log \theta_{\beta,\kappa}(\sigma(e))\Big]$$

= $|\operatorname{supp} \gamma|^{-1} \sum_{e \in \gamma} \mathbb{E}_{N,\infty,\kappa}\Big[\log \theta_{\beta,\kappa}(\sigma(e))\Big]$
= $|\operatorname{supp} \gamma|^{-1} \sum_{e \in \gamma} \mathbb{E}_{N,\infty,\kappa}\Big[\sum_{g \in G} \log \mathbb{1}_{\sigma(e)=g} \theta_{\beta,\kappa}(g)\Big] = \sum_{g \in G} \log \theta_{\beta,\kappa}(g)|\operatorname{supp} \gamma|^{-1}$
 $\sum_{e \in \gamma} \mathbb{E}_{N,\infty,\kappa}\Big[\mathbb{1}_{\sigma(e)=g}\Big].$

Consequently, by Lemma 10.26,

$$\left|\Theta_{N,\beta,\kappa}(\gamma) - \theta_{\beta,\kappa}(0)^{\sum_{e \in \gamma} \mathbb{E}_{N,\infty,\kappa}[\mathbb{1}_{\sigma(e)=0}]} \theta_{\beta,\kappa}(1)^{\sum_{e \in \gamma} \mathbb{E}_{N,\infty,\kappa}[\mathbb{1}_{\sigma(e)=1}]}\right| \leq 2\sqrt[3]{\frac{K_{13}\alpha_2(\beta,\kappa)}{|\operatorname{supp}\gamma|}}.$$

Next, by Lemma 10.6, we have

$$\begin{aligned} \left| \theta_{\beta,\kappa}(0)^{\sum_{e \in \gamma} \mathbb{E}_{N,\infty,\kappa} [\mathbbm{1}_{\sigma(e)=0}]} \theta_{\beta,\kappa}(1)^{\sum_{e \in \gamma} \mathbb{E}_{N,\infty,\kappa} [\mathbbm{1}_{\sigma(e)=1}]} \right| \\ &- e^{-2e^{-24\beta-4\kappa} \sum_{e \in \gamma} \mathbb{E}_{N,\infty,\kappa} [\mathbbm{1}_{\sigma(e)=0}] - 2e^{-24\beta+4\kappa} \sum_{e \in \gamma} \mathbb{E}_{N,\infty,\kappa} [\mathbbm{1}_{\sigma(e)=1}]} \right| \\ &\leq \sum_{e \in \gamma} \mathbb{E}_{N,\infty,\kappa} [\mathbbm{1}_{\sigma(e)=0}] \cdot \left| \theta_{\beta,\kappa}(0) - e^{-2e^{-24-4\kappa}} \right| + \sum_{e \in \gamma} \mathbb{E}_{N,\infty,\kappa} [\mathbbm{1}_{\sigma(e)=1}] \cdot \left| \theta_{\beta,\kappa}(1) - e^{-2e^{-24+4\kappa}} \right|. \end{aligned}$$

By combining Lemma 10.27 with Proposition 6.1, applied with M = 1, M' = 0, $\beta = \kappa_1 = \infty$, and $\kappa_2 = \kappa$, we can bound the previous equation from above by

$$|\operatorname{supp} \gamma| \cdot 4(e^{-24\beta - 4\kappa})^2 + K_1(\infty, \kappa) (K_4 \alpha_0(\kappa))^8 |\operatorname{supp} \gamma| \cdot 4(e^{-24\beta + 4\kappa})^2.$$

Combining the previous equations, we thus obtain

$$\begin{aligned} \left| \Theta_{N,\beta,\kappa}(\gamma) - e^{-2e^{-24\beta - 4\kappa} \sum_{e \in \gamma} \mathbb{E}_{N,\infty,\kappa} [\mathbbm{1}_{\sigma(e)=0}] - 2e^{-24\beta + 4\kappa} \sum_{e \in \gamma} \mathbb{E}_{N,\infty,\kappa} [\mathbbm{1}_{\sigma(e)=1}]} \right| \\ &\leq 2\sqrt[3]{\frac{K_{13}\alpha_2(\beta,\kappa)}{|\operatorname{supp}\gamma|}} + |\operatorname{supp}\gamma| \cdot 4(e^{-24\beta - 4\kappa})^2 + K_1(\infty,\kappa) (K_4\alpha_0(\kappa))^8 |\operatorname{supp}\gamma| \\ &\cdot 4(e^{-24\beta + 4\kappa})^2. \end{aligned}$$

Rearranging this equation, and recalling from (10.36) that $e^{-24\beta-4\kappa} = \alpha_2(\beta,\kappa)^6$ and $\alpha_0(\kappa) = e^{-4\kappa}$, we obtain (10.32) as desired.

10.9. A proof of Theorem 1.1. We now provide a proof of Theorem 1.1. Since this proof is very similar to the proof of Theorem 10.1, we will refer to this proof in order to avoid repetition.

Proof of Theorem 1.1. Let *N* be sufficiently large so that $dist_0(\gamma, \partial B_N) \ge 8$ and so that for each $e \in \gamma$, $\hat{\partial} e$ contains no boundary plaquettes of $C_2(B_N)$.

Using (10.35), it follows that if β and κ satisfy the assumptions of Theorem 1.1, then 3 hold.

By combining Propositions 10.18 and 10.32, using that $\left|\mathbb{E}_{N,\infty,\kappa}[L_{\gamma}(\sigma)]\right| \leq 1$, we obtain

$$\begin{split} & \mathbb{E}_{N,\beta,\kappa} \Big[L_{\gamma}(\sigma) \Big] - \mathbb{E}_{N,\infty,\kappa} \Big[L_{\gamma}(\sigma) \Big] \Theta'_{N,\beta,\kappa}(\gamma) \Big| \\ & \leq \Big| \mathbb{E}_{N,\beta,\kappa} \Big[L_{\gamma}(\sigma) \Big] - \mathbb{E}_{N,\infty,\kappa} \Big[L_{\gamma}(\sigma) \Big] \Theta_{N,\beta,\kappa}(\gamma) \Big| + \Big| \Theta_{N,\beta,\kappa}(\gamma) - \Theta'_{N,\beta,\kappa}(\gamma) \Big| \\ & \leq \Big(B + \frac{K_{14}\alpha_2(\beta,\kappa)^{12}}{2\alpha_5(\beta,\kappa)} \Big) \cdot 2 |\operatorname{supp} \gamma | \alpha_5(\beta,\kappa) + B' \cdot 2\sqrt{2} |\operatorname{supp} \gamma | \alpha_5(\beta,\kappa) \Big| \\ & + \sqrt[3]{\frac{K_{13}\alpha_2(\beta,\kappa)}{|\operatorname{supp} \gamma |^2\alpha_5(\beta,\kappa)}} \cdot 2\sqrt[3]{|\operatorname{supp} \gamma | \alpha_5(\beta,\kappa)}, \end{split}$$

where *B* and B' are given in (10.28).

Using that for x > 0, we have $x \le e^x$, $2\sqrt{x} \le e^x$, and $2\sqrt[3]{x} \le e^x$, it follows that

$$\begin{split} \left| \mathbb{E}_{N,\beta,\kappa} \Big[L_{\gamma}(\sigma) \Big] - \mathbb{E}_{N,\infty,\kappa} \Big[L_{\gamma}(\sigma) \Big] \Theta_{N,\beta,\kappa}'(\gamma) \Big| \\ & \leq \Big(B + \frac{K_{14}\alpha_2(\beta,\kappa)^{12}}{2\alpha_5(\beta,\kappa)} + B' + \sqrt[3]{\frac{K_{13}\alpha_2(\beta,\kappa)}{|\operatorname{supp}\gamma|^2\alpha_5(\beta,\kappa)}} \Big) e^{2|\operatorname{supp}\gamma|\alpha_5(\beta,\kappa)}. \end{split}$$

Combining this inequality with (10.30), we obtain

$$\begin{split} & \left| \mathbb{E}_{N,\beta,\kappa} [L_{\gamma}(\sigma)] - \mathbb{E}_{N,\infty,\kappa} [L_{\gamma}(\sigma)] \mathbb{E}_{N,\infty,\kappa} \Big[\prod_{e \in \gamma} \theta_{\beta,\kappa}(\sigma_{e}) \Big] \right|^{1+2|\operatorname{supp}\gamma|/|\operatorname{supp}(\gamma-\gamma_{c})|} \\ & \leq 2^{2|\operatorname{supp}\gamma|/|\operatorname{supp}(\gamma-\gamma_{c})|} \Big(K_{11} + \frac{K_{14}\alpha_{2}(\beta,\kappa)^{12}}{2\alpha_{5}(\beta,\kappa)} + K_{12} + \sqrt[3]{\frac{K_{13}\alpha_{2}(\beta,\kappa)}{|\operatorname{supp}\gamma|^{2}\alpha_{5}(\beta,\kappa)}} \Big). \end{split}$$

$$(10.38)$$

Now recall (10.36), and note that these expression imply that

$$\frac{\alpha_2(\beta,\kappa)^6}{\alpha_5(\beta,\kappa)} \le 1 \quad \text{and} \quad \frac{\alpha_4(\beta,\kappa)\alpha_0(\kappa)^2}{\alpha_5(\beta,\kappa)} \le 1.$$
(10.39)

Using these equations and inequalities and Proposition 10.23, it follows that

$$\begin{split} & B + \frac{K_{14}\alpha_{2}(\beta,\kappa)^{12}}{2\alpha_{5}(\beta,\kappa)} + B' + \sqrt[3]{\frac{K_{13}\alpha_{2}(\beta,\kappa)}{|\operatorname{supp}\gamma|^{2}\alpha_{5}(\beta,\kappa)}} \\ & \leq \mathbbm{1}(\partial\gamma \neq 0) \cdot \frac{2K_{3}K_{4}^{8}\alpha_{0}(\kappa)^{7}}{\sqrt{|\operatorname{supp}\gamma|}} \\ & \cdot \left(16 + \frac{2K_{4}\alpha_{0}(\kappa)}{1 - K_{4}\alpha_{0}(\kappa)} + \frac{|\operatorname{supp}\gamma|}{2} \left(K_{4}\alpha_{0}(\kappa)\right)^{\max(0,\max(0,\min(\ell_{1},\ell_{2})-7))}\right) \cdot \sqrt{\frac{1}{|\operatorname{supp}\gamma|}} \\ & + \frac{4K_{3}(K_{4}\alpha_{0}(\kappa))^{\operatorname{dist}_{1}(\operatorname{supp}\gamma,\partial C_{1}(B_{N}))}}{\alpha_{5}(\beta,\kappa)} + K_{2} \cdot \frac{|\operatorname{supp}\gamma_{c}|}{|\operatorname{supp}\gamma|} \\ & + K_{3}K_{4}^{2}\alpha_{0}(\kappa)^{5/6} \cdot \alpha_{2}(\beta,\kappa) + \frac{18^{4}K_{5}\alpha_{2}(\beta,\kappa)^{5}}{2} \cdot \alpha_{2}(\beta,\kappa) \\ & + \sqrt{\frac{2K_{9}\alpha_{4}(\beta,\kappa)(K_{4}\alpha_{0}(\kappa))^{\operatorname{dist}_{1}(\operatorname{supp}\gamma,\partial C_{1}(B_{N}))}}{|\operatorname{supp}\gamma|\alpha_{5}(\beta,\kappa)^{2}}} \\ & + 2\sqrt{\frac{2K_{3}\alpha_{4}(\beta,\kappa)(K_{4}\alpha_{0}(\kappa))^{\operatorname{dist}_{1}(\operatorname{supp}\gamma,\partial C_{1}(B_{N}))}{|\operatorname{supp}\gamma|\alpha_{5}(\beta,\kappa)^{2}}} \\ & + \left(\sqrt{2K_{8}\alpha_{0}(\kappa)^{3}\max(\alpha_{0}(\kappa),\alpha_{1}(\beta)^{6})} + 2\sqrt{2K_{7}\alpha_{0}(\kappa)^{6}} + \sqrt{12K_{2}}\right) \cdot \sqrt{\frac{1}{|\operatorname{supp}\gamma|}} \\ & + 2^{-1}K_{14}\alpha_{2}(\beta,\kappa)^{5} \cdot \alpha_{2}(\beta,\kappa) + \left(\sqrt{K_{10}\alpha_{0}(\kappa)^{6}} + 1\right) \cdot \sqrt{\frac{|\operatorname{supp}\gamma_{c}|}{|\operatorname{supp}\gamma_{c}|}} \\ & + \sqrt[6]{\frac{K_{13}^{2}}{|\operatorname{supp}\gamma|}} \cdot \sqrt{\frac{1}{|\operatorname{supp}\gamma|}}. \end{split}$$

Now note that since γ is a path along the boundary of a rectangle with side lengths $\ell_1, \ell_2 \geq 2$, we must have $|\operatorname{supp} \gamma_c| \leq 8$. Since 3 holds, we must have $2\alpha_0(\kappa) \leq K_4\alpha_0(\kappa) \leq 1$, and since $G = \mathbb{Z}_2$, we have $\alpha_2(\beta, \kappa), \alpha_0(\beta) \leq 1$.

Recalling Proposition 2.16 and Proposition 2.19, letting $N \to \infty$, and simplifying, we thus obtain

$$\begin{split} \left| \left\langle L_{\gamma}(\sigma,\phi) \right\rangle_{\beta,\kappa,\infty} - \left\langle L_{\gamma}(\sigma,\phi) \right\rangle_{\infty,\kappa,\infty} \Theta_{\beta,\kappa}'(\gamma) \right|^{1+2|\operatorname{supp}\gamma|/|\operatorname{supp}(\gamma-\gamma_c)|} \\ &\leq 2^{2|\operatorname{supp}\gamma|/|\operatorname{supp}(\gamma-\gamma_c)|} K_{15} \left(\alpha_2(\beta,\kappa) + |\operatorname{supp}\gamma|^{-1/2} \right), \end{split}$$

where

$$K_{15} := \mathbb{1}(\partial \gamma \neq 0) \cdot \left(32K_3K_4 + \frac{4K_3K_9^4\alpha_0(\kappa)^8}{1 - K_4\alpha_0(\kappa)} + K_3K_4|\operatorname{supp}\gamma|^{1/2}\left(K_4\alpha_0(\kappa)\right)^{\min(\ell_1,\ell_2)}\right) \\ + 8K_2 + K_3K_4^2 + 18^4K_5/2 + \sqrt{K_8} + \sqrt{K_7} + \sqrt{12K_2} + K_{14}/2 + \sqrt{8K_{10}} + \sqrt{8} + \sqrt[3]{K_{13}}.$$
(10.40)

Next, note that since γ is a path along the boundary of some rectangle with side lengths $\ell_1, \ell_2 \ge 2$, we have $|\operatorname{supp} \gamma_c| \le 8$. Since, by assumption, we have $|\operatorname{supp} \gamma| \ge 24$, it follows that $|\operatorname{supp} \gamma_c| / |\operatorname{supp} \gamma| \le 1/3$, and hence

$$\frac{1}{4} \leq \frac{1}{1+2|\operatorname{supp} \gamma|/(|\operatorname{supp} \gamma| - |\operatorname{supp} \gamma_c|)} \leq \frac{1}{3}.$$

If we in addition have $\alpha_2(\beta, \kappa) + \sqrt{\max(1, |\operatorname{supp} \gamma_c|)/|\operatorname{supp} \gamma|} \le 1$, then it follows that

$$\left| \left\langle L_{\gamma}(\sigma,\phi) \right\rangle_{\beta,\kappa,\infty} - \left\langle L_{\gamma}(\sigma,\phi) \right\rangle_{\infty,\kappa,\infty} \Theta_{\beta,\kappa}'(\gamma) \right| \le 2^{1-\frac{1}{4}} \cdot K_{15}^{1/3}$$
$$\cdot \left(\alpha_{2}(\beta,\kappa) + \sqrt{1/|\operatorname{supp}\gamma|} \right)^{\frac{1}{4}}.$$
(10.41)

Since $|\rho(g)| = 1$ for all $g \in G$, we always have

$$\left|\left\langle L_{\gamma}(\sigma,\phi)\right\rangle_{\beta,\kappa,\infty}-\left\langle L_{\gamma}(\sigma,\phi)\right\rangle_{\infty,\kappa,\infty}\Theta_{\beta,\kappa}'(\gamma)\right|\leq 2.$$

Consequently, if $\alpha_2(\beta, \kappa) + \sqrt{|\operatorname{supp} \gamma_c|/|\operatorname{supp} \gamma|} \ge 1$, then (10.41) automatically holds. If we let

$$K_0 := 2^{\frac{3}{4}} K_{15}^{1/3}, \tag{10.42}$$

we thus obtain (1.2). This completes the proof of Theorem 1.1.

Acknowledgements The author is grateful to Jonatan Lenells and Fredrik Viklund for many useful discussions.

Funding Open access funding provided by Chalmers University of Technology. The author acknowledges support from the Knut and Alice Wallenberg Foundation, from the Swedish Research Council, Grant Agreement No. 2015-05430, and from the European Research Council, Grant Agreement No. 682537.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

References

- Balian, R., Drouffe, J.M., Itzykson, C.: Gauge fields on a lattice. I. General outlook. Phys. Rev. D 10(10), 3376–3395 (1974)
- Balian, R., Drouffe, J.M., Itzykson, C.: Gauge fields on a lattice. II. Gauge-invariant Ising model. Phys. Rev. D 11(8), 2098–2103 (1975)
- Balian, R., Drouffe, J.M., Itzykson, C.: Gauge fields on a lattice. III. Strong-coupling expansions and transition points. Phys. Rev. D 11(8), 2104–2119 (1975)
- 4. Borgs, C.: Translation symmetry breaking in four dimensional lattice gauge theories. Commun. Math. Phys. **96**, 251–284 (1984)
- Bricmont, J., Frölich, J.: An order parameter distinguishing between different phases pf lattice gauge theories with matter fields. Phys. Lett. B 122(1), 73–77 (1983)
- Bricmont, J., Frölich, J.: Statistical mechanical methods in particle structure analysis of lattice field theories (III). Confinement and bound states in gauge theories. Nucl. Phys. B280(FS18), 385–444 (1987)
- 7. Cao, S.: Wilson loop expectations in lattice gauge theories with finite gauge groups. Commun. Math. Phys. **380**, 1439–1505 (2020)
- Chatterjee, S.: A probabilistic mechanism for quark confinement. Commun. Math. Phys. 385(2021), 1007–1039 (2021)
- 9. Chatterjee, S.: Wilson loop expectations in Ising lattice gauge theory. Commun. Math. Phys. **377**, 307–340 (2020)
- Chernodubab, M.N., Ilgenfritz, E., Schiller, A.: String breaking and monopoles: a case study in the 3D abelian Higgs model. Phys. Lett. B 547(3–4), 269–277 (2002)
- 11. Creutz, M.: Phase diagrams for coupled spin-gauge system. Phys. Rev. D 21(4), 1106-1112 (1980)
- Duminil-Copin, H.: Lectures on the ising and potts models on the hypercubic lattice. In: Random Graphs, Phase Transitions, and the Gaussian Free Field. Springer Proceedings in Mathematics & Statistics, Springer, New York (2017)
- Evertz, H.G., Grösch, V., Jansen, K., Jersak, J., Kastrup, H.A., Neuhaus, T.: Confined and free charges in compact scalar QED. Neucl. Phys. B285(FS19), 559–589 (1987)
- Evertz, H.G., Jansen, K., Jersák, J., Lang, C.B., Neuhaus, T.: Photon and Bosonium masses is scalar lattice QED. Nucl. Phys. B 285, 590–605 (1987)
- Filk, T., Marcu, M., Fredenhagen, K.: Line of second-order phase transitions in the four-dimensional Z₂ gauge theory with matter fields. Phys. Lett. B 169(4), 405–412 (1986)
- Forsström, M.P.: Decay of correlations in finite abelian lattice gauge theories. Commun. Math. Phys. 393, 1311–1346 (2022)
- Forsström, M.P., Lenells, J., Viklund, F.: Wilson loops in finite abelian lattice gauge theories. Ann. Inst. H. Poincaré Probab. Statist. 58(4), 2129–2164 (2022)
- Forsström, M.P., Lenells, J., Viklund, F.: Wilson loops in the abelian lattice Higgs model. Probab. Math. Phys. 4(2), 257–329 (2023)
- Fradkin, E., Shenker, S.H.: Phase diagrams of lattice gauge theories with Higgs fields. Phys. Rev. D 19(12), 3682–3697 (1979)
- Fredenhagen, K., Marcu, M.: Dual interpretation of order parameters for lattice gauge theories with matter fields. Nucl. Phys. B (Proc. Suppl.) 4, 352–357 (1988)
- Garban, C., Supelveda, A.: Improved spin-wave estimate for Wilson loops in U(1) lattice gauge gauge theory. Int. Math. Res. Not 2023(21), 18142–18198 (2023)
- 22. Ginibre, J.: General formulation of Griffiths' inequalities. Commun. Math. Phys. 16, 310-328 (1970)
- Gliozzi, F.: The functional form of open Wilson lines in gauge theories coupled to matter. Nuclear Phys. B Proc. Suppl. 153(1), 120–127 (2006)
- Gliozzi, F., Rago, A.: Monopole clusters, center vortices, and confinement in a Z2 gauge-Higgs system. Phys. Rev. D 66, 074511 (2002)
- Gregor, K., Huse, D.A., Moessner, R., Sondhi, S.L.: Diagnosing deconfinement and topological order. New J. Phys. 13, 025009 (2011)
- Jaffe, A., Witten, E.: Quantum Yang-Mills Theory. https://www.claymath.org/sites/default/files/ yangmills.pdf
- Jongeward, G.A., Stack, J.D., Jayaprakash, C.: Monte Carlo calculations on Z₂ gauge-Higgs theories. Phys. Rev. D 21(12), 3360–3368 (1980)
- Kanaya, K., Sugiyama, Y.: Meanfield Study of Z₂ Higgs model with radial excitations and mode correlation problem. Prog. Theor. Phys. 72(6), 1158–1175 (1984)
- 29. Marcu, M.: (Uses of) An Order Parameter for Lattice Gauge Theories with Matter Fields
- 30. Shrock, E.: Lattice Higgs models. Nucl. Phys. B (Proc. Suppl.) 4, 373-389 (1988)

- Wegner, F.J.: Duality in Generalized Ising Models and Phase Transitions without Local Order Parameters. J. Math. Phys. 12(10), 2259–2272 (1971)
- 32. Wilson, K.: Confinement of quarks. Phys. Rev. D 10, 2445–2459 (1974)

Communicated by S. Chatterjee