



**CHALMERS**  
UNIVERSITY OF TECHNOLOGY

## **Quasi-free (p,2p) reactions in inverse kinematics for studying the fission yield dependence on temperature**

Downloaded from: <https://research.chalmers.se>, 2024-12-20 15:34 UTC

Citation for the original published paper (version of record):

Graña-González, A., Rodriguez-Sanchez, J., Benlliure, J. et al (2023). Quasi-free (p,2p) reactions in inverse kinematics for studying the fission yield dependence on temperature. *Proceedings of Science*, 419. <http://dx.doi.org/10.22323/1.419.0017>

N.B. When citing this work, cite the original published paper.

## Quasi-free (p,2p) reactions in inverse kinematics for studying the fission yield dependence on temperature

A. Graña-González,<sup>1</sup> J. L. Rodríguez-Sánchez,<sup>1</sup> J. Benlliure,<sup>1</sup> G. García-Jiménez,<sup>1</sup> H. Alvarez-Pol,<sup>1</sup> D. Cortina-Gil,<sup>1</sup> L. Atar,<sup>2</sup> L. Audouin,<sup>3</sup> G. Authelet,<sup>4</sup> A. Besteiro,<sup>5</sup> G. Blanchon,<sup>6</sup> K. Boretzky,<sup>7</sup> P. Cabanelas,<sup>1</sup> E. Casarejos,<sup>5</sup> J. Cederkall,<sup>8</sup> A. Chatillon,<sup>6</sup> A. Corsi,<sup>4</sup> M. Feijoo,<sup>1</sup> D. Galaviz,<sup>9</sup> I. Gasparic,<sup>10</sup> R. Gernhäuser,<sup>11</sup> M. Heil,<sup>7</sup> A. Heinz,<sup>12</sup> M. Holl,<sup>12</sup> T. Jenegger,<sup>11</sup> L. Ji,<sup>2</sup> H.T. Johansson,<sup>12</sup> O.A. Kiselev,<sup>7</sup> P. Klenze,<sup>11</sup> A. Knyazev,<sup>8</sup> D. Körper,<sup>7</sup> T. Kröll,<sup>2</sup> I. Lihtar,<sup>10</sup> Y. A. Litvinov,<sup>7</sup> B. Löher,<sup>7</sup> P. Morfouace,<sup>6</sup> D. Mücher,<sup>13,14</sup> S. Murillo Morales,<sup>15</sup> A. Obertelli,<sup>2</sup> V. Panin,<sup>7</sup> J. Park,<sup>16</sup> S. Paschalis,<sup>15</sup> A. Perea,<sup>17</sup> M. Petri,<sup>15</sup> S. Pirrone,<sup>18</sup> L. Ponnath,<sup>11</sup> A. Revel,<sup>6</sup> H.-B. Rhee,<sup>2</sup> L. Rose,<sup>15</sup> D.M. Rossi,<sup>2</sup> P. Russotto,<sup>18</sup> H. Simon,<sup>7</sup> A. Stott,<sup>15</sup> Y. Sun,<sup>2</sup> C. Sürder,<sup>2</sup> J. Taieb,<sup>6</sup> R. Taniuchi,<sup>15</sup> O. Tengblad,<sup>17</sup> H.T. Törnqvist,<sup>2,7</sup> S. Velardita,<sup>2</sup> J. Vesic,<sup>19</sup> B. Voss<sup>7</sup> and the R<sup>3</sup>B collaboration.

<sup>1</sup>IGFAE, University of Santiago de Compostela, E-15782 Santiago de Compostela, Spain

<sup>2</sup>Technische Universität Darmstadt, Fachbereich Physik, Institut für Kernphysik, 64289 Darmstadt, Germany

<sup>3</sup>IPN Orsay, 15 rue Georges Clemenceau, 91406, Orsay, France

<sup>4</sup>CEA Saclay, IRFU/DPhN, Centre de Saclay, 91191, Gif-sur-Yvette, France

<sup>5</sup>CINTECX, Universidade de Vigo, E-36200 Vigo, Spain

<sup>6</sup>CEA DAM,DIF, F-91297 Arpajon, France

<sup>7</sup>GSI Helmholtzzentrum für Schwerionenforschung, Planckstraße 1, 64291, Darmstadt, Germany

<sup>8</sup>Department of Physics, Lund University, P.O. box 118, 221 00 Lund, Sweden

<sup>9</sup>LIP and Faculty of Sciences, University of Lisbon, 1000-149 Lisbon, Portugal

<sup>10</sup>RBI Zagreb, Bijenicka cesta 54, HR10000, Zagreb, Croatia

<sup>11</sup>Technische Universität München, James-Frank-Str 1, 85748, Garching, Germany

<sup>12</sup>Institutionen för Fysik, Chalmers Tekniska Högskola, 412 96 Göteborg, Sweden

<sup>13</sup>University of Guelph, 50 Stone Road E, N1G 2W1, Guelph, ON, Canada

<sup>14</sup>Institut für Kernphysik der Universität zu Köln, Zùlpicher Strasse 77, D-50937 Köln, Germany

<sup>15</sup>School of Physics, Engineering and Technology, University of York, YO10 5DD York, UK

<sup>16</sup>Institute for Basic Science, Center for Exotic Nuclear Studies, 34126, Daejeon, Korea (Republic of)

<sup>17</sup>Instituto de Estructura de la Materia, CSIC, E-28006 Madrid, Spain

<sup>18</sup>INFN Laboratori Nazionali del Sud, Via Santa Sofia 62, 95123, Catania, Italy

<sup>19</sup>Jozef Stefan Institute, Jamova cesta 39, 1000 Ljubljana, Slovenia

E-mail: [antia.grana.gonzalez@usc.es](mailto:antia.grana.gonzalez@usc.es)

FAIR next generation scientists - 7th Edition Workshop (FAIRness2022)

23-27 May 2022

Paralia (Pieria, Greece)

Despite the recent experimental and theoretical progress in the investigation of the nuclear fission process, a complete description still represents a challenge in nuclear physics because it is a very complex dynamical process, whose description involves the coupling between intrinsic and collective degrees of freedom, as well as different quantum-mechanical phenomena. To improve on the existing data on nuclear fission, we produce fission reactions of heavy nuclei in inverse kinematics by using quasi-free (p,2p) scattering, which induce fission through particle-hole excitations that can range from few to ten's of MeV. The measurement of the four-momenta of the two outgoing protons allows to reconstruct the excitation energy of the fissioning nucleus and therefore to study the evolution of the fission yields with temperature. The realization of this kind of experiment requires a complex experimental setup, providing full isotopic identification of both fission fragments and an accurate measurement of the momenta of the two outgoing protons. This was realized recently at the GSI/FAIR facility and here some preliminary results are presented.

## 1. Introduction


Nuclear fission plays an important role in the nucleosynthesis r-process, which was indicated as the mechanism responsible for the production of the heaviest elements in the universe. Therefore, fission studies in the neutron-rich region are required in order to measure fission yields and fission barrier heights to improve r-process calculations [1, 2].

In the last decade, unprecedented fission experiments have been carried out at the GSI/FAIR facility using the inverse-kinematics technique in combination with state-of-the-art detectors. For the first time in the history of fission, it was possible to simultaneously measure and identify both fission fragments in terms of their mass and atomic numbers [3, 4] and to extract correlations between fission observables sensitive to the dynamics of the fission process [5] and the nuclear structure at the scission point [6, 7]. However, a limitation of those previous experiments was the impossibility of measuring the excitation energy of the nucleus undergoing fission, an observable that plays a fundamental role in the description of the process. Depending on the amount of available energy, different regions of the potential landscape can be populated, leading to different fission paths and consequently to different fission yields distributions. The gradual weakening of the influence of shell structure on the fission yields with the increasing of excitation energies can be quantified with the so-called suppression function [8], which suppresses the microscopic term of the energy-dependent effective potential. However, the model parameters used in such an approach are still not constrained due to the lack of data at different excitation energies. This lack of data is caused by the use of Coulomb excitations to induce fission of exotic nuclei, where the Giant Dipole Resonance (GDR) is excited at  $\sim 11$  MeV and whose distribution has a width of few MeV. On the other hand, Transfer- and fusion-fission reactions give access to different excitation energies, but they are limited to stable nuclei. The experiment of this work went further by combining fission with quasi-free (p,2p) reactions, which allows for the first time to measure many excitation energies with stable and exotic nuclei providing therefore very valuable data to constrain the suppression function parameters. Another interesting question is the possible dependence of the energy sorting of the fission fragments on the excitation energies of the fissioning nucleus, since in the studies in direct kinematics with neutrons, the well known saw-tooth shape shows higher neutron multiplicities for higher incident neutron energies [9].

In this work, the main scientific goal is to investigate the energy dependence of the fission yields. Such data can be obtained by using quasi-free (p,2p) reactions and combining the previous experimental setup with a silicon tracker based on AMS-type detectors [10] and the calorimeter CALIFA (CALorimeter for In-Flight detection of gamma-rays and high energy charged pArticles) [11] developed by the R<sup>3</sup>B collaboration. Here we report on the analysis of the first fission experiment performed at the GSI/FAIR accelerator facilities in Darmstadt (Germany) using the quasi-free (p,2p) reactions as mechanism to induce fission of <sup>238</sup>U projectiles.

## 2. Experiment

The experiment was performed in March 2021 by using primary beams of <sup>238</sup>U at 560A MeV delivered by the SIS18 synchrotron, which were guided to the experimental area to impinge on a 15mm thick liquid hydrogen target in order to produce quasi-free (p,2p) reactions. The thickness of the target has been chosen in order to minimise the energy loss of the protons in the medium and the  $\delta$ -electrons production. The experiment is based on the inverse-kinematics technique, in which the heavy nucleus undergoing fission is the projectile. This methodology was developed at GSI and represents a major breakthrough in the characterization of the fission fragments. The major advantage is that the fission fragments are emitted in a narrow cone in the forward direction with very high velocities, permitting to detect both fragments simultaneously and to measure the isotopic yields.



**Figure 1:** Schematic view of the experimental setup used in the present work to study quasi-free (p,2p) induced fission reactions.


Using this methodology, previous fission experiments at GSI already provided for the first time the complete isotopic identification of both fission fragments. In the (p,2p)-fission experiment the fragments are identified by using the same detectors. The experimental setup (see the Fig 1) consists of several multi-wire proportional chambers (MWPCs) to track the trajectories before and

after the superconducting dipole GLAD, a Twin-MUSIC (Multi-Sampling Ionization Chamber) divided into two sections which are segmented into 16 anodes to measure the charges of the fission fragments and a ToF-Wall composed by 28 plastic scintillators to measure the time of flight. With all these measurements, employing the  $\Delta E - B\rho - ToF$  technique, the mass of the fission fragments can be obtained, allowing therefore for full isotopic identification. Neutrons emitted from the fission fragments are detected in NeuLAND (New Large-Area Neutron Detector) [12], located 15m downstream from the target.

Since our goal is to study the energy dependence of the fission yields, we access the excitation energy via the aforementioned quasi-free (p, 2p) reactions which are knockout reactions between protons of the liquid hydrogen target and protons bound in the heavy projectile nucleus. The products of the reaction are the two outgoing protons and the excited nucleus characterized by the particle-hole excitations created by the removed proton. This excited nucleus can then de-excite via fission or particle evaporation. From the measurement of the momenta of the two outgoing protons the excitation energy of the fissioning nucleus can be reconstructed using the missing energy method [13]. In order to obtain the four-momenta, the energy and the tracking of the protons are required. To reconstruct the protons trajectory a silicon tracker is located in front of the target, which was equipped with two double-plane arms consisting of an array of three AMS-type [10] 0.3 mm thick double-sided silicon-strip detectors. Then to measure the energy of the protons the CALIFA calorimeter [11] is surrounding the whole, which consists of 1504 CsI(Tl) crystal scintillators covering a polar angle range between 22 and 90 degrees.

### 3. Results

The first step of the analysis is to obtain the charge identification of both fission fragments given by the Twin-MUSIC, which is designed with two independent ionization chambers divided



**Figure 2:** Left: Two-dimensional cluster plot of the charges  $Z_1$  and  $Z_2$  of both fission fragments for all fissioning systems. The inset shows the fission fragments distribution for  $Z_1 + Z_2 = 91$ . Right: Fissioning systems distribution.

into two sections in order to measure the atomic number of both fission fragments simultaneously. The correlation of the derived charge for both fission fragments  $Z_1$  and  $Z_2$  is shown in the Fig. 2-left in a two-dimensional cluster plot. Each diagonal of clusters in this figure corresponds to a different fissioning system, and the fissioning system distribution is displayed in the Fig. 2-right. Since the beam is  $^{238}_{92}\text{U}$ , when the desired (p,2p) takes place the resulting nucleus is an excited  $^{237}_{91}\text{Pa}$ , which contributes to the  $Z_1 + Z_2 = 91$  peak, the one with the highest statistics of this plot. Nevertheless, it is important to notice that this peak also has contributions from other lighter Pa isotopes produced by the removal of one proton and several neutrons during the proton-nucleus collision, resulting in nuclear systems with higher excitation energies. Here, the kinematics of the two outgoing protons can be used to separate these types of contributions. The other two peaks to the right of the  $Z_1 + Z_2 = 91$  correspond to reactions in which neutrons might have been kicked out of the projectile nucleus, but no proton was removed. In particular, the systems with charge 93 correspond to the less likely case in which the projectile also captures one proton from the target. The peaks to the left of  $Z_1 + Z_2 = 91$  come from fissioning systems in which more than one proton was removed from the projectile.

The inset of the Fig. 2-left shows the fission fragment distribution for the production of  $_{91}\text{Pa}$ , with a charge resolution of 0.38 units (FWHM). The distribution fits well to an overall Gaussian shape, although two small humps around  $Z=40$  and  $Z=50$  are slightly noticeable. These two peaks are the asymmetric fission contribution and they appear because of the shell effects [14] which favour the formation of nuclei with magic numbers around  $Z=50$  in this case. Without any constrain on the energy (like in this plot) the symmetric contribution from the high energies will partially shadow the asymmetric humps and the total distribution will be the convolution of the three Gaussian distributions. Nevertheless, when this plot is constrained to low energies, the symmetric component should gradually vanish, since the fission barrier in  $_{91}\text{Pa}$  is supposed to be higher for symmetric fission. Therefore, the asymmetric distribution would start to be more and more noticeable and at some point it would be the only visible one. Once the excitation energy is reconstructed, it will be possible to plot this same graph for different excitation energies and to study the evolution of the fission yields from asymmetric to symmetric while going from low to high energies.

#### 4. Conclusions

The first (p,2p)-fission experiment has been carried out at GSI/FAIR with projectiles of  $^{238}\text{U}$  at energies of 560A MeV impinging on a liquid hydrogen target and using the R<sup>3</sup>B experimental setup to measure all the reaction products. These complete kinematics measurements will allow to correlate the fission yields with the excitation energy of the fissioning nucleus and to study the energy sharing between the two fission fragments. Additionally, this experiment permits to develop the data analysis and validate the methodology to use this approach in future experiments with exotic nuclei. The current state of the analysis is that the charge identification is completed and the next steps will be the identification in mass number and the reconstruction of the excitation energy spectrum. In the near future further (p,2p)-fission experiments are planned to be carried out at GSI/FAIR with exotic neutron-rich projectiles close to the neutron shell  $N=152$  to obtain their fission yields and fission barrier heights for constraining r-process calculations.

## 5. Acknowledgments

This work was partially supported by the Spanish Ministry for Science and Innovation under grants PGC2018-099746-B-C21 and PGC2018-099746-B-C22, by the Regional Government of Galicia under the program “Grupos de Referencia Competitiva” ED431C-2021-38 and by the “María de Maeztu” Units of Excellence program MDM-2016-0692. The experiment has been performed at GSI facilities within the framework of FAIR Phase-0. J.L.R.S. thanks the support from Xunta de Galicia under the postdoctoral fellowship ED481D-2021-018.

## References

- [1] N. Vassh et al. 2019 J. Phys. G: Nucl. Part. Phys. **46** 065202
- [2] Samuel A. Giuliani et al. 2018 Phys. Rev. C **97** 034323
- [3] J. L. Rodríguez-Sánchez et al. 2015 Phys. Rev. C **91** 064616
- [4] E. Pellereau et al. 2017 Phys. Rev. C **95** 054603
- [5] J. L. Rodríguez-Sánchez et al. 2016 Phys. Rev. C **94** 061601(R)
- [6] A. Chatillon et al. 2019 Phys. Rev. C **99** 054628
- [7] A. Chatillon et al. 2020 Phys. Rev. Lett. **102** 202502
- [8] J. Randrup et al. 2013 Phys. Rev. C **88** 064606
- [9] A.N. Andreyev et al. 2017 Reports Prog. Phys. **81** 016301
- [10] J. Alcaraz et al. 2008 Nucl. Instrum. Methods Phys. Res. A **593** 376
- [11] H. Alvarez Pol et al. 2014 Nucl. Instrum. Methods Phys. Res. A **767** 453
- [12] K. Boretzky et al. 2021 Nucl. Instrum. Method Phys. Res. A **1014** 165701
- [13] T. Noro et al. 2020 Prog. Theor. Exp. Phys. **093D02**
- [14] K.-H. Schmidt et al. 2000 Nucl. Phys. A **665** 221