



The salty emission of the intermediate-mass AGB star OH 30.1-0.7

Downloaded from: <https://research.chalmers.se>, 2026-04-14 23:13 UTC

Citation for the original published paper (version of record):

Danilovich, T., Richards, A., Van de Sande, M. et al (2025). The salty emission of the intermediate-mass AGB star OH 30.1-0.7. *Monthly Notices of the Royal Astronomical Society*, 536(1): 684-713. <http://dx.doi.org/10.1093/mnras/stae2584>

N.B. When citing this work, cite the original published paper.

The salty emission of the intermediate-mass AGB star OH 30.1–0.7

T. Danilovich^{1,2,3★}, A. M. S. Richards,⁴ M. Van de Sande⁵, C. A. Gottlieb,⁶ T. J. Millar⁷,
A. I. Karakas^{1,2}, H. S. P. Müller⁸, K. Justtanont,⁹ J. M. C. Plane¹⁰, S. Etoka⁴, S. H. J. Wallström,³
L. Decin,^{3,10} D. Engels,¹¹ M. A. T. Groenewegen¹², F. Kerschbaum,¹³ T. Khouri,⁹ A. de Koter,^{3,14}
H. Olofsson,⁹ C. Paladini¹⁵ and R. J. Stancliffe¹⁶

¹School of Physics & Astronomy, Monash University, Wellington Road, Clayton 3800, Victoria, Australia

²ARC Centre of Excellence for All Sky Astrophysics in 3 Dimensions (ASTRO 3D), Clayton 3800, Australia

³Institute of Astronomy, KU Leuven, Celestijnenlaan 200D, B-3001 Leuven, Belgium

⁴JBCA, Department Physics and Astronomy, University of Manchester, Manchester M13 9PL, UK

⁵Leiden Observatory, Leiden University, PO Box 9513, NL-2300 RA Leiden, The Netherlands

⁶Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, USA

⁷Astrophysics Research Centre, School of Mathematics and Physics, Queen's University Belfast, University Road, Belfast BT7 1NN, UK

⁸Astrophysik/I. Physikalisches Institut, Universität zu Köln, D-50937 Köln, Germany

⁹Department of Space, Earth and Environment, Chalmers University of Technology, Onsala Space Observatory, SE-43992 Onsala, Sweden

¹⁰School of Chemistry, University of Leeds, Leeds LS2 9JT, UK

¹¹Hamburger Sternwarte, Universität Hamburg, Gojenbergsweg 112, D-21029 Hamburg, Germany

¹²Koninklijke Sterrenwacht van België, Ringlaan 3, B-1180 Brussels, Belgium

¹³Department of Astrophysics, University of Vienna, Türkenschanzstrasse 17, A-1180 Vienna, Austria

¹⁴Anton Pannekoek Institute for Astronomy, University of Amsterdam, 1090 GE Amsterdam, the Netherlands

¹⁵European Southern Observatory, Alonso de Cordova 3107, Vitacura, Santiago, Chile

¹⁶H.H. Wills Physics Laboratory, University of Bristol, Tyndall Avenue, Bristol BS8 1TL, UK

Accepted 2024 November 12. Received 2024 November 11; in original form 2024 August 30

ABSTRACT

We analyse continuum and molecular emission, observed with Atacama Large Millimetre/submillimetre Array, from the dust-enshrouded intermediate-mass asymptotic giant branch (AGB) star OH 30.1–0.7. We find a secondary peak in the continuum maps, ‘feature B’, separated by 4.6 arcsec from the AGB star, which corresponds to a projected separation of 1.8×10^4 au, placing a lower limit on the physical separation. This feature is most likely composed of cold dust and is likely to be ejecta associated with the AGB star, though we cannot rule out that it is a background object. The molecular emission we detect includes lines of CO, SiS, CS, SO₂, NS, NaCl, and KCl. We find that the NS emission is off centre and arranged along an axis perpendicular to the direction of feature B, indicative of a UV-emitting binary companion (e.g. a G-type main sequence star or hotter), perhaps on an eccentric orbit, contributing to its formation. However, the NaCl and KCl emission constrain the nature of that companion to not be hotter than a late B-type main-sequence star. We find relatively warm emission arising from the inner wind and detect several vibrationally excited lines of SiS ($v = 1$), NaCl (up to $v = 4$), and KCl (up to $v = 2$), and emission from low-energy levels in the mid to outer envelope, as traced by SO₂. The CO emission is abruptly truncated around 3.5 arcsec or 14 000 au from the continuum peak, suggesting that mass loss at a high rate may have commenced as little as 2800 yr ago.

Key words: stars: AGB and post-AGB – circumstellar matter – stars: individual: OH 30.1–0.7.

1 INTRODUCTION

The asymptotic giant branch (AGB) phase is a late evolutionary stage that low- and intermediate-mass stars (with initial masses $0.8 M_{\odot} \lesssim M_i \lesssim 8 M_{\odot}$) pass through towards the end of their lives. During this phase, the stars eject a significant amount of material through a stellar wind, with mass-loss rates in the range $\sim 10^{-8}$ to $10^{-4} M_{\odot} \text{yr}^{-1}$. Some lower mass stars, with initial masses ~ 1.5 to $4 M_{\odot}$, are expected to end their lives as carbon stars, after dredging

up newly synthesized carbon from their interiors to their surfaces (Herwig 2005; Karakas & Lattanzio 2014). Intermediate-mass stars, with initial masses in the range ~ 4 to $8 M_{\odot}$, are not expected to end their lives as carbon stars (Herwig 2005; Karakas & Lattanzio 2014; Ventura et al. 2018; Marigo et al. 2020) because they are massive enough to initiate hot bottom burning (HBB), a phenomenon where the star’s convective envelope penetrates the hydrogen-burning shell and enables additional (proton-capture) nucleosynthesis processes (Karakas & Lattanzio 2014). In particular, HBB destroys ¹²C and ¹⁸O. The destruction of ¹²C prevents the formation of carbon stars and results in low ¹²C/¹³C ratios, and the destruction of ¹⁸O results in very large ¹⁷O/¹⁸O ratios, allowing us to observationally identify

* E-mail: taissa.danilovich@monash.edu

stars in which HBB has taken place (Justtanont et al. 2015). We also note that the precise mass ranges of stars that become carbon stars or which undergo HBB depends on their initial metallicity and on some numerical details of the models (such as the treatment of convection and mass-loss rates, Karakas & Lugaro 2016). The numbers given here are for solar metallicity.

A small sample of stars have been shown to have intermediate initial masses based on the prevalence of H_2^{17}O emission and the absence of H_2^{18}O emission in *Herschel* observations (Justtanont et al. 2015). Some of these stars have also been found to have low $^{12}\text{C}/^{13}\text{C}$ ratios (e.g. Delfosse, Kahane & Forveille 1997), providing further evidence for HBB. These intermediate-mass stars have all previously been found to have high mass-loss rates, though the precise values of the mass-loss rates have been the subject of much debate since different values are found using different methods (see e.g. Heske et al. 1990; Justtanont et al. 2006). Several explanations have been put forward to explain these discrepant results, including the concept of a ‘superwind’: a recently initiated period of enhanced mass loss (Heske et al. 1990; Delfosse et al. 1997; Justtanont et al. 2013). This scenario could explain the relatively high mass-loss rates obtained from spectral energy distributions (SEDs) and dust modelling, thought to represent the more recent mass loss, while the lower mass-loss rates from CO are thought to be the remnant of an earlier period of lower mass loss.

Thus far, no studies of the circumstellar chemistry of confirmed intermediate-mass AGB stars have been presented. In this work, we focus on the intermediate-mass AGB star OH 30.1–0.7, thought to have a very high mass-loss rate ($> 10^{-4} \text{ M}_{\odot}\text{yr}^{-1}$ based on SED modelling, Justtanont et al. 2013, 2015). This star has featured in several other studies of extremely dusty stars based on various (spatially unresolved) infrared and radio observations (Heske et al. 1990; Delfosse et al. 1997; Justtanont et al. 2006). Although several previous studies have considered OH 30.1–0.7 to have a superwind, Decin et al. (2019) argued for a binary scenario rather than a superwind.

Atacama Large Millimetre/submillimetre Array (ALMA) observations of CO around two intermediate-mass stars (OH 30.1–0.7 and OH 26.5+0.6) were presented by Decin et al. (2019) who attributed structures in their circumstellar envelopes (CSEs) to interactions with binary companions. Decin et al. (2019) further argue that the apparently high mass-loss rates could be explained by an equatorial density enhancement of gas and dust arising from gravitational shaping by a companion. This would then change the apparent mass-loss rate depending on the orientation of the system and the line of sight through the CSE. Certainly some of the structures observed in the winds of both of these stars are similar to structures seen around other AGB stars that are also attributed to binary interactions (e.g. Cernicharo et al. 2015; Kim et al. 2017; Decin et al. 2020).

For AGB stars more generally, it has recently been shown that binary companions, especially solar-like stars emitting notable UV flux, can have an impact on circumstellar chemistry (Van de Sande & Millar 2022). Observational studies have found some molecular emission distributions that are best explained through the presence of a solar-type companion (Siebert et al. 2022; Coenegrachts et al. 2023; Danilovich et al. 2024). In particular, Danilovich et al. (2024) showed that asymmetric distributions of SiN, SiC, and NS could be attributed to a main-sequence companion on a highly eccentric orbit passing through the inner wind of W Aql every ~ 1100 yr. Coenegrachts et al. (2023) found asymmetric clumps of NaCl emission forming an approximate spiral in the inner wind of the oxygen-rich AGB star IK Tau, which they attributed to an unseen companion close to the AGB star.

We present here a complete molecular inventory of the existing ALMA data of OH 30.1–0.7 (also known as V1362 Aql, IRAS 18460–0254, and AFGL 5535), where previously only CO observations were published. In Section 2, we describe our re-reduction of the ALMA data, present the continuum, and describe our analysis of the molecular lines. In Section 3, we present the detected molecular lines and discuss the implications of these results in Section 4, in particular how these observations support the presence of a companion star. We summarize our conclusions in Section 5.

2 OBSERVATIONS

OH 30.1–0.7 was observed with two ALMA receivers, Bands 3 and 6, in 2016 and 2017, as part of project 2016.1.00005.S. The frequency ranges, restoring beams, observation dates, and ALMA configurations are given for each tuning in Table A1. The Band 3 data were observed with a single ALMA configuration (angular resolution ~ 1.3 arcsec), whereas the Band 6 data were observed with two antenna configurations and, for convenience, we refer to these as the extended (angular resolution ~ 0.3 arcsec) and compact (angular resolution ~ 1.5 arcsec) configurations. The Band 3 data were observed with velocity resolutions of 2.6 to 2.9 km s^{-1} (1 MHz) and the Band 6 data were observed with a velocity resolution around 2.5 km s^{-1} (2 MHz). More precise velocity and frequency resolutions are noted in Table A1 for each tuning. At the time of the ALMA observations, the star was close to minimum light (phase $\phi \sim 0.45$) in its variability cycle.

2.1 Data reduction

2.1.1 Imaging

Rather than using the data products from the standard ALMA pipeline, we performed an additional self-calibration for OH 30.1–0.7. First we downloaded the standard ALMA data products for project 2016.1.00005.S from the ALMA Science Archive (Stoehr et al. 2017) and reconstituted the calibrated measurement sets for each array configuration and spectral tuning. We mainly followed the data processing steps as listed in section 3.2.1 of Gottlieb et al. (2022) for the ATOMIUM Large Programme (ALMA Tracing the Origins of Molecules from InG dUst in oxygen-rich M-type stars). In summary, we split out each spectral window of the calibrated target data, adjusting the frequencies to constant velocity with respect to the local standard of rest (v_{LSR}) in the target direction. We identified the line-free channels by inspection of the visibility spectra and (where available) the archive example image cubes. We made continuum-only images for each configuration and used these as a starting model for self-calibration. Continuum subtracted image cubes were made for each spectral window as listed in Table A1. The Band 6 data for OH 30.1–0.7 were observed in two configurations with equivalent spectral tunings, and these were combined as described in section 3.2.3 of Gottlieb et al. (2022) to make combined continuum images and combined spectral cubes.

The details of the various continuum images we made are given in Table A2. These include continua for each individual configuration, the combined images created using different weightings, and an additional Band 3 image created with the same beam parameters as the compact Band 6 continuum image, to allow for a more direct comparison, as discussed below. In all imaging, Briggs weighting (Briggs, Schwab & Sramek 1999) was used with robust 0.5 to interpolate across unsampled regions of the visibility plane, giving more weight to cells closer to measured samples. In combining

data from extended and compact configurations, we gave both data sets equal weight which gives a full resolution close to that of the extended configuration but with better sensitivity to large angular scales. However, the sampling of the visibility plane is sparse for the longest baselines, which can lead to slight artefacts. To mitigate this, we applied a Gaussian taper to the visibility weights as a function of projected baseline length to reduce the weight of data from the longest baselines. The FWHM (full width at half-maximum) of the taper is selected to correspond to a slightly smaller angular resolution than the desired synthesized beam. This gives a coarser resolution but reduces errors due to unsampled scales. The resulting beam sizes and tapers for the combined continuum data are given in Table A2. No taper was used for the line data and beam sizes for those cubes are given in Table A1.

2.1.2 Accuracy

The factors contributing to absolute flux scale and astrometric accuracy are described in the ALMA Technical Handbook (Cortes et al. 2023), as applied to similar data in section 3.2.2 of Gottlieb et al. (2022). We matched the continuum flux densities on baselines of similar length for the two Band 6 configurations to remove relative flux errors to the level of the noise (see Table A2). The ALMA flux scale in Band 6, for a single observation, is accurate to 5 per cent, or up to ~ 10 per cent for combined images, and somewhat better in Band 3.

After fitting a 2D Gaussian component to a compact source, the relative (stochastic) position error is proportional to the beam size divided by the signal-to-noise ratio (S/N). The constant of proportionality is 0.5 for a well-filled array (Condon 1997), rising for sparser uv coverage such as ALMA at higher frequencies and a more extended configuration. This gives stochastic errors for the position of the stellar peaks of ~ 12 , ~ 7 , and ~ 1 mas, for Band 3 and the Band 6 lower and higher resolutions, respectively (using the values in Table A2 and constants of 0.5, 0.75, and 1). These are relevant for relative separations e.g. from a putative companion. The phase-reference source was J1851+0035, 3.5 arcsec from the target. The main contribution to astrometric uncertainty is errors in transferring the phase reference solutions to the target (approximately [phase rms]/360 \times beam). The phase rms on a point source was about 25 $^\circ$, 15 $^\circ$, and 10 $^\circ$ at Band 3 and the Band 6 compact and extended configurations, respectively. This leads to astrometric errors of approximately [beam size]/15, /25, and /35, respectively. Using the mean beam sizes (Table A2), this corresponds to astrometric errors of approximately 90, 60, and 10 mas for Band 3 and the Band 6 lower and higher resolutions, respectively. This is in addition to S/N (stochastic) errors; other contributions such as antenna position errors are negligible except at the highest resolution, giving total astrometric errors for the continuum peaks of about 91, 63, and 10 mas, respectively. The position of OH 30.1–0.7 based on our observations, using a weighted mean and taking these astrometric uncertainties into account, is given in Table 1. We find that the measured positions are offset from the observing positions (also given in Table 1) by ~ 0.6 arcsec in each coordinate.

2.2 Continua

In Fig. 1, we plot the continuum images for each configuration, and the combined Band 6 continuum image. The central continuum emission is resolved in the Band 6 images, and we detect some

Table 1. Sky positions and key parameters of OH 30.1–0.7.

	Right ascension	Declination
Observed position B6	18:48:41.91	–2:50:28.3
Observed position B3	18:48:41.91	–2:50:28.3
Band 6 extended ^a	18:48:41.94961	–2:50:28.9241
Band 6 compact ^a	18:48:41.94932	–2:50:28.8653
Band 3 ^a	18:48:41.95096	–2:50:28.8333
Mean position ^b	18:48:41.9496 ± 1.3 mas	–2:50:28.921 ± 7.8 mas
Distance ^c	3900 \pm 800 pc	
v_{LSR} ^d	100.3 \pm 1.2 km s ^{–1}	

Notes. ^a Astrometric positions in ICRS J2000 obtained from fits to the continuum peaks in this work. Uncertainties as described in Section 2.1.2. ^b The weighted mean position, assuming negligible proper motion between epochs at the distance of OH 30.1–0.7, with the associated uncertainty calculated from the standard deviation. ^c Distance calculated using the maser phase-lag method by Engels et al. (2015). ^d LSR velocity calculated in Section 2.5.

extended emission. The red crosses in Fig. 1 indicate the peaks of the continuum emission, which we found through fitting a 2D Gaussian to the extended configuration continua using the CASA routine `imfit`. In the continuum plots (and all subsequent plots of molecular emission), the data are normalized such that the continuum peaks lie at (0,0). The precise location of this peak, in J2000 coordinates, is given in Table 1 as are the positions measured from the other ALMA configurations. We find that the slight differences between positions are close to the predicted astrometric errors.

Considering the combined Band 6 continuum emission, the 5σ contour shows some ellipticity with a major axis running from southeast to northwest and varying in extent from 0.57 to 0.73 arcsec. The 3σ contour has a less regular shape, encompassing more faint emission with a greater extent of 1.06 arcsec to the southeast and a minimum extent of 0.64 arcsec to the northwest. These measurements consider only regions of flux enclosing the continuum peak and neglect isolated islands of emission. We use a similar method to measure extents (including the extents of molecular emission) as that described in Danilovich et al. (2021) for angular sizes. Based on the distance of 3900 pc (Engels et al. 2015), these continuum extents correspond to dust at projected separations of 2200–4100 au from the continuum peak of OH 30.1–0.7. Although we mainly analysed the combined (equal weighting) continuum, we find essentially identical results when considering the continuum from the extended array only.

2.2.1 Secondary peak: feature B

We detect a secondary peak to the north of the primary peak, with a certainty $> 10\sigma$. This lies well beyond the extent of the emission centred on the primary peak. As can be seen in Fig. 1, there are some less intense regions of continuum emission (detected at $> 3\sigma$) lying between the primary and secondary peaks, hinting at a connection between them. To check whether the secondary peak is truly associated with the AGB star, rather than being a coincidental background or foreground source, we checked the individual configuration data observed with Band 6 and the Band 3 data. The secondary peak is clearly detected in data from both the extended and compact Band 6 configurations. However, the Band 3 continuum for OH 30.1–0.7 has a lower S/N ratio and we do not detect the secondary peak. In Fig. A1, we plot the Band 3 continuum




Figure 1. Various continuum images for OH 30.1–0.7, plotted using logarithmic flux scales. The configurations are given in the top-left corner of each image and refer to the Band 6 observations unless Band 3 is indicated. The contours indicate levels of 3σ (thinnest white line), and 5σ , 10σ , 30σ , and 100σ (thicker white lines). The crosses at (0,0) indicate the position of the continuum peak; see Table 1 for the astrometric precision of the position. The beams are shown as white ellipses in the bottom-left corners. North is up and east is left. Additional details are given in Table A2.

next to the compact configuration Band 6 continuum, with the former processed to have the same restoring beam as the Band 6 data. In the compact Band 6 data, there appears to be a bridge connecting the primary and secondary peaks, traces of which are seen in the combined and extended continuum data in Fig. 1. This suggests an association between the two peaks. For simplicity, we will henceforth refer to the secondary continuum peak as ‘feature B’ and will assume that the AGB star coincides with the primary continuum peak. We find the separation between the two peaks to be 4.6 arcsec. At a distance of 3.9 kpc, this gives a projected separation of 1.8×10^4 au (2.7×10^{17} cm), corresponding to a lower limit on the physical separation. Feature B lies beyond the extent of the CO $J = 2 - 1$ and $J = 1 - 0$ emission, as discussed in Section 4.3.

2.2.2 Spectral indices

Since we have observations from two different ALMA Bands for OH 30.1–0.7, we next calculate the spectral indices of the AGB star and feature B. This is done using the continuum images from the compact Band 6 configuration and the Band 3 data processed to have the same beam as the compact Band 6 data (see Fig. A1 and Table A2). The nominal centre frequencies for the Band 3 and Band 6 continua are 107.86 and 236.02 GHz, and the rms are 0.013

and 0.042 mJy, respectively. The continuum peaks are 0.788 and 7.26 mJy beam $^{-1}$. Allowing for the rms and a 7 per cent flux scale uncertainty as applied to the target, per band, we find a spectral index of 2.8 ± 0.3 for the AGB star.

For feature B, the Band 6 flux is 0.46 mJy beam $^{-1}$ and for Band 3, the flux in the same region is 0.028 mJy beam $^{-1}$ (i.e. $< 3\sigma$). Using the Band 3 flux as an upper limit, this gives a lower limit on the spectral index of > 3.6 , even steeper than for the AGB star itself. These results are discussed in more detail in Section 4.3.

2.3 Line identification

The line identification was initially carried out by extracting spectra in circular apertures centred on the stellar continuum peak. Spectra were extracted for apertures of different radii to ensure that lines excited in different parts of the CSE were not overlooked. For example, compact lines, excited close to the star, are most easily detected in spectra extracted for smaller apertures, whereas more extended lines, especially those with shell-like emitting regions, are more easily detected in larger extraction apertures, where the aperture encompasses large fractions of their fluxes. In cases where the line detection appeared marginal in the spectra, we constructed zeroth moment maps (integrated intensity maps) to check for emission. The




Figure 2. ALMA spectra of one tuning towards OH 30.1–0.7, extracted for two apertures, as indicated in the top right of each subplot. The molecular carriers of various spectral lines are indicated at the top of each plot; see Table A3 for further details. The narrow feature at 230.6 GHz is interstellar CO contamination (see Section 2.4).

lines identified towards OH 30.1–0.7 are listed in Table A3. Lines are included in our list of detections if they are seen at levels of $\geq 3\sigma$ above the noise in at least one of the spectra and/or the zeroth moment map. Lines marked as tentative either fulfill only one of these criteria or are blended with another line. Zeroth moment maps were constructed to include all channels with emission $> 3\sigma$ in the spectra plus an additional 1–2 channels on either side of the line. An exception are the partially blended lines Na³⁷Cl and ³⁰SiS at 227.560 and 227.590 GHz, for which we carefully chose a minimal channel between the two peaks that was not then included in either zeroth moment map, to avoid cross-contamination.

We used data from the Cologne Database of Molecular Spectroscopy (CDMS, Müller et al. 2001, 2005) to assign carriers to detected lines. Once we had constructed our initial list of identified lines, we checked which lines from each species were covered by our observations, including lines in vibrationally excited states. The detailed results of these checks are given in Section 3, but we note here that in some cases we identified additional lines in this manner that are blended with other, often brighter, lines. In those cases, we note the potentially blended lines in the notes columns of Table A3.

An example spectrum from the Band 6 data, plotted for two extraction apertures, is given in Fig. 2 with the identified lines labelled. The remaining Band 6 spectra are shown in Fig. A2 and the Band 3 spectra are shown in Fig. A3.

2.4 Interstellar contamination and resolved-out flux

Previous observations of CO towards OH 30.1–0.7 have encountered problems with contamination by interstellar features, see for example De Beck et al. (2010). In short, because this star is located in the direction of a densely populated region of the galactic disc, a larger telescope beam – such as one from a single antenna – is prone to capture emission, especially of low-energy CO lines, from background or foreground sources along the line of sight,

in addition to the emission originating from the target AGB star. The contaminating sources are commonly molecular clouds with relatively narrow lines, compared with the wide lines of AGB CSEs, making them easy to distinguish, though not necessarily easy to disentangle from the AGB emission. In the particular case of OH 30.1–0.7, we have identified the massive star-forming region W43 as the likely source of background contamination, which has been found to have gas velocities comparable to the velocities at which we see contamination (Nguyen Luong et al. 2011). (See also Saral et al. 2017, where OH 30.1–0.7 can be seen as a bright orange point source on the edge of W43 in their fig. 2.)

ALMA has an advantage because, although the ALMA field of view can be similar to the half-power beam width (HPBW) of a single antenna, the interferometer acts as a spatial filter, resolving out smooth large-scale emission, including that from large molecular clouds. Unfortunately, some contamination persists, as can be seen in our CO channel maps in Fig. 3 and Fig. A6. A more thorough discussion of the contamination is given in Appendix B.

A common problem with observing using an interferometer such as ALMA is the possibility of resolved-out flux – that is, smooth large-scale flux is filtered out of the observations. However, the maximum recoverable scale of our lowest resolution Band 6 observations is 12.8 arcsec, larger than the CO emitting region (see Fig. 3), and the spectra of the low- and high-resolution Band 6 observations are in good agreement, indicating that no flux has been resolved out.

2.5 Integrated line fluxes and central velocities

Having identified the carriers of the lines in our data, as described in Section 2.3, we performed a basic analysis of all the unambiguous detections of unblended lines. We fit a soft parabola line profile




Figure 3. Combined ALMA observations of CO $J = 2 - 1$ towards OH 30.1–0.7. The thicker white contours are drawn at levels of 10σ , 30σ , 50σ , 70σ , 100σ , 150σ , 200σ , 300σ , and 500σ , the thinner (grey) contours at levels of 3σ and 5σ , and the dashed (cyan) contours at levels of -3σ and -5σ . The crosses at (0,0) indicate the position of the continuum peak. The velocity of each channel is shown in the top-right corner of each panel and the beam is shown as a white ellipse in the bottom-left corners. North is up and east is left. Note that there is significant interstellar contamination, especially for channels with velocities lower than 100 km s^{-1} .

function (Olofsson et al. 1993) defined as

$$F(v) = F_0 \left(1 - \left[\frac{v - v_{\text{cent}}}{v_w} \right]^2 \right)^{\gamma/2}, \quad (1)$$

where the central velocity of the line is v_{cent} , the width of the line is v_w , and the flux at the centre of the line is F_0 . The fit is done over the free parameters F_0 , v_{cent} , v_w , and γ . The soft parabola fit is performed for all extraction apertures in which a given line is detected. In Table A3, we list the central velocities and velocity widths obtained from these fits and the extraction aperture for which we determined that the line was most clearly detected. This means that for lines with compact emission we fit soft parabolas to spectra extracted for smaller apertures, while for extended emission we used

spectra extracted for larger apertures. Generally, we were not able to fit soft parabolas to tentative lines or lines with hyperfine structure. After obtaining the line widths from the soft parabola fits, we used these as a guide to calculate the integrated line fluxes, over velocity ranges a few km s^{-1} wider than the line widths from the soft parabola fits, to better account for flux that may be in line wings. For each individual line, the same spectrum (extracted for the same aperture) was used to find both the central velocity and the integrated flux. The integrated fluxes are also listed in Table A3.

Some examples of lines with their corresponding soft parabola fit are shown in Fig. A4, where we plot CO ($J = 2 - 1$), low-energy lines of SiS and SO₂, and a higher energy ($v = 2$) line of NaCl. The fits to the SiS and SO₂ lines produce larger line widths than for CO: $v_w \sim 22 \text{ km s}^{-1}$ compared with $v_w = 18 \text{ km s}^{-1}$ for CO.

This is because the automated fit does not account for the blue wing of the CO emission where flux is strongly detected in the channels 76.5–81.5 km s⁻¹, despite there not being a clear CO detection in the channel at 84.1 km s⁻¹. The reason for this phenomena is discussed further in Section 4.2.2. To better account for the full velocity width of the CO line, and hence the expansion velocity of the CSE, we calculate the full width of the CO line from the channel maps (Fig. 3). By considering the reddest and bluest channels with CO detections centred on the continuum peak with a certainty > 30 σ , we determine the expansion velocity to be 23 km s⁻¹.

By averaging the central velocities found from the soft parabola fits, we calculate a systemic velocity of $v_{\text{LSR}} = 100.3 \pm 1.2$ km s⁻¹. This value is in agreement with the LSR velocity of 100.0 km s⁻¹ found by De Beck et al. (2010) from CO observations. The most significant outlier in our sample is the Na³⁵Cl ($v = 1$, $J = 8 - 7$) line at 103.415 GHz in Band 3, which has a central velocity of 103.7 km s⁻¹. We judge the line unlikely to be misidentified since several other NaCl lines are detected in our observations, including vibrationally excited lines. It is possible that this line might be blended with another, unidentified line. However, we note that the other vibrationally excited NaCl lines tend to lie redwards of the systemic velocity, while the lines in the ground vibrational state lie closer to the systemic velocity (within 1 km s⁻¹ but on the blue side). We also find that the KCl lines overall tend to lie slightly redwards of the systemic velocity (with the exception of one ⁴¹K³⁵Cl line). The CO lines also lie 0.6 to 1.6 km s⁻¹ redwards of the systemic velocity. In contrast, the SiS, SO₂, and CS lines tend to lie bluewards of the systemic velocity. These differences are likely because of differences in excitation conditions and regions and certainly contribute to the moderate uncertainty on our averaged systemic velocity.

3 DETECTED MOLECULES

In the following subsections, we discuss the detections of the various molecular species seen towards OH 30.1–0.7 in our observations. The observations, including the number of detected lines and the range of lower level energies they cover, are summarized in Table 2. Of particular note are the large number of NaCl and KCl lines (discussed in Section 3.6) and the NS lines detected (discussed in Section 3.5). After carefully checking our data, we were not left with any unidentified lines.

3.1 CO

CO is a well-studied density tracer of CSEs. In Fig. 3, we plot the combined channel maps of the $J = 2 - 1$ emission towards OH 30.1–0.7. Because of the large amount of interstellar contamination, we highlight contours at and above 10 σ and also show significant (3 σ and 5 σ) negative contours, which arise from resolved out (interstellar) flux. The CO emission originating from OH 30.1–0.7 has an almost circular distribution across channels, with the diameter of the emission shrinking towards more extreme red and blue velocities, as is typical of emission from a fairly spherical, radially expanding envelope. The most significant deviation from this is the absence of circumstellar emission in the channel at 84.1 km s⁻¹. This could be the result of CO self-absorption or a background interstellar medium (ISM) source with a similar flux level visually blended with the CSE and resulting in the CSE being resolved out in this channel. Single-dish observations (e.g. Heske et al. 1990; Justtanont et al. 2013, and Fig. B1) show significant contamination around this channel. There are also some internal arcs contributing to density

Table 2. An overview of all the molecules detected towards OH 30.1–0.7.

Molecule	E_{low} [K]	N	E_{low} [K]	N_{lines}	N_{iso}	Section	Ref.
CO	16	3	0–6	3	2	3.1	1
SiS	145–5431	11	66–1134	8	4	3.2	2
CS	49	1	24	1	1	3.3	3
SO ₂	19–152	4	3–108	10	3	3.4	4
NS	53–373	3*	3	2*	1	3.5	5
NaCl	199–2263	10	17–2143	9	2	3.6	6
KCl	333–1563	17	39–977	9	3	3.6	7

Notes. The E_{low} columns give the range of lower level energies among the detected lines of that molecule. The N_{lines} column indicates the total number of lines from that molecule, including isotopologues but excluding hyperfine components, that were detected. The N_{iso} column indicates the total number of isotopologues that were detected. The Sect column gives the reference to the section in this work where observations of that molecule are discussed in more detail. The Ref column gives the reference for the line frequencies included in this work. *indicates that the molecular lines are made up of multiple (hyperfine) components which are not counted as separate lines. References: 1: Winnewisser et al. (1997), 2: Müller et al. (2007), 3: Gottlieb, Myers & Thaddeus (2003), 4: Klisch et al. (1997); Belov et al. (1998); Müller & Brünken (2005) 5: Lee, Ozeki & Saito (1995) 6: Caris, Lewen & Winnewisser (2002), and 7: Caris et al. (2004).

structures within the CSE, notably in the channels from 86.6 to 114.0 km s⁻¹. The internal structures are more clearly seen in the (overall slightly less sensitive) extended configuration only channel maps, shown in Fig. A5, which also show less contamination owing to a smaller maximum recoverable scale. The $J = 1 - 0$ channel maps are shown in Fig. A6 and show a similar almost-circular distribution. However, the internal structures are not resolved, owing to a larger beam size (1.5 arcsec \times 1.1 arcsec, see Table A1).

The emission in the central channel of $J = 2 - 1$, which we defined as the channel with LSR velocity $v_{\text{LSR}} = 99.3$ km s⁻¹ because it is the closest to the systemic velocity, extends out to separations of 2.5–3.9 arcsec (9800 to 15 000 au) from the continuum peak when considering the 3 σ contour. The minimum extent occurs to the north-northwest, while the maximum extent is in the southeasterly direction (see Fig. 4). For the Band 3 data of CO $J = 1 - 0$, the channel closest to the systemic velocity (99.2 km s⁻¹) suffers from a large amount of interstellar contamination, so we measured the extent for the channel with $v_{\text{LSR}} = 101.7$ km s⁻¹, and found that the smallest and largest extents were 3.3 and 4.4 arcsec, in the same directions as for the $J = 2 - 1$ emission (see Fig. A6).

In Fig. 4, we plot integrated intensity maps of CO $J = 2 - 1$ and $J = 1 - 0$ emission. For the $J = 2 - 1$ map, we also overplot the combined continuum emission. For the $J = 1 - 0$ map, we overplot the continuum emission from the Band 6 compact array, which is of a comparable resolution and is sensitive to feature B, unlike the Band 3 continuum. For the $J = 1 - 0$ emission we only integrate over the red part of the CO line (i.e. for $v \geq 101.7$ km s⁻¹) because the ISM contamination is too large and confounds the data at lower velocities. The comparison in Fig. 4 shows that the ‘indentation’ seen in the CO emission to the north-northwest, corresponds well to the location of feature B in the continuum emission. The indentation is most clearly seen in the CO $J = 2 - 1$ emission in the velocity channels from 91.7 to 114.6 km s⁻¹ (Fig. 3). In the CO $J = 1 - 0$ emission the indentation is less apparent, owing to the larger beam size of those observations. Nevertheless, there is a slight asymmetry in the direction of feature B. This phenomenon is discussed in more detail in Section 4.3.




Figure 4. Left: the zeroth moment map of CO $J = 2 - 1$ from the combined data towards OH 30.1–0.7. The contours indicate flux levels of 5σ , 10σ , 30σ , and 50σ . The continuum peak is indicated by a cross at (0,0) and the beam is shown as a white ellipse in the bottom-left corner. North is up and east is left. Centre: the same CO $J = 2 - 1$ emission overlaid with red contours showing the continuum emission in Band 6 at levels of 5σ , 10σ , 30σ , and 100σ . The white ellipse in the bottom left corner corresponds to the CO beam and the red ellipse to the continuum beam. Right: the intensity map of CO $J = 1 - 0$ integrated over the red channels (from 101.7 to 124.6 km s^{-1} , refer to Fig. A6), overlaid with the Band 6 compact array continuum emission. The white contours indicated levels of 3σ , 5σ , 10σ , 30σ , and 50σ in the CO flux and the red contours indicated the 5σ , 10σ , and 30σ levels of the continuum flux. The white ellipse in the bottom-left corner corresponds to the CO beam and the red ellipse to the continuum beam.

One line of CO in the first vibrationally excited state was covered in our observations but not detected. In addition to the main isotopologue of CO, we tentatively detect the $\text{C}^{17}\text{O } J = 1 - 0$ line in the spectrum. The line is not clearly seen in the zeroth moment map and it is possible that there is some interstellar contamination around that frequency. No lines of ^{13}CO or C^{18}O were covered and we note that detections of C^{18}O were not expected, as previous studies have shown evidence of HBB (Justtanont et al. 2015), which destroys ^{18}O (Karakas & Lugaro 2016).

3.2 SiS

Our spectral coverage of SiS is uneven and misses lines from the most abundant isotopologue, $^{28}\text{Si}^{32}\text{S}$ (hereafter we drop the isotope labels for ^{28}Si and ^{32}S). Therefore, although we detect several lines from isotopologues of SiS, we do not perform an analysis of isotopic ratios in this work.

We detect all the covered lines of ^{29}SiS , ^{30}SiS , and Si^{34}S in the ground and first vibrationally excited states and note that these lines were not covered for Si^{33}S . The vibrational ground state $^{29}\text{Si}^{34}\text{S}$ line covered in our Band 6 observations was also detected. No lines in the lowest two vibrationally excited states were covered by our Band 3 observations. The Band 3 data were not sensitive enough to detect any lines of doubly substituted isotopologues.

Zeroth moment maps of our SiS detections are plotted in Fig. 5. The lines in the ground vibrational state of the singly substituted isotopologues are relatively featureless, with only a few small protrusions seen for the $^{30}\text{SiS } v = 0, J = 14 - 13$ line at a level of 3σ . The lines in the vibrational ground state extend out to 0.3 to 0.7 arcsec (1200 to 2700 au) from the continuum peak, and the peaks of the SiS zeroth moment maps tend to be offset slightly from the continuum peak in the southwest direction. The SiS lines in vibrationally excited states and for the doubly substituted isotopologue are either spatially unresolved or marginally resolved and do not exhibit any structure, though some are still slightly offset to the southwest of the continuum peak.

3.3 CS

One line of $^{12}\text{C}^{32}\text{S}$ (hereafter, CS) in the vibrational ground state was detected. The zeroth moment map for CS is shown in Fig. 6. The

emission is centrally peaked and the smoothed 3σ contour extends out to ~ 1 arcsec (3900 au). We smoothed the contour by binning and averaging it over 60° sections, to reduce the ragged edges of the contour. We also neglected isolated islands of emission for this measurement.

No lines in the vibrational ground state of ^{13}CS or C^{34}S were covered in our frequency setups. One line of C^{33}S in the ground vibrational state was covered and not detected, as expected given the likely isotopic ratio and the intensity of the corresponding C^{32}S line.

3.4 SO₂

SO_2 was detected towards OH 30.1–0.7 with a shell-like distribution, as expected for a star with a high mass-loss rate and a dense stellar wind (Danilovich et al. 2016; Wallström et al. 2024, see also Appendix C3). In general, the detected transitions span relatively low-energy levels ($E_{\text{low}} < 110$, see Table 2). This is despite higher energy transitions being covered by our observations, in line with SO_2 mainly being formed and excited further out in the wind. A notable non-detection was the $\text{SO}_2 J_{K_a, K_c} = 11_{5,7} - 12_{4,8}$ line, which has a lower level energy of 111 K, only 3 K above our highest energy detected line ($J_{K_a, K_c} = 14_{3,11} - 14_{2,12}$, $E_{\text{low}} = 108$ K). This result is in line with the trend reported by Wallström et al. (2024) for higher mass-loss rate AGB stars to exhibit lower energy SO_2 lines, and lower mass-loss rate AGB stars to exhibit higher energy SO_2 lines.

In Fig. 7, we plot the predicted line intensities of all the SO_2 lines (taken from CDMS) with lower level energies < 500 K. If we consider the predicted line intensities at 300 K (see right panel in Fig. 7), then all the covered lines with lower level energies between 100 and 500 K are expected to be brighter than at least one or two lines that were detected. We checked the predicted line intensities for several other temperatures and found that the undetected $J_{K_a, K_c} = 11_{5,7} - 12_{4,8}$ line is only predicted to be fainter than all the detected lines for temperatures $\lesssim 25$ K. The distribution for predicted line intensities at 25 K is shown in the left panel of Fig. 7. This suggests a surprisingly low excitation temperature for SO_2 around OH 30.1–0.7. This results from the most abundant isotopologue is in agreement with the exclusively very low-energy transitions (tentatively) detected for $^{34}\text{SO}_2$ and $^{33}\text{SO}_2$.

In Fig. 8, we plot zeroth moment maps of the most clearly detected SO_2 lines. In total, five lines of SO_2 were detected, one of which we




Figure 5. Zeroth moment maps of SiS towards OH 30.1–0.7. The details of the transitions are given in Table A3. The transitions are indicated in the top right of each panel with the details of the transitions given in Table A3. The contours are drawn at levels of 3σ , 5σ , 10σ , and 30σ . The beam is shown as a white ellipse in the bottom-left corner and the location of the continuum peak is indicated by the cross at (0,0). North is up and east is left.




Figure 6. Zeroth moment map of CS towards OH 30.1–0.7. The details of the transition are given in Table A3. The contours are drawn at levels of 3σ , 5σ , 10σ , and 30σ . The beam is shown as a white ellipse in the bottom-left corner and the location of the continuum peak is indicated by the cross at (0,0). North is up and east is left.

do not plot because it is blended with a line of KCl, and four lines of $^{34}\text{SO}_2$ were detected, two of which are bright enough to plot in Fig. 8. We tentatively detected one $^{33}\text{SO}_2$ line, which we do not plot because the emission is too faint in the zeroth moment map (but is tentatively detected in the spectrum). The two plotted Band 3 lines of SO_2 and $^{34}\text{SO}_2$ coincidentally have the lowest level energies with $E_{\text{low}} < 10$ K. The lines are resolved and show some internal structure in the zeroth moment maps, with visible emission peaks offset from the continuum peak. We find that the Band 3 emission extends out to a minimum of 1.5 arcsec and a maximum of 2.5 arcsec from the continuum peak for the SO_2 line (measured after binning the 3σ contours, as described for CS), and out to a minimum of 1.7 arcsec and a maximum of 2.2 arcsec for the $^{34}\text{SO}_2$ line. The slight difference between these extents and between the corresponding flux distribution patterns shown in Fig. 8 are most likely a result of different sensitivities and the slightly different gas temperatures being probed by the lines. The Band 6 observations of SO_2 show well-resolved emission and cover slightly higher energy lines (40–108 K for SO_2 and 8–82 K for $^{34}\text{SO}_2$). The $J_{K_a, K_c} = 14_{0,14} - 13_{1,13}$ line is the brightest covered line ($E_{\text{low}} = 82$ K) and exhibits well-resolved




Figure 7. The predicted relative integrated intensities for the SO_2 lines covered by our observations with lower level energies < 500 K. The detected lines are shown as blue squares and the other covered lines as orange circles. Filled markers are lines that were covered in Band 6 and unfilled markers in Band 3. On the left, we show the predictions for a gas temperature of 25 K and on the right, we show the same for a gas temperature of 300 K.

extended emission. As can be seen in the channel maps plotted in Fig. A7, the emission peak lies to the northwest of the continuum peak in most channels. The smoothed 3σ contour in the zeroth moment map (Fig. 8) extends out to 1.1–2.1 arcsec, comparable to the Band 3 emission. The $J_{K_a, K_c} = 10_{3,7} - 10_{2,8}$ ($E_{\text{low}} = 61$ K) has a very similar distribution, including the offset emission peak, and a slightly smaller extent for the 3σ contour, ranging from 0.7 to 1.9 arcsec. This difference most likely arises because the full (outer) extent of

the weaker line is below the noise. The highest energy detected line is $J_{K_a, K_c} = 14_{3,11} - 14_{2,12}$ with $E_{\text{low}} = 108$ K. The emission peak of this line is also offset slightly to the northwest. As can be seen in Fig. 8, it has the smallest extent of the well-detected SO_2 lines, out to only 0.4–0.8 arcsec. This could be a result of observational sensitivity and the line being intrinsically fainter, or could be an indication that the slightly warmer emission is not distributed as widely as for the lower energy lines. Given our low estimate for the SO_2 gas temperature, the former explanation is more likely.

We also note that the central velocities found for the sufficiently bright SO_2 lines (see Table A3) tend to fall on the blue side of the average systemic LSR velocity by 1–2 km s^{-1} for SO_2 , and by 0.4 km s^{-1} for $^{34}\text{SO}_2$. The two brightest Band 6 SO_2 lines are the most blueshifted. We do not calculate isotopic ratios here because we do not clearly detect any two pairs of lines with the same quantum numbers across isotopologues.

3.5 NS

NS was only covered by the less sensitive Band 3 observations. Two lines in the $^2\Pi_{1/2}$ ladder were covered, though only the e line was clearly detected. Spectra for both lines are shown in Fig. 9 with the hyperfine components marked. In Table A3, we list the three brightest hyperfine components for each line. For the f line, the brightest hyperfine component is separated from the next two brightest components by 38 and 41 km s^{-1} , further than the half-width of single-component lines towards this star. Whereas for the e line, the brightest hyperfine component is only 7.5 and 24 km s^{-1} from the next two brightest components. The overlap of the two brightest components means that their fluxes combine to make the e line more detectable. We plot a theoretical Gaussian profile with the spectra (red dotted line in Fig. 9, assuming Gaussian components




Figure 8. Zeroth moment maps of SO_2 towards OH 30.1–0.7. The top-left and centre lines were observed with ALMA Band 3 and the remaining lines were observed with ALMA Band 6. Note the different angular scales and resolutions for the two sets of observations. The transitions are indicated in the top right of each panel with the details of the transitions given in Table A3. The contours are drawn at levels of 3σ , 5σ , 10σ , and 30σ . The beam is shown as a white ellipse in the bottom left corner and the location of the continuum peak is indicated by the cross at (0,0). North is up and east is left.




Figure 9. NS emission towards OH 30.1–0.7. The details of the transitions are given in Table A3. Left: Zeroth moment map of e component of NS towards OH 30.1–0.7. The contours are drawn at levels of 3σ and 5σ . The pink dotted lines are drawn such that one connects the AGB continuum peak (red cross) with the peak of feature B (not shown) and the second line is perpendicular to this, passing through the AGB continuum peak. The beam is shown as a white ellipse in the bottom-left corner. North is up and east is left. Right: Spectra extracted from centred circular apertures with radii of 2 arcsec (black histograms) for NS lines observed towards OH 30.1–0.7. The positions of the hyperfine components are shown as vertical red lines with heights corresponding to their relative intensities. The dotted red lines show the predicted line profiles for Gaussian hyperfine components of width 5 km s^{-1} . The velocities are calculated with respect to the brightest hyperfine component.

with widths of 5 km s^{-1}) to emphasize the theoretical line shapes. The observed e component has a more double-peaked profile than the predicted Gaussian profile, suggesting that emission might originate in a shell around the star, rather than in a spherical region centred on the star.

The zeroth moment map in Fig. 9 shows that the NS emission is mostly located to the east and north of OH 30.1–0.7, and is only marginally spatially resolved. The emission is highly asymmetric; the averaged 3σ contour of the emission extends out to only 0.7 arcsec to the south-southwest and out to 1.8 arcsec to the east-northeast. The major axis of the emitting region lies perpendicular to the projected axis joining the AGB continuum peak and feature B. This is emphasized by overplotting two dotted lines on the zeroth moment map in Fig. 9. Although the S/N of the NS emission is rather low in individual channels (summing to a 5σ detection in the zeroth moment map), we constructed a position–velocity (PV) diagram along the major axis (69° east of north, as illustrated in Fig. 9), which shows emission offset from the continuum peak by around ~ 1.5 to 2 arcsec and $\sim 5 \text{ km s}^{-1}$ (see Fig. A8). The potential astrochemical implications of the NS distribution will be discussed in Section 4.1.1.

3.6 NaCl and KCl

We detect several lines of NaCl and KCl, including lines from the ^{35}Cl and ^{37}Cl isotopologues of both species. We detect lines up to the third and fourth vibrationally excited states of Na^{35}Cl and up to the third vibrationally excited state of K^{35}Cl . These data represent the first detections of highly excited vibrational states of both NaCl and KCl towards AGB stars. Zeroth moment maps of NaCl are given in Fig. 10, and of KCl in Fig. 11. The emitting regions of both molecules are generally featureless and uniform, even for the lower energy lines which are spatially resolved.

No lines of Na^{35}Cl in the ground vibrational state were covered by our observations. However, we clearly detect three lines of Na^{37}Cl in the ground vibrational state. All lines of both isotopologues covered in Band 6 were detected up to $v = 4$. Owing to the lower sensitivity of the Band 3 data, only one Na^{35}Cl line in the first vibrationally excited state was detected.

The $v = 0$, $J = 8 - 7$ line of Na^{37}Cl appears to be marginally spatially resolved, with the 3σ contour extended out to 1–1.6 arcsec from the continuum peak. This is likely an overestimate of the emission extent owing to the larger beam for the Band 3 observations compared with the Band 6 observations. We note that the $J = 9 - 8$ line appears slightly smaller, despite the line being a little brighter, but this is almost certainly a sensitivity effect since the line lies in a slightly noisier part of the spectrum. Most of the NaCl lines covered in Band 6 appear to be more spatially resolved than the Band 3 lines (with the exception of the $v = 4$ line). However, we note that even the two brightest lines, the Na^{37}Cl $v = 0$, $J = 18 - 17$ and $v = 1$, $J = 19 - 18$ lines, which appear several times larger than the beam, still have a slightly oblong shape along the same axis as the beam, suggesting that they are not truly resolved. The smoothed 3σ contour for the Na^{37}Cl $v = 0$, $J = 18 - 17$ line (the only ground vibrational state line observed in Band 6) extends out to 0.3–0.6 arcsec.

Our Band 3 data were only sensitive enough to detect the single covered $^{39}\text{K}^{35}\text{Cl}$ line in the ground vibrational state, which is partly blended with CO, and none of the lines in higher vibrational states. In the Band 6 data, we detect all covered lines in the ground, first and second vibrational states of $^{39}\text{K}^{35}\text{Cl}$ and $^{39}\text{K}^{37}\text{Cl}$. We detect two lines of $^{41}\text{K}^{35}\text{Cl}$ in the second vibrationally excited state, one tentatively, and we note that no lines in the ground or first vibrationally excited state were covered in the Band 6 observations.

As can be seen in Fig. 11, the KCl lines are not spatially resolved. Similar to the observed NaCl emission, the two KCl lines in the ground vibrational state appear larger than the beam but are still slightly elongated along the same axis as the beam. The 3σ contours for both lines extend out to 0.2–0.5 arcsec from the continuum peak.

The significance of detecting so many (high-energy) lines of NaCl and KCl are discussed in Section 4.1.2, as are the potential chemical formation pathways. Based on the data presented here, we constructed population diagrams for Na^{35}Cl and K^{35}Cl . The resultant plot is given in Fig. A9. We derive very similar temperatures for both species: approximately 600 K. This is consistent with the emission originating from the inner wind, unlike the low-energy SO_2 emission




Figure 10. Zeroth moment maps of NaCl towards OH 30.1–0.7. The lines in the top row were observed with ALMA Band 3 and the middle and bottom rows were observed with ALMA Band 6. Note the different angular scales and resolutions. The details of the transitions are given in Table A3. The contours are drawn at levels of 3σ , 5σ , 10σ , and 30σ . The beam is shown as a white ellipse in the bottom-left corner and the location of the continuum peak is indicated by the cross at (0,0). North is up and east is left.

(see Section 3.4). We also derive column densities of $1.4 \times 10^{12} \text{ cm}^{-2}$ for Na^{35}Cl and $1.6 \times 10^{11} \text{ cm}^{-2}$ for K^{35}Cl , resulting in $\text{Na}^{35}\text{Cl}/\text{K}^{35}\text{Cl} = 8.8$. This is higher than expected based on chemical equilibrium and may be an indication that the abundance ratio of Na/K is higher than solar (in agreement with models for intermediate-mass stars, especially if OH 30.1–0.7 had a higher initial metallicity than solar, Cinquegrana & Karakas 2022). This will be explored in more detail in a follow-up study.

4 DISCUSSION

4.1 The circumstellar chemistry of OH 30.1–0.7

Several of the molecules detected towards OH 30.1–0.7 are commonly seen around oxygen-rich AGB stars generally. These include CO, SiS, and CS. In the case of oxygen-rich AGB stars, SiS, CS are commonly seen for those with high mass-loss rates (or, equivalently, high wind densities; Danilovich et al. 2018). SO_2 is very commonly detected around oxygen-rich AGB stars (but has not been detected for carbon-rich or S-type stars), with a shell-like distribution for high mass-loss rate stars (Danilovich

et al. 2020; Wallström et al. 2024). The observations of all of these molecules towards OH 30.1–0.7 agree with previously established trends, as explained in more detail in Appendix C.

NS, NaCl, and KCl have been observed less frequently and hence studied in less detail thus far. It is on these three molecules that we will focus our discussion in the following subsections. Some additional background on past observations of these molecules is also given in Appendix C.

4.1.1 NS

Previous observations of the NS molecule towards AGB stars include the oxygen-rich AGB star IK Tau (Velilla Prieto et al. 2017), where resolved ALMA observations show its emitting region to be close to the continuum peak (within 170 au, Decin et al. 2018), and the S-type AGB star W Aql (Danilovich et al. 2024). For W Aql, Danilovich et al. (2024) found that NS may have been formed when the main-sequence (F9V) companion to the AGB star passed through the dense inner CSE, temporarily driving UV photochemistry in that region. This conclusion is based on the chemical models of Van de Sande &




Figure 11. Zeroth moment maps of KCl towards OH 30.1–0.7. The details of the transitions are given in Table A3. The contours are drawn at levels of 3σ , 5σ , 10σ , and 30σ . The beam is shown as a white ellipse in the bottom-left corner and the location of the continuum peak is indicated by the cross at (0,0). North is up and east is left.

Millar (2022), who found that the formation of NS was enhanced in the presence of UV photons from a companion star. Although NS towards IK Tau has not been analysed in detail (only some basic measurements are given in Decin et al. 2018), the recent analysis of NaCl in the same star found evidence that a companion may be driving or enhancing the formation of NaCl (Coenegrachts et al. 2023). It is also possible that the formation of NS around IK Tau could be driven by that companion, albeit primarily through UV chemistry rather than the shock-heating proposed by Coenegrachts et al. (2023) for NaCl.

The radial extent of NS in the case of OH 30.1–0.7 is much larger than the extents seen for IK Tau or W Aql. The integrated NS emission towards W Aql is irregular but is detected no more than $0.4 \text{ arcsec} \approx 160 \text{ au}$ from the star (Danilovich et al. 2024). For IK Tau, the angular width is reported to be $490\text{--}650 \text{ mas}$ (Decin et al. 2018), which corresponds to radial extents of ~ 130 to 170 au . In contrast, for OH 30.1–0.7, the most distant NS emission within the 3σ contour is seen at 1.8 arcsec . This corresponds to 7000 au , while the southeastern extent of 0.7 arcsec corresponds to 2700 au . These distances come with the caveat that the Band 3 emission is only marginally resolved, so the radial extents may be overestimated and only represent upper limits (e.g. see the difference between the Band

3 and 6 emission extents for NaCl in Section 3.6). Taken at face value, this emission appears to be 20–40 times further from the AGB star than the most distant NS emission seen around W Aql and IK Tau. The theoretical study of Van de Sande & Millar (2022) found that higher mass-loss rates result in NS peaks further from AGB star and that a relatively warm companion star is required to form NS; i.e. a solar-type or white dwarf companion will drive NS formation but a red dwarf will not.

The asymmetric NS emission in the zeroth moment map (Fig. 9) is reminiscent of the SiN emission around W Aql (Danilovich et al. 2024), which was determined to lie on the side of the AGB star that aligned with the periastron of the binary orbit. If a similar scenario can account for the NS around OH 30.1–0.7, then the alignment of the NS emission can help constrain the projection of the orbital plane on the sky. Another possibility is a companion on a circular but wider orbit such that NS is preferentially formed on the side of the CSE where the companion is presently located.

4.1.2 Chlorides

There have been enough adequately sensitive observations of a variety of AGB stars that we can firmly say that NaCl and KCl

are not ubiquitously present in CSEs. This is most clearly illustrated by the detection of one or both molecules in 3 AGB stars out of a sample of 14 in the ATOMIUM survey (Wallström et al. 2024). Some physical or chemical characteristic must set these stars apart from their unsalted counterparts. Similarly, something should also set stars like OH 30.1–0.7, with a large number of NaCl and KCl lines, apart from stars like IK Tau, with detections only in the ground vibrational state (Velilla Prieto et al. 2017; Decin et al. 2018).

Coenegrachts et al. (2023) proposed that an unseen companion could be contributing to the formation of NaCl around IK Tau, since the motion of the companion through the CSE would be supersonic, hence generating a shock. As discussed by Decin et al. (2019), and evidenced by our detection of NS, OH 30.1–0.7 is also thought to have a companion. Since the OH 30.1–0.7 shows evidence of HBB (Justtanont et al. 2015), we can conclude it had an intermediate initial mass, $\gtrsim 4 M_{\odot}$. In this mass range, the binary fraction is thought to be close to one or higher (that is, stars with masses from 4–8 M_{\odot} are expected to have one or more companions with mass ratios $q > 0.1$, Moe & Di Stefano 2017). In light of this, and the results of Coenegrachts et al. (2023), it is highly likely that a companion is present around OH 30.1–0.7 and that such a companion could be driving the formation of NaCl and KCl in a similar way. Unlike the case of IK Tau, we see many vibrationally excited NaCl and KCl lines around OH 30.1–0.7. This is consistent with the different temperatures found for NaCl around the two stars: 14–70 K for IK Tau (Velilla Prieto et al. 2017; Coenegrachts et al. 2023) and 600 K for OH 30.1–0.7 (see Section 3.6 and Fig. A9). This may be a result of more extreme shocks owing to the higher wind density of OH 30.1–0.7 and possibly a higher orbital velocity of the companion.

The formation pathways of NaCl and KCl are discussed in detail in Appendix C5.1. We conclude that a companion star hotter than around ~ 12000 K would cause too much photoionization of Na and K to reconcile our observations, unless densities are very high in that region of the wind. Hence, we can hypothesize that the companion to OH 30.1–0.7 cannot be hotter than a late B-type main-sequence star.

4.2 Our evolving understanding of OH 30.1–0.7

OH 30.1–0.7 has been classified as an extreme OH/IR star, which is to say, a dusty AGB star with a high mass-loss rate and bright OH masers (Justtanont et al. 2006). It is an example of a large-amplitude variable star with one of the longest periods known ($P = 2170$ d ~ 6 yr,¹ Engels et al. 2024). Such stars are thought to have evolved to the end of the AGB phase (e.g. Olofsson et al. 2022). From *Herschel* observations of H₂O isotopologues, OH 30.1–0.7 was determined to have an intermediate initial mass ($\gtrsim 4 M_{\odot}$, Justtanont et al. 2015). This is in contrast with several other OH/IR stars which have been found to have lower initial masses $\sim 1.5 M_{\odot}$ (Olofsson et al. 2022). An overview of past observational difficulties, including a discussion on the reliability of observations in light of ISM contamination towards OH 30.1–0.7, are given in Appendix D.

4.2.1 Past results from the literature

OH 30.1–0.7 was included in the CO study of Heske et al. (1990), which was the first to observe a sample of OH/IR stars in CO emission. Heske et al. (1990) found that their observed intensity

¹Obtained from its OH maser light curve: https://hsweb.hs.uni-hamburg.de/projects/nrt_monitoring/stockert.html

ratios of the CO $J = 1 - 0$ and $J = 2 - 1$ lines could be explained by variable mass-loss rates at different eras, which could be the reason why mass-loss rates derived from CO observations resulted in an order of magnitude lower mass-loss than those from dust. This idea, first dubbed the ‘superwind’ scenario by Renzini (1981), was taken up and developed in subsequent work (e.g. Justtanont & Tielens 1992; Delfosse et al. 1997; Justtanont et al. 2013). The idea, as it developed, being that in the last portion of the AGB phase, mass loss increases substantially (i.e. the stellar wind becomes a superwind) during a relatively short period immediately before the completion of the AGB phase.

Decin et al. (2019) presented the ALMA CO observations of OH 30.1–0.7 (from the data set also presented here, though we have performed a new reduction of the ALMA data using self-calibration, see Section 2.1) and of another extreme OH/IR star, OH 26.5 +0.6. A key argument of Decin et al.’s (2019) work was that it is statistically unlikely that a dozen extreme OH/IR stars should all be observed in the midst of the relatively short superwind phase. Rather, they interpret the structures within the CO emission of OH 30.1–0.7 (and the spiral structure seen for OH 26.5 +0.6) as interactions with a companion star. The ultimate result of these interactions, they argue, is an equatorial density enhancement; that is, the companion pulls some of the wind into a toroidal structure of denser gas and dust around the star. This equatorial density enhancement could then mimic the superwind by making the mass-loss rate appear high because of the line of sight alignment of this region.

Low $^{12}\text{CO}/^{13}\text{CO}$ ratios ~ 3.5 were found by both Delfosse et al. (1997) and Justtanont et al. (2013) for OH 30.1–0.7, using different observational data-sets and a similar approach involving radiative transfer modelling. This low $^{12}\text{CO}/^{13}\text{CO}$ ratio provides further evidence of HBB in OH 30.1–0.7, and hence that it must be an intermediate-mass star, with an initial mass $4 M_{\odot} \lesssim M_i \lesssim 8 M_{\odot}$.

The recent study of Marini et al. (2023) modelled the SED of OH 30.1–0.7 and compared various properties such as luminosity and pulsation period of the star to theoretical models. Based on their results (and considering the distance derived by Engels et al. 2015), their determination of the initial mass of OH 30.1–0.7 is likely between 4 and 5 M_{\odot} . However, if the luminosity is higher (i.e. $5 \times 10^4 L_{\odot}$ based on the luminosity adopted by Justtanont et al. (2013), scaled to our adopted distance of 3.9 kpc) then OH 30.1–0.7 could have an initial mass of 6 M_{\odot} .

4.2.2 New understanding from ALMA CO

Our ALMA observations show that the angular extent of the CO emission for both the $J = 2 - 1$ and $J = 1 - 0$ are essentially the same. Our measured maximum angular extents (see Section 3.1) are 3.5 arcsec for $J = 2 - 1$ from the combined data and 4.2 arcsec for $J = 1 - 0$. Although it may appear from this that the $J = 1 - 0$ extent is larger than the $J = 2 - 1$ extent, this is primarily a function of the different beam sizes (see Table A1). If we instead compare the data from only the compact array configuration for $J = 2 - 1$ (with a comparable beam to the Band 3 data), then we find a maximum angular extent of 4.2 arcsec, as for the $J = 1 - 0$. This unexpected result suggests that both the emission from CO $J = 1 - 0$ and $J = 2 - 1$ trace the outer extent of the CO envelope. Note also that in the channel maps close to the LSR velocity (and ignoring the regions with the most contamination) the lowest few contours are very close together (3σ , 5σ , and 10σ for $J = 1 - 0$, Fig. A6, and 5σ , 10σ , 30σ , and 50σ for $J = 2 - 1$, Fig. 3), indicating a sharp drop-off in CO emission and likely also CO abundance. This is also seen

in the reddest and bluest channels (more clearly for CO $J = 2 - 1$ because of higher contamination for CO $J = 1 - 0$) where the CO emission is detected above 30σ , then not detected in the subsequent channel at all. This behaviour is in contrast with other spatially resolved observations of AGB stars which typically have less abrupt edges, with gradually decreasing emission across the velocity and angular axes (see e.g. Cernicharo et al. 2015; Kim et al. 2017 and the channel maps in the supplementary material of Decin et al. 2020, for a selection of varied AGB stars).

If we take half the maximum velocity extent in the CO channel maps as the wind speed (23 km s^{-1}) and consider the well-constrained distance of 3.9 kpc, we find that it took around 2800 yr for the gas to expand to $3.5''$ ($2 \times 10^{17} \text{ cm} = 14000 \text{ au}$) from the star. This is around a factor of 3 longer than the superwind phase obtained for this star by Delfosse et al. (1997) and a factor of 8 larger than the superwind radius derived by Justtanont et al. (2013). We also do not detect any significant emission beyond this radius, aside from the background contamination, which is not evenly distributed around the star. In conjunction with the sharp drop-off in CO emission discussed above, this suggests that any earlier mass loss must have been at a much lower rate, if it was present at all. If we consider the photodissociation radius of CO based on the mass-loss rate, expansion velocity and interstellar radiation field, and take the measured envelope size to be the half-abundance radius for CO destruction, then the mass loss corresponding to the observed envelope size is $\dot{M} \approx 2 \times 10^{-5} M_{\odot} \text{ yr}^{-1}$ (Mamon, Glassgold & Huggins 1988; Saberi, Vlemmings & De Beck 2019). Essentially, any earlier mass loss at this rate or lower would not be detectable, because that CO would have photodissociated before expanding to larger radii. So if there was a lower mass-loss rate ($\leq 2 \times 10^{-5} M_{\odot} \text{ yr}^{-1}$) before the current high mass-loss rate period commenced 2800 yr ago, it would not be detectable.

If we take the mass-loss rate of $\dot{M} = 1.8 \times 10^{-4} M_{\odot} \text{ yr}^{-1}$ computed by Justtanont et al. (2013), and assume it has not changed over the past 2800 yr, we calculate that $0.5 M_{\odot}$ has been ejected by the star in this time. If we instead use a lower rate such as $\dot{M} = 2 \times 10^{-5} M_{\odot} \text{ yr}^{-1}$, close to the value found by Heske et al. (1990) and in agreement with the findings of Decin et al. (2019), we correspondingly find less mass ejected in that time frame: only $0.06 M_{\odot}$. Clearly the mass-loss rate will significantly affect the longevity of this object. A further analysis of the CO properties of OH 30.1–0.7 would require a detailed radiative transfer analysis and is beyond the scope of this work, but will be the subject of a follow-up paper.

4.2.3 Evidence from other molecular species

As discussed in Section 4.1, the circumstellar chemistry of OH 30.1–0.7, especially the detection of NS, strongly suggests the presence of a companion star emitting UV in the inner wind. However, while the companion star should emit enough UV to facilitate the formation of NS through the photodissociation of N_2 (it should not be a red dwarf), the companion should not be too hot ($< 12000 \text{ K}$) to avoid photoionizing Na and K and inhibiting the formation of NaCl and KCl. Hence, the companion ought not to be a massive star and is likely to be a late B- to G-type star. This would put a companion in the mass range of ~ 1 to $4 M_{\odot}$, which also fulfills the criteria of being less massive than the AGB star, since the AGB star has evolved off the main sequence first.

Based on the temperature we calculated for the SO_2 lines (see Fig. 7 and Section 3.4), we can make an estimate of the average radial

temperature profile of the CSE. For this we take the SO_2 temperature to be 25 K and the average outer radius of SO_2 to be $1 \times 10^{17} \text{ cm}$ (corresponding to $\sim 1.7 \text{ arcsec}$, close to the average maximal extent of the SO_2 observations, see Fig. 8, and in good agreement with the radius of the outer SO_2 shell visible in Fig. A7). We take the inner temperature to be 2000 K close to the surface of the star (at $1.3 \times 10^{14} \text{ cm}$). Assuming a power-law radial temperature profile, $T(r) = T_{\star} (r/R_{\star})^{-\epsilon}$, we then calculate the exponent, $\epsilon = 0.66$. This is very close to the values found for lower mass-loss rate oxygen-rich AGB stars such as R Dor (0.65, Van de Sande et al. 2018) and W Hya (0.65, Khouri et al. 2014). If we try to include the average NaCl and KCl temperature (600 K, Fig. A9) at the apparent size of the emitting region, we find an implausibly high value of $\epsilon = 2.4$. This is most likely because the NaCl and KCl lines are not spatially resolved in our observations, as noted in Section 3.6. If we instead use the estimated radial temperature profile with $\epsilon = 0.66$ to calculate the average radius of the salt extents, we find $R_{\text{salt}} = 8 \times 10^{14} \text{ cm}$, which corresponds to $\sim 0.014 \text{ arcsec}$ (and $\sim 50 \text{ au}$), much smaller than our restoring beams (Table A1).

The companion is most likely driving the formation of the salt molecules. Hence, the above indicates that the separation between the AGB star and the likely main-sequence companion is significantly smaller than our minimum resolution of $\sim 0.3 \text{ arcsec} = 1200 \text{ au} \approx 2 \times 10^{16} \text{ cm}$. From our estimate of the average salt extent, we predict that observations with an angular resolution of 5 mas or better are required to resolve the structures in the inner wind. Further observations targeting molecular tracers of UV-emitting companions such as NS and SiN (Van de Sande & Millar 2022), will help us to understand the nature and orbit of such a companion.

Finally, we note that as well as affecting the circumstellar chemistry and gravitationally shaping the wind, a companion may also contribute to dust formation. Even wide binary companions (with semimajor axes on the order of hundreds of au) are essentially guaranteed to have orbital velocities in excess of the local sound speed, resulting in additional shocks in the wind. Freytag & Höfner (2023) found, in the context of stellar pulsations, that dust forms in the wake of shocks. Some evidence for this was seen by Coenegrachts et al. (2023) who found that an asymmetric dust distribution in the inner CSE of IK Tau correlated with a clumpy spiral of NaCl emission that may have formed partly as a result of shocks and other interactions with an unseen binary companion. Indeed, a more extreme version of companion-enhanced dust formation has long been known to take place for Wolf–Rayet stars, where dust forms at the nexus of colliding winds from massive stars (Usov 1991). The lack of prior direct detection of such a companion is not evidence of its absence, since the optically thick circumstellar dust would very strongly attenuate the signal at optical wavelengths (e.g. see the strong silicate absorption in fig. 1 of Justtanont et al. 2013).

4.3 On the nature of feature ‘B’

Feature B is detected in the continuum and lies 4.6 arcsec north and slightly west of the primary continuum peak. The lower limit on the separation is $1.8 \times 10^4 \text{ au}$. We extracted spectra around this peak, using a circular aperture with a radius of 0.3 arcsec, and did not detect any line emission above the noise aside from CO. As shown in our channel maps (Figs 3 and A5), this emission is most likely dominated by contamination from the background molecular cloud associated with W43. Inspecting the channel maps, there are no localized features in the CO emission associated with the position of feature B. As can be clearly seen in the channel maps and in the integrated intensity (zerth moment) map in Fig. 4, feature B

lies beyond any of the molecular emission associated with the AGB star. The size of feature B, when considering the 5σ contours, is approximately 0.8×0.5 arcsec.

In Section 2.2.2 we calculated the spectral indices of both the primary continuum peak and feature B. The spectral index of the primary peak, 2.8 ± 0.3 is in general agreement with past results for OH 30.1–0.7. For example Nicolaes et al. (2018) found spectral indices ranging from 2.13 at shorter (far-IR) wavelengths to 3.42 at long wavelengths, based on *Herschel*/PACS and *Herschel*/SPIRE observations. The lower limit of 3.6 for feature B is very high compared with commonly found spectral indices (see for example Juvela et al. 2015, who find values around 1–2 for cold cores). Nicolaes et al. (2018) found some comparably steep slopes for other evolved stars in their sample, notably at longer wavelengths ($> 100\mu\text{m}$) and for stars classed as OH/IR stars, similar to OH 30.1–0.7. When it comes to dust sources not associated with evolved stars, Kuan, Mehringer & Snyder (1996) found very steep spectral indices (up to 4.6) for Sagittarius B2(N). They conclude that this is most likely a result of a cold dust temperature and ice-coated grains and that Sgr B2(N) is a young object in an early stage of star formation. Whether our feature B is a similar background source to Sgr B2(N) or more closely associated with OH 30.1–0.7, cold ice-coated dust grains are a feasible explanation for the spectral index. Given that Justtanont et al. (2006) found evidence of H_2O ice in the dust around OH 30.1–0.7, the high spectral index of feature B could indicate a similar dust composition (i.e. ice-coated grains) but at a lower temperature since the feature is located further from the AGB star with much of the stellar light obscured by the circumstellar dust.

4.3.1 A direct association with OH 30.1–0.7?

If we draw a line connecting the primary continuum peak and feature B, we find it lies 20.5° west of north (clockwise on the observed maps in Fig. 1). The CO emission appears mostly round in shape (Fig. 3), however, there is a slight indentation in the direction of feature B. The centre of the indentation sits about $2.4''$ from the primary continuum peak, whereas on the opposite side the 5σ contour of the CO emission extends to around 3.1 arcsec from the primary continuum peak. This alignment could be coincidental, but it is also possible that feature B is contributing to the shaping of the CO emission or was formed through the same interaction as the CO indentation. This would imply that feature B is an object at the same distance as OH 30.1–0.7. Given the large separation between the CO indentation and feature B (around 2.2 arcsec), it is unlikely that the source of feature B is (at present) contributing to significant gravitational shaping of the AGB circumstellar environment. Decin et al. (2019) suggested that the indentation in the CO was a knot caused by a companion and its density wake. Another possibility is that feature B is causing the indentation through non-gravitational means, perhaps through high-energy UV photons, although this is difficult to reconcile with the cold dust implied by the spectral index.

A recent study of the S-type AGB star, W Aql, found that the F-type main-sequence companion, located around 200 au from the AGB star, contributed to the enhanced photodissociation of several molecular species, including SiO, SiS, CS, and HCN. The channels exhibiting such photodissociation bear some resemblance to the CO towards OH 30.1–0.7, although the CO indentation is less extreme. At 6000 K, the companion to W Aql did not produce enough energetic UV photons to significantly photodissociate CO. However, a hotter stellar companion could potentially affect the CO distribution. For example,

when analysing the impact of increasing the interstellar radiation field on the size of the CO envelopes of AGB stars, Saberi et al. (2019) found that, for stars with similar mass-loss rates and velocities to OH 30.1–0.7, doubling the strength of the interstellar radiation field reduced the size of the CO envelope by 14 per cent–20 per cent. High mass-loss rates result in smaller effects because of CO self-shielding. For example, Van de Sande & Millar (2022) found that including a hot 80 000 K (white dwarf) inner companion in their model did not significantly impact the CO distribution for higher mass-loss rates. If feature B is a hot star, it could act as a localized increase in the UV radiation field. Counter to this and unlike a companion located within the dust envelope of OH 30.1–0.7, it is unclear why such a UV source would be too attenuated to be detected from Earth but not too attenuated to affect the OH 30.1–0.7 CO envelope. Or why such a hot source would not destroy ice-covered dust grains. We also note that if feature B does correspond to a companion, this would not negate the potential presence of a closer companion that does shape the wind, even though feature B is too distant to be such a companion itself.

A more likely possibility is that feature B is a clump of dust (and perhaps undetected gas) that was ejected from OH 30.1–0.7 itself, perhaps as a pre-cursor to the present-day mass loss (beginning 2800 y-r ago, according to the CO extent) or owing to a complex binary interaction. The fact that the extent of the NS emission lies perpendicular to the projected separation between the AGB star and feature B lends some weight to this scenario. If feature B were ejected with the same velocity as the observed CO envelope, that would suggest an ejection around 3700 yr ago. However, if the ejection took place at the same time that mass loss was initiated, that would imply a velocity of 30 km s^{-1} , which is not implausibly high. If feature B is rather a high-velocity bullet, such as those seen for the carbon star V Hya (which has a relatively close companion on an eccentric orbit; Sahai, Scibelli & Morris 2016), then feature B could have been ejected more recently at a higher velocity. In both of these cases we are assuming that feature B was moving in the plane of the sky, making our time and distance estimates lower limits. An ejection scenario could, however, account for the apparent bridge between the primary continuum peak and feature B, if some ejecta were not accelerated as strongly.

4.3.2 A background source?

It is possible that feature B is a coincident background (or foreground) source. Given the line-of-sight proximity of OH 30.1–0.7 to the star-forming region W43, that is the most likely origin of background sources. The distance to W43 has been determined to be 5.5 ± 0.3 kpc (Zhang et al. 2014), significantly further than the 3.9 kpc to OH 30.1–0.7 (Engels et al. 2015). At the distance of W43, the extent of feature B would correspond to around 2800–4400 au. Placing it at the distance of OH 30.1–0.7 gives an extent of around 2000–3000 au. Both of these sizes are upper limits as they are not significantly larger than the beam size.

5 CONCLUSIONS

We present newly reduced ALMA observations of the intermediate-mass AGB star OH 30.1–0.7, including continuum and line emission. In the continuum maps, we identified a secondary peak (feature B), detected with a certainty of $> 10\sigma$, and we speculate that it is most likely ejecta associated with the AGB star, but may be a background source or a companion object. With a high lower limit for its spectral

index, feature B is likely composed of cold dust with ice covering the dust grains. We do not conclusively find any molecular emission associated with the feature, aside from some weak CO emission which may originate from background contamination rather than from that source.

We identified all the molecular emission present in the ALMA maps and spectra, finding rotational transitions arising from CO, SiS, CS, SO₂, NS, NaCl, and KCl. In addition to transitions in the ground vibrational state, we find several rotational transitions in vibrationally excited states from SiS (up to $v = 1$), NaCl (up to $v = 4$), and KCl (up to $v = 2$). Based on a simplified molecular line excitation analysis, we estimate the radial temperature profile as a power law with a similar index to other oxygen-rich AGB stars.

We interpret the presence of NS as indicative of a UV-emitting main-sequence companion in the inner regions of the OH 30.1–0.7 wind, with a separation smaller than our resolution limit of ~ 1200 au. The distribution of the NS emission suggests an eccentric binary orbit with an orbital plane that may be perpendicular to the direction of feature B. While the presence of NS suggests a UV emitting companion, the presence of NaCl and KCl require that the companion not be too hot. From this, we determine that the companion cannot be hotter than a late B-type main-sequence star, but may be as cool as a G-type star, and that a red dwarf star would not be able to trigger adequate NS formation.

Past observations and studies of OH 30.1–0.7 have indicated optically thick dust and CO emission – which our observations support – and have suggested that the AGB star may have entered a superwind phase at the tail end of the AGB. Our observations reveal a very steep drop off at the edges of the CO emission, both in the plane of the sky and in velocity space. This suggests that the bounds of the CO envelope are not truncated by the interstellar radiation field, as is usually the case. Instead it seems that the observed CO corresponds to a period of high mass loss beginning around 2800 years ago. This is a longer period than predicted by the superwind scenario, and a detailed explanation is deferred to a later study. We also conclude that any earlier period of lower mass loss (with a mass-loss rate $\dot{M} \leq 2 \times 10^{-5} M_{\odot} \text{yr}^{-1}$) would not be detectable in light of the recent high mass loss. We suggest companion-enhanced dust production, leading to a higher mass-loss rate, as a possible explanation for the current high mass-loss rate.

Additional higher angular resolution observations are required to fully understand OH 30.1–0.7 and its high mass loss and dense wind. In particular, we recommend observations with angular resolution of 5 mas or better to resolve possible companion interactions in the inner wind.

ACKNOWLEDGEMENTS

The authors thank Elvire De Beck for sharing the original data from her 2010 paper (De Beck et al. 2010). TD is supported in part by the Australian Research Council through a Discovery Early Career Researcher Award (DE230100183). This research is supported in part by the Australian Research Council Centre of Excellence for All Sky Astrophysics in 3 Dimensions (ASTRO 3D), through project number CE170100013. MVdS acknowledges the Oort Fellowship at Leiden Observatory. TJM’s research at QUB is supported by grant no. ST/T000198/1 from the STFC. HSPM thanks the Deutsche Forschungsgemeinschaft (DFG) for support through the collaborative research center SFB 1601 (project ID 500700252) subprojects A4 and Inf. KJ acknowledges the support from the Swedish National Space Agency (SNSA). JMCP was supported by STFC grant no. ST/T000287/1. SHJW acknowledges support from

the Research Foundation Flanders (FWO) through grant 1285221N. LD acknowledges support from the KU Leuven C1 MAESTRO grant C16/17/007, the FWO research grant G099720N, and the KU Leuven C1 BRAVE grant C16/23/009. This paper makes use of the following ALMA data: ADS/JAO.ALMA#2016.1.00005.S. ALMA is a partnership of ESO (representing its member states), NSF (USA) and NINS (Japan), together with NRC (Canada), MOST and ASIAA (Taiwan), and KASI (Republic of Korea), in cooperation with the Republic of Chile. The Joint ALMA Observatory is operated by ESO, AUI/NRAO and NAOJ. We acknowledge excellent support from the UK ALMA Regional Centre (UK ARC), which is hosted by the Jodrell Bank Centre for Astrophysics (JBCA) at the University of Manchester. The UK ARC Node is supported by STFC grant ST/P000827/1. Based on observations with the Atacama Pathfinder EXperiment (APEX) telescope with project ID: O-0102.F-9301B. APEX is a collaboration between the Max Planck Institute for Radio Astronomy, the European Southern Observatory, and the Onsala Space Observatory. Swedish observations on APEX are supported through Swedish Research Council grant no. 2017–00648. This research made use of ASTROPY (Astropy Collaboration 2013, 2018), SCIPY (Virtanen et al. 2020), MATPLOTLIB (Hunter 2007), and NUMPY (Harris et al. 2020).

DATA AVAILABILITY

The observational data used here are openly available through the data archives for ALMA (<https://almascience.nrao.edu/aq/>) and ESO for the APEX data (<http://archive.eso.org>). Custom ALMA data products that were produced for this study are available from TD or AMSR upon reasonable request.

REFERENCES

- Agúndez M., Fonfría J. P., Cernicharo J., Kahane C., Daniel F., Guélin M., 2012, *A&A*, 543, A48
- Astropy Collaboration, 2013, *A&A*, 558, A33
- Astropy Collaboration, 2018, *AJ*, 156, 123
- Baudry A. et al., 2023, *A&A*, 674, A125
- Belov S. P., Tretyakov M. Y., Kozin I. N., Klisch E., Winnemisser G., Lafferty W. J., Flaud J. M., 1998, *J. Mol. Spectrosc.*, 191, 17
- Briggs D. S., Schwab F. R., Sramek R. A., 1999, in Taylor G. B., Carilli C. L., Perley R. A. eds, ASP Conf. Ser. Vol. 180, Synthesis Imaging in Radio Astronomy II, p. 127, http://www.asppubs.org/a/volumes/table_of_contents/180
- Caris M., Lewen F., Winnemisser G., 2002, *Z. Nat. Teil A*, 57, 663
- Caris M., Lewen F., Müller H. S. P., Winnemisser G., 2004, *J. Mol. Struct.*, 695–696, 243
- Cernicharo J., Guélin M., 1987, *A&A*, 183, L10
- Cernicharo J., Marcelino N., Agúndez M., Guélin M., 2015, *A&A*, 575, A91
- Cherchneff I., 2012, *A&A*, 545, A12
- Cinquegrana G. C., Karakas A. I., 2022, *MNRAS*, 510, 1557
- Coenegrachts A., Danilovich T., De Ceuster F., Decin L., 2023, *A&A*, 678, A85
- Condon J. J., 1997, *PASP*, 109, 166
- Cortes P. et al., 2023, *ALMA Cycle 10 Technical Handbook*
- Cutri R. M. et al., 2012, Explanatory Supplement to the WISE All-Sky Data Release Products. <http://wise2.ipac.caltech.edu/docs/release/allsky/expsup/index.html>
- Danilovich T., De Beck E., Black J. H., Olofsson H., Justtanont K., 2016, *A&A*, 588, A119
- Danilovich T., Lombaert R., Decin L., Karakas A., Maercker M., Olofsson H., 2017a, *A&A*, 602, A14
- Danilovich T., Van de Sande M., De Beck E., Decin L., Olofsson H., Ramstedt S., Millar T. J., 2017b, *A&A*, 606, A124