



Reliability and Accuracy of Standard Reference Procedures for Measurements of Trunk and Arm Postures in Ergonomics

Downloaded from: <https://research.chalmers.se>, 2025-03-22 07:16 UTC




Citation for the original published paper (version of record):

Lind, C., Rhen, I., Forsman, M. (2025). Reliability and Accuracy of Standard Reference Procedures for Measurements of Trunk and Arm Postures in Ergonomics. *Bioengineering*, 12(1). <http://dx.doi.org/10.3390/bioengineering12010050>

N.B. When citing this work, cite the original published paper.

Article

Reliability and Accuracy of Standard Reference Procedures for Measurements of Trunk and Arm Postures in Ergonomics

Carl M. Lind ^{1,*}, Ida-Märta Rhen ^{1,2,3} and Mikael Forsman ^{1,2,4}

¹ Unit of Occupational Medicine, Institute of Environmental Medicine, Karolinska Institutet, 171 77 Stockholm, Sweden

² Centre for Occupational and Environmental Medicine, Stockholm County Council, 113 65 Stockholm, Sweden

³ Department of Industrial and Materials Science, Chalmers University of Technology, 412 96 Gothenburg, Sweden

⁴ School of Engineering Sciences in Chemistry, Biotechnology and Health, KTH Royal Institute of Technology, 141 57 Stockholm, Sweden

* Correspondence: carl.lind@ki.se

Abstract: Adequate reference procedures for obtaining the reference zero-angle position are important for precise and accurate posture measurements, but few studies have systematically investigated these. A limited number of previous studies suggest differences in accuracy between procedures, with some causing an underestimation of the true arm elevation angle when sensors are taped to the skin. The reliability of commonly used reference procedures for the measurement of the trunk posture is also not well explored, and alternative procedures may improve precision. Based on this identified gap, this study evaluated the test–retest reliability of the N-position (I-pose), i.e., the standard procedure for recording trunk postures, and compared it with two new alternative procedures. Additionally, the accuracy of the N-position for measuring arm elevation angles was compared with one alternative procedure. A total of 40 participants (22 women and 18 men) aged 26–70 years performed the reference procedures in a laboratory setting. Postures were recorded using a smart workwear system equipped with two inertial measurement units (IMUs) embedded in pockets within the workwear. For the trunk posture, the N-position showed a slight lack of test–retest reliability, while one of the alternative procedures demonstrated better test–retest reliability. For the arm posture, the N-position, which does not include lateral trunk inclination, resulted in a substantial underestimation of the arm elevation angle of approximately 15°, which is a novel finding. In contrast, the posture involving trunk inclination closely matched the targeted reference, with a difference of less than 2°. This study underscores the importance of selecting appropriate reference procedures to ensure precise and accurate posture measurements.

Keywords: measurement error; measurement bias; work postures; calibration procedures; posture assessment; standard posture; plumb line test; normal standing position; inclinometry; wearables



Academic Editors: Christina Zong-Hao Ma and Hong Fu

Received: 4 December 2024

Revised: 3 January 2025

Accepted: 7 January 2025

Published: 9 January 2025

Citation: Lind, C.M.; Rhen, I.-M.; Forsman, M. Reliability and Accuracy of Standard Reference Procedures for Measurements of Trunk and Arm Postures in Ergonomics.

Bioengineering **2025**, *12*, 50.

<https://doi.org/10.3390/bioengineering12010050>

[bioengineering12010050](https://doi.org/10.3390/bioengineering12010050)

Copyright: © 2025 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Musculoskeletal disorders (MSDs) of the lower back and neck are significant global health concerns, accounting for approximately 95 million disability-adjusted life years (DALYs) annually, while other musculoskeletal disorders collectively account for over 33 million DALYs worldwide [1]. Shoulder pain is prevalent globally with a seven-day prevalence of up to approximately 34% and a 12-month prevalence of up to around 55% [2].

The economic burden of work-related musculoskeletal disorders (WMSDs) and occupational accidents and diseases has been estimated at 3.3% of the gross domestic product in the European Union and 3.9% of the global gross domestic product [3]. WMSDs contribute to increased risks of short- and long-term work absenteeism, a reduced work capacity, and a premature exit from the labor market [4–6]. WMSDs are multifactorial, with both physical and psychosocial risk factors playing a role [7–9]. Important physical risk factors include strenuous postures and movements [7,10–16], strenuous and repetitive manual handling [7,10,12,16–22], and whole-body and hand–arm vibrations [7,10,23,24]. Accurate and precise measurements of these risk factors are important for establishing exposure–effect relationships and for risk assessment [25–29].

1.1. Sensor-Based Motion Capture Devices in Risk Prevention

Recent advancements in microelectronics have facilitated the increased use of sensor-based measurement instruments, such as triaxial accelerometers and inertial measurement units (IMUs), in research and occupational health practice. These tools enable more accurate assessments of work postures [27], complementing the traditional assessment that primarily relies on observation-based tools [25,26,30–32], enabling extended measurement durations from a few minutes to several hours or full workdays, providing more representative exposure measurements.

To increase the ease of use, motion devices can be integrated into workwear and connected to systems that enable the automated analysis and real-time monitoring of workers [27,33,34]. Examples of such sensor-based solutions include stretchable textile sensors attached on the outer surface of the workwear [35,36] and IMUs placed in embedded pockets in the workwear [37–40]. Stretchable textile sensors can be used to distinguish gross physical activity types [35,36], but generally exhibit lower accuracy compared to IMUs embedded in workwear. Embedding motion devices such as IMUs into workwear eliminates the need for skin attachment using tape or straps, which sometimes can be inconvenient and sensitive, as it may require partial undressing, access to private spaces, and anatomical knowledge for precise sensor placement.

1.2. Review of Reference Procedures for Accurate and Reliable Sensor-Based Measurements of Sagittal Trunk Inclination and Arm Elevation

In addition to accurate instrumentation, precise reference postures (i.e., procedures for obtaining the reference zero-angle position, typically a neutral body posture) are essential for obtaining accurate and reliable measurements and to make studies inter-comparable. While several studies have assessed, e.g., the reliability and inter-rater reliability of instruments and methods to assess the range of motion, such as for lumbar flexion [41–43], and the accuracy of inertia-based wearable motion capture systems against marker-based motion capture systems [44–47], less attention has been given to evaluating the reliability of reference positions for a neutral position for the trunk and upper arm. Ideally, a reference procedure (i.e., reference positions or postures and other arrangements such as instructions, assistance, and the use of assisting objects like chairs and external loads) should not only provide accurate and reliable measurements but also be practical for use in both laboratory and field studies.

To record the neutral posture for the trunk, arms, and head (or neck), Aarås and Stranden [48] proposed an upright, well-balanced position where the shoulders are relaxed and the upper arms hang naturally alongside the body without holding an external load. This reference procedure allows for simultaneous zero-posture recordings of the trunk, head, and arms.

The posture is often referred to as the natural, normal, or standard posture, the normal stance [49,50], or the N-position, N-posture, or I-pose [51] (hereafter to be referred to as the

N-position). Variations of the N-position have been widely used in ergonomics research to establish the reference zero-angle position for both the trunk (see Table 1) and the upper arm (see Table 2).

As shown in Table 1, the N-position appears to be the most used zero posture. Among the variations of the N-position, some studies specify that the feet should be close together or shoulder width apart; while some describe subjects looking forward, others specify using a visual mark or point at eye level. An alternative variation of the N-position, referred to as the Toe-position in this article, was used in studies by Fan et al. [52] and Forsman et al. [53], where subjects returned to the N-position from a toe stand.

Table 1. Reference posture used to record a neutral trunk posture, indicating 0° trunk inclination or flexion. All studies were conducted in real work environments, and the selection was based on those identified by Lind et al. [27].

Reference Posture	
Standing upright	Reference posture not reported ¹
[13,40,54–89]	[46,90–102]

Note: ¹ not reported or information cannot be found.

The N-position appears to be the dominant posture for recording the reference zero-angle position for the trunk posture. It is also used to establish the reference zero-angle position for the upper arm, alongside other reference procedures, as summarized in Table 2. While the N-position does not involve lateral trunk inclination or typically require the use of an external dumbbell, most other reference procedures include these elements, typically using a 2 kg dumbbell. One of these procedures, referred to here as the “Stockholm procedure”, involves lateral trunk inclination and an external weight and is performed either standing or seated [44,89,103–105].

Table 2. Reference posture used to record a neutral arm posture, indicating 0° arm elevation. All studies were conducted in real work environments, and the selection was based on those identified by Lind et al. [27].

Lateral Trunk Inclination				No Lateral Trunk Inclination				Posture Not Reported	
Standing		Seated		Standing		Seated			
weight	no weight	weight	no weight	weight	no weight	not reported	weight	no weight	
[89]	[106]	[52–57,64,67,74,83,88,91,104,106–118]	-	[119]	[14,75,79,120]	[65,70,71,73,76–78,87,121,122]	[103,123–125]	-	[46,80,84,86,93,95,97–99,101,102,126–132] ¹

Note: ¹ a weight was used.

However, the peer-reviewed literature provides limited information on the reproducibility and accuracy of these reference postures and similar approaches for measuring neutral trunk- and arm postures. Additionally, alternative procedures may offer improved reliability. Aarås and Strandén [48] evaluated the reliability of the N-position reference posture for the head and arms in five participants, reporting an average deviation from the midpoint of less than 1° for the head posture, 1.2° for arm flexion (i.e., arm elevation in the sagittal plane, forward), and 0.9° for arm abduction (i.e., arm elevation in the frontal plane, sideways). In a more recent study [69], median group differences before and after measurements were 3.8° for trunk sagittal inclination and 1.8° for trunk lateral inclination (Nastaran, Raffler. 2020. e-mail message to author, August 25). In our research, we have observed that this posture can occasionally be challenging to consistently adopt

without postural divergence. While sensor-based inclinometers provide highly accurate measurements, calibration-related errors of approximately 4° in the reference posture can be unacceptably large, potentially obscuring actual improvements following an intervention [133]. To our knowledge, no study in the ergonomics literature has systematically evaluated the reliability of this reference posture for trunk angles on a larger sample or the reliability of alternative procedures.

With regard to the arm posture, Dahlqvist et al. [106,134] observed considerable differences between reference positions involving lateral (sideways) trunk inclination when a seated posture with an external weight was compared to standing reference positions without an external weight. For example, Dahlqvist et al. [134] reported group root mean square differences of 8.5° and 14° for the left and right arms, respectively. Furthermore, Jackson et al. [105] evaluated the accuracy of upper arm inclinometry against visually projected lines between 30° and 150° , with incremental steps of 30° . The inclinometers (triaxial accelerometers) were taped to the arm and a seated reference position with lateral trunk inclination with an external weight was used as the reference position (0° arm elevation). They reported that the inclinometers systematically underestimated the arm elevation angle, for example, by 13.5° and 11.1° on average at 90° arm abduction and arm flexion, respectively. The divergences were potentially attributed (at least partly) to soft tissue artifacts [105,135,136], and the authors presented conversion equations to allow for correction for the bias. Notably, this bias was not observed by Hansson [137] who, in a letter to the editor, reported an average divergence of $\leq 1^\circ$ for the right and left arms when the participants held their arms at what was judged to be 90° .

1.3. Summary of Identified Research Gap

The identified literature highlights a lack of research on the reliability of reference postures for arm and trunk measurements. Additionally, it remains unclear whether the bias in arm elevation measurements reported by Jackson et al. [105] persists when motion capture devices are placed in the pockets of workwear, allowing for slight movement, rather than being taped directly to the arm.

1.4. Research Objective

Given these research gaps, the aim of this study was to evaluate the reliability of three reference positions for sagittal (forward/backward) trunk inclination and the accuracy of two reference positions for arm inclinometry using motion capture devices embedded in workwear.

2. Materials and Methods

2.1. Subjects

The study population (Table 3) comprised 22 women and 18 men, recruited from a working population pool at three organizations located in the Stockholm Region, Sweden. To ensure a broad working-age sample, the inclusion criteria required participants to be 18 years or older, currently employed or enrolled as students, and free from physical conditions, such as pain, disorders, or impairments that could hinder the execution of the reference procedures. Thirty-seven participants reported right-hand dominance and three reported left-hand dominance. Twenty participants had a BMI ≥ 23.0 , and eleven had a BMI ≥ 25.0 . There were no statistically significant differences in age or stature between these groups and the rest of the population. The study was approved by the Regional Ethics Committee in Stockholm, Sweden (2017/1586-31/4 and 2019-01206), and all participants provided written informed consent prior to participation.

Table 3. Summary of participant characteristics (SD, standard deviation).

Demographic Variables	Women, Mean (SD)	Men, Mean (SD)
N	22	18
Age (years)	42.0 (13.4)	42.5 (10.7)
Body mass (kg)	61.3 (9.7)	76.8 (10.6)
Stature (cm)	164.0 (6.5)	178.7 (8.4)
BMI (kg/m ²)	22.7 (2.8)	24.0 (2.2)

2.2. Instrumentation

Postural data for the trunk and upper arms were recorded using the Smart Workwear System [37,38]. The system included a stretchy workwear shirt (Model 9418, LiteWork, Snickers Workwear, Hultafors Group Sverige AB, Bollebygd, Sweden; fabric: 58% polyamid and 42% polyester 37.5[®]; mass: 165 g/m²) customized into a short-sleeved shirt with embedded pockets on the trunk and arms for IMUs (LPMS-B2, LP Research, Tokyo, Japan; dimensions: 39 × 39 × 8 mm; mass: 12 g). Participants were individually fitted with appropriately sized shirts (XS–L) to accommodate their body dimensions. To measure the trunk inclination, a single IMU was placed in the trunk pocket positioned on the upper back at the level of thoracic vertebrae 1–2 [45,52] (Figure 1). To measure the right upper arm elevation angle, a single IMU was placed in a pocket of the workwear enabling the upper edge of the IMU to be positioned approximately distal to the medial deltoideus muscle insertion [39,40,138] (Figure 1). The IMUs (Figure 1D) transmitted real-time data via Bluetooth to an Android 8.0 smartphone (Galaxy A5 2017, Samsung, Seoul, Republic of Korea), and the data were analyzed using the ErgoRiskLogger application, as described elsewhere [38]. Since the study focused on static postures, the IMU sampling rate was set to 25 Hz. The 3-axis accelerometer in the IMUs was set to a range, per axis, of ±4 G, and the 3-axis gyroscope was set to ±500°/s. A 16-bit resolution was used, followed by an extensive complementary Kalman filter (that is, a sensor fusion filter) to obtain the orientation of the sensor. As this study concerned static postures, it mainly used the accelerometer signals. According to the specification of the IMU, the static angle error was below 0.5°. More information on the Smart Workwear System is reported elsewhere [38].




Figure 1. Positions of the inertial measurement units (IMUs) (here highlighted with red circles) from the anterior view (A), the lateral view (B), and the posterior view (C). The motion tracking device (i.e., the IMU) used (D).

2.3. Experimental Protocol

Four different reference positions were performed in a fixed order (Figures 2 and 3). Trunk postures were assessed using three of the four positions, and arm postures were assessed using two positions. As shown in Figure 2, the evaluation began with recordings of the N-position, followed by the Stockholm-position. Subsequently, static arm elevation angles were recorded at a set angle of 90°. The N-position was then repeated. Finally, the Balance-position was repeated twice, and the session concluded with two repetitions of the Toe-Position.




Figure 2. The order of the reference procedures and set angle trials.




Figure 3. Description of the four reference procedures.

The set angle procedure involved holding the upper arm still for 5 s in a 90° arm elevation (abduction), followed by another 90° arm elevation (flexion), with assistance from an instructor. The set angles were performed with a straight arm, maintaining a straight elbow and wrist, with the palm facing downward. The instructor (one of two physiotherapists) aligned the participants' right arm by positioning the midline of the humerus parallel to the targeted abduction and inclination angles guided by the set angles printed on a large paper sheet (100 × 135 cm). This approach was modeled after the procedure described by Jackson et al. [105], but instead of visual projection, printed lines were used to eliminate the spatial distortion observed during preliminary pilot tests. Consistent with Jackson et al. [105], this assisted positioning was assumed to represent the "true" angle of the upper arm.

To record the postures of the trunk and arm in the version of the ErgoRiskLogger application we used, a global (original) reference recording was first made (the first N-position shown in Figure 2), and the other recordings were derived from this as a base. For arm postures, the global (original) value for the Stockholm-position was derived by subtracting the angular elevation difference (delta) between the global reference recording (i.e., 1st N-position) and the Stockholm-position. To illustrate this, if an elevation angle of 15° was recorded for the Stockholm-position, this value was equal to the difference compared to the global reference recording (i.e., 1st N-position). In the same way, the measured elevation at the set angle of 90° arm abduction and 90° arm flexion for the Stockholm-position was estimated as the individually measured angle in the global reference recording, with the addition of the individual delta.

2.4. Data Analysis

The data were checked for normality using the Shapiro–Wilk test ($\alpha = 0.05$) and by visually inspecting histograms. For non-normally distributed data, the Wilcoxon signed-rank test was applied, while the paired samples *t*-test was used for normally distributed data.

For the trunk posture, the average trunk sagittal inclination angle was compared pairwise among the reference procedures using the Wilcoxon signed-rank test, as one of the three pairs was non-normally distributed.

The test–retest reliability was evaluated through pairwise comparisons among the three reference procedures using the Wilcoxon signed-rank test, as all three pairs were non-normally distributed.

For the arm posture, the differences in the arm elevation between the N-position and the Stockholm-position were compared pairwise at the set angles of 90° abduction and 90° flexion using the paired samples *t*-test.

Statistical analyses were performed using MATLAB R2023b Update 4 (The MathWorks, Inc., Natick, MA, USA), while normality tests were conducted using R (The R Foundation, version 2024.09.0). The significance level was set at $\alpha = 0.01$.

3. Results

The results are based on 240 recordings of trunk postures and 160 recordings of arm postures, encompassing all 40 participants with no missing data.

3.1. Trunk Posture

The group mean differences in trunk inclination angles in the sagittal plane relative to the N-position are shown in Table 4. At the group level, the Balance-position resulted in a slightly more forward trunk inclination angle relative to the N-position (0.44°), while the Toe-position resulted in a slightly more backward trunk inclination angle (−0.33°); however, these differences were not statistically significant ($p > 0.01$).

Table 4. Group mean differences in trunk angles in the sagittal plane (forward/backward) relative to the N-position and the statistical differences (Wilcoxon signed-rank test) of the group differences.

N-position	Angular Differences			Statistical Differences		
	Balance-position	Toe-position	N-position vs. Balance-position	N-position vs. Toe-position	Balance-position vs. Toe-position	
Ref.	0.44°	−0.33°	$p = 0.095$	$p = 0.622$	$p = 0.067$	

Notes: Angular differences with positive values indicate a forward sagittal trunk inclination relative to the reference, and negative values indicate a backward sagittal trunk inclination relative to the reference.

Trunk Posture—Reliability

The test–retest reliability of the reference postures for trunk angles in the sagittal plane is presented in Table 5, based on data from 40 subjects, each performing two repetitions of each position (no missing values). The group mean absolute difference for the N-position and the Balance-position was 1.60° and 1.56°, respectively, with individual maximum differences exceeding 5°. The Toe-position demonstrated the lowest group mean absolute difference at 0.72°, i.e., the highest test–retest reliability, significantly higher than the two other reference postures ($p < 0.005$).

Table 5. Average absolute test–retest differences in the sagittal trunk inclination (forward/backward) of the reference positions to record a neutral trunk angle, first versus second trial, expressed as the mean (standard deviation, SD), and the statistical differences (Wilcoxon signed-rank test) of the group mean differences. The results are based on 40 subjects, each performing two repetitions of each reference position.

Absolute Group Differences (SD) for 1st vs. 2nd Recording			Statistical Differences		
N-position	Balance-position	Toe-position	N-position vs. Balance-position	N-position vs. Toe-position	Balance-position vs. Toe-position
1.60° (1.37°)	1.56° (1.29°)	0.72° (0.54°)	$p = 0.717$	$p < 0.005$	$p < 0.001$

3.2. Arm Posture

Arm Posture—Accuracy

The corresponding values for the N-position and Stockholm-position at the 90° set angle for abduction and flexion are displayed in Figure 4 and Appendix A.

For arm abduction, using the N-position as a reference resulted in an average elevation angle of 74.1° (SD 4.7), which is 15.9° lower than the assumed true angle. In comparison, the Stockholm-position produced an average elevation angle of 88.3° (SD of 5.5), which was 1.7° lower than the assumed true angle.

For arm flexion, the N-position yielded an average elevation angle of 74.4° (SD of 3.9), which was 15.6° lower than the assumed true angle. In comparison, the Stockholm-position resulted in an average elevation angle of 88.6° (SD of 4.7), i.e., 1.4° lower than the assumed true angle. Hence, the Stockholm-position consistently yielded values that were substantially closer to the assumed true angle, and the angular differences between the N-position and the Stockholm-position at the 90° set angle were statistically significant (paired samples t-test) for both arm abduction and arm flexion ($p < 0.001$).




Figure 4. Box plots of arm elevation angles (abduction and flexion) at 90° set angles (yellow dotted line) for the two reference positions: N-position and Stockholm-position. The box represents the interquartile range (25th to 75th percentiles) and the median, while the mean is marked with an “x”. Vertical lines extending from the box indicate adjacent values, which are the most extreme values within 1.5 times the interquartile range for each group.

4. Discussion

4.1. General Summary of the Findings

This study, which included 40 subjects, evaluated commonly used reference procedures for obtaining the reference zero-angle positions for measurements of trunk and arm postures. The findings support the conclusion that the choice of reference procedure is critical, as it can significantly affect the reliability and accuracy of technical measurements using sensor-based inclinometers.

For recordings of trunk postures, the results of this study suggest that the test–retest reliability can be improved by replacing the commonly used reference posture (referred to as the N-position) with an alternative reference procedure (referred to as the Toe-position) where the participants arrive at a balanced upright posture from a toe stance.

For recordings of arm postures, the results show that the commonly used N-position, performed in an upright posture without lateral inclination, can lead to substantial and systematic underestimation of arm elevation angles relative to the reference, with errors of approximately 15° for both abduction and flexion if the bias is not corrected. In contrast, the use of an alternative reference procedure (position), i.e., the upright Stockholm-position, which involves lateral trunk inclination and the use of an external weight, resulted in group mean values that closely matched the reference angles, with deviations of less than 2°. These findings highlight the importance of incorporating lateral trunk inclination, and possibly an external weight, to improve measurement accuracy.

4.2. General Interpretation of the Results and Comparisons with Related Literature

This study contributes to the identified research gap by providing new information on the reliability of a commonly used procedure for achieving accurate recordings of a neutral trunk posture, based on a relatively large and diverse sample of men and women spanning various age groups.

4.2.1. Trunk Sagittal Inclination

Regarding trunk postures, the group mean (absolute) test–retest differences in the trunk sagittal inclination in the current study were smaller for all three positions evaluated (i.e., all $\leq 1.60^\circ$) compared to the median group differences of 3.8° in the study by Raffler et al. [69]. Notably, Raffler et al. [69] measured the postures before and after the measurement of work activities, which lasted about 1–2 h. One potential reason for the difference could be related to changes in the position of the motion device, which can occur, especially during prolonged use, dynamic activities, or when motion devices are insufficiently secured. The group mean (absolute) test–retest differences in the trunk sagittal inclination in the current study, especially for the Toe-position (0.72°), were close to the test–retest values for the neck (head) of 0.85° reported by Aarås and Strandén [48].

4.2.2. Arm Elevation

Concerning arm postures, contrary to findings from earlier studies by Bernmark and Wiktorin [44], Jackson et al. [105], Lind and Forsman [139], and the related study by Lind et al. [140], this study did not confirm a considerable bias in arm posture recordings at a 90° arm elevation. Instead, and in agreement with what was reported by Hansson [137], who used a seated reference posture, the observed values in the current study closely approximated the assumed true angle, with only slightly lower values of $<2^\circ$. This can be compared with the difference from the assumed true angle noted by Jackson et al. [105], which averaged 13.5° and 11.1° at a 90° arm abduction and arm flexion, respectively. Previous research has developed equations to harmonize data across different instruments and calculations [52,53,141] or to correct inaccuracies potentially caused by soft tissue artifacts and reference procedures [105]. The large underestimation of arm elevation angles (approximately 15°) is likely caused by the arm, although relaxed, being more abducted when leaning against the body.

4.3. Strengths and Limitations

It should be noted that this study is limited to the sagittal trunk inclination and arm elevation. Although these are the most common exposure metrics for postures of the trunk and upper arm in the ergonomics literature related to WMSD outcomes [27], future studies are needed to explore the reference position in relation to, e.g., trunk sagittal inclination.

Regarding reference procedures for obtaining the reference zero-angle position for the trunk, while the study's findings suggest that the reliability can be improved through alternative procedures, the accuracy of these reference positions was not directly evaluated. Furthermore, it is debated which posture truly represents the “natural” or “neutral” upright posture [49,50]. However, since high accuracy depends on precision, some conclusions can still be drawn regarding the potential advantages of adopting alternative procedures to enhance both the reliability and overall measurement accuracy. A potential limitation was that the reference procedures were not performed in a randomized order, which could introduce learning effects, particularly for reference positions performed later. However, all participants were already familiarized with the reference procedures from a prior trial [140], with the exception of the Balance-position and Toe-position. Therefore, it seems unlikely that the relatively higher test–retest reliability observed for the Toe-position was due to learning effects.

With respect to reference procedures for obtaining the reference zero-angle position for the arm, while the results provide important findings, the accuracy was evaluated only at a 90° arm elevation angle (flexion and abduction). Consequently, the bias may vary in magnitude at other angles. Furthermore, additional information is necessary to develop conversion equations for correcting bias, particularly for the N-position, which resulted

in substantial bias. Although only the standing version of the Stockholm procedure was included in this study, preliminary findings from a related study suggest that the differences between the standing and seated variations are small [140].

Additionally, the global (original) reference recording for arm abduction and flexion for the Stockholm-position was derived by subtracting the angular elevation difference between the first N-position and the Stockholm-position (as described in Section 2.3). This approach was necessitated by limitations in the version of the Smart Workwear System used. Preferably, another global (original) reference recording should have been obtained when having the participants stand in the Stockholm-position. The present method introduces an approximation in determining the angle between the Stockholm-position and the 90° arm elevations. Nevertheless, this adjustment likely has a negligible effect on the abduction angle recorded at the 90° set angle, and a negligible-to-minimal effect on the flexion angle. Future studies are, however, recommended to further investigate this.

A strength of this study is the inclusion of a relatively large and diverse sample of subjects, encompassing both women and men across a wide age range (26–70 years). This improves the generalizability to other populations. The stature and body mass of the women in this study were slightly lower than those reported among a Swedish female working population [142] with a median stature of 167.3 cm and median body mass of 64 kg. This stature approximates that of US women aged 40–49 years (162.6 cm) but with a higher body mass (75.2 kg) [143]. The men's stature and body mass were approximately equal to the median stature of 177.9 cm and body mass of 75 kg among a Swedish male working population [142]. This stature approximates that of US men aged 40–49 years (176.5 cm), who have a higher body mass (90.5 kg) [143].

All measurements were recorded with static (isometric) postures, without involving any movement. The potential improvement in the measurement accuracy that can be attributed to having the sensors embedded in tight workwear (as opposed to taped) may differ during tasks involving high movement velocities, especially if the fit of the workwear is loose in some parts where the sensors are embedded, as pointed out by Hoareau et al. [144].

4.4. Practical Implications and Future Research

The bias attributed to reference procedures seems to have been underestimated, especially as measurements from technical measurement instruments such as IMUs and accelerometers are often regarded as having superior accuracy and precision when compared to self-reports and systematic observations [54,145–148].

The main implication of this study is that an upright neutral position is not recommended for recording the neutral position of the upper arm. Instead, the reference position should include lateral trunk inclination. As a result, findings from studies that assume that the N-position represents zero arm elevation (without additional corrective measures) may not be directly comparable to those that use other reference positions. Furthermore, it is questionable whether the N-position should be used at all for recording arm postures without corrections. Given the relatively common use of the N-position, as shown in Table 2, caution is warranted when interpreting the results of studies that did not account for the bias associated with the N-position as observed in the current study. This also underscores the need for conversion methods to convert data obtained from different reference procedures, particularly when one involves trunk inclination and the other does not.

While the current study demonstrated that a posture including trunk inclination and an external load can yield relatively accurate arm elevation angles (both abduction and flexion), the divergent results from the related study by Lind et al. [140], as well as findings from Jackson et al. [105], Bernmark and Wiktorin [44], and Lind and Forsman [139],

underscore the need for further research to better understand the reasons behind these discrepancies. The results of the current study highlight the need for the conversion of postures obtained using different reference positions to ensure accuracy and comparability between studies.

To improve the test–retest reliability in recording a neutral trunk position, and consequently the accuracy of the N-position, future studies could explore the feasibility and efficiency of increasing the number of measurements, allowing for the calculation of an average value based on multiple recordings taken before and after the work activity, or alternatively, consider using the Toe-position.

5. Conclusions

The choice of reference procedures for obtaining reference zero-angle positions can significantly impact the reliability and accuracy of technical measurements using sensor-based inclinometers. For recordings of a zero-angle trunk position in the sagittal plane, the commonly used N-position (or N-posture, or I-pose) was observed to exhibit some imprecision. An alternative posture, where participants arrived at the N-position from a toe stand, improved the test–retest reliability.

For recordings of arm elevation, the N-position was compared with an alternative procedure (the Stockholm-position), which includes lateral trunk inclination. The N-position resulted in a substantial average bias of 15°, raising concerns about its use without adjustments to mitigate this error. In contrast, the Stockholm-position demonstrated higher accuracy, with deviations averaging less than 2°, making it the superior option of the two.

The findings of this study can guide the selection of appropriate reference procedures for both field and laboratory applications. Additionally, the findings highlight the importance of incorporating lateral trunk inclination when recording the reference zero-angle position of the arm.

Author Contributions: Conceptualization, C.M.L., I.-M.R. and M.F.; data curation, C.M.L.; methodology, C.M.L., I.-M.R. and M.F.; formal analysis, C.M.L. and M.F.; investigation, C.M.L. and I.-M.R.; writing—original draft preparation, C.M.L.; writing—review and editing, C.M.L., I.-M.R. and M.F.; project administration, C.M.L.; funding acquisition, C.M.L., I.-M.R. and M.F. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by AFA Insurance (Afa Försäkring), grant numbers 180098 and 180254.

Institutional Review Board Statement: The study was conducted in accordance with the Declaration of Helsinki and approved by the Regional Ethics Committee in Stockholm, Sweden (2017/1586–31/4 and 2019-01206).

Informed Consent Statement: Informed consent was obtained from all participants involved in the study and from the person in the figures showing the system and reference procedures.

Data Availability Statement: The raw data supporting the conclusions of this article will be made available by the authors on request.

Acknowledgments: Johan Eriksson is gratefully acknowledged for his contribution to the data collection. We would also like to express our gratitude to the participants.

Conflicts of Interest: The authors declare no conflicts of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

Appendix A

Table A1. Mean (SD) arm elevation angles (abduction and flexion) at 90° set angles for two reference positions: N-position and Stockholm-position.

Arm Abduction		Arm Flexion	
N-position	Stockholm-position	N-position	Stockholm-position
74.1 (4.7)	88.3 (5.5)	74.4 (3.9)	88.6 (4.7)

References

- Kassebaum, N.J.; Arora, M.; Barber, R.M.; Bhutta, Z.A.; Brown, J.; Carter, A.; Casey, D.C.; Charlson, F.J.; Coates, M.M.; Coggeshall, M.; et al. Global, regional, and national disability-adjusted life-years (DALYs) for 315 diseases and injuries and healthy life expectancy (HALE), 1990–2015: A systematic analysis for the Global Burden of Disease Study 2015. *Lancet* **2016**, *388*, 1603–1658. [[CrossRef](#)]
- Lucas, J.; van Doorn, P.; Hegedus, E.; Lewis, J.; van der Windt, D. A systematic review of the global prevalence and incidence of shoulder pain. *BMC Musculoskelet. Disord.* **2022**, *23*, 1073. [[CrossRef](#)]
- Tompa, E.; Mofidi, A.; van den Heuvel, S.; van Bree, T.; Michaelsen, F.; Jung, Y.; Porsch, L.; van Emmerik, M. *The Value of Occupational Safety and Health and the Societal Costs of Work-Related Injuries and Diseases*; Publications Office of the European Union: Luxembourg, 2019. [[CrossRef](#)]
- Sundstrup, E.; Andersen, L.L. Joint association of physical and psychosocial working conditions with risk of long-term sickness absence: Prospective cohort study with register follow-up. *Scand. J. Public Health* **2021**, *49*, 132–140. [[CrossRef](#)] [[PubMed](#)]
- Pedersen, J.; Bjorner, J.B.; Andersen, L.L. Physical work demands and expected labor market affiliation (ELMA): Prospective cohort with register-follow-up among 46 169 employees. *Scand. J. Public Health* **2022**, *48*, 641–650. [[CrossRef](#)] [[PubMed](#)]
- Badarin, K.; Hemmingsson, T.; Hillert, L.; Kjellberg, K. The impact of musculoskeletal pain and strenuous work on self-reported physical work ability: A cohort study of Swedish men and women. *Int. Arch. Occup. Environ. Health* **2022**, *95*, 939–952. [[CrossRef](#)]
- NRC. *Musculoskeletal Disorders and the Workplace: Low Back and Upper Extremities*; National Academies Press: Washington, DC, USA, 2001. [[CrossRef](#)]
- Widanarko, B.; Legg, S.; Devereux, J.; Stevenson, M. The combined effect of physical, psychosocial/organisational and/or environmental risk factors on the presence of work-related musculoskeletal symptoms and its consequences. *Appl. Ergon.* **2014**, *45*, 1610–1621. [[CrossRef](#)] [[PubMed](#)]
- Hauke, A.; Flintrop, J.; Brun, E.; Rugulies, R. The impact of work-related psychosocial stressors on the onset of musculoskeletal disorders in specific body regions: A review and metaanalysis of 54 longitudinal studies. *Work Stress* **2011**, *25*, 243–256. [[CrossRef](#)]
- NIOSH. *Musculoskeletal Disorders and Workplace Factors*; National Institute for Occupational Safety and Health: Cincinnati, OH, USA, 1997.
- Hoogendoorn, W.E.; Bongers, P.M.; de Vet, H.C.; Douwes, M.; Koes, B.W.; Miedema, M.C.; Ariens, G.A.; Bouter, L.M. Flexion and rotation of the trunk and lifting at work are risk factors for low back pain: Results of a prospective cohort study. *Spine* **2000**, *25*, 3087–3092. [[CrossRef](#)] [[PubMed](#)]
- da Costa, B.R.; Vieira, E.R. Risk factors for work-related musculoskeletal disorders: A systematic review of recent longitudinal studies. *Am. J. Ind. Med.* **2010**, *53*, 285–323. [[CrossRef](#)]
- Gupta, N.; Bjerregaard, S.S.; Yang, L.; Forsman, M.; Rasmussen, C.L.; Rasmussen, C.D.N.; Clays, E.; Holtermann, A. Does occupational forward bending of the back increase long-term sickness absence risk? A 4-year prospective register-based study using device-measured compositional data analysis. *Scand. J. Work Environ. Health* **2022**, *48*, 651–661. [[CrossRef](#)] [[PubMed](#)]
- Gupta, N.; Rasmussen, C.L.; Forsman, M.; Sogaard, K.; Holtermann, A. How does accelerometry-measured arm elevation at work influence prospective risk of long-term sickness absence? *Scand. J. Work Environ. Health* **2022**, *48*, 137–147. [[CrossRef](#)] [[PubMed](#)]
- Versloot, A.H.C.; Jackson, J.A.; van Rijn, R.M.; Elbers, R.G.; Sogaard, K.; Macri, E.M.; Koes, B.; Burdorf, A.; Chiarotto, A.; Gerger, H. Physical and psychosocial work-related exposures and the occurrence of disorders of the shoulder: A systematic review update. *Appl. Ergon.* **2024**, *118*, 104277. [[CrossRef](#)] [[PubMed](#)]
- Jahn, A.; Andersen, J.H.; Christiansen, D.H.; Seidler, A.; Dalbøge, A. Occupational mechanical exposures as risk factor for chronic low-back pain: A systematic review and meta-analysis. *Scand. J. Work Environ. Health* **2023**, *49*, 453–465. [[CrossRef](#)]
- van Rijn, R.M.; Huisstede, B.M.; Koes, B.W.; Burdorf, A. Associations between work-related factors and the carpal tunnel syndrome—a systematic review. *Scand. J. Work Environ. Health* **2009**, *35*, 19–36. [[CrossRef](#)] [[PubMed](#)]
- Garg, A.; Boda, S.; Hegmann, K.T.; Moore, J.S.; Kapellusch, J.M.; Bhojar, P.; Thiese, M.S.; Merryweather, A.; Deckow-Schaefer, G.; Bloswick, D.; et al. The NIOSH lifting equation and low-back pain, part 1: Association with low-back pain in the BackWorks prospective cohort study. *Hum. Factors* **2014**, *56*, 6–28. [[CrossRef](#)] [[PubMed](#)]

19. Garg, A.; Kapellusch, J.M.; Hegmann, K.T.; Moore, J.S.; Boda, S.; Bhojar, P.; Thiese, M.S.; Merryweather, A.; Deckow-Schaefer, G.; Bloswick, D.; et al. The NIOSH lifting equation and low-back pain, Part 2: Association with seeking care in the backworks prospective cohort study. *Hum. Factors* **2014**, *56*, 44–57. [[CrossRef](#)] [[PubMed](#)]
20. Hoozemans, M.J.; Knelange, E.B.; Frings-Dresen, M.H.; Veeger, H.E.; Kuijjer, P.P. Are pushing and pulling work-related risk factors for upper extremity symptoms? A systematic review of observational studies. *Occup. Environ. Med.* **2014**, *71*, 788–795. [[CrossRef](#)] [[PubMed](#)]
21. Kozak, A.; Schedlbauer, G.; Wirth, T.; Euler, U.; Westermann, C.; Nienhaus, A. Association between work-related biomechanical risk factors and the occurrence of carpal tunnel syndrome: An overview of systematic reviews and a meta-analysis of current research. *BMC Musculoskelet. Disord.* **2015**, *16*, 231. [[CrossRef](#)] [[PubMed](#)]
22. Fox, R.R.; Lu, M.L.; Occhipinti, E.; Jaeger, M. Understanding outcome metrics of the revised NIOSH lifting equation. *Appl. Ergon.* **2019**, *81*, 102897. [[CrossRef](#)] [[PubMed](#)]
23. Nilsson, T.; Wahlström, J.; Burström, L. Hand-arm vibration and the risk of vascular and neurological diseases—A systematic review and meta-analysis. *PLoS ONE* **2017**, *12*, e0180795. [[CrossRef](#)] [[PubMed](#)]
24. Burström, L.; Nilsson, T.; Wahlström, J. Whole-body vibration and the risk of low back pain and sciatica: A systematic review and meta-analysis. *Int. Arch. Occup. Environ. Health* **2015**, *88*, 403–418. [[CrossRef](#)]
25. Rose, L.M.; Eklund, J.; Nord Nilsson, L.; Barman, L.; Lind, C.M. The RAMP package for MSD risk management in manual handling—A freely accessible tool, with website and training courses. *Appl. Ergon.* **2020**, *86*, 103101. [[CrossRef](#)]
26. Lind, C.M.; Forsman, M.; Rose, L.M. Development and evaluation of RAMP II—A practitioner’s tool for assessing musculoskeletal disorder risk factors in industrial manual handling. *Ergonomics* **2020**, *63*, 477–504. [[CrossRef](#)]
27. Lind, C.M.; Abtahi, F.; Forsman, M. Wearable Motion Capture Devices for the Prevention of Work-Related Musculoskeletal Disorders in Ergonomics-An Overview of Current Applications, Challenges, and Future Opportunities. *Sensors* **2023**, *23*, 4259. [[CrossRef](#)] [[PubMed](#)]
28. Lind, C.M.; Rose, L.M. Shifting to proactive risk management: Risk communication using the RAMP tool. *Agron. Res.* **2016**, *14*, 513–524.
29. Cantley, L.F.; Taiwo, O.A.; Galusha, D.; Barbour, R.; Slade, M.D.; Tessier-Sherman, B.; Cullen, M.R. Effect of systematic ergonomic hazard identification and control implementation on musculoskeletal disorder and injury risk. *Scand. J. Work Environ. Health* **2014**, *40*, 57–65. [[CrossRef](#)] [[PubMed](#)]
30. Lind, C.M.; Forsman, M.; Rose, L.M. Development and evaluation of RAMP I—A practitioner’s tool for screening of musculoskeletal disorder risk factors in manual handling. *Int. J. Occup. Saf. Ergon.* **2019**, *25*, 165–180. [[CrossRef](#)]
31. Eliasson, K.; Lind, C.M.; Nyman, T. Factors influencing ergonomists’ use of observation-based risk-assessment tools. *Work* **2019**, *64*, 93–106. [[CrossRef](#)]
32. Lowe, B.D.; Dempsey, P.G.; Jones, E.M. Ergonomics assessment methods used by ergonomics professionals. *Appl. Ergon.* **2019**, *81*, 102882. [[CrossRef](#)]
33. Lind, C.M. A Rapid Review on the Effectiveness and Use of Wearable Biofeedback Motion Capture Systems in Ergonomics to Mitigate Adverse Postures and Movements of the Upper Body. *Sensors* **2024**, *24*, 3345. [[CrossRef](#)]
34. Lind, C.M. Effectiveness of Sensors-Based Augmented Feedback in Ergonomics to Reduce Adverse Biomechanical Exposure in Work-Related Manual Handling-A Rapid Review of the Evidence. *Sensors* **2024**, *24*, 6977. [[CrossRef](#)]
35. Esfahani, M.I.M.; Nussbaum, M.A.; Kong, Z.Y. Using a smart textile system for classifying occupational manual material handling tasks: Evidence from lab-based simulations. *Ergonomics* **2019**, *62*, 823–833. [[CrossRef](#)] [[PubMed](#)]
36. Mokhlespour Esfahani, M.I.; Nussbaum, M.A. Classifying Diverse Physical Activities Using “Smart Garments”. *Sensors* **2019**, *19*, 3133. [[CrossRef](#)] [[PubMed](#)]
37. Lind, C.M.; Sandsjö, L.; Mahdavian, N.; Högberg, D.; Hanson, L.; Diaz Olivares, J.A.; Yang, L.; Forsman, M. Prevention of Work-Related Musculoskeletal Disorders Using Smart Workwear—The Smart Workwear Consortium. In *Human Systems Engineering and Design*; Ahram, T., Karwowski, W., Taiar, R., Eds.; Springer Nature: Cham, Switzerland, 2019; Volume 876, pp. 477–483. [[CrossRef](#)]
38. Lind, C.M.; Diaz-Olivares, J.A.; Lindecrantz, K.; Eklund, J. A Wearable Sensor System for Physical Ergonomics Interventions Using Haptic Feedback. *Sensors* **2020**, *20*, 6010. [[CrossRef](#)]
39. Lind, C.M.; Yang, L.; Abtahi, F.; Hanson, L.; Lindecrantz, K.; Lu, K.; Forsman, M.; Eklund, J. Reducing postural load in order picking through a smart workwear system using real-time vibrotactile feedback. *Appl. Ergon.* **2020**, *89*, 103188. [[CrossRef](#)] [[PubMed](#)]
40. Lind, C.M.; De Clercq, B.; Forsman, M.; Grootaers, A.; Verbrugge, M.; Van Dyck, L.; Yang, L. Effectiveness and usability of real-time vibrotactile feedback training to reduce postural exposure in real manual sorting work. *Ergonomics* **2023**, *66*, 198–216. [[CrossRef](#)] [[PubMed](#)]
41. Saur, P.M.; Ensink, F.B.; Frese, K.; Seeger, D.; Hildebrandt, J. Lumbar range of motion: Reliability and validity of the inclinometer technique in the clinical measurement of trunk flexibility. *Spine* **1996**, *21*, 1332–1338. [[CrossRef](#)] [[PubMed](#)]

42. Tyson, S. A Systematic Review of Methods to Measure Posture. *Phys. Ther. Rev.* **2003**, *8*, 45–50. [[CrossRef](#)]
43. Fraeulin, L.; Holzgreve, F.; Brinkbäumer, M.; Dziuba, A.; Friebe, D.; Klemz, S.; Schmitt, M.; Theis, A.A.; Tenberg, S.; van Mark, A.; et al. Intra- and inter-rater reliability of joint range of motion tests using tape measure, digital inclinometer and inertial motion capturing. *PLoS ONE* **2020**, *15*, e0243646. [[CrossRef](#)] [[PubMed](#)]
44. Bernmark, E.; Wiktorin, C. A triaxial accelerometer for measuring arm movements. *Appl. Ergon.* **2002**, *33*, 541–547. [[CrossRef](#)] [[PubMed](#)]
45. Korshøj, M.; Skotte, J.H.; Christiansen, C.S.; Mortensen, P.; Kristiansen, J.; Hanisch, C.; Ingebrigtsen, J.; Holtermann, A. Validity of the Acti4 software using ActiGraph GT3X+accelerometer for recording of arm and upper body inclination in simulated work tasks. *Ergonomics* **2014**, *57*, 247–253. [[CrossRef](#)]
46. Schall, M.C., Jr.; Fethke, N.B.; Chen, H.; Oyama, S.; Douphrate, D.I. Accuracy and repeatability of an inertial measurement unit system for field-based occupational studies. *Ergonomics* **2016**, *59*, 591–602. [[CrossRef](#)]
47. Yang, L.; Grooten, W.J.A.; Forsman, M. An iPhone application for upper arm posture and movement measurements. *Appl. Ergon.* **2017**, *65*, 492–500. [[CrossRef](#)] [[PubMed](#)]
48. Aarås, A.; Strandén, E. Measurement of postural angles during work. *Ergonomics* **1988**, *31*, 935–944. [[CrossRef](#)]
49. Barra-López, M.E. The standard posture is a myth: A scoping review. *J. Rehabil. Med.* **2024**, *56*, jrm41899. [[CrossRef](#)] [[PubMed](#)]
50. Xue, R.; Liu, D.; Shen, Y. The differences in whole-body sagittal alignment between different postures in young, healthy adults. *BMC Musculoskelet. Disord.* **2020**, *21*, 696. [[CrossRef](#)]
51. Lee, W.; Lin, J.H.; Bao, S. Inter-rater reliability of an inertial measurement unit sensor-based posture-matching method: A pilot study. *Int. J. Ind. Ergon.* **2020**, *80*, 103025. [[CrossRef](#)]
52. Fan, X.; Lind, C.M.; Rhen, I.M.; Forsman, M. Effects of Sensor Types and Angular Velocity Computational Methods in Field Measurements of Occupational Upper Arm and Trunk Postures and Movements. *Sensors* **2021**, *21*, 5527. [[CrossRef](#)]
53. Forsman, M.; Fan, X.; Rhen, I.-M.; Lind, C.M. Mind the gap—Development of conversion models between accelerometer- and IMU-based measurements of arm and trunk postures and movements in warehouse work. *Appl. Ergon.* **2022**, *105*, 103841. [[CrossRef](#)]
54. Juul-Kristensen, B.; Hansson, G.A.; Fallentin, N.; Andersen, J.H.; Ekdahl, C. Assessment of work postures and movements using a video-based observation method and direct technical measurements. *Appl. Ergon.* **2001**, *32*, 517–524. [[CrossRef](#)] [[PubMed](#)]
55. Byström, J.U.; Hansson, G.A.; Rylander, L.; Ohlsson, K.; Källrot, G.; Skerfving, S. Physical workload on neck and upper limb using two CAD applications. *Appl. Ergon.* **2002**, *33*, 63–74. [[CrossRef](#)] [[PubMed](#)]
56. Christmansson, M.; Medbo, L.; Hansson, G.Å.; Ohlsson, K.; Unge Byström, J.; Möller, T.; Forsman, M. A case study of a principally new way of materials kitting—An evaluation of time consumption and physical workload. *Int. J. Ind. Ergon.* **2002**, *30*, 49–65. [[CrossRef](#)]
57. Arvidsson, I.; Arvidsson, M.; Axmon, A.; Hansson, G.Å.; Johansson, C.R.; Skerfving, S. Musculoskeletal disorders among female and male air traffic controllers performing identical and demanding computer work. *Ergonomics* **2006**, *49*, 1052–1067. [[CrossRef](#)] [[PubMed](#)]
58. Freitag, S.; Ellegast, R.; Dulon, M.; Nienhaus, A. Quantitative measurement of stressful trunk postures in nursing professions. *Ann. Occup. Hyg.* **2007**, *51*, 385–395. [[CrossRef](#)] [[PubMed](#)]
59. Hermanns, I.; Raffler, N.; Ellegast, R.P.; Fischer, S.; Göres, B. Simultaneous field measuring method of vibration and body posture for assessment of seated occupational driving tasks. *Int. J. Ind. Ergon.* **2008**, *38*, 255–263. [[CrossRef](#)]
60. Raffler, N.; Hermanns, I.; Sayn, D.; Göres, B.; Ellegast, R.; Rissler, J. Assessing combined exposures of whole-body vibration and awkward posture—further results from application of a simultaneous field measurement methodology. *Ind. Health* **2010**, *48*, 638–644. [[CrossRef](#)] [[PubMed](#)]
61. Ribeiro, D.C.; Sole, G.; Abbott, J.H.; Milosavljevic, S. Cumulative postural exposure measured by a novel device: A preliminary study. *Ergonomics* **2011**, *54*, 858–865. [[CrossRef](#)]
62. Fethke, N.B.; Gant, L.C.; Gerr, F. Comparison of biomechanical loading during use of conventional stud welding equipment and an alternate system. *Appl. Ergon.* **2011**, *42*, 725–734. [[CrossRef](#)]
63. Freitag, S.; Fincke-Junod, I.; Seddouki, R.; Dulon, M.; Hermanns, I.; Kersten, J.F.; Larsson, T.J.; Nienhaus, A. Frequent bending—An underestimated burden in nursing professions. *Ann. Occup. Hyg.* **2012**, *56*, 697–707. [[CrossRef](#)]
64. Jonker, D.; Rolander, B.; Balogh, I.; Sandsjö, L.; Ekberg, K.; Winkel, J. Rationalisation in public dental care—Impact on clinical work tasks and mechanical exposure for dentists—A prospective study. *Ergonomics* **2013**, *56*, 303–313. [[CrossRef](#)]
65. Ciccarelli, M.; Straker, L.; Mathiassen, S.E.; Pollock, C. Posture variation among office workers when using different information and communication technologies at work and away from work. *Ergonomics* **2014**, *57*, 1678–1686. [[CrossRef](#)] [[PubMed](#)]
66. Freitag, S.; Seddouki, R.; Dulon, M.; Kersten, J.F.; Larsson, T.J.; Nienhaus, A. The effect of working position on trunk posture and exertion for routine nursing tasks: An experimental study. *Ann. Occup. Hyg.* **2014**, *58*, 317–325. [[CrossRef](#)]
67. Wahlström, J.; Bergsten, E.; Trask, C.; Mathiassen, S.E.; Jackson, J.; Forsman, M. Full-Shift Trunk and Upper Arm Postures and Movements Among Aircraft Baggage Handlers. *Ann. Occup. Hyg.* **2016**, *60*, 977–990. [[CrossRef](#)]

68. Raffler, N.; Ellegast, R.; Kraus, T.; Ochsmann, E. Factors affecting the perception of whole-body vibration of occupational drivers: An analysis of posture and manual materials handling and musculoskeletal disorders. *Ergonomics* **2016**, *59*, 48–60. [[CrossRef](#)] [[PubMed](#)]
69. Raffler, N.; Rissler, J.; Ellegast, R.; Schikowsky, C.; Kraus, T.; Ochsmann, E. Combined exposures of whole-body vibration and awkward posture: A cross sectional investigation among occupational drivers by means of simultaneous field measurements. *Ergonomics* **2017**, *60*, 1564–1575. [[CrossRef](#)]
70. Yu, D.; Dural, C.; Morrow, M.M.; Yang, L.; Collins, J.W.; Hallbeck, S.; Kjellman, M.; Forsman, M. Intraoperative workload in robotic surgery assessed by wearable motion tracking sensors and questionnaires. *Surg. Endosc.* **2017**, *31*, 877–886. [[CrossRef](#)] [[PubMed](#)]
71. Vignais, N.; Bernard, F.; Touvenot, G.; Sagot, J.-C. Physical risk factors identification based on body sensor network combined to videotaping. *Appl. Ergon.* **2017**, *65*, 410–417. [[CrossRef](#)]
72. Arias, O.E.; Umukoro, P.E.; Stoffel, S.D.; Hopcia, K.; Sorensen, G.; Dennerlein, J.T. Associations between trunk flexion and physical activity of patient care workers for a single shift: A pilot study. *Work* **2017**, *56*, 247–255. [[CrossRef](#)] [[PubMed](#)]
73. Zare, M.; Biau, S.; Brunet, R.; Roquelaure, Y. Comparison of three methods for evaluation of work postures in a truck assembly plant. *Ergonomics* **2017**, *60*, 1551–1563. [[CrossRef](#)]
74. Heiden, M.; Zetterberg, C.; Mathiassen, S.E. Trunk and upper arm postures in paper mill work. *Appl. Ergon.* **2019**, *76*, 90–96. [[CrossRef](#)]
75. Jorgensen, M.B.; Gupta, N.; Korshoj, M.; Lagersted-Olsen, J.; Villumsen, M.; Mortensen, O.S.; Skotte, J.; Sogaard, K.; Madeleine, P.; Samani, A.; et al. The DPhacto cohort: An overview of technically measured physical activity at work and leisure in blue-collar sectors for practitioners and researchers. *Appl. Ergon.* **2019**, *77*, 29–39. [[CrossRef](#)] [[PubMed](#)]
76. Thamsuwan, O.; Galvin, K.; Tchong-French, M.; Kim, J.H.; Johnson, P.W. A feasibility study comparing objective and subjective field-based physical exposure measurements during apple harvesting with ladders and mobile platforms. *J. Agromedicine* **2019**, *24*, 268–278. [[CrossRef](#)] [[PubMed](#)]
77. Thamsuwan, O.; Galvin, K.; Tchong-French, M.; Aulck, L.; Boyle, L.N.; Ching, R.P.; McQuade, K.J.; Johnson, P.W. Comparisons of physical exposure between workers harvesting apples on mobile orchard platforms and ladders, part 1: Back and upper arm postures. *Appl. Ergon.* **2020**, *89*, 103193. [[CrossRef](#)]
78. Zare, M.; Bodin, J.; Sagot, J.C.; Roquelaure, Y. Quantification of Exposure to Risk Postures in Truck Assembly Operators: Neck, Back, Arms and Wrists. *Int. J. Environ. Res. Public Health* **2020**, *17*, 6062. [[CrossRef](#)] [[PubMed](#)]
79. Holtermann, A.; Fjeldstad Hendriksen, P.; Greby Schmidt, K.; Jagd Svendsen, M.; Nørregaard Rasmussen, C.D. Physical Work Demands of Childcare Workers in Denmark: Device-Based Measurements and Workplace Observations Among 199 Childcare Workers from 16 Day Nurseries. *Ann. Work Expo. Health* **2020**, *64*, 586–595. [[CrossRef](#)]
80. Robert-Lachaine, X.; Larue, C.; Denis, D.; Delisle, A.; Mecheri, H.; Corbeil, P.; Plamondon, A. Feasibility of quantifying the physical exposure of materials handlers in the workplace with magnetic and inertial measurement units. *Ergonomics* **2020**, *63*, 283–292. [[CrossRef](#)]
81. Khan, M.I.; Bath, B.; Kociolek, A.; Zeng, X.; Koehncke, N.; Trask, C. Trunk Posture Exposure Patterns among Prairie Ranch and Grain Farmers. *J. Agromedicine* **2020**, *25*, 210–220. [[CrossRef](#)] [[PubMed](#)]
82. Porta, M.; Pau, M.; Orrù, P.F.; Nussbaum, M.A. Trunk Flexion Monitoring among Warehouse Workers Using a Single Inertial Sensor and the Influence of Different Sampling Durations. *Int. J. Environ. Res. Public Health* **2020**, *17*, 7117. [[CrossRef](#)]
83. Wilhelmsson, S.; Andersson, M.; Arvidsson, I.; Dahlgvist, C.; Hemsworth, P.H.; Yngvesson, J.; Hultgren, J. Physical workload and psychosocial working conditions in Swedish pig transport drivers. *Int. J. Ind. Ergon.* **2021**, *83*, 103124. [[CrossRef](#)]
84. Schall, M.C., Jr.; Zhang, X.; Chen, H.; Gallagher, S.; Fethke, N.B. Comparing upper arm and trunk kinematics between manufacturing workers performing predominantly cyclic and non-cyclic work tasks. *Appl. Ergon.* **2021**, *93*, 103356. [[CrossRef](#)]
85. Brents, C.; Hischke, M.; Reiser, R.; Rosecrance, J. Trunk Posture during Manual Materials Handling of Beer Kegs. *Int. J. Environ. Res. Public Health* **2021**, *18*, 7380. [[CrossRef](#)] [[PubMed](#)]
86. Kim, J.H.; Vaughan, A.; Kincl, L. Characterization of Musculoskeletal Injury Risk in Dungeness Crab Fishing. *J. Agromedicine* **2022**, *28*, 309–323. [[CrossRef](#)]
87. Nunes, M.L.; Folgado, D.; Fujão, C.; Silva, L.; Rodrigues, J.; Matias, P.; Barandas, M.; Carreiro, A.V.; Madeira, S.; Gamboa, H. Posture Risk Assessment in an Automotive Assembly Line Using Inertial Sensors. *IEEE Access* **2022**, *10*, 83221–83235. [[CrossRef](#)]
88. Rolander, B.; Forsman, M.; Ghafouri, B.; Abtahi, F.; Wåhlin, C. Measurements and observations of movements at work for warehouse forklift truck operators. *Int. J. Occup. Saf. Ergon.* **2022**, *28*, 1840–1848. [[CrossRef](#)]
89. Fan, X.; Forsman, M.; Yang, L.; Lind, C.M.; Kjellman, M. Surgeons' physical workload in open surgery versus robot-assisted surgery and nonsurgical tasks. *Surg. Endosc.* **2022**, *36*, 8178–8194. [[CrossRef](#)] [[PubMed](#)]
90. Hess, J.A.; Hecker, S.; Weinstein, M.; Lunger, M. A participatory ergonomics intervention to reduce risk factors for low-back disorders in concrete laborers. *Appl. Ergon.* **2004**, *35*, 427–441. [[CrossRef](#)]

91. Kazmierczak, K.; Mathiassen, S.E.; Forsman, M.; Winkel, J. An integrated analysis of ergonomics and time consumption in Swedish “craft-type” car disassembly. *Appl. Ergon.* **2005**, *36*, 263–273. [[CrossRef](#)] [[PubMed](#)]
92. Nowak, J.; Erbe, C.; Hauck, I.; Groneberg, D.A.; Hermanns, I.; Ellegast, R.; Ditchen, D.; Ohlendorf, D. Motion analysis in the field of dentistry: A kinematic comparison of dentists and orthodontists. *BMJ Open* **2016**, *6*, e011559. [[CrossRef](#)]
93. Schall, M.C., Jr.; Fethke, N.B.; Chen, H. Working postures and physical activity among registered nurses. *Appl. Ergon.* **2016**, *54*, 243–250. [[CrossRef](#)] [[PubMed](#)]
94. Kozak, A.; Freitag, S.; Nienhaus, A. Evaluation of a Training Program to Reduce Stressful Trunk Postures in the Nursing Professions: A Pilot Study. *Ann. Work Expo. Health* **2017**, *61*, 22–32. [[CrossRef](#)]
95. Granzow, R.F.; Schall, M.C., Jr.; Smidt, M.F.; Chen, H.; Fethke, N.B.; Huangfu, R. Characterizing exposure to physical risk factors among reforestation hand planters in the Southeastern United States. *Appl. Ergon.* **2018**, *66*, 1–8. [[CrossRef](#)] [[PubMed](#)]
96. Hauck, I.; Erbe, C.; Nowak, J.; Hermanns, I.; Ditchen, D.; Ellegast, R.; Oremek, G.; Groneberg, D.A.; Ohlendorf, D. Kinematic posture analysis of orthodontists in their daily working practice. *J. Orofac. Orthop.* **2018**, *79*, 389–402. [[CrossRef](#)]
97. Merino, G.; da Silva, L.; Mattos, D.; Guimarães, B.; Merino, E. Ergonomic evaluation of the musculoskeletal risks in a banana harvesting activity through qualitative and quantitative measures, with emphasis on motion capture (Xsens) and EMG. *Int. J. Ind. Ergon.* **2019**, *69*, 80–89. [[CrossRef](#)]
98. Merkus, S.L.; Lunde, L.K.; Koch, M.; Waersted, M.; Knardahl, S.; Veiersted, K.B. Physical capacity, occupational physical demands, and relative physical strain of older employees in construction and healthcare. *Int. Arch. Occup. Environ. Health* **2019**, *92*, 295–307. [[CrossRef](#)]
99. Fethke, N.B.; Schall, M.C., Jr.; Chen, H.; Branch, C.A.; Merlino, L.A. Biomechanical factors during common agricultural activities: Results of on-farm exposure assessments using direct measurement methods. *J. Occup. Environ. Hyg.* **2020**, *17*, 85–96. [[CrossRef](#)] [[PubMed](#)]
100. Bijanzadeh, A.; Hermanns, I.; Ellegast, R.; Fraeulin, L.; Holzgreve, F.; Mache, S.; Groneberg, D.A.; Ohlendorf, D. A kinematic posture analysis of neurological assistants in their daily working practice—a pilot study. *J. Occup. Med. Toxicol.* **2020**, *15*, 36. [[CrossRef](#)] [[PubMed](#)]
101. Loske, D.; Klumpp, M.; Keil, M.; Neukirchen, T. Logistics Work, Ergonomics and Social Sustainability: Empirical Musculoskeletal System Strain Assessment in Retail Intralogistics. *Logistics* **2021**, *5*, 89. [[CrossRef](#)]
102. Zhao, J.; Obonyo, E.; Bilén, S.G. Wearable Inertial Measurement Unit Sensing System for Musculoskeletal Disorders Prevention in Construction. *Sensors* **2021**, *21*, 1324. [[CrossRef](#)]
103. Ettinger, L.; Kincl, L.; Johnson, P.; Carter, C.; Garfinkel, S.; Karduna, A. Workday Arm Elevation Exposure: A Comparison Between Two Professions. *IIEE Trans. Occup. Ergon. Hum. Factors* **2013**, *1*, 119–127. [[CrossRef](#)]
104. Hanvold, T.N.; Wærsted, M.; Mengshoel, A.M.; Bjertness, E.; Veiersted, K.B. Work with prolonged arm elevation as a risk factor for shoulder pain: A longitudinal study among young adults. *Appl. Ergon.* **2015**, *47*, 43–51. [[CrossRef](#)] [[PubMed](#)]
105. Jackson, J.A.; Mathiassen, S.E.; Wahlström, J.; Liv, P.; Forsman, M. Is what you see what you get? Standard inclinometry of set upper arm elevation angles. *Appl. Ergon.* **2015**, *47*, 242–252. [[CrossRef](#)] [[PubMed](#)]
106. Dahlqvist, C.; Nordander, C.; Forsman, M.; Enquist, H. Self-recordings of upper arm elevation during cleaning—Comparison between analyses using a simplified reference posture and a standard reference posture. *BMC Musculoskelet. Disord.* **2018**, *19*, 402. [[CrossRef](#)] [[PubMed](#)]
107. Hansson, G.Å.; Balogh, I.; Unge Byström, J.; Ohlsson, K.; Nordander, C.; Asterland, P.; Sjölander, S.; Rylander, L.; Winkel, J.; Skerfving, S. Questionnaire versus direct technical measurements in assessing postures and movements of the head, upper back, arms and hands. *Scand. J. Work Environ. Health* **2001**, *27*, 30–40. [[CrossRef](#)] [[PubMed](#)]
108. Balogh, I.; Ohlsson, K.; Hansson, G.Å.; Engström, T.; Skerfving, S. Increasing the degree of automation in a production system: Consequences for the physical workload. *Int. J. Ind. Ergon.* **2006**, *36*, 353–365. [[CrossRef](#)]
109. Jonker, D.; Rolander, B.; Balogh, I. Relation between perceived and measured workload obtained by long-term inclinometry among dentists. *Appl. Ergon.* **2009**, *40*, 309–315. [[CrossRef](#)] [[PubMed](#)]
110. Jonker, D.; Rolander, B.; Balogh, I.; Sandsjö, L.; Ekberg, K.; Winkel, J. Mechanical exposure among general practice dentists in Sweden and possible implications of rationalisation. *Ergonomics* **2011**, *54*, 953–960. [[CrossRef](#)] [[PubMed](#)]
111. Heilskov-Hansen, T.; Svendsen, S.W.; Thomsen, J.F.; Mikkelsen, S.; Hansson, G.Å. Sex differences in task distribution and task exposures among Danish house painters: An observational study combining questionnaire data with biomechanical measurements. *PLoS ONE* **2014**, *9*, e110899. [[CrossRef](#)] [[PubMed](#)]
112. Simonsen, J.G.; Dahlqvist, C.; Enquist, H.; Nordander, C.; Axmon, A.; Arvidsson, I. Assessments of Physical Workload in Sonography Tasks Using Inclinometry, Goniometry, and Electromyography. *Saf. Health Work* **2018**, *9*, 326–333. [[CrossRef](#)]
113. Unge, J.; Ohlsson, K.; Nordander, C.; Hansson, G.Å.; Skerfving, S.; Balogh, I. Differences in physical workload, psychosocial factors and musculoskeletal disorders between two groups of female hospital cleaners with two diverse organizational models. *Int. Arch. Occup. Environ. Health* **2007**, *81*, 209–220. [[CrossRef](#)]

114. Nordander, C.; Ohlsson, K.; Balogh, I.; Hansson, G.Å.; Axmon, A.; Persson, R.; Skerfving, S. Gender differences in workers with identical repetitive industrial tasks: Exposure and musculoskeletal disorders. *Int. Arch. Occup. Environ. Health* **2008**, *81*, 939–947. [[CrossRef](#)] [[PubMed](#)]
115. Wahlström, J.; Mathiassen, S.E.; Liv, P.; Hedlund, P.; Ahlgren, C.; Forsman, M. Upper arm postures and movements in female hairdressers across four full working days. *Ann. Occup. Hyg.* **2010**, *54*, 584–594. [[CrossRef](#)] [[PubMed](#)]
116. Åkesson, I.; Balogh, I.; Hansson, G.Å. Physical workload in neck, shoulders and wrists/hands in dental hygienists during a work-day. *Appl. Ergon.* **2012**, *43*, 803–811. [[CrossRef](#)] [[PubMed](#)]
117. Balogh, I.; Ohlsson, K.; Nordander, C.; Bjork, J.; Hansson, G.A. The importance of work organization on workload and musculoskeletal health—Grocery store work as a model. *Appl. Ergon.* **2016**, *53 Pt A*, 143–151. [[CrossRef](#)]
118. Wærsted, M.; Enquist, H.; Veiersted, K.B. Hairdressers' shoulder load when blow-drying—Studying the effect of a new blow dryer design on arm inclination angle and muscle pain. *Int. J. Ind. Ergon.* **2019**, *74*, 102839. [[CrossRef](#)]
119. Nourollahi-Darabad, M.; Afshari, D.; Dianat, I.; Jodakinia, L. Long-duration assessment of upper arm posture and motion and their association with perceived symptoms among bakery workers. *Int. J. Ind. Ergon.* **2020**, *80*, 103029. [[CrossRef](#)]
120. Palm, P.; Gupta, N.; Forsman, M.; Skotte, J.; Nordquist, T.; Holtermann, A. Exposure to Upper Arm Elevation During Work Compared to Leisure Among 12 Different Occupations Measured with Triaxial Accelerometers. *Ann. Work Expo. Health* **2018**, *62*, 689–698. [[CrossRef](#)] [[PubMed](#)]
121. Bruno Garza, J.L.; Eijkelhof, B.H.W.; Johnson, P.W.; Raina, S.M.; Rynell, P.W.; Huysmans, M.A.; van Dieën, J.H.; van der Beek, A.J.; Blatter, B.M.; Dennerlein, J.T. Observed differences in upper extremity forces, muscle efforts, postures, velocities and accelerations across computer activities in a field study of office workers. *Ergonomics* **2012**, *55*, 670–681. [[CrossRef](#)] [[PubMed](#)]
122. Thamsuwan, O.; Galvin, K.; Tchong-French, M.; Aulck, L.; Boyle, L.N.; Ching, R.P.; McQuade, K.J.; Johnson, P.W. Comparisons of physical exposure between workers harvesting apples on mobile orchard platforms and ladders, part 2: Repetitive upper arm motions. *Appl. Ergon.* **2020**, *89*, 103192. [[CrossRef](#)]
123. Moriguchi, C.S.; Carnaz, L.; Alencar, J.F.; Miranda Júnior, L.C.; Granqvist, L.; Hansson, G.; Gil Coury, H.J. Postures and movements in the most common tasks of power line workers. *Ind. Health* **2011**, *49*, 482–491. [[CrossRef](#)] [[PubMed](#)]
124. Moriguchi, C.S.; Carnaz, L.; Miranda Júnior, L.C.; Marklin, R.W.; Gil Coury, H.J. Are posture data from simulated tasks representative of field conditions? Case study for overhead electric utility workers. *Ergonomics* **2012**, *55*, 1382–1394. [[CrossRef](#)]
125. Moriguchi, C.S.; Carnaz, L.; Veiersted, K.B.; Hanvold, T.N.; Hæg, L.B.; Hansson, G.; Cote Gil Coury, H.J. Occupational posture exposure among construction electricians. *Appl. Ergon.* **2013**, *44*, 86–92. [[CrossRef](#)]
126. Arvidsson, I.; Hansson, G.Å.; Mathiassen, S.E.; Skerfving, S. Changes in physical workload with implementation of mouse-based information technology in air traffic control. *Int. J. Ind. Ergon.* **2006**, *36*, 613–622. [[CrossRef](#)]
127. Buchholz, B.; Park, J.S.; Gold, J.; Punnett, L. Subjective ratings of upper extremity exposures: Inter-method agreement with direct measurement of exposures. *Ergonomics* **2008**, *51*, 1064–1077. [[CrossRef](#)] [[PubMed](#)]
128. Hess, J.A.; Kincl, L.; Amasay, T.; Wolfe, P. Ergonomic evaluation of masons laying concrete masonry units and autoclaved aerated concrete. *Appl. Ergon.* **2010**, *41*, 477–483. [[CrossRef](#)] [[PubMed](#)]
129. Arvidsson, I.; Balogh, I.; Hansson, G.; Ohlsson, K.; Åkesson, I.; Nordander, C. Rationalization in meat cutting—Consequences on physical workload. *Appl. Ergon.* **2012**, *43*, 1026–1032. [[CrossRef](#)]
130. Douphrate, D.I.; Fethke, N.B.; Nonnenmann, M.W.; Rosecrance, J.C.; Reynolds, S.J. Full shift arm inclinometry among dairy parlor workers: A feasibility study in a challenging work environment. *Appl. Ergon.* **2012**, *43*, 604–613. [[CrossRef](#)]
131. Kersten, J.T.; Fethke, N.B. Radio frequency identification to measure the duration of machine-paced assembly tasks: Agreement with self-reported task duration and application in variance components analyses of upper arm postures and movements recorded over multiple days. *Appl. Ergon.* **2019**, *75*, 74–82. [[CrossRef](#)] [[PubMed](#)]
132. Veiersted, K.B.; Gould, K.S.; Osteras, N.; Hansson, G.A. Effect of an intervention addressing working technique on the biomechanical load of the neck and shoulders among hairdressers. *Appl. Ergon.* **2008**, *39*, 183–190. [[CrossRef](#)]
133. Mathiassen, S.E.; Paquet, V. The ability of limited exposure sampling to detect effects of interventions that reduce the occurrence of pronounced trunk inclination. *Appl. Ergon.* **2010**, *41*, 295–304. [[CrossRef](#)]
134. Dahlqvist, C.; Hansson, G.Å.; Forsman, M. Validity of a small low-cost triaxial accelerometer with integrated logger for uncomplicated measurements of postures and movements of head, upper back and upper arms. *Appl. Ergon.* **2016**, *55*, 108–116. [[CrossRef](#)]
135. Jackson, J.A.; Mathiassen, S.E.; Wahlström, J.; Liv, P.; Forsman, M. Digging deeper into the assessment of upper arm elevation angles using standard inclinometry. *Appl. Ergon.* **2015**, *51*, 102–103. [[CrossRef](#)]
136. Peters, A.; Galna, B.; Sangeux, M.; Morris, M.; Baker, R. Quantification of soft tissue artifact in lower limb human motion analysis: A systematic review. *Gait Posture* **2010**, *31*, 1–8. [[CrossRef](#)] [[PubMed](#)]
137. Hansson, G. Letter to the editor. Re: Is what you see what you get? Standard inclinometry of set upper arm elevation angles. *Appl. Ergon.* **2015**, *48*, 109–110. [[CrossRef](#)] [[PubMed](#)]

138. Hansson, G.A.; Arvidsson, I.; Ohlsson, K.; Nordander, C.; Mathiassen, S.E.; Skerfving, S.; Balogh, I. Precision of measurements of physical workload during standardised manual handling. Part II: Inclinometry of head, upper back, neck and upper arms. *J. Electromyogr. Kinesiol.* **2006**, *16*, 125–136. [[CrossRef](#)]
139. Lind, C.M.; Forsman, M. Accuracy of a posture measurement system for practitioners. In Proceedings of the 47th Nordic Ergonomics and Human Factors Society Conference, Lillehammer, Norway, 1–4 November 2015.
140. Lind, C.M.; Fan, X.; Rhen, I.M.; Forsman, M. *Does Posture Matter—Accuracy of Standard Calibration Procedures for Arm Inclinometry [Manuscript in Preparation]*; Institute of Environmental Medicine, Karolinska Institutet: Stockholm, Sweden, 2024.
141. Forsman, M.; Fan, X.; Rhen, I.M.; Lind, C.M. Concerning a Work Movement Velocity Action Level Proposed in “Action Levels for the Prevention of Work-Related Musculoskeletal Disorders in the Neck and Upper Extremities: A Proposal” by Inger Arvidsson et al. (2021). *Ann. Work Expo. Health* **2022**, *66*, 130–131. [[CrossRef](#)] [[PubMed](#)]
142. Hanson, L.; Sperling, L.; Gard, G.; Ipsen, S.; Olivares Vergara, C. Swedish anthropometrics for product and workplace design. *Appl. Ergon.* **2009**, *40*, 797–806. [[CrossRef](#)] [[PubMed](#)]
143. Fryar, C.D.; Carroll, M.D.; Gu, Q.; Afful, J.; Ogden, C.L. Anthropometric Reference Data for Children and Adults: United States, 2015–2018. In *Vital & Health Statistics. Series 3, Analytical and Epidemiological Studies*; Centers for Disease Control and Prevention National Center for Health Statistics: Hyattsville, MD, USA, 2021; pp. 1–44.
144. Hoareau, D.; Fan, X.; Abtahi, F.; Yang, L. Evaluation of In-Cloth versus On-Skin Sensors for Measuring Trunk and Upper Arm Postures and Movements. *Sensors* **2023**, *23*, 3969. [[CrossRef](#)] [[PubMed](#)]
145. Stock, S.R.; Fernandes, R.; Delisle, A.; Vézina, N. Reproducibility and validity of workers’ self-reports of physical work demands. *Scand. J. Work Environ. Health* **2005**, *31*, 409–437. [[CrossRef](#)]
146. Koch, M.; Lunde, L.K.; Gjulem, T.; Knardahl, S.; Veiersted, K.B. Validity of Questionnaire and Representativeness of Objective Methods for Measurements of Mechanical Exposures in Construction and Health Care Work. *PLoS ONE* **2016**, *11*, e0162881. [[CrossRef](#)]
147. Barriera-Viruet, H.; Sobeih, T.M.; Daraiseh, N.; Salem, S. Questionnaires vs observational and direct measurements: A systematic review. *Theor. Issues Ergon. Sci.* **2006**, *7*, 261–284. [[CrossRef](#)]
148. Nyman, T.; Rhen, I.M.; Johansson, P.J.; Eliasson, K.; Kjellberg, K.; Lindberg, P.; Fan, X.; Forsman, M. Reliability and Validity of Six Selected Observational Methods for Risk Assessment of Hand Intensive and Repetitive Work. *Int. J. Environ. Res. Public Health* **2023**, *20*, 5505. [[CrossRef](#)] [[PubMed](#)]

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.