

# **Cosmic Ray Irradiation of Interstellar Ices on Sulfur-Rich Grains: A Possible Source of Sulfur-Bearing Molecules**

Downloaded from: https://research.chalmers.se, 2025-06-08 01:09 UTC

Citation for the original published paper (version of record):

Mifsud, D., Kaňuchová, Z., Auriacombe, O. et al (2025). Cosmic Ray Irradiation of Interstellar Ices on Sulfur-Rich Grains: A Possible Source of Sulfur-Bearing Molecules. ACS Earth and Space Chemistry, 9(5): 1227-1242. http://dx.doi.org/10.1021/acsearthspacechem.5c00036

N.B. When citing this work, cite the original published paper.

research.chalmers.se offers the possibility of retrieving research publications produced at Chalmers University of Technology. It covers all kind of research output: articles, dissertations, conference papers, reports etc. since 2004. research.chalmers.se is administrated and maintained by Chalmers Library



Article

# Cosmic Ray Irradiation of Interstellar Ices on Sulfur-Rich Grains: A Possible Source of Sulfur-Bearing Molecules

Published as part of ACS Earth and Space Chemistry special issue "Harold Linnartz Festschrift".

Duncan V. Mifsud,\* Zuzana Kaňuchová, Olivier Auriacombe, Péter Herczku, Danna Qasim, Sándor T. S. Kovács, Richárd Rácz, Béla Sulik, Zoltán Juhász, István Rajta, István Vajda, Sándor Biri, Robert W. McCullough, Sergio Ioppolo, Ujjwal Raut, and Nigel J. Mason\*



**ABSTRACT:** The major reservoir of sulfur in dense interstellar clouds is still largely unknown, although a growing body of evidence suggests that it may exist in a refractory form (i.e., as minerals or allotropes of the element). Therefore, it is possible that the irradiation of sulfur-free interstellar ices on top of sulfur-rich refractory grain components by cosmic rays or stellar winds may result in the formation of simple inorganic sulfur molecules that could be readily detected by ground- or space-borne telescopes. In this study, we have irradiated neat ices of O<sub>2</sub>, CO, CO<sub>2</sub>, and H<sub>2</sub>O on top of layers of allotropic sulfur at 20 K using 1 MeV He<sup>+</sup> ions as a mimic of space radiation. Experiments with CO<sub>2</sub> and H<sub>2</sub>O ices were also repeated at 70 K to provide data obtained under conditions more relevant to icy bodies in the outer solar system for comparative purposes. We have found qualitative mid-infrared spectroscopic evidence for the synthesis of SO<sub>2</sub>, CS<sub>2</sub>, OCS,



and  $H_2SO_4$  hydrates, but not  $H_2S$ , in our experiments and have quantified the efficiency of their formation by calculating the *G*-value (i.e., the number of molecules formed per 100 eV of energy deposited) for each ice-refractory system. Overall,  $SO_2$  and  $CS_2$  are the most commonly observed products in our experiments, although the highest *G*-value was that for  $H_2SO_4$  hydrates formed as a result of the irradiation of  $H_2O$  ice on top of sulfur at 70 K. An important outcome of our study is that our experimental results are consistent with recent observational surveys that suggest  $SO_2$  formation in interstellar ices proceeds primarily *via* an "energetic" route involving radiolytic processes, while OCS forms as a result of "nonenergetic" processes such as atom or radical addition reactions. KEYWORDS: *astrochemistry, radiation chemistry, sulfur, interstellar ices, laboratory experiments, infrared spectroscopy* 

#### 1. INTRODUCTION

Downloaded via 213.89.182.114 on May 19, 2025 at 07:41:18 (UTC). See https://pubs.acs.org/sharingguidelines for options on how to legitimately share published articles.

The astrochemistry of sulfur has long posed problems to the scientific community.<sup>1</sup> In interstellar media, for instance, it is known that the observable abundance of sulfur in diffuse regions largely aligns with its expected cosmic abundance, but that it is depleted by about two orders of magnitude within dense molecular clouds.<sup>2-7</sup> The principal drivers and mechanisms behind this so-called sulfur depletion problem are not fully understood, though a number of previous experimental and computational studies have suggested various reservoirs that may accommodate this missing sulfur. For instance, modeling efforts by Laas and Caselli<sup>8</sup> suggested that organosulfur molecules embedded within interstellar icy grain mantles could account for a large percentage of sulfur. Other studies have proposed gas-phase atoms as the dominant sulfur species, which could be converted to H<sub>2</sub>S by surface-catalyzed reactions on dust grains or ice surfaces.<sup>9,10</sup> However, the as-yet nondetection of solid H<sub>2</sub>S within dense molecular clouds<sup>11,12</sup> seems to suggest that other potential interstellar sulfur

reservoirs should be sought. More recently, sulfur-bearing salts such as NH<sub>4</sub>SH have been suggested to be a potential sink for interstellar sulfur.<sup>13,14</sup>

Ruffle et al.<sup>6</sup> proposed that a potential major sink of gasphase sulfur during the evolution of diffuse interstellar clouds into denser structures is the Coulomb-enhanced freeze-out of  $S^+$  ions onto negatively charged dust grains. Although the subsequent chemical reactions in which grain-adsorbed sulfur atoms are still somewhat speculative, it is conceivable that atom addition reactions may proceed to yield various allotropes. Indeed, evidence for the reaction of sulfur atoms to yield  $S_2$  and thence  $S_{2n}$  allotropes can be found in previous

Received:February 3, 2025Revised:April 23, 2025Accepted:April 29, 2025Published:May 7, 2025



experimental studies.<sup>15–18</sup> Comprehensive studies by Shingledecker et al.<sup>19</sup> that incorporated both cosmic ray-induced chemistry and fast, nondiffusive radical combination reactions of the type previously described by Garrod et al.<sup>20</sup> have also demonstrated that sulfur chemistry in interstellar icy grain mantles could produce large quantities of S<sub>8</sub>. These theoretical results have been corroborated by experimental studies that have shown that the processing of astrophysical ice analogs containing simple inorganic sulfur-bearing molecules by ions, electrons, or ultraviolet photons results in the formation of refractory sulfur-rich residues, which likely include large allotropes.<sup>21–27</sup>

Although the observation of sulfur allotropes on interstellar dust grains is very challenging due to their low absorbances across the mid-infrared range,<sup>28</sup> their possible existence in such environs is perhaps reinforced by their detection in asteroid 162173 Ryugu and comet 67P/Churyumov-Gerasimenko.<sup>29,30</sup> An alternative possible sink for the depletion of interstellar sulfur is its inclusion in mineral phases, primarily as metal sulfides. Such minerals are thought to be among the earliest synthesized in the history of mineralogical evolution and, indeed, so-called GEMSs (silicate glass with embedded metal and sulfides) have been detected in interstellar dust grains found preserved in chondritic meteorites.<sup>31</sup> Moreover, observations of (post)asymptotic giant branch stars have revealed a high abundance of sulfide minerals, particularly those of iron and magnesium.<sup>32,33</sup> Work by Kama et al.<sup>34</sup> has also suggested that such minerals could account for a large majority (>80%) of the sulfur in protoplanetary disks around young stars.

Either way, the results of these recent studies suggest that a considerable fraction of the missing interstellar sulfur could be locked away as refractory solid components of dust grains (i.e., allotropes or minerals). Such an implication is important, as this sulfur may interact chemically with an overlying ice mantle to produce new molecules. This is especially true when galactic cosmic rays interact with interstellar icy grain mantles, since such an injection of energy is known to lead to the formation of a plethora of new species, some of which may be complex organic molecules that are relevant to the emergence of life.<sup>35–37</sup> Previous laboratory work has demonstrated that the irradiation of interstellar ice analogs on top of carbon-rich grain analogs leads to the formation of a number of carbonbearing inorganic molecules, most prominently CO and  $CO_2$ . For instance, Mennella et al.<sup>38</sup> demonstrated that the irradiation of H<sub>2</sub>O ice on top of hydrogenated carbon grains simulating interstellar dust particles using 30 keV He<sup>+</sup> ions resulted in the synthesis of both CO and CO<sub>2</sub>. This result was echoed by Raut et al.,<sup>39</sup> who observed the formation of CO<sub>2</sub> after the 100 keV proton irradiation of H<sub>2</sub>O ice on amorphous carbon. Gomis and Strazzulla<sup>40</sup> went on to demonstrate that the synthesis of CO and CO<sub>2</sub> in this manner does not depend on the chemical nature of the underlying residue, so long as it is carbon-rich. Finally, the studies of Mennella et al.<sup>41</sup> and Fulvio et al.42 showed that this chemistry could also be induced by ultraviolet photons, as well as ions.

Given that sulfur may be an important component of refractory interstellar grains, either as allotropes or minerals, and in light of the results of previous analogous studies making use of refractory carbon,  $^{38-42}$  it occurred to us that the interaction of galactic cosmic rays with icy material on top of sulfur-rich dust grains may lead to the synthesis of new molecules at the ice-refractory interface, which may be

detectable by ground- or space-borne observatories and telescopes. This idea is reinforced by the recent work of Ferrari et al.,<sup>43</sup> which revealed that the fragmentation of S<sub>8</sub> under simulated astrophysical conditions is facile, thus further hinting at a potentially important role for allotropic sulfur in interstellar sulfur chemistry. At present, only two sulfur-bearing molecules have been either definitively or tentatively detected within interstellar icy grain mantles: OCS and SO<sub>2</sub>.<sup>11,12</sup> Indeed, it is possible that this mechanism of refractory sulfur volatilization is an important contributor to the presence of sulfur-bearing molecules may represent the starting point for a relatively rich subsequent organic chemistry leading to ever more complex sulfur molecules.

In this study, we probed the formation of sulfur-bearing molecules of interest to interstellar chemistry as a result of the cosmic ray irradiation of sulfur-free ices on top of sulfur-rich dust grains. Specifically, we have irradiated neat ices of O<sub>2</sub>, CO, CO<sub>2</sub>, and H<sub>2</sub>O on top of allotropic sulfur at 20 K using 1 MeV He<sup>+</sup> ions as mimics of galactic cosmic rays. In the case of CO<sub>2</sub> and H<sub>2</sub>O ices deposited on top of sulfur, experiments were also performed at 70 K to allow for a comparison of the radiationinduced sulfur chemistry occurring under interstellar conditions to that occurring on the surfaces of higher-temperature outer solar system objects. Sulfur-bearing radiolytic products have been identified through mid-infrared absorption spectroscopy, and their formation efficiencies have been quantified through the so-called G-value, which is a measurement of the number of molecules formed per 100 eV of energy dosed into the irradiated target. We note that several related but somewhat distinct definitions exist for the G-value; however, we have followed the example of a number of previous studies that define it by considering the total energy dosed into both the adsorbate and the adsorbent materials, 49-53 which, in our case, correspond to the icy material and the underlying refractory sulfur. Finally, our results are discussed in the context of their applicability to sulfur astrochemistry and the detection of various sulfur molecules in interstellar and, to a lesser extent, outer solar system environments.

### 2. METHODOLOGY

Experiments were carried out using the Ice Chamber for Astrophysics-Astrochemistry (ICA), a laboratory setup for radiation astrochemistry studies located at the HUN-REN Institute for Nuclear Research (Atomki) in Debrecen, Hungary. This setup has been described in detail in previous publications,<sup>54,55</sup> and so, only the most salient features will be described here. The ICA (Figure 1) is an ultrahigh-vacuum stainless-steel chamber with an operational base pressure of a few  $10^{-9}$  mbar. Within the center of the chamber is a goldcoated, oxygen-free, high-conductivity copper sample holder into which up to four infrared-transparent ZnSe substrates may be mounted. The sample holder and deposition substrates may be cooled to 20 K by virtue of being held in thermal contact with the coldfinger of a closed-cycle helium cryostat; however, the temperature of the sample holder and the substrates may be regulated within the 20-300 K range by establishing an equilibrium between the cooling effect of the cryostat and the warming induced by an internal cartridge heater.

Sulfur-rich interstellar grain analogs were prepared by first cooling the sample holder and substrates down to 20 K (or 70 K if simulating outer solar system objects) and then rotating them to face the nozzle of a commercial effusive evaporator NCO

Gas Inlet

**Figure 1.** Simplified top-view schematic diagram of the ICA setup. Sulfur layers were prepared onto the cooled ZnSe substrates by rotating them to directly face the nozzle of an effusive evaporator, after which they were rotated back to face an incident mid-infrared spectroscopic beam. Ices were prepared on top of the sulfurous layers through the background condensation of dosed gases and vapors. Irradiation with 1 MeV He<sup>+</sup> ions was performed using the arrangement shown here, with ions impacting the targets at an angle of 36° to the surface normal. All mid-infrared spectra were acquired in transmission mode by using an external mercury–cadmium–telluride (MCT) detector.

(Createc OLED-40-10-WK-SHM) that had been fitted to the ICA and which had been preloaded with a few grams of pharmaceutical-grade sulfur. The exact allotropic composition of this sulfur is not known, but is likely to be dominated by cyclic S8 molecules, due to the known stability of this structure.<sup>56</sup> The sulfur was subsequently warmed to 78  $\pm$  3 °C and allowed to equilibrate at this temperature for a few minutes, before opening the shutter in front of the evaporator's nozzle to allow for the deposition of the sulfur on the ZnSe substrates. Sulfur layers were deposited to a thickness of approximately 4  $\mu$ m (determined through interferometric methods), after which the shutter was closed and the sample holder rotated to face an incident mid-infrared spectroscopic beam. At this point, a background spectrum was acquired to be used as a reference when collecting subsequent mid-infrared absorption spectra.

Interstellar ice analogs composed of neat O<sub>2</sub>, CO, CO<sub>2</sub>, or H<sub>2</sub>O were prepared on top of the sulfur layers by dosing gases or vapors from liquids into the main chamber though a fine regulating needle valve, which allowed for the background deposition of the ices. In the case of the  $O_2$ , CO, and  $CO_2$ experiments, gases from lecture bottles were used as supplied without any further purification  $(O_2 \text{ and } CO_2 \text{ purity} =$ 99.995%; CO purity = 99.97%; all supplied by Linde), whereas in the case of the H<sub>2</sub>O experiments, vapor was extracted from a deionized water sample that had been degassed through several iterations of the freeze-pump-thaw cycle. The deposition of the interstellar ice analogs could be followed in situ using Fourier-Transform mid-infrared transmission absorption spectroscopy over the 4000-650 cm<sup>-1</sup> range, which allowed for a quantitative assessment of the amount of icy material that was deposited. The column density N (molecules  $cm^{-2}$ ) of a deposited ice is related to the integrated absorbance  $S(cm^{-1})$ of one of its characteristic mid-infrared absorption bands as

$$N = \ln(10)\frac{S}{A_{\nu}} \tag{1}$$

where  $A_{\nu}$  is the so-called integrated strength constant (cm molecule<sup>-1</sup>) of that band. The constant factor ln(10) is included in eq 1 to allow for the relation of *S*, which is

measured on an absorbance scale, to  $A_{\nu}$ , which is measured on an optical depth scale. The column density of a deposited ice is related to its thickness h ( $\mu$ m) as

$$h = 10^4 \frac{NM}{\rho N_{\rm A}} \tag{2}$$

where *M* is the molar mass of the ice (g mol<sup>-1</sup>),  $\rho$  is the ice density (g cm<sup>-3</sup>), and  $N_A$  is the Avogadro constant (6.02 × 10<sup>23</sup> molecules mol<sup>-1</sup>). The constant 10<sup>4</sup> is included in eq 2 to express *h* in units of  $\mu$ m.

A list of the integrated band strength constants and densities used to quantify the molecular column densities and thicknesses of the astrophysical ice analogs considered in this study is given in Table 1. Of course, eqs 1 and 2 cannot be

Table 1. Absorption Band Strength Constants  $(A_{\nu})$  and Densities  $(\rho)$  of the Interstellar Ice Analogs and Sulfur-Rich Grain Analogs Considered in This Study

	absorption band	$A_{ u}$	ρ
molecule	cm <sup>-1</sup>	$10^{-17}$ cm molecule <sup>-1<i>a</i></sup>	g cm <sup>-3b</sup>
S <sub>8</sub>			2.07
O <sub>2</sub>			1.54
CO	2139 ( $\nu_{\rm s}$ )	1.1	0.88
CO <sub>2</sub>	2343 ( $\nu_3$ )	7.6	0.98 at 20 K, 1.48 at 70 K
$H_2O$	$\sim$ 3250 ( $\nu_{\rm s}$ )	20.0	0.93

<sup>*a*</sup>Integrated band strength constants taken from the work of Gerakines et al. <sup>62</sup> <sup>*b*</sup>Densities of S<sub>8</sub>, O<sub>2</sub>, CO, CO<sub>2</sub>, and H<sub>2</sub>O respectively taken from the works of Greenwood and Earnshaw, <sup>63</sup> Freiman and Jodl, <sup>64</sup> Luna et al., <sup>65</sup> Satorre et al., <sup>66</sup> and Narten et al. <sup>67</sup>

used to quantify the column density and thickness of the  $O_2$ ice, since this species is infrared-inactive due to it being a homonuclear diatomic molecule. Instead, we assumed that the deposition rate of  $O_2$  is similar to that of CO based on the similarities in the geometries, molecular masses, and chamber pumping speeds of these molecules; and so, depositing an  $O_2$ ice at the same chamber pressure and for the same duration as was done for the CO ice would result in a similar deposited column density, from which the ice thickness could be determined based on the known density of solid  $O_2$  (Table 1).

Once deposited to a satisfactory thickness, the interstellar and outer solar system ice analogs prepared on top of sulfurrich grain analogs were exposed to a 1 MeV He<sup>+</sup> ion beam supplied by a 2 MV Tandetron accelerator, 57,58 with incident ions impacting the ices at an angle of 36° to the surface normal. The choice to use 1 MeV He<sup>+</sup> ions in this study was motivated by two factors: first, such ions are a known component of galactic cosmic rays<sup>59,60</sup> and are thus relevant to astrochemistry. Second, calculations performed with the Stopping and Ranging of Ions in Matter (SRIM) software<sup>61</sup> revealed that in every case, the penetration range of these ions was less than the combined thickness of the ice and sulfur-rich grain analog ( $R_{max} < 3.4 \ \mu m$ ), and thus, the neutralized ions came to rest in the sulfur layer. By making use of helium ions, any chemical modifications of the sulfur layer as a result of its reaction with the implanted particle could be precluded due to the inert nature of the neutral helium atom. Overall, ices were irradiated to a total fluence of  $10^{15}$  ions cm<sup>-2</sup>, with several midinfrared absorption spectra being acquired at preselected fluence intervals. Spectra were acquired as an average 128

coadded scans and at a resolution of  $1 \text{ cm}^{-1}$ . A summary of the experiments performed is provided in Table 2.

Table 2. Summary of the Experiments Performed in This Study, Including Information on the Column Densities (N) and Thicknesses (h) of the Interstellar Ice Analogs Deposited on Top of Sulfur, as well as the Maximum Penetration Range  $(R_{max})$  of 1 MeV He<sup>+</sup> Ions into the Combined Thickness of the Ice and Sulfur-Rich Grain Analog<sup>a</sup>

		N	h	R <sub>max</sub>
experiment	interstellar ice analog	$10^{17}$ molecules cm <sup>-2</sup>	μm	μm
1	O <sub>2</sub> at 20 K	5.79	0.20	2.9
2	CO at 20 K	3.78	0.20	3.0
3	CO <sub>2</sub> at 20 K	0.94	0.07	3.0
4	CO <sub>2</sub> at 70 K	1.01	0.05	2.9
5	H <sub>2</sub> O at 20 K	37.32	1.20	3.3
6	H <sub>2</sub> O at 70 K	31.10	1.00	3.1

"Note that the temperature at which a particular ice-refractory system was irradiated was the same as the temperature at which it was prepared

## 3. RESULTS AND DISCUSSION

**3.1.**  $O_2$  lce on Top of Sulfur. The irradiation of a neat  $O_2$  interstellar ice analog on top of a sulfur-rich layer at 20 K resulted in the formation of SO<sub>2</sub>, as was determined by the appearance of two mid-infrared absorption features at about 1335 and 1150 cm<sup>-1</sup> (Figure 2). The bands have been



**Figure 2.** Mid-infrared absorption spectra acquired during the irradiation of a neat  $O_2$  ice  $(h = 0.2 \ \mu m)$  on top of a layer of allotropic sulfur  $(h \approx 4 \ \mu m)$  at 20 K using 1 MeV He<sup>+</sup> ions. A spectrum of pristine, unirradiated (1:6) SO<sub>2</sub>:O<sub>2</sub> mixed ice is also included for comparison. Spectra are vertically offset from one another in the interest of clarity. Oscillations in the spectra at lower wavenumbers are likely the result of changes in the ice thickness due to ion-induced sputtering of volatile material.

respectively attributed to the asymmetric and symmetric stretching modes of  $SO_2$  on the basis of the excellent agreement of their wavenumber positions with those of a reference  $SO_2:O_2$  (1:6) ice mixture, as well as their exhibiting the expected  $SO_2$  spectral profile in which the intensity of the asymmetric stretching mode is greater than that of the symmetric stretching mode.<sup>68</sup> By making use of an integrated



**Figure 3.** Column density of SO<sub>2</sub> measured during the irradiation of neat O<sub>2</sub> ice ( $h = 0.2 \ \mu m$ ) on top of a layer of allotropic sulfur ( $h \approx 4 \ \mu m$ ) at 20 K using 1 MeV He<sup>+</sup> ions. The red curve represents an exponential fit to the data ( $r^2 > 0.999$ ), while the blue line represents a linear trend ( $r^2 > 0.999$ ) fitted to the initial points of the exponential curve, from which a *G*-value of 0.0457 ± 0.0004 molecules per 100 eV was calculated.

fluence of  $4.2 \times 10^{13}$  ions cm<sup>-2</sup> had been delivered, and its measured column density could be fitted by an exponential growth function that plateaued at a value of about  $1.2 \times 10^{16}$  molecules cm<sup>-2</sup>. By fitting a linear trend line to the initial points of this exponential growth curve (Figure 3), where the only process likely contributing to the abundance of SO<sub>2</sub> in the ice is its own radiolytic formation, we could compute a *G*-value of 0.0457  $\pm$  0.0004 molecules per 100 eV.

A number of reactions may contribute to the synthesis of SO<sub>2</sub> in this experiment and, although it is not possible to confirm with any certainty the dominance of one reaction relative to another, it is possible to speculate on the most likely processes contributing to SO<sub>2</sub> formation. For instance, it is possible that the reaction of a sulfur allotrope  $S_n$  with electronically excited O2 could lead to the formation of a cyclic intermediate structure in which one sulfur atom is covalently bonded to the two oxygen atoms, as discussed previously by Mayer.<sup>70</sup> This sulfur atom then splits off from the intermediate to yield  $SO_2$  and  $S_{n-1}$ . Although low-temperature solid-phase studies on the addition of atomic or molecular oxygen to allotropic sulfur are generally lacking, previous atmospheric chemistry studies concerned with analogous gasphase reactions have determined that such reactions are kinetically favorable,<sup>71</sup> and thus, it is conceivable that they would also proceed under our experimental conditions. Matrix isolation spectroscopic chemiluminescence studies by Long and Pimentel<sup>72</sup> also demonstrated that electronically excited sulfur atoms may undergo an insertion-type reaction with molecular oxygen to directly yield SO<sub>2</sub>. This reaction is believed to proceed with a near-zero activation energy barrier at temperatures between 10 and 40 K.<sup>72,73</sup>

Oxygen atoms that are formed as a result of the radiolytic dissociation of  $O_2$  away from the ice-refractory interface may undergo atom addition reactions with  $O_2$  to yield  $O_3$ . Indeed, both the asymmetric stretching and combination  $(\nu_1 + \nu_3)$  modes of  $O_3$  were detected in acquired mid-infrared absorption spectra of the irradiated ice at 1040 and 2110 cm<sup>-1,74</sup> respectively (Figure 2). The chemistry of  $O_3$  formation in irradiated ices that are initially rich in  $O_2$  has been well documented by previous studies,<sup>75–79</sup> and so will not be discussed in greater detail here.

**3.2. CO Ice on Top of Sulfur.** The irradiation of a CO interstellar ice analog on top of a sulfur-rich layer resulted in the formation of several new products, as evidenced by the appearance of many new absorption features in the acquired mid-infrared absorption spectra (Figures 4 and 5). Many of



**Figure 4.** Mid-infrared absorption spectra acquired during the irradiation of neat CO ice  $(h = 0.2 \ \mu m)$  on top of a layer of allotropic sulfur  $(h \approx 4 \ \mu m)$  at 20 K using 1 MeV He<sup>+</sup> ions, illustrating the radiolytic synthesis of various oxocarbon molecules in the ice.



**Figure 5.** Mid-infrared absorption spectra acquired during the irradiation of a neat CO ice  $(h = 0.2 \ \mu\text{m})$  on top of a layer of allotropic sulfur  $(h \approx 4 \ \mu\text{m})$  at 20 K using 1 MeV He<sup>+</sup> ions. Spectra are vertically offset from one another in the interest of clarity. Oscillations in the spectra at lower wavenumbers are likely the result of changes in the ice thickness due to ion-induced sputtering of volatile material.

these new absorption features were clustered in the 2400–2000 cm<sup>-1</sup> wavenumber range and are associated with the C= O bond stretching modes of various oxocarbon molecules (Table 3).<sup>80-82</sup> The radiolytic synthesis of these oxocarbon

Table 3. Oxocarbon Molecules Detected after the Irradiation of CO Ice on Top of a Sulfur-Rich Layer at 20  $K^a$ 

na di abuti a	mana durat	malagula
radioivite	DFOULTEL	morecine

formula	name	mid-infrared band position $(cm^{-1})$
<sup>12</sup> CO <sub>2</sub>	carbon dioxide	2346
<sup>13</sup> CO <sub>2</sub>	carbon dioxide	2281
C <sub>3</sub> O	tricarbon monoxide	2247
$C_3O_2$	tricarbon dioxide (carbon suboxide)	2242
$C_5O_2$	pentacarbon dioxide	2211
$C_3O_2$	tricarbon dioxide (carbon suboxide)	2193
<sup>12</sup> CO	carbon monoxide	2138
$C_7O_2$	heptacarbon dioxide	2122
C <sub>5</sub> O	pentacarbon monoxide	2106
<sup>13</sup> CO	carbon monoxide	2092
C <sup>18</sup> O	carbon monoxide	2088
<sup>a</sup> Band as	signments were made on the ba	sis of those made by previou

"Band assignments were made on the basis of those made by previous studies."  $^{80-82}$ 

species in irradiated CO-rich interstellar ice analogs has been well documented by previous studies and is believed to proceed through carbon atom addition reactions. In this scheme, electronically excited CO first reacts with ground-state CO to yield CO<sub>2</sub> and atomic carbon, which may then add to various  $C_nO$  species in the ice to yield  $C_{n+1}O$  products.<sup>83,84</sup> It is also possible for two oxocarbon molecules  $C_nO$  and  $C_mO$  to combine to yield a cumulene dioxide  $C_{m+n}O_2$ .<sup>83,84</sup> It is interesting to note that our results suggest that oxygen atoms are not yielded as part of the mechanistic chemistry occurring during the irradiation of CO ice on top of allotropic sulfur. This is evidenced by the lack of  $O_3$  or any oxygenated sulfur molecule such as  $SO_2$  in this experiment, whose synthesis would require the radiolytic liberation and subsequent reaction of oxygen atoms.<sup>71,75–79,85</sup>

The only detected sulfur-bearing product was CS2, which was identified via the appearance of its  $\nu_3$  mode at 1509 cm<sup>-1</sup> (Figure 5).<sup>86</sup> It should be noted that the exact wavenumber position of this band has been reported to vary somewhat depending on the composition of the molecular ice: for instance, Sivaraman<sup>87</sup> measured its position to be  $1511 \text{ cm}^{-1}$  in a mixed  $CS_2$ :CO<sub>2</sub> ice deposited at  $\overline{85}$  K, while Ferrante et al.<sup>46</sup> measured it to be 1524 cm<sup>-1</sup> for CS<sub>2</sub> produced as a result of the irradiation of a H<sub>2</sub>S:CO mixed ice using 800 keV protons at 10 K. The chemical reactions responsible for the synthesis of CS<sub>2</sub> are somewhat challenging to determine, since literature studies on the formation of this molecule in astrophysical ice analogs are relatively sparse. Ferrante et al.46 proposed that CS<sub>2</sub> could result from the stepwise reaction of CO with two sulfur atoms, which would sequentially yield OCS and OCS<sub>2</sub>. This latter molecule is unstable and thus would dissociate to  $CS_2$  and atomic oxygen.

However, this reaction mechanism is at odds with the nondetection of the OCS in our experiment (Figure 5). Moreover, the reaction between atomic sulfur and OCS could

also conceivably yield CS and S<sub>2</sub>, as discussed by Ikeda et al.,<sup>88</sup> thereby reducing the formation efficiency of CS<sub>2</sub>. The presence of mobile oxygen atoms within the ice matrix should also have resulted in the synthesis of highly oxygenated products, such as O3, which were not detected. Instead, we speculate that an alternative pathway toward CS2 may rely on the abundance of free carbon atoms within the irradiated CO ice: these atoms could abstract a sulfur atom from an allotrope of sulfur  $S_n$  to yield CS which, being unstable, could then abstract a second sulfur atom to yield the CS2 product. To the best of our knowledge, the reaction between carbon atoms and sulfurcontaining molecules has not yet been experimentally investigated in the solid phase; however, previous gas-phase work by Deeyamulla and Husain<sup>89</sup> showed that carbon atoms may abstract sulfur atoms from various simple inorganic sulfur molecules.

By making use of an integrated band strength constant of 9.1  $\times 10^{-17}$  cm molecule<sup>-1</sup> for the asymmetric stretching mode of CS<sub>2</sub>,<sup>87</sup> its column density could be quantified as a function of fluence. As can be seen in Figure 5, CS<sub>2</sub> is unambiguously detectable in acquired mid-infrared spectra at fluences greater than  $10^{14}$  ions cm<sup>-2</sup>, and its column density was found to grow in a linear fashion with increasing ion fluence (Figure 6). The



**Figure 6.** Column density of CS<sub>2</sub> measured during the irradiation of neat CO ice ( $h = 0.2 \,\mu$ m) on top of a layer of allotropic sulfur ( $h \approx 4 \,\mu$ m) at 20 K using 1 MeV He<sup>+</sup> ions. The blue line represents a linear trend fitted to the data ( $r^2 > 0.94$ ) from which a *G*-value of (6.66 ± 0.50) × 10<sup>-4</sup> molecules per 100 eV was calculated.

overall abundance of CS<sub>2</sub> yielded is low when compared to the abundance of SO<sub>2</sub> produced during the analogous irradiation of O<sub>2</sub> ice on allotropic sulfur (Figure 3). As such, it is likely that the data in Figure 6 represent the earliest fluence points at which CS<sub>2</sub> synthesis occurs within the irradiated ice, and at which any other chemical processes that could impact the abundance of CS<sub>2</sub> within the ice are negligible. Therefore, a linear trend line could be fitted to the entire data set so as to calculate a *G*-value of  $(6.66 \pm 0.50) \times 10^{-4}$  molecules per 100 eV.

**3.3.**  $CO_2$  lce on Top of Sulfur. The irradiation of  $CO_2$  ice on top of allotropic sulfur was performed at two temperatures

(i.e., 20 and 70 K) to allow for a comparison of the possible chemistry occurring within dense molecular clouds in the interstellar medium to that on the surfaces of cold outer solar system objects. At 20 K, the irradiation resulted in the formation of a number of new product molecules (Figure 7),



**Figure 7.** Mid-infrared absorption spectra acquired during the irradiation of neat CO<sub>2</sub> ice  $(h = 0.07 \ \mu\text{m})$  on top of a layer of allotropic sulfur  $(h \approx 4 \ \mu\text{m})$  at 20 K using 1 MeV He<sup>+</sup> ions. Spectra are vertically offset from one another in the interest of clarity.

including those known to be yielded as a result of the radiolytic processing of solid CO<sub>2</sub>, such as CO, CO<sub>3</sub>, and O<sub>3</sub>.<sup>90–93</sup> Moreover, the absorption features of both SO<sub>2</sub> and CS<sub>2</sub> were detected in mid-infrared spectra acquired during irradiation. The radiation chemistry leading to the formation of all of these molecules is very rich, and multiple reaction pathways may contribute to the formation of common molecules. The irradiation of CO<sub>2</sub> results in the homolytic dissociation of one C=O bond, yielding CO and atomic oxygen as radiolytic products.<sup>94</sup> The addition of this oxygen atom to a parent CO<sub>2</sub> molecule may yield the CO<sub>3</sub> product, while the combination of free oxygen atoms may sequentially yield O<sub>2</sub> and O<sub>3</sub>.<sup>90</sup>

We propose that there are at least three reaction pathways through which SO<sub>2</sub> can be produced. Two of these routes are identical to those proposed to account for the formation of  $SO_2$  as a result of the irradiation of  $O_2$  ice on top of allotropic sulfur, namely, the insertion-type reaction of electronically excited sulfur atoms into O2 molecules generated via the combination of radiolytically derived oxygen atoms, and the addition of  $O_2$  to allotropic sulfur  $S_n$  to yield a cyclic intermediate that decomposes to yield SO<sub>2</sub> and  $S_{n-1}$ . The third possible route involves the reaction of CO<sub>2</sub> with a sulfur atom to yield CO and SO,<sup>95</sup> prior to the oxidation of the latter species to yield SO<sub>2</sub>.<sup>71,85</sup> On the other hand, the most likely reaction sequence leading to the formation of CS<sub>2</sub> is analogous to that proposed by Hawkins et al.<sup>96</sup> and Ferrante et al.<sup>46</sup> in which CO sequentially abstracts two atoms of sulfur from an allotropic structure to yield OCS and OCS<sub>2</sub>, after which this latter species dissociates to CS<sub>2</sub> and atomic oxygen.<sup>9</sup> However, as was demonstrated by Ferrante et al.,<sup>46</sup> OCS is not particularly stable when processed by ionizing radiation and therefore may not accumulate within the ice, hence explaining its nondetection in the experiment period. K. Aside from the reaction with sulfur to yield  $OCS_{22}^{46,96} OCS$ further contributing to the production of SO<sub>2</sub> via the rapid



**Figure 8.** Column density of SO<sub>2</sub> (left panel) and CS<sub>2</sub> (right panel) measured during the irradiation of neat CO<sub>2</sub> ice ( $h = 0.07 \ \mu$ m) on top of a layer of allotropic sulfur ( $h \approx 4 \ \mu$ m) at 20 K using 1 MeV He<sup>+</sup> ions. Note than an exponential curve could not be fitted to the SO<sub>2</sub> data, and thus, the blue line represents a linear trend fitted to the first four data points ( $r^2 > 0.90$ ) from which a *G*-value of 0.0078 ± 0.0014 molecules per 100 eV was calculated. Conversely, an exponential curve could be fitted to the CS<sub>2</sub> data (red curve;  $r^2 > 0.95$ ), to which a linear trend (blue line;  $r^2 > 0.99$ ) could be fitted to its initial points. From this linear trend, a *G*-value of (5.89 ± 0.13) × 10<sup>-4</sup> molecules per 100 eV was calculated.

oxidation of the SO intermediate.<sup>71,85</sup> It should also be noted that  $OCS_2$  need not necessarily dissociate to  $CS_2$  and atomic oxygen, but may also dissociate to CO and  $S_2$ .<sup>101</sup> Although  $S_2$  is a volatile sulfur allotrope,<sup>102</sup> and thus represents one outcome of the volatilization of the deposited sulfur layer, its formation could not be confirmed through mid-infrared spectroscopy due to it being a homonuclear diatomic molecule and thus infrared-inactive.

The column densities of both SO<sub>2</sub> and CS<sub>2</sub> were measured as a function of fluence throughout the 20 K experiment (Figure 8). Interestingly, the column density of SO<sub>2</sub> steadily increased until a fluence of  $6 \times 10^{14}$  ions cm<sup>-2</sup> had been delivered, after which point it began to decrease. Such a trend was also observed for the non-sulfur-bearing radiolytic products, whose absorption bands all began to decrease in intensity at higher fluences (Figure 7). CS<sub>2</sub> was first observed in the irradiated ice after a fluence of about  $10^{14}$  ions cm<sup>-2</sup> had been delivered and, in contrast to the trends observed for the other products of this irradiation, its column density steadily increased before plateauing at about  $2.7 \times 10^{15}$  molecules cm<sup>-2</sup> by the end of the experiment (Figure 8). G-values for the  $SO_2$ and  $CS_2$  products were calculated to be 0.0078 ± 0.0014 and  $(5.89 \pm 0.13) \times 10^{-4}$  molecules per 100 eV, respectively. The reason for the apparent decline in the column densities of all radiolytic products other than CS<sub>2</sub> at higher fluences is difficult to explain; however, a likely contributor to this observation was ice sputtering, which has been described in detail for CO2 ices bombarded by ions at low temperatures (i.e., <25 K).<sup>103</sup> It is also possible to speculate that the greater relative abundances of CS<sub>2</sub> within the ice at higher fluences allowed for the consumption of volatile ice components in the formation of carbon-sulfur polymers that could not be detected using midinfrared absorption spectroscopy. Indeed, prior research has documented the efficient formation of such complex polymeric structures after the irradiation of CS<sub>2</sub> and that these polymers exhibit very weak or difficult-to-observe absorptions over midinfrared wavelengths.<sup>104–106</sup>

The spectroscopic results of the 1 MeV He<sup>+</sup> ion irradiation of  $CO_2$  ice on top of sulfur at 20 K (Figure 7) contrast strongly with those of the analogous experiment performed at 70 K (Figure 9). Perhaps the most evident difference is that at this higher temperature, the irradiation of the ice results in higher



**Figure 9.** Mid-infrared absorption spectra acquired during the irradiation of neat CO<sub>2</sub> ice ( $h = 0.05 \ \mu\text{m}$ ) on top of a layer of allotropic sulfur ( $h \approx 4 \ \mu\text{m}$ ) at 70 K using 1 MeV He<sup>+</sup> ions. Spectra are vertically offset from one another in the interest of clarity.

abundances of the sulfur-bearing products SO2 and CS2, whose column densities were found to increase with higher fluences (Figure 10). Moreover, OCS was detected through the appearance of a broad absorption band centered at 2031  $cm^{-1}$  in acquired mid-infrared spectra (Figure 9), attributable to its  $\nu_3$  mode. The position of the peak of this broad band is known to shift by up to 30 cm<sup>-1</sup> based upon the composition of the ice;<sup>23,46,87</sup> however, our observation is in good agreement with that reported by Sivaraman,<sup>87</sup> who observed it at 2035 cm<sup>-1</sup> after the irradiation of a CO<sub>2</sub>:CS<sub>2</sub> mixed ice at 85 K, and is in reasonable agreement with that reported by Ferrante et al.,<sup>46</sup> who observed it at 2019 cm<sup>-1</sup> for a neat OCS ice prepared at 10 K and subsequently warmed to 25 K. It is to be further noted that the position of this OCS absorption feature is in close proximity to the  $\nu_1$  mode of CO<sub>3</sub>; however, these two bands may be distinguished from each other on the basis of their full-widths at half-maxima,<sup>46</sup> since the CO<sub>3</sub>  $\nu_1$ mode is known to present as a relatively narrow band. Indeed, in our experiment, the OCS absorption band at 2031 cm<sup>-1</sup> had a significantly larger full-width at half-maximum of 11 cm<sup>-1</sup>,



**Figure 10.** Column density of SO<sub>2</sub> (left panel), CS<sub>2</sub> (center panel), and OCS (right panel) measured during the irradiation of neat CO<sub>2</sub> ice ( $h = 0.05 \ \mu m$ ) on top of a layer of allotropic sulfur ( $h \approx 4 \ \mu m$ ) at 70 K using 1 MeV He<sup>+</sup> ions. Note that a *G*-value of 0.0104 ± 0.0003 molecules per 100 eV was calculated for SO<sub>2</sub> by fitting a linear trend line (blue line;  $r^2 > 0.99$ ) to the initial seven data points. A *G*-value of (9.99 ± 0.19) × 10<sup>-4</sup> molecules per 100 eV was calculated for CS<sub>2</sub> by fitting a linear trend line (blue line;  $r^2 > 0.997$ ) to the entire data set. Lastly, an exponential curve could be fitted to the OCS data (red curve;  $r^2 > 0.92$ ), to which a linear trend line (blue line;  $r^2 > 0.999$ ) could be fitted to its initial points. From this linear trend, a *G*-value of (0.89 ± 0.01) × 10<sup>-4</sup> molecules per 100 eV was calculated.

compared to the more modest 4 cm<sup>-1</sup> for the CO<sub>3</sub> band at 2045 cm<sup>-1</sup> (Figure 9).

Interestingly, although the irradiation of CO<sub>2</sub> ice on top of sulfur at 70 K led to higher abundances of sulfur-bearing products (i.e., SO<sub>2</sub>, CS<sub>2</sub>, and OCS) compared to the analogous experiment at 20 K, abundances of nonsulfur-bearing products (i.e., CO, CO<sub>3</sub>, and O<sub>3</sub>) were generally lower (Figures 7 and 9). Given that the initial abundances of both  $CO_2$  ice and allotropic sulfur were similar across both experiments (Table 2), the difference in these results must stem from the different reaction pathways being favored at these two temperatures. Since radicals are known to be more mobile within solid matrices at higher temperatures, it is likely that large sulfur atoms released from the underlying layer of allotropic sulfur as a result of its irradiation can migrate further into the overlying ice layer at 70 K than they could have done at 20 K. Indeed, the increased mobility of sulfur atoms in an ice matrix at higher temperatures has already been demonstrated in previous studies.<sup>107,108</sup> Therefore, the comparatively higher concentration of sulfur atoms intermixed with CO<sub>2</sub> ice could conceivably promote the reaction leading to SO and CO, which then produces  $SO_2$  upon oxidation of the SO.<sup>71,85</sup> Some tentative evidence for the enhancement of this reaction lies in the appearance of the SO<sub>2</sub> symmetric stretching mode which, unlike in the analogous 20 K experiment, appears to be split into a double-peaked structure with peaks at 1046 and 1036  $cm^{-1}$  (Figure 9). We speculate that this double-peaked structure is the result of contributions from the symmetric stretching modes of SO<sub>2</sub> (at 1046  $\text{cm}^{-1}$ ) and of SO (at 1036 cm<sup>-1</sup>).<sup>97,109</sup> Although SO is generally an unstable species that is typically rapidly oxidized to  $SO_{21}$  it is possible that the productivity of the sulfur chemistry occurring in our 70 K irradiation experiment allows for the sustained production of this radical and thus its tentative spectroscopic detection. Moreover, the higher concentration of sulfur atoms within the irradiated CO<sub>2</sub> ice would also compete more efficiently for oxygen atoms and CO molecules, which would have the net effect of reducing the abundance of these species and, by extension, of both O<sub>3</sub> and CO<sub>3</sub> within the irradiated ice.

The column densities of  $SO_2$ ,  $CS_2$ , and OCS were tracked throughout the 70 K irradiation experiment (Figure 10). In the case of  $SO_2$ , it was apparent that the column density initially increased linearly as a function of fluence; however, beginning

at a fluence of approximately  $9 \times 10^{13}$  ions cm<sup>-2</sup>, the gradient of this linear trend decreased. As such, the G-value, which was found to be 0.0104  $\pm$  0.0003 molecules per 100 eV, was calculated by fitting a trend line to the initial higher gradient linear data points at the earliest fluences. The evolution of the CS<sub>2</sub> column density was linear throughout the irradiation experiment (Figure 10) and, indeed, was qualitatively similar to the accumulation of CS2 observed during the irradiation of neat CO ice on top of sulfur at 20 K (Figure 6). The G-value for this product, which was calculated to be  $(9.99 \pm 0.19) \times$  $10^{-4}$  molecules per 100 eV, was therefore found by fitting a linear trend line to the entire data set. Finally, the column density of OCS was measured by making use of an integrated band strength constant of  $1.5 \times 10^{-16}$  cm molecule<sup>-1</sup> for the asymmetric stretching mode<sup>110</sup> and was found to follow a general exponential growth trend, albeit at low abundances (Figure 10). This is perhaps reflective of the comparative radiolytic instability of the OCS, which precludes it from accumulating within the ice during irradiation. By fitting a linear trend line to the initial points of this exponential growth curve, where radiolytic formation is likely the only process contributing to the abundance of OCS, a G-value of  $(0.89 \pm$  $(0.01) \times 10^{-4}$  molecules per 100 eV could be calculated.

3.4. H<sub>2</sub>O Ice on Top of Sulfur. The mid-infrared spectra acquired during the irradiation of H<sub>2</sub>O ice on top of allotropic sulfur at 20 K are shown in Figure 11. It is possible to note that irradiation does not result in many noticeable changes to the appearance of the spectra, other than the emergence of a weak but broad absorption band centered at 2863 cm<sup>-1</sup>, which has been attributed to the formation of  $H_2O_2$ .<sup>111–114</sup> Furthermore, two small absorption features at 3696 and 3721  $\text{cm}^{-1}$  that were apparent in the mid-infrared spectrum of the H<sub>2</sub>O ice upon its preparation at 20 K were observed to disappear very rapidly during irradiation. These bands were attributed to the OH dangling bonds that exist in microporous H<sub>2</sub>O ice at low temperatures, and their disappearance is evidence of the ice having undergone a process of radiation-induced compaction.<sup>115-117</sup> A particularly interesting result of this experiment is the fact that no sulfur-bearing products could be identified during irradiation. Comparisons with the results of previous studies are somewhat challenging, since most previous experimental work was performed with the aim of characterizing radiation-induced sulfur chemistry on the surfaces of the



**Figure 11.** Mid-infrared absorption spectra acquired during the irradiation of neat  $H_2O$  ice  $(h = 1.2 \ \mu m)$  on top of a layer of allotropic sulfur  $(h \approx 4 \ \mu m)$  at 20 K using 1 MeV He<sup>+</sup> ions. Spectra are vertically offset from one another in the interest of clarity.

icy Galilean moons of Jupiter and thus was carried out at higher temperatures.

Indeed, to the best of our knowledge, the only experiment performed at a comparable temperature to that of the present experiment is that of Kaňuchová et al.,<sup>118</sup> who investigated the 30 keV He<sup>+</sup> ion irradiation of a mixed H<sub>2</sub>O:SO<sub>2</sub> (1:2) ice at 16 K. Their work demonstrated that the irradiation of this ice resulted in the formation of H<sub>2</sub>SO<sub>4</sub> and its monohydrate and tetrahydrate forms, as inferred from mid-infrared absorption bands at 1105 and 1070 cm<sup>-1</sup> that were respectively attributed to the presence of  $HSO_4^{-}$  and  $SO_4^{2-}$  ions, as well as the broad  $H_3O^+$  band centered at about 1730 cm<sup>-1</sup>. This result contrasts strongly with that presented in Figure 11, where no such midinfrared absorption bands could be identified. The reason for this difference may, however, be fairly mundane: it is likely that the sulfur chemistry occurring in the experiment of Kaňuchová et al.<sup>118</sup> occurred to a greater extent due to the more homogeneous mixing of the SO<sub>2</sub> and H<sub>2</sub>O components of their ice, thus allowing for sulfur product formation to occur uniformly throughout the ice. This was not the case in the present experiment, where the radiolytic formation of sulfurbearing products could only occur at the ice-refractory interface and its immediate environs and was likely limited by the low mobilities of sulfur atoms at 20 K.<sup>107,108</sup>

The 1 MeV He<sup>+</sup> ion irradiation of  $H_2O$  ice on top of a layer of allotropic sulfur was also performed at 70 K, and, in this case, spectroscopic evidence for the radiation-induced formation of H<sub>2</sub>SO<sub>4</sub> and its monohydrate and tetrahydrate forms could be identified (Figure 12). In particular, absorption features at 1139, 1107, and 1060 cm<sup>-1</sup> were observed and could be respectively associated with the formation of  $H_2SO_4$ ,  $HSO_4^{-}$ , and  $SO_4^{2-}$  within the irradiated ice.<sup>118-120</sup> Although the microporous structure of  $H_2O$  ice, which has been previously described as a facilitator of radical and molecule mobility throughout the ice matrix, is known to vary with ice temperature,<sup>115-117,121,122</sup> we do not believe that this was the main driver behind the difference in the chemical productivities of our experiments conducted at 20 and 70 K. This is due to the fact that absorption bands associated with OH dangling bonds were initially detected in spectra of the H<sub>2</sub>O ices acquired before irradiation at both these temperatures, but quickly disappeared during irradiation due to ice compaction



**Figure 12.** Mid-infrared absorption spectra acquired during the irradiation of neat  $H_2O$  ice  $(h = 1.0 \ \mu\text{m})$  on top of a layer of allotropic sulfur  $(h \approx 4 \ \mu\text{m})$  at 70 K using 1 MeV He<sup>+</sup> ions. Spectra are vertically offset from one another in the interest of clarity.

and micropore destruction, even as the abundances of the acid hydrate products continued to increase. Moreover, bands associated with the OH dangling bonds were more intense in the 20 K ice compared with the 70 K ice, indicating a reduced porosity in the ice prepared at the higher temperature. Nevertheless, more extensive sulfur chemistry was observed in the H<sub>2</sub>O ice irradiated on top of sulfur at 70 K, thus indicating that the dominant driver of this chemistry is more likely to be the greater thermally driven mobility of radicals and molecules in the 70 K ice; particularly of large sulfur atoms released from the underlying layer of allotropic sulfur which would then migrate further into the ice structure and trigger more extensive chemistry at this higher temperature.<sup>107,108</sup> A similar argument was made to explain the enhanced formation of sulfur-bearing products as a result of the irradiation of CO<sub>2</sub> ice on top of sulfur at 70 K compared with 20 K.

The reactions leading to the formation of these acid hydrate products were described in detail by Della Guardia and Johnston,<sup>123</sup> who highlighted the important role played by OH radicals and H<sub>2</sub>SO<sub>2</sub>, a lower oxoacid of sulfur whose astrochemical importance was expounded upon more recently by Góbi et al.<sup>124</sup> In this reaction network (shown in Scheme 1), OH radicals produced from the radiolytic dissociation of H<sub>2</sub>O react with atomic sulfur to yield H<sub>2</sub>SO<sub>2</sub>, which then dissociates to  $H^+$  and  $HSO_2^-$ . The anion is subsequently neutralized via a charge exchange reaction with OH, before a subsequent reaction with OH oxidizes it to HSO<sub>3</sub><sup>-</sup>. Another round of neutralization via charge exchange followed by oxidation mediated by the reaction with OH radicals yields  $HSO_4^-$ . At this point,  $HSO_4^-$  may react in three different ways to yield H<sub>2</sub>SO<sub>4</sub>, the monohydrate acid, or the tetrahydrate acid. The H<sub>2</sub>SO<sub>4</sub> product is formed as a result of the simple combination reaction between  $H^+$  and  $HSO_4^-$ , while the monohydrate is formed via the combination reaction between  $H_3O^+$  and  $HSO_4^-$ . Finally, the tetrahydrate is produced in a two-step process in which  $HSO_4^-$  first dissociates to  $H^+$  and  $SO_4^{2-}$ , after which two  $H_3O^+$  ions and two neutral  $H_2O$  molecules associate with the  $SO_4^{2-}$  anion.

The integrated absorbances of the bands related to the acid hydrate products were not easy to measure in the mid-infrared spectra acquired during irradiation, and thus, we have made use of so-called difference spectra in order to facilitate these 11+

Scheme 1. Reaction Network Proposed by Della Guardia and Johnston<sup>123</sup> Leading to the Formation of H<sub>2</sub>SO<sub>4</sub> and Its Monohydrate and Tetrahydrate Forms after the Irradiation of H<sub>2</sub>O Ice on Top of Sulfur at 70 K Using 1 MeV He<sup>+</sup> Ions

$$H_{2}SO_{2} \xrightarrow{-H^{+}} HSO_{2}^{-} \xrightarrow{+OH} OH^{-} + HSO_{2}$$

$$\downarrow^{+} \stackrel{s}{}_{+OH} \qquad \qquad \downarrow^{+} OH$$

$$OH \qquad H^{+} + HSO_{3}^{-} \qquad \qquad \downarrow^{+} OH$$

$$H_{2}O \qquad OH^{-} + HSO_{3} \qquad \qquad \qquad \downarrow^{+} OH$$

$$H_{2}SO_{4} \xrightarrow{+H^{+}} HSO_{4}^{-} \xrightarrow{+H_{3}O^{+}} H_{2}SO_{4}.H_{2}O$$

$$\downarrow$$

$$SO_{4}^{2-} \qquad \qquad \downarrow^{+} 2H_{2}O_{4} + 2H_{3}O^{+}$$

$$H_{2}SO_{4}.4H_{2}O$$

measurements. Difference spectra were produced by subtracting the spectrum acquired at zero fluence from all spectra subsequently acquired during irradiation, as shown in the inset of Figure 12. Furthermore, in measuring the column densities of the acid hydrate products, we have followed the example of Ding et al.<sup>125</sup> and have integrated over all acid hydrate absorption bands across the 1350-760 cm<sup>-1</sup> wavenumber range, making use of a band strength constant of  $1.3 \times 10^{-17}$ cm molecule<sup>-1</sup>. The total abundance of the acid hydrate products increased in an exponential-like fashion (Figure 13), and by fitting a linear trend line to the initial points of the fitted curve, a G-value of  $0.4163 \pm 0.0031$  molecules per 100 eV could be calculated.



**Figure 13.** Total column density of acid hydrates (i.e.,  $H_2SO_4 \cdot nH_2O_3$ ; n = 0, 1, 4) measured during the irradiation of neat H<sub>2</sub>O ice (h = 1.0 $\mu$ m) on top of a layer of allotropic sulfur ( $h \approx 4 \mu$ m) at 70 K using 1 MeV He<sup>+</sup> ions. The red curve represents an exponential fit to the data  $(r^2 > 0.97)$ , while the blue line represents a linear trend  $(r^2 > 0.999)$ fitted to the initial points of the exponential curve, from which a Gvalue of  $0.4163 \pm 0.0031$  molecules per 100 eV was calculated.

#### 4. ASTROCHEMICAL IMPLICATIONS AND CONCLUSIONS

In this study, we investigated whether simple inorganic sulfurbearing molecules that could be detected by ground- or spaceborne telescopes can be formed in interstellar ices (and, to a lesser extent, outer solar system ices) as a result of the irradiation of sulfur-free icy material on top of sulfur-rich dust grains. Our experiments have demonstrated that several such molecules can be formed in this way, including SO<sub>2</sub>, CS<sub>2</sub>, OCS, and H<sub>2</sub>SO<sub>4</sub> hydrates. One particularly noteworthy result of our experiments performed with H<sub>2</sub>O ice was the nondetection of H<sub>2</sub>S among the radiolytic products. Indeed, the identification of this molecule in interstellar icy grain mantles has thus far also eluded observational astronomers. This is perhaps counterintuitive, as surface-catalyzed hydrogen atom addition reactions are known to be a feasible route toward the formation of simple molecules in dense molecular clouds,<sup>126</sup> and so the hydrogenation of grain-adsorbed sulfur atoms could in principle offer a route toward interstellar H<sub>2</sub>S. Although past theoretical studies have suggested that this hydrogenation reaction is accompanied by very high energy barriers when sulfur is adsorbed on a platinum surface,<sup>127</sup> it is important to note that, to the best of our knowledge, this reaction route has not been studied either theoretically or experimentally using surfaces that are more relevant to astrochemistry and so its possible contribution to interstellar H<sub>2</sub>S is unknown. Recent studies have demonstrated that H<sub>2</sub>S is very likely to undergo chemical desorption (also sometimes referred to as "reactive desorption") as a result of its formation via hydrogen atom addition to HS radicals,<sup>128,129</sup> which may account for its nondetection in interstellar icy grain mantles but its positive detection in the gas-phase within interstellar media.<sup>130–132</sup> In any case, our results suggest that the possible formation and retention of H<sub>2</sub>S in the ice phase as a result of the cosmic ray or stellar wind irradiation of H<sub>2</sub>O ice on allotropic sulfur is not efficient, and thus, alternative strategies to explain its hypothesized presence in icy astrophysical environments must be sought.

As can be noted in Figure 14 and Table 4, the most commonly observed sulfur-bearing products in our experiments were SO<sub>2</sub> and CS<sub>2</sub>. Comparing the G-values of these molecules, it is possible to note two distinct trends: first, SO<sub>2</sub> is in general more readily produced than CS<sub>2</sub> as demonstrated by its higher G-values, and second, the irradiation of a given ice on top of sulfur at 70 K is more productive than the analogous irradiation at 20 K, most likely due to the higher mobilities of radiolytically derived radicals at higher temperatures. Given that SO<sub>2</sub> and CS<sub>2</sub> were observed in experiments performed at both 20 and 70 K, our results therefore indicate that the irradiation of volatile ices on top of allotropic sulfur is a plausible strategy toward the synthesis of these molecules in both interstellar icy grain mantles as well as on the surfaces of icy outer solar system bodies. Therefore, it may be worthwhile for future observational studies to search for spectroscopic evidence of CS<sub>2</sub> in astrophysical ices, such as in interstellar icy grain mantles where it may coexist alongside the already detected SO<sub>2</sub>.<sup>11,12</sup>

Another result of interest was the comparatively low G-value of OCS, which was only detected after the irradiation of CO<sub>2</sub> ice on sulfur at 70 K. The fact that OCS could not be detected in our experiments conducted at the more interstellar-relevant temperature of 20 K suggests that the cosmic ray irradiation of



**Figure 14.** Calculated *G*-values for the sulfur-bearing products observed as a result of irradiating different ices on top of allotropic sulfur using 1 MeV He<sup>+</sup> ions. Note that circles represent data points acquired at 20 K, while diamonds represent those acquired at 70 K. Information about error bars can be found in Table 4.

Table 4. G-Values Calculated for the Sulfur-Bearing Products Observed as a Result of Irradiating Different Neat Ices on Top of Allotropic Sulfur Using 1 MeV He<sup>+</sup> Ions

	G-values (molecules per 100 eV) for sulfur products			
ice	SO <sub>2</sub>	CS <sub>2</sub>	ocs	H <sub>2</sub> SO <sub>4</sub> hydrates
O <sub>2</sub> ice at 20 K	$\begin{array}{r} 0.0457 \pm \\ 0.0004 \end{array}$			
CO ice at 20 K		$(6.66 \pm 0.50) \times 10^{-4}$		
CO <sub>2</sub> ice at 20 K	$\begin{array}{c} 0.0078 \pm \\ 0.0014 \end{array}$	$(5.89 \pm 0.13) \times 10^{-4}$		
CO <sub>2</sub> ice at 70 K	$\begin{array}{r} 0.0104 \pm \\ 0.0003 \end{array}$	$(9.99 \pm 0.19) \times 10^{-4}$	$(0.89 \pm 0.01) \times 10^{-4}$	
H <sub>2</sub> O ice at 20 K				
H <sub>2</sub> O ice at 70 K				$0.4163 \pm 0.0031$

volatile ices on top of allotropic sulfur is not a major contributor toward the formation of this species, strongly contrasting with the apparent ease with which SO<sub>2</sub> is formed in this way. This result complements the recent observational surveys of  $SO_2$  and OCS toward protostellar regions carried out by Santos et al.,<sup>133</sup> who measured a weak correlation between the gas-phase column densities of these two sulfurbearing molecules. This was interpreted as the result of these two molecules having different formation chemistries during distinct interstellar cloud evolutionary phases. Indeed, in that and subsequent studies, Santos et al.<sup>133,134</sup> proposed that solid  $SO_2$  is likely formed as a result of the energetic processing (e.g., by galactic cosmic rays) during the low-density phase of interstellar molecular clouds when the surface-catalyzed formation of oxygen-rich molecules (e.g., H2O, O2, and  $CO_2$ ) is a dominant process.<sup>135,136</sup> Conversely, the formation of OCS was proposed to occur after the catastrophic freeze-out of gas-phase CO as a result of a neutral-neutral reaction between CO molecules and HS radicals to yield HSCO as an intermediate species, which, upon hydrogenation, dissociates to OCS and  $H_2$ .<sup>134,137,138</sup> The notion of SO<sub>2</sub> in interstellar ices forming primarily via an "energetic" route and OCS forming via a "nonenergetic" route is not only in agreement with our present findings, but also with the experimentally demonstrated radiolytic instability of OCS in interstellar ice analogs.<sup>46,87,139</sup>

The production of H<sub>2</sub>SO<sub>4</sub> hydrates as a result of the irradiation of H<sub>2</sub>O ice on top of sulfur at 70 K was found to have the highest calculated G-value (Figure 14 and Table 4), thus highlighting this reaction as an especially productive one. The result of this experiment is in line with those of several previous studies; for instance, early studies by Donaldson and Johnston<sup>140</sup> and Johnston and Donaldson<sup>141</sup> investigated the irradiation of colloidal suspensions of elemental sulfur in deaerated H<sub>2</sub>O and found evidence for the solubilization of the sulfur and the subsequent formation of H<sub>2</sub>SO<sub>4</sub>. The mechanism behind this chemistry was later described by Della Guardia and Johnston<sup>123</sup> (Scheme 1). Perhaps the first work to consider the irradiation of H2O:S8 mixtures in astrophysical ice analogs was that conducted by Carlson,<sup>1</sup> who investigated the chemical evolution of S<sub>8</sub> that was mixed with liquid H<sub>2</sub>O and subsequently frozen prior to irradiation. Their work demonstrated that not only is H<sub>2</sub>SO<sub>4</sub> efficiently formed by solid-phase radiation chemistry, but that this synthesis is in fact more efficient at 77 K (which is similar to our 70 K irradiation temperature) than at 195 K. Taken together, these results indicate that sulfur allotropes are likely to play a pivotal role in the chemistry of cold outer solar system bodies whose surfaces are characterized by temperatures of circa 70 K, such as Jupiter's moon Europa.<sup>143</sup>

Indeed, on the surface of Europa, allotropic sulfur is thought to be the second-most abundant reservoir of the element, behind  $H_2SO_4$  hydrates.<sup>142,144</sup> Given that the surface of Europa is also dominated by H2O ice, our results suggest that the irradiation of this ice on top of deposits of allotropic sulfur could represent a possible route toward the formation of H<sub>2</sub>SO<sub>4</sub> hydrates and thus could contribute to the so-called radiolytic sulfur cycle taking place on Europa's surface.<sup>1,45,119,120,125,142,144,145,146</sup> It should be noted, however, that the current prevailing hypothesis as to the major source of H<sub>2</sub>SO<sub>4</sub> hydrates on Europa is the implantation of energetic sulfur ions from the giant Jovian magnetosphere,<sup>125,147,148,149</sup> which would not only explain the surface abundance of the acid hydrates observed but also their "bulls eye"-like distribution and enhanced presence on the trailing hemi-sphere.<sup>119,120,150,153,154</sup> Systematic laboratory work by Ding et al.<sup>125</sup> has strengthened this hypothesis by demonstrating that the implantation of sulfur ions with kinetic energies of 35-200 keV into H<sub>2</sub>O ice at 80 K results in the production of H<sub>2</sub>SO<sub>4</sub> with increasing efficiencies at higher ion energies. Sulfur ion implantations into other ices have also been studied both experimentally and theoretically,<sup>151,152,155–162</sup> and work by Mifsud et al.<sup>151,152</sup> demonstrated that the implantation of 400 keV sulfur ions into O<sub>2</sub> ice at 20 K results in the production of  $SO_{2}$ ; this result echoes our present finding with regard to the irradiation of O2 ice on top of allotropic sulfur at the same temperature.

In conclusion, our experiments have demonstrated that the irradiation of astrophysical ice analogs on top of refractory sulfur using 1 MeV He<sup>+</sup> ion beams (used as a mimic of space radiation) results in the formation of various simple inorganic molecules, including SO<sub>2</sub>, CS<sub>2</sub>, OCS, and H<sub>2</sub>SO<sub>4</sub> hydrates, but notably not H<sub>2</sub>S. The formation of these species from different ices on top of sulfur has been characterized by quantifying their respective *G*-values, and reactions leading to their radiolytic synthesis have been suggested. We emphasize that further work

is required to definitively identify the specific reactions leading to the sulfur-bearing products observed in our experiments, including further work on the possible contribution of ionmediated reactions compared with the neutral-neutral reactions proposed herein. Our results have demonstrated that SO<sub>2</sub> and CS<sub>2</sub> are the most commonly observed products among the different ices considered, but that the formation of  $H_2SO_4$  hydrates as a result of the irradiation of  $H_2O$  ice on sulfur at 70 K is associated with the highest G-value. Finally, we have discussed the astrochemical implications of our work in detail and propose that CS2 is a good candidate molecule for future identification in astrophysical ices by observational surveys. Furthermore, we have provided experimental evidence that SO<sub>2</sub> in interstellar ices could be primarily formed via energetic mechanisms while OCS is more likely formed as a result of nonenergetic processes, in agreement with recently published results from experimental and observational studies.

#### AUTHOR INFORMATION

#### **Corresponding Authors**

- Duncan V. Mifsud Centre for Astrophysics and Planetary Science, School of Physics and Astronomy, University of Kent, Canterbury CT2 7NH, United Kingdom; HUN-REN Institute for Nuclear Research (Atomki), Debrecen H-4026, Hungary; orcid.org/0000-0002-0379-354X; Email: mifsud.duncan@atomki.hu
- Nigel J. Mason Centre for Astrophysics and Planetary Science, School of Physics and Astronomy, University of Kent, Canterbury CT2 7NH, United Kingdom; HUN-REN Institute for Nuclear Research (Atomki), Debrecen H-4026, Hungary; Email: n.j.mason@kent.ac.uk

#### Authors

- Zuzana Kaňuchová Astronomical Institute, Slovak Academy of Sciences, Tatranská Lomnica SK-059 60, Slovakia
- **Olivier Auriacombe** Department of Microtechnology and Nanoscience, Chalmers University of Technology, Göteborg 412 96, Sweden
- Péter Herczku HUN-REN Institute for Nuclear Research (Atomki), Debrecen H-4026, Hungary; Octid.org/0000-0002-1046-1375
- Danna Qasim Southwest Research Institute, San Antonio, Texas 78238, United States
- Sándor T. S. Kovács HUN-REN Institute for Nuclear Research (Atomki), Debrecen H-4026, Hungary
- Richárd Rácz HUN-REN Institute for Nuclear Research (Atomki), Debrecen H-4026, Hungary
- Béla Sulik HUN-REN Institute for Nuclear Research (Atomki), Debrecen H-4026, Hungary
- Zoltán Juhász HUN-REN Institute for Nuclear Research (Atomki), Debrecen H-4026, Hungary
- István Rajta HUN-REN Institute for Nuclear Research (Atomki), Debrecen H-4026, Hungary
- István Vajda HUN-REN Institute for Nuclear Research (Atomki), Debrecen H-4026, Hungary
- Sándor Biri HUN-REN Institute for Nuclear Research (Atomki), Debrecen H-4026, Hungary
- **Robert W. McCullough** Department of Physics and Astronomy, School of Mathematics and Physics, Queen's University Belfast, Belfast BT7 1NN, United Kingdom
- Sergio Ioppolo Centre for Interstellar Catalysis, Department of Physics and Astronomy, Aarhus University, Aarhus DK-8000, Denmark; orcid.org/0000-0002-2271-1781

Ujjwal Raut – Southwest Research Institute, San Antonio, Texas 78238, United States

Complete contact information is available at:

https://pubs.acs.org/10.1021/acsearthspacechem.5c00036

#### **Author Contributions**

Duncan V. Mifsud: conceptualization, methodology, validation, formal analysis, investigation, data curation, writing-(original draft), writing (review and editing), visualization, project administration; Zuzana Kaňuchová: formal analysis, data curation, writing (review and editing); Olivier Auriacombe: conceptualization, resources, writing (review and editing), funding acquisition; Péter Herczku: methodology, validation, investigation, writing (review and editing); Danna Qasim: conceptualization, visualization, writing (review and editing), project administration; Sándor T. S. Kovács: methodology, validation, investigation, writing (review and editing); Richárd Rácz: methodology, validation, investigation, writing (review and editing); Béla Sulik: methodology, validation, investigation, writing (review and editing); Zoltán Juhász: methodology, validation, investigation, writing (review and editing); István Rajta: resources, writing (review and editing); István Vajda: resources, writing (review and editing); Sándor Biri: resources, writing (review and editing); Robert W. McCullough: writing (review and editing), supervision; Sergio Ioppolo: resources, writing (review and editing), supervision, project administration, funding acquisition; Ujjwal Raut: visualization, writing (review and editing); Nigel J. Mason: resources, writing (review and editing), supervision, project administration, funding acquisition.

#### Notes

The authors declare no competing financial interest.

#### ACKNOWLEDGMENTS

This article is dedicated to the memory of our dear friend and colleague Harold Linnartz (1965-2023), who is sorely missed by the team and the wider research community. The authors acknowledge support from the Europlanet 2024 RI, which has been funded by the European Union's Horizon 2020 Research Innovation Program under grant agreement no. 871149. The main components of the ICA setup were purchased using funds obtained from the Royal Society through grants UF130409, RGF/EA/180306, and URF/R/191018. Further developments of the installation were supported in part by the Eötvös Loránd Research Network through grants ELKH IF-2/ 2019 and ELKH IF-5/2020. This work has also received support from the European Union and the State of Hungary; cofinanced by the European Regional Development Fund through grant GINOP-2.3.3-15-2016-00005. Support has also been received from the Research, Development, and Innovation Fund of Hungary through grant nos. K128621 and ADVANCED-151196. This paper is also based on work from the COST Actions CA20129 "MultIChem" and CA22133 "PLANETS", supported by COST (European Cooperation in Science and Technology). Duncan V. Mifsud is the grateful recipient of a University of Kent Vice-Chancellor's Research Scholarship. The research of Zuzana Kaňuchová is supported by the Slovak Grant Agency for Science (grant no. 2/0059/22). Danna Qasim acknowledges the "joint" effort from SwRI and STScI. Support for programme no. 4810 was provided through a grant from the STScI under NASA contract no. NAS5-03127. Danna Qasim

would also like to extend her deepest gratitude to Harold Linnartz for his invaluable guidance as her Ph.D. supervisor. Zoltán Juhász is grateful for the support of the Hungarian Academy of Sciences through the János Bolyai Research Scholarship. Robert W. McCullough is the grateful recipient of an honorary visiting scholar post at Queen's University Belfast. Sergio Ioppolo acknowledges support from the Danish National Research Foundation through the Centre of Excellence "InterCat" (grant agreement no. DNRF150), as well as from the Royal Society.

#### REFERENCES

(1) Mifsud, D. V.; Kaňuchová, Z.; Herczku, P.; Ioppolo, S.; Juhász, Z.; Kovács, S. T. S.; Mason, N. J.; McCullough, R. W.; Sulik, B.; et al. Sulfur Ice Astrochemistry: A Review of Laboratory Studies. *Space Sci. Rev.* **2021**, *217*, 14.

(2) Geballe, T. R.; et al. New Infrared Absorption Features due to Solid Phase Molecules Containing Sulfur in W33A. *Astron. Astrophys.* **1985**, *146*, L6.

(3) Tieftrunk, A.; et al. SO and  $H_2S$  in Low Density Molecular Clouds. Astron. Astrophys. **1994**, 289, 579.

(4) Palumbo, M. E.; et al. Solid Carbonyl Sulfide (OCS) in W33A. Astrophysical Journal **1995**, 449, 674.

(5) Boogert, A. C. A.; et al. Infrared Observations and Laboratory Simulations of Interstellar  $CH_4$  and  $SO_2$ . *Astron. Astrophys.* **1997**, *317*, 929.

(6) Ruffle, D. P.; et al. The Sulfur Depletion Problem. Mon. Not. R. Astron. Soc. 1999, 306, 691.

(7) Jenkins, E. B. A Unified Representation of Gas-Phase Element Depletions in the Interstellar Medium. *Astrophysical Journal* **2009**, 700, 1299.

(8) Laas, J. C.; Caselli, P. Modeling Sulfur Depletion in Interstellar Clouds. *Astron. Astrophys.* 2019, 624, A108.

(9) Vidal, T. H. G.; et al. On the Reservoir of Sulphur in Dark Clouds: Chemistry and Elemental Abundance Reconciled. *Mon. Not. R. Astron. Soc.* **2017**, *469*, 435.

(10) Navarro-Almaida, D.; et al. Gas Phase Elemental Abundances in Molecular Clouds (GEMS)-II. On the Quest for the Sulfur Reservoir in Molecular Clouds: the  $H_2S$  Case. *Astron. Astrophys.* **2020**, 637, A39.

(11) Boogert, A. C. A.; et al. Observations of the Icy Universe. Annual Review of Astronomy and Astrophysics **2015**, 53, 541.

(12) McClure, M. K.; et al. An Ice Age JWST Inventory of Dense Molecular Clouds Ices. *Nature Astronomy* **2023**, *7*, 431.

(13) Altwegg, K.; et al. Abundant Ammonium Hydrosulfide Embedded in Cometary Dust Grains. *Mon. Not. R. Astron. Soc.* **2022**, *516*, 3900.

(14) Slavicinska, K.; et al. Ammonium Hydrosulfide ( $NH_4SH$ ) as a Potentially Significant Sulfur Sink in Interstellar Ices. Astron. Astrophys. **2025**, 693, A146.

(15) Gunning, H. E.; Strausz, O. P. The Reactions of Sulfur Atoms. *Adv. Photochem.* **1966**, *4*, 143.

(16) Brom, J. M.; Lepak, E. J. Afterglow from the Photodissociation of OCS in an Argon Matrix at 4 K. *Chem. Phys. Lett.* **1976**, *41*, 185.

(17) Smardzewski, R. R. Chemiluminescent Matrix Reactions of Atomic Oxygen, Sulfur, and O  $({}^{3}P)$  + H<sub>2</sub>S. J. Chem. Phys. **1978**, 68, 2878.

(18) Lee, Y. P.; Pimentel, G. C. Chemiluminescence of  $S_2$  in Solid Argon. J. Chem. Phys. **1979**, 70, 692.

(19) Shingledecker, C. N.; et al. The Effects of Cosmic-Ray-Driven Radiation Chemistry and Nondiffusive Bulk Reactions. *Astrophysical Journal* **2020**, 888, 52.

(20) Garrod, R. T.; et al. Formation of Complex Organic Molecules in Hot Molecular Cores Through Nondiffusive Grain-Surface and Ice-Mantle Chemistry. *Astrophysical Journal Supplement Series* **2022**, 259, 1.

(21) Cazaux, S.; et al. Photoprocessing of  $H_2S$  on Dust Grains: Building S Chains in Translucent Clouds and Comets. *Astron. Astrophys.* **2022**, 657, A100. (22) Carrascosa, H.; et al. Formation and Desorption of Sulfur Chains  $(H_2S_x \text{ and } S_x)$  in Cometary Ice: Effects of Ice Composition and Temperature. *Mon. Not. R. Astron. Soc.* **2024**, *533*, 967.

(23) Garozzo, M.; et al. The Fate of S-Bearing Species after Ion Irradiation of Interstellar Icy Grain Mantles. *Astron. Astrophys.* **2010**, *509*, A67.

(24) Gomis, O.; Strazzulla, G. Ion Irradiation Studies of  $H_2O$  Ice on Top of Sulfurous Solid Residues and its Relevance to the Galilean Satellites. *Icarus* **2008**, *194*, 146.

(25) Strazzulla, G.; et al. The Origin of Sulfur-Bearing Species on the Surfaces of Icy Satellites. *Adv. Space Res.* **2009**, *43*, 1442.

(26) Mahjoub, A.; et al. Production of Sulfur Allotropes in Electron Irradiated Jupiter Trojans Ice Analogs. *Astrophysical Journal* **2017**, *846*, 148.

(27) Mifsud, D. V.; Herczku, P.; Rácz, R.; Rahul, K. K.; Kovács, S. T. S.; Juhász, Z.; Sulik, B.; Biri, S.; McCullough, R. W.; Kaňuchová, Z.; Ioppolo, S.; Hailey, P. A.; Mason, N. J.; et al. Energetic Electron Irradiations of Amorphous and Crystalline Sulfur-Bearing Astrochemical Ices. *Front. Chem.* **2022**, *10*, 1003163.

(28) Trofimov, B. A.; et al. Vibrations of the S-S Bond in Elemental Sulfur and Organic Polysulfides: A Structural Guide. *Journal of Sulfur Chemistry* **2009**, *30*, 518.

(29) Aponte, J. C.; Dworkin, J. P.; Glavin, D. P.; Elsila, J. E.; Parker, E. T.; McLain, H. L.; Naraoka, H.; Okazaki, R.; Takano, Y.; Tachibana, S.; Dong, G.; Zeichner, S. S.; Eiler, J. M.; Yurimoto, H.; Nakamura, T.; Yabuta, H.; Terui, F.; Noguchi, T.; Sakamoto, K.; Yada, T.; Nishimura, M.; Nakato, A.; Miyazaki, A.; Yogata, K.; Abe, M.; Okada, T.; Usui, T.; Yoshikawa, M.; Saiki, T.; Tanaka, S.; Nakazawa, S.; Tsuda, Y.; Watanabe, S. i.; et al. PAHs, Hydrocarbons, and Dimethylsulfides in Asteroid Ryugu Samples A0106 and C0107 and the Orgueil (CI1) Meteorite. *Earth, Planets, and Space* **2023**, 75, 28.

(30) Calmonte, U.; et al. Sulfur-Bearing Species in the Coma of Comet 67P/Churyumov-Gerasimenko. *Mon. Not. R. Astron. Soc.* 2016, 462, S253.

(31) Hazen, R. M.; et al. Mineral Evolution. Am. Mineral. 2008, 93, 1693.

(32) Hony, S.; et al. The Carrier of the 30  $\mu$ m Emission Feature in Evolved Stars: A Simple Model Using Magnesium Sulfide. *Astron. Astrophys.* **2002**, 390, 533.

(33) Hony, S.; et al. The Detection of Iron Sulfides in Planetary Nebulae. *Astron. Astrophys.* **2002**, 393, L103.

(34) Kama, M.; et al. Abundant Refractory Sulfur in Protoplanetary Disks. *Astrophysical Journal* **2019**, *885*, 114.

(35) Öberg, K. I. Photochemistry and Astrochemistry: Photochemical Pathways to Interstellar Complex Organic Molecules. *Chem. Rev.* 2016, *116*, 9631.

(36) Arumainayagam, C. R.; et al. Extraterrestrial Prebiotic Molecules: Photochemistry vs. Radiation Chemistry of Interstellar Ices. *Chem. Soc. Rev.* 2019, 48, 2293.

(37) Herbst, E.; Garrod, R. T. Synthetic Approaches to Complex Organic Molecules in the Cold Interstellar Medium. *Frontiers in Astronomy and Space Sciences* **2022**, *8*, No. 789428.

(38) Mennella, V.; et al. Formation of CO and  $CO_2$  Molecules by Ion Irradiation of Water Ice-Covered Hydrogenated Carbon Grains. *Astrophysical Journal* **2004**, *615*, 1073.

(39) Raut, U.; et al. Radiation Synthesis of Carbon Dioxide in Ice-Coated Carbon: Implications for Interstellar Grains and Icy Moons. *Astrophysical Journal* **2012**, *752*, 159.

(40) Gomis, O.; Strazzulla, G.  $CO_2$  Production by Ion Irradiation of  $H_2O$  on Top of Carbonaceous Materials and its Relevance to the Galilean Satellites. *Icarus* **2005**, *177*, 570.

(41) Mennella, V.; et al. Synthesis of CO and  $CO_2$  Molecules by UV Irradiation of Water Ice-Covered Hydrogenated Carbon Grains. Astrophysical Journal **2006**, 643, 923.

(42) Fulvio, D.; et al. Photosynthesis of Carbon Dioxide from Carbon Surfaces Coated with Oxygen: Implications for Interstellar Molecular Clouds and the Outer Solar System. *Astrophysical Journal Letters* **2012**, *752*, L33. (43) Ferrari, P.; Berden, G.; Redlich, B.; Waters, L. B. F. M.; Bakker, J. M.; et al. Laboratory Infrared Spectra and Fragmentation Chemistry of Sulfur Allotropes. *Nat. Commun.* **2024**, *15*, 5928.

(44) Yamada, M.; et al. A Comprehensive Investigation on the Formation of Organosulfur Molecules in Dark Clouds via Neutral-Neutral Reactions. *Astron. Astrophys.* **2002**, *395*, 1031.

(45) Moore, M. H.; et al. The Radiolysis of  $SO_2$  and  $H_2S$  in Water Ice: Implications for the Icy Jovian Satellites. *Icarus* 2007, 189, 409.

(46) Ferrante, R. F.; et al. Formation of Interstellar OCS: Radiation Chemistry and IR Spectra of Precursor Ices. *Astrophysical Journal* **2008**, 684, 1210.

(47) Wang, J.; et al. Formation of Thioformic Acid (HCOSH): The Simplest Thioacid in Interstellar Ice Analogs. *J. Phys. Chem. A* 2022, 126, 9699.

(48) Santos, J. C.; et al. Formation of S-Bearing Complex Organic Molecules in Interstellar Clouds via Ice Reactions with  $C_2H_2$ , HS, and Atomic H. ACS Earth and Space Chemistry **2024**, 8, 1646.

(49) Caffrey, J. M.; Allen, A. O. Radiolysis of Pentane Adsorbed on Mineral Solids. J. Phys. Chem. **1958**, 62, 33.

(50) Sutherland, J. W.; Allen, A. O. Radiolysis of Pentane in the Adsorbed State. J. Am. Chem. Soc. 1961, 83, 1040.

(51) Hentz, R. R. Radiation-Induced Reactions of Isopropylbenzene on Silica-Alumina. J. Phys. Chem. **1964**, 68, 2889.

(52) Rabe, J. G.; et al. Radiolysis and Energy Transfer in the Adsorbed State. J. Phys. Chem. 1966, 70, 1098.

(53) LaVerne, J. A.; Tandon, L. H<sub>2</sub> Production in the Radiolysis of Water on UO<sub>2</sub> and Other Oxides. *J. Phys. Chem. B* **2003**, *107*, 13623.

(54) Herczku, P.; Mifsud, D. V.; Ioppolo, S.; Juhász, Z.; Kaňuchová, Z.; Kovács, S. T. S.; Traspas Muinña, A.; Hailey, P. A.; Rajta, I.; Vajda, I.; Mason, N. J.; McCullough, R. W.; Paripás, B.; Sulik, B.; et al. The Ice Chamber for Astrophysics-Astrochemistry (ICA): A New Experimental Facility for Ion Impact Studies of Astrophysical Ice Analogs. *Rev. Sci. Instrum.* **2021**, *92*, No. 084501.

(55) Mifsud, D. V.; Juhász, Z.; Herczku, P.; Kovács, S. T. S.; Ioppolo, S.; Kaňuchová, Z.; Czentye, M.; Hailey, P. A.; Muiña, A. T.; Mason, N. J.; McCullough, R. W.; Paripás, B.; Sulik, B.; et al. Electron Irradiation and Thermal Chemistry Studies of Interstellar and Planetary Ice Analogs at the ICA Astrochemistry Facility. *Eur. Phys. J. D* **2021**, 75, 182.

(56) Meyer, B. Elemental Sulfur. Chem. Rev. 1976, 76, 367.

(57) Rajta, I.; et al. Accelerator Characterization of the New Ion Beam Facility at MTA Atomki in Debrecen, Hungary. *Nuclear Instruments and Methods in Physics Research A* **2018**, 880, 125.

(58) Biri, S.; Vajda, I. K.; Hajdu, P.; Rácz, R.; Csík, A.; Kormány, Z.; Perduk, Z.; Kocsis, F.; Rajta, I.; et al. The Atomki Accelerator Center. *Eur. Phys. J. Plus* **2021**, *136*, 247.

(59) Simpson, J. A. Elemental and Isotopic Composition of the Galactic Cosmic Rays. *Annual Review of Nuclear and Particle Science* **1983**, 33, 323.

(60) Mewaldt, R. A. Galactic Cosmic Ray Composition and Energy Spectra. *Adv. Space Res.* **1994**, *14*, 737.

(61) Ziegler, J. F.; et al. SRIM – The Stopping and Range of Ions in Matter. Nuclear Instruments and Methods in Physics Research B 2010, 268, 1818.

(62) Gerakines, P. A.; et al. The Infrared Band Strengths of  $H_2O$ , CO, and  $CO_2$  in Laboratory Simulations of Astrophysical Ice Mixtures. *Astron. Astrophys.* **1995**, *296*, 810.

(63) Greenwood, N. N.; Earnshaw, A. "Sulfur." In: *Chemistry of the Elements*, pp 645–746. Butterworth-Heinemann, 1997.

(64) Freiman, Y. A.; Jodl, H. J. Solid Oxygen. Phys. Rep. 2004, 401, 1.

(65) Luna, R.; et al. Density and Refractive Index of Carbon Monoxide Ice at Different Temperatures. *Astrophysical Journal* **2022**, 935, 134.

(66) Satorre, MA; et al. Density of  $CH_4$ ,  $N_2$ , and  $CO_2$  Ices at Different Temperatures of Deposition. *Planetary and Space Science* **2008**, 56, 1748.

(67) Narten, A. H.; et al. Diffraction Pattern and Structure of Amorphous Solid Water at 10 and 77 K. J. Chem. Phys. **1976**, 64, 1106.

(68) Mifsud, D. V.; et al. A Systematic Mid-Infrared Spectroscopic Study of Thermally Processed SO<sub>2</sub> Ices. *Phys. Chem. Chem. Phys.* **2023**, 25, 26278.

(69) Garozzo, M.; et al. H-Implantation in  $SO_2$  and  $CO_2$  Ices. Planetary and Space Science **2008**, 56, 1300.

(70) Mayer, R. Organic Chemistry of Sulfur, pp 33-69. Springer, 1977.

(71) Hills, A. J.; et al. Kinetics of the Reactions of Diatomic Sulfur with Atomic Oxygen, Molecular Oxygen, Ozone, Nitrous Oxide, Nitric Oxide, and Nitrogen Dioxide. *J. Phys. Chem.* **1987**, *91*, 1199.

(72) Long, S. R.; Pimentel, G. C. Chemiluminescent Reactions of Sulfur  $({}^{3}P_{2})$  Atoms in Cryogenic Matrices: S+O<sub>2</sub>  $\rightarrow$  SO<sub>2</sub>  $(a^{3}B_{1})$ . J. Chem. Phys. 1977, 66, 2219.

(73) Lotnik, S. V.; Kazakov, V. P. Chemiluminescence in Cryogenic Matrices. *Russ. Chem. Rev.* **1989**, *58*, 337.

(74) Chaabouni, H.; et al. Infrared Spectroscopy of Neat Solid Ozone and That of Ozone in Interaction with Amorphous and Crystalline Water Ice. J. Phys. Chem. A **2000**, 104, 6962.

(75) Bennett, C. J.; Kaiser, R. I. Laboratory Studies on the Formation of Ozone on Icy Satellites and on Interstellar and Cometary Ices. *Astrophysical Journal* **2005**, *635*, 1362.

(76) Sivaraman, B.; et al. Temperature-Dependent Formation of Ozone in Solid Oxygen by 5 keV Electron Irradiation and Implications for Solar System Ices. *Astrophysical Journal* **2007**, *669*, 1414.

(77) Ennis, C. P.; et al. On the Formation of Ozone in Oxygen-Rich Solar System Ices via Ionizing Radiation. *Phys. Chem. Chem. Phys.* **2011**, *13*, 9469.

(78) Sivaraman, B.; Raja Sekhar, B. N.; Fulvio, D.; Hunniford, A.; McCullough, B.; Palumbo, M. E.; Mason, N.; et al. Ozone and Carbon Trioxide Synthesis by Low Energy Ion Implantation onto Solid Carbon Dioxide and Implications to Astrochemistry. *J. Chem. Phys.* **2013**, *139*, No. 074706.

(79) Mifsud, D. V.; et al. Ozone Production in Electron Irradiated CO<sub>2</sub>:O<sub>2</sub> Ices. *Phys. Chem. Chem. Phys.* **2022**, *24*, 18169.

(80) Trottier, A.; Brooks, R. L. Carbon-Chain Oxides in Proton-Irradiated CO Ice Films. *Astrophysical Journal* **2004**, *612*, 1214.

(81) Loeffler, M. J.; et al.  $CO_2$  Synthesis in Solid CO by Lyman- $\alpha$  Photons and 200 keV Protons. *Astron. Astrophys.* **2005**, 435, 587.

(82) Palumbo, M. E.; et al. Detection of  $C_3O$  in the Low-Mass Protostar Elias 18. Astrophysical Journal 2008, 685, 1033.

(83) Jamieson, C. S.; et al. Understanding the Kinetics and Dynamics of Radiation-Induced Reaction Pathways in Carbon Monoxide Ice at 10 K. Astrophysical Journal Supplement Series 2006, 163, 184.

(84) Bennett, C. J.; et al. Mechanistical Studies on the Formation of Carbon Dioxide in Extraterrestrial Carbon Monoxide Ice Analog Samples. *Phys. Chem. Chem. Phys.* **2009**, *11*, 4210.

(85) Rolfes, T. R.; et al. The Chemiluminescent Reaction of Oxygen Atoms with Sulfur Monoxide at Low Pressures. *J. Phys. Chem.* **1965**, 69, 849.

(86) Jiménez-Escobar, A.; et al. Sulfur Depletion in Dense Clouds and Circumstellar Regions: Organic Products Made from UV Photoprocessing of Realistic Ice Analogs Containing H<sub>2</sub>S. *Mon. Not. R. Astron. Soc.* **2014**, *443*, 343.

(87) Sivaraman, B. Electron Irradiation of Carbon Dioxide-Carbon Disulfide Ice Analog and its Implication on the Identification of Carbon Disulfide on Moon. *Journal of Chemical Sciences* **2016**, *128*, 159.

(88) Ikeda, A.; et al. Photodissociation Dynamics of OCS And  $CS_2$  Adsorbed on Water Ice Films at 193 nm. *Journal of Photochemistry and Photobiology A* **2008**, 195, 330.

(89) Deeyamulla, M. P.; Husain, D. Kinetic Studies of Atomic Carbon,  $C[2p^{2}(^{3}P_{J})]$ , with Small Sulfur-Containing Molecules by Time-Resolved Atomic Resonance Absorption Spectroscopy in the

Vacuum Ultra-Violet. Journal of Photochemistry and Photobiology A 2006, 184, 347.

(90) Martín-Doménech, R.; et al. UV Photoprocessing of  $CO_2$  Ice: A Complete Quantification of Photochemistry and Photon-Induced Desorption Processes. *Astron. Astrophys.* **2015**, *584*, A14.

(91) Mifsud, D. V.; et al. Mid-IR and VUV Spectroscopic Characterization of Thermally Processed and Electron Irradiated  $CO_2$  Astrophysical Ice Analogs. J. Mol. Spectrosc. 2022, 385, No. 111599.

(92) Pilling, S.; et al. Chemical Evolution of  $CO_2$  Ices Under Processing by Ionizing Radiation: Characterization of Nonobserved Species and Chemical Equilibrium Phase with the Employment of PROCODA Code. *Astrophysical Journal* **2022**, *925*, 147.

(93) Pilling, S.; et al. Mapping the Evolution of Pure  $CO_2$  Ices Irradiated by Ions, UV, and Electrons Using the Upgraded PROCODA Code (Employing an Effective Rate Constant Ordering by Thermochemistry Data). *Adv. Space Res.* **2023**, *71*, 5466.

(94) Bennett, C. J.; et al. Mechanistical Studies on the Formation and Destruction of Carbon Monoxide (CO), Carbon Dioxide (CO<sub>2</sub>), and Carbon Trioxide (CO<sub>3</sub>) in Interstellar Ice Analog Samples. *Phys. Chem. Chem. Phys.* **2010**, *12*, 4032.

(95) Froese, R. D. J.; Goddard, J. D. *Ab Initio* Studies of the Lowest Singlet and Triplet Potential Energy Surfaces of CO<sub>2</sub>S. *Mol. Phys.* **1993**, *79*, 685.

(96) Hawkins, M.; et al. Photochemistry of Low-Temperature Matrixes Containing Carbonyl Sulfide: Reactions of Sulfur Atoms with the Phosphorus Trihalides Phosphorus Trifluoride ( $PF_3$ ) and Phosphorus Trichloride ( $PCl_3$ ) and the Hydrocarbons Methane, Ethene, and Ethyne. *J. Phys. Chem.* **1985**, *89*, 3326.

(97) Lo, W. J.; et al. Isomers of  $OCS_2$ : IR Absorption Spectra of OSCS and  $O(CS_2)$  in Solid Ar. J. Chem. Phys. **2004**, 121, 12371.

(98) Jones, P. R.; Taube, H. Photochemical Studies on Ozone with Carbon Disulfide and with Carbonyl Sulfide in Low-Temperature Matrices. *J. Phys. Chem.* **1973**, *77*, 1007.

(99) Hsu, D. S. Y.; et al. Dynamics of Reactions of O ( $^{3}P$ ) Atoms with CS, CS<sub>2</sub>, and OCS. *Chem. Phys.* **1979**, 44, 143.

(100) Chen, X.; et al. Internal Energy Distributions of the SO( $X^{3}\Sigma^{-}$ ) Product from the O(<sup>3</sup>P)+OCS Reaction. *Chem. Phys. Lett.* **1995**, 247, 313.

(101) van Veen, N.; et al. The Adiabatic and Diabatic Reactions of  $S(^{1}D)$  Atoms with OCS: Internal State Distribution of the  $S_{2}$  Products. *J. Chem. Phys.* **1983**, 79, 4295.

(102) Moore, M. H.; et al. Vaporization of Ices Containing  $S_2$  – Implications for Comets. *Icarus* 1988, 74, 399.

(103) Mejía, C.; et al. Radiolysis and Sputtering of Carbon Dioxide Ice Induced by Swift Ti, Ni, and Xe Ions. *Nuclear Instruments and Methods in Physics Research B* **2015**, 365, 477.

(104) Cataldo, F. On the Photopolymerization of Carbon Disulfide. *Inorg. Chim. Acta* **1995**, 232, 27.

(105) Cataldo, F.; Heymann, D. Carbon Disulfide Sonopolymer: A Comparison with Other Carbon Disulfide Polymers. *European Journal of Solid State and Inorganic Chemistry* **1998**, *35*, 619.

(106) Cataldo, F. On the Radiopolymerization of Carbon Disulfide. *Radiat. Phys. Chem.* **2000**, 58, 217.

(107) Khriachtchev, L.; et al. Low-Temperature Thermoluminescence in Solid Argon: Short-Range Mobility of Atoms. *J. Chem. Phys.* **1999**, *111*, 1650.

(108) Kiljunen, T.; et al. Electronic Structure and Short-Range Recombination Dynamics of  $S_2$  in Solid Argon. J. Chem. Phys. 2000, 112, 7475.

(109) Hopkins, A. G.; Brown, C. W. Infrared Spectrum of Sulfur Monoxide. J. Chem. Phys. 1975, 62, 2511.

(110) Hudgins, D. M.; et al. Mid- and Far-Infrared Spectroscopy of Ices: Optical Constants and Integrated Absorbances. *Astrophysical Journal Supplement Series* **1993**, *86*, 713.

(111) Giguère, P. A.; Harvey, K. B. An Infrared Study of Hydrogen Bonding in Solid  $H_2O_2$  and  $H_2O:H_2O_2$  Mixtures. J. Mol. Spectrosc. **1959**, 3, 36.

(112) Moore, M. H.; Hudson, R. L. IR Detection of  $H_2O_2$  at 80 K in Ion-Irradiated Laboratory Ices Relevant to Europa. *Icarus* **2000**, *145*, 282.

(113) Loeffler, M. J.; Hudson, R. L. Descent Without Modification? The Thermal Chemistry of  $H_2O_2$  on Europa and Other Icy Worlds. *Astrobiology* **2015**, *15*, 453.

(114) Mifsud, D. V.; Hailey, P. A.; Herczku, P.; Juhász, Z.; Kovács, S. T. S.; Sulik, B.; Ioppolo, S.; Kaňuchová, Z.; McCullough, R. W.; Paripás, B.; Mason, N. J.; et al. Laboratory Experiments on the Radiation Astrochemistry of Water Ice Phases. *European Physical Journal D* **2022**, *76*, 87.

(115) Baragiola, R. A. Water Ice on Outer Solar System Surfaces: Basic Properties and Radiation Effects. *Planetary and Space Science* **2003**, *51*, 953.

(116) Palumbo, M. E. Formation of Compact Solid Water After Ion Irradiation at 15 K. *Astron. Astrophys.* **2006**, *453*, 903.

(117) Boduch, P.; et al. Radiation Effects in Astrophysical Ices. *Journal of Physics Conference Series* **2015**, *629*, No. 012008.

(118) Kaňuchová, Z.; et al. Thermal and Energetic Processing of Astrophysical Ice Analogs Rich in SO<sub>2</sub>. Astron. Astrophys. **2017**, 604, A68.

(119) Loeffler, M. J.; et al. Radiolysis of Sulfuric Acid, Sulfuric Acid Monohydrate, and Sulfuric Acid Tetrahydrate and its Relevance to Europa. *Icarus* **2011**, *215*, 370.

(120) Loeffler, M. J.; et al. Thermal Regeneration of Sulfuric Acid Hydrates After Irradiation. *Icarus* **2012**, *219*, 561.

(121) Bossa, J. B.; et al. Thermal Collapse of Porous Interstellar Ice. *Astron. Astrophys.* **2012**, *545*, A82.

(122) Isokoski, K.; et al. Porosity and Thermal Collapse Measurements of  $H_2O$ ,  $CH_3OH$ ,  $CO_2$ , and  $H_2O:CO_2$  Ices. *Phys. Chem. Chem. Phys.* **2014**, *16*, 3456.

(123) Della Guardia, R. A.; Johnston, F. J. Radiation-Induced Reaction of Sulfur and Water. *Radiat. Res.* **1980**, *84*, 259.

(124) Góbi, S.; et al. Successive Hydrogenation of SO and SO<sub>2</sub> in Solid *para*-H<sub>2</sub>: Formation of Elusive Small Oxoacids of Sulfur. ACS Earth and Space Chemistry **2021**, 5, 1180.

(125) Ding, J. J.; et al. Implantation of Multiply Charged Sulfur Ions in Water Ice. *Icarus* **2013**, *226*, 860.

(126) Linnartz, H.; et al. Atom Addition Reactions in Interstellar Ice Analogs. *Int. Rev. Phys. Chem.* **2015**, *34*, 205.

(127) Michaelides, A.; Hu, P. Hydrogenation of S to  $H_2S$  on Pt(111): A First-Principles Study. J. Chem. Phys. **2001**, 115, 8570.

(128) Oba, Y.; et al. An Infrared Measurement of Chemical Desorption from Interstellar Ice Analogs. *Nature Astronomy* **2018**, *2*, 228.

(129) Oba, Y.; et al. Physico-Chemical Behavior of Hydrogen Sulfide Induced by Reactions with H and D Atoms on Different Types of Ice Surfaces at Low Temperature. *Astrophysical Journal* **2019**, 874, 124.

(130) Thaddeus, P.; et al. Interstellar Hydrogen Sulfide. Astrophysical Journal 1972, 176, L73.

(131) Ukita, N.; et al. Hydrogen Sulfide in a Circumstellar Envelope. *Astron. Astrophys.* **1983**, *121*, 15.

(132) Minh, Y. C.; et al. Detection of Interstellar Hydrogen Sulfide in Cold, Dark Clouds. *Astrophysical Journal* **1989**, 345, L63.

(133) Santos, J. C.; et al.  $SO_2$  and OCS Toward High-Mass Protostars: A Comparative Study of Ice and Gas. *Astron. Astrophys.* **2024**, 689, A248.

(134) Santos, J. C.; et al. Fromation of Carbonyl Sulfide (OCS) *via* SH Radicals in Interstellar CO-Rich Ice Under Dense Cloud Conditions. *Astron. Astrophys.* **2024**, *690*, A24.

(135) Ioppolo, S.; et al. Laboratory Evidence for Efficient Water Formation in Interstellar Ices. *Astrophysical Journal* **2008**, 686, 1474. (136) Ioppolo, S.; et al. Surface Formation of  $CO_2$  Ice at Low Temperatures. *Mon. Not. R. Astron. Soc.* **2011**, 413, 2281.

(137) Adriaens, D.; et al. Computational Study of Carbonyl Sulfide Formation on Model Interstellar Dust Grains. J. Phys. Chem. C 2010, 114, 1892. (138) Nguyen, T.; et al. Successive H-Atom Addition to Solid OCS on Compact Amorphous Solid Water. *Astrophysical Journal* **2021**, *922*, 146.

(139) Maity, S.; Kaiser, R. I. Electron Irradiation of Carbon Disulfide-Oxygen Ices: Toward the Formation of Sulfur-Bearing Molecules in Interstellar Ices. *Astrophysical Journal* **2013**, *773*, 184. (140) Donaldson, G. W.; Johnston, F. J. Radiolysis of Colloidal

Sulfur. J. Phys. Chem. 1968, 72, 3552.

(141) Johnston, F. J.; Donaldson, G. W. Radiolysis of Colloidal Sulfur. Mechanism for Solubilization. J. Phys. Chem. 1971, 75, 756.

(142) Carlson, R. W.; et al. Sulfuric Acid Production on Europa: The Radiolysis of Sulfur in Water Ice. *Icarus* **2002**, *157*, 456.

(143) Ashkenazy, Y. Heliyon 2019, 5, No. e01908.

(144) Carlson, R. W., Europa's Surface Composition. In: *Europa*, pp 283–328. Eds: Pappalardo, R. T.; McKinnon, W.B.; Khurana, K.. The University of Arizona Press, 2009.

(145) Carlson, R. W.; et al. Sulfuric Acid on Europa and the Radiolytic Sulfur Cycle. *Science* **1999**, *286*, 97.

(146) Carlson, R. W.; et al. Distribution of Hdyrate on Europa: Further Evidence for Sulfuric Acid Hydrate. *Icarus* **2005**, *177*, 461.

(147) Strazzulla, G.; et al. Hydrate Sulfuric Acid After Sulfur Implantation in Water Ice. *Icarus* **200**7, *192*, 623.

(148) Li, J.; Li, C. Sulfur Chemistry on the Surface of Europa. *Icarus* **2023**, *394*, No. 115438.

(149) Cassidy, T.; et al. Radiolysis and Photolysis of Icy Satellite Surfaces: Experiments and Theory. *Space Science Reviews* **2010**, *153*, 299.

(150) Dalton, J. B.; et al. Exogenic Controls on Sulfuric Acid Hydrate Production at the Surface of Europa. *Planetary and Space Science* **2013**, *77*, 45.

(151) Mifsud, D. V.; Kaňuchová, Z.; Herczku, P.; Juhász, Z.; Kovács, S. T. S.; Lakatos, G.; Rahul, K. K.; Rácz, R.; Sulik, B.; Biri, S.; Rajta, I.; Vajda, I.; Ioppolo, S.; McCullough, R. W.; Mason, N. J.; et al. Sulfur Ion Implantations into Condensed CO<sub>2</sub>: Implications for Europa. *Geophys. Res. Lett.* **2022**, 49, No. e2022GL100698.

(152) Mifsud, D. V.; et al. Sulfur Ion Implantation into  $O_2$ , CO, and  $CO_2$  Ices: Implications for the Formation of Sulfur-Bearing Molecules in the Kuiper Belt. *Icarus* **2024**, *411*, No. 115926.

(153) Hendrix, A. R.; et al. Europa's Disk-Resolved Ultraviolet Spectra: Relationships with Plasma Flux and Surface Terrains. *Icarus* **2011**, *212*, 736.

(154) Becker, T. M.; et al. Mid-Ultraviolet Hubble Observations of Europa and the Global Surface Distribution of  $SO_2$ . Planetary Science Journal **2022**, 3, 129.

(155) Lv, X. Y.; et al. Sulfur Implantation in CO and  $CO_2$  Ices. Mon. Not. R. Astron. Soc. **2014**, 438, 922.

(156) Lv, X. Y.; et al. Thermal and Energetic Processing of Ammonia and Carbon Dioxide Bearing Solid Mixtures. *Phys. Chem. Chem. Phys.* 2014, 16, 3433.

(157) Boduch, P.; et al. Ion Processing of Ices and the Origin of  $SO_2$  and  $O_3$  on the Icy Surfaces of the Icy Jovian Satellites. *Icarus* **2016**, 277, 424.

(158) Anders, C.; Urbassek, H. M. Energetic Sulfur Ion Impacts into Cometary Ice Surfaces: A Molecular Dynamics Study. *Mon. Not. R. Astron. Soc.* **2019**, 482, 2374.

(159) Anders, C.; Urbassek, H. M. High-Energy Ion Impacts into the Sulfur-Bearing Ice Surface of Europa: An Atomistic Study of Chemical Transformations. *Astron. Astrophys.* **2019**, *625*, A140.

(160) Ruf, A.; et al. Organosulfur Compounds Formed by Sulfur Ion Bombardment of Astrophysical Ice Analogs: Implications for Moons, Comets, and Kuiper Belt Objects. *Astrophysical Journal Letters* **2019**, 885, L40.

(161) Ruf, A.; et al. Sulfur Ion Irradiation Experiments Simulating Space Weathering of Solar System Body Surfaces: Organosulfur Compound Formation. *Astron. Astrophys.* **2021**, *655*, A74.

(162) Bouquet, A.; et al. Sulfur Implantation into Water Ice with Propane: Implications for Organic Chemistry on the Surface of Europa. *Planetary Science Journal* **2024**, *5*, 102.