



On the Futaki invariant of Fano threefolds

Downloaded from: <https://research.chalmers.se>, 2026-05-15 20:50 UTC

Citation for the original published paper (version of record):

Sektnan, L., Tipler, C. (2024). On the Futaki invariant of Fano threefolds. *Annali dell'Universita di Ferrara*, 70(3): 811-837. <http://dx.doi.org/10.1007/s11565-024-00503-x>

N.B. When citing this work, cite the original published paper.



On the Futaki invariant of Fano threefolds

Lars Martin Sektnan¹ · Carl Tipler²

Received: 29 August 2023 / Accepted: 4 February 2024 / Published online: 24 February 2024
© The Author(s) 2024

Abstract

We study the zero locus of the Futaki invariant on K -polystable Fano threefolds, seen as a map from the Kähler cone to the dual of the Lie algebra of the reduced automorphism group. We show that, apart from families 3.9, 3.13, 3.19, 3.20, 4.2, 4.4, 4.7 and 5.3 of the Iskovskikh–Mori–Mukai classification of Fano threefolds, the Futaki invariant of such manifolds vanishes identically on their Kähler cone. In all cases, when the Picard rank is greater or equal to two, we exhibit explicit 2-dimensional differentiable families of Kähler classes containing the anti-canonical class and on which the Futaki invariant is identically zero. As a corollary, we deduce the existence of non Kähler–Einstein cscK metrics on all such Fano threefolds.

Keywords Futaki invariant · Fano 3-folds · Constant scalar curvature Kähler metrics

Mathematics Subject Classification 32Q20 · 14J30 · 14J45 · 14J50

1 Introduction

The Futaki invariant was introduced by Futaki [9, 10] as an obstruction to the existence of Kähler–Einstein metrics on Fano manifolds. Its definition extends to any compact polarised Kähler manifold, and its vanishing is a necessary condition for the existence of a constant scalar curvature Kähler metric (cscK for short) in a given Kähler class.

In this note, we study the zero locus of the Futaki invariant, seen as a map from the Kähler cone to the dual of the Lie algebra of the reduced automorphism group (see

Lars Martin Sektnan and Carl Tipler have contributed equally to this work.

✉ Lars Martin Sektnan
sektnan@chalmers.se

Carl Tipler
Carl.Tipler@univ-brest.fr

¹ Department of Mathematical Sciences, University of Gothenburg, 412 96 Gothenburg, Sweden

² LMBA, UMR CNRS 6205, Université de Bretagne Occidentale, 6, avenue Victor Le Gorgeu, 29238 Brest, France

Sect. 2 for the definitions). This locus is fully understood for Fano surfaces from the works [21, 23, 24], which we recall in Sect. 2.1. Here we will focus on K -polystable Fano threefolds. The description of this class of manifolds has seen recently great progress, in particular with [1] (see also references therein).

Relying on a case by case analysis, our little contribution to “Fanography” is the following:

Theorem 1 *Let $(X, -K_X)$ be a K -polystable Fano threefold that belongs to family $N^\circ\mathcal{N}$, with*

$$\mathcal{N} \notin \{3.9, 3.13, 3.19, 3.20, 4.2, 4.4, 4.7, 5.3\}.$$

Then, the Futaki invariant of X vanishes identically on its Kähler cone.

The numbering in the above is the same as the numbering of the families of the Iskovskikh–Mori–Mukai classification of Fano threefolds given in [1].

Note that when $\text{Aut}(X)$ is finite or when the Picard rank $\rho(X) = 1$, the Futaki invariant vanishes identically on the Kähler cone, as soon as X is K -polystable in the second case. From the classification in [1], there exists 33 families of Fano threefolds with $\rho(X) \geq 2$ that admit members which are K -polystable with respect to the anti-canonical polarisation and which have infinite automorphism group. We verify that of these, all but 8 families have vanishing Futaki invariant for every K -polystable member. Further, for every K -polystable member of the remaining 8 families, we provide explicit 2-dimensional families of Kähler classes that contain $c_1(X)$ and on which the Futaki invariant vanishes.

Theorem 2 *Let $(X, -K_X)$ be a K -polystable Fano threefold that belongs to family $N^\circ\mathcal{N}$, with*

$$\mathcal{N} \in \{3.9, 3.13, 3.19, 3.20, 4.2, 4.4, 4.7, 5.3\}.$$

Then, there is at least a 2-dimensional family of Kähler classes on X , containing $c_1(X)$, where the Futaki invariant vanishes.

Note that this is a 2-dimensional family in the actual Kähler cone of X . The Futaki invariant of a vector field ν with respect to a class Ω rescales as the scaling factor to the dimension of the manifold. Thus one always knows there is at least a 1-dimensional family of Kähler classes near a given class with vanishing Futaki invariant, on which the Futaki invariant vanishes. For the smooth Fano varieties considered in the above, we show that the actual locus around the anti-canonical class where the Futaki invariant vanishes is at least one dimension higher than this.

From the LeBrun–Simanca openness theorem ([13, Section 2.3]), we deduce the following corollary.

Corollary 3 *Let X be a K -polystable Fano threefold with Picard rank $\rho(X) \geq 2$. Then X admits a 2-dimensional family of cscK metrics parametrised by a 2-dimensional family of Kähler classes containing $c_1(X)$.*

The above uses LeBrun–Simanca's openness of the extremal cone, together with the fact that an extremal metric is cscK precisely when the classical Futaki invariant vanishes. From the algebro-geometric perspective, assuming the Yau–Tian–Donaldson conjecture to be true in full generality [7, 22, 26], this means that for the members of the families considered in Theorem 1 with an arbitrary polarisation, and for the polarisations considered in Theorem 2, relative K -stability is equivalent to K -polystability (see e.g. [18] for the relative version of K -stability and its relation to extremal Kähler metrics).

Remark 1 For K -polystable members of the families $\mathcal{N} \in \{4.2, 4.4, 4.7\}$ with infinite automorphism group, we actually show that there is a 3-dimensional family of Kähler classes near $-K_X$ with vanishing Futaki invariant.

Remark 2 It would be interesting to find an example of Kähler–Einstein Fano manifold with Picard rank greater than two but with *no* non Kähler–Einstein cscK metric in Kähler classes near the anticanonical one.

Remark 3 Our results should be compared with the recent [16], where the Futaki invariant of Bott manifolds is studied. In contrast to our results, which guarantee the vanishing of the Futaki invariant in many cases, it is shown in [16] that the only Bott manifolds for which the Futaki invariant vanishes on the whole Kähler cone are isomorphic to products of projective lines. The key observation to prove our results is the existence of enough discrete symmetries that preserve every Kähler class on Fano threefolds, which in the majority of the cases considered here will be responsible for the vanishing of the Futaki invariant.

Notations and conventions

Throughout the paper, for a compact Kähler manifold X , we will denote by $\text{Aut}(X)$ (respectively $\text{Aut}_0(X)$) its automorphism group (respectively the connected component of the identity of the reduced automorphism group of X), and by $\mathfrak{aut}(X)$ the Lie algebra of $\text{Aut}(X)$. If $Z \subset X$ is a subvariety (not necessarily connected), $\text{Aut}(X, Z)$ stands for elements in $\text{Aut}(X)$ that leave Z globally invariant. We denote by \mathcal{K}_X the Kähler cone of X . We will identify a divisor D with $\mathcal{O}(D)$, and use the notation $c_1(D)$ for its first Chern class.

2 Preliminaries

Let X be a compact Kähler manifold, and $\Omega \in \mathcal{K}_X$ a Kähler class on X . We denote the Futaki invariant of (X, Ω) by

$$\begin{aligned} \text{Fut}_{(X, \Omega)} : \mathfrak{aut}_0(X) &\rightarrow \mathbb{C} \\ v &\mapsto -\int_X f_{v, g} \text{scal}_g d\mu_g, \end{aligned}$$

where $\mathfrak{aut}_0(X)$ is the Lie algebra of the reduced automorphism group of X , g denotes a Kähler metric with Kähler form in Ω and volume form $d\mu_g$, $f_{v, g}$ is the normalised

holomorphy potential of v with respect to g , and scal_g denotes the scalar curvature of g (see e.g. [4], [14, Section 3.1] or [11, Chapter 4] for this formulation of the Futaki invariant, initially introduced in [9]).

We will be interested in K -polystable Fano manifolds, or equivalently Fano manifolds admitting a Kähler–Einstein metric of positive curvature by the resolution of the Yau–Tian–Donaldson conjecture [6, 7, 22, 26]. For such manifolds, by Matsushima’s result [15], and from Bochner’s formula (see [11, Section 3.6]), we have $\text{aut}_0(X) = \text{aut}(X)$. We will therefore consider the Futaki invariant as a map

$$\text{Fut}_X : \mathcal{K}_X \rightarrow \text{aut}(X)^*.$$

By construction, Fut_X vanishes on any class that admits a cscK metric, and it is then straightforward that $\text{Fut}_X \equiv 0$ whenever X is a K -polystable Fano manifold with Picard rank 1, or when the automorphism group of X is finite.

2.1 The case of smooth Del Pezzo surfaces

We refer here the reader to [17, Section 2] and [1, Section 2]. If X is a smooth Del Pezzo surface with infinite automorphism group, then $K_X^2 \in \{6, 7, 8, 9\}$. Moreover, it is K -polystable and of Picard rank $\rho(X) \geq 2$ if and only if $X = \mathbb{P}^1 \times \mathbb{P}^1$ or $K_X^2 = 6$, i.e. when X is a blow-up of \mathbb{P}^2 along three non-collinear points [21, 23]. In the first case, X admits a product cscK metric in each class, and $\text{Fut}_X \equiv 0$, while in the latter case, the vanishing locus of Fut_X is described in [24, Section 5] (see Sect. 4.2 for the exact description).

2.2 Further properties of the Futaki invariant

The key property that we will use is the invariance of Fut_X under the $\text{Aut}(X)$ -action. This was already used in [10, Chapter 3] to show the vanishing of Fut_X on specific examples.

We will use the following proposition repeatedly.

Proposition 4 *Let (X, Ω) be a polarised Fano manifold. Assume that there is $\tau \in \text{Aut}(X)$ and $v \in \text{aut}(X)$ such that*

- (i) $\tau^*\Omega = \Omega$,
- (ii) *there is $c \in \mathbb{C}^* \setminus \{1\}$ with $\text{Ad}_\tau(v) = c \cdot v$.*

Then $\text{Fut}_{(X, \Omega)}(v) = 0$.

Proof This follows from the Ad-invariance of the Futaki invariant, which implies that

$$\text{Fut}_{(X, (\tau^{-1})^*\Omega)}(\text{Ad}_\tau(v)) = \text{Fut}_{(X, \Omega)}(v),$$

see [10, Chapter 3] or [14, Section 3.1]. □

Remark 4 The anti-canonical class $c_1(X)$ is always $\text{Aut}(X)$ -invariant.

As an application, we have the following useful corollary:

Corollary 5 *Let $\pi : X \rightarrow Y$ be the blow-up of a smooth Fano manifold Y along smooth and disjoint subvarieties $Z_i \subset Y$. Assume that there is a finite group $G \subset \text{Aut}(Y)$ such that:*

- (i) *Each Z_i is G -invariant;*
- (ii) *Each class $\Omega \in H^{1,1}(Y, \mathbb{R})$ is G -invariant;*
- (iii) *For any $v \in \text{aut}(Y)$ that lifts to X , there is $\tau \in G$ and $c \in \mathbb{C}^* \setminus \{1\}$ such that $\text{Ad}_\tau(v) = c \cdot v$.*

Then $\text{Fut}_X \equiv 0$.

Proof From hypothesis (i), the G -action on Y lifts to a G -action on X . The vector space $H^{1,1}(X, \mathbb{R})$ is spanned by the pullback of the classes in $H^{1,1}(Y, \mathbb{R})$ and the exceptional divisors of π . By hypothesis (i) and (ii), any class in $H^{1,1}(X, \mathbb{R})$ is then G -invariant. The Lie algebra $\text{aut}(X)$ is spanned by lifts of elements in $\text{aut}(Y)$ that preserve the Z_i 's. For any such element, the identity $\text{Ad}_\tau(v) = c \cdot v$ holds on $X \setminus \bigcup_i \pi^{-1}(Z_i)$, hence on X , by continuity. The result follows from Proposition 4. □

Remark 5 In practice, we will mainly use Corollary 5 with

$$G \simeq \mathbb{Z}/2\mathbb{Z}, \text{ aut}(X) \simeq \mathbb{C}, c = -1.$$

To prove item (i) of Proposition 4 or item (ii) of Corollary 5, we will use the fact that in homogeneous coordinates, the Fubini–Study metric

$$\omega_{FS} = \frac{i}{2} \partial \bar{\partial} \log(|z|^2),$$

and hence its class $[\omega_{FS}] \in H^{1,1}(\mathbb{P}^n, \mathbb{R})$, is invariant under the \mathfrak{S}_{n+1} -action on \mathbb{P}^n by permutation of the homogeneous coordinates.

2.3 The list to check

From the discussion in the beginning of this section, to prove Theorem 1, it is enough to consider K -polystable Fano threefolds with infinite automorphism group and Picard rank $\rho(X) \geq 2$. From [1, Section 6], this reduces to Fano threefolds in family $N^\circ \mathcal{N}$, for

$$\mathcal{N} \in \left\{ \begin{array}{l} 2.20, 2.21, 2.22, 2.24, 2.27, 2.29, 2.32, 2.34, 3.5, 3.8, 3.9, \\ 3.10, 3.12, 3.13, 3.15, 3.17, 3.19, 3.20, 3.25, 3.27, 4.2, 4.3, \\ 4.4, 4.6, 4.7, 4.13, 5.1, 5.3, 6.1, 7.1, 8.1, 9.1, 10.1 \end{array} \right\}.$$

The strategy of the proof is then direct – we will use the invariance of Fut_X to show its vanishing on \mathcal{K}_X using a case by case study. For X belonging to family $N^\circ \mathcal{N}$ with

$$\mathcal{N} \in \left\{ \begin{array}{l} 2.20, 2.21, 2.22, 2.24, 2.27, 2.29, 2.32, 2.34, \\ 3.5, 3.8, 3.10, 3.12, 3.15, 3.17, 3.25, 3.27, \\ 4.3, 4.6, 4.13, 5.1, 6.1, 7.1, 8.1, 9.1, 10.1 \end{array} \right\},$$

we will see that $\text{Fut}_X \equiv 0$.

For the remaining 8 families, we will obtain explicit Kähler classes of the form

$$c_1(X) + \varepsilon c_1(D) \in \mathcal{K}_X$$

with $D \subset X$ a divisor and $\varepsilon \in \mathbb{R}$ a parameter such that $\text{Fut}_{(X, c_1(X) + \varepsilon c_1(D))} = 0$. As the vanishing of the Futaki invariant is preserved under scaling of the Kähler metric, this provides the 2-dimensional families of Kähler classes with vanishing Futaki invariant alluded to in the introduction. Corollary 3 then follows from LeBrun–Simanca's openness theorem [13], which asserts that the locus in the Kähler cone of Kähler classes that admit an extremal metric in the sense of Calabi [3] is open, together with the characterisation of cscK metrics amongst extremal metrics as the ones with zero Futaki invariant [4].

3 Families with $\text{aut}(X) \simeq \mathfrak{sl}_n(\mathbb{C})$

Here we will consider families $N^\circ \mathcal{N}$, with

$$\mathcal{N} \in \{2.27, 2.32, 3.17, 4.6, 6.1, 7.1, 8.1, 9.1, 10.1\}.$$

The Lie algebra $\mathfrak{sl}_n(\mathbb{C})$ is simple, hence equal to its derived ideal $[\mathfrak{sl}_n(\mathbb{C}), \mathfrak{sl}_n(\mathbb{C})]$. As the Futaki invariant is a character from $\text{aut}(X)$ to \mathbb{C} (see e.g. [10, Chapter 3] or [14, Section 3.1]), it vanishes on the derived ideal $[\text{aut}(X), \text{aut}(X)]$. Hence, if $\text{aut}(X) \simeq \mathfrak{sl}_n(\mathbb{C})$, $[\text{aut}(X), \text{aut}(X)] = \text{aut}(X)$ and the Futaki invariant vanishes identically on the whole Kähler cone of X . From

$$\text{PGL}_n(\mathbb{C}) \simeq \text{SL}_n(\mathbb{C})/\mu_n,$$

the Lie algebra of $\text{PGL}_n(\mathbb{C})$ is $\mathfrak{sl}_n(\mathbb{C})$. From [1, Section 6, Big Table], this settles the case of all the K -polystable Fano threefolds in families $N^\circ \mathcal{N}$, with

$$\mathcal{N} \in \{2.27, 2.32, 3.17, 4.6, 6.1, 7.1, 8.1, 9.1, 10.1\},$$

and also some cases in families $\{2.21, 3.13\}$.

4 Products

Next, we consider families $N^\circ 2.34$, $N^\circ 3.27$ and $N^\circ 5.3$, which are products of lower dimensional Fano manifolds.

4.1 Families 2.34 and 3.27

The unique members in these two families are $\mathbb{P}^1 \times \mathbb{P}^2$ and $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$, which both carry a product of cscK metrics in any class, and thus has vanishing Futaki character for any Kähler class.

4.2 Family 5.3

The unique Fano threefold in family 5.3 is $\mathbb{P}^1 \times S_6$, where S_6 is the Del Pezzo surface with $K_{S_6}^2 = 6$. It is K -polystable as a product of Kähler–Einstein manifolds from [21, 23]. The surface S_6 is the unique (up to isomorphism) toric surface obtained by blowing-up \mathbb{P}^2 in the three fixed points under the torus action. We denote by H (the strict transform of) a generic hyperplane and D_1, D_2 and D_3 the three exceptional divisors in S_6 . From [24, Section 5, Proposition 5.2 and Remark 5.1.(iii)], the Futaki invariant of S_6 vanishes exactly in the following families of Kähler classes

$$3c_1(H) - ac_1(D_1) - bc_1(D_2) - (3 - a - b)c_1(D_3)$$

and

$$3c_1(H) - c(c_1(D_1) + c_1(D_2) + c_1(D_3)),$$

where a, b, c are positive constants satisfying $a + b < 3$ and $c < \frac{3}{2}$. As the Futaki invariant vanishes on \mathbb{P}^1 , we easily deduce the vanishing locus of the Futaki invariant on $X = \mathbb{P}^1 \times S_6$. In particular, as $c_1(X) = c_1(\mathbb{P}^1) + c_1(S_6)$, and as $c_1(S_6) = 3c_1(H) - (c_1(D_1) + c_1(D_2) + c_1(D_3))$, we deduce the existence of differentiable families of Kähler classes on X containing $c_1(X)$ for which the Futaki invariant vanishes identically.

5 Blow-ups of projective space

In this section we address families $N^\circ \mathcal{N}$, with

$$\mathcal{N} \in \{2.22, 3.12, 3.25\}.$$

All the members of these families are obtained by blowing up certain curves in projective space \mathbb{P}^3 .

5.1 Family 2.22

Members of the family 2.22 of Fano threefolds are obtained as blowups of certain curves in \mathbb{P}^3 . More precisely, let $\Phi : \mathbb{P}^1 \times \mathbb{P}^1 \rightarrow \mathbb{P}^3$ be the Segre embedding

$$([x : y], [u : v]) \mapsto [xu : xv : yu : yv].$$

The image of Φ is the surface

$$S = \{z_0 z_3 - z_1 z_2 = 0\}.$$

A Fano threefold X is in the family 2.22 if it is the blowup of the image via Φ of a curve \check{C} with $\mathcal{O}(\check{C}) = \mathcal{O}(3, 1)$. Such X have Picard rank 2, generated by the line bundle associated to the proper transform of a hyperplane and of that generated by the exceptional divisor E of the blowup. The K -polystability of members of this family (with respect to the anticanonical polarisation) is discussed in detail in [5].

Up to biholomorphism, there is a unique member X_0 of this family with infinite automorphism group. It is K -polystable, and can be obtained by picking the curve \check{C} to be $\check{C}_0 = \{ux^3 - vy^3 = 0\}$, so that

$$X_0 = \text{Bl}_{C_0} \mathbb{P}^3,$$

where $C_0 = \Phi(\check{C}_0)$. The \mathbb{C}^* -action

$$\lambda \cdot ([z_0 : z_1 : z_2 : z_3]) = [\lambda z_0 : \lambda^4 z_1 : z_2 : \lambda^3 z_3]$$

preserves C_0 and so lifts to X_0 . This generates $\text{Aut}_0(X_0)$ (see [17, Lemma 6.13]).

The curve C_0 is a rational curve, which can e.g. be seen by applying the Riemann–Hurwitz formula to the restriction to $\check{C}_0 \subset \mathbb{P}^1 \times \mathbb{P}^1$ of the projection to the second factor. An explicit parametrisation $\phi : \mathbb{P}^1 \rightarrow \mathbb{P}^3$ is given by

$$[\tau_0 : \tau_1] \mapsto [\tau_0 \tau_1^3 : \tau_0^4 : \tau_1^4 : \tau_1 \tau_0^3].$$

Note that the action of the involution τ given by

$$\tau \cdot ([z_0 : z_1 : z_2 : z_3]) = [z_3 : z_2 : z_1 : z_0]$$

on \mathbb{P}^3 preserves C_0 and so lifts to X_0 . We will then apply Corollary 5 to the blow-up $X_0 \rightarrow \mathbb{P}^3$ with the group $G = \langle \tau \rangle \simeq \mathbb{Z}/2\mathbb{Z}$. First, hypothesis (i) of Corollary 5 can be checked by using the change of variables $[\tau_0 : \tau_1] \rightarrow [\tau_1 : \tau_0]$ for the parametrisation of the curve

$$\begin{aligned} \tau(\phi([\tau_0 : \tau_1])) &= \tau[\tau_0 \tau_1^3 : \tau_0^4 : \tau_1^4 : \tau_1 \tau_0^3] \\ &= [\tau_1 \tau_0^3 : \tau_1^4 : \tau_0^4 : \tau_0 \tau_1^3] \\ &= \phi([\tau_1 : \tau_0]), \end{aligned}$$

so C_0 is invariant under the action. Then, as the variety we are blowing up is \mathbb{P}^3 , hypothesis (ii) comes from the invariance of the Fubini–Study metric, cf Remark 5. Finally, a direct computation gives $\tau \lambda \tau^{-1} = \lambda^{-1}$, so that $\text{Ad}_\tau(v) = -v$ for v a generator of the λ -action. Hence we can apply Corollary 5, which implies the vanishing of the Futaki invariant on the Kähler cone of X_0 .

5.2 Family 3.12

From [1, Section 5.18], the only element in Family 3.12 with infinite automorphism group is given, up to isomorphism, by $X = \text{Bl}_{LUC}(\mathbb{P}^3)$ the blow up of \mathbb{P}^3 along the disjoint curves

$$L = \{x_0 = x_3 = 0\} \subset \mathbb{P}^3$$

and

$$C = \{[s^3 : s^2t : st^2 : t^3], [s : t] \in \mathbb{P}^1\} \subset \mathbb{P}^3.$$

The reduced automorphism group of X is isomorphic to \mathbb{C}^* , and its action is given by the lift of the \mathbb{C}^* -action on \mathbb{P}^3 described by

$$\lambda \cdot ([x_0 : x_1 : x_2 : x_3]) = [x_0 : \lambda x_1 : \lambda^2 x_2 : \lambda^3 x_3].$$

Then, we can consider the $\mathbb{Z}/2\mathbb{Z}$ -action given by

$$\tau([x_0 : x_1 : x_2 : x_3]) = [x_3 : x_2 : x_1 : x_0].$$

The group generated by τ in $\text{Aut}(\mathbb{P}^3)$ satisfies hypothesis (i) – (iii) from Corollary 5, and we deduce that the Futaki invariant of X vanishes on the whole Kähler cone.

5.3 Family 3.25

The Fano threefold X in family 3.25 is the blow-up of \mathbb{P}^3 in two disjoint lines. It is K -polystable from [2, 25]. We can assume the two blown-up lines are $\{x_1 = x_2 = 0\} \subset \mathbb{P}^3$ and $\{x_3 = x_4 = 0\} \subset \mathbb{P}^3$. One has

$$\text{Aut}_0(X) \simeq \text{PGL}_{(2,2)}(\mathbb{C}) \simeq \text{GL}_2(\mathbb{C}) \times \text{GL}_2(\mathbb{C})/\mathbb{C}^*,$$

where the first (resp. second) $\text{GL}_2(\mathbb{C})$ factor acts linearly on the coordinates (x_1, x_2) (resp. on (x_3, x_4)) while the \mathbb{C}^* -action corresponds to homotheties on \mathbb{C}^4 (see [17, Section 4]). The Lie algebra $\text{aut}(X)$ of $\text{Aut}_0(X)$ fits in an exact sequence

$$0 \rightarrow \mathbb{C} \rightarrow \mathfrak{gl}_2(\mathbb{C}) \oplus \mathfrak{gl}_2(\mathbb{C}) \rightarrow \text{aut}(X) \rightarrow 0.$$

We also have the sequence induced by the trace map $\mathfrak{gl}_2(\mathbb{C}) \rightarrow \mathbb{C}$:

$$0 \rightarrow \mathfrak{sl}_2(\mathbb{C}) \rightarrow \mathfrak{gl}_2(\mathbb{C}) \rightarrow \mathbb{C} \rightarrow 0,$$

from which we deduce the sequence of vector spaces

$$0 \rightarrow \mathbb{C} \rightarrow (\mathfrak{sl}_2(\mathbb{C}) \oplus \mathbb{C}) \oplus (\mathfrak{sl}_2(\mathbb{C}) \oplus \mathbb{C}) \rightarrow \text{aut}(X) \rightarrow 0.$$

From the discussion in Sect. 3, the Futaki invariant of X will vanish on the $\mathfrak{sl}_2(\mathbb{C})$ -factors that project to $\text{aut}(X)$. Hence, it is enough to test the vanishing of the Futaki invariant on the generators of the remaining two \mathbb{C}^* -actions modulo homotheties, which are induced by:

$$(\lambda, \mu) \cdot ([x_0 : x_1 : x_2 : x_3]) = [\lambda x_0 : x_1 : \mu x_2 : x_3],$$

where $(\lambda, \mu) \in (\mathbb{C}^*)^2$. We can consider the finite group G generated by the reflections

$$\tau([x_0 : x_1 : x_2 : x_3]) = [x_1 : x_0 : x_2 : x_3]$$

and

$$\sigma([x_0 : x_1 : x_2 : x_3]) = [x_0 : x_1 : x_3 : x_2].$$

This group preserves the two blown-up lines, while the adjoint action of τ (resp. σ) sends the generator of the λ -action (resp. the μ -action) to its inverse. Hence, we conclude as before by using Corollary 5.

6 Blow-ups of products of projective spaces

In this section, we will consider families $N^\circ \mathcal{N}$, with

$$\mathcal{N} \in \{3.5, 4.3, 4.13\}.$$

These are obtained as blowups of products of projective spaces.

6.1 Family 3.5

From [1, Section 5.14], the only element in Family 3.5 with infinite automorphism group is given, up to isomorphism, by $X = \text{Bl}_C(\mathbb{P}^1 \times \mathbb{P}^2)$ the blow up of $\mathbb{P}^1 \times \mathbb{P}^2$ along the curve $C = \psi(\check{C})$ given by the image of

$$\check{C} = \{ux^5 + vy^5 = 0\} \subset \mathbb{P}^1 \times \mathbb{P}^1$$

via the map

$$\begin{aligned} \psi : \quad \mathbb{P}^1 \times \mathbb{P}^1 &\rightarrow \mathbb{P}^1 \times \mathbb{P}^2 \\ ([u : v], [x : y]) &\mapsto ([u : v], [x^2 : xy : y^2]). \end{aligned}$$

Then, $\text{Aut}_0(X) \simeq \mathbb{C}^*$, where the \mathbb{C}^* -action is generated by the lift to X of the action

$$\lambda \cdot ([u : v], [x_0 : x_1 : x_2]) = ([\lambda^5 u : v], [x_0 : \lambda x_1 : \lambda^2 x_2]).$$

We also have a $\mathbb{Z}/2\mathbb{Z}$ -action induced by

$$\tau([u : v], [x_0 : x_1 : x_2]) = ([v : u], [x_2 : x_1 : x_0]).$$

Those actions come respectively from the actions

$$\lambda \cdot ([u : v], [x : y]) = ([\lambda^5 u : v], [x : \lambda y])$$

and

$$\tau([u : v], [x : y]) = ([v : u], [y : x])$$

on $\mathbb{P}^1 \times \mathbb{P}^1$, with respect to which ψ is equivariant. Then, we see that C is τ -invariant, as well as the classes $\pi_i^*[\omega_{FS}^i]$, where $\pi_1 : \mathbb{P}^1 \times \mathbb{P}^2 \rightarrow \mathbb{P}^1$ and $\pi_2 : \mathbb{P}^1 \times \mathbb{P}^2 \rightarrow \mathbb{P}^2$ denote the projections and ω_{FS}^i stands for the Fubini–Study metric on \mathbb{P}^i . Finally, identifying $\lambda \in \mathbb{C}^*$ with its action, we have $\tau \circ \lambda \circ \tau^{-1} = \lambda^{-1}$. Hence, hypothesis (i) – (iii) from Corollary 5 are satisfied, and the Futaki invariant of X vanishes for any Kähler class.

6.2 Family 4.3

Following [1, Section 5.21], up to isomorphism, the unique Fano threefold in Family 4.3 is the blow-up of $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ along

$$C = \{x_0y_1 - x_1y_0 = x_0z_1^2 + x_1z_0^2 = 0\}$$

where $[x_0 : x_1]$, $[y_0 : y_1]$ and $[z_0 : z_1]$ denote the homogeneous coordinates on the first, second and last factor respectively. We have $\text{Aut}_0(X) \simeq \mathbb{C}^*$ where the action is given by the lift of the \mathbb{C}^* -action on $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ given by

$$\lambda \cdot ([x_0 : x_1], [y_0 : y_1], [z_0 : z_1]) = ([x_0 : \lambda^2x_1], [y_0 : \lambda^2y_1], [z_0 : \lambda z_1]).$$

The involution

$$\tau([x_0 : x_1], [y_0 : y_1], [z_0 : z_1]) = ([x_1 : x_0], [y_1 : y_0], [z_1 : z_0])$$

preserves C and the $(1, 1)$ -classes on C given by $\iota_j^*[\omega_{FS}]$, for ι_j the composition of the inclusion $C \subset \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ and the projection on the j -th factor. The adjoint action of τ maps the generator of the \mathbb{C}^* -action to its inverse, so Proposition 4 applies and the Futaki invariant of X vanishes identically.

6.3 Family 4.13

From [1, Section 5.22], the only element in Family 4.13 with infinite automorphism group is given, up to isomorphism, by $X = \text{Bl}_C(\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1)$ the blow up of

$\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ along the curve

$$C = \{x_0y_1 - x_1y_0 = x_0^3z_0 + x_1^3z_1 = 0\}.$$

The reduced automorphism group of X is isomorphic to \mathbb{C}^* , and its action is given by the lift of the \mathbb{C}^* -action on $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ described by

$$\lambda \cdot ([x_0 : x_1], [y_0 : y_1], [z_0 : z_1]) = ([\lambda x_0 : x_1], [\lambda y_0 : y_1], [\lambda^{-3} z_0 : z_1]).$$

Then, we can consider the $\mathbb{Z}/2\mathbb{Z}$ -action given by

$$\tau([x_0 : x_1], [y_0 : y_1], [z_0 : z_1]) = ([x_1 : x_0], [y_1 : y_0], [z_1 : z_0]).$$

Clearly, this action satisfies hypothesis (i) – (iii) from Corollary 5 (notice that $\tau \circ \lambda \circ \tau^{-1} = \lambda^{-1}$, identifying λ with the induced action), from which we deduce the vanishing of the Futaki invariant of X for any Kähler class.

7 Blow-ups of a smooth quadric

In this section, we consider families $N^\circ \mathcal{N}$, with

$$\mathcal{N} \in \{2.21, 2.29, 3.10, 3.15, 3.19, 3.20, 4.4, 5.1\}.$$

7.1 Family 2.21

This family is somewhat similar to the Mukai–Umemura family 1.10. In addition to members of the family with discrete automorphism group, there is a one-dimensional family with automorphism group containing a semi-direct product of \mathbb{C}^* and $\mathbb{Z}/2\mathbb{Z}$, one member which admits an effective PGL_2 -action and one member which has a reduced automorphism group \mathbb{G}_a . The first two of these are K -polystable for the anti-canonical polarisation, whereas the last does not have a reductive automorphism group and is therefore not K -polystable.

The members that admit an effective \mathbb{G}_m -action can be described as follows (see [1, Section 5.9]). Let C be the quartic rational curve in \mathbb{P}^4 given as the image of the map $\mathbb{P}^1 \rightarrow \mathbb{P}^4$ given by

$$[p : q] \mapsto [p^4 : p^3q : p^2q^2 : pq^3 : q^4].$$

For $t \notin \{0, \pm 1\}$, let Q_t be the smooth hypersurface

$$Q_t = V(z_1z_3 - t^2z_0z_4 + (t^2 - 1)z_2^2).$$

Note that $C \subset Q_t$ for any t . Let $X_t = \mathrm{Bl}_C(Q_t)$. Then X_t is one of the members that admit an effective \mathbb{C}^* -action (including the member with an effective PGL_2 -

action, which corresponds to $t = \pm \frac{1}{2}$). Note that X_t has Picard rank 2, generated by a hyperplane H and the exceptional divisor E of the blowup.

The \mathbb{C}^* -action given by

$$\lambda \cdot ([z_0 : z_1 : z_2 : z_3 : z_4]) = [z_0 : \lambda z_1 : \lambda^2 z_2 : \lambda^3 z_3 : \lambda^4 z_4]$$

preserves C and Q_t , as does the involution

$$\tau([z_0 : z_1 : z_2 : z_3 : z_4]) = [z_4 : z_3 : z_2 : z_1 : z_0].$$

The lifts of these generate the effective actions of $\mathbb{C}^* \rtimes \mathbb{Z}/2\mathbb{Z}$ on X_t . As τ preserves C , the class $[\omega_{FS}]|_{X_t}$, and sends a generator of the \mathbb{C}^* -action to its inverse by conjugation, Proposition 4 shows that the Futaki invariant of X_t vanishes on its whole Kähler cone (note that the case $t = \pm \frac{1}{2}$, with $\text{aut}(X_t) \simeq \mathfrak{sl}_2(\mathbb{C})$, was dealt with in Sect. 3).

7.2 Family 2.29

There is a unique smooth Fano threefold X in family 2.29. It is isomorphic to the blow-up of

$$Q = \{x_0^2 + x_1x_2 + x_3x_4 = 0\} \subset \mathbb{P}^4.$$

along the smooth conic

$$C = \{x_0^2 + x_1x_2 = x_3 = x_4 = 0\} \subset Q.$$

It is K -polystable (see [12, 19, 20]) and the group $\text{Aut}_0(X)$ is isomorphic to $\mathbb{C}^* \times \text{PGL}_2(\mathbb{C})$ (see [17, Lemma 5.8]). We then have $\text{aut}(X) \simeq \mathbb{C} \oplus \mathfrak{sl}_2(\mathbb{C})$. From the discussion in Sect. 3, the Futaki invariant of $(X, [\omega])$ vanishes on the $\mathfrak{sl}_2(\mathbb{C})$ -component of $\text{aut}(X)$ for any Kähler class $[\omega]$. Thus, to check the vanishing of the Futaki invariant, it remains to check the vanishing on the \mathbb{C} -component of $\text{aut}(X)$. From [17, Lemma 5.7], the \mathbb{C}^* -component of $\text{Aut}_0(X)$ can be identified with the pointwise stabiliser of C in $\text{Aut}_0(Q)$. This is then the \mathbb{C}^* -action induced by

$$\lambda \cdot ([x_0 : x_1 : x_2 : x_3 : x_4]) = [x_0 : x_1 : x_2 : \lambda x_3 : \lambda^{-1} x_4].$$

We then introduce the involution

$$\tau([x_0 : x_1 : x_2 : x_3 : x_4]) = [x_0 : x_2 : x_1 : x_4 : x_3].$$

This automorphism of \mathbb{P}^4 preserves Q and C and lifts to an automorphism of X . Its adjoint action maps a generator of the \mathbb{C}^* -action of interest to its inverse, and by Corollary 5, we deduce the vanishing of the Futaki invariant of X for any Kähler class.

7.3 Family 3.10

Let X be a K -polystable element in the family 3.10 such that $\text{Aut}(X)$ is infinite. Then, from [1, Section 5.17], up to isomorphism, we may assume that $X = \text{Bl}_{C_1 \cup C_2}(Q_a)$ is the blow-up of the quadric

$$Q_a = \{w^2 + xy + zt + a(xt + yz) = 0\} \subset \mathbb{P}^4$$

along the two disjoint smooth irreducible conics $C_1 \subset Q_a$ and $C_2 \subset Q_a$ given by

$$C_1 = \{w^2 + zt = x = y = 0\}$$

and

$$C_2 = \{w^2 + xy = z = t = 0\}$$

where $[x, y, z, t, w]$ stand for the homogeneous coordinates on \mathbb{P}^4 and where $a \in \mathbb{C} \setminus \{-1, +1\}$ is a complex parameter. Moreover, for $a = 0$, $\text{Aut}_0(X) \simeq (\mathbb{C}^*)^2$ and for $a \neq 0$, $\text{Aut}_0(X) \simeq \mathbb{C}^*$.

7.3.1 Case $a = 0$

In this situation, the $(\mathbb{C}^*)^2$ -action on X is the lift of the action on Q_0 induced by the following formula, for $(\alpha, \beta) \in (\mathbb{C}^*)^2$:

$$(\alpha, \beta) \cdot ([x : y : z : t : w]) = [\alpha x : \alpha^{-1} y : \beta z : \beta^{-1} t : w].$$

Consider the group $G = \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ generated by (σ, τ) defined by

$$\sigma([x : y : z : t : w]) = [y : x : z : t : w]$$

and

$$\tau([x : y : z : t : w]) = [x : y : t : z : w].$$

Then $G \subset \text{Aut}(Q_0)$, and G preserves C_1 and C_2 . It also leaves invariant the class $\iota^*[\omega_{FS}]$ on Q_0 , where $\iota : Q_0 \rightarrow \mathbb{P}^4$ denotes the inclusion and ω_{FS} the Fubini–Study metric. Hence, hypothesis (i) and (ii) of Corollary 5 are satisfied. Finally, $\text{Ad}_\sigma(v_1) = -v_1$ and $\text{Ad}_\tau(v_2) = -v_2$, where v_1 generates the \mathbb{C}^* -action $\alpha \mapsto [\alpha x : \alpha^{-1} y : z : t : w]$ while v_2 generates the \mathbb{C}^* -action $\beta \mapsto [x : y : \beta z : \beta^{-1} t : w]$. Then, Corollary 5 implies the vanishing of the Futaki invariant on X for any class.

7.3.2 Case $a \neq 0$

The same argument as in the previous case applies, where this time the \mathbb{C}^* -action of $\text{Aut}_0(X)$ is induced by the diagonal of the above, given by

$$\alpha \cdot ([x : y : z : t : w]) = ([\alpha x : \alpha^{-1}y : \alpha z : \alpha^{-1}t : w]).$$

and the group $G \simeq \mathbb{Z}/2\mathbb{Z}$ is generated by

$$\varsigma([x : y : z : t : w]) = ([y : x : t : z : w]).$$

7.4 Family 3.15

From [1, Section 5.20], the only smooth K -polystable Fano threefold in family 3.15 is given by the blow-up $X = \text{Bl}_{L \cup C}(Q) \rightarrow Q$ of the quadric

$$Q = \{x_0^2 + 2x_1x_2 + 2x_1x_4 + 2x_2x_3 = 0\} \subset \mathbb{P}^4$$

along the line

$$L = \{x_0 = x_1 = x_2 = 0\}$$

and the smooth conic (disjoint from L)

$$C = \{x_0^2 + 2x_1x_2 = x_3 = x_4 = 0\}.$$

The automorphism group of X satisfies $\text{Aut}_0(X) \simeq \mathbb{C}^*$ with \mathbb{C}^* -action given, for $\lambda \in \mathbb{C}^*$, by (the lift of)

$$\lambda \cdot ([x_0 : x_1 : x_2 : x_3 : x_4]) = [\lambda x_0 : \lambda^2 x_1 : x_2 : \lambda^2 x_3 : x_4].$$

The involution

$$\tau([x_0 : x_1 : x_2 : x_3 : x_4]) = [x_0 : x_2 : x_1 : x_4 : x_3]$$

preserves Q , L and C . It also leaves the class $\iota^*[\omega_{FS}]$ invariant, where $\iota : Q \rightarrow \mathbb{P}^4$ is the inclusion. Then, Corollary 5 applies to $X \rightarrow Q$ and $G = \langle \tau \rangle \simeq \mathbb{Z}/2\mathbb{Z}$, so that the Futaki invariant of X identically vanishes on \mathcal{K}_X .

7.5 Families 3.19 and 3.20

Consider the smooth quadric Fano threefold

$$Q = \{x_0^2 + x_1x_2 + x_3x_4 = 0\} \subset \mathbb{P}^4.$$

The family 3.19 (resp. 3.20) is obtained by blowing-up Q in two points (respectively two disjoint lines). More precisely, we can obtain the unique Fano threefold in family 3.19 by considering X_1 to be the blow-up of Q along the points

$$P_1 = [0 : 0 : 0 : 1 : 0]$$

and

$$P_2 = [0 : 0 : 0 : 0 : 1].$$

The unique Fano threefold X_2 in family 3.20 is the blow-up of Q along the two disjoint lines

$$L_1 = \{x_0 = x_1 = x_3 = 0\}$$

and

$$L_2 = \{x_0 = x_2 = x_4 = 0\}.$$

In both cases, the Fano threefold X_i is K -polystable (see [12, 19, 20]) and the group $\text{Aut}_0(X_i)$ is isomorphic to $\mathbb{C}^* \times \text{PGL}_2(\mathbb{C})$ (see [17, Section 5]). We then have $\text{aut}(X_i) = \mathbb{C} \oplus \mathfrak{sl}_2(\mathbb{C})$. From the discussion in Sect. 3, the Futaki invariant of $(X_i, [\omega_i])$ vanishes on the $\mathfrak{sl}_2(\mathbb{C})$ -component of $\text{aut}(X_i)$ for any Kähler class $[\omega_i]$. Therefore, to check the vanishing of the Futaki invariant on $(X_i, [\omega_i])$, it remains to check the vanishing on the \mathbb{C} -component of $\text{aut}(X_i)$.

To this aim we introduce the involution

$$\tau([x_0 : x_1 : x_2 : x_3 : x_4]) = [x_0 : x_2 : x_1 : x_4 : x_3].$$

This automorphism of \mathbb{P}^4 preserves Q , and swaps the two connected components of the blown-up locus in both cases. Therefore, τ lifts to an automorphism of X_i , still denoted τ , for $i \in \{1, 2\}$. Note that on X_i , any Kähler class of the form

$$[\omega_\varepsilon] := c_1(X_i) + \varepsilon(c_1(\mathcal{O}(E_1^i)) + c_1(\mathcal{O}(E_2^i)))$$

is τ -invariant, where $\varepsilon \in \mathbb{R}$ is chosen so that

$$c_1(X_i) + \varepsilon(c_1(\mathcal{O}(E_1^i)) + c_1(\mathcal{O}(E_2^i))) > 0$$

and the E_j^i 's denote the exceptional divisors of the blow-up $X_i \rightarrow Q$.

Remark 6 While $[\omega_\varepsilon]$ is τ -invariant, it is not true that every Kähler class on X_i is τ -invariant. This is the reason we are not able to conclude the vanishing of the Futaki invariant for every Kähler class in these families. This is a general phenomenon for all the varieties considered in Theorem 2 – the reason we are not able to conclude the vanishing of the Futaki invariant on the whole Kähler cone is that there are some Kähler classes where we have not been able to find an involution preserving that class.

Next, we investigate how this action interacts with the generator of the \mathbb{C}^* -component in $\text{Aut}_0(X_i)$, to verify that we can apply Proposition 4 to deduce the vanishing of the Futaki invariant. We do this for the two families separately.

7.5.1 Family 3.19

We follow the discussion in [17, Lemma 5.13]. An automorphism of X_1 comes from an automorphism of \mathbb{P}^4 that leaves Q and $\{P_1\} \cup \{P_2\}$ invariant. By linearity, such an automorphism preserves the line spanned by the two points, and thus its orthogonal complement $\Pi = \{x_3 = x_4 = 0\}$. It then leaves the conic $C = Q \cap \Pi$ invariant. From [17, Lemma 5.7], the \mathbb{C}^* -component of $\text{Aut}_0(X_1)$ can be identified with the pointwise stabiliser of C in $\text{Aut}_0(Q)$. This is then the \mathbb{C}^* -action given by

$$\lambda \cdot ([x_0 : x_1 : x_2 : x_3 : x_4]) = [x_0 : x_1 : x_2 : \lambda x_3 : \lambda^{-1} x_4].$$

The adjoint action of τ maps a generator of this action to its inverse, and by Proposition 4, we deduce the vanishing of the Futaki invariant of $(X_1, [\omega_\varepsilon])$.

7.5.2 Family 3.20

Following the discussion in [17, Lemma 5.14], the \mathbb{C}^* -component of $\text{Aut}_0(X_2)$ is obtained as follows. An element in $\text{Aut}_0(Q, L_1 \cup L_2)$ must preserve the linear span of L_1 and L_2 , that is $Q \cap \{x_0 = 0\}$. It then leaves invariant

$$Q' = \{x_0 = x_1 x_2 + x_3 x_4 = 0\}.$$

The group $\text{Aut}_0(Q, L_1 \cup L_2)$ acts on the family of lines $(\ell_t)_{t \in \mathbb{P}^1}$ in Q' given by

$$[x_1 : x_3] \mapsto [0 : x_1 : t x_3 : x_3 : -t x_1] \subset Q'.$$

The \mathbb{C}^* -component of $\text{Aut}_0(X)$ then corresponds to the stabiliser of the lines $L_1 = \ell_\infty$ and $L_2 = \ell_0$ under this action. In coordinates, the action is given by

$$\lambda \cdot ([x_0 : x_1 : x_2 : x_3 : x_4]) = [\lambda x_0 : x_1 : \lambda^2 x_2 : x_3 : \lambda^2 x_4].$$

As with family 3.19, using the τ -action and the Ad-invariance of the Futaki invariant, we can conclude that the Futaki invariant of $(X_2, [\omega_\varepsilon])$ vanishes.

7.6 Family 4.4

Up to isomorphism, there is a unique smooth Fano threefold X in family 4.4. Its automorphism group satisfies $\text{Aut}_0(X) \simeq (\mathbb{C}^*)^2$, and it is K -polystable from [12, 19, 20]. Recall that the smooth Fano threefold X_1 in family 3.19 can be obtained as a blow-up along two points of a smooth quadric $Q \subset \mathbb{P}^4$. We can then realise the manifold X as the blow-up of X_1 along the proper transform of the conic that passes

through the blown-up points in Q . Coming back to our parametrisation in Sect. 7.5.1, we can take $Q \subset \mathbb{P}^4$ to be

$$Q = \{x_0^2 + x_1x_2 + x_3x_4 = 0\} \subset \mathbb{P}^4$$

and the blown-up points to be

$$P_1 = [0 : 0 : 0 : 1 : 0]$$

and

$$P_2 = [0 : 0 : 0 : 0 : 1].$$

Then, the conic in Q joining P_1 and P_2 is

$$C_1 = \{x_1 = x_2 = x_0^2 + x_3x_4 = 0\} \subset Q.$$

The $(\mathbb{C}^*)^2$ -action on Q that lifts to X through the two blow-up maps $X \rightarrow X_1 \rightarrow Q$ is given in coordinates by

$$(\lambda, \mu) \cdot ([x_0 : x_1 : x_2 : x_3 : x_4]) = [x_0 : \lambda x_1 : \lambda^{-1} x_2 : \mu x_3 : \mu^{-1} x_4].$$

Again, the involution

$$\tau([x_0 : x_1 : x_2 : x_3 : x_4]) = [x_0 : x_2 : x_1 : x_4 : x_3]$$

preserves C_1 and swaps the blown-up points. Arguing as before, we see that the Futaki invariant of X will vanish in classes of the form

$$c_1(X) + \varepsilon c_1(\mathcal{E}) + \delta (c_1(E_1) + c_1(E_2))$$

for $(\varepsilon, \delta) \in \mathbb{R}^2$ small enough and where \mathcal{E} is the exceptional divisor of $X \rightarrow X_1$, while E_1 and E_2 are the strict transforms of the exceptional divisors of $X_1 \rightarrow Q$. Note that after scaling, this gives a 3-dimensional family in the Kähler cone of X .

7.7 Family 5.1

From [1, Section 5.23], the unique smooth Fano threefold X in family 5.1 is K -polystable. It can be described as follows. Consider first the smooth quadric in \mathbb{P}^4

$$Q = \{x_1x_2 + x_2x_3 + x_3x_1 + x_4x_5 = 0\} \subset \mathbb{P}^4$$

where we denote by $[x_1 : x_2 : x_3 : x_4 : x_5]$ the homogeneous coordinates on \mathbb{P}^4 . We then fix a smooth conic $C = Q \cap \{x_4 = x_5 = 0\} \subset Q$ and points $P_1 = [1 : 0 : 0 : 0 : 0]$, $P_2 = [0 : 1 : 0 : 0 : 0]$ and $P_3 = [0 : 0 : 1 : 0 : 0]$ in Q . Let $Y \rightarrow Q$

the blow-up of Q in the three points $(P_i)_{1 \leq i \leq 3}$ and \check{C} the strict transform of C in Y . Then, X is obtained as the blow-up of Y along \check{C} . Its automorphism group satisfies $\text{Aut}_0(X) \simeq \mathbb{C}^*$, where the \mathbb{C}^* -action is the lift of the action defined on Q by

$$\lambda \cdot ([x_1 : x_2 : x_3 : x_4 : x_5]) = [\lambda x_1 : \lambda x_2 : \lambda x_3 : \lambda^2 x_4 : x_5].$$

The manifold X also admits an involution which is the lift of the involution τ defined on Q by

$$\tau([x_1 : x_2 : x_3 : x_4 : x_5]) = [x_1 : x_2 : x_3 : x_5 : x_4].$$

We observe that τ preserves the Kähler class associated to the hyperplane section $H \cap Q$ and fixes C , as well as the points P_1, P_2 and P_3 . Hence, all the $(1, 1)$ -classes on X are invariant under the (lifted) involution. As the adjoint action of τ maps the generator of the \mathbb{C}^* -action to its inverse, we conclude as in 5 the vanishing of the Futaki invariant of X for all its Kähler classes.

8 Hypersurfaces in $\mathbb{P}^2 \times \mathbb{P}^2$ and their blow-ups

In this section, we consider families $N^\circ \mathcal{N}$, with

$$\mathcal{N} \in \{2.24, 3.8, 4.7\}.$$

8.1 Family 2.24

From [12, 19, 20] (see also [1, Section 4.7]), the only K -polystable element in Family 2.24 with infinite automorphism group is given, up to isomorphism, by

$$X = \{xu^2 + yv^2 + zw^2\} \subset \mathbb{P}^2 \times \mathbb{P}^2.$$

It has $\text{Aut}_0(X) \simeq (\mathbb{C}^*)^2$, where the action of $(\alpha, \beta) \in (\mathbb{C}^*)^2$ is given by

$$(\alpha, \beta) \cdot ([x : y : z], [u : v : w]) = ([\alpha^2 x : \beta^2 y : z], [\alpha^{-1} u : \beta^{-1} v : w]).$$

The group $G = \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ acts on $\mathbb{P}^2 \times \mathbb{P}^2$, with the action of $(\sigma, \tau) \in G$ generated by

$$\sigma([x : y : z], [u : v : w]) = ([z : y : x], [w : v : u])$$

and

$$\tau([x : y : z], [u : v : w]) = ([x : z : y], [u : w : v]).$$

Note that $G \subset \text{Aut}(X)$ and that the inclusion $\iota : X \rightarrow \mathbb{P}^2 \times \mathbb{P}^2$ is G -equivariant. Hence we deduce that $\iota^*[\omega_{FS}^i]$ is G -invariant, where ω_{FS}^i denote the Fubini–Study metric on

the i -th factor. Then, any Kähler class on X is G -invariant. Denote by v_1 (resp. v_2) the generator of the \mathbb{C}^* -action $\alpha \cdot ([x : y : z], [u : v : w]) = ([\alpha^2 x : y : z], [\alpha^{-1} u : v : w])$ (resp. $\beta \cdot ([x : y : z], [u : v : w]) = ([x : \beta^2 y : z], [u : \beta^{-1} v : w])$) on X . A direct computation shows

$$\begin{cases} \text{Ad}_\sigma(v_1) = -(v_1 + v_2) \\ \text{Ad}_\tau(v_2) = -(v_1 + v_2). \end{cases}$$

Using Ad-invariance of the Futaki invariant, as discussed in Proposition 4, we deduce that for any Kähler class Ω on X :

$$\begin{cases} \text{Fut}_{(X,\Omega)}(v_1) = -\text{Fut}_{(X,\Omega)}(v_1) - \text{Fut}_{(X,\Omega)}(v_2) \\ \text{Fut}_{(X,\Omega)}(v_2) = -\text{Fut}_{(X,\Omega)}(v_1) - \text{Fut}_{(X,\Omega)}(v_2), \end{cases}$$

hence Fut_X is identically zero.

8.2 Family 3.8

From [1, Section 5.16], the only element in Family 3.8 with infinite automorphism group is given, up to isomorphism, by $X = \text{Bl}_C(Y)$ the blow up of Y along the curve C , where

$$Y = \{(vw + u^2)x + v^2y + w^2z = 0\} \subset \mathbb{P}^2 \times \mathbb{P}^2$$

is a smooth divisor of degree $(1, 2)$ and where $C = \pi_1^{-1}([1 : 0 : 0])$, with π_1 the projection onto the first factor of $\mathbb{P}^2 \times \mathbb{P}^2$. The variety Y is the only element in Family 2.24 with infinite automorphism group, and

$$\text{Aut}(X) \simeq \text{Aut}(Y) \simeq \mathbb{C}^* \rtimes \mathbb{Z}/2\mathbb{Z}.$$

More explicitly, the \mathbb{C}^* -action is for $\lambda \in \mathbb{C}^*$ given by

$$\lambda \cdot ([x : y : z], [u : v : w]) = ([x : \lambda^{-2}y : \lambda^2z], [\lambda u : \lambda^2v : w]),$$

while the $\mathbb{Z}/2\mathbb{Z}$ -action is generated by τ :

$$\tau([x : y : z], [u : v : w]) = ([x : z : y], [u : w : v]).$$

Identifying λ with the corresponding element in $\text{Aut}(Y)$, we have $\tau \circ \lambda \circ \tau^{-1} = \lambda^{-1}$, so that item (iii) in Corollary 5 is satisfied. The inclusion $\iota : Y \rightarrow \mathbb{P}^2 \times \mathbb{P}^2$ is τ -equivariant, and then the classes $\iota^*[\omega_{FS}^i]$ are τ -invariant, for ω_{FS}^i the Fubini–Study metric on each factor of $\mathbb{P}^2 \times \mathbb{P}^2$. This shows that hypothesis (ii) from Corollary 5 holds as well. Finally, the curve C is τ -invariant, and by Corollary 5, the Futaki character of X is identically zero on its Kähler cone.

8.3 Family 4.7

Let X be a smooth Fano threefold in family 4.7. Then it is a blow-up of a smooth divisor W of bidegree $(1, 1)$ on $\mathbb{P}^2 \times \mathbb{P}^2$ along two disjoint curves of bidegrees $(1, 0)$ and $(0, 1)$, and it is K -polystable [12, 19, 20]. To perform computations, we will assume that

$$W = \{xu + yv + zw = 0\} \subset \mathbb{P}^2 \times \mathbb{P}^2,$$

where $[x, y, z]$ and $[u, v, w]$ stand for homogeneous coordinates on the first and second factors respectively. We will denote by $\pi_i : W \rightarrow \mathbb{P}^2$ the natural projection on the i -th factor. We then let $C_i = \pi_i^{-1}([0 : 0 : 1]) \subset W$. Then, $X = \text{Bl}_{C_1 \cup C_2}(W)$ and from [17, Lemmas 7.1 and 7.7], we have

$$\text{Aut}_0(X) \simeq \text{GL}_2(\mathbb{C}).$$

The isomorphism is defined as follows. First, automorphisms of X are induced by automorphisms of W that leave $C_1 \cup C_2$ invariant. Arguing as in [17, Lemma 7.7], they correspond to lift of isomorphisms of \mathbb{P}^2 that leave the set

$$\pi_1(C_1 \cup C_2) = \{[0 : 0 : 1]\} \cup \{[x : y : 0], (x, y) \in \mathbb{C}^2 \setminus \{0\}\}$$

invariant. Those elements are easily identified to elements in $\text{GL}_2(\mathbb{C})$. From Sect. 3, the Futaki invariant vanishes on the $\mathfrak{sl}_2(\mathbb{C})$ -component in $\text{aut}(X)$. We can identify a supplementary subspace of $\mathfrak{sl}_2(\mathbb{C})$ in $\text{aut}(X)$ by considering the lift to X of the generators of the \mathbb{C}^* -action on \mathbb{P}^2 given by

$$\lambda \cdot ([x : y : z]) = ([\lambda x : y : z]).$$

The lift of this action to W is given by

$$\lambda \cdot ([x : y : z], [u : v : w]) = ([\lambda x : y : z], [\lambda^{-1}u : v : w]). \quad (1)$$

We introduce the involution

$$\tau([x : y : z], [u : v : w]) = ([u : v : w], [x : y : z]).$$

This preserves W , and swaps the curves C_1 and C_2 . It also swaps the $(1, 1)$ -classes $\pi_1^*[\omega_{FS}]$ and $\pi_2^*[\omega_{FS}]$. Finally, its adjoint actions maps a generator of the \mathbb{C}^* -action (1) to its inverse. Then, following Sect. 2, we deduce the vanishing of the Futaki invariant on X for any Kähler class of the form

$$c_1(X) + \varepsilon \pi^*(\pi_1^*[\omega_{FS}] + \pi_2^*[\omega_{FS}]) + \eta(c_1(\mathcal{O}(E_1)) + c_1(\mathcal{O}(E_2))),$$

where $\pi : X \rightarrow W$ denotes the blow-down map, E_1 and E_2 the exceptional divisors, and $(\varepsilon, \eta) \in \mathbb{R}^2$ are chosen so that the class is positive.

9 Remaining cases

We finish with families $N^\circ\mathcal{N}$, with

$$\mathcal{N} \in \{2.20, 3.9, 3.13, 4.2\}.$$

9.1 Family 2.20

Consider the Plücker embedding of $\text{Gr}(2, 5)$ in \mathbb{P}^9 . Any smooth intersection of this embedded sixfold with a linear subspace of codimension 3 is a Fano manifold. We call this Fano threefold V_5 and it is the unique member of family 1.15 of Fano threefolds.

Now, let C be a twisted cubic in V_5 and let $X = \text{Bl}_C(V_5)$. Then X is a member of the family 2.20 of Fano threefolds. Up to isomorphism, there is a unique choice of curve such that X has infinite automorphism group [17, Lemma 6.10]. In this case, $\text{Aut}(X)$ is a semidirect product $\mathbb{C}^* \rtimes \mathbb{Z}/2\mathbb{Z}$.

In [1, Section 5.8], it is shown that the unique element in family 2.20 with infinite automorphism group is K -polystable. Moreover, the following explicit description of X is given. First, V_5 can be realised as the subvariety of \mathbb{P}^6 cut out by the equations

$$\begin{cases} x_4x_5 - x_0x_2 + x_1^2 = 0 \\ x_4x_6 - x_1x_3 + x_2^2 = 0 \\ x_4^2 - x_0x_3 + x_1x_2 = 0 \\ x_1x_4 - x_0x_6 - x_2x_5 = 0 \\ x_2x_4 - x_3x_5 - x_1x_6 = 0. \end{cases}$$

We will then identify V_5 with this variety. Then, we can chose C to be the twisted cubic parametrised by

$$([r : s]) \mapsto ([r^3 : r^2s : rs^2 : s^3 : 0 : 0 : 0]) \in V_5.$$

We consider $X = \text{Bl}_C(V_5)$ with this parametrisation. The $\mathbb{C}^* \rtimes \mathbb{Z}/2\mathbb{Z}$ -action on \mathbb{P}^6 generated by

$$\lambda \cdot ([x_0 : x_1 : x_2 : x_3 : x_4 : x_5 : x_6]) = [\lambda^3x_0 : \lambda^5x_1 : \lambda^7x_2 : \lambda^9x_3 : \lambda^6x_4 : \lambda^4x_5 : \lambda^8x_6]$$

for $\lambda \in \mathbb{C}^*$ and the involution

$$\tau([x_0 : x_1 : x_2 : x_3 : x_4 : x_5 : x_6]) = [x_3 : x_2 : x_1 : x_0 : x_4 : x_6 : x_5]$$

preserves V_5 and C , hence lifts to X . This provides the isomorphism

$$\text{Aut}(X) \simeq \mathbb{C}^* \rtimes \mathbb{Z}/2\mathbb{Z}.$$

Note that conjugation by τ sends a generator of the \mathbb{C}^* -action to its inverse. As $H^{1,1}(V_5, \mathbb{R})$ is generated by the class of a hyperplane section in \mathbb{P}^6 , and as the class

of the Fubini–Study metric on \mathbb{P}^6 is τ -invariant, we can apply Corollary 5 to X , and we deduce the vanishing of the Futaki invariant on the Kähler cone of X .

9.2 Families 3.9 and 4.2

We now consider families 3.9 and 4.2. Any member of one of these families have $\text{Aut}_0(X) \simeq \mathbb{C}^*$ and is K -polystable by [1, Section 4.6], which we will follow closely.

Let S be either \mathbb{P}^2 or $\mathbb{P}^1 \times \mathbb{P}^1$ and let $\mathcal{C} \subset S$ be a smooth irreducible curve given by a quartic if $S = \mathbb{P}^2$ and a $(2, 2)$ -curve in the other case. Denote by pr_i the projection of $\mathbb{P}^1 \times S$ onto the i -th factor. We then set $\mathcal{B} = \text{pr}_2^*(\mathcal{C}) \simeq \mathbb{P}^1 \times \mathcal{C}$, $\mathcal{E} = \text{pr}_1^*([1 : 0])$ and $\mathcal{E}' = \text{pr}_1^*([0 : 1])$. We consider

$$G = \mathbb{C}^* \rtimes \mathbb{Z}/2\mathbb{Z}$$

acting on \mathbb{P}^1 by

$$\lambda \cdot [u : v] = [u : \lambda v]$$

and

$$\tau([u : v]) = [v : u].$$

The G -action lifts to $\mathbb{P}^1 \times S$, with the involution τ swapping \mathcal{E} and \mathcal{E}' . We then introduce $\eta : W \rightarrow \mathbb{P}^1 \times S$ a double cover branched over $\mathcal{E} + \mathcal{E}' + \mathcal{B}$, and $\overline{E}, \overline{E}'$ and \overline{B} the preimages on W of the surfaces $\mathcal{E}, \mathcal{E}'$ and \mathcal{B} respectively. Then, set $\hat{X} \rightarrow W$ the blow-up of W along the curves $\overline{E} \cap \overline{B}$ and $\overline{E}' \cap \overline{B}$ with exceptional surfaces \hat{S} and \hat{S}' . We denote the proper transforms of $\overline{E}, \overline{E}'$ and \overline{B} by $\hat{E}, \hat{E}', \hat{B}$ respectively. Finally, X is obtained as the image of a contraction $\hat{X} \rightarrow X$ of \hat{B} to a curve isomorphic to \mathcal{C} . We set E, E', S and S' the proper transforms on X of $\hat{E}, \hat{E}', \hat{S}$ and \hat{S}' respectively.

One can check that all the birational maps involved in producing X are G -equivariant, and we obtain $\text{Aut}_0(X) \simeq \mathbb{C}^*$. Moreover, the involution on X induced by τ (that we will still denote τ) swaps E and E' , and also swaps S and S' . Hence, the Kähler classes $c_1(E) + c_1(E')$ and $c_1(S) + c_1(S')$ are both τ -invariant. Clearly, on $\mathbb{P}^1 \times S$, the adjoint action of the involution τ maps a generator of the \mathbb{C}^* -action to its inverse. This remains true on W by equivariance, and thus on X that is birationally equivalent to W by continuity of holomorphic vector fields away from the exceptional loci. Then, Proposition 4 applies to show that the Futaki invariant of X vanishes in any Kähler class of the form

$$c_1(X) + \varepsilon(c_1(E) + c_1(E')) + \delta(c_1(S) + c_1(S')),$$

where $(\varepsilon, \delta) \in \mathbb{R}^2$ is chosen so that the class is positive.

To understand the subset of the Kähler cone these classes generate, we use the following alternative description of X , still following [1, Section 4.6].

9.2.1 Family 3.9

This is the case when $\mathcal{S} = \mathbb{P}^2$. X can then also be obtained as the blow-up $\phi : X \rightarrow V$ of V along a curve $C \subset V$ where

$$\pi : V = \mathbb{P}(\mathcal{O} \oplus \mathcal{O}(2)) \rightarrow \mathbb{P}^2 = \mathcal{S}$$

is a \mathbb{P}^1 -bundle, and $C = \pi^*\mathcal{C} \cap E_V$, where E_V is the zero section of π . We also have that the strict transform of E_V (resp. of the infinity section E'_V , and of $\pi^*\mathcal{C}$) on X is E (resp. E' and S'), while the exceptional divisor of ϕ is S . Hence we get the relation $c_1(E) + c_1(E') = 0$ in this case (but $c_1(S) + c_1(S') \neq 0$), and we obtain a 2-dimensional family of classes that admit cscK metrics given by

$$(\delta, r) \rightarrow r(c_1(X) + \delta(c_1(S) + c_1(S'))).$$

9.2.2 Family 4.2

This is the case when $\mathcal{S} = \mathbb{P}^1 \times \mathbb{P}^1$. Again, we can recover X from the maps

$$\pi : X \rightarrow V$$

and

$$\phi : V \rightarrow \mathcal{S},$$

with π the contraction of S to a curve isomorphic to \mathcal{C} and ϕ a \mathbb{P}^1 -bundle over $\mathbb{P}^1 \times \mathbb{P}^1$. According to [8, Section 10], we have

$$\text{Pic}(X) = \mathbb{Z}[H_1] \oplus \mathbb{Z}[H_2] \oplus \mathbb{Z}[E] \oplus \mathbb{Z}[E']$$

where $H_i = (\pi \circ \phi)^*(\ell_i)$ and ℓ_1, ℓ_2 denote two different rulings of $\mathbb{P}^1 \times \mathbb{P}^1$. There are relations $-K_X \sim 2(H_1 + H_2) + E + E'$, $S \sim H_1 + H_2 - E + E'$ and $S' \sim H_1 + H_2 + E - E'$, so that the Kähler classes described above can be written

$$2(1 + \delta)(c_1(H_1) + c_1(H_2)) + (1 + \varepsilon)(c_1(E) + c_1(E')).$$

Together with scaling we therefore obtain a 3-dimensional family of classes with vanishing Futaki invariant.

9.3 Family 3.13

Let X be a smooth K -polystable Fano threefold in family 3.13. From [1, Section 5.19], either $\text{Aut}_0(X) \simeq \text{PGL}_2(\mathbb{C})$, and so from Sect. 3 the Futaki invariant vanishes identically, or $\text{Aut}_0(X) \simeq \mathbb{C}^*$. In the latter case, denoting $[x_0 : x_1 : x_2]$, $[y_0 : y_1 : y_2]$

and $[z_0 : z_1 : z_3]$ the homogeneous coordinates on the first, second and third factors of $\mathbb{P}^2 \times \mathbb{P}^2 \times \mathbb{P}^2$, X is given by the equations

$$\begin{cases} x_0y_0 + x_1y_1 + x_2y_2 & = 0 \\ y_0z_0 + y_1z_1 + y_2z_2 & = 0 \\ (1 + s)x_0z_1 + (1 - s)x_1z_0 - 2x_2z_2 & = 0 \end{cases}$$

in $\mathbb{P}^2 \times \mathbb{P}^2 \times \mathbb{P}^2$, for $s \notin \{-1, 0, 1\}$, and

$$\text{Aut}(X) \simeq \mathbb{C}^* \rtimes \mathfrak{S}_3.$$

The \mathbb{C}^* -action for $\lambda \in \mathbb{C}^*$ is given on a point P with homogeneous coordinates $([x_0 : x_1 : x_2], [y_0 : y_1 : y_2], [z_0 : z_1 : z_3])$ by

$$\lambda \cdot (P) = ([\lambda x_0 : \lambda^{-1}x_1 : x_2], [\lambda^{-1}y_0 : \lambda y_1 : y_2], [\lambda z_0 : \lambda^{-1}z_1 : z_3]).$$

Further, there are two involutions $\tau_{x,z}$ and $\tau_{y,z}$ in $\text{Aut}(X)$, whose actions are given by

$$\tau_{x,z}(P) = ([z_1 : z_0 : z_2], [y_1 : y_0 : y_2], [x_1 : x_0 : x_2])$$

and

$$\tau_{y,z}(P) = \left([x_1 : x_0 : -x_2], [(1 - s)z_0 : (1 + s)z_1 : 2z_2], \left[\frac{y_0}{1 - s} : \frac{y_1}{1 + s} : \frac{y_2}{2} \right] \right).$$

Note that $\tau_{x,z} \circ \lambda \circ \tau_{x,z}^{-1} = \lambda^{-1}$ and $\tau_{y,z} \circ \lambda \circ \tau_{y,z}^{-1} = \lambda^{-1}$ (where we identified λ with the corresponding element in $\text{Aut}(X)$). From [1, Diagram 5.19.1], the projection maps $\eta_x, \eta_y, \eta_z : \mathbb{P}^2 \times \mathbb{P}^2 \times \mathbb{P}^2 \rightarrow \mathbb{P}^2$ induce holomorphic maps, still denoted η_x, η_y and η_z , from X to \mathbb{P}^2 . If we denote $\alpha_i := \eta_i^*[\omega_{FS}] \in H^{1,1}(X, \mathbb{R})$ the pullback of the class of the Fubini–Study form, for $i \in \{x, y, z\}$, by equivariance of the projections, we see that α_y is $\tau_{x,z}$ -invariant while α_x is $\tau_{y,z}$ -invariant. Hence, for any $\varepsilon > 0$ small enough, the class $c_1(X) + \varepsilon\alpha_x$ is $\tau_{y,z}$ -invariant and the class $c_1(X) + \varepsilon\alpha_y$ is $\tau_{x,z}$ -invariant. From Proposition 4, the Futaki invariants of $(X, c_1(X) + \varepsilon\alpha_x)$ and $(X, c_1(X) + \varepsilon\alpha_y)$ vanish. Hence, X will carry cscK deformations of its Kähler–Einstein metrics in the classes $c_1(X) + \varepsilon\alpha_y$ and $c_1(X) + \varepsilon\alpha_x$ for ε small enough by LeBrun–Simanca’s openness theorem.

Remark 7 We have used two different involutions $\tau_{x,z}$ and $\tau_{y,z}$ in the above to deduce the vanishing of the Futaki invariant in the classes $c_1(X) + \varepsilon\alpha_y$ and $c_1(X) + \varepsilon\alpha_x$. We are therefore not able from these arguments to deduce that the Futaki invariant vanishes on the sums of these classes. Hence we still only get a 2-dimensional family of classes with vanishing Futaki invariant in this case.

Acknowledgements The authors would like to thank Hendrik Süß for kindly answering our questions on complexity one Fano threefolds. We also thank Jesus Martinez-Garcia for helpful comments, and the referee for their suggestions that helped improve the paper.

Funding Open access funding provided by University of Gothenburg. CT is partially supported by the Grants MARGE ANR-21-CE40-0011 and BRIDGES ANR-FAPESP ANR-21-CE40-0017. LMS is funded by a Marie Skłodowska-Curie Individual Fellowship, funded from the European Union's Horizon 2020 research and innovation programme under Grant Agreement No 101028041.

Declarations

Conflict of interest The authors declare that there are no conflicts of interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Araujo, C., Castravet, A.M., Cheltsov, I.A., Fujita, K., Kaloghiros, A.S., Martinez-Garcia, J., Shramov, C., Süß, H., Viswanathan, N.: The Calabi Problem for Fano threefolds. *MPIM Preprint Series*, 31, (2021)
2. Batyrev, V.V., Selivanova, E.N.: Einstein-Kähler metrics on symmetric toric Fano manifolds. *J. Reine Angew. Math.* **512**, 225–236 (1999)
3. Calabi, E.: Extremal Kähler metrics. In: Seminar on differential geometry. *Ann. Math. Stud.*, vol. 102, pp. 259–290, Princeton Univ. Press, Princeton, NJ, (1982) (ISBN: 0-691-08268-5)
4. Calabi, E. Extremal Kähler metrics. II. In: *Differential geometry and complex analysis: a volume dedicated to the memory of Harry Ernest Rauch*, pp. 95–114, Springer, Berlin (1985) (ISBN: 3-540-13543-X)
5. Cheltsov, I.A., Park, J.: K-stable Fano threefolds of rank 2 and degree 30. *Eur. J. Math.* **8**(3), 834–852 (2022)
6. Chen, X., Donaldson, S.K., Sun, S.: Kähler-Einstein metrics on Fano manifolds. I, II, III. *J. Am. Math. Soc.* **28**(1), 183–197 (2015)
7. Donaldson, S.K.: Scalar curvature and stability of toric varieties. *J. Differ. Geom.* **62**(2), 289–349 (2002)
8. Fujita, K.: On K -stability and the volume functions of \mathbb{Q} -Fano varieties. *Proc. Lond. Math. Soc.* **113**(5), 541–582 (2016)
9. Futaki, A.: An obstruction to the existence of Einstein Kaehler metrics. *Invent. Math.* **73**, 437–443 (1983)
10. Futaki, A.: Kähler-Einstein Metrics and Integral Invariants, vol. 1314. Springer-Verlag, Berlin (1988)
11. Gauduchon, P.: Calabi's extremal Kähler metrics: An elementary introduction
12. Ilten, N., Süß, H.: K-stability for Fano manifolds with torus action of complexity 1. *Duke Math. J.* **166**(1), 177–204 (2017)
13. LeBrun, C., Simanca, S.R.: On the Kähler classes of extremal metrics. In *Geometry and global analysis. Report of the 1st MSJ (Mathematical Society of Japan) International Research Institute, Sendai, Japan, July 12–23, 1993*, pp. 255–271. Sendai: Tôhoku University, Mathematical Institute, (1993)
14. LeBrun, C., Simanca, S.R.: Extremal Kähler metrics and complex deformation theory. *Geom. Funct. Anal.* **4**(3), 298–336 (1994)
15. Matsushima, Y.: Sur la structure du groupe d'homéomorphismes analytiques d'une certaine variété kaehlérinne. *Nagoya Math. J.* **11**, 145–150 (1957)
16. Ono, H., Sano, Y., Yotsutani, N.: Bott manifolds with vanishing Futaki invariant for all Kähler classes. *ArXiv preprint 2305.05924*, (2023)
17. Przyjalkowski, V.V., Cheltsov, I.A., Shramov, C.A.: Fano threefolds with infinite automorphism groups. *Izv. Ross. Akad. Nauk Ser. Mat.* **83**(4), 226–280 (2019)

18. Stoppa, J., Székelyhidi, G.: Relative K-stability of extremal metrics. *J. Eur. Math. Soc. (JEMS)* **13**(4), 899–909 (2011)
19. Süß, H.: Kähler-Einstein metrics on symmetric Fano T -varieties. *Adv. Math.* **246**, 100–113 (2013)
20. Süß, H.: Fano threefolds with 2-torus action. *Doc. Math.* **19**, 905–940 (2014)
21. Tian, G.: On Calabi's conjecture for complex surfaces with positive first Chern class. *Invent. Math.* **101**(1), 101–172 (1990)
22. Tian, G.: Kähler-Einstein metrics with positive scalar curvature. *Invent. Math.* **130**(1), 1–37 (1997)
23. Tian, G., Yau, S.-T.: Kähler-Einstein metrics on complex surfaces with $C_1 > 0$. *Commun. Math. Phys.* **112**, 175–203 (1987)
24. Wang, X., Zhou, B.: On the existence and nonexistence of extremal metrics on toric Kähler surfaces. *Adv. Math.* **226**(5), 4429–4455 (2011)
25. Wang, X.-J., Zhu, X.: Kähler-Ricci solitons on toric manifolds with positive first Chern class. *Adv. Math.* **188**(1), 87–103 (2004)
26. Yau, S.T.: Open problems in geometry. In *Differential geometry. Part 1: Partial differential equations on manifolds. Proceedings of a summer research institute, held at the University of California, Los Angeles, CA, USA, July 8–28, 1990*, pp. 1–28. Providence, RI: American Mathematical Society, (1993)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.