



Do 6–9-year-old children in Denmark adhere to national dietary recommendations and are there sociodemographic disparities? The

Downloaded from: <https://research.chalmers.se>, 2026-03-05 08:53 UTC

Citation for the original published paper (version of record):

Holmegaard, F., Eilersen, A., Lauritzen, L. et al (2026). Do 6–9-year-old children in Denmark adhere to national dietary recommendations and are there sociodemographic disparities? The Generation Healthy Kids study. *European Journal of Nutrition*, 65(1). <http://dx.doi.org/10.1007/s00394-025-03863-y>

N.B. When citing this work, cite the original published paper.



Do 6–9-year-old children in Denmark adhere to national dietary recommendations and are there sociodemographic disparities? The Generation Healthy Kids study

Frederik Holmegaard¹ · Anna Gro Eilersen¹ · Lotte Lauritzen¹ · Christian Mølgaard¹ · Ming Rong Liu² · Ken D. Stark² · Rikard Landberg³ · Rikke Fredenslund Krølner⁴ · Ulla Toft⁵ · Camilla Trab Damsgaard¹

Received: 29 September 2025 / Accepted: 3 December 2025

© The Author(s) 2026

Abstract

Purpose Diet in childhood is important for growth, brain development, and long-term health. Thus, assessing children's adherence to dietary recommendations and identifying sociodemographic groups with low adherence is of great public health relevance. We investigated dietary intake, adherence to recommendations, and sociodemographic differences in a large cohort of Danish children.

Methods We analyzed baseline data from 1094 children aged 6–9 years from 23 schools across Denmark participating in the Generation Healthy Kids study. Diet was assessed by three-day dietary records and food frequency questionnaires for fish and supplements, focusing on key food groups, macronutrients, and iron. Fasting blood samples were collected from $n=347$ and analyzed for nutritional biomarkers of fish, wholegrains, and iron for validation.

Results Overall adherence to dietary recommendations showed a mean±SD score of 4.6 ± 1.0 out of 7.0. However, <15% adhered to the recommendations for fruit+vegetables, fish, and meat, and <33% to recommendations for saturated fat (SFA) and iron. Adherence decreased with age and shorter parental education, due to lower intakes of fruit+vegetables and dairy with age, and lower wholegrain and higher meat intake with shorter parental education. Also, rural children had lower adherence and consumed less fruit+vegetables and more added sugar and SFA than urban. Non-Danish descendants consumed less wholegrains and sugar than Danish, and weight status was not associated with adherence.

Conclusions Danish children had relatively good dietary adherence, but intakes of fruit+vegetables, fish, meat, SFA, and iron remain a concern. Attention should be given to children of older age, short parental education, and rural backgrounds.

Keywords Children's dietary intake · Dietary recommendations · Sociodemographic disparities · Public health

Frederik Holmegaard and Anna Gro Eilersen contributed equally to this paper and share first authorship.

✉ Frederik Holmegaard
fhj@nexs.ku.dk

¹ Department of Nutrition, Exercise and Sports, Faculty of Science, University of Copenhagen, Frederiksberg, Denmark

² Department of Kinesiology and Health Sciences, University of Waterloo, Waterloo, ON, Canada

³ Department of Life Sciences, Chalmers University of Technology, Gothenburg, Sweden

⁴ National Institute of Public Health, University of Southern Denmark, Copenhagen, Denmark

⁵ Department of Prevention, Health Promotion and Community Care, Steno Diabetes Center Copenhagen, Herlev, Denmark

Abbreviations

BMR	Basal metabolic rate
DHA	Docosahexaenoic acid
EI	Energy intake
EPA	Eicosapentaenoic acid
FBDG	Food based dietary guidelines
FFQ	Food frequency questionnaire
FRIDA	The Danish food composition database
GHK	Generation Healthy Kids
NNR2023	Nordic Nutrition Recommendations
n-3 LCPUFA	n-3 long chain polyunsaturated fatty acids
SFA	Saturated fat
vitD	Vitamin D

Introduction

Diet and nutrient intake in childhood are important for optimal growth, cognitive and bone development, and long-term health [1–3]. In Western countries children from the age of 2 years are typically advised to follow the same food based dietary guidelines (FBDG) and nutrient recommendations as adults but adjusted to fit their lower total energy requirements [4–7]. Most dietary recommendations emphasize high intakes of fruit and vegetables, wholegrains, and fish while limiting red and processed meats, saturated fat (SFA), and added sugar. Such a dietary pattern has been shown to be associated with reduced overweight and improved cardiometabolic risk markers in children [8–11], which may influence later risk of lifestyle diseases. Thus, it is of great public health relevance to evaluate children's adherence to dietary recommendations.

Previous dietary surveys have shown that children in Western countries have low adherence to dietary recommendations [4, 12]. A report from the European Food Safety Authority concluded that young children in the EU are at risk of insufficient dietary intakes of iron, vitamin D (vitD), and n-3 long-chain polyunsaturated fatty acids (n-3 LCP-UFAs) [13], a concern also evident among Danish children and adolescents [12]. However, the most recent national dietary survey among Danish children was conducted more than 10 years ago, and it did not include objective nutrient biomarkers, which are important to verify reported intakes and identify nutrient deficiencies.

Children's dietary intake and adherence to recommendations may be influenced by sociodemographic factors. For instance, children of parents with lower education levels have been shown to consume less fruit and vegetables, fish, and dietary fiber than children of parents with higher education [14, 15], and adherence to the recommendations tends to decline as children approach adolescence [15, 16]. Children's dietary intake may also be associated with other factors such as ethnic origin [17, 18] and rurality [19–21], although previous findings are inconsistent. As many sociodemographic factors are interrelated, comprehensive investigations that account for multiple factors simultaneously are needed to better understand social disparities in children's food and nutrient intake. Finally, despite the common perception that children with overweight have unhealthy diets, studies on associations between children's diets and weight status or adiposity remain inconclusive [15, 17, 22].

The present study used baseline data from 1094 6–9-year-old children from 23 schools in three of the five regions in Denmark, who participated in the Generation Healthy Kids (GHK) study. GHK was a cluster-randomized, multi-setting, multi-component intervention designed to promote

healthy weight development in 1st and 2nd grade children by focusing on healthy diets, physical activity, sleep, and screen media habits [23].

The primary aim of the present study was to investigate the dietary intake of the children and their adherence to the Danish FBDG and Nordic Nutrition Recommendations (NNR2023). We focused on intake of fruit and vegetables, wholegrains, meat, milk and dairy, fish, SFA, added sugar, n-3 LCPUFA, vitD, and iron. In a subgroup of children, blood biomarkers were used to assess n-3 LCPUFA and iron status and to validate reported intakes of fish and wholegrains. Finally, we explored whether dietary intake and adherence differed by the children's age, sex, parental education level, country of origin, rurality, and weight status.

Methods

Study design and participants

The present study is a cross-sectional investigation based on baseline data from the GHK study, collected during September–December 2023. The core components of the GHK intervention included a free-of-charge school lunch program, three weekly 40-min sessions of vigorous physical activity during school hours, and classroom exercises and parent workshops on sleep and screen media use, as previously described [23]. The study was approved by the Ethics Committee of Southern Denmark (S-20220094) and registered at clinicaltrials.gov (NCT 05940675).

Initial contact was established with 496 Danish schools from December 2022 to April 2023. Inclusion criteria for schools were: 1) location in The Capital Region, the Zealand Region, or the Region of Southern Denmark, 2) suitable kitchen facilities for school meal preparation. A total of 24 schools were recruited and randomized in clusters 1:1 to the intervention and control. One control school withdrew after randomization and 2006 children in 1st and 2nd grade were invited from the remaining 23 schools. There were no exclusion criteria for the children; if parents, teachers, or the clinically responsible physician (CM) judged that a child could not participate in certain parts of the intervention or measurements, they could still participate in the rest of the study. All children received oral information, and parents received both written and oral information about the study. Written informed consent for participation was obtained from the parents of 1350 children. Among these, 517 children (from the Capital and Zealand Regions only) also provided separate consent for blood sampling. The present study included the 1094 participating children who had ≥ 1 complete day of dietary recording and measurements of height and weight at baseline (Supplementary Fig. 1).

Background information

The children's age was collected from school records, and parents reported their education level and country of origin in an online questionnaire. Education level was reported in seven categories using the International Standard Classification of Education (ISCED97) adapted to a Danish context; 1) Primary education (1–6 years), 2) Lower secondary education (7–10 years), 3) Upper secondary education in high school (10–13 years), 4) Upper secondary vocational education (10–13 years), 5) Short tertiary education (14–16 years), 6) Bachelor's degree or similar (14–17 years), and 7) Master's degree or PhD (18–21 years). These categories were pooled into Short (≤ 13 y), Medium (14–17 y), and Long (≥ 18 y) education, and we used the highest level of education in the household for analyses. Children were classified as Danish descendants if one or both parents were born in Denmark, and as non-Danish descendants if both parents were born in other countries. Most non-Danish descendants originated from Eastern Europe (44%) and the Middle East (26%). When data was only available from one parent ($n=48$), education level and descentance were based on this information. Children's rurality was based on the geographical location of the schools, i.e. urban when located in cities or areas with $\geq 10,000$ inhabitants or rural when located in towns or areas with $< 10,000$ inhabitants.

Dietary records

Information about the children's diet was collected sequentially at each school during the same period as the clinical measurements, but never on the same day. Parents were asked to record their child's dietary intake prospectively for three consecutive days (two weekdays and one weekend day) using an online self-administered dietary assessment tool (myfood24®, Dietary Assessment Ltd., Leeds, UK.). 24-h dietary recalls in myfood24 have previously been validated against an interviewer-administered 24-h dietary recall in English adolescents [24] and against weighed dietary records and biomarkers in German adults [25]. Each day, parents received a link to the food diary with written guidelines in multiple languages and a video guide in Danish and English, and they had access to support by phone and e-mail. If all three days were not completed, reminder e-mails and text messages were sent to the parents 1–2 weeks later, on the same day(s) of the week.

Parents entered all food and drink items consumed by their child by searching a joint food database consisting of a Danish (FRIDA), a Swedish (Livsmedelsverket), an English (myfood24), and an American (USDA FoodData central) food composition database. Portion sizes were estimated

using portion images, descriptions, household measures, or by weighing if the parents wished so.

Estimation of dietary intake

From the dietary records, daily intakes of the food groups: fruit and vegetables, wholegrains, milk and dairy products, as well as meat and meat products were calculated. For mixed dishes, the contribution from each food group was estimated based on common recipes (e.g. for a stir fry with vegetables 50% was estimated to be vegetables). Wholegrain content was calculated using food specific wholegrain factors (e.g. 0.25 for wholegrain bread and buns, and 0.54 for wholegrain muesli and other breakfast cereals) based on the definitions of the Danish wholegrain labelling [26]. Daily intakes of SFA, dietary fiber, and iron were derived directly from myfood24. Intakes of added sugar were estimated as total sugar minus natural sugars (e.g. lactose in dairy) across the categories: sugar and jam, confectionary and chocolate, cakes and desserts, soft drinks, sauces and dips, refined grains, breakfast cereals, and dairy. For food items with missing database information on sugar content, added sugar were estimated from the Danish food composition database (FRIDA) [27] or product labels.

Under-, acceptable, and over-reporters of dietary intake

Children were classified as either under-reporters, acceptable reporters, or over-reporters of energy based on the ratio of mean reported energy intake (EI) to estimated basal metabolic rate (BMR) using the cut-offs suggested by Black [28]; under-reporters: $EI:BMR < 1.09$, acceptable reporters: $1.09 < EI:BMR < 2.26$, and over-reporters: $EI:BMR > 2.26$, assuming a physical activity level of 1.6 [7]. BMR was calculated from the Oxford equations [29] using the child's height, weight, sex, and age.

Food frequency questionnaire

To better assess habitual fish and n-3 LCPUFA intake, we used a self-administered, online food frequency questionnaire (FFQ) that asked specifically about the intake of fish. This fish FFQ was based on a semi-quantitative FFQ developed to assess overall food, energy, and nutrient intake in Denmark [30]. The original FFQ was validated in the Danish Cancer Cohort [31], and the derived fish specific FFQ has previously been validated among children in the FiSK trial [32].

Parents reported the overall frequency of their child's fish intake during the last month as bread topping and as hot meals, separately, along with intake frequencies for 11

and 13 specific fish types within each meal type, respectively. Each question had seven response options ranging from “rarely or never” to “twice or more per day”, which were converted to average daily frequencies. To avoid overestimation, we multiplied the daily frequencies for each fish type with a correction factor, calculated as the recorded frequency of overall fish intake divided by the sum of frequencies for intake of all fish types. Average daily intakes of each fish type were then estimated by multiplying the adjusted frequencies with fish-specific portion sizes (30–50 g for bread toppings and 50–125 g for hot meals) based on validated Danish standard portion sizes [33]. Daily intakes of total, oily, and lean fish were calculated, and intakes of n-3 LCPUFA from fish were estimated using nutritional data from FRIDA.

The FFQ also included questions about frequency, type, and brand of n-3 LCPUFA and vitD supplements used in the past month and during winter (October–April), respectively. Daily intake of n-3 LCPUFA from supplements was estimated and added to the intake from fish to derive total daily intake. Children were categorized as vitD supplement users during winter when they consumed supplements with $\geq 10 \mu\text{g}$ of vitD on ≥ 5 days per week.

Comparison with dietary recommendations

Children’s dietary intake was compared with the official dietary recommendations in Denmark at the time of data collection: the Danish FBDG 2021 and the NNR2023. The FBDG provide recommended intakes of food groups in a 10 MJ diet, i.e. $\geq 600 \text{ g/d}$ fruit and vegetables, $\geq 75 \text{ g/d}$ wholegrains, $\geq 350 \text{ g/week}$ fish of which $\geq 200 \text{ g}$ should be oily fish, $\geq 250 \text{ g/d}$ milk and dairy, and $\leq 350 \text{ g/week}$ meat and meat products. To compare directly with the FBDG and account for differences in EI, all food group intakes except fish and dairy were converted to $\text{g}/10 \text{ MJ}$. For fish the recommended intake for 2-10-year-olds is 175-275 g/week , which we defined as 245 g/week for our cohort (6-9 years), and for dairy, Danish children are recommended to consume the same amount as adults despite a lower EI [6]. Nutrient intakes were converted to E% and g/MJ , as appropriate, for comparison with the NNR2023, e.g. $< 10 \text{ E\%}$ added sugar, $< 10 \text{ E\% SFA}$, $\geq 2 \text{ g}/\text{MJ}$ dietary fiber, and $\geq 9 \text{ mg/d}$ iron [7].

To assess overall adherence to the dietary recommendations, a dietary adherence score was constructed based on seven key components: fruit and vegetables, wholegrains, fish, milk and dairy products, meat and meat products, added sugar, and SFA. For each component, a continuous score was calculated as the ratio of reported to recommended intake, capped at 1 for intakes meeting or exceeding the recommendations. For moderation components, i.e. meat, added

sugar, and SFA, the scores were inverted, so lower intakes resulted in higher scores, and the score 0 was assigned to intakes at or above the 90th percentile of the distributions reported for 4–10-year-olds in the latest national dietary survey [12], similar to the approach used in the Healthy Eating Index [34]. The overall adherence score was calculated as the sum of scores for all components, ranging from 0 to 7, with higher scores indicating better adherence to the dietary recommendations.

Clinical measurements and blood sampling

Clinical measurements and blood sampling were conducted in accordance with standard operating procedures on two separate school days by trained members of the research team. On the first day, children’s height and weight were measured. Height was determined as the mean of three measurements to the nearest 0.1 cm using a portable stadiometer (Seca 217, Seca, Germany), and weight was measured to the nearest 0.1 kg using a digital scale (InBody 270, InBody Co Ltd., California, USA). Children wore light clothing and were instructed to empty their bladder before being weighed. Sex- and age-adjusted z-scores for body mass index (BMI) were calculated using the WHO AnthroPlus software (v1.0.4, Department of Nutrition, WHO, Geneva, Switzerland) and the International Obesity Task Force (IOTF) LMS-curves [35]. Children were categorized as underweight, normal weight, and overweight as determined by age- and sex-specific cut-offs [35] for the IOTF BMI z-score.

On the second day, blood samples were drawn from an antecubital vein before 10:30 am after an overnight fast. In total, 347 children had a blood sample drawn and 325 of them (93.7%) confirmed fasting except for 1–2 glasses of water. Local anesthetic patches (EMLA, Astra Zeneca AB, Södertälje, Sweden) were provided beforehand and applied by the parents in the morning.

Whole blood for fatty acid analysis was collected with lithium heparin, and 500 μL was transferred to a cryotube with the antioxidant 2,6-di-tert-butyl-4-methylphenol (butylated hydroxytoluene; Sigma-Aldrich, St. Louis, MO, USA) and kept on dry ice. Additional blood collected in tubes with EDTA or clot activator was kept at room temperature and centrifuged at $2000 \times g$ at $4 \text{ }^\circ\text{C}$ for 10 min within 6 h after blood sampling at the Department of Nutrition, Exercise, and Sports (NEXS), University of Copenhagen. All samples were stored at $-80 \text{ }^\circ\text{C}$ until analysis.

Analysis of blood samples

Whole blood EPA + DHA: Whole blood fatty acid composition was analyzed within 7 months by gas chromatography

at the Department of Kinesiology and Health, University of Waterloo, ON, Canada, as previously described [36]. Briefly, lipids were extracted with chloroform and methanol containing an internal standard (22:3n-3 ethyl ester, Nu-Chek Prep Inc., Elysian, MN), and fatty acid methyl esters were generated using 14% boron trifluoride (BF_3) in methanol and heating for 1 h at 95 °C on a heating block. The fatty acid methyl esters were then determined using a Scion 8300 gas chromatograph (Scion Instruments Canada, Edmonton, AB, Canada) with a DB-FFAP capillary column (15 m \times 0.1 mm ID \times 0.1 μm film thickness; J&W Scientific from Agilent Technologies, Mississauga, ON, Canada). In the initial analysis, some samples had very low fatty acid concentrations ($<0.5 \mu\text{g}/\text{mg}$) with high SFA:LCPUFA ratios (>6.0), indicating sample oxidation, as previously described [37]. Therefore, all samples were reanalyzed or reintegrated, and the results with the lowest SFA:LCPUFA ratio for each sample were included in the statistical analysis. Samples with SFA:LCPUFA ratios >1 SD above or below the mean in both runs ($n=15$, 4%) were excluded. The relative whole blood content of Eicosapentaenoic Acid (EPA) and Docosahexaenoic Acid (DHA) as a percentage of all fatty acids (FA%) was used as a measure of n-3 LCPUFA status. To allow comparison with omega-3-index cut-offs ($<4\%$ [undesirable], 6%, and $>8\%$ [optimal]) [38], whole blood EPA+DHA was converted to red blood cell equivalents, as suggested by Stark et al. [39].

Plasma alkylresorcinols: Plasma alkylresorcinol concentrations were analyzed at the Department of Life Sciences at Chalmers University of Technology, Sweden by liquid chromatography and tandem mass spectrometry on a QTRAP 6500+ (AB SCIEX, Marlborough, MA, USA), as previously described [40]. Total alkylresorcinol concentration – a marker of wholegrain wheat and rye intake – was used as a biomarker for validating the reported wholegrain intake. The intra- and inter-assay variability were 4.7–7.8% and 10.5–19.0% for each alkylresorcinol homologue, respectively.

Iron status: Whole blood hemoglobin was analyzed on a Sysmex KX 21N Analyzer (Sysmex Corporation, Hyogo, Japan) at NEXS within 6 h after blood sampling. Serum ferritin was analyzed on an IMMULITE 2000 Analyzer (Siemens Healthcare GmbH, Erlangen, Germany) at the Department of Clinical Biochemistry at Rigshospitalet, Copenhagen within one year after sampling.

Statistical analysis

Data were analyzed using R (version 4.2.1). Crude odds ratios (OR) and a logistic regression model for age were used to assess sociodemographic differences between children included and not included in the present study. The

sociodemographic characteristics of included children were compared to those of the Danish background population using chi-square tests.

Dietary intake, nutrient status, and adherence to dietary recommendations are presented as mean \pm SD for normally distributed data and median (IQR) for skewed data. All mean and median intakes include children with zero intakes. Food and nutrient intakes were compared between boys and girls by Student's unpaired t-test and Mann–Whitney U-test when data were normally distributed and skewed, respectively.

Children with acceptable dietary reporting and data on parental education and country of origin were included in the primary multivariable linear regression models to explore associations between sociodemographic factors (age, sex, weight status, parental education level, country of origin, and rurality) and intakes of key food groups and nutrients. The dietary components included in the models – fruit and vegetables, wholegrains, fish, milk and dairy products, meat and meat products, added sugar, and SFA – represent major targets for dietary improvement among children and are linked to both potential nutrient inadequacies and chronic disease prevention goals. All sociodemographic factors were included as fixed effects to assess the independent association of each predictor with dietary intakes. In secondary sensitivity analyses, under-reporters and over-reporters were included in all models first, and subsequently children with unanswered parental education and ethnic origin were included. Among the subgroup with blood samples, models of nutrient biomarkers were used as sensitivity analyses for associations with wholegrains and fish. Model assumptions were evaluated by visual inspection of Q-Q plots and scatter plots of residuals.

The FFQ-reported intakes of fish and n-3 LCPUFA and the registered intakes of wholegrains were validated against whole blood EPA+DHA and total plasma alkylresorcinols, respectively, using Spearman rank correlations. Furthermore, the dose–response relationships between estimated intakes of n-3 LCPUFA and wholegrains were fitted against their respective biomarkers using both linear and logarithmic regression. The resulting plots were compared by visual inspection and R^2 -values.

Results

Children's characteristics and representativeness

Of the 1350 children originally included in GHK, 1142 children had at least one complete day of dietary recording at baseline. Ten were excluded from the analyses due to extreme dietary recordings (mean EI <2000 or $>15,000$ kJ/

day), and 38 were excluded due to missing anthropometric measurements (Supplementary Fig. 1). The 256 non-included children were more likely to have parents with short education (OR, 95% CI 2.83, 1.75–4.61, $P<0.001$), be non-Danish descendants (2.55, 1.58–4.04, $P<0.001$), reside in urban areas (1.37, 1.03–1.83, $P=0.031$) and have overweight (1.65, 1.09–2.45, $P=0.015$), but did not differ with regard to sex and age ($P>0.65$).

Boys and girls were equally represented among the included children (Table 1). Most children had normal weight, and the prevalence of underweight and overweight were comparable to the national numbers reported among Danish 6–7-year-olds ($\chi^2=0.5$, $P=0.795$) [41]. Only 15.5% were from households with ≤ 13 years of education compared to 39.9% among all Danish families with children ($\chi^2=12.3$, $P=0.002$) [42]. There was also a lower proportion of non-Danish descendants (7.8% vs. 16.2% among Danish 6–9-year-olds, $\chi^2=43.5$, $P<0.001$) [43], whereas the proportion of children residing in urban areas was similar to the general population (60.1% vs. 58.3%, $\chi^2=1.2$, $P=0.265$) [44].

In total, 870 (80%), 128 (12%), and 96 (9%) children had three, two, and one complete dietary recording days, respectively, and more than 80% of the children ($n=886$) were

Table 1 Characteristics of all children ($n=1094$)

Demographics and anthropometrics	
Sex, % girls / % boys	48.8 / 51.2
Age, y	7.8 \pm 0.6
Height, cm	130.3 \pm 6.7
Weight, kg	26.7 (24.1–30.4)
BMI, kg/m ²	15.7 (14.8–17.2)
BMI z-score	0.2 \pm 1.1
Weight status	
Underweight	95 (8.7)
Normal weight	858 (78.4)
Overweight	141 (12.9)
Parental education level	
Short education (≤ 13 y)	170 (15.5)
Medium education (14–17 y)	468 (42.8)
Long education (≥ 18 y)	364 (33.3)
Unanswered	92 (8.4)
Origin	
Danish descendants	921 (84.2)
Non-Danish descendants	85 (7.8)
Unanswered	88 (8.0)
Rurality	
Urban	657 (60.1)
Rural	437 (39.9)
Dietary reporting	
Under-reporters	184 (16.8)
Acceptable reporters	886 (81.0)
Over-reporters	24 (2.2)

Data are presented as mean \pm SD, median (IQR), or n (%) when data were normally distributed, skewed, or categorical, respectively

classified as acceptable reporters ($1.09<EI:BMR<2.26$). Over- and under-reporters were more likely to have parents with short education (OR, 95% CI 2.14, 1.37–3.32, $P<0.001$), be non-Danish descendants (1.73, 1.03–2.82, $P=0.030$), be boys (1.45, 1.08–1.96, $P=0.014$), and have overweight (2.55, 1.73–3.74, $P<0.001$).

Dietary intake and nutrient status

Acceptable reporters consumed 7.0 ± 1.3 MJ/d (Table 2), which is within the estimated energy requirement range (6.3–7.8 MJ) for 6–9-year-old children with average physical activity levels (PAL=1.6) according to NNR2023 [7]. Boys had higher EI than girls, but the relative macronutrient intakes were similar in boys and girls and aligned with the 45–60 E%, 25–40 E% and 10–20 E% recommended for carbohydrates, fat and protein [7], respectively. Boys had higher absolute intakes of wholegrains, dairy, meat, and iron, but did not differ from girls with regard to the other investigated food groups or n-3 LCPUFA intake (Table 2). In the subgroup with blood samples ($n=285$), nutrient biomarkers did not differ by sex (Table 2). Iron deficiency (serum ferritin <15 μ g/L) was observed in 7.8% of the children, and 0.4% ($n=1$) had anemia (hemoglobin <6.5 mmol/L). n-3 LCPUFA status was generally low as 39.3% had an omega-3 index $<4\%$ (undesirable), and only 0.4% had an index $>8\%$ (optimal). Including over- and under-reporters did not change the results (Supplementary Table 1).

Validation of parent-reported wholegrain, fish, and n-3 LCPUFA intakes

In the subgroup with blood samples, there was a dose–response relationship between wholegrain intake and plasma total alkylresorcinols (Supplementary Fig. 2), and parent-reported wholegrain intake was moderately correlated with total alkylresorcinols ($r=0.33$, $P<0.001$).

Absolute intakes of fish, oily fish and total n-3 LCPUFA from fish and supplements were moderately correlated with whole blood EPA+DHA ($r=0.30$ – 0.32 , all $P<0.001$), whereas the correlation for lean fish, which contains little n-3 LCPUFA, was weaker ($r=0.21$, $P<0.05$). The dose–response relationship between total n-3 LCPUFA intake and whole blood EPA+DHA was best fitted by a logarithmic function (Supplementary Fig. 3).

Adherence to dietary recommendations

The average adherence score was 4.6 ± 1.0 out of 7.0, indicating that children’s overall adherence to the dietary recommendations was fairly good. However, as shown in Table 3, less than 15% of the children adhered to the dietary

Table 2 Dietary intake (n=886) and nutrient biomarkers (n=264–283) of acceptable reporters

	Boys	Girls	Total
Energy and macronutrient intake			
Energy, MJ	7.3±1.4	6.5±1.2***	7.0±1.3
Carbohydrates, E%	49.9±5.5	50.4±5.6	50.2±5.6
Dietary fiber, g/MJ	2.7±0.7	2.6±0.7	2.7±0.7
Added sugar, E%	8.0 (5.2–11.2)	8.8 (5.4–12.2)*	8.5 (5.4–11.7)
Fat, E%	32.9±5.2	32.7±5.0	32.8±5.1
SFA, E%	11.2±2.5	11.2±2.5	11.2±2.5
Monounsaturated fat, E%	11.2±2.6	11.0±2.4	11.1±2.5
Polyunsaturated fat, E%	5.5±1.4	5.4±1.3	5.5±1.4
Protein, E%	15.0±2.9	14.8±2.8	14.9±2.8
Intake of food groups			
Fruit and vegetables, g/d	218 (130–327)	228 (143–326)	222 (137–326)
Wholegrains, g/d	53 (33–76)	39 (22–60)***	47 (28–70)
Fish ^a , g/d	12 (6–19)	12 (6–18)	12 (6–19)
Oily fish ^a , g/d	4 (0–8)	4 (1–7)	4 (1–7)
Milk and dairy products, g/d	217 (120–363)	188 (90–316)*	203 (104–343)
Meat and meat products, g/d	82 (52–124)	67 (44–101)***	75 (47–114)
Micronutrient intake			
n-3 LCPUFA ^a (mg/d)	120 (42–242)	117 (54–231)	117 (44–238)
Iron (mg/d)	7.7±2.0	6.7±1.7***	7.2±1.9
Food and nutrient biomarkers			
Total plasma alkylresorcinols ^b , nmol/L	275 (165–426)	234 (156–350)	248 (160–391)
Whole blood EPA+DHA ^c , FA%	3.2±0.8	3.3±0.8	3.3±0.8
Whole blood hemoglobin ^d , mmol/L	8.1±0.5	8.1±0.6	8.1±0.5
Serum ferritin ^e , µg/L	28 (21–38)	28 (21–40)	28 (21–39)

Data are presented as mean±SD or median (IQR) when data were normally distributed or skewed, respectively

^aData from FFQ where n=780,

^bn=264,

^cn=270,

^dn=277,

^en=283

*P<0.05, ***P<0.001 for differences between boys and girls

Table 3 Children's adherence to dietary recommendations (n=886)

Food group or nutrient	Danish FBDG or NNR2023	Daily intake	Adherence, n (%)
Fruit and vegetables, g/10 MJ	≥600 g/10 MJ	322 (196–472)	104 (11.7)
Wholegrains, g/10 MJ	≥75 g/10 MJ	68 (41–99)	394 (44.5)
Fish ^a , g/d	≥35 g/d (Min. 245 g/w)	12 (6–19)	56 (7.2)
Oily fish ^a , g/d	≥20 g/d (Min. 140 g/w)	3.7 (0.7–7.2)	46 (5.9)
Milk and dairy products, g/d	≥250 g/d	203 (104–343)	356 (40.2)
Meat and meat products, g/10 MJ	≤50 g/10 MJ (Max. 350 g/w)	110 (72–164)	125 (14.1)
SFA, E%	<10 E%	11.1 (9.4–12.8)	288 (32.5)
Dietary fiber, g/MJ	≥2 g/MJ	2.7±0.7	731 (82.5)
Added sugar, E%	<10 E%	8.5 (5.4–11.7)	568 (64.1)
Iron, mg/d	≥8 mg/d	7.2±1.9	272 (30.7)
VitD supplement during winter ^b	Supplement during winter (Oct-Apr)	487 yes, 307 no	487 (61.3)

Daily intakes are presented as mean±SD, median (IQR), or n when data were normally distributed, skewed, or binomial, respectively

^aData was derived from the food frequency questionnaire, n=780

^bData was derived from the supplements' questionnaire, n=794

recommendations for fruit and vegetables and fish. The low intake of fish, especially oily fish, was in line with the low whole blood n-3 LCPUFA observed (Table 2). The adherence with recommendations for intakes of meat and SFA was also low, as less than one third of the children consumed less than 50 g/10 MJ and 10 E%, respectively. Almost half of the children met the recommendations for wholegrains and dairy, and more than 60% adhered to the recommendations for added sugar, dietary fiber, and vitD supplementation during winter (Table 3). Including children with over- and underreporting in the analyses did not change the results (Supplementary Table 2).

Sociodemographic determinants of dietary intake

In the primary multivariable linear regression models with acceptable reporters, overall adherence to dietary recommendations decreased with age (Table 4). This was mainly due to lower intakes of fruit and vegetables and dairy as well as tendencies to lower fish and higher sugar intake with age. The associations for fruit and vegetables, dairy, and fish were supported in the sensitivity analyses with inclusion of under- and over-reporters and those with missing sociodemographic data, but the significance of the association for sugar decreased somewhat (Supplementary Table 3 and 4). However, the sensitivity analysis among the subgroup with blood samples did not support a decline in fish intake with age, as whole blood EPA+DHA tended to increase with age (Supplementary Table 5).

Longer parental education was associated with better overall adherence to the dietary recommendations, mainly due to higher intakes of wholegrains and lower intakes of meat. The association with wholegrain intake was supported by a positive association between parental education and plasma total alkylresorcinols (Supplementary Table 5). The sensitivity analyses including more children also supported the association with wholegrains (Supplementary Table 3 and 4), but they did not provide clear support for the association with meat.

Children residing in urban areas had higher adherence to the dietary recommendations than rural children. All analyses showed that this was due to a higher intake of fruit and vegetables and lower intakes of added sugar and SFA (Table 4; Supplementary Table 4 and 5). However, urban children consumed less wholegrains than rural children.

The overall adherence score did not differ by sex, weight status, or origin in the primary or any of the sensitivity analyses (Table 4; Supplementary Table 4 and 5). All of the analyses did, however, show that non-Danish descendants consumed less wholegrains, but also less added sugar than Danish descendants. Consistent associations were also seen for sex, with higher consumption of wholegrains and less

added sugar, but a concomitantly reduced intake of fruit and vegetables among boys relative to girls. Weight status was not associated with intake of any of the food groups or nutrients in the primary analysis, but in the sensitivity analyses meat intake was higher, whereas added sugar and SFA intake tended to be lower with increasing weight status.

Discussion

In a large group of Danish 6–9-year-olds, we showed that most children consumed less fruit and vegetables and fish as well as more meat than recommended, whereas almost half of the children reached the recommended level for wholegrain intake and more than 60% adhered to the recommendation for added sugar. Overall adherence to the dietary recommendations improved with higher parental education level and decreased with age. Children living in rural areas had lower adherence to the recommendations for fruit and vegetables, added sugar, and SFA, whereas the wholegrain intake was lower among urban children and Danish descendants.

The low intake of fruit and vegetables aligns with previous findings among Western children [12, 17, 45]. This is a concern as fruit and vegetables are key sources of micronutrients and dietary fiber and have been linked to lower risk of overweight [46] and improved cardiometabolic risk profile in children [10]. In contrast to the relatively high fish intake and n-3 LCPUFA status among children in Norway [47, 48], fish intake in the present study was very low as previously seen in Denmark [12, 32, 45], the US [4], and the UK [17]. This finding was supported by the low whole blood EPA+DHA which also aligned with previous reports in Danish children [32, 49]. Increasing the n-3 LCPUFA status of Danish children would be desirable, as this has been linked to better cognitive function [50] and a more favorable cardiometabolic risk profile [49], and both of these outcomes were found to be improved by an oily fish intake of ≈ 300 g/wk in our previous randomized trial in Danish children [8, 51]. The persistent low intakes of fruit and vegetables and fish may reflect barriers such as cost, accessibility, preferences, time constraints, and cooking skills [32, 52]. In contrast, intake of meat was high, as also seen in the latest national dietary survey [12], suggesting a continued predominance of animal-based foods in the Danish diet despite increased trends towards plant-based diets.

In general, wholegrain intake was high with a median intake approximately three times greater than among children in the UK and the US [4, 53]. Even though Denmark has one of the highest wholegrain recommendations globally [54], nearly half of the children met the recommendation, consistent with the latest national dietary survey [12].

Table 4 Associations between sociodemographic characteristics and dietary intake among children with acceptable dietary reporting and complete data on parental education and country of origin (n = 824)

	Fruit and vegetables, g/10 MJ		Wholegrains, g/10 MJ		Fish ^a , g/d		Milk and dairy, g/10 MJ		Meat and meat products, g/10 MJ		Added sugar, E%		SFA, E%		Adherence score	
	β (95% CI)	P	β (95% CI)	P	β (95% CI)	P	β (95% CI)	P	β (95% CI)	P	β (95% CI)	P	β (95% CI)	P	β (95% CI)	P
Sex																
Girls	Ref	<0.001	Ref	0.727	Ref	0.943	Ref	0.091	Ref	0.008	Ref	0.409	Ref	0.230		
Boys	-59.2 (-86.3, -32.1)		13.6 (7.7, 19.5)		-0.4 (-2.6, 1.8)		-1.3 (-36.4, 33.8)		8.2 (-1.3, 17.6)		-0.1 (-0.5, 0.2)		0.1 (-0.1, 0.2)			
Age, y	-27.6 (-50.5, -4.7)	0.018	-4.1 (-9.1, 0.9)	0.110	-1.7 (-3.6, 0.1)		-67.7 (-97.4, -38.1)		2.8 (-5.2, 10.8)		0.075 (-0.1, 1.1)		0.648 (-0.3, -0.1)		0.002	
Weight status																
Underweight	9.5 (-45.6, 64.5)	0.494	8.9 (-3.2, 20.9)	0.198	3.4 (-1.2, 8.0)		36.1 (-35.3, 107.6)		-15.6 (-34.8, 3.7)		0.0 (-1.4, 1.3)		0.3 (-0.5, 1.0)		0.557	0.1 0.585
Normal weight	Ref		Ref		Ref		Ref		Ref		Ref		Ref		Ref	
Overweight	-25.1 (-79.5, 29.3)		-1.9 (-13.8, 10.0)		0.0 (-4.4, 4.4)		-17.0 (-87.6, 53.7)		-2.3 (-21.3, 16.7)		-0.4 (-1.7, 1.0)		-0.2 (-0.9, 0.5)		-0.1 (-0.4, 0.2)	
Parental education																
Long	Ref	0.148	Ref	0.124	Ref	0.820	Ref	0.003	Ref	0.652	Ref	0.923	Ref	<0.001		
Medium	-23.2 (-59.3, 12.9)		-8.2 (-16.1, -0.4)		-2.1 (-5.0, -0.8)		-3.1 (-49.9, 43.8)		15.2 (2.6, 27.8)		0.0 (-0.8, 0.9)		-0.1 (-0.5, 0.4)		-0.2 (-0.4, -0.1)	
Short	-38.7 (-89.5, 12.0)		-15.2 (-26.3, -4.1)		-3.2 (-7.3, 1.0)		13.6 (-52.2, 79.5)		23.0 (5.3, 40.8)		0.5 (-0.8, 1.7)		0.0 (-0.7, 0.6)		-0.4 (-0.6, -0.1)	
Origin																
Danish	Ref	0.498	Ref	0.827	Ref	0.233	Ref	0.666	Ref	<0.001	Ref	0.051	Ref	0.185		
Non-Danish	17.5 (-33.2, 68.2)		-28.6 (-39.6, -17.5)		0.5 (-3.7, 4.6)		40.0 (-25.8, 105.8)		3.9 (-13.8, 21.6)		-2.5 (-3.7, -1.2)		0.6 (0.0, 1.3)		-0.2 (-0.4, 0.1)	
Residential area																
Urban	Ref	<0.001	Ref	0.255	Ref	0.444	Ref	0.362	Ref	0.002	Ref	0.026	Ref	0.028		
Rural	-52.9 (-81.7, -24.1)		7.7 (1.4, 13.9)		-1.4 (-3.7, 1.0)		14.6 (-51.9, 22.8)		4.7 (-5.4, 14.7)		1.1 (0.4, 1.8)		0.4 (0.0, 0.8)		-0.2 (-0.3, 0.0)	

Regression coefficients (β), 95% confidence intervals (CI), and P-values were derived from multivariable linear regression models examining the independent associations of sex, age, weight status, parental education level, country of origin, and residential area with intake of selected food groups and nutrients, as well as an overall dietary adherence score.

^aFish intake data were derived from the food frequency questionnaire, n = 776

This is likely beneficial for long-term health, as a high wholegrain intake has been shown to improve the plasma lipid profile in children [9] and to be linked to reduced risk of cardiovascular disease and mortality in adults [55]. Wholegrain intakes have increased substantially in Denmark since the early 2000's [26], likely due to initiatives such as the Danish Wholegrain Partnership, which has improved the availability, labeling, and public awareness of wholegrain products. However, wholegrain intake among non-Danish descendants remained particularly low in the present study, with only 21.5% meeting the recommendation. One explanation may be that wholegrain products, such as sour dough rye bread and rolled oats, are distinctive characteristics of the Danish food culture, which may take time to adopt among immigrant populations. The Danish recommendation for wholegrains was raised to 90 g/10 MJ in June 2024 [56], and when compared to this update, 31.3% of the children still adhered.

Surprisingly, 64% of the children adhered to the recommendation of limiting added sugar to <10 E%. This is a marked improvement compared to the 49% and 34% of 4–14-year-olds reported in the Danish national surveys from 2011–2013 and 2003–2008, respectively [12, 57]. Despite efforts to expand the myfood24 database with added sugar values to minimize underestimation, some underreporting cannot be ruled out [58]. Nevertheless, similar downward trends in sugar intake have been observed in the US and Germany since the early 2000s, especially between 2010–2016 [59, 60]. However, unlike in the present study, children's mean and median intakes of added sugar still exceeded the recommendation in these populations [59, 60].

The reported iron intake was somewhat low considering the children's high meat consumption, and when compared to the recommended intake (8.0 mg/d for 7–8-year-olds [7]) and the latest national dietary survey (6.7±1.7 mg/d vs. 8.9±2.1 mg/d and 7.7±2.0 mg/d vs. 9.6±2.4 mg/d in 6–9-year-old girls and boys, respectively [12]). However, both hemoglobin and ferritin levels were within the recommended ranges and only 8% of the children had iron deficiency, suggesting that the intake was adequate for most children. The intake of iron might be underestimated, but the findings may also indicate that the high bioavailability of iron from meat, or increased iron absorption seen at low intakes [61], compensated for the low total intake.

Consistent with previous studies [14–16], adherence to dietary recommendations improved with higher parental education level and decreased with age. Similar to the present study, parental education level was also associated with wholegrain intake in the Danish national survey 20 years ago, although only among girls [14], and the observed tendency for reduced meat intake with longer parental

education aligns with findings from the EU and the UK, particularly for processed meats [62, 63]. These associations may reflect greater health- and environmental awareness among highly educated parents. The decline in adherence to dietary recommendations with increasing age was evident despite a narrow age range in the present study [15, 16]. Early school years may be a period of rapid dietary changes, as children have to adapt to new environments and engage in new social relationships, which may prompt them to align their food preferences with perceived peer norms to ensure social acceptance [64].

The lower overall adherence to dietary recommendations, lower fruit and vegetable intake, and higher sugar intake among rural compared to urban children are consistent with studies from Europe [19] and Canada [20], whereas the higher SFA intake has not previously been reported. Rural children also had a higher wholegrain intake, which may reflect a stronger adherence to the Danish tradition of having wholegrain rye bread for lunch in rural areas, as also seen in Finland [65]. In contrast, an Australian study found that children residing in remote areas had overall healthier eating habits and consumed more fruit, and less takeaway foods and unhealthy snacks [21]. Besides differences in the definitions of urban and rural areas and the remoteness of the rural areas in the studies, the discrepancies could relate to country-specific differences in rural versus urban food environments.

A major strength of the present study was the large sample size of 1094 children which exceeded that of the national Danish dietary surveys [12]. We included participants from a wide geographic area across Denmark, who were representative of the general population in terms of sex, weight status, and residential area. Non-Danish descendants were better represented than in the national surveys [12], but they were still under-represented as were children from shorter educational backgrounds. This may have led to an overestimation of the adherence to the dietary recommendations and limited our ability to detect differences between groups, particularly by origin. The detailed dietary records of the whole diet combined with the fish-specific FFQ allowed for a comprehensive assessment of habitual dietary intake, which was validated against objective biomarkers. A three-day recording period was selected to ensure feasibility, particularly among less advantaged families, but it was not suited for evaluating micronutrient intake [66]. The online recording tool helped structure the recordings, had built-in reminders (e.g. for snacks and toppings), and allowed parents to edit entries during the day, thereby reducing reliance on memory. To support non-Danish families, we provided text instructions in Arabic and English and video instructions in English, and the software interface was available in 10 languages. However, the food item list remained in

Danish which may have challenged families with other language backgrounds.

In conclusion, the overall adherence to the Danish FBDG and NNR2023 was relatively high in the examined large group of Danish 6–9-year-old children, but challenges remain particularly in terms of intakes of fruit and vegetables, fish, meat, SFA, and iron. Sociodemographic differences in dietary intake were especially related to children's age, parental education level, and residential area, but not to weight status. These findings highlight the need for early interventions to improve children's dietary quality while also considering strategies for specific sociodemographic groups where improvements are needed the most.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s00394-025-03863-y>.

Acknowledgements The authors would like to thank the study participants and local partners within schools and local communities for fruitful collaborations. We also acknowledge the valuable contribution of other community members, GHK colleagues and students.

Author contributions Conceptualization: RFK, UT, CTD. Methodology: FH, AGE, LL, CM, CTD. Formal analysis and investigation: FH, AGE, MRL, KDS, RL. Writing of original draft: FH, AGE. Review and editing: LL, CM, MRL, KDS, RL, RFK, UT, CTD. Funding acquisition: CM, RFK, UT, CTD. Supervision: LL, CM, CTD.

Funding Open access funding provided by Copenhagen University. The Generation Healthy Kids study was funded by the Novo Nordisk Foundation (grant no. NNF22SA0077224).

Data availability The datasets analyzed in the current study are not publicly available due to ethical restrictions, but they are available from the corresponding author upon reasonable request.

Declarations

Conflict of interest The authors declare no competing interests.

Ethical approval The study was approved by the Ethics Committee of Southern Denmark (S-20220094).

Consent to participate All children received oral information, and custody holders received both written and oral information about the study before custody holders gave written informed consent for their child to participate.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright

holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Nguyen AN, Jen V, Jaddoe VWV et al (2020) Diet quality in early and mid-childhood in relation to trajectories of growth and body composition. *Clin Nutr* 39:845–852. <https://doi.org/10.1016/j.clnu.2019.03.017>
2. Cohen JFW, Gorski MT, Gruber SA et al (2016) The effect of healthy dietary consumption on executive cognitive functioning in children and adolescents: a systematic review. *Br J Nutr* 116:989–1000
3. Funtikova AN, Navarro E, Bawaked RA et al (2015) Impact of diet on cardiometabolic health in children and adolescents. *Nutr J* 14:118. <https://doi.org/10.1186/s12937-015-0107-z>
4. USDA (2020) Dietary Guidelines for Americans 2020 - 2025
5. Public Health England (2018) The Eatwell Guide
6. Danish Veterinary and Food Administration (2021) The Official Dietary Guidelines - good for health and climate. Glostrup, Denmark
7. Blomhoff R, Andersen R, Arnesen EK et al (2023) Nordic Nutrition Recommendations 2023. Denmark, Copenhagen
8. Vuholm S, Rantanen JM, Teisen MN et al (2019) Effects of oily fish intake on cardiometabolic markers in healthy 8- to 9-year-old children: the FiSK Junior randomized trial. *Am J Clin Nutr* 110:1296–1305. <https://doi.org/10.1093/ajcn/nqz233>
9. Madsen MTB, Landberg R, Nielsen DS et al (2024) Effects of wholegrain compared to refined grain intake on cardiometabolic risk markers, gut microbiota, and gastrointestinal symptoms in children: a randomized crossover trial. *Am J Clin Nutr* 119:18–28. <https://doi.org/10.1016/j.ajcnut.2023.10.025>
10. Damsgaard CT, Dalskov SM, Laursen RP et al (2014) Provision of healthy school meals does not affect the metabolic syndrome score in 8–11-year-old children, but reduces cardiometabolic risk markers despite increasing waist circumference. *Br J Nutr* 112:1826–1836. <https://doi.org/10.1017/S0007114514003043>
11. Siddiqui NZ, Nguyen AN, Santos S, Voortman T (2022) Diet quality and cardiometabolic health in childhood: the Generation R Study. *Eur J Nutr* 61:729–736. <https://doi.org/10.1007/s00394-021-02673-2>
12. Pedersen AN, Christensen T, Matthiessen J, et al (2015) Danskernes kostvaner 2011–2013 Hovedresultater [Dietary Habits in Denmark 2011–2013 Main results]. DTU Fødevareinstituttet, Afdeling for Ernæring, Søborg, Denmark.
13. EFSA Panel on Dietetic Products N and A (NDA) (2013) Scientific Opinion on nutrient requirements and dietary intakes of infants and young children in the European Union. *EFSA J* 11:3408. <https://doi.org/10.2903/j.efsa.2013.3408>
14. Groth MV, Christensen LM, Knudsen VK, et al (2013) Sociale forskelle - Børns kostvaner, fysiske aktivitet og overvægt & voksnes kostvaner [Social differences in children's dietary habits, physical activity and overweight and adult's dietary habits]
15. Golley RK, Hendrie GA, McNaughton SA (2011) Scores on the dietary guideline index for children and adolescents are associated with nutrient intake and socio-economic position but not adiposity. *J Nutr* 141:1340–1347. <https://doi.org/10.3945/jn.110.136879>
16. Banfield EC, Liu Y, Davis JS et al (2016) Poor adherence to US dietary guidelines for children and adolescents in the National Health and Nutrition Examination Survey population. *J Acad Nutr Diet* 116:21–27. <https://doi.org/10.1016/j.jand.2015.08.010>
17. Buckland G, Northstone K, Emmett PM, Taylor CM (2023) Adherence to UK dietary guidelines in school-aged children from

- the Avon Longitudinal Study of Parents and Children (ALSPAC) cohort. *Br J Nutr* 130:454–466. <https://doi.org/10.1017/S0007114522003336>
18. Arriscado D, Muros JJ, Zabala M, Dalmau JM (2014) Factors associated with low adherence to a Mediterranean diet in healthy children in northern Spain. *Appetite* 80:28–34. <https://doi.org/10.1016/j.appet.2014.04.027>
 19. Heinen MM, Bel-Serrat S, Kelleher CC et al (2021) Urban and rural differences in frequency of fruit, vegetable, and soft drink consumption among 6–9-year-old children from 19 countries from the WHO European region. *Obes Rev* 22:e13207. <https://doi.org/10.1111/obr.13207>
 20. Simen-Kapeu A, Kuhle S, Veugeliers PJ (2010) Geographic differences in childhood overweight, physical activity, nutrition and neighbourhood facilities: implications for prevention. *Can J Public Health* 101:128–132
 21. Jacobs J, Strugnell C, Becker D et al (2023) Understanding weight status and dietary intakes among Australian school children by remoteness: a cross-sectional study. *Public Health Nutr* 26:1185–1193. <https://doi.org/10.1017/S1368980023000198>
 22. Jennings A, Welch A, van Sluijs EMF et al (2011) Diet quality is independently associated with weight status in children aged 9–10 years. *J Nutr* 141:453–459. <https://doi.org/10.3945/jn.110.131441>
 23. Thomsen LT, Schmidt-Persson J, Damsgaard CT et al (2024) Generation Healthy Kids: Protocol for a cluster-randomized controlled trial of a multi-component and multi-setting intervention to promote healthy weight and wellbeing in 6–11-year-old children in Denmark. *PLoS ONE* 19:e0308142. <https://doi.org/10.2196/rs.3.rs-3423677/v1>
 24. Albar SA, Alwan NA, Evans CEL et al (2016) Agreement between an online dietary assessment tool (myfood24) and an interviewer-administered 24-h dietary recall in British adolescents aged 11–18 years. *Br J Nutr* 115:1678–1686. <https://doi.org/10.1017/S0007114516000593>
 25. Koch SAJ, Conrad J, Cade JE et al (2021) Validation of the web-based self-administered 24-h dietary recall myfood24-Germany: comparison with a weighed dietary record and biomarkers. *Eur J Nutr* 60:4069–4082. <https://doi.org/10.1007/s00394-021-02547-7>
 26. Mejborn H, Biltoft-Jensen A, Trolle E, Tetens I (2008) Fuldkorn - Definition og vidensgrundlag for anbefaling af fuldkornsindtag i Danmark [Wholegrain - Definition and scientific background for recommendations of wholegrain intake in Denmark]. Søborg, Denmark
 27. DTU National Food Institute (2024) Fødevaredata, version 5.2 [National food database]. The Technical University of Denmark, <http://www.frida.fooddata.dk>. Accessed June 2024.
 28. Black AE (2000) Critical evaluation of energy intake using the Goldberg cut-off for energy intake: basal metabolic rate. A practical guide to its calculation, use and limitations. *Int J Obes* 24:1119–1130
 29. Henry C (2005) Basal metabolic rate studies in humans: measurement and development of new equations. *Public Health Nutr* 8:1133–1152. <https://doi.org/10.1079/phn2005801>
 30. Overvad K, Tjønneland A, Haraldsdóttir J et al (1991) Development of a semiquantitative food frequency questionnaire to assess food, energy and nutrient intake in Denmark. *Int J Epidemiol* 20:900–905
 31. Tjønneland A, Overvad K, Haraldsdóttir J et al (1991) Validation of a semiquantitative food frequency questionnaire developed in Denmark. *Int J Epidemiol* 20:906–912
 32. Vuholm S, Teisen MN, Buch NG et al (2020) Is high oily fish intake achievable and how does it affect nutrient status in 8–9-year-old children? the FiSK Junior trial. *Eur J Nutr* 59:1205–1218. <https://doi.org/10.1007/s00394-019-01981-y>
 33. Ygil KH (2013) Mål, vægt og portionsstørrelser på fødevarer [Dimensions, weight and portion sizes of foods]
 34. Shams-White MM, Pannucci TE, Lerman JL et al (2023) Healthy eating index-2020: review and update process to reflect the dietary guidelines for Americans, 2020–2025. *J Acad Nutr Diet* 123:1280–1288. <https://doi.org/10.1016/j.jand.2023.05.015>
 35. Cole TJ, Lobstein T (2012) Extended international (IOTF) body mass index cut-offs for thinness, overweight and obesity. *Pediatr Obes* 7:284–294. <https://doi.org/10.1111/j.2047-6310.2012.00064.x>
 36. Metherel AH, Armstrong JM, Patterson AC, Stark KD (2009) Assessment of blood measures of n-3 polyunsaturated fatty acids with acute fish oil supplementation and washout in men and women. *Prostaglandins Leukot Essent Fatty Acids* 81:23–29. <https://doi.org/10.1016/j.plefa.2009.05.018>
 37. Metherel AH, Hogg RC, Buzikievich LM, Stark KD (2013) Butylated hydroxytoluene can protect polyunsaturated fatty acids in dried blood spots from degradation for up to 8 weeks at room temperature. *Lipids Health Dis* 12:22. <https://doi.org/10.1186/1476-511X-12-22>
 38. Harris WS, von Schacky C (2004) The Omega-3 Index: a new risk factor for death from coronary heart disease? *Prev Med (Baltim)* 39:212–220. <https://doi.org/10.1016/j.ypmed.2004.02.030>
 39. Stark KD, Aristizabal Henao JJ, Metherel AH, Pilote L (2016) Translating plasma and whole blood fatty acid compositional data into the sum of eicosapentaenoic and docosahexaenoic acid in erythrocytes. *Prostaglandins Leukot Essent Fatty Acids* 104:1–10. <https://doi.org/10.1016/j.plefa.2015.11.002>
 40. Ross AB, Svelander C, Savolainen OI et al (2016) A high-throughput method for liquid chromatography-tandem mass spectrometry determination of plasma alkylresorcinols, biomarkers of whole grain wheat and rye intake. *Anal Biochem* 499:1–7. <https://doi.org/10.1016/j.ab.2015.12.023>
 41. Andersen MB, Thorsted A, Jezek AH, et al (2020) Overvægt og svær overvægt blandt danske børn og unge - Forekomst og sociodemografisk fordeling [Overweight and obesity among Danish children and adolescents - Prevalence and sociodemographic distribution]
 42. Statistics Denmark (2024) StatBank Denmark: Income and education in the family. INDKF107 2023. <https://www.statistikbanken.dk/statbank5a/default.asp?w=2560> (accessed January 2025)
 43. Statistics Denmark (2024) StatBank Denmark: The population, sex, age, and origin quarterly. FOLK1E 2023 Q4. <https://www.statistikbanken.dk/statbank5a/default.asp?w=2560> (accessed January 2025)
 44. Statistics Denmark (2024) The population, age, city size, and time. BY2 2023. <https://www.statistikbanken.dk/by2> (accessed January 2025)
 45. Andersen R, Biltoft-Jensen A, Christensen T et al (2015) What do Danish children eat, and does the diet meet the recommendations? Baseline data from the OPUS School Meal Study. *J Nutr Sci* 4:e29. <https://doi.org/10.1017/jns.2015.17>
 46. Verduci E, Bronsky J, Embleton N et al (2021) Role of dietary factors, food habits, and lifestyle in childhood obesity development: a position paper from the European Society for Paediatric Gastroenterology, Hepatology and Nutrition Committee on Nutrition. *J Pediatr Gastroenterol Nutr* 72:769–783. <https://doi.org/10.1097/MPG.0000000000003075>
 47. Handeland K, Skotheim S, Baste V et al (2018) The effects of fatty fish intake on adolescents' nutritional status and associations with attention performance: results from the FINS-TEENS randomized controlled trial. *Nutr J* 17:30. <https://doi.org/10.1186/s12937-018-0328-z>
 48. Hysing M, Kvestad I, Kjelleveid M et al (2018) Fatty fish intake and the effect on mental health and sleep in preschool children in

- FINS-KIDS, a randomized controlled trial. *Nutrients* 10:1478. <https://doi.org/10.3390/nu10101478>
49. Damsgaard CT, Eidner MB, Stark KD et al (2014) Eicosapentaenoic acid and docosahexaenoic acid in whole blood are differentially and sex-specifically associated with cardiometabolic risk markers in 8-11-year-old Danish children. *PLoS ONE* 9:e109368. <https://doi.org/10.1371/journal.pone.0109368>
 50. Sørensen LB, Damsgaard CT, Dalskov S-M et al (2015) Diet-induced changes in iron and n-3 fatty acid status and associations with cognitive performance in 8-11-year-old Danish children: secondary analyses of the Optimal Well-Being, Development and Health for Danish Children through a Healthy New Nordic Diet School Meal Study. *Br J Nutr* 114:1623–1637. <https://doi.org/10.1017/S0007114515003323>
 51. Teisen MN, Vuholm S, Niclausen J et al (2020) Effects of oily fish intake on cognitive and socioemotional function in healthy 8–9-year-old children: the FiSK Junior randomized trial. *Am J Clin Nutr* 112:74–83. <https://doi.org/10.1093/ajcn/nqaa050>
 52. Livingstone KM, Burton M, Brown AK, McNaughton SA (2020) Exploring barriers to meeting recommendations for fruit and vegetable intake among adults in regional areas: a mixed-methods analysis of variations across socio-demographics. *Appetite* 153:104750. <https://doi.org/10.1016/j.appet.2020.104750>
 53. Mann KD, Pearce MS, McKeivith B et al (2015) Low whole grain intake in the UK: results from the National Diet and Nutrition Survey rolling programme 2008–11. *Br J Nutr* 113:1643–1651
 54. Miller KB (2020) Review of whole grain and dietary fiber recommendations and intake levels in different countries. *Nutr Rev* 78:29–36
 55. Aune D, Keum N, Giovannucci E et al (2016) Whole grain consumption and risk of cardiovascular disease, cancer, and all cause and cause specific mortality: Systematic review and dose-response meta-analysis of prospective studies. *BMJ* 353:i2716. <https://doi.org/10.1136/bmj.i2716>
 56. Trolle E, Christensen LM, Lassen AD (2024) Opdatering af det faglige grundlag for De officielle Kostråd i forhold til Nordiske Næringsstofanbefalinger 2023 [An update of the scientific background for the Official Dietary Guidelines in relation to the Nordic Nutrient Recommendations 2023]
 57. Rothausen BW, Matthiessen J, Hoppe C et al (2012) Differences in Danish children's diet quality on weekdays v. weekend days. *Public Health Nutr* 15:1653–1660. <https://doi.org/10.1017/S1368980012002674>
 58. Macdiarmid J, Blundell J (1998) Assessing dietary intake: who, what and why of under-reporting. *Nutr Res Rev* 11:231–253. <https://doi.org/10.1079/nrr19980017>
 59. Welsh JA, Sharma AJ, Grellinger L, Vos MB (2011) Consumption of added sugars is decreasing in the United States. *Am J Clin Nutr* 94:726–734. <https://doi.org/10.3945/ajcn.111.018366>
 60. Perrar I, Schmitting S, Della Corte KW et al (2020) Age and time trends in sugar intake among children and adolescents: results from the DONALD study. *Eur J Nutr* 59:1043–1054. <https://doi.org/10.1007/s00394-019-01965-y>
 61. Cook JD, Lipschitz DA, Miles LE, Finch CA (1974) Serum ferritin as a measure of iron stores in normal subjects. *Am J Clin Nutr* 27:681–687
 62. Fernández-Alvira JM, Mouratidou T, Bammann K et al (2013) Parental education and frequency of food consumption in European children: the IDEFICS study. *Public Health Nutr* 16:487–498. <https://doi.org/10.1017/S136898001200290X>
 63. Cribb VL, Jones LR, Rogers IS et al (2011) Is maternal education level associated with diet in 10-year-old children? *Public Health Nutr* 14:2037–2048. <https://doi.org/10.1017/S136898001100036X>
 64. Landwehr SC, Hartmann M (2024) Is it all due to peers? The influence of peers on children's snack purchase decisions. *Appetite* 192:107111. <https://doi.org/10.1016/j.appet.2023.107111>
 65. Prättälä R, Helasoja V, Mykkänen H (2001) The consumption of rye bread and white bread as dimensions of health lifestyles in Finland. *Public Health Nutr* 4:813–819. <https://doi.org/10.1079/pbn2000120>
 66. Basiotis PP, Welsh SO, Cronin FJ et al (1987) Number of days of food intake records required to estimate individual and group nutrient intakes with defined confidence. *J Nutr* 117:1638–1641