



CHALMERS
UNIVERSITY OF TECHNOLOGY

Scalable quantum simulator with an extended gate set in giant atoms

Downloaded from: <https://research.chalmers.se>, 2026-02-28 01:22 UTC

Citation for the original published paper (version of record):

Chen, G., Frisk Kockum, A. (2026). Scalable quantum simulator with an extended gate set in giant atoms. QUANTUM, 10

N.B. When citing this work, cite the original published paper.

Scalable quantum simulator with an extended gate set in giant atoms

Guangze Chen and Anton Frisk Kockum

Department of Microtechnology and Nanoscience, Chalmers University of Technology, 41296 Gothenburg, Sweden

Quantum computation and quantum simulation require a versatile gate set to optimize circuit compilation for practical applications. However, existing platforms are often limited to specific gate types or rely on parametric couplers to extend their gate set, which compromises scalability. Here, we propose a scalable quantum simulator with an extended gate set based on giant-atom three-level systems, which can be implemented with superconducting circuits. Unlike conventional small atoms, giant atoms couple to the environment at multiple points, introducing interference effects that allow exceptional tunability of their interactions. By leveraging this tunability, our setup supports both CZ and iSWAP gates through simple frequency adjustments, eliminating the need for parametric couplers. This dual-gate capability enhances circuit efficiency, reducing the overhead for quantum simulation. As a demonstration, we showcase the simulation of spin dynamics in dissipative Heisenberg XXZ spin chains, highlighting the setup's ability to tackle complex open quantum many-body dynamics. Finally, we discuss how a two-dimensional extension of our system could enable fault-tolerant quantum computation, paving the way for a universal quantum processor.

1 Introduction

Scalable universal quantum simulators are powerful tools for exploring complex quantum systems, including many-body physics in condensed matter physics and open quantum many-body dynamics [1–3]. A universal gate set for a quantum simulator or quantum computer can be realized using a complete set of single-qubit gates combined with an entangling two-qubit gate, such as iSWAP or CZ [4]. Most existing quantum simulators are optimized for implementing only specific two-qubit gates [3, 5, 6], limiting their versatility. Expanding the available gate set beyond the bare minimum facilitates more efficient quantum circuit compilation [7–9], reducing circuit depth and

improving performance. Notably, having access to both iSWAP and CZ gates enables any Clifford operation to be performed using single-qubit gates and no more than two two-qubit gates [10]. Moreover, introducing tunable qubit decay would open the door to simulating open quantum many-body dynamics. However, achieving a larger gate set typically requires additional resources, such as parametric couplers [10–18], other coupling elements [19–24], or complicated drive schemes [25–31], which limit the feasibility of building large-scale quantum simulators.

In this article, we show how giant artificial atoms [32] can be used to build a scalable quantum simulator for open quantum systems with both iSWAP and controlled-phase (CZ_φ) two-qubit gates in the gate set. Unlike traditional small atoms, which couple to their environment at a single point, giant atoms couple at multiple discrete points, often separated by wavelengths. The consequences of having multiple coupling points have been explored in many articles in recent years, both theoretically in, e.g., Refs. [32–55] and in experiments mostly using superconducting circuits, e.g., in Refs. [56–67]. The main point is that the multiple coupling points produce interference effects, enabling frequency-dependent control of relaxation rates [33, 64] and qubit-qubit interaction strengths [35]. This tunability makes giant atoms well-suited for implementing diverse gate operations without overhead like parametric couplers. For instance, iSWAP gates and tunable decays can be achieved in giant-atom systems by tuning qubit frequencies [61], enabling applications such as the Trotterized simulation of quantum Zeno dynamics in open quantum systems [68]. However, an extended gate set including CZ operations has not been studied previously in giant-atom setups.

Here, to enable the execution of CZ_φ gates in addition to iSWAP gates, we introduce three-level giant atoms with additional coupling points to eliminate unwanted interactions and reduce errors. We demonstrate the ability of our simulator built from such systems to simulate the dynamics of the dissipative XXZ model, illustrating the advantages of an extended gate set in reducing simulation errors. Additionally, we propose a two-dimensional extension of our quantum simulator, which enables the execution of long-range two-qubit operations. This extension further allows the implementation of surface codes for quantum er-

Guangze Chen: guangze@chalmers.se

Anton Frisk Kockum: anton.frisk.kockum@chalmers.se

arXiv:2503.04537v3 [quant-ph] 20 Jan 2026

ror correction [69], thereby supporting fault-tolerant quantum computation and positioning our setup as a potential universal quantum processor.

The rest of this article is organized as follows. In Section 2, we put forward a setup with two three-level giant atoms as the building block of our scalable quantum simulator. We demonstrate how this building block can realize both iSWAP and CZ gates, as well as the more general R_{XY} and controlled-phase (CZ_φ) gates, through simple frequency tuning of the giant atoms. We then analyze the average gate fidelity of these gates under realistic noise conditions, showing that state-of-the-art techniques yield fidelities of $\geq 98.8\%$ for both iSWAP and CZ operations; these fidelities can quite easily be increased by coupling the giant atoms more strongly to the waveguide. Then, in Section 3, we combine such building blocks into a scalable simulator architecture where the two-giant-atom structure is repeated to form a one-dimensional chain. We provide a protocol for tuning the atomic frequencies to implement nearest-neighbor R_{XY} and CZ_φ gates across this simulator, demonstrating the scalability of the extended gate set.

To highlight the simulator's potential, we showcase in Section 4 its simulation of spin dynamics in a dissipative XXZ spin chain [70–73], illustrating its ability to handle complex open quantum many-body system. To further enhance the capability of our simulator to execute long-range two-qubit gates, we propose in Section 5 a two-dimensional extension of it. We discuss how long-range two-qubit gates can be operated in such a setup by tuning the frequencies of the giant atoms. In particular, this extension allows to perform surface codes, enabling fault-tolerant quantum computation, and offering our setup as a universal quantum processor. We provide a discussion on the physical realization of our setup and the influence of non-Markovian effects in Section 6, followed by concluding remarks in Section 7. Some additional details on the error analysis for the quantum simulation in Section 4 are provided in Appendix A.

2 Two-qubit gates with giant atoms

We begin by studying a two-giant-atom setup that allows us to perform both iSWAP and CZ gates by simple frequency tuning. This setup serves as the building block for our scalable giant-atom-based simulator. The giant atoms we consider are Ξ -type three-level systems with an anharmonicity $\chi_k = \omega_{12,k} - \omega_{01,k} < 0$, where $\omega_{ij,k}$ is the transition frequency between states $|i\rangle$ and $|j\rangle$ in atom k ; for simplicity, we use in the following the notation $\omega_{01,k} \equiv \omega_k$. Such negative anharmonicity and ladder-type level structure are typical for superconducting transmon qubits [74], which is the most common platform for experiments on giant atoms so far. The giant atoms are coupled to a waveguide at multiple spatially separated points




Figure 1: A two-giant-atom setup for performing both iSWAP and CZ gates. (a) Sketch of the setup. The two giant atoms, with frequencies $\omega_{1,2}$ and detunings $\chi_{1,2}$, are coupled to the waveguide (black line) at multiple points with different coupling strengths γ_{kn} and spacings Δx_n . The coupling points are organized in a braided fashion. (b) Frequency dependence of the individual decay rates $\Gamma_{1,2}$, inter-atomic coupling strength g_{12} , and collective decay rates $\Gamma_{\text{coll},12}$ of the giant atoms. (c,d) The protocol to perform (c) an iSWAP gate and (d) a CZ gate in this setup.

each (with coordinate x_{kn} for the n th coupling point of atom k), and these coupling points are organized in a braided fashion, as shown in Fig. 1(a).

Due to the coupling to the waveguide, the two atoms acquire individual decay rates $\Gamma_{\text{ind},k}(\omega)$, a coherent interaction with strength $g_{jk}(\omega)$ between atoms j and k , and a collective decay rate $\Gamma_{\text{coll},jk}(\omega)$ for atoms j and k . The dependence on the frequency ω of the transition that these rates hold for is a consequence of the interference effects arising from the multiple coupling points of giant atoms. Assuming Markovianity, i.e., that the travel time between coupling points is negligible, the decay rates and interaction strengths are given by [35]

$$\begin{aligned}\Gamma_{\text{ind},k}(\omega) &= \sum_{n=1}^{N_k} \sum_{m=1}^{N_k} \sqrt{\gamma_{kn}\gamma_{km}} \cos \phi_{kn,km}(\omega), \\ g_{jk}(\omega) &= \sum_{n=1}^{N_j} \sum_{m=1}^{N_k} \frac{\sqrt{\gamma_{jn}\gamma_{km}}}{2} \sin \phi_{jn,km}(\omega), \\ \Gamma_{\text{coll},jk}(\omega) &= \sum_{n=1}^{N_j} \sum_{m=1}^{N_k} \sqrt{\gamma_{jn}\gamma_{km}} \cos \phi_{jn,km}(\omega),\end{aligned}\quad (1)$$

where N_k is the number of coupling points of atom k , γ_{kn} is the coupling strength at the n th coupling point of atom k , $\phi_{jn,km}(\omega) = \omega \Delta x_{jn,km} / v$ is the accumulated phase between the coupling points, $\Delta x_{jn,km} = |x_{jn} - x_{km}|$ the distance between the coupling points, and v the speed of light in the waveguide, which we assume to have linear dispersion. We note that using nonlinear structured waveguides [48] would not qualitatively change our results.

The interference effects in the giant atoms result in decoherence-free frequencies where the atomic decay into the waveguide vanishes. To illustrate this, we consider that all the coupling strengths are identical (γ), and $\Delta x_1/2 = \Delta x_7/2 = \Delta x_2 = \Delta x_3 = \Delta x_4 = \Delta x_5 = \Delta x_6 = \Delta x$ in Fig. 1(a). This leads to a periodic dependence of $\Gamma_{\text{ind},k}(\omega)$, $g_{jk}(\omega)$, and $\Gamma_{\text{coll},jk}(\omega)$ on ω , with a periodicity of $\omega_0 = 2\pi v / \Delta x$. Notable are the decoherence-free frequencies $\omega_{\text{DF},nm} = (n + m/8)\omega_0$ ($n \in \mathcal{N}, m = 1, 2, 3, 5, 6, 7$), at which the interaction g_{12} is non-zero, as shown in Fig. 1(b). These frequencies are essential for performing high-fidelity two-qubit gates.

Moreover, the individual decay rates are minimal within the frequency ranges $[\omega_{\text{DF},n1}, \omega_{\text{DF},n3}]$ and $[\omega_{\text{DF},n5}, \omega_{\text{DF},n7}]$, as highlighted in the grey regions in Fig. 1(b). This behavior results from having four coupling points for each giant atom; the regions can be extended by having more coupling points. As we will see in the following, this property reduces qubit decay during gate operations and provides a broad operational frequency range. To enable both iSWAP and CZ gates with high fidelity, we set $\chi_1 = -\chi = -\omega_0/8$.

We now show how different gates can be performed with this setup. The first thing to notice is that when

the two atoms are at different decoherence-free frequencies, they will have negligible coupling due to the detuning between them, and thus the system remains in a steady state. On top of this, single-qubit decay can be implemented by tuning the qubit's frequency to a non-decoherence-free value, allowing controlled dissipation. This feature is particularly beneficial for simulating open quantum systems, as explored in Section 4.

2.1 iSWAP gate with giant atoms

Decoherence-free interactions between braided giant atoms facilitate the realization of the iSWAP gate [61]. The action of an iSWAP gate leaves the states $|00\rangle$ and $|11\rangle$ unchanged, while the states $|01\rangle$ and $|10\rangle$ are swapped and acquire a phase factor i . This outcome is achieved in our setup by setting $\omega_1 = \omega_2 = \omega_{\text{DF},nm}$ such that the $|0\rangle \leftrightarrow |1\rangle$ transitions of the two atoms are resonant [Fig. 1(c)].

We now analyze the fidelity of the giant-atom iSWAP gate in realistic cases, when the qubits are subject to decay and dephasing. To be concrete, we consider the case of transmon qubits, where the decay and dephasing of the $|2\rangle$ state are twice those for the $|1\rangle$ state [74]. Additionally, for transmon qubits, a direct decay from $|2\rangle$ to $|0\rangle$ is prevented due to a vanishingly small matrix element connecting the two states. Thus, the Lindblad dissipators for qubit decay and dephasing are

$$L_- = \sqrt{\Gamma_{\text{ind}} + \Gamma_{\text{ex}}} (|1\rangle\langle 0| + \sqrt{2}|2\rangle\langle 1|), \quad (2)$$

$$L_\phi = \sqrt{2\Gamma_\phi} (|1\rangle\langle 1| + 2|2\rangle\langle 2|), \quad (3)$$

respectively, where Γ_{ex} is the extra decay rate of the qubit to environments other than the waveguide, and Γ_ϕ is the qubit's dephasing rate. For simplicity, we further assume that the two qubits have the same decay and dephasing rates. We also neglect the effect of tuning the atoms in and out of the conditions enabling the gate; we return to that topic in Section 4.

Taking into account these decoherence processes, we show in Fig. 2(a)-(b) the effect of decay and dephasing on the average process fidelity [75] of the iSWAP gate,

$$F_{\text{iSWAP}} = \left[\text{tr} \left(\sqrt{\sqrt{\Phi} \Phi_0 \sqrt{\Phi}} \right) \right]^2, \quad (4)$$

where Φ and Φ_0 are the Choi matrices [76] of our gate and the perfect iSWAP gate, respectively. The Choi matrix Φ of a process \mathcal{E} is defined as

$$\Phi = \sum_n |n\rangle\langle n| \otimes \mathcal{E}(|n\rangle\langle n|). \quad (5)$$

where n runs over the basis states of the computational subspace. We observe an expected linear dependence [77] of the process fidelity on both the decay




Figure 2: Average process fidelity of two-qubit (a,b) iSWAP (c,d) CZ and (e) CZ_φ gates performed with the setup in Fig. 1(a), as a function of qubit decay rate Γ_{ex} and dephasing rate Γ_ϕ . Here, $g \approx 2.1\gamma$ is the qubit-qubit coupling strength used in the gates.

and dephasing: $F_{\text{iSWAP}} \approx 1 - 1.57\Gamma_{\text{ex}}/g - 1.57\Gamma_\phi/g$, where g is the qubit-qubit coupling. This means the average gate fidelity of the iSWAP gate is given by $F_{\text{ave, iSWAP}} \approx 1 - 1.26\Gamma_{\text{ex}}/g - 1.26\Gamma_\phi/g$, in agreement with previous results¹. Assuming that the iSWAP gate is performed with $\omega_1 = \omega_2 = \omega_{\text{DF},n3}$, which yields $g \approx 2.1\gamma$, and taking an experimentally accessible value of $\gamma/(2\pi) = 2$ MHz, $\Gamma_{\text{ex}} = 0.02$ MHz $\approx 0.76 \cdot 10^{-3}g$ and $\Gamma_\phi = 0.05$ MHz $\approx 1.89 \cdot 10^{-3}g$ [78–84], the average gate fidelity is 99.67%. To achieve even higher fidelity, we can increase the qubit-waveguide coupling γ and consequently the qubit-qubit coupling g , such that the gate time becomes smaller; for example, with $\gamma/(2\pi) = 4$ MHz, an average gate fidelity of 99.83% can be achieved.

As the iSWAP gate in our setup is performed via the XY interactions between the giant atoms, an $R_{\text{XY}}(\theta)$ gate can in general be performed with our setup. This can be achieved by setting the interacting time $\tau = \theta/g$ for $g > 0$ and $\tau = (2\pi - \theta)/|g|$ for $g < 0$. In particular, the iSWAP gate is $R_{\text{XY}}(\pi/2)$.

2.2 CZ and CZ_φ gates with giant atoms

Our setup also allows the implementation of CZ and CZ_φ gates between giant atoms. The CZ gate, which adds a phase of π to the $|11\rangle$ state of the two-qubit system and leaves all other states unchanged, can be achieved by bringing the population of the $|11\rangle$ state to $|02\rangle$ or $|20\rangle$ and back [85]. This process requires a resonant transition between these two states. In our setup, this resonance is achieved by having $\omega_2 = \omega_1 + \chi_1$ (or $\omega_1 = \omega_2 + \chi_2$) such that there is an interaction between the $|11\rangle$ and $|20\rangle$ (or $|02\rangle$) states. To ensure that at the same time no extra decay happens for the involved levels, we can take $\omega_1 = \omega_{\text{DF},n2}$

¹The average gate fidelity and the process fidelity are related as [75]: $F_{\text{ave}} = 1 - dF/(d+1)$ where F_{ave} and F are the average gate and process fidelities, and d is the dimension of the computational space. For two-qubit iSWAP and CZ gates, we take $d = 4$. The average gate fidelity of a two-qubit iSWAP gate is $F_{\text{ave, iSWAP}} \approx 1 - 0.8\Gamma_{\text{ex}}\tau - 0.8\Gamma_\phi\tau$, where τ is the gate time [77]. For two qubits coupled with interaction strength g , the gate time is $\tau = \pi/(2g)$. Combining these equations, we find that it agrees with the equation in the main text.

and $\omega_2 = \omega_{\text{DF},n1}$ [Fig. 1(d)]; this choice yields a qubit-qubit coupling $g \approx 2.1\gamma$. The coupling between the corresponding levels is then $\sqrt{2}g$, where the factor of $\sqrt{2}$ originates from the fact that transmons are close to harmonic oscillators [33, 74].

The average process fidelity of the CZ gate implemented in this way is shown in Fig. 2(c)-(d). Due to the fact that CZ involves higher levels with higher decay and dephasing rates, and that it takes $\sqrt{2}$ longer time than iSWAP for the same coupling strength g , the process fidelity is lower than for iSWAP: $F_{\text{CZ}} \approx 1 - 2.19\Gamma_{\text{ex}}/g - 2.97\Gamma_\phi/g$. Correspondingly, the average gate fidelity is $F_{\text{ave, CZ}} \approx 1 - 1.75\Gamma_{\text{ex}}/g - 2.34\Gamma_\phi/g$. The average gate fidelity shows a stronger dependence on dephasing, in agreement with previous results [86, 87]. Taking a typical value of $\gamma/(2\pi) = 2$ MHz, $\Gamma_{\text{ex}} = 0.02$ MHz and $\Gamma_\phi = 0.05$ MHz, the average gate fidelity is 99.42% for our setup. The gate fidelity can be improved by increasing the qubit-waveguide coupling γ ; for example, with $\gamma/(2\pi) = 4$ MHz, the average gate fidelity becomes 99.71%.

We can also perform a generalized controlled-phase gate CZ_φ by introducing a detuning from the resonance condition used in the CZ gate. Specifically, setting $\omega_2 = \omega_1 + \chi_1 + \Delta$ produces the following Hamiltonian in the subspace spanned by $|11\rangle$ and $|20\rangle$:

$$H_{\text{CZ}_\varphi} = \begin{pmatrix} \Delta/2 & \sqrt{2}g \\ \sqrt{2}g & -\Delta/2 \end{pmatrix}, \quad (6)$$

where $\sqrt{2}g$ is the coupling between $|11\rangle$ and $|20\rangle$. Starting in $|11\rangle$, the shortest evolution time that returns the state to $|11\rangle$ (up to a phase) is $\tau = \pi/g'$, where $g' = \sqrt{2g^2 + \Delta^2}/4$. This yields

$$\exp(iH_{\text{CZ}_\varphi}\pi/g')|11\rangle = \exp(i\varphi)|11\rangle \quad (7)$$

with a Δ -dependent phase φ [19]:

$$\varphi = \pi \left(1 + \frac{\Delta}{\sqrt{8g^2 + \Delta^2}} \right). \quad (8)$$

The detuning Δ in qubit 2 with respect to the decoherence-free frequency results in the decay of its $|1\rangle$ level to the waveguide, which depends on Δ/ω_0 .

The process fidelity of the CZ_φ gate is thus influenced by both the shorter gate time and the decay into the waveguide induced by the detuning.

To analyze the process fidelity for the CZ_φ gate, we consider a realistic situation of $\chi/(2\pi) = 200$ MHz, which yields $\omega_0/(2\pi) = 1.6$ GHz $= 800\gamma$; we also fix the dephasing rate to a value of $\Gamma_\phi = 0.05$ MHz without loss of generality. The process fidelity for different φ and Γ_{ex} is shown in Fig. 2(e). We observe that, for $|\varphi - \pi| < \varphi_c$, where $\varphi_c \approx 0.9\pi$, the process fidelity is sensitive to Γ_{ex} , while for φ close to 0 and 2π it is not. This is because to have φ close to 0 and 2π , the detuning Δ in Eq. (8) will be large, resulting in a large decay into the waveguide that dominates over the intrinsic qubit decay Γ_{ex} . For smaller $|\varphi - \pi|$, the intrinsic qubit decay Γ_{ex} dominates over decay into the waveguide, and thus the fidelity is mostly influenced by Γ_{ex} and the gate time τ . In particular, when $|\varphi - \pi|$ increases, the gate time decreases, and a higher fidelity is obtained. Unlike the CZ gate, whose fidelity increases with γ due to the shorter gate time, the fidelity of the CZ_φ gate does not increase monotonically with γ . While a larger γ speeds up the gate, it also enhances decay of the detuned qubit into the waveguide, which lowers the fidelity. As this decay increases with Δ/ω_0 , a larger ω_0 reduces it and lowers γ_c , allowing high-fidelity CZ_φ gates over a wider range of φ .

We finally note that, taking $\omega_1 = \omega_{\text{DF},n3}$, different two-qubit operations can be achieved by solely tuning ω_2 within $[\omega_{\text{DF},n1}, \omega_{\text{DF},n3}]$: to have no evolution, $\omega_2 = \omega_{\text{DF},n1}$; to have CZ, $\omega_2 = \omega_{\text{DF},n2}$, and to have iSWAP, $\omega_2 = \omega_{\text{DF},n3}$. As the individual decay rates within this frequency range are small, highly tunable two-qubit gates with high fidelity can be realized here. We also note that negative couplings $g < 0$ can be achieved in our setup by switching to the frequency regime of $[\omega_{\text{DF},n5}, \omega_{\text{DF},n7}]$ — this allows to perform inverse operations of $R_{XY}(\theta)$. The setup's versatility and tunability establish it as a robust building block for scalable quantum simulators.

3 Scalable giant-atom-based quantum simulator

The tunability of giant atoms enables controlled single-qubit decay, iSWAP gates, and CZ gates via coupling to a waveguide, forming an extended gate set. All these operations can be realized using the simple structure depicted in Fig. 1(a), making the setup inherently scalable toward a many-body quantum simulator for open quantum systems. In this section, we demonstrate how such a scalable simulator can be constructed, in a configuration optimized for nearest-neighbor two-qubit iSWAP and CZ gates. Additionally, we note that alternative architectures, such as one supporting all-to-all tunable couplings,




Figure 3: Scalable giant-atom-based quantum simulator. (a) The architecture of the quantum simulator, where neighboring giant atoms are coupled to the waveguide in a braided configuration. (b,c) The frequency dependence of individual decay rates Γ_{ind} , coupling strength g , and collective decay rates Γ_{coll} for (b) neighboring and (c) non-neighboring giant atoms.

are feasible using a similar approach as in Ref. [68].

3.1 Setup

The architecture of the scalable simulator is shown in Fig. 3(a), where neighboring giant atoms are coupled to the waveguide in a braided fashion. Compared to Fig. 1(a), two more coupling points per atom have been added to create more decoherence-free frequencies for the performance of two-qubit operations between different neighbors. To minimize individual atom decay within the operational frequency range, the coupling strength at the middle connection points of each atom is set slightly higher than that of the outer ones: $\gamma' = 1.4\gamma$.

The frequency dependence of the coupling strength g , individual decay rates Γ_{ind} , and collective decay rates Γ_{coll} between neighboring and non-neighboring giant atoms, is depicted in Fig. 3(b)-(c). A set of decoherence-free frequencies $\omega_{\text{DF},nm}$ ($n \in \mathcal{N}, m = 1, \dots, 10$) allows coupling between neighboring atoms ($g_{12} \neq 0$), while suppressing unwanted coupling be-

tween non-neighboring atoms ($g_{13} = 0$). This feature is crucial for ensuring that only intended qubits interact during gate operations. The frequency regime $[\omega_{\text{DF},n1}, \omega_{\text{DF},n5}]$, which includes five decoherence-free frequencies, is ideal for operating the two-qubit gates due to minimal individual decay in this range.

The qubits in the simulator are arranged with odd-site qubits placed at fixed frequencies $\omega_{4k-3} = \omega_{\text{DF},n2}$, $\omega_{4k-1} = \omega_{\text{DF},n5}$ ($k \in \mathcal{N}$), where n is chosen such that $\omega_{\text{DF},n2}$ and $\omega_{\text{DF},n5}$ lie in the optimal frequency regime for the qubits. The even-site qubits are tunable, enabling diverse gate operations with their neighbors by adjusting their frequencies. This design minimizes errors from decoherence during frequency tuning, since only half of the qubits require tunability, and then only in a limited range.

3.2 Performing gates

We now discuss how different qubit operations can be executed in the scalable simulator. To maintain a steady state of the simulator, we can set $\omega_{2k} = \omega_{\text{DF},n3}$ such that all atoms are decoherence-free, while neighboring atoms are detuned and will not couple to each other. On top of this, single-qubit decay can be achieved by tuning the targeted qubit to a non-decoherence-free frequency.

To perform iSWAP gates, and in general $R_{XY}(\theta)$ gates, between neighboring atoms, there are two possibilities: (i) $R_{XY}(\theta)$ between qubits $2k-1$ and $2k$ and (ii) $R_{XY}(\theta)$ between qubits $2k$ and $2k+1$. To achieve (i), we set ($k \in \mathcal{N}$) [Fig. 4(a)]

$$\omega_{4k-2} = \omega_{\text{DF},n2}, \quad \omega_{4k} = \omega_{\text{DF},n5}, \quad (9)$$

such that qubits $4k-3$ and $4k-2$ are coupled with a strength of $g'_1 \approx 1.79\gamma$, qubits $4k-1$ and $4k$ are coupled with a strength of $g'_2 \approx 2.05\gamma$, and qubits $2k$ and $2k+1$ are detuned and thus are decoupled. Similarly, (ii) can be achieved with ($k \in \mathcal{N}$) [Fig. 4(b)]

$$\omega_{4k-2} = \omega_{\text{DF},n5}, \quad \omega_{4k} = \omega_{\text{DF},n2}. \quad (10)$$

Similar to iSWAP gates, there are also two possibilities to perform CZ gates: (i) CZ between qubits $2k-1$ and $2k$ and (ii) CZ between qubits $2k$ and $2k+1$. To achieve (i), we set [Fig. 4(c)]

$$\omega_{4k-2} = \omega_{\text{DF},n1}, \quad \omega_{4k} = \omega_{\text{DF},n4}, \quad (11)$$

and similarly for (ii) we set [Fig. 4(d)]

$$\omega_{4k-2} = \omega_{\text{DF},n4}, \quad \omega_{4k} = \omega_{\text{DF},n1}. \quad (12)$$

On top of this, to perform CZ_φ gates, a detuning Δ given by Eq. (8) can be added to the qubits on even sites.

We have thus demonstrated an architecture for a scalable quantum simulator with an extended gate set of $R_{XY}(\theta)$ and CZ_φ gates with giant atoms. Here,




Figure 4: Protocol to perform different two-qubit operations on the giant-atom-based simulator. The qubits' frequencies are tuned to achieve: (a) $R_{XY}(\theta)$ between qubits $2k-1$ and $2k$, (b) $R_{XY}(\theta)$ between qubits $2k$ and $2k+1$, (c) CZ between qubits $2k-1$ and $2k$ and (d) CZ between qubits $2k$ and $2k+1$.

the interference effects mediated by the waveguide not only enable different qubit operations with simple frequency tuning, but also eliminate unwanted couplings between non-neighboring qubits. The extended gate set reduces circuit depth for simulation tasks requiring a combination of these operations, and thus reduces simulation errors. Furthermore, the simulator's controllable qubit decay is uniquely suited for simulating open quantum dynamics. We illustrate these benefits in the next section with a concrete example of quantum simulation of the dynamics of an open quantum system.

4 Application in quantum simulation

The extended gate set of our giant-atom-based simulator makes it versatile for simulating a broad range of open quantum dynamics, such as the dynamics of dissipative XXZ chains [70–73] and the quantum contact process [88–90]. In this section, we illustrate the versatility of our simulator by showcasing the simulation of a dissipative XXZ model for N spins,

$$H = \sum_{k=1}^{N-1} [J(\sigma_k^x \sigma_{k+1}^x + \sigma_k^y \sigma_{k+1}^y) + J_z \sigma_k^z \sigma_{k+1}^z], \quad (13)$$

where J and J_z are coupling strengths and we add dissipation on the last site: $L = \sqrt{\Gamma} \sigma_N^-$. Here $\sigma^{x,y,z}$ are Pauli matrices and $\sigma^- = \sigma^x - i\sigma^y$.

To simulate the dynamics given by the above model, we employ the Trotter-Suzuki decomposition [91, 92]

$$\exp(\mathcal{L}t) = \left[\prod_{j=1}^n \exp(\mathcal{L}_j t/l) \right]^l + O\left(\frac{t^2}{l}\right), \quad (14)$$

where $\mathcal{L}[\rho] = -i[H, \rho] + L\rho L^\dagger - \frac{1}{2}(L^\dagger L\rho + \rho L^\dagger L)$ is the Liouvillian superoperator governing the dynamics of this model, l is the number of Trotter steps, and $\mathcal{L} = \sum_j \mathcal{L}_j$. We divide \mathcal{L} into components $\mathcal{L}_{1,2,3,4}[\rho] = -i[H_{1,2,3,4}, \rho]$ and $\mathcal{L}_5[\rho] = L\rho L^\dagger - \frac{1}{2}(L^\dagger L\rho + \rho L^\dagger L)$, where

$$\begin{aligned} H_1 &= J \sum_{k=1}^{N/2} (\sigma_{2k-1}^x \sigma_{2k}^x + \sigma_{2k-1}^y \sigma_{2k}^y), \\ H_2 &= J_z \sum_{k=1}^{N/2} \sigma_{2k-1}^z \sigma_{2k}^z, \\ H_3 &= J \sum_{k=1}^{N/2-1} (\sigma_{2k}^x \sigma_{2k+1}^x + \sigma_{2k}^y \sigma_{2k+1}^y), \\ H_4 &= J_z \sum_{k=1}^{N/2-1} \sigma_{2k}^z \sigma_{2k+1}^z. \end{aligned} \quad (15)$$

This decomposition enables the simulation of the original model by sequentially applying the dynamics of each component. In particular, $\exp(-iH_1 t/l)$ and $\exp(-iH_3 t/l)$ correspond to $R_{XY}(\theta)$ gates between the corresponding qubits with $\theta = -2Jt/l$. Similarly, $\exp(-iH_2 t/l)$ and $\exp(-iH_4 t/l)$ yield $R_{ZZ}(\varphi_0)$ gates with $\varphi_0 = -J_z t/l$. This can be achieved with a CZ_φ gate with $\varphi = -4J_z t/l$ and two single-qubit $R_z(-\varphi/4)$ gates. Lastly, \mathcal{L}_5 represents single-qubit decay at the chain's end. We note that additional single-qubit R_z gates are required to compensate for phase shifts from frequency tuning before the performance of $R_{XY}(\theta)$ gates. Combining all these considerations, the protocol for a single Trotter step is shown in Fig. 5(a).

As detailed in Section 3, the circuit in Fig. 5(a) can be implemented by simply adjusting the qubit




Figure 5: Protocol to simulate the dynamics of the dissipative XXZ spin chain using the quantum simulator in Fig. 3(a). (a) The operations that need to be performed within a single Trotter step to simulate the dynamics. (b) The protocol to tune the frequencies of the giant atoms to achieve the operations in (a).

frequencies. Specifically, the frequencies of odd site qubits are fixed as $\omega_{4k-3} = \omega_{DF,n2}$ and $\omega_{4k-1} = \omega_{DF,n5}$. During the execution of the $R_{XY}(\theta)$ gate between qubits $4k-2$ and $4k-3$, the frequency of qubit $4k-2$ is tuned to $\omega_{DF,n2}$, and the system evolves for a duration of $t_1 = \theta/g'_1$, where g'_1 denotes the coupling strength between these qubits. Concurrently, the $R_{XY}(\theta)$ gate between qubits $4k-1$ and $4k$ is implemented by tuning the frequency of qubit $4k$ to $\omega_{DF,n5}$ and evolving for $t'_1 = \theta/g'_2$, where g'_2 represents the coupling strength between qubits $4k-1$ and $4k$. Given that $t'_1 < t_1$, the frequency tuning for qubit $4k$ starts later than that for qubit $4k-2$ by a time interval $t_0 = t_1 - t'_1$ [Fig. 5(b)].

Similarly, other gates in the circuit of Fig. 5(a) are executed by frequency tuning, enabling the full circuit for a single Trotter step to be realized as shown in Fig. 5(b). In this process, $t_2 = \varphi/g'_1$ denotes the duration for executing the CZ gate on qubits $4k-1$ and $4k$, $t'_0 = \varphi/g'_1 - \varphi/g'_2$ represents the time offset between CZ gates on different qubits, t_3 is the duration for a single-qubit R_Z gate, and $t_4 = \Gamma t/(\Gamma_0)$ is the simulation time for single-qubit decay, with $\Gamma_0 \approx 1.36\gamma$ being the decay rate to the waveguide when the qubit frequency is set to a value $\omega_{\text{decay}} \in [\omega_{DF,n2}, \omega_{DF,n3}]$.

We now present concrete simulation results using typical parameter values: $\gamma/(2\pi) = 2$ MHz, $\Gamma_{\text{ex}} = 0.02$ MHz and $\Gamma_\phi = 0.05$ MHz. ω_0 is set to $\omega_0/(2\pi) = 3.2$ GHz such that $(\omega_{DF,n2} - \omega_{DF,n1})/(2\pi) \approx 200$ MHz is achievable as the detuning of the qubits. A con-




Figure 6: Simulation of the dynamics of the dissipative XXZ spin chain [Eq. (13)] with $N = 4$ sites, using our giant-atom-based quantum simulator. In (a) and (b), $l = 30$ Trotter steps were used; in (c) and (d), $l = 10$ Trotter steps were used. The results faithfully capture the slowdown of the spin transport as J_z increases.

servative single-qubit gate time of 30 ns is used [93]. The time required for tuning qubit frequencies is negligible (~ 1 ns) compared to the simulation duration, as tuning rates of ~ 0.1 GHz/ns are achievable [94]. Thus, we do not include it in our simulations. We simulate the spin dynamics for $N = 4$ sites with the initial state $|\psi_0\rangle = \sigma_1^+|\Omega\rangle$, where $|\Omega\rangle$ is the all-spin-down state. The XXZ Hamiltonian drives spin excitations across the chain, captured by the site populations $n_k(t) = (\langle\psi(t)|\sigma_k^z|\psi(t)\rangle + 1)/2$. This quantity corresponds to the qubit population in the simulator and is experimentally measurable.

The simulation results are presented in Fig. 6, with the number of Trotter steps optimized to balance Trotter and gate errors (see Appendix A). Without J_z coupling or dissipation, the spin excitation oscillates between sites 1 and 4 [Fig. 6(a)]. Dissipation at site 4 reduces the oscillation amplitude [Fig. 6(b)]. The inclusion of J_z interaction decreases spin current, with $J_z = J$ marking the transition between ballistic and diffusive transport regimes [72]. For $J_z = 5J > J$, the reduced spin current diminishes the impact of dissipation on the dynamics [Fig. 6(d)].

We note that the extended gate set in our simulator minimizes circuit depth compared to alternatives using only iSWAP gates: at least two iSWAP gates (among other single-qubit gates) are required to perform a R_{ZZ} gate. Instead, in our case, the R_{ZZ} gate is performed with only one CZ_φ and single-qubit gates. The efficient compilation provided by our simulator

reduces the simulation time and enhances the fidelity of the simulation; as discussed in Section 2, the increase in fidelity is approximately linear in the decrease in simulation time [77, 86].

5 Extension to simulations in higher dimensions and a universal quantum processor

The one-dimensional structure of our simulator as it is laid out in Section 3 introduces significant overhead when executing long-range two-qubit gates. Consequently, the implementation of quantum algorithms and protocols requiring such interactions—such as those used in the surface code [69]—becomes inefficient, restricting our simulator’s ability to function as a universal quantum processor. One potential solution to this limitation involves implementing a multi-braided configuration by having the waveguide cross itself once to enable tunable all-to-all couplings between giant atoms [35, 68]. However, this approach introduces a frequency-crowding challenge: distinct two-qubit gates must operate at sufficiently separated frequencies to prevent unwanted interactions during execution. Since qubits function within a constrained frequency range, increasing the number of qubits reduces the available frequency spacing, ultimately limiting scalability.

To alleviate this frequency-crowding issue, a square-lattice arrangement [95, 96] can be employed to suppress unwanted interactions between non-neighboring qubits while still enabling both iSWAP and CZ gates. Achieving this with giant atoms necessitates extending our simulator into two dimensions, incorporating an array of waveguides that mediate interactions between different giant atoms, as shown in Fig. 7(a). In this setup, the giant atoms (except the ones on the first and last row) are coupled to two waveguides in an identical fashion. This arrangement ensures that the giant atoms can be decoherence-free with respect to both waveguides at decoherence-free frequencies, enabling decoherence-free interactions between giant atoms that allows to execute two-qubit gates.

Since every giant atom is braided with at most four other giant atoms, we need at least nine decoherence-free frequencies (four for executing R_{XYs} , four for executing CZ_φ , and one for a decoupled regime) within the operational frequency range of the atoms. To minimize individual atom decay within this frequency range, we choose ten coupling points to each waveguide for each giant atom. The coupling points are evenly spaced by $2\Delta x$ and have the same coupling strength γ . This yields a set of decoherence-free frequencies $\omega_{DF,nm} = (n/2 + m/20)\omega_0$ with $n \in \mathcal{N}$, $m = 1, \dots, 9$. At $\omega_{DF,nm}$, the braided giant atoms have a non-zero coupling while the unwanted cou-




Figure 7: A two-dimensional extension of the giant-atom-based simulator from Fig. 3(a) into a universal quantum processor. (a) Sketch of the structure of the quantum processor. In this design, each giant atom—except those at the boundary—is braided with four neighboring atoms via two waveguides, enabling tunable interactions. This structure supports various two-qubit gate operations through frequency tuning. (b,c) The frequency dependence of individual decay rates Γ_{ind} , coupling strengths g and collective decay rates Γ_{coll} for (b) neighboring and (c) non-neighboring giant atoms.

pling between non-braided giant atoms is eliminated [see Fig. 7(b)-(c)]. Additionally, within the operational frequency regime $[\omega_{\text{DF},n1}, \omega_{\text{DF},n9}]$, the individual decay rates of the giant atoms are small, and can further be minimized by optimizing the coupling strengths. Thus, the giant-atom-based quantum processor in Fig. 7(a) maintains the benefits of the giant-atom-based quantum simulator in Fig. 3(a), and furthermore offers possibilities for long-range two-qubit gates owing to its two-dimensional structure.

We now demonstrate how different gates can be executed in the quantum processor in Fig. 7(a), focusing on its building block containing five qubits (numbered 1, 2, 4, 6, 7) indicated with the dashed orange rectangle. We set the anharmonicities of the atoms to $\chi_{1,2,6,7} = -\omega_0/20$ to ensure minimal decay. To avoid decoherence and interactions while performing single-qubit gates, the frequencies of the qubits can be set to $\omega_{1(2,4,6,7)} = \omega_{\text{DF},n2(4,5,7,9)}$, such that there are no couplings between the giant atoms. To perform an R_{XY} gate between qubits 1 and 4, we can tune $\omega_4 \rightarrow \omega_{\text{DF},n2}$; to perform a CZ_φ gate between qubits 1 and 4, we can tune $\omega_4 \rightarrow \omega_{\text{DF},n1}$. Similarly, two-qubit R_{XY} and CZ_φ gates can be performed between qubit 4 and other qubits.

We have thus demonstrated the ability of our two-dimensional quantum processor to execute the extended gate set between neighboring atoms on the square lattice. This in particular allows to perform algorithms such as the surface code [97, 98] to enable fault-tolerant quantum computation on our setup, offering it as a universal quantum processor. A detailed analysis of implementing specific algorithms within this structure would require extensive many-body calculations, which we leave for future investigation.

6 Discussion

We have presented a scalable quantum simulator for open quantum systems. The simulator is based on giant atoms and features an extended gate set that enhances its versatility for simulating open quantum many-body dynamics. Furthermore, we have discussed how to extend the structure in two dimensions for a universal quantum processor. We now move on to discuss (i) the physical realization of the proposed setups and (ii) the potential impact of non-Markovian effects as the system scales up.

6.1 Physical realization

A promising platform for implementing our quantum simulator is superconducting qubits [93, 99], such as transmons [74], coupled to a waveguide. There have already been several experiments demonstrating that this platform can be used for giant atoms [61, 64, 66, 67]. With a typical waveguide speed of light $v \approx 1.3 \times 10^8$ m/s and $\omega_0/(2\pi) = 3.2$ GHz, the required

coupling-point spacing is $\Delta x = 2\pi v/\omega_0 \approx 41$ nm. This means that adding a single qubit necessitates an additional waveguide length of $5\Delta x \approx 0.21$ m. State-of-the-art fabrication techniques can produce waveguides up to 30 m in length [100, 101], which could accommodate approximately 140 qubits, demonstrating the feasibility of our proposed architecture.

In addition to superconducting qubits, other physical platforms could support the realization of this setup. For example, cold atoms coupled to an optical lattice [36] present an intriguing alternative, offering distinct advantages in terms of coherence times and system scalability. Exploring such platforms could pave the way for diverse implementations of giant-atom-based quantum simulators.

6.2 Non-Markovian effects

Scaling up our simulator enhances non-Markovian effects, which could challenge the validity of Eq. (1). The primary source of non-Markovianity in this system is the time delay associated with photons traveling between coupling points. The Markovian assumption holds as long as $\gamma L_w/v \ll 1$, where L_w is the length of the waveguide between coupling points and γ is the coupling rate. With $\gamma/(2\pi) = 2$ MHz and $v = 1.3 \times 10^8$ m/s, this condition is satisfied for $L_w \ll 130$ m, exceeding current state-of-the-art waveguide lengths.

However, as waveguides approach these lengths or coupling rates increase, deviations from the Markovian regime may emerge. In such cases, incorporating time delays into the theoretical framework and exploring non-Markovian models is necessary. As solving non-Markovian many-body systems remains an open challenge, we leave this for future work.

7 Conclusion

We have proposed a scalable quantum simulator with an extended gate set, leveraging the unique properties of giant atoms to simulate open quantum systems. The fundamental building block of this processor consists of two giant three-level atoms coupled to the same waveguide in a braided configuration. We demonstrated that this setup enables the realization of both R_{XY} and controlled-phase (CZ_φ) gates through simple frequency tuning of some of the giant atoms. This capability arises from the decoherence-free interaction characteristic of giant atoms, eliminating the need for additional hardware components like parametric couplers.

The scalability of this building block facilitates the construction of a many-body quantum simulator, where nearest-neighbor R_{XY} and CZ_φ gates can be implemented efficiently by controlling the qubit frequencies. To showcase the simulator's potential, we performed a Trotterized simulation of the dynamics of

a dissipative XXZ spin chain, demonstrating its capability to tackle complex problems in open quantum many-body dynamics.

Our work provides a versatile and scalable platform for quantum simulation, featuring an extended gate set that enhances circuit compilation efficiency while maintaining scalability. The inclusion of both iSWAP-like and controlled-phase gates positions this simulator as a promising candidate for addressing state-of-the-art challenges in quantum simulation, particularly in open quantum many-body physics. Furthermore, extending our simulator into two dimensions could provide a pathway toward a scalable universal quantum processor.

As an outlook for future work, we have already mentioned a detailed analysis of the simulation of specific quantum systems (including ones featuring non-Markovian effects) or of implementation of specific quantum algorithms (including the surface code for error correction), as well as actual experimental implementation with superconducting circuits or other platforms. We also note the possibility of extending the gate set for giant-atom-based simulators even further, e.g., by incorporating three-qubit gates using

schemes similar to those in Refs. [102, 103].

Acknowledgements

We thank Liangyu Chen, Akshay Gaikwad, and Laura García Álvarez for fruitful discussions. G.C. is supported by European Union's Horizon Europe programme HORIZON-MSCA-2023-PF-01-01 via the project 101146565 (SING-ATOM). A.F.K. acknowledges support from the Swedish Research Council (grant number 2019-03696), the Swedish Foundation for Strategic Research (grant numbers FFL21-0279 and FUS21-0063), the Horizon Europe programme HORIZON-CL4-2022-QUANTUM-01-SGA via the project 101113946 OpenSuperQPlus100, and from the Knut and Alice Wallenberg Foundation through the Wallenberg Centre for Quantum Technology (WACQT).

A Error analysis for the quantum simulation of the dissipative XXZ model

Here we present the numerical details for the simulations presented in Fig. 6. The simulator dynamics, which involve 3-level atoms, are governed by the equation

$$\begin{aligned}
\partial_t \rho = & -i \left[\sum_{j=1}^N (\omega_j (|1\rangle\langle 1|)_j + (2\omega_j + \chi_j) (|2\rangle\langle 2|)_j) + \sum_{j=1}^N \sum_{k=1}^N g(\omega_j, \omega_k) \left(\sigma_j^{+, (01)} \sigma_k^{-, (01)} + \text{H.c.} \right) \right. \\
& + \sum_{j=1}^N \sum_{k=1}^N \sqrt{2} g(\omega_j + \chi_j, \omega_k) \left(\sigma_j^{+, (12)} \sigma_k^{-, (01)} + \text{H.c.} \right) + \sum_{j=1}^N \sum_{k=1}^N 2g(\omega_j + \chi_j, \omega_k + \chi_k) \left(\sigma_j^{+, (12)} \sigma_k^{-, (12)} + \text{H.c.} \right), \rho \\
& + \sum_{j=1}^N \Gamma(\omega_j) \mathcal{D}[\sigma_j^{-, (01)}] \rho + \sum_{j=1}^N \sqrt{2} \Gamma(\omega_j + \chi_j) \mathcal{D}[\sigma_j^{-, (12)}] \rho \\
& + \sum_{j=1}^N \sum_{k=1}^N \Gamma_{\text{coll}}(\omega_j, \omega_k) \left[\left(\sigma_j^{-, (01)} \rho \sigma_k^{+, (01)} - \frac{1}{2} \left\{ \sigma_j^{+, (01)} \sigma_k^{-, (01)}, \rho \right\} \right) + \text{H.c.} \right] \\
& + \sum_{j=1}^N \sum_{k=1}^N \sqrt{2} \Gamma_{\text{coll}}(\omega_j, \omega_k + \chi_k) \left[\left(\sigma_j^{-, (01)} \rho \sigma_k^{+, (12)} - \frac{1}{2} \left\{ \sigma_j^{+, (01)} \sigma_k^{-, (12)}, \rho \right\} \right) + \text{H.c.} \right] \\
& + \sum_{j=1}^N \sum_{k=1}^N \sqrt{2} \Gamma_{\text{coll}}(\omega_j + \chi_j, \omega_k) \left[\left(\sigma_j^{-, (12)} \rho \sigma_k^{+, (01)} - \frac{1}{2} \left\{ \sigma_j^{+, (12)} \sigma_k^{-, (01)}, \rho \right\} \right) + \text{H.c.} \right] \\
& + \sum_{j=1}^N \sum_{k=1}^N 2\Gamma_{\text{coll}}(\omega_j + \chi_j, \omega_k + \chi_k) \left[\left(\sigma_j^{-, (12)} \rho \sigma_k^{+, (12)} - \frac{1}{2} \left\{ \sigma_j^{+, (12)} \sigma_k^{-, (12)}, \rho \right\} \right) + \text{H.c.} \right]. \tag{16}
\end{aligned}$$

Here ω_j , χ_j are the transition frequency and anharmonicity of qubit j , respectively, while $\sigma_j^{+, (01)}$

$(\sigma_j^{-, (01)})$ is the raising (lowering) operator of the $|0\rangle$ ($|1\rangle$) level of qubit j . The coupling strength between the qubits mediated by the waveguide is denoted by




Figure 8: Simulation error for the dynamics of the dissipative XXZ spin chain [Eq. (13)] with $J_z = 0$, $\Gamma = J$, and $N = 4$ sites with our quantum simulator. With a larger number of Trotter steps l , the Trotter error is decreased while a larger error stems from the need for more gates to be performed. At short times, gate error dominates and thus a smaller number of Trotter steps results in a smaller error; at long times, Trotter error dominates and a larger number of Trotter steps results in a smaller error.

g . $\Gamma(\omega_j)$ and Γ_{coll} represent the individual and collective decay rates of the qubits, respectively, and H.c. denotes Hermitian conjugate. The parameter dependencies on the qubit frequencies are illustrated in Fig. 3(b).

Our simulations operate within the decoherence-free frequency regime, ensuring that non-neighboring qubits do not interact [Fig. 3(c)]. During the simulation, the time evolution of the qubit frequencies follows the profile shown in Fig. 5(b), leading to a time-dependent master equation as in Eq. (16), which we solve numerically using QuTiP [104–106].

We define the simulation error in the simulated population $n_k(t) := (\langle \psi(t) | \sigma_k^z | \psi(t) \rangle + 1)/2$ as

$$\Delta n_k(t) = [n_k(t)]_{\text{exact}} - n_k(t), \quad (17)$$

where $[n_k(t)]_{\text{exact}}$ is the exact result. We analyze the simulation error as a function of the number of Trotter steps l .

We first consider the case of $J_z = 0$ and $\Gamma = J$ in Eq. (13), where no CZ_φ gates are needed in the simulation. The results are shown in Fig. 8. There are two main sources of error: (i) the Trotter error stemming from the Trotterization, which scales as t^2/l , and (ii) the gate error, which increases with l since the number of gates grows with l . Thus, there is an optimal number of Trotter steps $l_{\text{opt}}(t)$ for the simulation of




Figure 9: Simulation error for the dynamics of the dissipative XXZ spin chain [Eq. (13)] with $J_z = 5J$, $\Gamma = J$, and $N = 4$ sites with our quantum simulator. Compared to the case of $J_z = 0$ in Fig. 8, the inclusion of CZ_φ gate results in faster growth of gate error with respect to l . Thus, the optimal number of Trotter steps has decreased for the same time t . In particular, for $t = 2$, $l = 10$ results in smaller errors than $l = 20$.

the dynamics at time t , and in particular, $l_{\text{opt}}(t)$ increases with t . This explanation is in agreement with the simulation error shown in Fig. 8, which decreases with l for large t , and increases with l for small t .

We next analyze the simulation error for $J_z = 5J$ and $\Gamma = J$ in Eq. (13). Here, CZ_φ gates are necessary, which increases the number of gates per Trotter step. Consequently, the gate error grows more rapidly with l compared to the $J_z = 0$ case, resulting in a smaller $l_{\text{opt}}(t)$. This is illustrated in Fig. 9, where the decreased $l_{\text{opt}}(t)$ leads to a higher Trotter error and thus greater overall simulation error.

The above analysis underscores the significance of minimizing circuit depth to ensure high simulation accuracy. In particular, a shorter circuit slows the growth of gate error with increasing l , enabling a larger $l_{\text{opt}}(t)$ and reducing simulation error. The extended gate set provided by our simulator effectively reduces circuit depth, enhancing the accuracy of quantum simulations.

References

- [1] I. M. Georgescu, S. Ashhab, and Franco Nori. “Quantum simulation”. *Reviews of Modern Physics* **86**, 153–185 (2014).
- [2] Ehud Altman et al. “Quantum Simulators: Ar-

- chitectures and Opportunities”. *PRX Quantum* **2**, 017003 (2021).
- [3] Benedikt Fauseweh. “Quantum many-body simulations on digital quantum computers: State-of-the-art and future challenges”. *Nature Communications* **15**, 2123 (2024).
- [4] Adriano Barenco, Charles H. Bennett, Richard Cleve, David P. DiVincenzo, Norman Margolus, Peter Shor, Tycho Sleator, John A. Smolin, and Harald Weinfurter. “Elementary gates for quantum computation”. *Physical Review A* **52**, 3457–3467 (1995).
- [5] Karen Wintersperger, Florian Dommert, Thomas Ehmer, Andrey HOURSANOV, Johannes Klepsch, Wolfgang Maurer, Georg Reuber, Thomas Strohm, Ming Yin, and Sebastian Luber. “Neutral atom quantum computing hardware: performance and end-user perspective”. *EPJ Quantum Technology* **10**, 32 (2023).
- [6] Thomas Strohm, Karen Wintersperger, Florian Dommert, Daniel Basilewitsch, Georg Reuber, Andrey HOURSANOV, Thomas Ehmer, Davide Vodola, and Sebastian Luber. “Ion-Based Quantum Computing Hardware: Performance and End-User Perspective” (2024). [arXiv:2405.11450](https://arxiv.org/abs/2405.11450).
- [7] Frank Leymann and Johanna Barzen. “The bitter truth about gate-based quantum algorithms in the NISQ era”. *Quantum Science and Technology* **5**, 044007 (2020).
- [8] Justin Kalloor, Mathias Weiden, Ed Younis, John Kubiatiowicz, Bert De Jong, and Costin Iancu. “Quantum Hardware Roofline: Evaluating the Impact of Gate Expressivity on Quantum Processor Design” (2024). [arXiv:2403.00132](https://arxiv.org/abs/2403.00132).
- [9] Yan Ge, Wu Wenjie, Chen Yuheng, Pan Kaisen, Lu Xudong, Zhou Zixiang, Wang Yuhan, Wang Ruocheng, and Yan Junchi. “Quantum Circuit Synthesis and Compilation Optimization: Overview and Prospects” (2024). [arXiv:2407.00736](https://arxiv.org/abs/2407.00736).
- [10] Christian Križan, Janka Biznárová, Liangyu Chen, Emil Hogedal, Amr Osman, Christopher W Warren, Sandoko Kosen, Hang-Xi Li, Tahereh Abad, Anuj Aggarwal, Marco Caputo, Jorge Fernández-Pendás, Akshay Gaikwad, Leif Grönberg, Andreas Nylander, Robert Rehammar, Marcus Rommel, Olga I Yuzepovich, Anton Frisk Kockum, Joonas Govenius, Giovanna Tancredi, and Jonas Bylander. “Quantum swap gate realized with cz and iswap gates in a superconducting architecture”. *New Journal of Physics* **27**, 074507 (2025).
- [11] Deanna M. Abrams, Nicolas Didier, Blake R. Johnson, Marcus P. da Silva, and Colm A. Ryan. “Implementation of xy entangling gates with a single calibrated pulse”. *Nature Electronics* **3**, 744–750 (2020).
- [12] M. Ganzhorn, G. Salis, D. J. Egger, A. Fuhrer, M. Mergenthaler, C. Müller, P. Müller, S. Paredes, M. Pechal, M. Werninghaus, and S. Filipp. “Benchmarking the noise sensitivity of different parametric two-qubit gates in a single superconducting quantum computing platform”. *Physical Review Research* **2**, 033447 (2020).
- [13] Eyob A. Sete, Nicolas Didier, Angela Q. Chen, Shobhan Kulshreshtha, Riccardo Manenti, and Stefano Poletto. “Parametric-Resonance Entangling Gates with a Tunable Coupler”. *Physical Review Applied* **16**, 024050 (2021).
- [14] Rigetti QCS. “Aspen-m-3 quantum processor” (2023).
- [15] R. Barends, C. M. Quintana, A. G. Petukhov, Yu Chen, D. Kafri, K. Kechedzhi, R. Collins, O. Naaman, S. Boixo, F. Arute, K. Arya, D. Buell, B. Burkett, Z. Chen, B. Chiaro, A. Dunsworth, B. Foxen, A. Fowler, C. Gidney, M. Giustina, R. Graff, T. Huang, E. Jeffrey, J. Kelly, P. V. Klimov, F. Kostritsa, D. Landhuis, E. Lucero, M. McEwen, A. Megrant, X. Mi, J. Mutus, M. Neeley, C. Neill, E. Ostby, P. Roushan, D. Sank, K. J. Satzinger, A. Vainsencher, T. White, J. Yao, P. Yeh, A. Zalcman, H. Neven, V. N. Smelyanskiy, and John M. Martinis. “Diabatic gates for frequency-tunable superconducting qubits”. *Physical Review Letters* **123**, 210501 (2019).
- [16] Atsushi Noguchi, Alto Osada, Shumpei Masuda, Shingo Kono, Kentaro Heya, Samuel Piotr Wolski, Hiroki Takahashi, Takanori Sugiyama, Dany Lachance-Quirion, and Yasunobu Nakamura. “Fast parametric two-qubit gates with suppressed residual interaction using the second-order nonlinearity of a cubic transmon”. *Physical Review A* **102**, 062408 (2020).
- [17] Eyob A. Sete, Nicolas Didier, Angela Q. Chen, Shobhan Kulshreshtha, Riccardo Manenti, and Stefano Poletto. “Parametric-resonance entangling gates with a tunable coupler”. *Physical Review Applied* **16**, 024050 (2021).
- [18] S. A. Caldwell, N. Didier, C. A. Ryan, E. A. Sete, A. Hudson, P. Karalekas, R. Manenti, M. P. da Silva, R. Sinclair, E. Acala, N. Alidoust, J. Angeles, A. Bestwick, M. Block, B. Bloom, A. Bradley, C. Bui, L. Capeluto, R. Chilcott, J. Cordova, G. Crossman, M. Curtis, S. Deshpande, T. El Bouayadi, D. Girshovich, S. Hong, K. Kuang, M. Lenihan, T. Manning, A. Marchenkov, J. Marshall, R. Maydra, Y. Mohan, W. O’Brien, C. Osborn, J. Otterbach, A. Papageorge, J.-P. Paquette, M. Pelstring, A. Polloreno, G. Prawiroatmodjo, V. Rawat, M. Reagor, R. Renzas, N. Rubin, D. Russell, M. Rust, D. Scarabelli,

- M. Scheer, M. Selvanayagam, R. Smith, A. Staley, M. Suska, N. Tezak, D. C. Thompson, T.-W. To, M. Vahidpour, N. Vodrahalli, T. Whyland, K. Yadav, W. Zeng, and C. Rigetti. “Parametrically activated entangling gates using transmon qubits”. *Physical Review Applied* **10**, 034050 (2018).
- [19] Nathan Lacroix, Christoph Hellings, Christian Kraglund Andersen, Agustin Di Paolo, Ants Remm, Stefania Lazar, Sebastian Krinner, Graham J. Norris, Mihai Gabureac, Johannes Heinsoo, Alexandre Blais, Christopher Eichler, and Andreas Wallraff. “Improving the Performance of Deep Quantum Optimization Algorithms with Continuous Gate Sets”. *PRX Quantum* **1**, 110304 (2020).
- [20] Youngkyu Sung, Leon Ding, Jochen Braumüller, Antti Vepsäläinen, Bharath Kannan, Morten Kjaergaard, Ami Greene, Gabriel O. Samach, Chris McNally, David Kim, Alexander Melville, Bethany M. Niedzielski, Mollie E. Schwartz, Jonilyn L. Yoder, Terry P. Orlando, Simon Gustavsson, and William D. Oliver. “Realization of High-Fidelity CZ and ZZ-Free iSWAP Gates with a Tunable Coupler”. *Physical Review X* **11**, 021058 (2021).
- [21] Helin Zhang, Chunyang Ding, D.K. Weiss, Ziwen Huang, Yuwei Ma, Charles Guinn, Sara Sussman, Sai Pavan Chitta, Danyang Chen, Andrew A. Houck, Jens Koch, and David I. Schuster. “Tunable Inductive Coupler for High-Fidelity Gates Between Fluxonium Qubits”. *PRX Quantum* **5**, 020326 (2024).
- [22] Zhen Chen, Weiyang Liu, Yanjun Ma, Weijie Sun, Ruixia Wang, He Wang, Huikai Xu, Guangming Xue, Haisheng Yan, Zhen Yang, Jiayu Ding, Yang Gao, Feiyu Li, Yujia Zhang, Zikang Zhang, Yirong Jin, Haifeng Yu, Jianxin Chen, and Fei Yan. “Efficient implementation of arbitrary two-qubit gates using unified control”. *Nature Physics* **21**, 1489–1496 (2025).
- [23] Tian-Ming Li, Jia-Chi Zhang, Bing-Jie Chen, Kaixuan Huang, Hao-Tian Liu, Yong-Xi Xiao, Cheng-Lin Deng, Gui-Han Liang, Chi-Tong Chen, Yu Liu, Hao Li, Zhen-Ting Bao, Kui Zhao, Yueshan Xu, Li Li, Yang He, Zheng-He Liu, Yi-Han Yu, Si-Yun Zhou, Yan-Jun Liu, Xiaohui Song, Dongning Zheng, Zhongcheng Xiang, Yun-Hao Shi, Kai Xu, and Heng Fan. “High-precision pulse calibration of tunable couplers for high-fidelity two-qubit gates in superconducting quantum processors”. *Physical Review Applied* **23**, 024059 (2025).
- [24] Hao-Tian Liu, Bing-Jie Chen, Jia-Chi Zhang, Yong-Xi Xiao, Tian-Ming Li, Kaixuan Huang, Ziting Wang, Hao Li, Kui Zhao, Yueshan Xu, Cheng-Lin Deng, Gui-Han Liang, Zheng-He Liu, Si-Yun Zhou, Cai-Ping Fang, Xiaohui Song, Zhongcheng Xiang, Dongning Zheng, Yun-Hao Shi, Kai Xu, and Heng Fan. “Direct implementation of high-fidelity three-qubit gates for superconducting processor with tunable couplers”. *Physical Review Letters* **135**, 050602 (2025).
- [25] K. X. Wei, E. Magesan, I. Lauer, S. Srinivasan, D. F. Bogorin, S. Carnevale, G. A. Keefe, Y. Kim, D. Klaus, W. Landers, N. Sundaresan, C. Wang, E. J. Zhang, M. Steffen, O. E. Dial, D. C. McKay, and A. Kandala. “Hamiltonian Engineering with Multicolor Drives for Fast Entangling Gates and Quantum Crosstalk Cancellation”. *Physical Review Letters* **129**, 060501 (2022).
- [26] Long B. Nguyen, Yosep Kim, Akel Hashim, Noah Goss, Brian Marinelli, Bibek Bhandari, Debmalya Das, Ravi K. Naik, John Mark Kreikebaum, Andrew N. Jordan, David I. Santiago, and Irfan Siddiqi. “Programmable Heisenberg interactions between Floquet qubits”. *Nature Physics* **20**, 240–246 (2024).
- [27] K. X. Wei, E. Magesan, I. Lauer, S. Srinivasan, D. F. Bogorin, S. Carnevale, G. A. Keefe, Y. Kim, D. Klaus, W. Landers, N. Sundaresan, C. Wang, E. J. Zhang, M. Steffen, O. E. Dial, D. C. McKay, and A. Kandala. “Hamiltonian engineering with multicolor drives for fast entangling gates and quantum crosstalk cancellation”. *Physical Review Letters* **129**, 060501 (2022).
- [28] Helin Zhang, Chunyang Ding, D.K. Weiss, Ziwen Huang, Yuwei Ma, Charles Guinn, Sara Sussman, Sai Pavan Chitta, Danyang Chen, Andrew A. Houck, Jens Koch, and David I. Schuster. “Tunable inductive coupler for high-fidelity gates between fluxonium qubits”. *PRX Quantum* **5**, 020326 (2024).
- [29] Simon J. Evered, Dolev Bluvstein, Marcin Kalinowski, Sepehr Ebadi, Tom Manovitz, Hengyun Zhou, Sophie H. Li, Alexandra A. Geim, Tout T. Wang, Nishad Maskara, Harry Levine, Giulia Semeghini, Markus Greiner, Vladan Vuletić, and Mikhail D. Lukin. “High-fidelity parallel entangling gates on a neutral-atom quantum computer”. *Nature* **622**, 268–272 (2023).
- [30] B. Foxen, C. Neill, A. Dunsworth, P. Roushan, B. Chiaro, A. Megrant, J. Kelly, Zijun Chen, K. Satzinger, R. Barends, F. Arute, K. Arya, R. Babbush, D. Bacon, J. C. Bardin, S. Boixo, D. Buell, B. Burkett, Yu Chen, R. Collins, E. Farhi, A. Fowler, C. Gidney, M. Giustina, R. Graff, M. Harrigan, T. Huang, S. V. Isakov, E. Jeffrey, Z. Jiang, D. Kafri, K. Kechedzhi, P. Klimov, A. Korotkov, F. Kostritsa, D. Landhuis, E. Lucero, J. McClean, M. McEwen, X. Mi, M. Mohseni, J. Y. Mutus, O. Naaman,

- M. Neeley, M. Niu, A. Petukhov, C. Quintana, N. Rubin, D. Sank, V. Smelyanskiy, A. Vainsencher, T. C. White, Z. Yao, P. Yeh, A. Zalcman, H. Neven, and J. M. Martinis. “Demonstrating a continuous set of two-qubit gates for near-term quantum algorithms”. *Physical Review Letters* **125**, 120504 (2020).
- [31] Ken Xuan Wei, Isaac Lauer, Emily Pritchett, William Shanks, David C. McKay, and Ali Javadi-Abhari. “Native two-qubit gates in fixed-coupling, fixed-frequency transmons beyond cross-resonance interaction”. *PRX Quantum* **5**, 020338 (2024).
- [32] A. F. Kockum. “Quantum Optics with Giant Atoms - the First Five Years”. In International Symposium on Mathematics, Quantum Theory, and Cryptography (Mathematics for Industry, vol 33). Pages 125–146. Springer (2021).
- [33] Anton Frisk Kockum, Per Delsing, and Göran Johansson. “Designing frequency-dependent relaxation rates and Lamb shifts for a giant artificial atom”. *Physical Review A* **90**, 013837 (2014).
- [34] L. Guo, A. L. Grimsmo, A. F. Kockum, M. Pletyukhov, and G. Johansson. “Giant acoustic atom: A single quantum system with a deterministic time delay”. *Physical Review A* **95**, 053821 (2017).
- [35] Anton Frisk Kockum, Göran Johansson, and Franco Nori. “Decoherence-Free Interaction between Giant Atoms in Waveguide Quantum Electrodynamics”. *Physical Review Letters* **120**, 140404 (2018).
- [36] A González-Tudela, C Sánchez Muñoz, and J I Cirac. “Engineering and Harnessing Giant Atoms in High-Dimensional Baths: A Proposal for Implementation with Cold Atoms”. *Physical Review Letters* **122**, 203603 (2019).
- [37] Lingzhen Guo, Anton Frisk Kockum, Florian Marquardt, and Göran Johansson. “Oscillating bound states for a giant atom”. *Physical Review Research* **2**, 043014 (2020).
- [38] P. O. Guimond, B. Vermersch, M. L. Juan, A. Sharafiev, G. Kirchmair, and P. Zoller. “A unidirectional on-chip photonic interface for superconducting circuits”. *npj Quantum Information* **6**, 32 (2020).
- [39] A. Ask, Y.-L. L. Fang, and A. F. Kockum. “Synthesizing electromagnetically induced transparency without a control field in waveguide QED using small and giant atoms” (2020). [arXiv:2011.15077](https://arxiv.org/abs/2011.15077).
- [40] Dario Cilluffo, Angelo Carollo, Salvatore Lorenzo, Jonathan A Gross, G Massimo Palma, and Francesco Ciccarello. “Collisional picture of quantum optics with giant emitters”. *Physical Review Research* **2**, 043070 (2020).
- [41] Xin Wang, Tao Liu, Anton Frisk Kockum, Hong-Rong Li, and Franco Nori. “Tunable Chiral Bound States with Giant Atoms”. *Physical Review Letters* **126**, 043602 (2021).
- [42] Lei Du and Yong Li. “Single-photon frequency conversion via a giant Λ -type atom”. *Physical Review A* **104**, 023712 (2021).
- [43] Ariadna Soro and Anton Frisk Kockum. “Chiral quantum optics with giant atoms”. *Physical Review A* **105**, 023712 (2022).
- [44] Xin Wang and Hong-rong Li. “Chiral quantum network with giant atoms”. *Quantum Science and Technology* **7**, 035007 (2022).
- [45] Lei Du, Yan Zhang, Jin Hui Wu, Anton Frisk Kockum, and Yong Li. “Giant Atoms in a Synthetic Frequency Dimension”. *Physical Review Letters* **128**, 223602 (2022).
- [46] Lei Du, Yao-Tong Chen, Yan Zhang, and Yong Li. “Giant atoms with time-dependent couplings”. *Physical Review Research* **4**, 023198 (2022).
- [47] Sergi Terradas-Briansó, Carlos A. González-Gutiérrez, Franco Nori, Luis Martín-Moreno, and David Zueco. “Ultrastrong waveguide QED with giant atoms”. *Physical Review A* **106**, 063717 (2022).
- [48] Ariadna Soro, Carlos Sánchez Muñoz, and Anton Frisk Kockum. “Interaction between giant atoms in a one-dimensional structured environment”. *Physical Review A* **107**, 013710 (2023).
- [49] Lei Du, Lingzhen Guo, Yan Zhang, and Anton Frisk Kockum. “Giant emitters in a structured bath with non-Hermitian skin effect”. *Physical Review Research* **5**, L042040 (2023).
- [50] Emil Raaholt Ingelsten, Anton Frisk Kockum, and Ariadna Soro. “Avoiding decoherence with giant atoms in a two-dimensional structured environment”. *Physical Review Research* **6**, 043222 (2024).
- [51] Luca Leonforte, Xuejian Sun, Davide Valenti, Bernardo Spagnolo, Fabrizio Illuminati, Angelo Carollo, and Francesco Ciccarello. “Quantum optics with giant atoms in a structured photonic bath”. *Quantum Science and Technology* **10**, 015057 (2025).
- [52] Xin Wang, Huai-Bing Zhu, Tao Liu, and Franco Nori. “Realizing quantum optics in structured environments with giant atoms”. *Physical Review Research* **6**, 013279 (2024).
- [53] Federico Roccati and Dario Cilluffo. “Controlling Markovianity with Chiral Giant Atoms”. *Physical Review Letters* **133**, 063603 (2024).
- [54] Rui-Yang Gong, Zi-Yu He, Cheng-He Yu, Ge-Fei Zhang, Franco Nori, and Ze-Liang Xiang. “Tunable quantum router with giant atoms, implementing quantum gates, teleportation, non-reciprocity, and circulators” (2024). [arXiv:2411.19307](https://arxiv.org/abs/2411.19307).

- [55] Lei Du and Anton Frisk Kockum. “Unconventional and robust light-matter interactions based on the non-Hermitian skin effect”. *Physical Review Research* **7**, 013140 (2025).
- [56] Martin V Gustafsson, Thomas Aref, Anton Frisk Kockum, Maria K Ekström, Göran Johansson, and Per Delsing. “Propagating phonons coupled to an artificial atom”. *Science* **346**, 207–211 (2014).
- [57] R Manenti, A F Kockum, A Patterson, T Behrle, J Rahamim, G Tancredi, F Nori, and P J Leek. “Circuit quantum acoustodynamics with surface acoustic waves”. *Nature Communications* **8**, 975 (2017).
- [58] K J Satzinger, Y P Zhong, H.-S. Chang, G A Peairs, A Bienfait, M.-H. Chou, A Y Cleland, C R Conner, É Dumur, J Grebel, I Gutierrez, B H November, R G Povey, S J Whiteley, D D Awschalom, D I Schuster, and A N Cleland. “Quantum control of surface acoustic-wave phonons”. *Nature* **563**, 661 (2018).
- [59] A Bienfait, K J Satzinger, Y P Zhong, H.-S. Chang, M.-H. Chou, C R Conner, É Dumur, J Grebel, G A Peairs, R G Povey, and A N Cleland. “Phonon-mediated quantum state transfer and remote qubit entanglement”. *Science* **364**, 368 (2019).
- [60] Gustav Andersson, Baladitya Suri, Lingzhen Guo, Thomas Aref, and Per Delsing. “Non-exponential decay of a giant artificial atom”. *Nature Physics* **15**, 1123–1127 (2019).
- [61] Bharath Kannan, Max J. Ruckriegel, Daniel L. Campbell, Anton Frisk Kockum, Jochen Braumüller, David K. Kim, Morten Kjaergaard, Philip Krantz, Alexander Melville, Bethany M. Niedzielski, Antti Vepsäläinen, Roni Winik, Jonilyn L. Yoder, Franco Nori, Terry P. Orlando, Simon Gustavsson, and William D. Oliver. “Waveguide quantum electrodynamics with superconducting artificial giant atoms”. *Nature* **583**, 775–779 (2020).
- [62] A Bienfait, Y P Zhong, H.-S. Chang, M.-H. Chou, C R Conner, É Dumur, J Grebel, G A Peairs, R G Povey, K J Satzinger, and A N Cleland. “Quantum Erasure Using Entangled Surface Acoustic Phonons”. *Physical Review X* **10**, 021055 (2020).
- [63] G Andersson, M K Ekström, and P Delsing. “Electromagnetically Induced Acoustic Transparency with a Superconducting Circuit”. *Physical Review Letters* **124**, 240402 (2020).
- [64] A. M. Vadiraj, Andreas Ask, T. G. McConkey, I. Nsanzineza, C. W. Sandbo Chang, Anton Frisk Kockum, and C. M. Wilson. “Engineering the level structure of a giant artificial atom in waveguide quantum electrodynamics”. *Physical Review A* **103**, 023710 (2021).
- [65] Zi Qi Wang, Yi Pu Wang, Jiguang Yao, Rui Chang Shen, Wei Jiang Wu, Jie Qian, Jie Li, Shi Yao Zhu, and J. Q. You. “Giant spin ensembles in waveguide magnonics”. *Nature Communications* **13**, 7580 (2022).
- [66] Chaitali Joshi, Frank Yang, and Mohammad Mirhosseini. “Resonance Fluorescence of a Chiral Artificial Atom”. *Physical Review X* **13**, 021039 (2023).
- [67] Jingjing Hu, Dengfeng Li, Yufan Qie, Zelong Yin, Anton Frisk Kockum, Franco Nori, and Shuoming An. “Engineering the Environment of a Superconducting Qubit with an Artificial Giant Atom” (2024). [arXiv:2410.15377](https://arxiv.org/abs/2410.15377).
- [68] Guangze Chen and Anton Frisk Kockum. “Simulating open quantum systems with giant atoms”. *Quantum Science and Technology* **10**, 025028 (2025).
- [69] A. G. Fowler, M. Mariantoni, J. M. Martinis, and A. N. Cleland. “Surface codes: Towards practical large-scale quantum computation”. *Physical Review A* **86**, 032324 (2012).
- [70] X. Mi et al. “Stable quantum-correlated many-body states through engineered dissipation”. *Science* **383**, 1332–1337 (2024).
- [71] E. Rosenberg et al. “Dynamics of magnetization at infinite temperature in a Heisenberg spin chain”. *Science* **384**, 48–53 (2024).
- [72] T Prosen. “Open XXZ Spin Chain: Nonequilibrium Steady State and a Strict Bound on Ballistic Transport”. *Physical Review Letters* **106**, 217206 (2011).
- [73] Marko Žnidarič. “Spin Transport in a One-Dimensional Anisotropic Heisenberg Model”. *Physical Review Letters* **106**, 220601 (2011).
- [74] J. Koch, T. M. Yu, J. Gambetta, A. A. Houck, D. I. Schuster, J. Majer, A. Blais, M. H. Devoret, S. M. Girvin, and R. J. Schoelkopf. “Charge-insensitive qubit design derived from the Cooper pair box”. *Physical Review A* **76**, 042319 (2007).
- [75] Alexei Gilchrist, Nathan K. Langford, and Michael A. Nielsen. “Distance measures to compare real and ideal quantum processes”. *Physical Review A* **71**, 062310 (2005).
- [76] Man-Duen Choi. “Completely positive linear maps on complex matrices”. *Linear Algebra and its Applications* **10**, 285–290 (1975).
- [77] Tahereh Abad, Jorge Fernández-Pendás, Anton Frisk Kockum, and Göran Johansson. “Universal Fidelity Reduction of Quantum Operations from Weak Dissipation”. *Physical Review Letters* **129**, 150504 (2022).
- [78] Morten Kjaergaard, Mollie E. Schwartz, Jochen Braumüller, Philip Krantz, Joel I.-J. Wang, Simon Gustavsson, and William D. Oliver. “Superconducting qubits: Current state of play”. *Annual Review of Condensed Matter Physics* **11**, 369–395 (2020).

- [79] Alexander P. M. Place, Lila V. H. Rodgers, Pranav Mundada, Basil M. Smitham, Mattias Fitzpatrick, Zhaoqi Leng, Anjali Premkumar, Jacob Bryon, Andrei Vrajitoarea, Sara Sussman, Guangming Cheng, Trisha Madhavan, Harshvardhan K. Babla, Xuan Hoang Le, Youqi Gang, Berthold Jäck, András Gyenis, Nan Yao, Robert J. Cava, Nathalie P. de Leon, and Andrew A. Houck. “New material platform for superconducting transmon qubits with coherence times exceeding 0.3 milliseconds”. *Nature Communications* **12**, 1779 (2021).
- [80] Aaron Somoroff, Quentin Ficheux, Raymond A. Mencia, Haonan Xiong, Roman Kuzmin, and Vladimir E. Manucharyan. “Millisecond Coherence in a Superconducting Qubit”. *Physical Review Letters* **130**, 267001 (2023).
- [81] Youngseok Kim, Andrew Eddins, Sajant Anand, Ken Xuan Wei, Ewout van den Berg, Sami Rosenblatt, Hasan Nayfeh, Yantao Wu, Michael Zaletel, Kristan Temme, and Abhinav Kandala. “Evidence for the utility of quantum computing before fault tolerance”. *Nature* **618**, 500 (2023).
- [82] Janka Biznárová, Amr Osman, Emil Rehnman, Lert Chayanun, Christian Krizán, Per Malmberg, Marcus Rommel, Christopher Warren, Per Delsing, August Yurgens, Jonas Bylander, and Anita Fadavi Roudsari. “Mitigation of interfacial dielectric loss in aluminum-on-silicon superconducting qubits”. *npj Quantum Information* **10**, 78 (2024).
- [83] Shingo Kono, Jiahe Pan, Mahdi Chegnizadeh, Xuxin Wang, Amir Youssefi, Marco Scigliuzzo, and Tobias J. Kippenberg. “Mechanically induced correlated errors on superconducting qubits with relaxation times exceeding 0.4 ms”. *Nature Communications* **15**, 3950 (2024).
- [84] Mustafa Bal et al. “Systematic improvements in transmon qubit coherence enabled by niobium surface encapsulation”. *npj Quantum Information* **10**, 43 (2024).
- [85] F. W. Strauch, P. R. Johnson, A. J. Dragt, C. J. Lobb, J. R. Anderson, and F. C. Wellstood. “Quantum Logic Gates for Coupled Superconducting Phase Qubits”. *Physical Review Letters* **91**, 167005 (2003).
- [86] Tahereh Abad, Yoni Schattner, Anton Frisk Kockum, and Göran Johansson. “Impact of decoherence on the fidelity of quantum gates leaving the computational subspace”. *Quantum* **9**, 1684 (2025).
- [87] The average gate fidelity of a two-qubit CZ gate is $F_{\text{ave, CZ}} \approx 1 - 0.8\Gamma_{\text{ex}}\tau - 1.15\Gamma_{\phi}\tau$, where τ is the gate time [86] and we have assumed that the two qubits have the same decay and dephasing rates. For two qubits coupled with interaction strength g , the gate time is $\tau = \pi/(\sqrt{2}g)$. Combining these equations, we find that it agrees with the equation in the main text up to a second-order correction.
- [88] Eli Chertkov, Zihan Cheng, Andrew C. Potter, Sarang Gopalakrishnan, Thomas M. Gatterman, Justin A. Gerber, Kevin Gilmore, Dan Gresh, Alex Hall, Aaron Hankin, Mitchell Matheny, Tanner Mengle, David Hayes, Brian Neyenhuis, Russell Stutz, and Michael Fossfeig. “Characterizing a non-equilibrium phase transition on a quantum computer”. *Nature Physics* **19**, 1799–1804 (2023).
- [89] Federico Carollo, Edward Gillman, Hendrik Weimer, and Igor Lesanovsky. “Critical Behavior of the Quantum Contact Process in One Dimension”. *Physical Review Letters* **123**, 100604 (2019).
- [90] Matteo Marcuzzi, Michael Buchhold, Sebastian Diehl, and Igor Lesanovsky. “Absorbing State Phase Transition with Competing Quantum and Classical Fluctuations”. *Physical Review Letters* **116**, 245701 (2016).
- [91] Masuo Suzuki. “Fractal decomposition of exponential operators with applications to many-body theories and Monte Carlo simulations”. *Physics Letters A* **146**, 319–323 (1990).
- [92] M. Kliesch, T. Barthel, C. Gogolin, M. Kastoryano, and J. Eisert. “Dissipative Quantum Church-Turing Theorem”. *Physical Review Letters* **107**, 120501 (2011).
- [93] Alexandre Blais, Arne L. Grimsmo, S. M. Girvin, and Andreas Wallraff. “Circuit quantum electrodynamics”. *Reviews of Modern Physics* **93**, 025005 (2021).
- [94] Michele C. Collodo, Johannes Herrmann, Nathan Lacroix, Christian Kraglund Andersen, Ants Remm, Stefania Lazar, Jean-Claude Besse, Theo Walter, Andreas Wallraff, and Christopher Eichler. “Implementation of Conditional Phase Gates Based on Tunable ZZ Interactions”. *Physical Review Letters* **125**, 240502 (2020).
- [95] Amr Osman, Jorge Fernández-Pendás, Christopher Warren, Sandoko Kosen, Marco Scigliuzzo, Anton Frisk Kockum, Giovanna Tancredi, Anita Fadavi Roudsari, and Jonas Bylander. “Mitigation of frequency collisions in superconducting quantum processors”. *Physical Review Research* **5**, 043001 (2023).
- [96] Sandoko Kosen, Hang-xi Li, Marcus Rommel, Robert Rehammar, Marco Caputo, Leif Grönberg, Jorge Fernández-Pendás, Anton Frisk Kockum, Janka Biznárová, Liangyu Chen, Christian Krizán, Andreas Nylander, Amr Osman, Anita Fadavi Roudsari, Daryoush Shiri, Giovanna Tancredi, Joonas Govenius, and Jonas Bylander. “Signal Crosstalk in a Flip-

- Chip Quantum Processor”. *PRX Quantum* **5**, 030350 (2024).
- [97] Christian Kraglund Andersen, Ants Remm, Stefania Lazar, Sebastian Krinner, Nathan Lacroix, Graham J. Norris, Mihai Gabureac, Christopher Eichler, and Andreas Wallraff. “Repeated quantum error detection in a surface code”. *Nature Physics* **16**, 875–880 (2020).
- [98] Rajeev Acharya et al. “Suppressing quantum errors by scaling a surface code logical qubit”. *Nature* **614**, 676–681 (2023).
- [99] X. Gu, A. F. Kockum, A. Miranowicz, Y.-X. Liu, and F. Nori. “Microwave photonics with superconducting quantum circuits”. *Physics Reports* **718–719**, 1–102 (2017).
- [100] Neereja M. Sundaresan, Yanbing Liu, Darius Sadri, László J. Szócs, Devin L. Underwood, Moein Malekakhlagh, Hakan E. Türeci, and Andrew A. Houck. “Beyond Strong Coupling in a Multimode Cavity”. *Physical Review X* **5**, 021035 (2015).
- [101] Simon Storz, Josua Schär, Anatoly Kulkov, Paul Magnard, Philipp Kurpiers, Janis Lütolf, Theo Walter, Adrian Copetudo, Kevin Reuer, Abdulkadir Akin, Jean-Claude Besse, Mihai Gabureac, Graham J. Norris, Andrés Rosario, Ferran Martin, José Martinez, Waldimar Amaya, Morgan W. Mitchell, Carlos Abellan, Jean-Daniel Bancal, Nicolas Sangouard, Baptiste Royer, Alexandre Blais, and Andreas Wallraff. “Loophole-free Bell inequality violation with superconducting circuits”. *Nature* **617**, 265–270 (2023).
- [102] Xiu Gu, Jorge Fernández-Pendás, Pontus Vikstål, Tahereh Abad, Christopher Warren, Andreas Bengtsson, Giovanna Tancredi, Vitaly Shumeiko, Jonas Bylander, Göran Johansson, and Anton Frisk Kockum. “Fast Multiqubit Gates through Simultaneous Two-Qubit Gates”. *PRX Quantum* **2**, 040348 (2021).
- [103] Christopher W. Warren, Jorge Fernández-Pendás, Shahnawaz Ahmed, Tahereh Abad, Andreas Bengtsson, Janka Biznárová, Kamanaish Debnath, Xiu Gu, Christian Krizan, Amr Osman, Anita Fadavi Roudsari, Per Delsing, Göran Johansson, Anton Frisk Kockum, Giovanna Tancredi, and Jonas Bylander. “Extensive characterization and implementation of a family of three-qubit gates at the coherence limit”. *npj Quantum Information* **9**, 44 (2023).
- [104] J.R. Johansson, P.D. Nation, and Franco Nori. “QuTiP: An open-source Python framework for the dynamics of open quantum systems”. *Computer Physics Communications* **183**, 1760–1772 (2012).
- [105] J.R. Johansson, P.D. Nation, and Franco Nori. “QuTiP 2: A Python framework for the dynamics of open quantum systems”. *Computer Physics Communications* **184**, 1234–1240 (2013).
- [106] Neill Lambert, Eric Giguère, Paul Menczel, Boxi Li, Patrick Hopf, Gerardo Suárez, Marc Gali, Jake Lishman, Rushiraj Gadhvi, Rochisha Agarwal, Asier Galicia, Nathan Shammah, Paul Nation, J.R. Johansson, Shahnawaz Ahmed, Simon Cross, Alexander Pitchford, and Franco Nori. “Qutip 5: The quantum toolbox in python”. *Physics Reports* **1153**, 1–62 (2026).