



## On Unsolvable Equations of Prime Degree

Downloaded from: <https://research.chalmers.se>, 2026-06-18 03:41 UTC

Citation for the original published paper (version of record):

Brzezinski, J., Stevens, J. (2026). On Unsolvable Equations of Prime Degree. *American Mathematical Monthly*, In Press. <http://dx.doi.org/10.1080/00029890.2026.2660614>

N.B. When citing this work, cite the original published paper.



# The American Mathematical Monthly

ISSN: 0002-9890 (Print) 1930-0972 (Online) Journal homepage: [www.tandfonline.com/journals/uamm20](http://www.tandfonline.com/journals/uamm20)

## On Unsolvable Equations of Prime Degree

Juliusz Brzeziński & Jan Stevens

To cite this article: Juliusz Brzeziński & Jan Stevens (20 May 2026): On Unsolvable Equations of Prime Degree, The American Mathematical Monthly, DOI: [10.1080/00029890.2026.2660614](https://doi.org/10.1080/00029890.2026.2660614)

To link to this article: <https://doi.org/10.1080/00029890.2026.2660614>



© 2026 The Author(s). Published with license by Taylor & Francis Group, LLC.



Published online: 20 May 2026.



Submit your article to this journal [↗](#)



Article views: 349



View related articles [↗](#)



View Crossmark data [↗](#)

---

# On Unsolvable Equations of Prime Degree

---

Juliusz Brzeziński  and Jan Stevens 

---

 OPEN ACCESS

**Abstract.** Leopold Kronecker observed that either all roots or only one root of a solvable irreducible equation of odd prime degree with integer coefficients are real. This gives a possibility to construct specific examples of equations not solvable by radicals. A relatively elementary proof without using the full power of Galois theory is due to Heinrich Weber. We give a rather short proof of Kronecker's Theorem with an argument that is slightly different from Weber's. Several modern presentations of Weber's proof contain inaccuracies, which can be traced back to an error in the original proof. We discuss this error and how it can be corrected.

**Introduction.** One of the main objectives of any course in Galois theory is a proof of the existence of polynomials that can not be solved by radicals, showing that similar formulas as for quadratic, cubic, or quartic equations cannot exist for higher degree equations. Today this kind of problem mainly has a pedagogical and historical value. Galois theory is an extremely important part of mathematics and therefore one of the fundamental parts of mathematical education. Solvability of polynomial equations is used as an illustration of the efficacy of abstract algebraic methods in solving concrete, important, and interesting mathematical problems. The fact that most equations cannot be solved by radicals is usually presented using the full power of Galois theory: that is, the Galois correspondence and the relations between solvable equations and solvable groups. A hint that a more elementary approach is possible is given by the circumstance that the original impossibility proof of Niels-Henrik Abel predates the work of Évariste Galois.

Specific examples of unsolvable equations are easily obtained from the fact that for an irreducible equation of odd prime degree with integer coefficients all of its roots or only one are real, an observation made by Leopold Kronecker [1]. A relatively elementary proof, using rather limited knowledge related to the field extensions, was given by Heinrich Weber [2].

Weber's proof occurs in the algebra textbook [3] by Trygve Nagell. Thanks to this book in Swedish, the first author became aware of the possibility of proving the unsolvability of the quintic in an "elementary" way and transformed it into Exercise 13.6 in [4]. Unfortunately, the included solution is incorrect. We were led to a closer study of Nagell's proof and to a search for its origin. We found several recent presentations of Weber's proof in textbooks [5, 6], which are based on the English translation [7] of the popular account by Heinrich Dörrie [8]. Several objections to [7] and expositions following it have been raised; see [9, 10]. The ultimate source for the inaccuracies observed is an error in Weber's proof. The problem lies in reducible radicals (radicals of the form  $\alpha = \sqrt[q]{a}$  with  $X^q - a$  reducible). The need to deal with them also complicates the variant of the proof we developed.

---

[doi.org/10.1080/00029890.2026.2660614](https://doi.org/10.1080/00029890.2026.2660614)

MSC: 12F10, 12-03, 01A55

© 2026 The Author(s). Published with license by Taylor & Francis Group, LLC.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The terms on which this article has been published allow the posting of the Accepted Manuscript in a repository by the author(s) or with their consent.

In Section 1, we place Kronecker’s Theorem in a historical context. We discuss Kronecker’s arguments for the theorem and trace the history of “elementary” proofs. In Section 2, we discuss the criteria that make the proofs of Kronecker’s Theorem elementary. We also recall a number of algebraic results that, according to these criteria, are needed in its elementary proofs. Section 3 contains our proof of Kronecker’s Theorem. Finally, in Section 4, we discuss the mistake in Weber’s proof and in proofs inspired by his idea, as well as ways in which the proofs can be corrected.

**1. HISTORY.** After Gerolamo Cardano published the solution of cubic and quartic equations in 1545, attempts were made to find solutions of quintic equations, expressing their roots in a similar way as function of the coefficients, by formulas involving only arithmetical operations and radicals. The work of E. Waring, A.-T. Vandermonde and especially J.-L. Lagrange around 1770 suggested that such formulas for equations of degree 5 probably do not exist. This was proven to be so in the work of Paulo Ruffini and Niels Henrik Abel between 1799 and 1826, see the expository paper by Michael Rosen [11].

Much of the effort of Abel, Galois, and the mathematicians continuing their work concentrated on the characterization of the equations solvable by radicals. The problem was clearly formulated by Abel in his unfinished memoir “Sur la resolution algébrique des équations” [12]. He also indicated the way to arrive at its solution. According to Abel one should find all equations of a given degree that are algebraically solvable. In the course of his investigations he arrives at “several general propositions about the solvability of equations and about the form of their roots” [12, p. 219].

The investigation of the form of the roots of a solvable equation was taken up by Kronecker; see the interesting paper by Birgit Petri and Norbert Schappacher [13]. Three years after his first note on the subject [14], Kronecker observes in [1] the following:

Wenn eine irreductible Gleichung mit ganzzahligen Coëfficienten auflösbar und der Grad derselben eine ungrade Primzahl ist, so sind entweder *alle* ihre Wurzeln oder nur *eine* reell. <sup>1</sup>

This is a special case of the general result:

Wenn eine Gleichung — deren Grad eine ungrade Primzahl  $\mu$  ist, deren Coëfficienten rationale Functionen irgend welcher reeller Größen  $A, B, C, \dots$  also selbst reell sind und welche endlich nicht in Factoren niederen Grades zerlegt werden kann, so daß deren Coëfficienten wiederum rationale Functionen von  $A, B, C, \dots$  wären — durch eine explicite algebraische Function jener Größen  $A, B, C, \dots$  erfüllt wird, so sind entweder *alle* ihre Wurzeln, oder nur *eine* derselben reell. <sup>2</sup>

At the time Kronecker wrote this, the concept of field was not yet defined. With it Kronecker’s result can be formulated in the following way.

**Theorem 1 (Kronecker’s Theorem).** *Let  $K$  be a subfield of the real numbers. Suppose that an irreducible polynomial  $f(X) \in K[X]$  of odd prime degree  $p$  is solvable by radicals. Then exactly one root or all roots of the polynomial are real.*

As Kronecker clearly states, he found his results from the study of the form of the roots of solvable polynomials. This study was continued by Heinrich Weber [15],

<sup>1</sup>If an irreducible equation with integer coefficients is solvable and its degree is an odd prime, then either *all* of its roots or only *one* are real.

<sup>2</sup>If an equation—whose degree is an odd prime  $\mu$ , whose coefficients are rational functions of any real quantities  $A, B, C, \dots$  and therefore themselves real and which finally cannot be decomposed in factors of lower degree so that their coefficients are again rational functions of  $A, B, C, \dots$ —is satisfied by an explicit algebraic function of those quantities  $A, B, C, \dots$ , then either *all* of its roots, or only *one* of them are real.

Anders Wiman [16] and recently by Harold Edwards [17]. In the penultimate section of the first volume of his algebra textbook [15], Weber connects the number of real roots to the number of real roots of an auxiliary equation of degree  $p - 1$ , and this seems to establish Kronecker's Theorem, but Weber starts out from the result, referring to a simple proof earlier in his book. This simple proof was already mentioned by Kronecker himself [1]:

Ich bemerke ferner, daß die angegebene Eigenschaft der irreductibeln auflösbaren Gleichungen  $\mu$ ten Grades nicht bloß aus der allgemeinen Form ihrer Wurzeln hervorgeht, sondern auch aus dem schon von Galois herrührenden Satze "daß jede Wurzel einer solchen Gleichung sich als rationale Function von irgend zwei andern darstellen läßt". Wenn nämlich diese Function nur reelle Coefficienten enthält, so folgt hieraus unmittelbar, daß alle Wurzeln reell sein müssen, sobald nur zwei derselben reell sind.<sup>3</sup>

Galois' result can be formulated in modern language as follows.

**Theorem 2 (Galois' Theorem).** *An irreducible polynomial equation of prime degree  $p$  over a subfield of  $\mathbb{C}$  is solvable by radicals if and only if its splitting field is generated by any two of its roots.*

Kronecker's Theorem follows immediately, just as Kronecker remarked, since a solvable irreducible polynomial of prime degree  $p$  over a field  $K \subset \mathbb{R}$  having at least two real roots  $\alpha, \beta$  splits in the field  $K(\alpha, \beta)$ , so all its roots are real.

Galois proved this result in his famous "Mémoire sur les conditions de résolubilité des équations par radicaux" that was rejected in 1831 by the Académie des Sciences, and only published by Joseph Liouville in 1846, see [18]; for a proof see [4, Ex. 13.9]. S.-D. Poisson, the referee of the memoir, noted in his report (see [18, p. 146]) that an analogous proposition of Abel was printed after Abel's death in *Crelle's Journal* [19]. The excerpt in question of Abel's letter to A.L. Crelle from October 18, 1828 reads:

Si trois racines d'une équation quelconque irréductible, dont le degré est un nombre premier, sont liées entre elles de sorte que l'une de ces racines peut être exprimée rationnellement par les deux autres, l'équation en question sera toujours résoluble à l'aide de radicaux.<sup>4</sup>

It gives only a sufficient condition. The report goes on to say that no proof has been published and that Galois' proof of his necessary and sufficient condition is not satisfactory. Poisson complained that the memoir did not contain, as the title promised, the condition of solvability of equations by radicals. For, to decide whether a given equation of prime degree is solvable, one should first have to determine if this equation is irreducible, and then if one of its roots can be expressed as a rational function of two others. The condition, if it exists, should be verifiable by inspecting the coefficients of a given equation, or, at most, by solving other equations of lower degree (see [18, p. 148]).

Kronecker's Theorem gives a possibility to construct equations that are not solvable by radicals: any irreducible polynomial equation of odd prime degree with integral coefficients having more than one real root and also complex conjugate roots will do. One of the simplest examples is the polynomial  $X^5 - 4X - 2$ ; see Figure

<sup>3</sup>Furthermore I remark, that the stated property of irreducible solvable equations of degree  $\mu$  does not only follow from the general shape of their roots, but also from the Theorem due to Galois, "that every root of such an equation can be represented as a rational function of any two others." Namely, if this function only contains real coefficients, then it follows immediately that all roots must be real as soon as only two of them are real.

<sup>4</sup>If three roots of an arbitrary equation of which the degree is a prime number are related to each other in such a way that any one of these roots may be expressed rationally by the other two, the equation in question will always be solvable with the help of radicals.

1. The function  $f(X)$  has only two real extrema and the polynomial is irreducible by the Schönemann–Eisenstein criterion, published by Th. Schönemann in 1846 and G. Eisenstein in 1850 (see [20]); this criterion was of course unknown to Poisson.




Figure 1. The graph of  $f(X) = X^5 - 4X - 2$ .

A similar example is given in Ian Stewart’s textbook *Galois Theory* [21], but on the basis of Weber’s Theorem [15, §186], that an irreducible polynomial  $f(X) \in K[X] \subset \mathbb{R}[X]$  of prime degree  $p$  with  $p - 2$  real and 2 complex nonreal roots has as Galois group the symmetric group  $S_p$ , see [4, Ex. 13.4]. We note that Galois’ Theorem as stated above does not occur in [21], as in many other modern textbooks.

A proof of Kronecker’s Theorem was given by Weber in the first volume of Weber–Wellstein’s *Encyklopädie der Elementar-Mathematik* [2, § 101], originally for the case  $p = 5$ , and in later editions for general  $p$ . Elementary mathematics here means roughly the mathematics taught in secondary school, and this “handbook for teachers and students” is mainly directed at teachers. A fourth, heavily revised edition [22] was prepared by Paul Epstein. He made almost no changes in the chapter on solvable equations, but modernized the terminology.

Weber’s proof is suitable for a university course in algebra that treats the solution of equations, but not Galois theory. Such a course was given at Uppsala University by Trygve Nagell since 1931. In his textbook [3] Nagell treats the case  $p = 5$  following the first edition of Weber–Wellstein.

The insolubility of the quintic was included as one of the hundred problems in Heinrich Dörrie’s book *Triumph der Mathematik* [8], intended for a general audience. It contains a variety of problems, like Kirkman’s schoolgirl problem, or the length of the polar night, but also the transcendence of  $\pi$ . Dörrie was one of the first Ph.D students of David Hilbert with a dissertation on the quadratic reciprocity law in quadratic number fields with class number one. Dörrie’s book has been translated into Japanese, Hungarian, and English. The English translation [7] is unfortunately marred by strange terminology: *Körper* is not translated as field but as group, *Adjunktion* as substitution. For the insolubility of equations Dörrie follows the 1922 edition of Weber–Wellstein [22]. His version is also the basis for the expositions in [9], [6], and [5]. The last text inspired the proof in [10]. None of the sources mentioned cites Weber.<sup>5</sup>

## 2. PRELIMINARIES.

**What is an elementary proof?** Elementary certainly means avoiding deep general theory. In the context of Kronecker’s Theorem this excludes the fundamental theorem

<sup>5</sup>The latest version of the preprint [9] contains a footnote stating that Weber earlier gave the same proof.

of Galois theory. What can be used depends on the target group of the exposition. Almost a century ago Dörrie wrote for a general audience. Consequently he started his chapter on Kronecker's Theorem by explaining what a field (consisting of numbers) is and proved the facts about field extensions needed for Weber's proof, but avoided the concept of group. The proof uses the theory of field extensions within the scope defined by the contents of a second course in algebra at a university level. For our notion of elementary we have students taking such a course in mind. The considerations can always be limited to subfields  $K$  of the complex numbers  $\mathbb{C}$ , which we assume throughout this article. Thus, it is possible to include field extensions by the roots of polynomials (consequently, their splitting fields), the description of simple extensions, and the behavior of the degree according to the Tower law. It is essentially no problem to discuss automorphism groups of fields, since it is simply one more example of groups, which are always discussed in such courses. Thus, the notion of the Galois group of a polynomial  $f(X) \in K[X]$  as the group of all automorphisms of its splitting field  $K_f = K(\alpha_1, \dots, \alpha_n)$  that fix every element of  $K$ , where the  $\alpha_i$  are all the roots of  $f(X) = 0$  in  $\mathbb{C}$ , may also be accepted as elementary. This group we denote by  $G(K_f/K)$ . What is essential to avoid is the fundamental theorem of Galois theory (together with the Galois correspondence, in particular, between normal subgroups and normal extensions of the ground field), and as regards the notion of solvability, the correspondence between solvable groups and solvable extensions. But the notion of a solvable extension of fields is, of course, elementary. It is often presented as an example of the power of (abstract) algebraic language when one wants to formulate the problem of algebraic solvability and, in particular, when one wants to encourage the students to read a course in Galois theory in order to find out why there is no formula for an algebraic solution of a general quintic.

We can freely use modern concepts such as dimension and groups of field automorphisms, replacing the old fashioned arguments of Weber and Dörrie. We note that not only they, but also Nagel, deal with field automorphisms, looking at their actions on rational functions of the roots and considering "transformations" of these rather complicated expressions, and therefore with Galois groups, without mentioning them explicitly.

**Field extensions.** In this subsection, we recall several well-known results concerning the "elementary part" of the theory of field extensions. As a general reference for the notions and results that we mention, we use [4]. Let  $K \subseteq L$  be a field extension. An element  $\alpha \in L$  is *algebraic* over  $K$  if there is a nonzero polynomial  $f(X) \in K[X]$  such that  $f(\alpha) = 0$ . Zeros of univariate polynomials are called *roots*. Recall that a field extension  $K \subseteq L$  is called *finite* if  $L$  has finite dimension as a vector space over  $K$ . This dimension is denoted by  $[L : K]$  and is called the *degree* of  $L$  over  $K$ . The field extension  $K \subseteq L$  is *simple* if  $L$  is obtained by adjoining one element  $\alpha$ , so  $L$  is of the form  $K(\alpha)$ . If  $\alpha$  is algebraic over  $K$ , then the *minimal polynomial* of  $\alpha$  is the monic polynomial of smallest degree with coefficients in  $K$  that has  $\alpha$  as a root. For the reader's convenience and easier reference, we formulate two elementary properties of finite field extensions.

**Lemma 3 (Simple extension theorem [4, T.4.2]).** *If  $\alpha$  is algebraic over a field  $K$  and  $n$  is the degree of its minimal polynomial over  $K$ , then  $1, \alpha, \dots, \alpha^{n-1}$  is a basis of  $K(\alpha)$  over  $K$ . Thus  $[K(\alpha) : K] = n$  and each element in  $K(\alpha)$  can be uniquely represented as  $b_0 + b_1\alpha + \dots + b_{n-1}\alpha^{n-1}$ , where  $b_i \in K$ .*

**Lemma 4 (Tower law [4, T.4.2]).** *Let  $K \subseteq L$  and  $L \subseteq M$  be finite field extensions. Then  $K \subseteq M$  is a finite field extension and  $[M : K] = [M : L][L : K]$ .*

A *splitting field*  $L = K_f$  for a polynomial  $f(X) \in K[X]$  is a field  $L$  that contains all roots  $\alpha_1, \dots, \alpha_n$  of  $f$  and is generated over  $K$  by these roots, that is,  $L = K(\alpha_1, \dots, \alpha_n)$ . Recall that a field extension  $K \subseteq L$  is *normal* if every irreducible polynomial over  $K$  that has one root in  $L$  necessarily has all the remaining roots in  $L$ . A proof that splitting fields of polynomials over a field  $K$  coincide with finite normal extensions  $L$  of this field is usually not included in “elementary” algebra courses, but it is straightforward and uses only the structure of simple extensions and the Tower law together with basic properties of isomorphism of fields (see the proof [4, p. 120] of [4, T.7.1]).

Let  $K \subseteq L$  be a finite extension. The *Galois group* of  $L$  over  $K$ , denoted by  $G(L/K)$ , is the group of all automorphisms of  $L$  that fix  $K$  elementwise. If  $H$  is any subgroup of  $G(L/K)$ , then  $L^H$  is the subfield of  $L$  consisting of all elements, which are fixed by all automorphism belonging to  $H$ .

Two properties of isomorphisms of fields play an essential role in several arguments. The first is often proved in the context of a theorem usually attributed to Kronecker, stating that for any field and any non-constant polynomial, there exists an extension field containing a root of this polynomial. The second lemma is less known, but its proof is closely related to the first one.

**Lemma 5.** (a) *Let  $K$  be a field and let  $\alpha, \alpha'$  be two roots of an irreducible polynomial  $f(X) \in K[X]$ . Then there is an isomorphism  $\sigma: K(\alpha) \rightarrow K(\alpha')$  such that  $\sigma(\alpha) = \alpha'$  and  $\sigma$  is the identity on  $K$ .*

(b) *If  $\tau: K \rightarrow K'$  is an isomorphism of fields,  $L$  a splitting field of a polynomial  $f \in K[X]$ , and  $L'$  a splitting field of the polynomial  $\tau(f) \in K'[X]$ , then there exists an isomorphism  $\sigma: L \rightarrow L'$  extending  $\tau$  (that is,  $\sigma|_K = \tau$ ).*

For a proof of (a) we refer to [4, T.5.1], and for a proof of (b) to [4, T.5.2]. It follows by induction from [4, T.5.1], which is a more general version of (a), where  $\sigma$  extends an isomorphism  $\tau$ .

**Lemma 6.** *Let  $K \subset L$  be a normal extension and let  $\alpha, \alpha' \in L$  be two roots of an irreducible polynomial  $f(X) \in K[X]$ . Then there is an automorphism  $\sigma$  of  $L$  over  $K$  such that  $\sigma(\alpha) = \alpha'$ .*

We sketch the argument from [4, Ex. 7.4 (a)]. Since  $L$  is a finite and normal extension of  $K$ , it is a splitting field of a polynomial, say  $g(X) \in K[X]$ , and also a splitting field of the same polynomial over both  $K(\alpha)$  and  $K(\alpha')$ . By Lemma 5 (b) the isomorphism  $\tau: K(\alpha) \rightarrow K(\alpha')$  over  $K$  mapping  $\alpha$  onto  $\alpha'$  has an extension to an isomorphism of  $L$ .

**Remark 7.** Notice that Lemma 6 says that the automorphism group of a normal field extension  $L$  of a field  $K$  acts transitively on the roots of any irreducible polynomial in  $K[X]$  that has its roots in  $L$ . Moreover, it follows that  $L^{G(L/K)} = K$ : if  $\alpha$  and  $\alpha'$  are two different roots of the minimal polynomial  $f(X)$  of  $\alpha$  over  $K$ , then there is an automorphism  $\sigma$  of  $L$  over  $K$  such that  $\sigma(\alpha) = \alpha'$ . Therefore  $\alpha$  is not contained in the field  $L^{G(L/K)}$ , which of course contains  $K$ . This also applies to any intermediate field  $K \subseteq M \subseteq L$ , since  $L$  is also normal over  $M$  as the splitting field of the same polynomial whose splitting field over  $K$  is equal to  $L$ . This gives the “trivial” part of the Galois correspondence, which says that if we go from a subfield  $M$  to the subgroup  $G(L/M)$  and back to the corresponding field  $L^{G(L/M)}$ , then we come back to the same subfield  $M$ . In particular, this says that different subfields  $M$  go to different subgroups  $G(L/M)$  of  $G(L/K)$ . The difficult part of the Galois correspondence says that  $G(L/L^H) = H$  for every subgroup  $H$  of  $G(L/K)$ .

We are interested in algebraic solvability of polynomials, which means expressing the roots using arithmetical operations and radicals. The problem is best formulated in terms of fields. A *simple radical extension* is an extension  $K \subseteq K(\alpha)$ , where  $\alpha^q = a \in K$  for some positive integer  $q$ , that is,  $\alpha$  is a root of the binomial polynomial  $X^q - a$ . A field extension  $K \subseteq L$  is *radical* if there is a chain

$$K = K_0 \subseteq K_1 \subseteq \cdots \subseteq K_{i-1} \subseteq K_i \subseteq \cdots \subseteq K_n = L \quad (1)$$

of simple radical extensions  $K_i = K_{i-1}(\alpha_i)$ . We may assume that the  $q_i$  are prime numbers, by using the fact that  $\sqrt[q]{\sqrt[s]{a}} = \sqrt[r]{\sqrt[s]{a}}$ . A polynomial  $f(X) \in K[X]$  is *solvable by radicals* if a splitting field of  $f(X)$  is contained in a radical extension  $K \subseteq L$  [4, p. 78].

We stress the fact that in the definition of a simple radical extension we do not assume that the polynomial  $X^q - \alpha^q$  is irreducible over  $K$ . If  $X^q - \alpha^q$  is irreducible, then  $\alpha = \sqrt[q]{a}$  is called an *irreducible radical*. Otherwise, that is, if  $X^q - a$  is reducible over  $K$ , then any  $\alpha = \sqrt[q]{a}$  is called a *reducible radical*. A polynomial  $f(X) \in K[X]$  is *solvable by irreducible radicals* if  $K_f \subseteq L$ , where  $K \subseteq L$  is a radical extension such that in the chain of simple radical extensions every  $X^{q_i} - \alpha_i^{q_i}$  is irreducible over  $K_{i-1}$ . From the (constructive) point of view of writing down radical expressions for the roots of an equation it is quite natural to require irreducibility in the definition of solvability. But this gives the same class of polynomials: a solvable polynomial is also solvable by irreducible radicals, see e.g., [21, Ch. 21].

In the following sections, we need a few properties of the splitting fields of the binomials  $X^q - a$ . We gather them in the following lemma.

**Lemma 8.** *Let  $K$  be a subfield of  $\mathbb{C}$ ,  $q$  a prime number, and  $a \in K \setminus \{0\}$ .*

- (a) *The splitting field of  $X^q - a \in K[X]$  over  $K$  is  $K' = K(\alpha, \varepsilon)$ , where  $\alpha \in K'$  is a root of  $X^q - a$  and  $\varepsilon$  a primitive  $q$ -th root of unity.*
- (b) *The binomial  $X^q - a$  is reducible over  $K$  if and only if it has a root in  $K$ .*
- (c) *If  $\varepsilon \in K$  and  $X^q - a$  is irreducible over  $K$ , then  $K' = K(\alpha)$  and  $[K' : K] = q$ . All automorphisms of  $K'$  over  $K$  are mappings  $\varphi(\alpha) = \varepsilon^r \alpha$ ,  $r = 0, 1, \dots, q - 1$ . Thus the Galois group is the cyclic group  $G(K'/K) = \mathbb{Z}_q$ .*
- (d) *If  $\varepsilon \notin K$  and  $X^q - a$  is reducible over  $K$ , then  $K' = K(\varepsilon)$  is the splitting field of the polynomial  $X^q - 1$ . Every automorphism of  $K'$  maps  $\varepsilon$  onto a power of  $\varepsilon$ .*

We do not present detailed proofs, but we comment on the points above.

(a) The equation  $X^q - a = 0$  has  $q$  different solutions  $\varepsilon^r \sqrt[q]{a}$ , where  $\varepsilon \neq 1$  denotes a (primitive)  $q$ -th root of unity,  $r = 0, \dots, q - 1$  and  $\alpha = \sqrt[q]{a}$  is any fixed solution to the equation  $X^q - a = 0$ . Thus the splitting field of the binomial  $X^q - a$  over  $K$  is equal to  $K' = K(\alpha, \varepsilon)$  ([4, Ex. 5.11 (c)]).

(b) This is a result of Abel. The fact that the polynomial  $X^q - a$  is reducible over  $K$  if and only if  $a$  is a  $q$ -th power in  $K$  is proved in [4, Ex. 5.12 (a)].

(c) If  $\varepsilon \in K$  and  $X^q - a$  is irreducible, then according to (b),  $\alpha \notin K$ , since  $a$  is not a  $q$ -th power in  $K$  ( $\alpha^q = a$ ). The extension  $K' = K(\alpha)$  of  $K$  is simple, of degree  $q$  by the Simple extension theorem (Lemma 3). According to (a), the polynomial  $X^q - a$  has  $q$  different roots  $\varepsilon^r \alpha$  for  $r = 0, 1, \dots, q - 1$ . Every automorphism of  $K'$  maps  $\alpha$  on a root of the same polynomial. Thus, we have  $q$  different automorphisms mapping  $\alpha$  onto  $\varepsilon^r \alpha$ . The exponents  $r$  form the group  $\mathbb{Z}_q$  of all residues  $r$  modulo  $q$  (as  $\varepsilon^q = 1$ ).

(d) If  $X^q - a$  has a root  $\alpha \in K$ , then it follows from (a) that  $K' = K(\varepsilon)$ , where as before  $\varepsilon$  denotes a primitive  $q$ -th root of unity. Thus  $K'$  is a splitting field of the polynomial  $X^q - 1$ , since all  $q$ -th roots of unity are powers of  $\varepsilon$ . The minimal polynomial of  $\varepsilon$  over  $K$  divides the minimal polynomial of  $\varepsilon$  over  $\mathbb{Q}$ , which has degree  $q - 1$ ; it

is  $(X^q - 1)/(X - 1)$  ([4, Ex. 4.3 (f)]; for the history of this result see [20]). Every automorphism maps  $\varepsilon$  onto another  $q$ -th root of unity, which is a power  $\varepsilon^k$  for some  $1 \leq k \leq q - 1$ .

**Auxiliary results.** The proofs of the following results, which are needed for Weber's and our proof of Kronecker's Theorem, do not depend on Galois theory.

**Lemma 9 (Nagell's Lemma).** *If  $f \in K[X]$  has prime degree  $p$  and is irreducible over  $K$ , but reducible over a field extension  $K \subset L$ , then  $p \mid [L : K]$ .*

*Proof.* Let  $r = [L : K]$  and let  $\alpha$  be a root of  $f(X)$  in a field extension of  $L$ . As  $f(X)$  is reducible in  $L$ , we have  $[L(\alpha) : L] = m < p = [K(\alpha) : K]$ . Because  $L(\alpha)$  is an extension of  $K(\alpha)$ ,  $p = [K(\alpha) : K] \mid [L(\alpha) : K] = mr$ , where the last equality holds by the Tower law (Lemma 4). Therefore  $p \mid r$ . ■

The lemma is formulated differently in [3, §98, p. 259] and [4, Ex. 4.13], but the proofs given there establish the present result. In fact, it follows from the general fact that if the degree of an irreducible polynomial in  $K[X]$  and the degree of a finite field extension  $L$  of  $K$  are relatively prime, then the polynomial remains irreducible over  $L$  (see [4, Ex. 4.2]). Weber [22] and Dörrie [8] use Proposition 16 with the same effect.

The following lemma can be found in [4, Ex. 7.8]. It only depends on Lemma 5 and the Tower law.

**Lemma 10.** *Let  $K \subset L$  be a normal extension and let  $f(X)$  be a monic polynomial irreducible over  $K$  but reducible over  $L$ . Then all irreducible factors of  $f(X)$  in  $L[X]$  have the same degree.*

*Proof.* If  $\alpha_i$  and  $\alpha_j$  are roots (in an extension of  $L$ ) of two irreducible factors  $f_i(X)$  and  $f_j(X)$  of  $f(X)$  over  $L$ , then there is an isomorphism  $\tau : K(\alpha_i) \rightarrow K(\alpha_j)$  over  $K$ , by Lemma 5 (a) (since both  $\alpha_i, \alpha_j$  are roots of  $f(X)$ ). Let  $L$  be a splitting field of a polynomial  $g(X) \in K[X]$ . Then  $L(\alpha_i)$  is a splitting field of  $g(X)$  over  $K(\alpha_i)$ , and  $L(\alpha_j)$  is a splitting field of  $g(X)$  over  $K(\alpha_j)$ . By Lemma 5 (b) this automorphism can be extended to an isomorphism  $\sigma : L(\alpha_i) \rightarrow L(\alpha_j)$  over  $K$ . Therefore the degrees  $[L(\alpha_i) : K]$  and  $[L(\alpha_j) : K]$  are equal and thus also the degrees  $[L(\alpha_i) : L]$  and  $[L(\alpha_j) : L]$  are equal, which are just the degrees of  $f_i(X)$  and  $f_j(X)$ . ■

Kronecker's Theorem is about the solvability of polynomials with real coefficients. We need terminology and results adapted to this situation. We say that a field  $K \subseteq \mathbb{C}$  is *conjugation invariant* if complex conjugation is an automorphism of this field. Notice that if  $K$  is a conjugation invariant field, then its extension  $K(\alpha)$ , where  $\alpha \in \mathbb{C}$ , is conjugation invariant if and only if  $\bar{\alpha} \in K(\alpha)$ .

**Lemma 11.** *Let  $K \subseteq L$  be a radical extension of a conjugation invariant field  $K$ . Then there exists a chain  $K = K_0 \subseteq K_1 \subseteq \dots \subseteq K_m$  of simple radical extensions (of prime degree) with  $L \subseteq K_m$ , where all fields  $K_i$  are conjugation invariant.*

*Proof.* We use induction with respect to the number  $n$  of different simple radical extensions between  $K$  and  $L$ . As already noted, we may assume that every simple radical extension is by a root of a binomial of prime degree. Assume that the Lemma is true for every radical extension of  $K$  by such a chain of at most  $n \geq 0$  simple radical extensions. The base case, when  $L = K$  (that is,  $n = 0$ ), is trivially true.

Let  $L$  be a radical extension of  $K$  by a chain of  $n + 1$  simple radical extensions (each by a root of a binomial of prime degree). Let  $L' \subset L$  be the last subextension in the chain. Then  $K \subset L'(\alpha) = L$ , where  $\alpha^q = a \in L'$  for a prime number  $q$ . The field

$L'$  is a radical extension of  $K$  by  $n - 1$  simple radical extensions. By the induction hypothesis, we have  $L' \subset K_{m'}$ , where  $K = K_0 \subset K_1 \subset \cdots \subset K_{m'}$  is a chain of simple radical extensions with all fields  $K_i$  conjugation invariant and  $K_i = K_{i-1}(\alpha_i)$  with  $\alpha_i^{q_i} = a_i \in K_{i-1}$  and  $q_i$  a prime number for  $i = 1, \dots, m'$ .

We construct a chain for  $L$ . We have that  $L \subset K_{m'}(\alpha)$ . If  $\bar{\alpha} \in K_{m'}(\alpha)$ , then  $K_{m'}(\alpha)$  is conjugation invariant and we obtain the required chain by putting  $K_{m'+1} = K_{m'}(\alpha)$ . If  $\bar{\alpha} \notin K_{m'}(\alpha)$  we consider the real number  $\rho = \alpha\bar{\alpha}$ . The fields  $K_{m'+1} = K_{m'}(\rho)$  and  $K_{m'+2} = K_{m'+1}(\alpha) = K_{m'}(\rho, \alpha) = K_{m'}(\alpha, \bar{\alpha})$  are conjugation invariant. Moreover, we have  $\rho^q = \alpha^q \bar{\alpha}^q = a\bar{a} \in K_{m'}$  and  $\alpha^q = a \in L' \subseteq K_{m'} \subseteq K_{m'+1}$ , which shows that  $K_{m'} \subseteq K_{m'+1}$  and  $K_{m'+1} \subseteq K_{m'+2}$  are radical extensions by roots of prime degree binomials. Consider now the chain

$$K = K_0 \subseteq \cdots \subseteq K_{m'} \subseteq K_{m'+1} \subseteq K_{m'+2}.$$

Clearly, we have that  $L = L'(\alpha) \subseteq K_{m'}(\alpha) \subseteq K_{m'}(\alpha, \bar{\alpha}) = K_{m'+2}$ . If we have an equality of fields in the chain, we can simply remove repeating copies. The constructed chain satisfies the induction step. ■

The number  $\rho$  in the above proof plays an important role in Weber's proof. In the following lemma we collect some properties of it, which easily follow from Lemma 8.

**Lemma 12.** *Let  $K$  be a conjugation invariant field and consider the prime degree binomial  $X^q - a \in K[X]$ . The real root  $\rho = \sqrt[q]{a\bar{a}}$  satisfies  $\rho = \alpha\bar{\alpha}$  for any root  $\alpha$  of  $X^q - a$ . Therefore  $\rho \in K(\alpha, \bar{\alpha})$ , and if  $X^q - a$  is reducible, that is, has a root in  $K$ , then  $\rho \in K$ .*

**Remark 13.** Lemma 11 is not formulated as such by Weber, but the idea of the construction is due to him. Weber [2, 22] just says that he adjoins  $\alpha$  and  $\bar{\alpha}$  at the same time. He considers this clearly as one step, and only when needed he first adjoins  $\rho$  and then  $\alpha$ . In particular, the first adjunction in a chain that makes  $f(X)$  reducible is an extension  $L \subset L(\alpha, \bar{\alpha})$  of conjugation invariant fields, which subsequently is studied in greater detail. The same holds for Dörrie [8]. The objection that sometimes has been raised to his proof, that not all extensions in the chain are conjugation invariant, is therefore not founded.

**3. A PROOF OF KRONECKER'S THEOREM.** In this section, we give a proof of Kronecker's Theorem, which, in principle allows us to treat both reducible and irreducible radicals using the same arguments. However, there is one point where a deeper knowledge of Galois theory for cyclic field extensions seems to be needed for such a unification. We comment on this after the proof but we choose to give an elementary proof, where we consider the two cases (irreducible and reducible) separately. For the second case we use an argument that more closely follows Weber's ideas. We formulate Kronecker's Theorem for polynomials with real coefficients in a conjugation invariant field.

**Theorem 14 (Kronecker's Theorem).** *Let  $K$  be a conjugation invariant subfield of the complex numbers. Suppose that an irreducible polynomial equation  $f(X) \in K[X]$  of odd prime degree  $p$  with real coefficients is solvable by radicals over  $K$ . Then exactly one root or all roots of the polynomial are real.*

*Proof.* We first remark that  $f(X)$  has at least one real root, since it has real coefficients. As  $f(X)$  is solvable, we may assume by Lemma 11 that there is a conjugation invariant chain  $K = K_0 \subseteq K_1 \subseteq \cdots \subseteq K_n = L$  of simple radical extensions such that  $K_i$  is an

extension of  $K_{i-1}$  by a root of binomial of prime degree for  $i = 1, \dots, n$  and  $K_f \subseteq L$ . There exists an  $i$  such that  $f(X)$  is irreducible in  $K_{i-1}$  but reducible in  $K_i = K_{i-1}(\alpha_i)$ , where  $\alpha_i^q = a_i \in K_{i-1}$  and  $q$  is a prime number. For ease of notation we write  $\alpha$  for  $\alpha_i$  and  $a$  for  $a_i$ . By Nagell's Lemma 9 the degree  $p$  of  $f(X)$  divides  $d = [K_i : K_{i-1}]$ . We distinguish between two cases depending on whether  $X^q - a$  is irreducible or  $X^q - a$  is reducible over  $K_{i-1}$ .

*Case I.* If  $X^q - a$  is irreducible over  $K_{i-1}$ , then  $d = q$  is prime, which implies that  $q = p$ . We may assume that  $K_{i-1}$  contains a primitive  $p$ -th root of unity  $\varepsilon_p$ , for otherwise we can replace  $K_{i-1}$  by  $K_{i-1}(\varepsilon_p)$ . Both polynomials  $X^p - a$  and  $f(X)$  remain irreducible over  $K_{i-1}(\varepsilon_p)$  by Nagell's Lemma, as the degree  $p$  is relatively prime to the degree of the minimal polynomial of  $\varepsilon_p$  over  $K_{i-1}$ . In fact, this polynomial divides  $X^{p-1} + X^{p-2} + \dots + 1$  (the minimal polynomial of  $\varepsilon_p$  over  $\mathbb{Q}$ ), so its degree is at most equal to  $p - 1$ . Thus  $K_i = K_{i-1}(\alpha)$  is a splitting field of the polynomial  $X^p - a$  (see Lemma 8 (c)), which says that the extension  $K_{i-1} \subset K_i$  is normal. Notice that in this case  $[K_i : K_{i-1}] = p$ .

Since  $f(X)$  is reducible over  $K_i$  (and irreducible over  $K_{i-1}$ ), the degrees of all irreducible factors of  $f(X)$  are equal by Lemma 10. Thus the factors of  $f(X)$  over  $K_i$  are linear and all roots  $\beta_1, \dots, \beta_p$  of  $f(X)$  belong to  $K_i$ . Hence, the splitting field  $(K_{i-1})_f = K_{i-1}(\beta_1, \beta_2, \dots, \beta_p)$  of  $f(X)$  over  $K_{i-1}$  is a subfield of  $K_i$ . For every  $r = 1, \dots, p$ , we have  $[K_{i-1}(\beta_r) : K_{i-1}] = p$ . It implies that  $K_{i-1}(\beta_r) = K_i$ , since also  $K_i$  has degree  $p$  over  $K_{i-1}$ . Thus  $K_i = (K_{i-1})_f = K_{i-1}(\beta_r)$  for every  $r = 1, \dots, p$ .

Suppose that at least two roots,  $\beta_1$  and  $\beta_2$ , are real. By Lemma 8 (c) the Galois group  $G((K_{i-1})_f/K_{i-1})$  is cyclic of order  $p$ . Every nontrivial automorphism has order  $p$  and since the group acts transitively on the roots  $\beta_r$  (see Lemma 6), its elements are cycles of length  $p$ . We may choose as a generator the cycle  $\sigma = (\beta_1, \beta_2, \dots, \beta_p)$  starting with the real roots  $\beta_1, \beta_2$ . Because the powers  $\beta_1^j$  form a basis of  $(K_{i-1})_f$  over  $K_{i-1}$  (Lemma 3), we can write

$$\beta_2 = c_0 + c_1\beta_1 + \dots + c_{p-1}\beta_1^{p-1}, \quad (2)$$

with uniquely determined coefficients  $c_j \in K_{i-1}$ . Since  $\beta_1$  and  $\beta_2$  are real, complex conjugation gives

$$\beta_2 = \bar{c}_0 + \bar{c}_1\beta_1 + \dots + \bar{c}_{p-1}\beta_1^{p-1},$$

so  $\bar{c}_i = c_i$  for  $i = 0, 1, \dots, p - 1$ . Thus all the coefficients  $c_i$  are real.

Applying the automorphism  $\sigma$  to equation (2), we get

$$\beta_3 = c_0 + c_1\beta_2 + \dots + c_{p-1}\beta_2^{p-1},$$

showing that  $\beta_3$  is also real. By repeating this last argument, we find that all roots of  $f(X)$  are real, if more than one root is real.

*Case II.* If  $X^q - a$  is reducible, then  $K_i = K_{i-1}(\alpha) = K_{i-1}(\varepsilon_q)$  for a primitive  $q$ -th root of unity according to Lemma 8 (d). In order to simplify notation, we denote  $\varepsilon_q$  by  $\varepsilon$ . Also in this case, the extension  $K_{i-1} \subset K_i$  is normal as a splitting field of  $X^q - 1$  and  $[K_i : K_{i-1}] = d > 1$ .

Since the extension  $K_{i-1} \subset K_i$  is normal and  $f(X)$  is reducible over  $K_i$  (and irreducible over  $K_{i-1}$ ), we get as in Case I that all roots  $\beta_1, \dots, \beta_p$  of  $f(X)$  belong to  $K_i$ . Hence the splitting field  $(K_{i-1})_f = K_{i-1}(\beta_1, \beta_2, \dots, \beta_p)$  of  $f(X)$  over  $K_{i-1}$  is a subfield of  $K_i$ . We work in the field  $K_i$  instead of  $(K_{i-1})_f$ , which we did in Case I.

As we know, the polynomial  $f(X)$  has a real root. Assume that  $\beta_1$  is real. Since  $K_i = K_{i-1}(\alpha)$ , we can write

$$\beta_1 = c_0 + c_1\alpha + \dots + c_{d-1}\alpha^{d-1} = \psi(\alpha), \quad (3)$$

where  $\psi(X) = c_0 + c_1X + \cdots + c_{d-1}X^{d-1} \in K[X]$ . By Lemma 12 we have  $\rho = \sqrt[d]{a\bar{a}} \in K$ , and  $\rho = \alpha_i \bar{\alpha}_i$  for all roots  $\alpha_i = \varepsilon^i \alpha$  of  $X^q - a$ . We conjugate the equality (3). We write  $\bar{\psi}(X)$  for the polynomial in  $K[X]$  with coefficients the complex conjugates of the coefficients of the polynomial  $\psi(X)$ . As  $\beta_1$  is real, we get  $\psi(\alpha) = \bar{\psi}(\alpha) = \bar{\psi}(\bar{\alpha}) = \bar{\psi}(\frac{\rho}{\alpha})$ . In the equation  $\psi(\alpha) = \bar{\psi}(\frac{\rho}{\alpha})$  only  $\alpha$  does not lie in  $K$ . Thus the equality  $\psi(\alpha) = \bar{\psi}(\frac{\rho}{\alpha})$  still holds when  $\alpha$  is replaced by another root, which is the image of  $\alpha$  by any automorphism of the field  $K_{i-1}(\alpha) = K_{i-1}(\varepsilon)$ . From Lemma 8 (d) it follows that every automorphism of this field maps  $\varepsilon$  onto a power of this number, and  $\alpha$  onto another root of  $X^q - a$ . At the same time such an automorphism maps  $\beta_1$  onto another root of the polynomial  $f(X)$ . Since this polynomial is irreducible over  $K_{i-1}$ , any root can be mapped at any other root (see Lemma 6), so the equality  $\beta_r = \bar{\beta}_r$  is valid for all roots of  $f(X)$ , showing that they are real numbers. ■

**Remark 15.** In the proof above we have the situation that all roots  $\beta_1, \dots, \beta_p$  of an irreducible polynomial  $f(X) \in K[X]$  generate subfields  $K(\beta_i)$  of an extension field  $L \supset K$  of the same degree  $p$  over  $K$ . In Case I, we conclude that all these fields are equal (and equal to  $L$ ), since also  $L$  has degree  $p$  over  $K$ . In case II, it is not possible to claim the equality of all subfields  $K(\beta_i)$  using the same argument. But assuming a little more knowledge than we needed in Case II, we can establish equality. Also in this case, the extension  $K \subset L$  is normal and cyclic, so the Galois group is cyclic. A known property of finite cyclic groups says that for every divisor  $\delta$  of the group order, there is exactly one subgroup whose order is equal to  $\delta$  (see [4, Prop. A.2.2 (b)]). Since for any two roots  $\beta_i$  and  $\beta_j$  there is an automorphism  $\sigma$  of  $L$  over  $K$  such that  $\sigma(\beta_i) = \beta_j$  (see Lemma 6), the groups  $G(L/K(\beta_i))$  and  $G(L/K(\beta_j))$  are isomorphic (in fact, conjugated by  $\sigma$ —see [4, Prop. A.9.2]), so they have the same order  $\delta$ . Thus the group  $H = G(L/K(\beta_i))$  is the same for all  $i$ . The “trivial part” of Galois correspondence (see Remark 7) says that for intermediate fields  $M$  of a Galois extension  $K \subseteq L$ , we have  $L^{G(L/M)} = M$ . Applying this to  $M = K(\beta_i)$ , we get  $K(\beta_i) = L^H$ . Thus all the fields  $K(\beta_i)$  corresponding to the roots of  $f(X)$  are equal, and we can use the argument of Case I. Treating in this way both cases simultaneously results in a unified and short proof of Kronecker’s Theorem.

**4. WEBER’S PROOF.** In this Section, we return to the history of “elementary” proofs and discuss the errors that such proofs had until recently, and proposed corrections.

As our proof is inspired by the presentation of Nagell [3], it follows the main lines of Weber’s proof. Given a chain  $K = K_0 \subset K_1 \subset \cdots \subset K_n = L$  of simple radical extensions, the essential point is to study what happens in the first extension  $K_{i-1} \subset K_i$  such that the solvable polynomial  $f(X)$  of degree  $p$  is irreducible in  $K_{i-1}$  but reducible in  $K_i$ , where  $K_i = K_{i-1}(\alpha)$  with  $\alpha$  a root of the binomial  $X^q - a$ ,  $q$  prime. Weber (and Epstein) [2, 22] as well as Nagell [3] claim that  $q = p$ , but this does not hold if  $X^q - a$  is reducible over  $K_{i-1}$ .

The main point in Weber’s proof (for the irreducible case) that differs from our proof, is the following. As before, it can be assumed that  $\varepsilon := \varepsilon_p \in K_{i-1}$ . The real root  $\beta_1$  can be written as

$$\beta_1 = c_0 + c_1\alpha + \cdots + c_{p-1}\alpha^{p-1} = \psi(\alpha), \quad (4)$$

where  $\psi(X) = c_0 + c_1X + \cdots + c_{p-1}X^{p-1} \in K_{i-1}(X)$  (Lemma 3). Weber distinguishes two cases depending on whether  $\alpha \in \mathbb{R}$  or  $\alpha \notin \mathbb{R}$ .

If  $\alpha \in \mathbb{R}$ , then complex conjugation gives that  $\bar{c}_i = c_i$ , or in other words, all  $c_i$  are real, by uniqueness of the coefficients in equation (4). As the automorphisms of the

field  $K_i$  over  $K_{i-1}$  map  $\alpha$  onto  $\varepsilon^r \alpha$  for  $r = 0, \dots, p-1$ , the other roots of  $f(X)$  are

$$\beta_r = c_0 + c_1 \varepsilon^r \alpha + \dots + c_{p-1} \varepsilon^{r(p-1)} \alpha^{p-1},$$

so they come in complex conjugate pairs and  $\beta_1$  is the only real root.

If  $\alpha \notin \mathbb{R}$ , then consider  $K_{i-1}(\rho)$  with  $\rho = \sqrt[p]{a\bar{\alpha}}$  (by construction  $K_{i-1}$  is conjugation invariant, see Remark 13). If  $f(X)$  is reducible in  $K_{i-1}(\rho)$ , we are back in the previous case. Otherwise  $f(X)$  becomes reducible in the extension  $K_{i-1}(\rho) \subset K_{i-1}(\rho, \alpha)$ . Then complex conjugation gives  $\beta_1 = \psi(\alpha) = \bar{\psi}(\bar{\alpha}) = \bar{\psi}(\frac{\rho}{\alpha})$  and Weber argues that, as  $\bar{\alpha}_s = \frac{\rho}{\alpha_s}$  for any root of  $X^p - a$ , the equality  $\psi(\alpha) = \bar{\psi}(\bar{\alpha})$  still holds when  $\alpha$  is replaced by another root when we apply the automorphism mapping  $\beta_1$  onto this root (that is, we map  $\alpha$  onto  $\varepsilon^s \alpha$ ). Therefore all roots are real.

The wrong claim  $q = p$  is based on an application of the following variant of Nagell's Lemma 9, which is proved by Weber [22, § 111, 5] and Dörrie [8, § 24, Satz IV]. Interestingly, the proof of this relatively elementary result by both Weber and Dörrie does not use the concept of the degree of a field extension, which makes it relatively long and somewhat complicated. Nagell [3] gives a short proof of Lemma 9, using similar arguments to ours.

**Proposition 16.** *An irreducible polynomial  $f(X)$  of prime degree  $p$  can only become reducible through adjunction of a root of an [irreducible] equation  $\varphi(X) = 0$ , whose degree is divisible by  $p$ .*

We write the word *irreducible* in brackets, because it occurs in [8], but it is left out in [22], although the proof in [22] uses the assumption that the equation  $\varphi(X) = 0$  is irreducible. In [22] the Proposition is applied to the equation  $X^q = a$ , with the conclusion that  $q = p$ . Irreducibility of this equation is not assumed, not even implicitly, as the next step argues that  $a$  is not the  $p$ -th power of a number  $\beta \in K_{i-1}$ , because otherwise  $\alpha$  has the form  $\varepsilon^k \beta$  and would belong to  $K_{i-1}$ . This is clearly a circular argument.

Dörrie [8] apparently noticed the error. To be able to use the correctly stated Proposition 16 he requires the radicals in the chain to be irreducible. Dörrie does not realize that he might introduce reducible radicals when making the chain conjugation invariant:

Ferner wollen wir mit jedem adjungierten Radikal unsere Kette, das noch keine Zerlegung von  $f(x)$  ermöglicht, auch gleich das komplex konjugierte Radikal adjungieren. Das ist vielleicht überflüssig, sicher aber nicht schädlich.<sup>6</sup>

But actually harm can be done, as the following example shows.

**Example 17.** Let  $\zeta$  be a primitive 7-th root of unity and let  $f(X)$  be the minimal polynomial of the real number  $\beta = \zeta + \bar{\zeta}$ . One easily computes that

$$f(X) = X^3 + X^2 - 2X - 1.$$

In fact,  $\beta$  generates the maximal real subfield of the cyclotomic field  $\mathbb{Q}(\zeta)$  whose degree over  $\mathbb{Q}$  is equal to 6. Thus its maximal real subfield has degree 3 over  $\mathbb{Q}$ . A similar construction is possible for any prime number  $p$  such that  $2p+1$  is also a prime number (Sophie Germain primes).

Let a chain of simple radical extensions start with  $K_0 = \mathbb{Q} \subset K_1 = \mathbb{Q}(\alpha)$ , where e.g.,  $\alpha = \zeta \sqrt[7]{11}$  and  $\sqrt[7]{11}$  is the real root. Then  $\alpha$  is an irreducible radical (the root of

<sup>6</sup>Furthermore, with each adjoined radical of our chain, which not yet makes it possible to factorize  $f(x)$ , we will also adjoin at the same time the complex conjugate radical. That is maybe superfluous, but certainly not harmful.

$X^7 - 11 = 0$ ) and the polynomial  $f(X)$  of degree 3 is irreducible over  $K_1$  by Nagell's Lemma 9, as 3 does not divide the degree  $[K_1 : \mathbb{Q}] = 7$ . But when we extend  $K_1 = \mathbb{Q}(\alpha)$  by the complex conjugate  $\bar{\alpha} = \zeta^6 \sqrt[7]{11}$ , then  $\bar{\alpha}$  is a reducible radical over  $K_1$  (it satisfies the equation  $X^7 - 11 = 0$  and the polynomial  $X^7 - 11$  is reducible over  $K_1$ ). Moreover, we have  $\zeta \in K_2 = \mathbb{Q}(\alpha, \bar{\alpha})$  as  $\alpha/\bar{\alpha} = \zeta^2$  and  $\zeta^2$  is a primitive 7-th root of unity. Hence  $\beta = \zeta + \bar{\zeta} \in \mathbb{Q}(\alpha, \bar{\alpha})$  and  $f(X)$  is reducible over this field. According to Weber and Dörrie and the construction in Lemma 11, we should in this case first adjoin  $\rho = \alpha\bar{\alpha} = \sqrt[7]{121}$  and then  $\alpha$ , but  $X^7 - 11$  is also reducible over  $\mathbb{Q}(\sqrt[7]{121})$ , as  $\sqrt[7]{11} = (\sqrt[7]{121})^4/11$  is a root in  $\mathbb{Q}(\sqrt[7]{121})$ .

It is possible to express  $\beta$  in terms of  $\alpha$ , which Case II in the proof of Theorem 14 suggests, but it is easier to adjoin  $\zeta$  in the first place, something that is allowed according to Weber and Nagell. The polynomial  $f(X)$  becomes reducible by the conjugation invariant extension  $\mathbb{Q} \subset \mathbb{Q}(\zeta)$  of degree 6. The automorphism  $\sigma_k$  of  $\mathbb{Q}(\zeta)$  that sends  $\zeta$  to  $\zeta^k$ , maps  $\beta = \zeta + \zeta^6$  to  $\beta_k = \zeta^k + \zeta^{7-k}$ , for  $k = 1, \dots, 6$ . As  $\beta_k = \beta_{7-k}$ , we obtain all three roots of  $f(X)$  twice.

It is easy to solve the equation  $f(X) = 0$  using a computer algebra system. The three roots are

$$\frac{\sqrt[3]{28 + 84\sqrt{-3}}}{6} + \frac{14}{3\sqrt[3]{28 + 84\sqrt{-3}}} + \frac{1}{3},$$

where one should always take the same values of the multiply occurring square and cubic roots. This answer avoids the ambiguities in Cardano's formulas. We note that the primitive 7-th root of unity  $\zeta$  is a root of the polynomial  $X^2 - \beta X + 1$ , but we refrain from giving an explicit expression for  $\zeta$ .

Weber's and Dörrie's argument can be rescued if the adjunction of a reducible radical making the polynomial  $f(X)$  reducible is replaced by a chain of adjunctions of irreducible radicals, using Gauss' result that the roots of unity are expressible by irreducible radicals [21, Ch. 21]. This is the approach chosen by Yan Pan and Yuzhen Chen [9] and A.B. Skopenkov [10], who pointed out an error related to reducibility in the proof by V.V. Prasolov [5]. Instead of the conjugation invariant chain provided by our Lemma 11, they construct a chain where in addition all prime degree binomials are irreducible ([9, Thm. 5.1] and [10, Lemma 8.4.15]). The proof is by induction, using the fact that there exists for any prime  $p \geq 3$  numbers  $\beta_0, \dots, \beta_{p-2}$  such that  $\beta_i \in \mathbb{Z}[\varepsilon_{p-1}]$  with  $\varepsilon_{p-1}$  a primitive  $(p-1)$ -th root of unity and  $\varepsilon_p \in \mathbb{Q}(\varepsilon_{p-1}, \beta_0, \dots, \beta_{p-2})$  (see [21, (21.9)]). In Example 17 the construction of a chain of adjunctions of irreducible ideals would lead to the adjunction of the radicals expressing the primitive root of unity  $\zeta$ .

The complications caused by making all radicals irreducible can be avoided by using the same argument in the reducible case as in our proof of Theorem 14. In fact, this argument is similar to the one Weber uses to treat equation (4) in the case that  $f(X)$  is still reducible over  $K_{i-1}(\rho)$ , although more complicated. In the irreducible case we have  $\varepsilon_p \in K$  and it is immediate from (4) that varying the roots of  $X^p - a$ , that is, multiplying  $\alpha$  with powers of  $\varepsilon_p$ , gives  $p$  different roots. In the reducible case adjunction of  $\alpha$  also adjoins  $\varepsilon_q$  and typically  $p < d \leq q$ , as in Example 17. The fact that all roots of  $f(X)$  are obtained follows from Lemma 6. Of course this lemma can also be applied in the easier case  $q = p$ . Therefore both cases, where all roots are real, can be combined. They are characterized by the properties of the number  $\rho = \sqrt[p]{a\bar{a}}$ .

**Proposition 18.** *Let  $f(X) \in K(X)$  be an irreducible polynomial of odd prime degree  $p$  with real coefficients lying in a conjugation invariant field  $K$  and let  $K \subset L = K(\alpha)$ , where  $\alpha^q = a$ ,  $a \in K$ , be a normal and conjugation invariant extension such*

that  $f(X)$  is reducible over  $L$ . In case  $q = p$ , assume that  $K$  contains a primitive  $p$ -th root of unity  $\varepsilon$ . Let  $\rho = \sqrt[p]{a\bar{a}} = \alpha\bar{\alpha}$ . If  $\rho \in K$ , then all roots of  $f(X)$  are real, while if  $\rho \notin K$ , then  $f(X)$  has exactly one real root.

*Sketch of proof.* If  $\rho \in K$ , then we write the real root  $\beta_1$  as in equation (3), where the case  $q = p$  is allowed. The argument for Case II in the proof of Theorem 14 applies, as remarked above. If  $\rho \notin K$ , then  $X^q - a$  is irreducible, and  $q = p$ . As  $\rho \in K(\alpha) = K(\alpha, \bar{\alpha})$  we have  $K(\alpha) = K(\rho)$  by the Tower law (Lemma 4). Therefore we may replace  $\alpha$  by the real number  $\rho$ , and Weber's argument for the case  $\alpha \in \mathbb{R}$  applies. ■

As Kronecker [1] already remarked, the two cases, only one or all roots real, are for  $p \equiv 3 \pmod{4}$  distinguished by the sign of the discriminant of the polynomial (the square of the product of all differences between roots). This follows from the fact that the discriminant of a polynomial with real coefficients (without multiple factors) is positive if the number of pairs of complex conjugate roots is even and negative if this number is odd. It also follows that for  $p \equiv 1 \pmod{4}$  only irreducible equations with positive discriminant can be solvable.

**ACKNOWLEDGMENTS.** We express our gratitude to the referee for all remarks and comments, which contributed to a significant improvement of the paper.

**DISCLOSURE STATEMENT.** No potential conflict of interest was reported by the authors.

## ORCID

Jan Stevens  <http://orcid.org/0000-0001-9099-419X>

Juliusz Brzeziński  <http://orcid.org/0000-0002-5512-4313>

## REFERENCES

- [1] Kronecker L. Über die algebraisch auflösbaren Gleichungen (II. Abhandlung). Monatsberichte der Königlich Preussischen Akademie der Wissenschaften zu Berlin. 1856; p. 203–215. Also in: Werke Bd. IV. Herausgegeben auf Veranlassung der Preußischen Akademie der Wissenschaften von K. Hensel. Leipzig: B. G. Teubner; 1929. p. 27–37.
- [2] Weber H, Wellstein J. Elementare Algebra und Analysis bearbeitet von Heinrich Weber. Vol. 1, Encyclopädie der Elementar-Mathematik; ein Handbuch für Lehrer und Studierende. Leipzig: B.G. Teubner; 1903.
- [3] Nagell T. Lärbok i algebra. Stockholm: Almqvist & Wiksells; 1949.
- [4] Brzeziński J. Galois theory through exercises. Cham: Springer International Publishing; 2018.
- [5] Prasolov VV. Problems in algebra, arithmetic, and analysis (Russian). Moscow Center for Continuous Mathematical Education; 2007.
- [6] Morris SA, Jones A, Pearson KR. Abstract algebra and famous impossibilities. Squaring the circle, doubling the cube, trisecting an angle, and solving quintic equations. 2nd ed. Undergraduate Texts Math. Cham: Springer; 2022.
- [7] Dörrie H. 100 great problems of elementary mathematics. Their history and solution. Transl. from the German by David Antin. New York: Dover Publications; 1965.
- [8] Dörrie H. Triumph der Mathematik. Hundert berühmte Probleme aus zwei Jahrtausenden mathematischer Kultur. Breslau: Ferdinand Hirt; 1933.
- [9] Pan Y, Chen Y. On Kronecker's solvability theorem. arXiv:1912.07489v7; 2025.
- [10] Skopenkov A. Mathematics via problems. Part 1. Algebra. Vol. 25 of MSRI Mathematical Circles Library. Providence (RI): American Mathematical Society; 2021. Translated from the Russian original by Paul Zeitz and Sergei G. Shubin, With a foreword by Paul Zeitz.
- [11] Rosen MI. Niels Hendrik Abel and equations of the fifth degree. Amer Math Monthly. 1995;102:495–505.
- [12] Abel NH. Sur la resolution algébrique des équations. In: Sylow L, Lie S, editors. Œuvres complètes de Niels Henrik Abel, Vol. II. Christiania: Grøndahl & Søn; 1881. p. 217–243.

- [13] Petri B, Schappacher N. From Abel to Kronecker: Episodes from 19th Century algebra. In: Laudal OA, Piene R, editors. The legacy of Niels Henrik Abel. Berlin, Heidelberg: Springer; 2004. p. 227–266.
- [14] Kronecker L. Über die algebraisch auflösbaren Gleichungen (I. Abhandlung). Monatsberichte der Königlich Preussischen Akademie der Wissenschaften zu Berlin. 1853;p. 365–374. Also in: Werke Bd. IV. Herausgegeben auf Veranlassung der Preußischen Akademie der Wissenschaften von K. Hensel. Leipzig: B. G. Teubner; 1929. p. 3–11.
- [15] Weber H. Lehrbuch der Algebra. 2nd ed. Braunschweig: F. Vieweg und Sohn; 1898.
- [16] Wiman A. Über die metacyklischen Gleichungen von Primzahlgrad. Acta Math. 1903;27:163–175.
- [17] Edwards HM. Roots of solvable polynomials of prime degree. Expo Math. 2014;32:79–91.
- [18] Neumann PM. The mathematical writings of Évariste Galois. Heritage of European Mathematics. Zürich: European Mathematical Society (EMS); 2011.
- [19] Abel NH. Mathematische Bruchstücke aus Herrn N. H. Abels Briefen. J Reine Angew Math. 1830;5:336–343.
- [20] Cox DA. Why Eisenstein proved the Eisenstein criterion and Why Schönemann discovered it first. Amer Math Mon. 2011;118:3–21.
- [21] Stewart I. Galois theory. 4th ed. Boca Raton (FL): CRC Press; 2015.
- [22] Weber H, Wellstein J. Arithmetik, Algebra und Analysis, von Heinrich Weber. vol. 1 of Enzyklopädie der Elementar-Mathematik; ein Handbuch für Lehrer und Studierende. Vierte Auflage, neu bearbeitet von Paul Epstein ed. Leipzig-Berlin: B.G. Teubner; 1922.

**JULIUSZ BRZEZIŃSKI** Juliusz Brzeziński received his Ph.D. in mathematics from the University of Warsaw in 1967. He is now professor emeritus at the University of Gothenburg.

*Department of Mathematical Sciences, Chalmers University of Technology and University of Gothenburg. SE 412 96 Gothenburg, Sweden*  
[jub@chalmers.se](mailto:jub@chalmers.se)

**JAN STEVENS** Jan Stevens received his Ph.D. in mathematics from the University of Leiden in 1985. He is now associate professor emeritus at the University of Gothenburg.

*Department of Mathematical Sciences, Chalmers University of Technology and University of Gothenburg. SE 412 96 Gothenburg, Sweden*  
[stevens@chalmers.se](mailto:stevens@chalmers.se)