



Centrally pure C^* -algebras

Downloaded from: <https://research.chalmers.se>, 2026-06-26 22:14 UTC

Citation for the original published paper (version of record):

Perera, F., Thiel, H., Vilalta Vila, E. (2026). Centrally pure C^* -algebras. *Journal of Functional Analysis*, 291(8). <http://dx.doi.org/10.1016/j.jfa.2026.111578>

N.B. When citing this work, cite the original published paper.



Contents lists available at ScienceDirect

Journal of Functional Analysis

journal homepage: www.elsevier.com/locate/jfa



Regular Article

Centrally pure C^* -algebras [☆]

Francesc Perera ^{a,b,*}, Hannes Thiel ^c, Eduard Vilalta ^d



^a *Departament de Matemàtiques, Universitat Autònoma de Barcelona, 08193 Bellaterra, Barcelona, Spain*

^b *Centre de Recerca Matemàtica, Edifici Cc, Campus de Bellaterra, 08193 Cerdanyola del Vallès, Barcelona, Spain*

^c *Department of Mathematical Sciences, Chalmers University of Technology and the University of Gothenburg, SE-412 96 Gothenburg, Sweden*

^d *Department de Matemàtiques, Universitat Politècnica de Catalunya - BarcelonaTech (UPC), 08028 Barcelona, Barcelona, Spain*

ARTICLE INFO

Article history:

Received 24 December 2025

Accepted 11 May 2026

Available online 22 May 2026

Communicated by J. Gabe

MSC:

primary 46L05

secondary 19K14, 46L80, 46L85

Keywords:

C^* -algebras

Pureness

\mathcal{Z} -stability

Central sequence algebra

ABSTRACT

We show that a separable C^* -algebra A is \mathcal{Z} -stable if and only if its uncorrected central sequence algebra $A' \cap A_{\mathcal{U}}$ is pure, and only if Kirchberg's central sequence algebra $F(A)$ is pure. More generally, we show that a C^* -algebra A is separably \mathcal{Z} -stable if and only if the relative central sequence algebra $B' \cap A_{\mathcal{U}}$ is pure for every separable subalgebra $B \subseteq A_{\mathcal{U}}$.

© 2026 Published by Elsevier Inc.

[☆] FP and EV were partially supported by the Spanish State Research Agency through the grant PID2023-147110NB-I00 and by the Comissionat per Universitats i Recerca de la Generalitat de Catalunya (grant No. 2021-SGR-01015). FP was also supported by the Spanish State Research Agency through the Severo Ochoa and Maria de Maeztu Program for Centers and Units of Excellence in R&D (CEX2020-001084-M). HT and EV were partially supported by the Knut and Alice Wallenberg Foundation (KAW 2021.0140). HT was partially supported by the Swedish Research Council project grant 2024-04200.

* Corresponding author.

E-mail addresses: francesc.perera@uab.cat (F. Perera), hannes.thiel@chalmers.se (H. Thiel), eduard.vilalta@upc.edu (E. Vilalta).

URLs: <https://mat.uab.cat/web/perera> (F. Perera), <https://www.hannesthiel.org> (H. Thiel), <https://www.eduardvilalta.com> (E. Vilalta).

1. Introduction

Pureness and \mathcal{Z} -stability have emerged as decisive regularity properties in the structure and classification theory of C^* -algebras. Their central roles became apparent in the development of the Elliott program, which seeks to classify simple, nuclear C^* -algebras with an invariant encoding K -theoretic and tracial data. Toms [31] constructed examples of such algebras with the same invariant that are nevertheless nonisomorphic, since one is \mathcal{Z} -stable (i.e., it absorbs tensorially the Jiang-Su algebra \mathcal{Z}), while the other is not. These groundbreaking examples showed that additional regularity assumptions are indispensable in the classification program.

Subsequent work has established that \mathcal{Z} -stability is exactly the additional regularity assumption required. Indeed, building on decades of research, a series of landmark results [30,11,15] (see also the recent approach [8] and the survey [34]) established that simple, separable, nuclear, \mathcal{Z} -stable C^* -algebras satisfying the Universal Coefficient Theorem (UCT) are classified up to isomorphism by their Elliott invariants. Moreover, it is conceivable that the UCT condition is superfluous, or that it can be replaced by a classification framework based on KK-theory [28].

Pureness, introduced by Winter in his foundational work [36] on the structure of simple, nuclear C^* -algebras, is defined as the combination of strict comparison and almost divisibility in the Cuntz semigroup (see Section 3, and also [5,6] for a more systematic study of pure C^* -algebras). It is closely related to \mathcal{Z} -stability: Rørdam [24] showed that \mathcal{Z} -stability implies pureness in general, while the famous Toms–Winter conjecture [37] predicts that for simple, separable, nonelementary, nuclear C^* -algebras \mathcal{Z} -stability and strict comparison are equivalent, and this has been verified in many important cases [36,26,17,32,29]. Since pureness lies between strict comparison and \mathcal{Z} -stability, the conjecture in particular predicts that pureness and \mathcal{Z} -stability coincide in this setting. More broadly, the nonsimple Toms–Winter conjecture [5] proposes that a separable, nowhere scattered, nuclear C^* -algebra is \mathcal{Z} -stable if and only if it is pure. This has been verified in some cases, for example in [22, Theorems 7.10, 7.15].

Beyond the nuclear setting, pureness continues to play an important role. For instance, reduced free group C^* -algebras are pure [1] although not \mathcal{Z} -stable, showing that pureness can provide regularity where \mathcal{Z} -stability does not apply. Moreover, the solution to the rank problem in the stable rank one setting [29,2] shows that for simple, nonelementary C^* -algebras of stable rank one—and more generally for nowhere scattered C^* -algebras of stable rank one—pureness is equivalent to strict comparison.

In this paper we establish another close connection between \mathcal{Z} -stability and pureness. For a free ultrafilter \mathcal{U} on \mathbb{N} , denote by $A_{\mathcal{U}}$ the ultrapower of A . Our main result is:

Theorem A (see 3.6). *Let A be a separable C^* -algebra. Then the following are equivalent:*

- (1) A is \mathcal{Z} -stable.

- (2) The central sequence algebra $A' \cap A_{\mathcal{U}}$ is pure.
- (3) Kirchberg’s central sequence algebra $F(A) := (A' \cap A_{\mathcal{U}})/(A^{\perp} \cap A_{\mathcal{U}})$ is pure.

This shows that \mathcal{Z} -stability can be characterized as *central pureness*. It also suggests that pureness may be interpreted as a noncentral analog of \mathcal{Z} -stability.

We also obtain a nonseparable generalization. Recall that a C^* -algebra A is *separably \mathcal{Z} -stable* if any separable sub- C^* -algebra B_0 is contained in a separable, \mathcal{Z} -stable sub- C^* -algebra B ; see Definition 2.3 and the comments preceding it.

Theorem B (see 3.8). *Let A be a C^* -algebra. Then the following are equivalent:*

- (1) A is separably \mathcal{Z} -stable.
- (2) The relative central sequence algebra $B' \cap A_{\mathcal{U}}$ is pure for every separable subalgebra $B \subseteq A_{\mathcal{U}}$.
- (3) The quotient $(B' \cap A_{\mathcal{U}})/(B^{\perp} \cap A_{\mathcal{U}})$ is pure for every separable subalgebra $B \subseteq A_{\mathcal{U}}$.

Along the way (Section 2), we collect and streamline a number of well-known results on absorption of strongly self-absorbing C^* -algebras, some of which had not previously appeared in the literature in full generality. In the final section (Section 4), we investigate divisibility in central sequence algebras. Our results show that good central comparison follows from good central divisibility, so that \mathcal{Z} -stability is already detected by central divisibility properties. In fact, for a C^* -algebra to be \mathcal{Z} -stable it suffices for its central sequence algebra to satisfy n -almost divisibility for some fixed n (Proposition 4.4). It remains open whether even weaker divisibility properties, such as functional divisibility, are enough (Question 4.5).

2. Absorption of a strongly self-absorbing C^* -algebra

In this section, we present basic results that characterize when a separable C^* -algebra tensorially absorbs a strongly self-absorbing C^* -algebra \mathcal{D} ; see Theorem 2.2. These results are well-known, but certain aspects seem to not have appeared in the literature so far, or only under additional assumptions like unitality. We also give characterizations of separable \mathcal{D} -stability in the not necessarily separable case; see Theorem 2.7.

Throughout the paper, \mathcal{U} denotes a free ultrafilter on \mathbb{N} . Given a C^* -algebra A , we use $\prod_{\mathbb{N}} A$ to denote the C^* -algebra of bounded sequences in A , and the *ultrapower* of A , denoted by $A_{\mathcal{U}}$, is defined as the quotient $\prod_{\mathbb{N}} A/c_{\mathcal{U}}(A)$, where $c_{\mathcal{U}}(A)$ is the closed, two-sided ideal given by

$$c_{\mathcal{U}}(A) := \left\{ (a_n)_n \in \prod_{\mathbb{N}} A : \lim_{n \rightarrow \mathcal{U}} \|a_n\| = 0 \right\}.$$

We view A as a subalgebra of $A_{\mathcal{U}}$ via the natural embedding $A \rightarrow A_{\mathcal{U}}$ given by mapping an element $a \in A$ to the image of the constant sequence $(a, a, \dots) \in \prod_{\mathbb{N}} A$ in $A_{\mathcal{U}}$. The *central sequence algebra* of A is defined as the commutant of A inside $A_{\mathcal{U}}$, denoted by $A' \cap A_{\mathcal{U}}$. Note that the image of a sequence $(a_n)_n \in \prod_{\mathbb{N}} A$ in $A_{\mathcal{U}}$ belongs to $A' \cap A_{\mathcal{U}}$ if and only if

$$\lim_{n \rightarrow \mathcal{U}} \|a_n b - b a_n\| = 0$$

for every $b \in A$.

The *annihilator* of A inside $A_{\mathcal{U}}$ (denoted by $A^{\perp} \cap A_{\mathcal{U}}$) is the set of elements $b \in A_{\mathcal{U}}$ satisfying $bA = \{0\} = Ab$. This is a closed, two-sided ideal in $A' \cap A_{\mathcal{U}}$, and *Kirchberg’s central sequence algebra* is $F(A) := (A' \cap A_{\mathcal{U}})/(A^{\perp} \cap A_{\mathcal{U}})$; see [16, Definition 1.1]. To stress the difference between $F(A)$ and $A' \cap A_{\mathcal{U}}$, we sometimes refer to the latter as the *uncorrected central sequence algebra*.

If A is unital, then $A^{\perp} \cap A_{\mathcal{U}} = \{0\}$ and we get $F(A) = A' \cap A_{\mathcal{U}}$, but in general $F(A)$ is a proper quotient of $A' \cap A_{\mathcal{U}}$. If A is σ -unital, then $F(A)$ is unital.

A unital, separable C^* -algebra \mathcal{D} is *strongly self-absorbing* if $\mathcal{D} \neq \mathbb{C}$ and there is a $*$ -isomorphism $\mathcal{D} \rightarrow \mathcal{D} \otimes \mathcal{D}$ which is approximately unitarily equivalent to $\text{id}_{\mathcal{D}} \otimes 1_{\mathcal{D}}$; see [33, Definition 1.3]. It is known that strongly self-absorbing C^* -algebras are automatically simple and nuclear, and the only known examples are the Jiang-Su algebra \mathcal{Z} , the Cuntz algebras \mathcal{O}_2 and \mathcal{O}_{∞} , UHF-algebras of infinite type, and the tensor product of \mathcal{O}_{∞} with an UHF-algebra of infinite type. For \mathcal{D} strongly self-absorbing, a C^* -algebra A is said to be \mathcal{D} -stable provided $A \cong A \otimes \mathcal{D}$.

The next result is folklore. Its converse holds whenever A is separable; see Theorem 2.2.

Lemma 2.1. *Let \mathcal{D} be a strongly self-absorbing C^* -algebra, and let A be a (not necessarily separable) unital C^* -algebra that is \mathcal{D} -stable. Then there exists a unital embedding $\mathcal{D} \rightarrow A' \cap A_{\mathcal{U}}$.*

Proof. The sequence of unital embeddings

$$\varphi_n : \mathcal{D} \rightarrow A \otimes \mathcal{D} \otimes \mathcal{D} \otimes \dots, \quad d \mapsto 1_A \otimes 1_{\mathcal{D}} \otimes \dots \otimes 1_{\mathcal{D}} \otimes d \otimes 1_{\mathcal{D}} \otimes \dots$$

is approximately central in the sense that

$$\lim_{n \rightarrow \infty} \|\varphi_n(d)a - a\varphi_n(d)\| = 0,$$

for every $d \in \mathcal{D}$ and $a \in A \otimes \mathcal{D} \otimes \mathcal{D} \otimes \dots$. Using that

$$A \cong A \otimes \mathcal{D} \cong A \otimes \mathcal{D} \otimes \mathcal{D} \otimes \dots,$$

the sequence $(\varphi_n)_n$ induces a unital embedding $\mathcal{D} \rightarrow A' \cap A_{\mathcal{U}}$. \square

The next result is well-known to experts. The equivalence of (1) and (3) was shown in [33, Theorem 2.2] and is based on Rørdam’s approximate intertwining theorem [23, Theorem 7.2.2]; see also [18, Theorem 2.5]. To the best of our knowledge, the equivalence of (1) and (4) for non-unital C^* -algebras has only appeared in the literature for tensorial absorption of the Jiang-Su algebra in [19, Proposition 5.1], and we adapt the argument of Nawata to obtain the general result. A crucial ingredient is Winter’s result that strongly self-absorbing C^* -algebras are K_1 -injective [35]; see also Kirchberg’s result on tensorial absorption [16, Proposition 4.11]. For a C^* -algebra A , we denote by $M(A)$ its multiplier algebra.

Theorem 2.2. *Let A be a separable C^* -algebra, and let \mathcal{D} be a strongly self-absorbing C^* -algebra. Then the following are equivalent:*

- (1) *The C^* -algebra A is \mathcal{D} -stable.*
- (2) *There is a \mathcal{D} -stable C^* -algebra B with $A \subseteq B \subseteq M(A)$ and $1_{M(A)} \in B$.*
- (3) *There is a unital $*$ -homomorphism $\mathcal{D} \rightarrow A' \cap M(A)_{\mathcal{U}}$.*
- (4) *There is a unital $*$ -homomorphism $\mathcal{D} \rightarrow F(A)$.*

Proof. We show the implications ‘(1) \Rightarrow (2) \Rightarrow (3) \Rightarrow (4) \Rightarrow (1)’.

(1) \Rightarrow (2): Assume that A is \mathcal{D} -stable, and fix a $*$ -isomorphism $\varphi: A \rightarrow A \otimes \mathcal{D}$. This induces a $*$ -isomorphism $M(\varphi): M(A) \rightarrow M(A \otimes \mathcal{D})$. Let $\iota: A \rightarrow M(A)$ and $\iota': A \otimes \mathcal{D} \rightarrow M(A \otimes \mathcal{D})$ denote the natural inclusions. Then $\iota \otimes \text{id}_{\mathcal{D}}$ identifies $A \otimes \mathcal{D}$ with an essential, closed ideal in $M(A) \otimes \mathcal{D}$. By the universal property of multiplier algebras, we obtain a (unique) unital, injective $*$ -homomorphism $\psi: M(A) \otimes \mathcal{D} \rightarrow M(A \otimes \mathcal{D})$ such that $\iota' = \psi \circ (\iota \otimes \text{id}_{\mathcal{D}})$. The situation is shown in the following commutative diagram:

$$\begin{array}{ccc}
 A & \xrightarrow{\iota} & M(A) \\
 \cong \downarrow \varphi & & \cong \downarrow M(\varphi) \\
 A \otimes \mathcal{D} & \xrightarrow{\iota \otimes \text{id}_{\mathcal{D}}} M(A) \otimes \mathcal{D} \xrightarrow{\psi} & M(A \otimes \mathcal{D}).
 \end{array}$$

The desired algebra is $B := M(\varphi)^{-1}(\psi(M(A) \otimes \mathcal{D}))$.

(2) \Rightarrow (3): Let B be a \mathcal{D} -stable C^* -algebra with $A \subseteq B \subseteq M(A)$ and $1_{M(A)} \in B$. Applying Lemma 2.1, we get a unital $*$ -homomorphism $\mathcal{D} \rightarrow B' \cap B_{\mathcal{U}}$. Composing with the natural inclusion $A' \cap B_{\mathcal{U}} \subseteq A' \cap M(A)_{\mathcal{U}}$, we obtain the desired map

$$\mathcal{D} \rightarrow A' \cap B_{\mathcal{U}} \subseteq A' \cap M(A)_{\mathcal{U}}.$$

(3) \Rightarrow (4): Let $(e_n)_n$ be a contractive, approximate identity in A . This induces a (well-defined) contractive, linear map $M(A)_{\mathcal{U}} \rightarrow A_{\mathcal{U}}$ by mapping $[(x_n)_n] \in M(A)_{\mathcal{U}}$ to $[(e_n x_n)_n] \in A_{\mathcal{U}}$, which restricts to a map $\pi_0: A' \cap M(A)_{\mathcal{U}} \rightarrow A' \cap A_{\mathcal{U}}$.

While π_0 is not necessarily multiplicative, we have $\pi_0(xy) - \pi_0(x)\pi_0(y) \in A^\perp$ for all $x, y \in A' \cap M(A)_\mathcal{U}$. It follows that the composition of π_0 with the quotient map $A' \cap A_\mathcal{U} \rightarrow F(A)$ is a $*$ -homomorphism $\pi: A' \cap M(A)_\mathcal{U} \rightarrow F(A)$. We claim that π is surjective (and moreover independent of the choice of a contractive, approximate identity in A). To see that π is surjective, let $c \in F(A)$, represented by $b = [(b_n)_n] \in A' \cap A_\mathcal{U}$ for some bounded sequence $(b_n)_n$ in A . Viewing b as an element of $A' \cap M(A)_\mathcal{U}$, we claim that $b - \pi_0(b) \in A^\perp \cap A_\mathcal{U}$. Indeed, for $a \in A$ and $n \in \mathbb{N}$, we have

$$\|a(1 - e_n)b_n\| \leq \|a(1 - e_n)\| \|b_n\|$$

and

$$\begin{aligned} \|(1 - e_n)b_n a\| &\leq \|(1 - e_n)(b_n a - ab_n)\| + \|(1 - e_n)ab_n\| \\ &\leq \|b_n a - ab_n\| + \|(1 - e_n)a\| \|b_n\| \end{aligned}$$

and therefore $\lim_{n \rightarrow \mathcal{U}} \|a(1 - e_n)b_n\| = 0$ and $\lim_{n \rightarrow \mathcal{U}} \|(1 - e_n)b_n a\| = 0$ since $(e_n)_n$ is an approximate identity in A , and $(b_n)_n$ is bounded, and since $(b_n)_n$ approximately commutes with a . This shows that $a(b - \pi_0(b)) = (b - \pi_0(b))a = 0$, and since this holds for every $a \in A$, we get $b - \pi_0(b) \in A^\perp \cap A_\mathcal{U}$. Thus, b and $\pi_0(b)$ have the same image $F(A)$, and thus $c = \pi(b)$, proving surjectivity.

Now, if there is a unital $*$ -homomorphism $\mathcal{D} \rightarrow A' \cap M(A)_\mathcal{U}$, then by composing with π we obtain a unital $*$ -homomorphism $\mathcal{D} \rightarrow F(A)$.

(4) \Rightarrow (1): By assumption, there exists a unital $*$ -homomorphism $\alpha: \mathcal{D} \rightarrow F(A)$. To verify $A \cong A \otimes \mathcal{D}$, we consider the subalgebra $A \otimes 1$ of $M(A \otimes \mathcal{D})$, which induces embeddings $A \otimes 1 \subseteq (A \otimes 1)_\mathcal{U} \subseteq M(A \otimes \mathcal{D})_\mathcal{U}$. We will show that there exists a sequence of unitaries $(u_n)_n$ in $(A \otimes 1)' \cap M(A \otimes \mathcal{D})_\mathcal{U}$ such that

$$\lim_{n \rightarrow \infty} \text{dist}(u_n b u_n^*, (A \otimes 1)_\mathcal{U}) = 0$$

for all $b \in A \otimes \mathcal{D} \subseteq M(A \otimes \mathcal{D}) \subseteq M(A \otimes \mathcal{D})_\mathcal{U}$. It then follows from Rørdam’s approximate intertwining theorem [23, Proposition 7.2.1] that $A \cong A \otimes \mathcal{D}$.

To find the unitaries in $(A \otimes 1)' \cap M(A \otimes \mathcal{D})_\mathcal{U}$, we consider the quotient

$$C := \left((A \otimes 1)' \cap M(A \otimes \mathcal{D})_\mathcal{U} \right) / \left((A \otimes 1)^\perp \cap M(A \otimes \mathcal{D})_\mathcal{U} \right),$$

and we let $\pi: (A \otimes 1)' \cap M(A \otimes \mathcal{D})_\mathcal{U} \rightarrow C$ denote the quotient map. The subalgebra $1 \otimes \mathcal{D} \subseteq M(A \otimes \mathcal{D}) \subseteq M(A \otimes \mathcal{D})_\mathcal{U}$ lies in the commutant of $A \otimes 1$, which defines a unital $*$ -homomorphism

$$\beta_0: \mathcal{D} \xrightarrow{\cong} 1 \otimes \mathcal{D} \rightarrow (A \otimes 1)' \cap M(A \otimes \mathcal{D})_\mathcal{U}.$$

Composed with π , we obtain a unital $*$ -homomorphism $\beta := \pi \circ \beta_0: \mathcal{D} \rightarrow C$.

We can also interpret α as a map with codomain C . Specifically, denote by $\tilde{\pi}: A' \cap M(A)_{\mathcal{U}} \rightarrow F(A)$ the surjective $*$ -homomorphism constructed in the proof of the implication ‘(3) \Rightarrow (4)’. One can show that the kernel of $\tilde{\pi}$ is $A^{\perp} \cap M(A)_{\mathcal{U}}$, which establishes a natural $*$ -isomorphism

$$F(A) \cong (A' \cap M(A)_{\mathcal{U}})/(A^{\perp} \cap M(A)_{\mathcal{U}}).$$

This allows us to view $F(A)$ as a natural subalgebra of C and, via this identification, we have $\alpha: D \rightarrow C$. The situation is shown in the following diagram:

$$\begin{array}{ccc}
 (A \otimes 1)' \cap (A \otimes 1)_{\mathcal{U}} & \longrightarrow & ((A \otimes 1)' \cap (A \otimes 1)_{\mathcal{U}})/((A \otimes 1)^{\perp} \cap (A \otimes 1)_{\mathcal{U}}) \cong F(A) \\
 \downarrow & & \downarrow \\
 (A \otimes 1)' \cap M(A \otimes \mathcal{D})_{\mathcal{U}} & \xrightarrow{\pi} & ((A \otimes 1)' \cap M(A \otimes \mathcal{D})_{\mathcal{U}})/((A \otimes 1)^{\perp} \cap M(A \otimes \mathcal{D})_{\mathcal{U}}) =: C \\
 \beta_0 \uparrow & \nearrow \beta & \\
 \mathcal{D} \cong 1 \otimes \mathcal{D} & &
 \end{array}$$

One verifies that the maps $\alpha, \beta: \mathcal{D} \rightarrow C$ have commuting images, and since \mathcal{D} is nuclear we obtain a unital $*$ -homomorphism $\alpha \otimes \beta: \mathcal{D} \otimes \mathcal{D} \rightarrow C$. It follows from the definition of strong self-absorption that there is a sequence $(w_n)_n$ of unitaries in $\mathcal{D} \otimes \mathcal{D}$ such that $\lim_{n \rightarrow \infty} w_n(x \otimes y)w_n^* = y \otimes x$ for all $x, y \in \mathcal{D}$ (one says that \mathcal{D} has approximately inner flip). By [33, Proposition 1.13], the unitaries w_n may be chosen to be trivial in $K_1(\mathcal{D} \otimes \mathcal{D})$. By [35, Theorem 3.1], every strongly self-absorbing C^* -algebra is itself \mathcal{Z} -stable, and therefore it is K_1 -injective. This implies that the unitaries w_n lie in $\mathcal{U}_0(\mathcal{D} \otimes \mathcal{D})$, the connected component of the unit in the unitary group.

Setting $v_n := (\alpha \otimes \beta)(w_n)$, we thus have a sequence $(v_n)_n$ of unitaries in $\mathcal{U}_0(C)$ such that

$$\lim_{n \rightarrow \infty} v_n \alpha(x) \beta(y) v_n^* = \alpha(y) \beta(x)$$

for all $x, y \in \mathcal{D}$. Since unitaries in the connected component of the identity can be lifted along quotient maps, we can find unitaries u_n in $(A \otimes 1)' \cap M(A \otimes \mathcal{D})_{\mathcal{U}}$ with $\pi(u_n) = v_n$.

To show that $(u_n)_n$ have the desired properties, let $a \in A$ and $d \in \mathcal{D}$. Since $\alpha(\mathcal{D}) \subseteq \pi((A \otimes 1)' \cap (A \otimes 1)_{\mathcal{U}})$, we can pick $x \in (A \otimes 1)' \cap (A \otimes 1)_{\mathcal{U}}$ such that $\pi(x) = \alpha(d)$. We then have

$$\pi(u_n \beta_0(d) u_n^*) = v_n \beta(d) v_n^* \rightarrow \alpha(d) = \pi(x).$$

Since the kernel of π is the annihilator of $A \otimes 1$, we deduce that

$$u_n(a \otimes d) u_n^* = u_n(a \otimes 1) \beta_0(d) u_n^* = (a \otimes 1) u_n \beta_0(d) u_n^* \rightarrow (a \otimes 1) x \in (A \otimes 1)_{\mathcal{U}}.$$

This shows that $\lim_{n \rightarrow \infty} \text{dist}(u_n b u_n^*, (A \otimes 1)u) = 0$ for every elementary tensor $b = a \otimes d$ in $A \otimes \mathcal{D}$, and consequently for every $b \in A \otimes \mathcal{D}$. \square

While the definition of \mathcal{D} -stability makes sense for arbitrary C^* -algebras, it is less useful for nonseparable C^* -algebras since even natural candidates such as the ultrapower of \mathcal{D} fail to be \mathcal{D} -stable. Instead, the model-theoretically natural concept is to require that ‘sufficiently many’ separable sub- C^* -algebras are \mathcal{D} -stable. Variations of this concept have appeared under different names in the literature. We follow the terminology of Schafhauser [27, Definition 1.4]; see also [13, Proposition 1.6].

Definition 2.3. Let \mathcal{D} be a strongly self-absorbing C^* -algebra. A C^* -algebra A is said to be *separably \mathcal{D} -stable* if for every separable sub- C^* -algebra $B_0 \subseteq A$ there exists a separable sub- C^* -algebra $B \subseteq A$ that is \mathcal{D} -stable and such that $B_0 \subseteq B$.

We use $\text{Sep}(A)$ to denote the collection of separable sub- C^* -algebras of a given C^* -algebra A . Restricting the model-theoretic definition to our case of interest, a family $F \subseteq \text{Sep}(A)$ is a *club* if it is *cofinal* (that is, for every $B_0 \in \text{Sep}(A)$ there exists $B \in F$ such that $B_0 \subseteq B$) and σ -*complete* (that is, for every countable directed subset $F' \subseteq F$ we have that $\overline{\bigcup_{B \in F'} B} \in F$).

Let \mathcal{D} be a strongly self-absorbing C^* -algebra. Using that \mathcal{D} -stability passes to sequential inductive limits of separable C^* -algebras [33, Corollary 3.4], we see that a C^* -algebra A is separably \mathcal{D} -stable if and only if the separable, \mathcal{D} -stable sub- C^* -algebras of A form a club in $\text{Sep}(A)$.

Toms and Winter [33, Section 3] proved permanence properties of \mathcal{D} -stability among separable C^* -algebras, in particular passage to quotients, to hereditary sub- C^* -algebras, and to extensions. Farah and Szabó [13] showed that analogous permanence properties hold for separable \mathcal{D} -stability, which we record for later use.

Lemma 2.4. *Let \mathcal{D} be a strongly self-absorbing C^* -algebra. The following statements hold:*

- (1) *If $0 \rightarrow I \rightarrow E \rightarrow B \rightarrow 0$ is an extension of C^* -algebras, then E is separably \mathcal{D} -stable if and only if I and B are separably \mathcal{D} -stable.*
- (2) *If A is separably \mathcal{D} -stable, and $B \subseteq A$ is a hereditary sub- C^* -algebra, then B is separably \mathcal{D} -stable.*

Proof. The first statement was proved in [13, Lemma 1.14], and the second statement was shown in [13, Proposition 1.12]. \square

The following concept was introduced in [13, Definition 1.11].

Definition 2.5. Let \mathcal{D} be a unital C^* -algebra. We say that a unital C^* -algebra A is *\mathcal{D} -saturated* if for every separable sub- C^* -algebra $B \subseteq A$ there is a unital $*$ -embedding $\mathcal{D} \rightarrow B' \cap A$.

Remark 2.6. Let \mathcal{D} be a strongly self-absorbing C^* -algebra. Then every unital \mathcal{D} -saturated C^* -algebra B is separably \mathcal{D} -stable. Indeed, let $C_0 \subseteq B$ be a separable sub- C^* -algebra. By \mathcal{D} -saturation, there exists a unital $*$ -homomorphism $\varphi: \mathcal{D} \rightarrow C'_0 \cap B$. Since \mathcal{D} is separable, $\mathcal{D}_0 := \varphi(\mathcal{D})$ is a separable sub- C^* -algebra of B commuting with C_0 , and $C_1 := C^*(C_0 \cup \mathcal{D}_0) \subseteq B$ is a separable sub- C^* -algebra containing C_0 . The map $\mathcal{D} \otimes C_0 \rightarrow C_1$ given by $d \otimes c \mapsto \varphi(d)c$ is a $*$ -isomorphism, showing that C_1 is \mathcal{D} -stable. Thus B is separably \mathcal{D} -stable.

Theorem 2.7. Let \mathcal{D} be a strongly self-absorbing C^* -algebra, and let A be a C^* -algebra. Then the following are equivalent:

- (1) The C^* -algebra A is separably \mathcal{D} -stable.
- (2) The ultrapower $A_{\mathcal{U}}$ is separably \mathcal{D} -stable.
- (3) The relative commutant $B' \cap A_{\mathcal{U}}$ is separably \mathcal{D} -stable for every separable sub- C^* -algebra $B \subseteq A_{\mathcal{U}}$.
- (4) The quotient $(B' \cap A_{\mathcal{U}})/(B^\perp \cap A_{\mathcal{U}})$ is \mathcal{D} -saturated for every separable sub- C^* -algebra $B \subseteq A_{\mathcal{U}}$.
- (5) There exists a unital $*$ -homomorphism $\mathcal{D} \rightarrow (B' \cap A_{\mathcal{U}})/(B^\perp \cap A_{\mathcal{U}})$ for every separable sub- C^* -algebra $B \subseteq A_{\mathcal{U}}$.

If A is σ -unital, then the above statements are also equivalent to

- (6) The multiplier algebra $M(A)$ is separably \mathcal{D} -stable.
- (7) The C^* -algebra $M(A)_{\mathcal{U}}$ is \mathcal{D} -saturated.

Proof. We show the implications ‘(1) \Leftrightarrow (2)’ and ‘(1) \Leftrightarrow (5)’ and ‘(3) \Rightarrow (4) \Rightarrow (5) \Rightarrow (3)’ and ‘(1) \Leftrightarrow (6)’ and ‘(6) \Rightarrow (7) \Rightarrow (6)’.

(1) \Leftrightarrow (2): A property \mathcal{P} of C^* -algebras is said to pass to ultrapowers if, for every C^* -algebra B and ultrafilter \mathcal{V} , whenever B has \mathcal{P} , so does $B_{\mathcal{V}}$. It is said to pass to ultraroots if, for every C^* -algebra B and ultrafilter \mathcal{V} , whenever $B_{\mathcal{V}}$ has \mathcal{P} , so does B ; see, for example, [12, Section 2.4].

It follows from [13, Corollary 1.10] that separable \mathcal{D} -stability is axiomatizable; see also [12, Theorem 2.5.2(21)]. Since axiomatizable properties pass to ultrapowers and ultraroots ([12, Theorem 2.4.1]), we see that (1) and (2) are equivalent.

(1) \Leftrightarrow (5): We use that by [13, Corollary 1.10(2) and Definition 1.8], the C^* -algebra A is separably \mathcal{D} -stable if and only if A satisfies condition (1) in [13, Proposition 1.7], namely, for any sub- C^* -algebra B_0 of A , there is a unital $*$ -homomorphism $\mathcal{D} \rightarrow (B'_0 \cap A_\infty)/(B^\perp_0 \cap A_\infty)$.

For the forward implication, assuming that A is separably \mathcal{D} -stable, let $B \subseteq A_{\mathcal{U}}$ be a separable sub- C^* -algebra. We consider the natural quotient map $\pi: A_\infty \rightarrow A_{\mathcal{U}}$ from the sequence algebra $A_\infty := \prod_{\mathbb{N}} A/c_0(A)$ where $c_0(A)$ is the closed ideal of sequences $(a_n)_n$ in A with $\lim_{n \rightarrow \infty} \|a_n\| = 0$. Choose a separable sub- C^* -algebra $B_0 \subseteq A_\infty$ with

$\pi(B_0) = B$. By [13, Proposition 1.7 (1)], there exists a unital $*$ -homomorphism $\mathcal{D} \rightarrow (B'_0 \cap A_\infty)/(B_0^\perp \cap A_\infty)$. The map π restricts to a map $B'_0 \cap A_\infty \rightarrow B' \cap A_{\mathcal{U}}$, which induces a unital $*$ -homomorphism from $(B'_0 \cap A_\infty)/(B_0^\perp \cap A_\infty)$ to $(B' \cap A_{\mathcal{U}})/(B^\perp \cap A_{\mathcal{U}})$. We obtain the desired map as the composition

$$\mathcal{D} \rightarrow (B'_0 \cap A_\infty)/(B_0^\perp \cap A_\infty) \rightarrow (B' \cap A_{\mathcal{U}})/(B^\perp \cap A_{\mathcal{U}}).$$

For the backward implication, we notice that statement (5) immediately implies condition (3) in [13, Proposition 1.7], which then implies that A is separably \mathcal{D} -stable.

(3) \Rightarrow (4): To verify (4), let $B \subseteq A_{\mathcal{U}}$ be a separable sub- C^* -algebra. Set $Q := (B' \cap A_{\mathcal{U}})/(B^\perp \cap A_{\mathcal{U}})$ and let $\pi: B' \cap A_{\mathcal{U}} \rightarrow Q$ denote the quotient map. To show that Q is \mathcal{D} -saturated, let $C \subseteq Q$ be separable. We need find a unital $*$ -homomorphism $\mathcal{D} \rightarrow C' \cap Q$.

Choose a separable sub- C^* -algebra $C_0 \subseteq B' \cap A_{\mathcal{U}}$ with $\pi(C_0) = C$. Since the C^* -algebra E generated by C_0 and B is separable, it follows from the assumption that $E' \cap A_{\mathcal{U}}$ is separably \mathcal{D} -stable, and hence so is the quotient $\pi(E' \cap A_{\mathcal{U}})$ by Lemma 2.4. Since $\pi(E' \cap A_{\mathcal{U}})$ is unital and separably \mathcal{D} -stable, it admits a unital $*$ -homomorphism from \mathcal{D} , and we obtain the desired map as the composition

$$\mathcal{D} \rightarrow \pi(E' \cap A_{\mathcal{U}}) \subseteq \pi(C_0)' \cap \pi(B' \cap A_{\mathcal{U}}) = C' \cap Q.$$

(4) \Rightarrow (5): This follows using that every unital, \mathcal{D} -saturated C^* -algebra admits a unital $*$ -homomorphism from \mathcal{D} .

(5) \Rightarrow (3): Assume that (5) holds. With a similar argument as in the implication '(3) \Rightarrow (4)', we see that $(B' \cap A_{\mathcal{U}})/(B^\perp \cap A_{\mathcal{U}})$ is \mathcal{D} -saturated (and consequently separably \mathcal{D} -stable, by Remark 2.6) for every separable sub- C^* -algebra $B \subseteq A_{\mathcal{U}}$.

To verify (3), let $B \subseteq A_{\mathcal{U}}$ be a separable sub- C^* -algebra, and consider the extension

$$0 \rightarrow B^\perp \cap A_{\mathcal{U}} \rightarrow B' \cap A_{\mathcal{U}} \rightarrow (B' \cap A_{\mathcal{U}})/(B^\perp \cap A_{\mathcal{U}}) \rightarrow 0.$$

We have already shown that (5) implies (2), which shows that $A_{\mathcal{U}}$ is separably \mathcal{D} -stable. Since $B^\perp \cap A_{\mathcal{U}}$ is a hereditary sub- C^* -algebra of $A_{\mathcal{U}}$, it follows from Lemma 2.4(2) that $B^\perp \cap A_{\mathcal{U}}$ is separably \mathcal{D} -stable. Further, as noted above, the quotient $(B' \cap A_{\mathcal{U}})/(B^\perp \cap A_{\mathcal{U}})$ is also separably \mathcal{D} -stable. Now it follows from Lemma 2.4(1) that $B' \cap A_{\mathcal{U}}$ is separably \mathcal{D} -stable.

(1) \Leftrightarrow (6): Assuming that A is σ -unital, the equivalence of (1) and (6) is shown in [13, Theorem B].

(6) \Rightarrow (7): Assume that $M(A)$ is separably \mathcal{D} -stable. To show that $M(A)_{\mathcal{U}}$ is \mathcal{D} -saturated, let $B \subseteq M(A)_{\mathcal{U}}$ be separable. Applying the equivalence of (1) and (3) for $M(A)$, it follows that $B' \cap M(A)_{\mathcal{U}}$ is separably \mathcal{D} -stable, and in particular admits a unital $*$ -homomorphism $\mathcal{D} \rightarrow B' \cap M(A)_{\mathcal{U}}$. Namely, if B_0 is a unital separable sub- C^* -algebra of $B' \cap M(A)_{\mathcal{U}}$, then there is a separable \mathcal{D} -stable sub- C^* -algebra \tilde{B}_0 with $B_0 \subseteq \tilde{B}_0$.

(7) \Rightarrow (6): Assume that $M(A)_{\mathcal{U}}$ is \mathcal{D} -saturated. By Remark 2.6, we have that $M(A)_{\mathcal{U}}$ is separably \mathcal{D} -stable. Applying the equivalence of (1) and (2) for $M(A)$, it follows that $M(A)$ is separably \mathcal{D} -stable. \square

Applying Theorem 2.7 for a separable C^* -algebra, we obtain:

Corollary 2.8. *Let A be a separable C^* -algebra, and let \mathcal{D} be a strongly self-absorbing C^* -algebra. Then the following are equivalent:*

- (1) *The C^* -algebra A is \mathcal{D} -stable.*
- (2) *The algebra $M(A)_{\mathcal{U}}$ is \mathcal{D} -saturated.*
- (3) *The algebra $A' \cap M(A)_{\mathcal{U}}$ is \mathcal{D} -saturated.*
- (4) *The (uncorrected) central sequence algebra $A' \cap A_{\mathcal{U}}$ is separably \mathcal{D} -stable.*
- (5) *Kirchberg’s central sequence algebra $F(A)$ is \mathcal{D} -saturated.*

Proof. We show the implications ‘(1) \Leftrightarrow (2) \Rightarrow (3) \Rightarrow (1)’ and ‘(1) \Leftrightarrow (4)’ and ‘(1) \Leftrightarrow (5)’.

(1) \Leftrightarrow (2): This follows from the equivalence of (1) and (7) in Theorem 2.7.

(2) \Rightarrow (3): In general, if B is a \mathcal{D} -saturated C^* -algebra and $C \subseteq B$ is a separable sub- C^* -algebra, then $C' \cap B$ is \mathcal{D} -saturated. For, if $E \subseteq C' \cap B$ is a separable sub- C^* -algebra, then $C^*(E \cup C)$ is also separable and by assumption there is a unital $*$ -embedding $\mathcal{D} \rightarrow C^*(E \cup C)' \cap B \subseteq E' \cap C' \cap B$. Now, since A is separable, we see that (2) implies (3).

(3) \Rightarrow (1): Assuming that $A' \cap M(A)_{\mathcal{U}}$ is \mathcal{D} -saturated, it follows that there is a unital $*$ -homomorphism $\mathcal{D} \rightarrow A' \cap M(A)_{\mathcal{U}}$. Applying the implication ‘(3) \Rightarrow (1)’ in Theorem 2.2, we see that A is \mathcal{D} -stable.

(1) \Leftrightarrow (4): Since A is separable, the forward implication follows from the implication ‘(1) \Rightarrow (3)’ in Theorem 2.7. For the backward implication, assume that $A' \cap A_{\mathcal{U}}$ is separably \mathcal{D} -stable. Since $F(A)$ is a quotient of $A' \cap A_{\mathcal{U}}$, and using Lemma 2.4(1), we deduce that $F(A)$ is separably \mathcal{D} -stable and therefore admits a unital $*$ -homomorphism $\mathcal{D} \rightarrow F(A)$. Applying the implication ‘(4) \Rightarrow (1)’ in Theorem 2.2, we see that A is \mathcal{D} -stable.

(1) \Leftrightarrow (5): Since A is separable, the forward implication follows from the implication ‘(1) \Rightarrow (4)’ in Theorem 2.7. Conversely, if $F(A)$ is \mathcal{D} -saturated, then it admits a unital $*$ -homomorphism $\mathcal{D} \rightarrow F(A)$, which by the implication ‘(4) \Rightarrow (1)’ in Theorem 2.2 shows that A is \mathcal{D} -stable. \square

3. Purenness of central sequence algebras

In this section, we prove the main result of this paper (Theorem 3.6): a separable C^* -algebra A is \mathcal{Z} -stable if and only if its central sequence algebra $A' \cap A_{\mathcal{U}}$ is pure. In Theorem 3.8, we obtain a similar characterization of separable \mathcal{Z} -stability (for not necessarily separable C^* -algebras) in terms of pureness of the relative central sequence algebras $B' \cap A_{\mathcal{U}}$ for separable sub- C^* -algebras $B \subseteq A_{\mathcal{U}}$.

Pureness is defined as the combination of strict comparison and almost divisibility, which can be thought of as good comparison and divisibility properties in the Cuntz semigroup; see below for the precise definitions. From this point of view, Theorem 3.6 shows that a separable C^* -algebra is \mathcal{Z} -stable if and only if its central sequence algebra enjoys good comparison and good divisibility properties. We refer the reader to [14] for a survey on various aspects on the Cuntz semigroup $\text{Cu}(A)$ of a C^* -algebra A . We use \ll to denote the sequential way-below relation on $\text{Cu}(A)$, which is defined by setting $x \ll y$ if for every increasing sequence $(y_n)_n$ in $\text{Cu}(A)$ with $y \leq \sup_n y_n$, there exists n such that $x \leq y_n$.

The relation $<_s$ on an ordered monoid is defined by setting $x <_s y$ if there exists $n \in \mathbb{N}$ such that $(n + 1)x \leq ny$. Then, by definition, A has *strict comparison* if $x <_s y$ in $\text{Cu}(A)$ implies $x \leq y$. This is also referred to by saying that $\text{Cu}(A)$ is *almost unperforated*. One says that A is *n-almost divisible* if, for any pair $x'x \in \text{Cu}(A)$ with $x' \ll x$, $k \in \mathbb{N}$, there is $y \in \text{Cu}(A)$ with $ky \leq x$ and $x' \leq (n + 1)(k + 1)y$. The C^* -algebra A is *almost divisible* if it is 0-almost divisible.

We now record the following observation.

Proposition 3.1. *Every separably \mathcal{Z} -stable C^* -algebra is pure.*

Proof. Rørdam showed in [24] that every \mathcal{Z} -stable C^* -algebra has almost unperforated Cuntz semigroup. It follows from [7, Theorem 5.35] that it is also almost divisible, and hence pure; see [5, Proposition 5.2]. (Let us remark that said properties are established for the uncompleted Cuntz semigroup $W(A)$, and they also pass to $\text{Cu}(A)$ since the latter is the completion of $W(A)$; see [4, Theorem 3.2.8].)

Now let A be a separably \mathcal{Z} -stable C^* -algebra. Then the family of separable, \mathcal{Z} -stable sub- C^* -algebras of A forms an inductive system (indexed over itself) whose inductive limit is (isomorphic to) A . By [6, Theorem 3.8], pureness of C^* -algebras is separably determined (a term also introduced in [6, Definition 2.2]), so in particular it passes to inductive limits, which shows that A is pure. \square

For $n \geq 1$, the *dimension-drop algebra* $Z_{n,n+1}$ is defined as the sub- C^* -algebra of $C([0, 1], M_n \otimes M_{n+1})$ consisting of the continuous functions $f: [0, 1] \rightarrow M_n \otimes M_{n+1}$ that satisfy

$$f(0) \in M_n \otimes 1_{n+1}, \quad \text{and} \quad f(1) \in 1_n \otimes M_{n+1}.$$

In [25, Proposition 5.1], Rørdam and Winter showed that a unital C^* -algebra A of stable rank one admits a unital $*$ -homomorphism $Z_{n,n+1} \rightarrow A$ if and only if the unit can be ‘almost’ divided by n in the Cuntz semigroup $\text{Cu}(A)$ in the sense that there exists a Cuntz class $x \in \text{Cu}(A)$ such that $nx \leq [1] \leq (n + 1)x$. (Again, this was established in $W(A)$ and the result also holds in $\text{Cu}(A)$ by density.)

In [10, Theorem 3.6], Dadarlat and Toms proved the same result for separable, unital C^* -algebras with nonempty tracial state space and with strict comparison of positive

elements by tracial states. The next result generalizes the result of Dadarlat and Toms by removing the assumptions of separability and of nonempty tracial state space, and by weakening strict comparison of positive elements by tracial states to strict comparison of positive elements by quasitraces. A proof can essentially be found in [20, Proposition 4.9], noting that the argument can be adapted to only require almost divisibility of the unit. We reproduce the (adapted) argument here for the convenience of the reader.

Proposition 3.2. *Let A be a unital C^* -algebra with strict comparison (of positive elements by quasitraces). Then, the following are equivalent:*

- (1) *The element $[1]$ is almost divisible in $\text{Cu}(A)$.*
- (2) *For each $n \geq 1$, there exists a unital $*$ -homomorphism $Z_{n,n+1} \rightarrow A$.*

Proof. Assuming (2), let us verify (1). To show that $[1]$ is almost divisible, let $n \geq 1$. We need to find $y \in \text{Cu}(A)$ such that $ny \leq [1] \leq (n + 1)y$. By assumption, there is a unital $*$ -homomorphism $\varphi: Z_{n,n+1} \rightarrow A$. By [24, Lemma 4.2], there exists a positive element $e \in Z_{n,n+1}$ such that $n[e] \leq [1] \leq (n + 1)[e]$ in $\text{Cu}(Z_{n,n+1})$. Now $y := [\varphi(e)]$ has the desired properties.

Assuming (1), let us prove (2). For $a, b \in A_+$, write $a \simeq b$ whenever $a = xx^*$ and $b = x^*x$ for some $x \in A$. We will apply [25, Proposition 5.1], which states that there exists a unital $*$ -homomorphism $Z_{n,n+1} \rightarrow A$ whenever one can find pairwise orthogonal elements $a_1, \dots, a_n \in A_+$ such that $a_i \simeq a_j$ for each pair i, j and $1 - \sum_i a_i \precsim (a_1 - \varepsilon)_+$ for some $\varepsilon > 0$ (where \precsim denotes Cuntz subequivalence).

To verify (2), let $n \geq 1$ and set $m = 2n$. By (1), there exists $c \in (A \otimes \mathcal{K})_+$ such that $m[c] \leq [1] \leq (m+1)[c]$. Since $[1] \ll [1]$, we can pick $\varepsilon > 0$ such that $[1] \leq (m+1)[(c - \varepsilon)_+]$. We thus have

$$m[c] \leq [1] \leq (m + 1)[(c - \varepsilon)_+].$$

Applying [21, Lemma 2.4] (or, rather, its proof) to the first inequality, one obtains pairwise orthogonal elements $c_1, \dots, c_m \in A_+$ such that

$$(c - \frac{\varepsilon}{2})_+ \simeq c_1 \simeq \dots \simeq c_m.$$

Let $f: \mathbb{R} \rightarrow [0, 1]$ be a continuous function that takes the value 0 on $(-\infty, 0]$ and that takes the value 1 on $[\frac{\varepsilon}{2}, \infty)$. Using continuous functional calculus, set $d'_j := (c_j - \frac{\varepsilon}{2})_+$ and $\underline{d}_j := f(c_j)$, and note that then $d'_j = d'_j \underline{d}_j$ for $j = 1, \dots, m$. Since d_j belongs to $c_j A c_j$, and since c_1, \dots, c_m are pairwise orthogonal, it follows that d_1, \dots, d_m are pairwise orthogonal as well.

For each $j = 1, \dots, m$, we claim that $d_j \simeq f((c - \frac{\varepsilon}{2})_+)$ and $d'_j \simeq (c - \varepsilon)_+$. Since $(c - \frac{\varepsilon}{2})_+ \simeq c_j$, we can choose $x_j \in A$ such that $(c - \frac{\varepsilon}{2})_+ = x_j x_j^*$ and $c_j = x_j^* x_j$. Let $x_j = v_j |x_j|$ be the polar decomposition of x_j in A^{**} , and set

$$y_j := f\left((c - \frac{\varepsilon}{2})_+\right)^{\frac{1}{2}} v_j, \quad \text{and} \quad z_j := (c - \varepsilon)_+^{\frac{1}{2}} v_j.$$

Since $f\left((c - \frac{\varepsilon}{2})_+\right)^{\frac{1}{2}}$ and $(c - \varepsilon)_+^{\frac{1}{2}}$ belong to $\overline{A|x_j^*|}$ and $|x_j^*|v_j = x_j \in A$, we deduce that y_j and z_j belong to A . We have

$$y_j y_j^* = f\left((c - \frac{\varepsilon}{2})_+\right), \quad \text{and} \quad z_j z_j^* = (c - \varepsilon)_+.$$

Note that $v_j^*(c - \frac{\varepsilon}{2})_+ v_j = v_j^* x_j x_j^* v_j = |x_j| |x_j| = c_j$. Since the map $b \mapsto v_j^* b v_j$ defines a $*$ -isomorphism from $(c - \frac{\varepsilon}{2})_+ A (c - \frac{\varepsilon}{2})_+$ onto $c_j A c_j$, and since applying functional calculus is compatible with $*$ -homomorphisms, we deduce that

$$y_j^* y_j = v_j^* f\left((c - \frac{\varepsilon}{2})_+\right) v_j = f\left(v_j^*(c - \frac{\varepsilon}{2})_+ v_j\right) = f(c_j) = d_j$$

and

$$\begin{aligned} z_j^* z_j &= v_j^*(c - \varepsilon)_+ v_j \\ &= v_j^*\left((c - \frac{\varepsilon}{2})_+ - \frac{\varepsilon}{2}\right)_+ v_j \\ &= \left(v_j^*(c - \frac{\varepsilon}{2})_+ v_j - \frac{\varepsilon}{2}\right)_+ \\ &= (c_j - \frac{\varepsilon}{2})_+ = d'_j. \end{aligned}$$

It follows that

$$d_j = y_j^* y_j \simeq y_j y_j^* = f\left((c - \frac{\varepsilon}{2})_+\right), \quad \text{and} \quad d'_j = z_j^* z_j \simeq z_j z_j^* = (c - \varepsilon)_+,$$

as claimed.

Given a quasitrace τ on A , we define $d_\tau: (A \otimes \mathcal{K})_+ \rightarrow [0, \infty]$ by $d_\tau(a) := \lim_n \tau(a^{1/n})$. Using at the second step that $1 - \sum_{j=1}^m d_j$ is orthogonal to each d'_1, \dots, d'_m , and using at the last step that $(c - \varepsilon)_+$ is in particular Cuntz equivalent to d'_1 , one gets

$$\begin{aligned} d_\tau \left(1 - \sum_{j=1}^m d_j \right) + m d_\tau(d'_1) &= d_\tau \left(1 - \sum_{j=1}^m d_j \right) + \sum_{j=1}^m d_\tau(d'_j) \leq d_\tau(1) \\ &\leq (m + 1) d_\tau((c - \varepsilon)_+) = (m + 1) d_\tau(d'_1). \end{aligned}$$

Since $d_\tau(d'_1) < \infty$, we can cancel $m d_\tau(d'_1)$ and obtain

$$d_\tau \left(1 - \sum_{j=1}^m d_j \right) \leq d_\tau(d'_1) = \frac{1}{2} d_\tau(d'_1) + \frac{1}{2} d_\tau(d'_2) = \frac{1}{2} d_\tau(d'_1 + d'_2).$$

Pair the elements d_j by setting $a_1 = d_1 + d_2, \dots, a_n = d_{2n-1} + d_{2n}$. We have

$$d'_1 = (c_1 - \frac{\varepsilon}{2})_+ \preceq (f(c_1) - \frac{\varepsilon}{2})_+ = (d_1 - \frac{\varepsilon}{2})_+$$

and similarly $d'_2 \lesssim (d_2 - \frac{\epsilon}{2})_+$. Using that d_1 and d_2 are orthogonal, we deduce that

$$d'_1 + d'_2 \lesssim (d_1 - \frac{\epsilon}{2})_+ + (d_2 - \frac{\epsilon}{2})_+ = (a_1 - \frac{\epsilon}{2})_+.$$

We thus have

$$d_\tau \left(1 - \sum_{j=1}^n a_j \right) = d_\tau \left(1 - \sum_{j=1}^m d_j \right) \leq \frac{1}{2} d_\tau (d'_1 + d'_2) \leq \frac{1}{2} d_\tau ((a_1 - \frac{\epsilon}{2})_+).$$

Using strict comparison, one deduces from the previous functional inequality that $1 - \sum_{j=1}^n a_j \lesssim (a_1 - \frac{\epsilon}{2})_+$. This finishes the proof. \square

Corollary 3.3 ([20, Proposition 4.9]). *Every unital pure C^* -algebra A admits unital $*$ -homomorphisms $Z_{n,n+1} \rightarrow A$ for every $n \in \mathbb{N}$.*

We expect a positive answer to the following question. Under the additional assumption of stable rank one, this is shown in [4, Proposition 7.6.5].

Question 3.4. Does every unital pure C^* -algebra admit a unital $*$ -homomorphism from \mathcal{Z} ?

We will use the following consequence of [16, Proposition 1.12], which was stated under the additional assumption that A is unital in [18, Proposition 2.2, Corollary 2.3].

Proposition 3.5 (Kirchberg). *Let A be a separable C^* -algebra, and let $B, D \subseteq F(A)$ be separable sub- C^* -algebras with $1_{F(A)} \in D$. Then there exists a unital, injective $*$ -homomorphism $D \rightarrow B' \cap F(A)$.*

Consequently, there exists a unital $*$ -homomorphism $\bigotimes_{\max}^\infty D \rightarrow F(A)$.

Theorem 3.6. *Let A be a separable C^* -algebra. Then the following are equivalent:*

- (1) *The C^* -algebra A is \mathcal{Z} -stable.*
- (2) *The (uncorrected) central sequence algebra $A' \cap A_{\mathcal{U}}$ is pure.*
- (3) *Kirchberg’s central sequence algebra $F(A)$ is pure.*

Proof. Assuming (1), it follows from the implication ‘(1) \Rightarrow (4)’ in Corollary 2.8 that $A' \cap A_{\mathcal{U}}$ is separably \mathcal{Z} -stable, and therefore pure by Proposition 3.1, thus verifying (2). The implication ‘(2) \Rightarrow (3)’ follows since $F(A)$ is a quotient of $A' \cap A_{\mathcal{U}}$, and since pureness passes to quotients by [6, Theorem 4.11].

Assuming (3), let us verify (1). Since $F(A)$ is pure, there exists a unital $*$ -homomorphism $Z_{2,3} \rightarrow F(A)$ by Corollary 3.3. This is well-known to imply that A is \mathcal{Z} -stable; see, for example, [18, Theorem 2.5] in case A is unital. For completeness, we include

the argument: Using [9, Theorem 1.1] and Proposition 3.5, we obtain unital $*$ -homomorphisms

$$\mathcal{Z} \rightarrow \bigotimes_{\min}^{\infty} Z_{2,3} = \bigotimes_{\max}^{\infty} Z_{2,3} \rightarrow F(A).$$

Applying the implication ‘(4) \Rightarrow (1)’ in Theorem 2.2, it follows that A is \mathcal{Z} -stable. \square

Remark 3.7. Analogous to Theorem 3.6, one can show that \mathcal{O}_{∞} -stability of a separable C^* -algebra A is encoded in the Cuntz semigroups of $A' \cap A_{\mathcal{U}}$ and $F(A)$. Specifically, the following are equivalent:

- (1) The C^* -algebra A is \mathcal{O}_{∞} -stable.
- (2) The (uncorrected) central sequence algebra $A' \cap A_{\mathcal{U}}$ is purely infinite.
- (3) Kirchberg’s central sequence algebra $F(A)$ is purely infinite.

This relies on the fact that a separably \mathcal{O}_{∞} -absorbing C^* -algebra is purely infinite (analogous to Proposition 3.1), and that a unital, purely infinite C^* -algebra admits a unital $*$ -homomorphism from \mathcal{O}_{∞} .

Theorem 3.8. *Let A be a C^* -algebra. Then the following are equivalent:*

- (1) *The C^* -algebra A is separably \mathcal{Z} -stable.*
- (2) *The relative commutant $B' \cap A_{\mathcal{U}}$ is pure for every separable $B \subseteq A_{\mathcal{U}}$.*
- (3) *The quotient $(B' \cap A_{\mathcal{U}})/(B^{\perp} \cap A_{\mathcal{U}})$ is pure for every separable $B \subseteq A_{\mathcal{U}}$.*

Proof. We show the implications ‘(1) \Rightarrow (2) \Rightarrow (3) \Rightarrow (1)’.

(1) \Rightarrow (2): Assume (1). To verify (2), let $B \subseteq A_{\mathcal{U}}$ be a separable sub- C^* -algebra. It follows from the implication ‘(1) \Rightarrow (3)’ in Theorem 2.7 that $B' \cap A_{\mathcal{U}}$ is separably \mathcal{Z} -stable, and therefore pure by Proposition 3.1.

(2) \Rightarrow (3): This follows since pureness passes to quotients by [6, Theorem 4.11].

(3) \Rightarrow (1): Assume (3). We will verify condition (5) in Theorem 2.7, which then implies that A is separably \mathcal{Z} -stable. Let $B \subseteq A_{\mathcal{U}}$ be a separable sub- C^* -algebra. We need to find a unital $*$ -homomorphism $\mathcal{Z} \rightarrow (B' \cap A_{\mathcal{U}})/(B^{\perp} \cap A_{\mathcal{U}})$.

Set $Q := (B' \cap A_{\mathcal{U}})/(B^{\perp} \cap A_{\mathcal{U}})$ and let $\pi: B' \cap A_{\mathcal{U}} \rightarrow Q$ denote the quotient map. We first show that Q is $Z_{2,3}$ -saturated. Let $C \subseteq Q$ be a separable sub- C^* -algebra. Choose a separable preimage, that is, a separable sub- C^* -algebra $C_0 \subseteq B' \cap A_{\mathcal{U}}$ with $\pi(C_0) = C$. Since the sub- C^* -algebra E of $A_{\mathcal{U}}$ generated by C_0 and B is separable, it follows from the assumption that $(E' \cap A_{\mathcal{U}})/(E^{\perp} \cap A_{\mathcal{U}})$ is pure and therefore admits a unital $*$ -homomorphism from $Z_{2,3}$ by Corollary 3.3.

The map π induces a $*$ -homomorphism $E' \cap A_{\mathcal{U}} \rightarrow C' \cap Q$ whose image contains the unit of $C' \cap Q$. Since the annihilator of $C_0 \cup B$ is contained in the annihilator of B ,

this map factors through the quotient by $E^\perp \cap A_{\mathcal{U}}$. We obtain the desired map as the composition of the unital $*$ -homomorphisms

$$Z_{2,3} \rightarrow (E' \cap A_{\mathcal{U}})/(E^\perp \cap A_{\mathcal{U}}) \rightarrow C' \cap (B' \cap A_{\mathcal{U}})/(B^\perp \cap A_{\mathcal{U}}) = C' \cap Q.$$

This shows that Q is $Z_{2,3}$ -saturated.

Using this, we first find any unital $*$ -homomorphism $\varphi_1: Z_{2,3} \rightarrow Q$, and then successively unital $*$ -homomorphisms $\varphi_n: Z_{2,3} \rightarrow Q$ for $n \geq 2$ such that

$$\varphi_n(Z_{2,3}) \subseteq (\varphi_1(Z_{2,3}) \cup \dots \cup \varphi_{n-1}(Z_{2,3}))' \cap Q.$$

This induces a unital $*$ -homomorphism $\bigotimes_{\max}^\infty Z_{2,3} \rightarrow Q$. Using [9, Theorem 1.1], we obtain unital $*$ -homomorphisms

$$\mathcal{Z} \rightarrow \bigotimes_{\min}^\infty Z_{2,3} = \bigotimes_{\max}^\infty Z_{2,3} \rightarrow Q. \quad \square$$

4. Divisibility of central sequence algebras

In this section, we show that \mathcal{Z} -stability is equivalent to good central divisibility (that is, the central sequence algebra is *almost divisible*). One could phrase this by saying that good central comparison is a consequence of good central divisibility. In fact, \mathcal{Z} -stability already follows if the central sequence algebra satisfies the weaker property of *n-almost divisibility* for some n ; see Proposition 4.4. It remains unclear if it suffices to assume the even weaker property of *functional divisibility* in the central sequence algebras; see Question 4.5.

Given $N \in \mathbb{N}$ with $N \geq 1$, we define the relation \leq_N on a Cu-semigroup by setting $x \leq_N y$ if $nx \leq ny$ for all $n \geq N$. We recall the definition of controlled comparison from [5, Definition 3.6]; see also [3, Definition 6.1]. For this, for any $[a] \in \text{Cu}(A)$, we denote as customary by $\widehat{[a]}$ the (lower semicontinuous) function defined on $QT(A)$ by $\widehat{[a]}(\tau) = d_\tau(a)$.

Definition 4.1. A C^* -algebra A is said to have *controlled comparison* if for every $\gamma \in (0, 1)$ and $d \geq 1$, there exists $N = N(\gamma, d) \geq 1$ such that the following statement holds:

For every $a, b \in M_d(A)_+$, if $\widehat{[a]} \leq \gamma \widehat{[b]}$ then $[a] \leq_N [b]$.

We remark that controlled comparison was originally termed *locally bounded comparison amplitude* in [3, Definition 6.1].

We will use the characterization of controlled comparison for *stable* C^* -algebras given below, which is an easy consequence of [3, Corollary 7.6]. Recall the relation $<_s$ from Section 3.

Proposition 4.2. *Let A be a stable C^* -algebra. Then the following are equivalent:*

- (1) The C^* -algebra A has controlled comparison.
- (2) There exists $N \in \mathbb{N}$ such that $\widehat{x} \leq \widehat{y}$ implies $x \leq Ny$, for all $x, y \in \text{Cu}(A)$.
- (3) There exists $M \in \mathbb{N}$ such that $x <_s y$ implies $x \leq My$, for all $x, y \in \text{Cu}(A)$.

Proof. The equivalence of (1) and (2) is shown in [3, Corollary 7.6]. More concretely, it follows from the characterization given in [3, Theorem 7.5] that A having controlled comparison is equivalent to condition (i) in [3, Corollary 7.6], which in turn is equivalent to (2) above (that is, condition (iii) in [3, Corollary 7.6]).

Using that $x <_s y$ implies $\widehat{x} \leq \widehat{y}$, we deduce that (2) implies (3) with $M = N$. Conversely, assume that (3) holds with some $M \in \mathbb{N}$. We verify (2) for $N = 2M$. Let $x, y \in \text{Cu}(A)$ satisfy $\widehat{x} \leq \widehat{y}$. Then $\widehat{x} <_s 2\widehat{y}$. Let $x' \in \text{Cu}(A)$ with $x' \ll x$. By [4, Theorem 5.2.18], we deduce that $x' <_s 2y$, and thus $x' \leq 2My$ by assumption. Since this holds for every element x' way-below x , we deduce that $x \leq 2My$. \square

Proposition 4.3. *Let A be a C^* -algebra (not necessarily stable). Assume that there exists $M \in \mathbb{N}$ such that $x <_s y$ implies $x \leq My$, for all $x, y \in \text{Cu}(A)$. Then A has controlled comparison.*

Proof. By Proposition 4.2, we deduce that $A \otimes \mathcal{K}$ has controlled comparison. It follows easily from Definition 4.1 that a C^* -algebra has controlled comparison whenever its stabilization does. \square

Proposition 4.4. *Let A be a separable C^* -algebra. Then the following are equivalent:*

- (1) The C^* -algebra A is \mathcal{Z} -stable.
- (2) The algebra $A' \cap A_{\mathcal{U}}$ is almost divisible.
- (3) The unit of $F(A)$ is n -almost divisible for some $n \in \mathbb{N}$.

Proof. We show the implications ‘(1) \Rightarrow (2) \Rightarrow (3) \Rightarrow (1)’.

(1) \Rightarrow (2): If A is \mathcal{Z} -stable, then $A' \cap A_{\mathcal{U}}$ is pure by Theorem 3.6, and thus in particular almost divisible.

(2) \Rightarrow (3): Assume that $A' \cap A_{\mathcal{U}}$ is almost divisible. Since $F(A)$ is a quotient of $A' \cap A_{\mathcal{U}}$, and using that almost divisibility passes to quotients by [6, Lemma 4.9], it follows that $F(A)$ is almost divisible. In particular, the unit of $F(A)$ is n -almost divisible for some n (namely $n = 0$).

(3) \Rightarrow (1): Let $n \in \mathbb{N}$ be such that the unit in $F(A)$ is n -almost divisible. We first verify the following claim:

Claim: *Let $x, y \in \text{Cu}(F(A))$ satisfy $x <_s y$. Then $x \leq (n + 1)y$.*

To prove the claim, pick $m \in \mathbb{N}$ such that $(m + 1)x \leq my$. Using that [1] is n -almost divisible, we obtain $z \in \text{Cu}(F(A))$ such that

$$mz \leq [1] \leq (n + 1)(m + 1)z.$$

Since the elements x and y , and the Cuntz subequivalence $(m + 1)x \leq my$ are realized by countably many elements in $F(A) \otimes \mathcal{K}$, we can find a separable, unital sub- C^* -algebra $B \subseteq F(A)$ such that $\text{Cu}(B)$ contains elements \bar{x}, \bar{y} that are sent to x and y by the map $\text{Cu}(B) \rightarrow \text{Cu}(F(A))$ induced by the inclusion $B \rightarrow F(A)$, and such that $(m + 1)\bar{x} \leq m\bar{y}$ in $\text{Cu}(B)$. To elaborate on the construction of B , choose positive elements $a, b \in (F(A) \otimes \mathcal{K})_+$ such that $[a] = x$ and $[b] = y$ in $\text{Cu}(F(A))$. Since $(m + 1)x \leq my$, there exists a sequence $(v_k)_k$ in $M_{m+1,m}(F(A) \otimes \mathcal{K})$ such that $\|v_k^* \text{diag}(b, \dots, b)v_k - \text{diag}(a, \dots, a)\| \rightarrow 0$ as $k \rightarrow \infty$, where $\text{diag}(b, \dots, b)$ contains m copies of b and $\text{diag}(a, \dots, a)$ contains $m + 1$ copies of a . Let $S \subseteq F(A)$ be the countable set consisting of all matrix entries of a, b , and each v_k . Then $B_0 := C^*(S)$ is a separable sub- C^* -algebra of $F(A)$; let $\bar{a}, \bar{b} \in (B_0 \otimes \mathcal{K})_+$ denote the corresponding elements. By construction, $(m + 1)[\bar{a}] \leq m[\bar{b}]$ holds in $\text{Cu}(B_0)$. Finally, set $B := B_0 + \mathbb{C} \cdot 1_{F(A)}$, which is separable, unital, and contains elements $\bar{x} := [\bar{a}]$ and $\bar{y} := [\bar{b}]$ with the required properties.

Similarly, we can find a separable, unital sub- C^* -algebra $D \subseteq F(A)$ such that $\text{Cu}(D)$ contains an element \bar{z} such that $m\bar{z} \leq [1] \leq (n + 1)(m + 1)\bar{z}$ in $\text{Cu}(D)$.

By Proposition 3.5, we may assume that D is contained in $B' \cap F(A)$. We obtain a unital $*$ -homomorphism $\pi: B \otimes_{\max} D \rightarrow F(A)$ such that $\pi(b \otimes 1) = b$ for $b \in B$. As explained in [4, Paragraph 6.4.10], there exists a natural map $\varphi: \text{Cu}(B) \times \text{Cu}(D) \rightarrow \text{Cu}(B \otimes_{\max} D)$ such that $\varphi([b], [d]) = [b \otimes d]$ for $b \in B_+$ and $d \in D_+$. We consider the composition $\psi := \text{Cu}(\pi) \circ \varphi$, as shown in the following diagram:

$$\begin{array}{ccc} \text{Cu}(B) \times \text{Cu}(D) & \xrightarrow{\varphi} & \text{Cu}(B \otimes_{\max} D) & \xrightarrow{\text{Cu}(\pi)} & \text{Cu}(F(A)). \\ & & \searrow \psi & \nearrow & \\ & & & & \end{array}$$

We note that ψ is additive and order-preserving in each entry, and that $\psi(\bar{x}, [1]) = x$ and $\psi(\bar{y}, [1]) = y$. We deduce that

$$\begin{aligned} x = \varphi(\bar{x}, [1]) &\leq \varphi(\bar{x}, (n + 1)(m + 1)z) \\ &= \varphi((n + 1)(m + 1)\bar{x}, z) \\ &\leq \varphi((n + 1)m\bar{y}, z) \\ &= \varphi((n + 1)\bar{y}, mz) \leq \varphi((n + 1)\bar{y}, [1]) = (n + 1)y, \end{aligned}$$

which proves the claim.

Now, it follows from Proposition 4.3 that $F(A)$ has controlled comparison. By similar arguments as in the proof of the claim, one shows that the n -almost divisibility of the class of the unit in $F(A)$ propagates to all Cuntz classes, which means that $F(A)$ is n -almost divisible. By [5, Proposition 4.9], we get that $F(A)$ is functionally divisible. It then follows from [5, Theorem 5.7] that $F(A)$ is pure. Applying Theorem 3.6, we deduce that A is \mathcal{Z} -stable. \square

In [5], the authors introduced *functional divisibility* as a property weaker than n -almost divisibility for every n and that, together with controlled comparison, still implies pureness. With a view towards Proposition 4.4, the following question arises:

Question 4.5. Is a separable C^* -algebra \mathcal{Z} -stable whenever its central sequence algebra is functionally divisible?

Acknowledgments

The authors wish to thank the anonymous referee for a careful reading and for making suggestions that have improved the manuscript.

Data availability

No data was used for the research described in the article.

References

- [1] T. Amrutam, D. Gao, S. Kunnawalkam Elayavalli, G. Patchell, Strict comparison in reduced group C^* -algebras, *Invent. Math.* 242 (2025) 639–657.
- [2] R. Antoine, F. Perera, L. Robert, H. Thiel, C^* -algebras of stable rank one and their Cuntz semigroups, *Duke Math. J.* 171 (2022) 33–99.
- [3] R. Antoine, F. Perera, L. Robert, H. Thiel, Traces on ultrapowers of C^* -algebras, *J. Funct. Anal.* 286 (2024) 110341.
- [4] R. Antoine, F. Perera, H. Thiel, Tensor products and regularity properties of Cuntz semigroups, *Mem. Am. Math. Soc.* 251 (2018), viii+191.
- [5] R. Antoine, F. Perera, H. Thiel, E. Vilalta, Pure C^* -algebras, preprint, arXiv:2406.11052 [math.OA], 2024.
- [6] R. Antoine, F. Perera, H. Thiel, E. Vilalta, Extensions of pure C^* -algebras, *Trans. Am. Math. Soc.* (2026), in press, arXiv:2506.10529 [math.OA].
- [7] P. Ara, F. Perera, A.S. Toms, K -theory for operator algebras. Classification of C^* -algebras, in: *Aspects of Operator Algebras and Applications*, in: *Contemp. Math.*, vol. 534, Amer. Math. Soc., Providence, RI, 2011, pp. 1–71.
- [8] J.R. Carrión, J. Gabe, C. Schafhauser, A. Tikuisis, S. White, Classifying $*$ -homomorphisms I: unital simple nuclear C^* -algebras, preprint, arXiv:2307.06480 [math.OA], 2023.
- [9] M. Dadarlat, A.S. Toms, \mathcal{Z} -stability and infinite tensor powers of C^* -algebras, *Adv. Math.* 220 (2009) 341–366.
- [10] M. Dadarlat, A.S. Toms, Ranks of operators in simple C^* -algebras, *J. Funct. Anal.* 259 (2010) 1209–1229.
- [11] G.A. Elliott, G. Gong, H. Lin, Z. Niu, On the classification of simple amenable C^* -algebras with finite decomposition rank, II, *J. Noncommut. Geom.* 19 (2025) 73–104.
- [12] I. Farah, B. Hart, M. Lupini, L. Robert, A. Tikuisis, A. Vignati, W. Winter, Model theory of C^* -algebras, *Mem. Am. Math. Soc.* 271 (2021), viii+127.
- [13] I. Farah, G. Szabó, Coronas and strongly self-absorbing C^* -algebras, preprint, arXiv:2411.02274, 2024.
- [14] E. Gardella, F. Perera, The modern theory of Cuntz semigroups of C^* -algebras, *EMS Surv. Math. Sci.* 13 (2026) 133–214.
- [15] G. Gong, H. Lin, Z. Niu, A classification of finite simple amenable \mathcal{Z} -stable C^* -algebras, II: C^* -algebras with rational generalized tracial rank one, *C. R. Math. Acad. Sci. Soc. R. Can.* 42 (2020) 451–539.
- [16] E. Kirchberg, Central sequences in C^* -algebras and strongly purely infinite algebras, in: *Operator Algebras: The Abel Symposium 2004*, in: *Abel Symp.*, vol. 1, Springer, Berlin, 2006, pp. 175–231.

- [17] E. Kirchberg, M. Rørdam, Central sequence C^* -algebras and tensorial absorption of the Jiang-Su algebra, *J. Reine Angew. Math.* 695 (2014) 175–214.
- [18] E. Kirchberg, M. Rørdam, When central sequence C^* -algebras have characters, *Int. J. Math.* 26 (2015) 1550049.
- [19] N. Nawata, Picard groups of certain stably projectionless C^* -algebras, *J. Lond. Math. Soc. (2)* 88 (2013) 161–180.
- [20] P.W. Ng, L. Robert, Sums of commutators in pure C^* -algebras, *Münster J. Math.* 9 (2016) 121–154.
- [21] L. Robert, M. Rørdam, Divisibility properties for C^* -algebras, *Proc. Lond. Math. Soc. (3)* 106 (2013) 1330–1370.
- [22] L. Robert, A. Tikuisis, Nuclear dimension and \mathcal{Z} -stability of non-simple C^* -algebras, *Trans. Am. Math. Soc.* 369 (2017) 4631–4670.
- [23] M. Rørdam, Classification of nuclear, simple C^* -algebras, in: *Classification of Nuclear C^* -Algebras. Entropy in Operator Algebras*, in: *Encyclopaedia Math. Sci.*, vol. 126, Springer, Berlin, 2002, pp. 1–145.
- [24] M. Rørdam, The stable and the real rank of \mathcal{Z} -absorbing C^* -algebras, *Int. J. Math.* 15 (2004) 1065–1084.
- [25] M. Rørdam, W. Winter, The Jiang-Su algebra revisited, *J. Reine Angew. Math.* 642 (2010) 129–155.
- [26] Y. Sato, Trace spaces of simple nuclear C^* -algebras with finite-dimensional extreme boundary, preprint, arXiv:1209.3000 [math.OA], 2012.
- [27] C. Schafhauser, Subalgebras of simple AF-algebras, *Ann. Math. (2)* 192 (2020) 309–352.
- [28] C. Schafhauser, KK-rigidity of simple nuclear C^* -algebras, preprint, arXiv:2408.02745 [math.OA], 2024.
- [29] H. Thiel, Ranks of operators in simple C^* -algebras with stable rank one, *Commun. Math. Phys.* 377 (2020) 37–76.
- [30] A. Tikuisis, S. White, W. Winter, Quasidiagonality of nuclear C^* -algebras, *Ann. Math. (2)* 185 (2017) 229–284.
- [31] A.S. Toms, On the classification problem for nuclear C^* -algebras, *Ann. Math. (2)* 167 (2008) 1029–1044.
- [32] A.S. Toms, S. White, W. Winter, \mathcal{Z} -stability and finite-dimensional tracial boundaries, *Int. Math. Res. Not.* (2015) 2702–2727.
- [33] A.S. Toms, W. Winter, Strongly self-absorbing C^* -algebras, *Trans. Am. Math. Soc.* 359 (2007) 3999–4029.
- [34] S. White, Abstract classification theorems for amenable C^* -algebras, in: *ICM—International Congress of Mathematicians*, vol. 4. Sections 5–8, EMS Press, Berlin, 2023, pp. 3314–3338.
- [35] W. Winter, Strongly self-absorbing C^* -algebras are \mathcal{Z} -stable, *J. Noncommut. Geom.* 5 (2011) 253–264.
- [36] W. Winter, Nuclear dimension and \mathcal{Z} -stability of pure C^* -algebras, *Invent. Math.* 187 (2012) 259–342.
- [37] W. Winter, Structure of nuclear C^* -algebras: from quasidiagonality to classification and back again, in: *Proceedings of the International Congress of Mathematicians—Rio de Janeiro 2018*, vol. III. Invited Lectures, World Sci. Publ., Hackensack, NJ, 2018, pp. 1801–1823.