



LIE IDEALS IN PROPERLY INFINITE C^* -ALGEBRAS

Downloaded from: <https://research.chalmers.se>, 2026-07-12 16:24 UTC

Citation for the original published paper (version of record):

Thiel, H. (2026). LIE IDEALS IN PROPERLY INFINITE C^* -ALGEBRAS. *Publicacions Matemàtiques*, 70(2): 525-548. <http://dx.doi.org/10.5565/PUBLMAT7022609>

N.B. When citing this work, cite the original published paper.

LIE IDEALS IN PROPERLY INFINITE C^* -ALGEBRAS

HANNES THIEL

Abstract: We show that every Lie ideal in a unital, properly infinite C^* -algebra is commutator-equivalent to a unique two-sided ideal. It follows that the Lie ideal structure of such a C^* -algebra is concisely encoded by its lattice of two-sided ideals. This answers a question of Robert in this setting.

We obtain similar structure results for Lie ideals in unital, real rank zero C^* -algebras without characters. As an application, we show that every Lie ideal in a von Neumann algebra is related to a unique two-sided ideal, which solves a problem of Brešar, Kissin, and Shulman.

2020 Mathematics Subject Classification: Primary: 46L05, 46L10. Secondary: 16W10, 17B60, 47B47.

Key words: Lie ideals, C^* -algebras, properly infinite, von Neumann algebras, commutators, square-zero elements.

1. Introduction

A *Lie ideal* in a C^* -algebra A is a linear subspace $L \subseteq A$ such that for every $a \in A$ and $x \in L$ the commutator $[a, x] := ax - xa$ belongs to L . The study of Lie ideals in C^* -algebras has a long history, and we refer to Section 2 in [25] for an overview of available results and directions of active research.

Immediate examples of Lie ideals in a C^* -algebra A are all subspaces contained in the center $Z(A)$ and all subspaces containing the commutator subspace $[A, A]$. If A is unital and simple, then this does in fact describe all Lie ideals—a consequence of Herstein’s description of Lie ideals in simple rings [20]. Since the center of a unital, simple C^* -algebra A contains only scalar multiples of the identity, a subspace L is a Lie ideal in A if and only if $L = \{0\}$, $L = \mathbb{C}1$, or $[A, A] \subseteq L$.

Pop showed that a unital C^* -algebra satisfies $A = [A, A]$ if and only if A has no tracial states [30]. Consequently, a simple, unital C^* -algebra A without tracial states has exactly three Lie ideals: $\{0\}$, $\mathbb{C}1$, and A . Cuntz and Pedersen gave a description of the closure $\overline{[A, A]}$ as the intersection of the kernels of all tracial states on A [10]. In particular, a simple, unital C^* -algebra A with a unique tracial state has exactly four closed Lie ideals: $\{0\}$, $\mathbb{C}1$, $\overline{[A, A]}$, and A . This was first observed by Marcoux and Murphy [26], generalizing the case of UHF algebras, which was first proved by Marcoux [24].

Ng and Robert showed that $[A, A]$ is closed in every pure, exact C^* -algebra [28]. It follows that a simple, unital, exact, pure C^* -algebra A with a unique tracial state has exactly four Lie ideals: $\{0\}$, $\mathbb{C}1$, $[A, A]$, and A . This applies for example to UHF algebras, to irrational rotation algebras, and to the Jiang–Su algebra (where it was first shown by Ng [27]). It also applies to many, and possibly all, simple, reduced group C^* -algebras. Indeed, if G is a countable, exact, acylindrically hyperbolic group with trivial finite radical and with the rapid decay property, then $C_{\text{red}}^*(G)$ has strict comparison [1], and therefore is pure [3], and has a unique tracial state [9].

Turning towards Lie ideals in nonsimple C^* -algebras, it is clear that the structure of the lattice of two-sided ideals will play a role. A systematic study of the interplay between Lie and two-sided ideals was initiated by Brešar, Kissin, and Shulman [8]. We first introduce the necessary notation. Following the usual convention, given subspaces $M, N \subseteq A$, we use $[M, N]$ to denote the subspace of A generated by the set of commutators $[x, y]$ for $x \in M$ and $y \in N$. Further, given a subspace $K \subseteq A$, we set

$$T(K) := \{a \in A : [A, a] \subseteq K\}.$$

This ‘derived subspace’ was first considered by Herstein [20], and it is denoted by $N(K)$ in [8].

Given a two-sided ideal $I \subseteq A$, Brešar, Kissin, and Shulman observed that a subspace $L \subseteq A$ is a Lie ideal whenever

$$[A, I] \subseteq L \subseteq T([A, I]).$$

We say that a Lie ideal L is *embraced by* I if it satisfies these inclusions. The Lie ideals that are embraced by some two-sided ideal are thought of as ‘unsurprising’. A major question is if there exist any Lie ideals that do not arise this way:

Problem A (Brešar, Kissin, and Shulman [8, p. 74]). Given a C^* -algebra A , is every Lie ideal of A embraced by some two-sided ideal of A ?

To study this problem, two further notions were introduced in [8]: Given a Lie ideal $L \subseteq A$ and a two-sided ideal $I \subseteq A$, we say that L is *commutator-equivalent to* I if

$$[A, I] = [A, L].$$

Further, we say that L is *related to* I if

$$[A, I] \subseteq L \subseteq T(I).$$

It is easy to see that these constitute a sufficient and a necessary conditions for embracement. Indeed, if L is commutator-equivalent to I , then L is embraced by I . Further, if L is embraced by I , then L is related to I .

It was shown in [8, Theorem 5.27] that every Lie ideal L in a C^* -algebra A is topologically commutator-equivalent to a two-sided ideal I in the sense that $[A, L]$ and $[A, I]$ have the same closure. This leads to a description of all *closed* Lie ideals in terms of *closed* two-sided ideals. However, this does not show that all closed Lie ideals in a C^* -algebra are commutator-equivalent to (let alone embraced by or related to) a two-sided ideal. For general nonclosed Lie ideals even less is known, and Problem A as well as the following stronger problem remain open:

Problem B (Robert [31, Question 1.12]). Given a C^* -algebra A , is every Lie ideal of A commutator-equivalent to some two-sided ideal of A ?

The results mentioned at the beginning of the introduction show that every Lie ideal in a unital, simple C^* -algebra A is commutator-equivalent to a two-sided ideal, which gives positive solutions to Problems A and B in this setting. Indeed, the only two-sided ideals are $\{0\}$ and A , and a Lie ideal is commutator-equivalent to $\{0\}$ if and only if it is contained in $Z(A)$, while a Lie ideal is commutator-equivalent to A if and only if it contains $[A, A]$. (This uses that $[A, A] = [A, [A, A]]$, which follows for example from [17, Corollary 3.4].)

In [13], Fong, Miers, and Sourour showed that every Lie ideal in the von Neumann algebra of bounded, linear operators on a separable Hilbert space is embraced by a two-sided ideal. This was substantially extended by Brešar, Kissin, and Shulman, who showed in [8, Theorem 5.19] that every Lie ideal in a von Neumann algebra is

commutator-equivalent to a two-sided ideal, thereby providing positive solutions to Problems A and B for von Neumann algebras.

Our main result is a positive solution to Problem B for unital, properly infinite C^* -algebras.

Theorem C (5.3). *Let L be a Lie ideal in a unital, properly infinite C^* -algebra A . Then L is commutator-equivalent to the two-sided ideal $I := A[A, L]A$, and hence also embraced by I , and related to I . Further, I is the only two-sided ideal of A to which L is related.*

The proof of Theorem C is inspired by a result of Marcoux [24], who showed that if B is a unital C^* -algebra and $n \geq 2$, then every Lie ideal in the matrix algebra $A = M_n(B)$ is related to a two-sided ideal in A . Our main technical advancement is a generalization of this result to unital C^* -algebras A that admit a unital $*$ -homomorphism $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow A$.

The next result summarizes Theorems 2.7, 2.9, 3.8, and 3.10.

Theorem D. *Let A be a unital C^* -algebra that admits a unital $*$ -homomorphism $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow A$. The following statements hold:*

- (1) *Given a Lie ideal $L \subseteq A$, for the two-sided ideal $I = A[A, L]A$, we have*

$$[A, I] = [A, [A, L]] \subseteq [A, L] \subseteq I \quad \text{and} \quad [A, I] \subseteq L \subseteq T(I).$$

Further, I is the only two-sided ideal of A to which L is related.

- (2) *A Lie ideal $L \subseteq A$ is embraced by some two-sided ideal (which then necessarily is $A[A, L]A$) if and only if L is commutator-equivalent to some two-sided ideal, if and only if $[A, L] = [A, [A, L]]$.*
- (3) *Given a two-sided ideal $I \subseteq A$, we have*

$$I = A[A, I]A \quad \text{and} \quad [A, I] = [A, [A, I]].$$

- (4) *There is a natural bijection between the set $\mathcal{I}$ of two-sided ideals in A and the set*

$$\mathcal{L} = \{L \subseteq A \text{ Lie ideal} : L = [A, L]\}$$

given by the maps $I \mapsto [A, I]$ and $L \mapsto A[A, L]A$.

We note that Theorem D applies to all unital, properly infinite C^* -algebras, which then yields Theorem C. It also applies to unital, real rank zero C^* -algebras without characters, and in particular to all von Neumann algebras with zero commutative summand, which we use to recover the solution to Problem B for von Neumann algebras from [8]. Our approach also shows that a Lie ideal in a von Neumann algebra without commutative summand is related to a unique two-sided ideal, which solves [8, Problem 5.21].

Theorem E (6.1). *Let L be a Lie ideal in a von Neumann algebra M . Then L is commutator-equivalent to the two-sided ideal $I := M[M, L]M$, and hence also embraced by I , and related to I .*

If M has zero commutative summand, then I is the only two-sided ideal of M to which L is related.

Conventions. Given a C^* -algebra, we use A_+ to denote its collection of positive elements, and we use $\tilde{A}$ to denote the minimal unitization. The absolute value of an element $x \in A$ is defined as $|x| := (x^*x)^{\frac{1}{2}}$.

2. Lie ideals in C^* -algebras with weakly divisible unit

In this section, we prove that every Lie ideal L in a C^* -algebra A that admits a unital $*$ -homomorphism $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow A$ is related to a unique two-sided ideal I in A ; see Theorem 2.7. We further show that L is commutator-equivalent to I if and only if $[A, L] = [A, [A, L]]$; see Theorem 2.9.

Following [29, Definition 5.1], a nonzero projection p in a C^* -algebra is said to be *weakly divisible of degree n* if there exists a unital $*$ -homomorphism

$$M_{n_1}(\mathbb{C}) \oplus M_{n_2}(\mathbb{C}) \oplus \cdots \oplus M_{n_r}(\mathbb{C}) \rightarrow pAp$$

for some natural numbers $r \geq 1$ and $n_1, \dots, n_r \geq n$. The $*$ -homomorphism may additionally be assumed to be injective.

As noted in [29, p. 164], every matrix algebra $M_n(\mathbb{C})$ for $n \geq 2$ admits a unital (not necessarily injective) $*$ -homomorphism $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow M_n(\mathbb{C})$, and it follows that a projection p weakly divisible of degree 2 if and only if there exists a unital (not necessarily injective) $*$ -homomorphism $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow pAp$. We deduce that the unit in a C^* -algebra A is weakly divisible of degree 2 if and only if there exists an injective $*$ -homomorphism

$$M_2(\mathbb{C}) \rightarrow A, \quad M_3(\mathbb{C}) \rightarrow A, \quad \text{or} \quad M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow A.$$

In the first two cases, we get $A \cong M_2(B)$ and $A \cong M_3(C)$ for suitable unital C^* -algebras B and C , and for such matrix algebras it was shown by Marcoux ([24, Theorem 2.6]) that every Lie ideal is related to a two-sided ideal (which is also unique, as shown in [8, Corollary 4.18]).

Our main contribution to Theorem 2.7 is therefore in the case that A admits a unital (injective) $*$ -homomorphism $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow A$ but no unital $*$ -homomorphism $M_n(\mathbb{C}) \rightarrow A$ for $n = 2$ or $n = 3$ (or actually any $n \geq 2$). The Cuntz algebra $\mathcal{O}_\infty$ is an important such example. (Using K -theory, we see that there exists no unital $*$ -homomorphism $M_n(\mathbb{C}) \rightarrow \mathcal{O}_\infty$ for any $n \geq 2$. On the other hand, $\mathcal{O}_\infty$ admits a unital $*$ -homomorphism from $M_2(\mathbb{C}) \oplus M_3(\mathbb{C})$ as explained in the proof of Theorem 5.3.)

Our approach is inspired by the methods developed by Marcoux ([24]) to describe Lie ideals in matrix algebras, as well as the techniques of Brešar, Kissin, and Shulman ([8, Section 3]) to study Lie ideals in algebras that decompose with respect to an idempotent. However, the result does not simply follow by combining these methods, for example by applying [8, Theorem 3.4] for a suitable idempotent in the C^* -algebra. Specifically, given a unital $*$ -homomorphism $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow A$, we do not decompose A with respect to the projection p given by the image of the unit of $M_2(\mathbb{C})$, in which case $pAp + (1-p)A(1-p)$ is a direct sum of matrix algebras (whence its Lie ideals are related to two-sided ideals), but it is unclear if (3.17) in [8, Theorem 3.4] holds (or if p is locally cyclic in the sense of [8, Definition 3.5]). Instead, we use a ‘mixed’ decomposition with respect to a projection g that is given by the image of the sum of a rank one projection in $M_2(\mathbb{C})$ with a rank one projection in $M_3(\mathbb{C})$, as explained in Paragraph 2.1. In this case, one can show that g is locally cyclic, which is the idea underlying the proof that $A_{12}LA_{12} \subseteq [A, [A, L]]$ in Lemma 2.2. On the other hand, the algebra $gAg + (1-g)A(1-g)$ is not a sum of matrix algebras and it is therefore not immediate that its Lie ideals are related to two-sided ideals.

2.1. Let A be a unital C^* -algebra that admits a unital $*$ -homomorphism

$$\varphi: M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow A.$$

Let $(e_{ij})_{i,j=1,2}$ denote the image of matrix units in $M_2(\mathbb{C})$ under φ , and similarly let $(f_{ij})_{i,j=1,2,3}$ denote the image of matrix units in $M_3(\mathbb{C})$ under φ . We do not assume that φ is injective, and therefore the e_{ij} or f_{ij} may be zero.

Set

$$g := e_{11} + f_{11} \quad \text{and} \quad h := e_{22} + f_{22} + f_{33}.$$

Then $1 = g + h$, and we view elements in A as operator matrices with respect to the decomposition induced by g and h . More formally, we set

$$A_{11} := gAg, \quad A_{12} := gAh, \quad A_{21} := hAg, \quad \text{and} \quad A_{22} := hAh,$$

and every element a in A can be written uniquely as a sum $a = a_{11} + a_{12} + a_{21} + a_{22}$ with $a_{ij} \in A_{ij}$ for $i, j = 1, 2$.

Lemma 2.2. *Retain the notation from Paragraph 2.1, and let $L \subseteq A$ be a Lie ideal. Then the sets*

$$A_{11}LA_{22}, \quad A_{22}LA_{11}, \quad A_{12}LA_{12}, \quad \text{and} \quad A_{21}LA_{21}$$

are all contained in $[A, [A, L]]$, and thus also in $[A, L]$ and in L .

Proof: We verify that $A_{11}LA_{22} \subseteq [A, [A, L]]$. We first show that $gLh \subseteq L$. To see this, let $x \in L$. Then

$$gx - xg = [g, x] \in [A, L] \quad \text{and} \quad gx - 2gxg + xg = [g, [g, x]] \in [A, [A, L]] \subseteq [A, L].$$

Adding these expressions, we get

$$2gxh = 2gx - 2gxg \in [A, L] \subseteq L,$$

and using that $hg = 0$ we get

$$gxh = [g, gxh] = [\tfrac{1}{2}g, 2gxh] \in [A, L] \subseteq L.$$

We have shown that $gLh \subseteq L$.

Now, given $x \in L$ and $a, b \in A$, we have

$$gagxhbh = [[gag, gxh], hbh] \in [[A, L], A] = [A, [A, L]].$$

We have shown that $A_{11}LA_{22} \subseteq [A, [A, L]]$.

Analogously, one verifies that $A_{22}LA_{11} \subseteq [A, [A, L]]$.

We verify that $A_{12}LA_{12} \subseteq [A, [A, L]]$. Given $v \in A_{12}$ and $y \in L$, we have

$$(2.1) \quad v y v = [-\tfrac{1}{2}v, [v, y]] \in [A, [A, L]].$$

Set

$$v_1 := e_{12} + f_{12}, \quad w_1 := e_{21} + f_{21}, \quad v_2 := e_{12} + f_{13}, \quad \text{and} \quad w_2 := e_{21} + f_{31}.$$

Then $v_1, v_2 \in A_{12}$ and $w_1, w_2 \in A_{21}$, and we have

$$v_1 w_1 = v_2 w_2 = e_{11} + f_{11} = g, \quad w_1 v_1 = e_{22} + f_{22}, \quad \text{and} \quad w_2 v_2 = e_{22} + f_{33}.$$

Now, given $x \in L$ and $a, b \in A_{12} = gAh$, set

$$b' = b(\tfrac{1}{2}e_{22} + f_{22} + f_{33}).$$

Then b' belongs to gAh and we have

$$b = b'(2e_{22} + f_{22} + f_{33}).$$

We have already proved that $hAhLgAg \subseteq L$, which gives

$$w_1 a x b' w_1 \in L \quad \text{and} \quad w_2 a x b' w_2 \in L.$$

Applying (2.1) for $y = w_1axb'w_1$ and $v = v_1$, we get

$$axb'(e_{22} + f_{22}) = v_1(w_1axb'w_1)v_1 \in [A, [A, L]].$$

Analogously, we obtain that

$$axb'(e_{22} + f_{33}) = v_2(w_2axb'w_2)v_2 \in [A, [A, L]],$$

and consequently

$$axb = axb'(2e_{22} + f_{22} + f_{33}) \in [A, [A, L]].$$

We have shown that $A_{12}LA_{12} \subseteq [A, [A, L]]$.

Analogously, one verifies that $A_{21}LA_{21} \subseteq [A, [A, L]]$. $\square$

Lemma 2.3. *Retain the notation from Paragraph 2.1, and let $L \subseteq A$ be a Lie ideal. Then*

$$AgLhA = AhLgA.$$

Proof: Using at the first step that $A = AgA = AhA$ since g and h are full, and applying at the third step that $hAgLhAg \subseteq L$ by Lemma 2.2, we get

$$AgLhA = AhAgLhAgA = Ah(hAgLhAg)gA \subseteq AhLgA.$$

Analogously, we have

$$AhLgA = AgAhLgAhA = Ag(gAhLgAh)hA \subseteq AgLhA,$$

as desired. $\square$

Lemma 2.4. *Retain the notation from Paragraph 2.1, and let $L \subseteq A$ be a Lie ideal. Then*

$$[A, L] \subseteq AgLhA.$$

Proof: To simplify notation, we set $I = AgLhA$. We have

$$[A, gLh] \subseteq AgLh + gLhA \subseteq I.$$

Further, using Lemma 2.3 at the third step, we have

$$[A, hLg] \subseteq AhLg + hLgA \subseteq AhLgA = AgLhA = I.$$

Thus, given $x \in L$, we have $[A, gxh + hxx] \subseteq I$. Note that $gxh, hxx \in L$ by Lemma 2.2. It follows that

$$gxg + hxx = x - gxh - hxx \in L.$$

We need to show that $[A, gxg + hxx] \subseteq I$. Using additivity of the Lie bracket and that I is an additive subgroup, it suffices to verify that $[A_{ij}, gxg + hxx] \subseteq I$ for $i, j = 1, 2$.

We show that $[A_{12}, gxg + hxx] \subseteq I$. Given $a \in A$, we have

$$[gah, gxg + hxx] = gahxh - gxgah.$$

Using that $gxg + hxx \in L$, we have $[gah, gxg + hxx] \in L$, and thus

$$[gah, gxg + hxx] \in L \cap gAh = gLh \subseteq AgLhA = I.$$

We have shown that $[A_{12}, gxg + hxx] \subseteq I$.

Similarly, and using also Lemma 2.3 at the last step, we get

$$[hag, gxg + hxx] = hagxg - hxxag \in L \cap hAg = hLg \subseteq AhLgA = I$$

for every $a \in A$. This shows that $[A_{21}, gxg + hxx] \subseteq I$.

We show that $[A_{11}, g x g + h x h] \subseteq I$. Given $a \in A$, set $y = g a g x g - g x g a g$. Then

$$y = g a g x g - g x g a g = [g a g, g x g + h x h] \in [A, L] \subseteq L.$$

Then

$$y(e_{12} + f_{12}) = [y, e_{12} + f_{12}] \in [L, A] \subseteq L$$

and thus

$$y(e_{12} + f_{12}) \in L \cap g A h = g L h \subseteq I.$$

We deduce that

$$y = y(e_{12} + f_{12})(e_{21} + f_{21}) \in I A = I.$$

We show that $[A_{22}, g x g + h x h] \subseteq I$. Given $b \in A$, set $z = h b h x h - h x h b h$. Then

$$z = h b h x h - h x h b h = [h b h, g x g + h x h] \in L \cap A_{22}.$$

Then

$$(e_{12} + f_{12})z = [e_{12} + f_{12}, z] \in L \cap g A h \subseteq I \quad \text{and} \quad f_{13}z = [f_{13}, z] \in I.$$

We deduce that

$$z = (e_{21} + f_{21})(e_{12} + f_{12})z + f_{31}f_{13}z \in A I + A I = I,$$

as desired. $\square$

Proposition 2.5. *Retain the notation from Paragraph 2.1, and let $L \subseteq A$ be a Lie ideal. Then*

$$A[A, L] = A[A, L]A = A g L h A = A h L g A.$$

Proof: In general, if $M \subseteq A$ is a Lie ideal, then $A M A = A M$; see, for example, [15, Lemma 3.2]. Since $[A, L]$ is a Lie ideal, we get $A[A, L] = A[A, L]A$.

By Lemma 2.4, we have $[A, L] \subseteq A g L h A$, and thus $A[A, L]A \subseteq A g L h A$. By Lemma 2.3, we have $A g L h A = A h L g A$. Finally, by Lemma 2.2, we have $g L h \subseteq [A, L]$, and thus $A g L h A \subseteq A[A, L]A$. $\square$

Lemma 2.6. *Retain the notation from Paragraph 2.1, and let $L \subseteq A$ be a Lie ideal, and set $I = A[A, L]A$. Then*

$$[A, I] \subseteq [A, [A, L]].$$

Proof: For each $i, j = 1, 2$, set $I_{ij} := I \cap A_{ij}$. Since I is a two-sided ideal, we have $I = I_{11} + I_{12} + I_{21} + I_{22}$. By Proposition 2.5, we have $I = A g L h A$, and thus

$$I = A g L h A = \underbrace{g A g L h A g}_{I_{11}} + \underbrace{g A g L h A h}_{I_{12}} + \underbrace{h A g L h A g}_{I_{21}} + \underbrace{h A g L h A h}_{I_{22}}.$$

By additivity of the Lie bracket and since $[A, L]$ is an additive subgroup, it suffices to show that $[A, I_{ij}] \subseteq [A, [A, L]]$ for each $i, j = 1, 2$.

By Lemma 2.2, we have

$$I_{12} = g A g L h A h \subseteq [A, L] \quad \text{and} \quad I_{21} = h A g L h A g \subseteq [A, L],$$

and thus $[A, I_{12}] \subseteq [A, [A, L]]$ and $[A, I_{21}] \subseteq [A, [A, L]]$.

We show that $[A, I_{11}] \subseteq [A, [A, L]]$. It suffices to verify that $[A_{ij}, I_{11}] \subseteq [A, [A, L]]$ for each $i, j = 1, 2$. Using that $h g = 0$, and using Lemma 2.2 at the last step, we get

$$[A_{12}, I_{11}] = [g A h, g A g L h A g] = -(g A g L h A g)(g A h) = g A g L h A g A h \subseteq [A, [A, L]].$$

Similarly, we have

$$[A_{21}, I_{11}] = [hAg, gAgLhAg] = (hAg)(gAgLhAg) = hAgAgLhAg \subseteq [A, [A, L]].$$

We also have $[A_{22}, I_{11}] = \{0\} \subseteq [A, [A, L]]$.

Further, using Lemma 2.2 at the last step, we have

$$I_{11} = (gAgLh)Ag \subseteq [gAgLh, Ag] + Ag(gAgLh) \subseteq [A, L] + AgLh.$$

Using again Lemma 2.2 at the last step, we get

$$\begin{aligned} [A_{11}, I_{11}] &\subseteq [A_{11}, [A, L]] + [A_{11}, AgLh] \subseteq [A, [A, L]] + [gAg, AgLh] \\ &\subseteq [A, [A, L]] + gAgAgLh \subseteq [A, [A, L]]. \end{aligned}$$

We have verified that $[A, I_{11}] \subseteq [A, [A, L]]$.

Analogously, one shows that $[A, I_{22}] \subseteq [A, [A, L]]$, which completes the proof. $\square$

The next result verifies statement (1) in Theorem D.

Theorem 2.7. *Let A be a unital C^* -algebra that admits a unital $*$ -homomorphism $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow A$, and let $L \subseteq A$ be a (not necessarily closed) Lie ideal. Then, for the two-sided ideal $I := A[A, L]A$, we have*

$$(2.2) \quad [A, I] = [A, [A, L]] \subseteq [A, L] \subseteq I \quad \text{and} \quad [A, I] \subseteq L \subseteq T(I).$$

Further, I is the only two-sided ideal of A to which L is related.

Proof: Using Lemma 2.4 at the first step, and Proposition 2.5 at the second step, we have

$$[A, L] \subseteq AgLhA = I.$$

This implies that $[A, [A, L]] \subseteq [A, I]$, and the converse inclusion $[A, I] \subseteq [A, [A, L]]$ is shown in Lemma 2.6. Using that L is a Lie ideal, we have $[A, L] \subseteq L$, and therefore $[A, [A, L]] \subseteq [A, L]$. This shows the left chain of inclusions in (2.2).

The inclusion $[A, I] \subseteq L$ follows using that $[A, I] \subseteq [A, L]$ and $[A, L] \subseteq L$. Finally, since $[A, L] \subseteq I$, we have $L \subseteq T(I)$.

To show the uniqueness of I , let J be a two-sided ideal that is related to L , that is, such that

$$[A, J] \subseteq L \subseteq T(J).$$

Then $[A, L] \subseteq J$, and since I is the two-sided ideal generated by $[A, L]$ we get $I \subseteq J$.

Applying Lemma 2.2 at the first step for J , considered as a Lie ideal, we get

$$gJh \subseteq [A, J] \subseteq L,$$

and thus

$$gJh \subseteq L \cap gAh = gLh.$$

Since g and h are full, we have $A = AgA$ and $A = AhA$, and consequently

$$J = AJA = AgAJAhA = AgJhA \subseteq AgLhA = I.$$

In conclusion, we get $I = J$, as desired. $\square$

Corollary 2.8. *Let A be a unital C^* -algebra that admits a unital $*$ -homomorphism $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow A$, and let $L \subseteq A$ be a closed Lie ideal. Then, the two-sided ideal $I := A[A, L]A$ is closed.*

Proof: By Theorem 2.7, L is related to I , that is, we have $[A, I] \subseteq L \subseteq T(I)$. Using that L is closed, it follows that the closure $\bar{I}$ satisfies $[A, \bar{I}] \subseteq L \subseteq T(\bar{I})$. Since I is the only two-sided ideal of A to which L is related (Theorem 2.7), we get $I = \bar{I}$. $\square$

The next result verifies statement (2) in Theorem D.

Theorem 2.9. *Let A be a unital C^* -algebra that admits a unital $*$ -homomorphism $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow A$, and let $L \subseteq A$ be a (not necessarily closed) Lie ideal. Then the following statements are equivalent:*

- (1) *The Lie ideal L is commutator-equivalent to the two-sided ideal $A[A, L]A$.*
- (2) *The Lie ideal L is embraced by some two-sided ideal.*
- (3) *We have $[A, L] = [A, [A, L]]$.*

Proof: It is clear that (1) implies (2). Assuming (2), let us show that (3) holds. Set $I := A[A, L]A$, and let $J \subseteq A$ be some two-sided ideal that embraces L . Then L is also related to J . It follows that $I = J$, since I is the unique two-sided ideal to which L is related, by Theorem 2.7. It follows that $L \subseteq T([A, I])$. Using this at the first step, and using Theorem 2.7 at the second step, we get

$$[A, L] \subseteq [A, I] = [A, [A, L]],$$

which verifies (3).

Assuming (3), let us show that (1) holds. Using that assumption at the first step, and Theorem 2.7 at the second step, we get

$$[A, L] = [A, [A, L]] = [A, I],$$

which verifies (1). □

The next result is essentially contained in [29, Proposition 5.7]. We only show how to remove the separability assumption. A *character* on a C^* -algebra A is a one-dimensional, irreducible representation, that is, a surjective $*$ -homomorphism $A \rightarrow \mathbb{C}$.

Proposition 2.10 (Perera, Rørdam [29]). *Let A be a unital C^* -algebra of real rank zero. Then A has no characters if and only if A admits a unital $*$ -homomorphism $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow A$.*

Proof: The backwards implication is clear. To show the forward implication, assume that A has no characters. By [22, Lemma 3.5], there exists a *separable* sub- C^* -algebra $A_0 \subseteq A$ containing the unit of A . Using that real rank zero is separably inheritable (see [6, Section II.8.5]), we obtain a *separable* sub- C^* -algebra $B \subseteq A$ of real rank zero with $A_0 \subseteq B$. It follows that B has no characters, which allows us to apply [29, Proposition 5.7] for B . We obtain a unital $*$ -homomorphism from $M_2(\mathbb{C}) \oplus M_3(\mathbb{C})$ to B , and thus also to A . □

Example 2.11. Let A be a unital C^* -algebra of real rank zero that has no characters. Then Theorem 2.7 applies to A . In particular, every Lie ideal in A is related to a (unique) two-sided ideal.

We expect that the following question has a positive answer.

Problem 2.12. Let A be a unital C^* -algebra that has no characters. Is every Lie ideal in A related to a (unique) two-sided ideal?

3. Two-sided ideals in C^* -algebras with weakly divisible unit

In this section, we prove that every two-sided ideal I in a C^* -algebra A that admits a unital $*$ -homomorphism $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow A$ satisfies $I = A[A, I]A$, and we describe $[A, I]$ as the subspace spanned by certain square-zero elements in I ; see Theorem 3.8. If I is a Dixmier ideal (in which case there is a well-defined ideal $I^{\frac{1}{2}}$ satisfying $(I^{\frac{1}{2}})^2 = I$), then $I = [A, I] + [A, I^{\frac{1}{2}}]^2$; see Theorem 3.14. In this case every element in I is a sum of square-zero elements in I and of products of pairs of square-zero elements in $I^{\frac{1}{2}}$.

The next definition generalizes the notion of orthogonally factorizable square-zero elements in a ring to a relative notion with respect to a two-sided ideal; see [16, Definition 5.1].

Definition 3.1. Let I be a two-sided ideal in a C^* -algebra A . We let $N_2(I)$ denote the square-zero elements in I , that is,

$$N_2(I) := \{x \in I : x^2 = 0\}.$$

We further set

$$\text{FN}_2(A, I) := \{x \in I : \text{there exist } a, b \in A, y \in I \text{ with } x = ayb \text{ and } ba = 0\}.$$

Lemma 3.2. Let $I \subseteq A$ be a two-sided ideal in a C^* -algebra, and let $x \in \text{FN}_2(A, I)$. Then there exist $a, b \in A_+$ and $y \in \text{FN}_2(A, I) \subseteq I$ such that

$$x = ayb \quad \text{and} \quad ab = ba = 0.$$

Proof: By definition, there are $c, d \in A$ and $z \in I$ such that

$$x = czd \quad \text{and} \quad dc = 0.$$

Let $c = v|c|$ and $d = w|d|$ the polar decompositions in A^{**} and set

$$a := |c^*|^{\frac{1}{2}}, \quad b := |d|^{\frac{1}{2}}, \quad \text{and} \quad y := |c^*|^{\frac{1}{2}}vzw|d|^{\frac{1}{2}}.$$

By properties of the polar decomposition (see, for example, [14, Proposition 2.1]), we have $|c^*|^{\frac{1}{2}}v, w|d|^{\frac{1}{2}} \in A$, and thus $y \in A$. Further, we have $x = ayb$.

Since $dc = 0$, we have $(d^*d)^n(cc^*)^m = 0$ for every $m, n \geq 1$. It follows that $p(d^*d)q(cc^*) = 0$ for all polynomials p and q with vanishing constant terms. Using functional calculus, we get $f(d^*d)g(cc^*) = 0$ for all continuous functions $f, g: \mathbb{R} \rightarrow \mathbb{R}$ with $f(0) = g(0) = 0$. In particular, we get $|d|^{\frac{1}{2}}|c^*|^{\frac{1}{2}} = 0$. Thus, we have $ba = 0$, and then also $ab = 0$.

To show that y belongs to $\text{FN}_2(A, I)$, consider the decomposition

$$y = |c^*|^{\frac{1}{4}}(|c^*|^{\frac{1}{4}}vzw|d|^{\frac{1}{4}})|d|^{\frac{1}{4}}.$$

Then $|c^*|^{\frac{1}{4}}vzw|d|^{\frac{1}{4}}$ belongs to I , and we have also seen above that $|c^*|^{\frac{1}{4}}$ and $|d|^{\frac{1}{4}}$ have zero product, whence $y \in \text{FN}_2(A, I)$. $\square$

The next result is analogous to [16, Lemma 5.2].

Lemma 3.3. Let $I \subseteq A$ be a two-sided ideal in a C^* -algebra. Then

$$\text{FN}_2(A, I) \subseteq [A, \text{FN}_2(A, I)] \subseteq [A, [A, I]] \subseteq [A, I] \quad \text{and} \quad \text{FN}_2(A, I) \subseteq N_2(I).$$

Proof: Let $x \in \text{FN}_2(A, I)$. By Lemma 3.2, we can pick $a, b \in A_+$ and $y \in I$ such that $x = ayb$ and $ba = 0$. As in the proof of Lemma 3.2, we see that $ba^{\frac{1}{2}} = 0$. It follows that $a^{\frac{1}{2}}yb \in \text{FN}_2(A, I)$ and then

$$x = ayb = [a^{\frac{1}{2}}, a^{\frac{1}{2}}yb] \in [A, \text{FN}_2(A, I)].$$

Since $\text{FN}_2(A, I)$ is a subset of I , we have $[A, \text{FN}_2(A, I)] \subseteq [A, I]$. Combining these results, and using at the last step that $[A, I]$ is a Lie ideal, we get

$$\text{FN}_2(A, I) \subseteq [A, \text{FN}_2(A, I)] \subseteq [A, [A, \text{FN}_2(A, I)]] \subseteq [A, [A, I]] \subseteq [A, I].$$

Given $x = ayb \in \text{FN}_2(A, I)$, with $a, b \in A$ and $y \in I$ such that $ba = 0$, we have

$$x^2 = (ayb)(ayb) = 0,$$

and thus $x \in N_2(I)$. $\square$

Remark 3.4. We note that Definition 3.1 and [16, Definition 5.1] agree when considering the ideal $I = A$ in a C^* -algebra A . Indeed, an element x in A belongs to $\text{FN}_2(A)$ in the sense of [16, Definition 5.1] if there exist $s, t \in A$ such that $x = st$ and $ts = 0$, while x belongs to $\text{FN}_2(A, A)$ as defined in Definition 3.1 if there exist $a, y, b \in A$ such that $x = ayb$ and $ba = 0$.

Given a factorization $x = ayb$ with $ba = 0$, we can use $s = a$ and $t = yb$. Conversely, given a factorization $x = st$ with $ts = 0$, it follows that $|t|^{\frac{1}{2}}s = 0$. We consider the polar decomposition $t = v|t|$ in A^{**} . Then $v|t|^{\frac{1}{2}}$ belongs to A , and we can use $a = s$, $y = v|t|^{\frac{1}{2}}$, and $b = |t|^{\frac{1}{2}}$.

Given a subset V in a complex vector space, we use $\text{span}_{\mathbb{C}} V$ to denote the linear subspace generated by V . Following [16, Definition 2.6], we say that a $\mathbb{C}$ -algebra A is *zero-product balanced* if for all $a, b, c \in A$ we have

$$ab \otimes c - a \otimes bc \in \text{span}_{\mathbb{C}}\{u \otimes v \in A \otimes_{\mathbb{C}} A : uv = 0\} \subseteq A \otimes_{\mathbb{C}} A.$$

This is closely related to the concept of ‘zero-product determination’, for which we refer to [7]. By [16, Corollary 2.12], a (not necessarily unital) C^* -algebra is zero-product balanced as a $\mathbb{C}$ -algebra if and only if it is zero-product determined.

We see that the class of C^* -algebras considered in this section are zero-product balanced:

Lemma 3.5. *Let A be a unital C^* -algebra that admits a unital $*$ -homomorphism $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow A$. Then A is zero-product balanced as a $\mathbb{C}$ -algebra.*

Proof: As in Paragraph 2.1, let $(e_{ij})_{i,j=1,2}$ and $(f_{ij})_{i,j=1,2,3}$ be the images in A of matrix units in $M_2(\mathbb{C})$ and $M_3(\mathbb{C})$. Then $e_{11} + f_{11}$ and $e_{22} + f_{22}$ are two full orthogonal projections in A . This allows us to apply [31, Corollary 3.8], which shows that A is generated by its projections as a $\mathbb{C}$ -algebra, and thus is zero-product determined (hence, zero-product balanced) by [7, Theorem 2.3]. $\square$

A two-sided ideal $I \subseteq A$ in a C^* -algebra is said to be *idempotent* if $I = I^2$, that is, if every element in I is of the form $x_1y_1 + \cdots + x_ny_n$ for some $n \in \mathbb{N}$ and suitable $x_j, y_j \in I$.

Proposition 3.6. *Let A be a C^* -algebra that is zero-product balanced as a $\mathbb{C}$ -algebra, and let $I \subseteq A$ be a two-sided ideal such that $I = AIA$ (for example, A is unital or I is idempotent). Then*

$$[A, I] = [A, [A, I]] = \text{span}_{\mathbb{C}} \text{FN}_2(A, I).$$

Proof: The inclusions ‘ $\supseteq$ ’ hold by Lemma 3.3. The proof of the converse inclusion is based on a combination of the methods used in the proofs of Proposition 2.14 and Theorem 5.3 in [16], which in turn are inspired by that of [7, Theorem 9.1]. To verify that $[A, I] \subseteq \text{span}_{\mathbb{C}} \text{FN}_2(A, I)$, let $x \in A$ and $y \in I$. Set $F := \text{span}_{\mathbb{C}} \text{FN}_2(A, I)$, and define

$$\varphi_y: A \times A \rightarrow I/F, \quad \varphi_y(a, b) := bya + F \quad (a, b \in A).$$

Note that φ_y is a well-defined $\mathbb{C}$ -bilinear map. Further, φ_y preserves zero-products since if $a, b \in A$ satisfy $ab = 0$, then $bya \in \text{FN}_2(A, I) \subseteq F$. Using [16, Proposition 2.8] at the second step, we obtain that

$$cyab + F = \varphi_y(ab, c) = \varphi_y(a, bc) = bcya + F$$

and thus

$$bcya - cyab \in F = \text{span}_{\mathbb{C}} \text{FN}_2(A, I)$$

for all $a, b, c \in A$. The assumption that $I = AIA$ allows us to pick n and $c_j, a_j \in A$ and $y_j \in I$ for $j = 1, \dots, n$ such that $y = \sum_j c_j y_j a_j$. Then

$$[x, y] = \sum_{j=1}^n [x, c_j y_j a_j] = \sum_{j=1}^n (x c_j y_j a_j - c_j y_j a_j x) \in \text{span}_{\mathbb{C}} \text{FN}_2(A, I),$$

which implies that $[A, I] \subseteq \text{span}_{\mathbb{C}} \text{FN}_2(A, I)$. $\square$

Problem 3.7. Does there exist a two-sided ideal I in a C^* -algebra A such that $[A, [A, I]] \neq [A, I]$?

The next result verifies statement (3) in Theorem D.

Theorem 3.8. *Let A be a unital C^* -algebra that admits a unital $*$ -homomorphism $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow A$, and let $I \subseteq A$ be a (not necessarily closed) two-sided ideal. Then*

$$I = A[A, I]A = A[A, I] \quad \text{and} \quad [A, I] = [A, [A, I]] = \text{span}_{\mathbb{C}} \text{FN}_2(A, I).$$

Proof: Considering I as a Lie ideal in A , it follows from Theorem 2.7 that $A[A, I]A$ is the unique two-sided ideal in A to which I is related. Since I is clearly related to itself, we get $I = A[A, I]A$. The inclusion $A[A, I] \subseteq A[A, I]A$ holds since A is unital. The converse inclusion holds since $[A, I]$ is a Lie ideal in A ; see, for example, [15, Lemma 3.2].

Now the equality $[A, I] = [A, [A, I]]$ also follows from Theorem 2.7. Further, since A is zero-product balanced by Lemma 3.5, we have $[A, I] = \text{span}_{\mathbb{C}} \text{FN}_2(A, I)$ by Proposition 3.6. $\square$

Remark 3.9. Let A be a unital C^* -algebra that admits a unital $*$ -homomorphism $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow A$. Applying Theorem 3.8 for the ideal A , we get

$$A = A[A, A]A,$$

which means that A is generated by its commutators as an ideal.

In fact, A is even generated by its commutators as a ring. Indeed, by [18, Theorem 5.15], every element in A is a sum of a commutator and the product of two commutators, and also the sum of three elements that are products of commutators.

The next result verifies statement (4) in Theorem D.

Theorem 3.10. *Let A be a unital C^* -algebra that admits a unital $*$ -homomorphism $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow A$. There is a natural bijection between the set $\mathcal{I}$ of two-sided ideals in A and the set*

$$\mathcal{L} := \{L \subseteq A \text{ Lie ideal} : L = [A, L]\}$$

given by the maps $I \mapsto [A, I]$ and $L \mapsto A[A, L]A$.

Proof: We denote the maps by $\alpha: \mathcal{I} \rightarrow \mathcal{L}$ and $\beta: \mathcal{L} \rightarrow \mathcal{I}$. Given $I \in \mathcal{I}$, using Theorem 3.8 at the second step, we have

$$\alpha(I) = [A, I] = [A, [A, I]] = [A, \alpha(I)]$$

which shows that α is well defined.

Next, we show that α and β are inverses of each other. First, given $I \in \mathcal{I}$, using Theorem 3.8 at the last step, we get

$$\beta(\alpha(I)) = \beta([A, I]) = A[A, I]A = I.$$

Second, given $L \in \mathcal{L}$, it follows from $L = [A, L]$ that $[A, L] = [A, [A, L]]$, and thus $L = [A, [A, L]]$. Using this at the last step, and using Theorem 2.7 at the third step, we get

$$\alpha(\beta(L)) = \alpha(A[A, L]A) = [A, A[A, L]A] = [A, [A, L]] = L,$$

as desired. $\square$

Robert showed in [31, Lemma 2.1] that every nilpotent element in a C^* -algebra is a sum of commutators. With a view to Theorem 3.8, we ask:

Problem 3.11. Let I be a (not necessarily closed) two-sided ideal in a C^* -algebra A . Does every nilpotent element in I belong to $[A, I]$?

We give positive answers to Problem 3.11 for arbitrary two-sided ideals in von Neumann algebras (Proposition 6.2), and for semiprime two-sided ideals in C^* -algebras (Proposition 4.5).

Following [14, Definition 3.2], a *Dixmier ideal* in a C^* -algebra A is a two-sided ideal $I \subseteq A$ such that I is positively spanned (that is, $I = \text{span}_{\mathbb{C}}(I \cap A_+)$) and hereditary (that is, if $a, b \in A_+$ satisfy $a \leq b$ and $b \in I$, then $a \in I$) and strongly invariant (that is, if $x \in A$ satisfies $x^*x \in I$, then $xx^* \in I$). Two-sided ideals in von Neumann algebras are automatically Dixmier ideals, which was essentially shown by Dixmier (see [14, Proposition 3.4]).

Given a Dixmier ideal I and $s \in (0, \infty)$, there is a unique Dixmier ideal whose set of positive elements is $\{a^s : a \in I \cap A_+\}$, and we use I^s to denote this ideal; see Proposition 3.7 and Definition 3.8 in [14]. We note that for $n \in \mathbb{N}$ the ring-theoretic definition of I^n as the two-sided ideal generated by $x_1 \cdots x_n$ for $x_1, \dots, x_n \in I$ agrees with the ‘new’ definition of I^n as the linear span of elements a^n for $a \in I \cap A_+$. This leads to a well-behaved theory of roots and powers for Dixmier ideals ([14, Theorem 3.9]), which we recall for the convenience of the reader.

Theorem 3.12 (Gardella, Kitamura, Thiel [14]). *Let $I \subseteq A$ be a Dixmier ideal in a C^* -algebra. Then*

$$(I^s)^t = I^{st} = (I^t)^s \quad \text{and} \quad I^s I^t = I^{s+t} = I^t I^s$$

for all $s, t \in (0, \infty)$.

Lemma 3.13. *Let $I \subseteq A$ be a Dixmier ideal in a C^* -algebra, let $x \in I$ with polar decomposition $x = v|x|$ in A^{**} , and let $s \in (0, \infty)$. Then*

$$v|x|^s, |x|^s, |x^*|^s, |x|^s v^* \in I^s.$$

Proof: Since I is a Dixmier ideal, we have $x^* \in I$, and thus $x^*x, xx^* \in I^2$. Then by Theorem 3.12 we get

$$|x| = (x^*x)^{\frac{1}{2}} \in (I^2)^{\frac{1}{2}} = I \quad \text{and} \quad |x^*| = (xx^*)^{\frac{1}{2}} \in (I^2)^{\frac{1}{2}} = I.$$

It follows that $|x|^s, |x^*|^s \in I^s$.

By [14, Proposition 3.10], given a Dixmier ideal $J \subseteq A$, an element $y \in A$ belongs to $J^{\frac{1}{2}}$ if and only if $y^*y \in J$. Applying this for the Dixmier ideal I^{2s} , and using that

$$(v|x|^s)^*(v|x|^s) = |x|^{2s} \in I^{2s},$$

we deduce that $v|x|^s \in (I^{2s})^{\frac{1}{2}} = I^s$. Since I^s is a Dixmier ideal, we further have $|x|^s v^* = (v|x|^s)^* \in I^s$. $\square$

The next result showcases a situation when every element in a Dixmier ideal I is a sum of square-zero elements in I and products of pairs of square-zero elements in $I^{\frac{1}{2}}$. In Theorem 6.4, we strengthen this result for ideals in von Neumann algebras.

Theorem 3.14. *Let A be a unital C^* -algebra that admits a unital $*$ -homomorphism $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow A$, and let $I \subseteq A$ be a Dixmier ideal. Then*

$$I = [A, I] + [A, I^{\frac{1}{2}}]^2.$$

Proof: We first show that

$$A \cdot \text{FN}_2(A, I) \subseteq [A, I] + [A, I^{\frac{1}{2}}]^2.$$

Let $c \in A$ and $x \in \text{FN}_2(A, I)$. By Lemma 3.2, we can pick $a, b \in A_+$ and $y \in I$ such that $x = ayb$ and $ab = ba = 0$. Set $z := yb \in I$, and let $z = v|z|$ be the polar decomposition in A^{**} . Then

$$cx = [ca^{\frac{1}{2}}, a^{\frac{1}{2}}z] + a^{\frac{1}{2}}zca^{\frac{1}{2}} = [ca^{\frac{1}{2}}, a^{\frac{1}{2}}z] + (a^{\frac{1}{2}}v|z|^{\frac{1}{2}})(|z|^{\frac{1}{2}}ca^{\frac{1}{2}}).$$

By Lemma 3.13, $v|z|^{\frac{1}{2}}$ and $|z|^{\frac{1}{2}}$ belong to $I^{\frac{1}{2}}$. Further, since $za = 0$, we get $|z|^{\frac{1}{2}}a^{\frac{1}{2}} = 0$, and hence

$$a^{\frac{1}{2}}v|z|^{\frac{1}{2}} = [a^{\frac{1}{2}}, v|z|^{\frac{1}{2}}] \in [A, I^{\frac{1}{2}}] \quad \text{and} \quad |z|^{\frac{1}{2}}ca^{\frac{1}{2}} = [-ca^{\frac{1}{2}}, |z|^{\frac{1}{2}}] \in [A, I^{\frac{1}{2}}].$$

It follows that cx belongs to $[A, I] + [A, I^{\frac{1}{2}}]^2$.

Using Theorem 3.8 at the first and second step, and using the above at the last step, we get

$$I = A[A, I] = \text{span}_{\mathbb{C}}(A \cdot \text{FN}_2(A, I)) \subseteq [A, I] + [A, I^{\frac{1}{2}}]^2.$$

On the other hand, we clearly have $[A, I] \subseteq I$. Using Theorem 3.12, we also have $[A, I^{\frac{1}{2}}]^2 \subseteq (I^{\frac{1}{2}})^2 = I$. $\square$

4. Square-zero elements in semiprime ideals in C^* -algebras

In this section, we study the subspace N generated by the square-zero elements in a semiprime ideal in a C^* -algebra A . We show that ANA , the two-sided ideal generated by N , satisfies $ANA = [A, N] + N^2$; see Theorem 4.3. This generalizes results from [17, Section 4].

We then characterize semiprimeness for two-sided ideals in a C^* -algebra whose unit is weakly divisible of degree 2; see Theorem 4.6. In particular, such an ideal is semiprime if and only if it is generated by its commutators as an ideal (or as a ring).

We begin with some preparatory results about commutators involving Dixmier ideals, which might be of use in future work.

Lemma 4.1. *Let $I \subseteq A$ be a Dixmier ideal in a C^* -algebra. Then*

$$[A, I] \subseteq [I^{\frac{1}{2}}, I^{\frac{1}{2}}] \subseteq I.$$

Proof: By Theorem 3.12, we have $I = I^{\frac{1}{2}}I^{\frac{1}{2}}$. Using that $[a, xy] = [ax, y] + [ya, x]$ for $a \in A$ and $x, y \in I$, it follows that

$$[A, I] = [A, I^{\frac{1}{2}}I^{\frac{1}{2}}] \subseteq [AI^{\frac{1}{2}}, I^{\frac{1}{2}}] + [I^{\frac{1}{2}}A, I^{\frac{1}{2}}] \subseteq [I^{\frac{1}{2}}, I^{\frac{1}{2}}].$$

Further, we have $[I^{\frac{1}{2}}, I^{\frac{1}{2}}] \subseteq I^{\frac{1}{2}}I^{\frac{1}{2}} = I$. $\square$

Recall that $N_2(I)$ denotes the set of square-zero elements in an ideal I .

Lemma 4.2. *Let $I \subseteq A$ be a Dixmier ideal in a C^* -algebra. Then*

$$N_2(I) \subseteq \text{span}_{\mathbb{C}}[N_2(I^{\frac{1}{2}}), N_2(I^{\frac{1}{2}})].$$

Proof: Let $x \in N_2(I)$, and let $x = v|x|$ be the polar decomposition in A^{**} . We have $x = |x^*|v = |x^*|^{\frac{1}{2}}v|x|^{\frac{1}{2}}$. Since $x^2 = 0$, the elements $|x|$ and $|x^*|$ are orthogonal, and hence so are $|x|^{\frac{1}{2}}$ and $|x^*|^{\frac{1}{2}}$. Therefore

$$x = [\tfrac{1}{2}v|x|^{\frac{1}{2}}, |x|^{\frac{1}{2}} - |x^*|^{\frac{1}{2}}].$$

By Lemma 3.13, the element $\tfrac{1}{2}v|x|^{\frac{1}{2}}$ belongs to $I^{\frac{1}{2}}$, and thus $\tfrac{1}{2}v|x|^{\frac{1}{2}} \in N_2(I^{\frac{1}{2}})$. Set

$$y := |x|^{\frac{1}{2}} - |x^*|^{\frac{1}{2}} + v|x|^{\frac{1}{2}} - v^*|x^*|^{\frac{1}{2}}.$$

It is elementary (but tedious) to check that $y^2 = 0$. Note that y belongs to $I^{\frac{1}{2}}$ by Lemma 3.13, and thus $y \in N_2(I^{\frac{1}{2}})$. Using that $v|x|^{\frac{1}{2}}$ and $v^*|x^*|^{\frac{1}{2}}$ also belong to $N_2(I^{\frac{1}{2}})$, it follows that

$$|x|^{\frac{1}{2}} - |x^*|^{\frac{1}{2}} = y - v|x|^{\frac{1}{2}} + v^*|x^*|^{\frac{1}{2}} \in \text{span}_{\mathbb{C}} N_2(I^{\frac{1}{2}}),$$

and thus

$$x = [\tfrac{1}{2}v|x|^{\frac{1}{2}}, |x|^{\frac{1}{2}} - |x^*|^{\frac{1}{2}}] \in \text{span}_{\mathbb{C}}[N_2(I^{\frac{1}{2}}), N_2(I^{\frac{1}{2}})],$$

as desired. $\square$

A two-sided ideal I in a ring R is said to be *semiprime* if for all two-sided ideals $J \subseteq R$ we have $J \subseteq I$ whenever $J^2 \subseteq I$. We refer to [23, Section 10] for details. In [14, Section 5], together with Gardella and Kitamura, we showed that a two-sided ideal I in a C^* -algebra is semiprime if and only if it is idempotent (that is, $I = I^2$), if and only if it is a Dixmier ideal and satisfies $I = I^{\frac{1}{2}}$.

The next result is a generalization of [17, Theorem 4.2] from the study of square-zero elements in C^* -algebras to square-zero elements in semiprime ideals in C^* -algebras. Recall that $\tilde{A}$ denotes the minimal unitization of a C^* -algebra A .

Theorem 4.3. *Let $I \subseteq A$ be a semiprime ideal in a C^* -algebra, let N denote the subspace generated by $N_2(I)$, and let $J := \tilde{A}N\tilde{A}$ denote the two-sided ideal of A generated by N . Then J is semiprime, and we have*

$$N = [N, N] \subseteq N^2, \quad J = [J, J]^2 = AN = [A, N]^2 = [A, N] + N^2,$$

and

$$[J, J] = [A, J] = [A, [A, N]] = [[J, J], [J, J]] = [[A, N], [A, N]].$$

Proof: We will use that I is a Dixmier ideal satisfying $I = I^{\frac{1}{2}}$. Applying Lemma 4.2 at the second step, we have

$$N = \text{span}_{\mathbb{C}} N_2(I) \subseteq \text{span}_{\mathbb{C}}[N_2(I^{\frac{1}{2}}), N_2(I^{\frac{1}{2}})] = \text{span}_{\mathbb{C}}[N_2(I), N_2(I)] = [N, N].$$

The inclusion $[N, N] \subseteq N^2$ follows since every commutator in $[N, N]$ belongs to N^2 . To verify that $[N, N] \subseteq N$, let $x, y \in N_2(I)$. Then

$$[x, y] = (1+x)y(1-x) - y + xyx.$$

Each of the three summands is a square-zero element in I , which shows $[x, y] \in N$.

We have

$$J := \tilde{A}N\tilde{A} \subseteq \tilde{A}N^2\tilde{A} \subseteq (\tilde{A}N)(N\tilde{A}) \subseteq J^2,$$

which shows that J is idempotent and therefore semiprime by [14, Theorem A].

The inclusion $AN \subseteq \tilde{A}N\tilde{A} = J$ is clear. Conversely, given $a \in \tilde{A}$, $x \in N_2(I)$, and a unitary $u \in \tilde{A}$, using that $u^*xu \in N_2(I)$ and that $N \subseteq N^2$, we have

$$axu = au(u^*xu) \in \tilde{A}N_2(I) \subseteq \tilde{A}N \subseteq \tilde{A}N^2 \subseteq AN.$$

Since every element in $\tilde{A}$ is a linear combination of four unitaries, we get $J = AN$.

The inclusion $[A, N] + N^2 \subseteq \tilde{A}N\tilde{A} = J$ is clear. For the converse inclusion, we use a similar argument as in the proof of Theorem 3.14 to show that $AN \subseteq [A, N] + N^2$. Given $a \in A$ and $x \in N_2(I)$, let $x = v|x|$ be the polar decomposition in A^{**} . Since I is semiprime, it is Dixmier and $I = I^{\frac{1}{2}} = I^{\frac{1}{4}}$. By Lemma 3.13, we have $v|x|^{\frac{1}{2}} \in I^{\frac{1}{2}}$ and thus $v|x|^{\frac{1}{2}} \in N_2(I)$. Similarly, we have $v|x|^{\frac{1}{4}}, |x|^{\frac{1}{4}}a|x^*|^{\frac{1}{2}} \in N_2(I)$. Using that $x = |x^*|^{\frac{1}{2}}v|x|^{\frac{1}{2}}$, we get

$$ax = [a|x^*|^{\frac{1}{2}}, v|x|^{\frac{1}{2}}] + (v|x|^{\frac{1}{4}})(|x|^{\frac{1}{4}}a|x^*|^{\frac{1}{2}}) \in [A, N_2(I)] + N_2(I)^2.$$

Using that $J = AN$, we get $J = [A, N] + N^2$.

The inclusion $[A, N]^2 \subseteq \tilde{A}N\tilde{A} = J$ is clear. Conversely, using that $N = [N, N]$, we have

$$[A, N] = [A, [N, N]] \subseteq [N, [A, N]] \subseteq [[A, N], [A, N]] \subseteq [A, N]^2.$$

We also have $N^2 = [N, N]^2 \subseteq [A, N]^2$ and thus

$$J = [A, N] + N^2 \subseteq [A, N]^2.$$

The inclusion $[J, J] \subseteq [A, J]$ is clear. Using that $J = J^2$ and $[a, xy] = [ax, y] + [ya, x]$ for $a \in A$ and $x, y \in J$, we have

$$[A, J] = [A, J^2] \subseteq [AJ, J] + [JA, J] \subseteq [J, J].$$

This shows that $[J, J] = [A, J]$. We get

$$[[A, N], [A, N]] \subseteq [A, [A, N]] \subseteq [A, J] = [J, J].$$

We have

$$[[A, N], N^2] \subseteq [A, N^2] \subseteq [AN, N] + [NA, N] \subseteq [A, N],$$

and similarly $[N^2, N^2] \subseteq [A, N^2] \subseteq [A, N]$. We also have

$$[A, N] = [A, [N, N]] \subseteq [N, [A, N]] = [[N, N], [A, N]] \subseteq [[A, N], [A, N]].$$

Using that $J = [A, N] + N^2$, it follows that

$$\begin{aligned} [J, J] &\subseteq [[A, N], [A, N]] + [N^2, [A, N]] + [[A, N], N^2] + [N^2, N^2] \\ &\subseteq [[A, N], [A, N]]. \end{aligned}$$

Using that $[A, N] \subseteq [A, J] = [J, J]$, we also have

$$[J, J] = [[A, N], [A, N]] \subseteq [[J, J], [J, J]].$$

Since the converse inclusion is clear, we get

$$[J, J] = [A, J] = [A, [A, N]] = [[J, J], [J, J]] = [[A, N], [A, N]].$$

Finally, since $[A, N] \subseteq [J, J]$ and $J = [A, N]^2$, we have $J \subseteq [J, J]^2$ and thus $J = [J, J]^2$. $\square$

Lemma 4.4. *Let $I \subseteq A$ be a Dixmier ideal in a C^* -algebra, and let $x \in I$ be nilpotent. Then $x \in [I^{\frac{1}{2}}, I^{\frac{1}{2}}]$.*

Proof: As in the proof of [31, Lemma 2.1], we proceed by induction on the degree of nilpotency. To start, let $x \in I$ be a square-zero element, and let $x = v|x|$ be the polar decomposition of x in A^{**} . By Lemma 3.13, the elements $v|x|^{\frac{1}{2}}$ and $|x|^{\frac{1}{2}}$ belong to $I^{\frac{1}{2}}$, and thus

$$x = [v|x|^{\frac{1}{2}}, x^{\frac{1}{2}}] \in [I^{\frac{1}{2}}, I^{\frac{1}{2}}].$$

Next, assume that for some $k \geq 2$ we have shown that every element $x \in I$ with $x^k = 0$ belongs to $[I^{\frac{1}{2}}, I^{\frac{1}{2}}]$. Let $x \in I$ satisfy $x^{k+1} = 0$, and let $x = v|x|$ be the polar decomposition of x in A^{**} . Set $y := |x|^{\frac{1}{2}}v|x|^{\frac{1}{2}}$, which belongs to A . As in the proof of [31, Lemma 2.1], one shows that $y^k = 0$. By Lemma 3.13, we have $v|x|^{\frac{1}{2}}, |x|^{\frac{1}{2}} \in I^{\frac{1}{2}}$. By assumption of the induction, we have $y \in [I^{\frac{1}{2}}, I^{\frac{1}{2}}]$, and thus

$$x = [v|x|^{\frac{1}{2}}, |x|^{\frac{1}{2}}] + y \in [I^{\frac{1}{2}}, I^{\frac{1}{2}}],$$

as desired. $\square$

Proposition 4.5. *Let $I \subseteq A$ be a semiprime ideal in a C^* -algebra, and let $x \in I$ be nilpotent. Then $x \in [A, I]$.*

Proof: Since I is semiprime, it is a Dixmier ideal and satisfies $I = I^{\frac{1}{2}}$. Applying Lemma 4.4, we have

$$x \in [I^{\frac{1}{2}}, I^{\frac{1}{2}}] \subseteq [A, I^{\frac{1}{2}}] = [A, I],$$

as desired. $\square$

Theorem 4.6. *Let A be a unital C^* -algebra that admits a unital $*$ -homomorphism $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow A$, and let $I \subseteq A$ be a two-sided ideal. The following statements are equivalent:*

- (1) *The ideal $I \subseteq A$ is semiprime.*
- (2) *We have $I = A[I, I]A$, that is, I is generated by its commutators as an ideal in A .*
- (3) *We have $I = \widetilde{I}[I, I]\widetilde{I}$, that is, I is generated by its commutators as an ideal.*
- (4) *We have $I = [I, I]^2$.*
- (5) *We have $[I, I] = [[I, I], [I, I]]$.*

Proof: To show that (2) implies (1), assume that I is generated by $[I, I]$ as an ideal in A . We clearly have $[I, I] \subseteq I^2$, and therefore

$$I = A[I, I]A \subseteq AI^2A = I^2.$$

It follows that I is idempotent, and thus semiprime by [14, Theorem A].

It is clear that (4) implies (3), which in turn implies (2).

Assume that I is semiprime, and let N denote the subspace generated by the square-zero elements in I . Applying Theorem 3.8, we have

$$I = A[A, I]A = \text{span}_{\mathbb{C}} A \text{FN}_2(A, I)A \subseteq ANA.$$

Now it follows from Theorem 4.3 that

$$I = ANA = [ANA, ANA]^2 = [I, I]^2,$$

which verifies (4), and that

$$[I, I] = [ANA, ANA] = [[ANA, ANA], [ANA, ANA]] = [[I, I], [I, I]],$$

which verifies (5).

Finally, assuming (5), let us verify (1). Using that $[a, xy] = [ax, y] + [ya, x]$ for $a \in A$ and $x, y \in I$, we have

$$[A, I^2] \subseteq [AI, I] + [IA, I] \subseteq [I, I].$$

Applying Theorem 3.8 for the ideal I^2 , we get

$$I^2 = A[A, I^2]A \subseteq A[I, I]A \subseteq A[[I, I], [I, I]]A \subseteq I^4.$$

It follows that $I^2 = I^4$, and is thus semiprime by [14, Theorem A]. $\square$

5. Lie ideals in properly infinite C^* -algebras

Let A be a C^* -algebra with a unit that is weakly divisible of degree 2, that is, that admits a unital $*$ -homomorphism $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow A$, and let $L \subseteq A$ be a (not necessarily closed) Lie ideal. We have seen in Theorem 2.7 that L is related to the two-sided ideal $I := A[A, L]A$. Further, by Theorem 2.9, L is commutator-equivalent to I if and only if $[A, L] = [A, [A, L]]$.

This raises the question of when Lie ideals L in A satisfy $[A, L] = [A, [A, L]]$. A sufficient condition is that $A = Z(A) + [A, A]$, where $Z(A)$ denotes the center of A ; see Lemma 5.1. We observe in Proposition 5.2 that this condition is satisfied in a number of situations, in particular for all properly infinite C^* -algebras.

We then prove the main result of the paper: Every Lie ideal in a unital, properly infinite C^* -algebra is commutator-equivalent to a (unique) two-sided ideal; see Theorem 5.3.

Lemma 5.1. *Let A be a unital C^* -algebra such that $A = Z(A) + [A, A]$, and let $L \subseteq A$ be a Lie ideal. Then*

$$[A, L] = [A, [A, L]].$$

Proof: The inclusion ‘ $\supseteq$ ’ is clear. Conversely, using the assumption at the first step, and using the Jacobi identity at the last step, we get

$$[A, L] = [Z(A) + [A, A], L] = [[A, A], L] \subseteq [A, [A, L]],$$

as desired. $\square$

Following Winter [32], a C^* -algebra A is said to be *pure* if its Cuntz semigroup $\text{Cu}(A)$ is almost unperforated and almost divisible. Pure C^* -algebras form a robust class ([3]) that includes every C^* -algebra that tensorially absorbs the Jiang–Su algebra $\mathcal{Z}$, that is, $A \cong A \otimes \mathcal{Z}$. By [2, Theorem 7.3.11], a C^* -algebra A is pure if and only if $\text{Cu}(A) \cong \text{Cu}(A) \otimes \text{Cu}(\mathcal{Z})$.

By a *quasitrace* of a C^* -algebra A we mean a lower-semicontinuous, $[0, \infty]$ -valued 2-quasitrace. By Haagerup’s theorem [19], every bounded quasitrace on a unital, exact C^* -algebra is a trace, and this was later extended by Kirchberg [21], who showed that arbitrary quasitraces on exact C^* -algebras are traces.

A unital C^* -algebra A is *weakly central* if the center separates maximal ideals: Maximal ideals $M_1, M_2 \subseteq A$ satisfy $M_1 = M_2$ whenever $M_1 \cap Z(A) = M_2 \cap Z(A)$. By Vesterstrøm’s theorem, A is weakly central if and only if A has the *center-quotient property*: For every closed, two-sided ideal $I \subseteq A$, the quotient map $A \rightarrow A/I$ maps $Z(A)$ onto $Z(A/I)$.

A unital C^* -algebra A has the *Dixmier property* if for every element $a \in A$ the closed convex hull of the unitary orbit $\{uau^* : u \in A \text{ unitary}\}$ has a nonempty intersection with the center $Z(A)$. It was shown in [5, Theorem 1.1] that this property holds if and only if A is weakly central, every simple quotient of A has at most one tracial state, and every extreme tracial state of A factors through a simple quotient. Dixmier showed that every von Neumann algebra has the property named after him.

Proposition 5.2. *Let A be a unital C^* -algebra. Then $A = Z(A) + [A, A]$ in each of the following cases:*

- (1) *If A is pure, every quasitrace on A is a trace, and A has the Dixmier property.*
- (2) *If A has no tracial states (in particular, if A is properly infinite).*
- (3) *If A is a von Neumann algebra.*

Proof: (1) If A has the Dixmier property, then every element in A is a Dixmier element in the sense of [4], and thus it follows from [4, Lemma 4.10] that $A = Z(A) + [A, A]$.

If A is pure and every quasitrace on A is a trace, then $[A, A]$ is closed by [28, Theorem 1.1]. Combining both results, we get $A = Z(A) + [A, A]$.

(2) By [30, Theorem 1], we have $A = [A, A]$ if (and only if) A has no tracial states.

(3) It was shown in the proof of [8, Theorem 5.19] that every von Neumann algebra A satisfies $A = Z(A) + [A, A]$. $\square$

Theorem 5.3. *Let A be a unital, properly infinite C^* -algebra, and let $L \subseteq A$ be a Lie ideal. Then for the two-sided ideal $I := A[A, L]A$ we have*

$$[A, L] = [A, [A, L]] = [A, I].$$

In particular, L is commutator-equivalent to I , and hence also embraced by I , and related to I .

Further, I is the only two-sided ideal of A to which L is related.

Proof: Since A is unital and properly infinite, there exists a unital $*$ -homomorphism from the Cuntz algebra $\mathcal{O}_\infty$ to A ; see [6, Proposition III.1.3.3]. Further, it is well known that $\mathcal{O}_\infty$ admits a unital $*$ -homomorphism from $M_2(\mathbb{C}) \oplus M_3(\mathbb{C})$, using for example Proposition 2.10 and that $\mathcal{O}_\infty$ is a unital C^* -algebra of real rank zero that admits no characters. It follows that A admits a unital $*$ -homomorphism $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow A$.

We may therefore apply Theorem 2.7 and obtain that L is related to I , and that I is the only two-sided ideal to which L is related. Further, since A is unital and properly infinite, we have $A = [A, A]$ by Pop's theorem [30], and thus $[A, L] = [A, [A, L]]$ by Lemma 5.1. It therefore follows from Theorem 2.9 that L is commutator-equivalent to I (and thus also embraced by I). $\square$

Corollary 5.4. *Let A be a unital, properly infinite C^* -algebra. Then an additive subgroup $L \subseteq A$ is a Lie ideal if and only if $[A, I] \subseteq L \subseteq T([A, I])$ for some (unique) two-sided ideal $I \subseteq A$. We get a decomposition of the family $\mathcal{L}$ of all Lie ideals in A as:*

$$\mathcal{L} = \bigsqcup_{I \in \mathcal{I}} \{L \subseteq A \text{ additive subgroup} : [A, I] \subseteq L \subseteq T([A, I])\},$$

where $\mathcal{I}$ denotes the lattice of two-sided ideals in A .

Remark 5.5. The proof of Theorem 5.3 can easily be adapted to show the following: In a C^* -algebra A that admits a unital $*$ -homomorphism $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow A$ and such that $A = Z(A) + [A, A]$, every Lie ideal L is commutator-equivalent to the two-sided ideal $A[A, L]A$.

We note that a unital C^* -algebra admits no characters whenever it is pure or has no tracial states. Therefore, with a view to Problem 2.12 and Proposition 5.2, we expect that every Lie ideal L is commutator-equivalent to $A[A, L]A$ whenever A is a unital, pure C^* -algebra with the Dixmier property and such that every quasitrace on A is a trace, or whenever A is unital and has no tracial states.

Problem 5.6. Given Lie ideals $L, M \subseteq A$ in a unital, properly infinite C^* -algebra, what is the relationship between the two-sided ideals generated by $[L, M]$, by $[A, [L, M]]$, and by $[A, L]$ and $[A, M]$?

6. Lie ideals and two-sided ideals in von Neumann algebras

In this section, we show that every (not necessarily closed) Lie ideal L in a von Neumann algebra M is commutator-equivalent to the two-sided ideal $I := M[M, L]M$, that is, $[M, L] = [M, I]$; see Theorem 6.1. This recovers [8, Theorem 5.19]. Moreover, if M has zero commutative summand, then we show that I is the only two-sided ideal to which L is related, which solves [8, Problem 5.21].

Given a two-sided ideal I in a von Neumann algebra with zero commutative summand, every element in I is a sum of products of pairs of square-zero elements in $I^{\frac{1}{2}}$; see Theorem 6.4.

Theorem 6.1. *Let M be a von Neumann algebra, and let $L \subseteq M$ be a Lie ideal. Then, for the two-sided ideal $I := M[M, L]M$, we have*

$$[M, L] = [M, [M, L]] = [M, I].$$

In particular, L is commutator-equivalent to I , and hence also embraced by I , and related to I .

If M has zero commutative summand, then I is the only two-sided ideal of M to which L is related.

Proof: Let $M = M_0 \oplus M_1$ be the (unique) decomposition of M into a commutative summand M_0 and a summand M_1 that admits no characters. By Proposition 2.10, there is a unital $*$ -homomorphism $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow M_1$. Set

$$L_1 := M_1 \cap L \quad \text{and} \quad I_1 := M_1 \cap I.$$

Then L_1 is a Lie ideal in M_1 , and we have

$$[M, L] = [M_1, L_1] \quad \text{and} \quad I = I_1 = M_1[M_1, L_1]M_1.$$

We may apply Theorem 2.7 for the Lie ideal L_1 in M_1 and obtain that L_1 is related to I_1 , and that I_1 is the only two-sided ideal of M_1 to which L_1 is related. It follows that L is related to I , and if M_0 is zero, then I is also unique with this property. (If M_0 is nonzero, then $M_0 \oplus I$ is another two-sided ideal of M to which L is related.)

As shown in the proof of [8, Theorem 5.19], we have $M = Z(M) + [M, M]$. It follows that $[M, L] = [M, [M, L]]$ by Lemma 5.1, and then

$$[M, L] = [M, [M, L]] = [M, I]$$

by Theorem 2.9. □

Recall that $N_2(I)$ denotes the square-zero elements, and we use $\text{Nil}(I)$ to denote the set of nilpotent elements in an ideal I . Recall the definition of $\text{FN}_2(M, I)$ from Definition 3.1.

Proposition 6.2. *Let $I \subseteq M$ be a two-sided ideal in a von Neumann algebra. Then*

$$[M, I] = \text{span}_{\mathbb{C}} N_2(I) = \text{span}_{\mathbb{C}} \text{Nil}(I) = \text{span}_{\mathbb{C}} \text{FN}_2(M, I).$$

Proof: Let $M = M_0 \oplus M_1$ denote the (unique) decomposition of M into a commutative summand M_0 and a summand M_1 that admits no characters. Then

$$\begin{aligned} [M, I] &= [M_1, M_1 \cap I], & N_2(I) &= N_2(M_1 \cap I), \\ \text{Nil}(I) &= \text{Nil}(M_1 \cap I), & \text{FN}_2(M, I) &= \text{FN}_2(M_1, M_1 \cap I), \end{aligned}$$

which allows us to reduce the problem to studying the ideal $M_1 \cap I$ in M_1 .

Without loss of generality, we may therefore assume that M has zero commutative summand. Then M admits a unital $*$ -homomorphism $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow M$ by Proposition 2.10. Using Theorem 3.8 at the first step, we get

$$[M, I] = \text{span}_{\mathbb{C}} \text{FN}_2(M, I) \subseteq \text{span}_{\mathbb{C}} \text{N}_2(I) \subseteq \text{span}_{\mathbb{C}} \text{Nil}(I).$$

It remains to show that $\text{Nil}(I) \subseteq [M, I]$. As in the proof of Lemma 4.4 (and [31, Lemma 2.1]), we proceed by induction on the degree of nilpotency. To start, let $x \in I$ be a square-zero element. Let $x = v|x|$ be the polar decomposition in M . Then $|x| = v^*x \in I$. Since $x^2 = 0$, we have $|x|v = 0$, and thus

$$x = [v, |x|] \in [M, I].$$

Next, assume that for some $k \geq 2$ we have shown that every element $x \in I$ with $x^k = 0$ belongs to $[M, I]$. Let $x \in I$ satisfy $x^{k+1} = 0$. Again, let $x = v|x|$ be the polar decomposition in M . We will show that the element $y := |x|v$ satisfies $y^k = 0$. Since $x^{k+1} = 0$ and $|x| = v^*v|x|$, we have

$$y^k|x| = |x|v \cdots |x|v|x| = v^*v|x|v \cdots |x|v|x| = v^*x^{k+1} = 0.$$

For every polynomial p with vanishing constant term it follows that $y^k p(|x|) = 0$. Using functional calculus, we get $y^k |x|^{\frac{1}{n}}$ for every $n \geq 1$. The sequence $(|x|^{\frac{1}{n}})_n$ converges in the weak*-topology of M to v^*v , which is the support projection of $|x|$. We deduce that $y^k v^*v = 0$. Using that $v = vv^*v$, we get

$$y^k = y^{k-1}|x|v = y^{k-1}|x|vv^*v = y^k v^*v = 0.$$

By assumption of the induction, we have $y \in [M, I]$, and thus

$$x = [v, |x|] + |x|v \in [M, I],$$

as desired. $\square$

Remark 6.3. Let H be a separable, infinite-dimensional Hilbert space, and let I be a two-sided ideal in the von Neumann algebra $\mathcal{B}(H)$. Given a nilpotent element $x \in I$, it follows from Proposition 6.2 that $x \in [\mathcal{B}(H), I]$, that is, x is a finite sum of commutators of an element in $\mathcal{B}(H)$ and an element in I . Dykema and Krishnaswamy-Usha showed in [12, Proposition 3.1] that one summand suffices, that is, we have $x = [y, z]$ for some $y \in \mathcal{B}(H)$ and $z \in I$. Does their result hold for nilpotent elements in two-sided ideals in arbitrary von Neumann algebras?

Theorem 6.4. *Let M be a von Neumann algebra with zero commutative summand, and let $I \subseteq M$ be a two-sided ideal. Then*

$$I = [M, I^{\frac{1}{2}}]^2 = [I^{\frac{1}{4}}, I^{\frac{1}{4}}]^2,$$

and

$$[M, I] \subseteq [[M, I^{\frac{1}{2}}], [M, I^{\frac{1}{2}}]] \subseteq [[I^{\frac{1}{4}}, I^{\frac{1}{4}}], [I^{\frac{1}{4}}, I^{\frac{1}{4}}]] \subseteq [I^{\frac{1}{2}}, I^{\frac{1}{2}}].$$

Proof: We will use that I is a Dixmier ideal, since two-sided ideals in von Neumann algebras are automatically Dixmier ideals ([14, Proposition 3.4]). Using Proposition 6.2 at the first and last step, and using Lemma 4.2 at the second step, we have

$$[M, I] = \text{span}_{\mathbb{C}} \text{N}_2(I) \subseteq \text{span}_{\mathbb{C}} [\text{N}_2(I^{\frac{1}{2}}), \text{N}_2(I^{\frac{1}{2}})] = [[M, I^{\frac{1}{2}}], [M, I^{\frac{1}{2}}]].$$

Using Lemma 4.1, we then get

$$[M, I] \subseteq [[M, I^{\frac{1}{2}}], [M, I^{\frac{1}{2}}]] \subseteq [[I^{\frac{1}{4}}, I^{\frac{1}{4}}], [I^{\frac{1}{4}}, I^{\frac{1}{4}}]] \subseteq [I^{\frac{1}{2}}, I^{\frac{1}{2}}].$$

By Proposition 2.10, there is a unital $*$ -homomorphism $M_2(\mathbb{C}) \oplus M_3(\mathbb{C}) \rightarrow M$. We can therefore apply Theorem 3.14 at the first step, and together with the above we obtain that

$$I = [M, I] + [M, I^{\frac{1}{2}}]^2 \subseteq [M, I^{\frac{1}{2}}]^2.$$

Applying Lemma 4.1 for the Dixmier ideal $I^{\frac{1}{2}}$, and using Theorem 3.12 at the last step, we get

$$I \subseteq [M, I^{\frac{1}{2}}]^2 \subseteq [I^{\frac{1}{4}}, I^{\frac{1}{4}}]^2 \subseteq (I^{\frac{1}{2}})^2 = I,$$

as desired. $\square$

Remark 6.5. It is likely that the inclusions involving $[M, I]$ in Theorem 6.4 are equalities. The key question is whether $[I^{\frac{1}{2}}, I^{\frac{1}{2}}]$ is a subset of $[M, I]$. Using that I is a Dixmier ideal and therefore admits a well-behaved theory of roots and powers (Theorem 3.12), we have

$$[I^{\frac{1}{2}}, I^{\frac{1}{2}}] = [I^{\frac{1}{4}} I^{\frac{1}{4}}, I^{\frac{1}{2}}] \subseteq [I^{\frac{1}{4}}, I^{\frac{1}{4}} I^{\frac{1}{2}}] + [I^{\frac{1}{2}}, I^{\frac{1}{2}} I^{\frac{1}{4}}] = [I^{\frac{1}{4}}, I^{\frac{3}{4}}],$$

and by an analogous argument $[I^{\frac{1}{4}}, I^{\frac{3}{4}}] \subseteq [I^{\frac{1}{8}}, I^{\frac{7}{8}}]$. Inductively, it follows that $[I^{\frac{1}{2}}, I^{\frac{1}{2}}] \subseteq [I^{\frac{1}{2^n}}, I^{\frac{2^n-1}{2^n}}]$ for every $n \geq 1$, and thus

$$[I^{\frac{1}{2}}, I^{\frac{1}{2}}] \subseteq \bigcap_{n \geq 1} [I^{\frac{1}{2^n}}, I^{\frac{2^n-1}{2^n}}] \subseteq \bigcap_{\epsilon > 0} [M, I^{1-\epsilon}].$$

It is, however, not clear if $\bigcap_{\epsilon > 0} [M, I^{1-\epsilon}] = [M, I]$.

For two-sided ideals $I, J \subseteq \mathcal{B}(H)$, it was shown in [11, Theorem 5.10] that $[\mathcal{B}(H), IJ] = [I, J]$. In particular, we have $[\mathcal{B}(H), I] = [I^{\frac{1}{2}}, I^{\frac{1}{2}}]$, and we obtain the following consequence of Theorem 6.4:

Corollary 6.6. *Let H be a separable, infinite-dimensional Hilbert space, and let I be a two-sided ideal in $\mathcal{B}(H)$. Then*

$$[\mathcal{B}(H), I] = [[\mathcal{B}(H), I^{\frac{1}{2}}], [\mathcal{B}(H), I^{\frac{1}{2}}]] = [[I^{\frac{1}{4}}, I^{\frac{1}{4}}], [I^{\frac{1}{4}}, I^{\frac{1}{4}}]] = [I^{\frac{1}{2}}, I^{\frac{1}{2}}].$$

Problem 6.7. Let $I, J \subseteq M$ be two-sided ideals in a von Neumann algebra. Do we have $[M, IJ] = [I, J]$?

Acknowledgments. The author was partially supported by the Knut and Alice Wallenberg Foundation (KAW 2021.0140).

The author thanks Leonel Robert for valuable comments regarding the Dixmier property for C^* -algebras, and Matej Brešar for helpful feedback on an earlier version of this paper. The author is also grateful to the anonymous referees for their insightful suggestions, in particular for pointing out a consequence of the results that is now included as Corollary 2.8.

References

- [1] T. AMRUTAM, D. GAO, S. KUNNAWALKAM ELAYAVALLI, AND G. PATCHELL, Strict comparison in reduced group C^* -algebras, *Invent. Math.* **242(3)** (2025), 639–657. DOI: 10.1007/s00222-025-01366-5.
- [2] R. ANTOINE, F. PERERA, AND H. THIEL, Tensor products and regularity properties of Cuntz semigroups, *Mem. Amer. Math. Soc.* **251(1199)** (2018), 191 pp. DOI: 10.1090/memo/1199.
- [3] R. ANTOINE, F. PERERA, H. THIEL, AND E. VILALTA, Pure C^* -algebras, Preprint (2024). arXiv:2406.11052.
- [4] R. J. ARCHBOLD, I. GOGIĆ, AND L. ROBERT, Local variants of the Dixmier property and weak centrality for C^* -algebras, *Int. Math. Res. Not. IMRN* **2023(2)** (2023), 1483–1513. DOI: 10.1093/imrn/rnab275.

- [5] R. ARCHBOLD, L. ROBERT, AND A. TIKUISIS, The Dixmier property and tracial states for C^* -algebras, *J. Funct. Anal.* **273**(8) (2017), 2655–2718. DOI: 10.1016/j.jfa.2017.06.026.
- [6] B. BLACKADAR, *Operator Algebras. Theory of C^* -algebras and von Neumann algebras*, Encyclopaedia Math. Sci. **122**, Oper. Alg. Non-commut. Geom. III, Springer-Verlag, Berlin, 2006. DOI: 10.1007/3-540-28517-2.
- [7] M. BREŠAR, *Zero Product Determined Algebras*, Front. Math., Birkhäuser/Springer, Cham, 2021. DOI: 10.1007/978-3-030-80242-4.
- [8] M. BREŠAR, E. KISSIN, AND V. S. SHULMAN, Lie ideals: from pure algebra to C^* -algebras, *J. Reine Angew. Math.* **623** (2008), 73–121. DOI: 10.1515/CRELLE.2008.073.
- [9] E. BREUILLARD, M. KALANTAR, M. KENNEDY, AND N. OZAWA, C^* -simplicity and the unique trace property for discrete groups, *Publ. Math. Inst. Hautes Études Sci.* **126** (2017), 35–71. DOI: 10.1007/s10240-017-0091-2.
- [10] J. CUNTZ AND G. K. PEDERSEN, Equivalence and traces on C^* -algebras, *J. Functional Analysis* **33**(2) (1979), 135–164. DOI: 10.1016/0022-1236(79)90108-3.
- [11] K. DYKEMA, T. FIGIEL, G. WEISS, AND M. WODZICKI, Commutator structure of operator ideals, *Adv. Math.* **185**(1) (2004), 1–79. DOI: 10.1016/S0001-8708(03)00141-5.
- [12] K. DYKEMA AND A. KRISHNASWAMY-USHA, Nilpotent elements of operator ideals as single commutators, *Proc. Amer. Math. Soc.* **146**(7) (2018), 3031–3037. DOI: 10.1090/proc/13987.
- [13] C. K. FONG, C. R. MIERS, AND A. R. SOUROUR, Lie and Jordan ideals of operators on Hilbert space, *Proc. Amer. Math. Soc.* **84**(4) (1982), 516–520. DOI: 10.2307/2044026.
- [14] E. GARDELLA, K. KITAMURA, AND H. THIEL, Semiprime ideals in C^* -algebras, *J. Eur. Math. Soc.* (2025), published online first. DOI: 10.4171/JEMS/1699.
- [15] E. GARDELLA, T.-K. LEE, AND H. THIEL, Fully noncentral Lie ideals and invariant additive subgroups in rings, *J. Lond. Math. Soc. (2)* **111**(3) (2025), Paper no. e70127, 22 pp. DOI: 10.1112/jlms.70127.
- [16] E. GARDELLA AND H. THIEL, Zero-product balanced algebras, *Linear Algebra Appl.* **670** (2023), 121–153. DOI: 10.1016/j.laa.2023.04.015.
- [17] E. GARDELLA AND H. THIEL, Prime ideals in C^* -algebras and applications to Lie theory, *Proc. Amer. Math. Soc.* **152**(9) (2024), 3647–3656. DOI: 10.1090/proc/16808.
- [18] E. GARDELLA AND H. THIEL, Rings and C^* -algebras generated by commutators, *J. Algebra* **662** (2025), 214–241. DOI: 10.1016/j.jalgebra.2024.08.020.
- [19] U. HAAGERUP, Quasitraces on exact C^* -algebras are traces, *C. R. Math. Acad. Sci. Soc. R. Can.* **36**(2-3) (2014), 67–92.
- [20] I. N. HERSTEIN, On the Lie and Jordan rings of a simple associative ring, *Amer. J. Math.* **77**(2) (1955), 279–285. DOI: 10.2307/2372531.
- [21] E. KIRCHBERG, On the existence of traces on exact stably projectionless simple C^* -algebras, in: *Operator Algebras and Their Applications*, Fields Inst. Commun. **13**, American Mathematical Society, Providence, RI, 1997, pp. 171–172. DOI: 10.1090/fic/013/09.
- [22] E. KIRCHBERG AND M. RØRDAM, When central sequence C^* -algebras have characters, *Internat. J. Math.* **26**(7) (2015), 1550049, 32 pp. DOI: 10.1142/S0129167X15500494.
- [23] T. Y. LAM, *A First Course in Noncommutative Rings*, Second edition, Grad. Texts in Math. **131**, Springer-Verlag, New York, 2001. DOI: 10.1007/978-1-4419-8616-0.
- [24] L. MARCOUX, On the closed Lie ideals of certain C^* -algebras, *Integral Equations Operator Theory* **22**(4) (1995), 463–475. DOI: 10.1007/BF01203386.
- [25] L. W. MARCOUX, Projections, commutators and Lie ideals in C^* -algebras, *Math. Proc. R. Ir. Acad.* **110A**(1) (2010), 31–55. DOI: 10.3318/PRIA.2010.110.1.31.
- [26] L. W. MARCOUX AND G. J. MURPHY, Unitarily-invariant linear spaces in C^* -algebras, *Proc. Amer. Math. Soc.* **126**(12) (1998), 3597–3605. DOI: 10.1090/S0002-9939-98-04934-X.
- [27] P. W. NG, Commutators in the Jiang–Su algebra, *Internat. J. Math.* **23**(11) (2012), 1250113, 29 pp. DOI: 10.1142/S0129167X12501133.
- [28] P. W. NG AND L. ROBERT, Sums of commutators in pure C^* -algebras, *Münster J. Math.* **9**(1) (2016), 121–154. DOI: 10.17879/35209721075.
- [29] F. PERERA AND M. RØRDAM, AF-embeddings into C^* -algebras of real rank zero, *J. Funct. Anal.* **217**(1) (2004), 142–170. DOI: 10.1016/j.jfa.2004.05.001.
- [30] C. POP, Finite sums of commutators, *Proc. Amer. Math. Soc.* **130**(10) (2002), 3039–3041. DOI: 10.1090/S0002-9939-02-06484-5.
- [31] L. ROBERT, On the Lie ideals of C^* -algebras, *J. Operator Theory* **75**(2) (2016), 387–408. DOI: 10.7900/jot.2015may17.2070.
- [32] W. WINTER, Nuclear dimension and $\mathcal{Z}$ -stability of pure C^* -algebras, *Invent. Math.* **187**(2) (2012), 259–342. DOI: 10.1007/s00222-011-0334-7.

Department of Mathematical Sciences, Chalmers University of Technology and University of Gothenburg, Gothenburg SE-412 96, Sweden

E-mail address: `hannes.thiel@chalmers.se`

URL: `www.hannesthiel.org`

ORCID: 0000-0003-0388-6495

Received on January 7, 2025.

Accepted on June 16, 2025.